

End-of-Life Use of Systemic Therapy in Patients With Advanced Melanoma

Citation for published version (APA):

van Breeschoten, J., Ismail, R. K., Wouters, M. W. J. M., Hilarius, D. L., de Wreede, L. C., Haanen, J. B., Blank, C. U., Aarts, M. J. B., van den Berkmortel, F. W. P. J., de Groot, J. W. B., Hospers, G. A. P., Kapiteijn, E., Piersma, D., van Rijn, R. S., Stevense-den Boer, M. A., van der Veldt, A. A. M., Vreugdenhil, G., Boers-Sonderen, M. J., Suijkerbuijk, K. P. M., & van den Eertwegh, A. J. M. (2022). End-of-Life Use of Systemic Therapy in Patients With Advanced Melanoma: A Nationwide Cohort Study. *JCO Oncology Practice*, 18(10), e1611-e1620. <https://doi.org/10.1200/OP.22.00061>

Document status and date:

Published: 01/10/2022

DOI:

[10.1200/OP.22.00061](https://doi.org/10.1200/OP.22.00061)

Document Version:

Publisher's PDF, also known as Version of record

Document license:

Taverne

Please check the document version of this publication:

- A submitted manuscript is the version of the article upon submission and before peer-review. There can be important differences between the submitted version and the official published version of record. People interested in the research are advised to contact the author for the final version of the publication, or visit the DOI to the publisher's website.
- The final author version and the galley proof are versions of the publication after peer review.
- The final published version features the final layout of the paper including the volume, issue and page numbers.

[Link to publication](#)

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal.

If the publication is distributed under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license above, please follow below link for the End User Agreement:

www.umlib.nl/taverne-license

Take down policy

If you believe that this document breaches copyright please contact us at:

repository@maastrichtuniversity.nl

providing details and we will investigate your claim.

End-of-Life Use of Systemic Therapy in Patients With Advanced Melanoma: A Nationwide Cohort Study

Jesper van Breeschoten, PharmD^{1,2}; Rawa K. Ismail, PharmD^{1,3}; Michel W.J.M. Wouters, MD, PhD^{1,4,5}; Doranne L. Hilarius, PharmD, PhD⁶; Liesbeth C. de Wreede, PhD⁵; John B. Haanen, MD, PhD^{7,8,9}; Christian U. Blank, MD, PhD^{7,8}; Maureen J.B. Aarts, MD, PhD¹⁰; Franchette W.P.J. van den Berkmortel, MD, PhD¹¹; Jan Willem B. de Groot, MD, PhD¹²; Geke A.P. Hospers, MD, PhD¹³; Ellen Kapiteijn, MD, PhD⁹; Djura Piersma, MD, PhD¹⁴; Rozemarijn S. van Rijn, MD, PhD¹⁵; Marion A. Stevense-den Boer, MD, PhD¹⁶; Astrid A.M. van der Veldt, MD, PhD¹⁷; Gerard Vreugdenhil, MD, PhD¹⁸; Marye J. Boers-Sonderen, MD, PhD¹⁹; Karijn P.M. Suijkerbuijk, MD, PhD²⁰; and Alfons J.M. van den Eertwegh, MD, PhD²

QUESTION ASKED: What percentage of Dutch patients with advanced melanoma start a new systemic therapy within 45 and 90 days of death and does this percentage differ between melanoma centers?

SUMMARY ANSWER: A total of 503 (13.2%) patients with advanced melanoma, who died between 2013 and 2019, received a new systemic therapy \leq 45 days before death and 29% of all patients started a new systemic therapy within the last 90 days before death (N = 1,120). The percentage of patients receiving a new systemic therapy within 45 and 90 days before death was significantly different between Dutch melanoma centers.

WHAT WE DID: We selected all patients diagnosed with unresectable stage IIIc or IV melanoma that were registered in the Dutch Melanoma Treatment Registry who died between 2013 and 2019. Retrospectively, we investigated what percentage of patients received a new systemic therapy in the last 45 and 90 days before death. Practice variation between centers was assessed using funnel plots using with 95% and 99% confidence limits. We also investigated the type of systemic therapies started, adverse events (AEs) in the last phase of life, and costs associated with these systemic therapies.

WHAT WE FOUND: In the Netherlands, a minority of Dutch patients with advanced melanoma started a new systemic therapy in the last phase of life. However, the percentages varied between Dutch melanoma centers.

BIAS AND CONFOUNDING FACTORS: The current treatment landscape is different from 2013 to 2014 when anti-programmed cell death protein-1 and BRAF/MEK inhibitors were not available. Secondary, this study does not provide insights on which patients are unlikely to benefit from starting a new systemic therapy as we retrospectively selected patients who died.

REAL-LIFE IMPLICATIONS: The results of this study are important for both clinicians and patients. Clinicians might be unaware of the percentage of patients they are treating in the last phase of life with a new systemic therapy and how this percentage compares with other centers. To better understand the rationale of starting a new systemic therapies, comparisons between individual centers should take place. For patients, it is important to know that severe AEs in the last phase of life seem rare. If the patient understands that treatment may have a positive effect on the course of disease, and agrees with the risk of potentially severe AEs, there is no reason to not start a new systemic therapy.

ASSOCIATED CONTENT

Data Supplement

Author affiliations and disclosures are available with the complete article at ascopubs.org/journal/op.

Accepted on June 17, 2022 and published at ascopubs.org/journal/op on August 9, 2022.

Full-length article available online at DOI <https://doi.org/10.1200/OP.22.00061>

CORRESPONDING AUTHOR

Jesper van Breeschoten, PharmD, Rijnsburgerweg 10, 2333AA Leiden, the Netherlands; e-mail: j.vanbreeschoten@dica.nl.

End-of-Life Use of Systemic Therapy in Patients With Advanced Melanoma: A Nationwide Cohort Study

Jesper van Breeschoten, PharmD^{1,2}; Rawa K. Ismail, PharmD^{1,3}; Michel W.J.M. Wouters, MD, PhD^{1,4,5}; Doranne L. Hilarius, PharmD, PhD⁶; Liesbeth C. de Wreede, PhD⁵; John B. Haanen, MD, PhD^{7,8,9}; Christian U. Blank, MD, PhD^{7,8}; Maureen J.B. Aarts, MD, PhD¹⁰; Franchette W.P.J. van den Berkortel, MD, PhD¹¹; Jan Willem B. de Groot, MD, PhD¹²; Geke A.P. Hospers, MD, PhD¹³; Ellen Kapiteijn, MD, PhD⁹; Djura Piersma, MD, PhD¹⁴; Rozemarijn S. van Rijn, MD, PhD¹⁵; Marion A. Stevense-den Boer, MD, PhD¹⁶; Astrid A.M. van der Veldt, MD, PhD¹⁷; Gerard Vreugdenhil, MD, PhD¹⁸; Marye J. Boers-Sonderen, MD, PhD¹⁹; Karijn P.M. Suijkerbuijk, MD, PhD²⁰; and Alfons J.M. van den Eertwegh, MD, PhD²

abstract

PURPOSE The introduction of immune checkpoint inhibitors and targeted therapies improved the overall survival of patients with advanced melanoma. It is not known how often these costly treatments with potential serious side effects are ineffectively applied in the last phase of life. This study aimed to investigate the start of a new systemic therapy within 45 and 90 days of death in Dutch patients with advanced melanoma.

METHODS We selected patients who were diagnosed with unresectable IIIC or stage IV melanoma, registered in the Dutch Melanoma Treatment Registry, and died between 2013 and 2019. Primary outcome was the probability of starting a new systemic therapy 45 and 90 days before death. Secondary outcomes were type of systemic therapy started, grade 3/4 adverse events (AEs), and the total costs of systemic therapies.

RESULTS Between 2013 and 2019, 3,797 patients with unresectable IIIC or stage IV melanoma were entered in the registry and died. The percentage of patients receiving a new systemic therapy within 45 and 90 days before death was significantly different between Dutch melanoma centers (varying from 6% to 23% and 20% to 46%, respectively). Thirteen percent of patients (n = 146) developed grade 3/4 AEs in the last period before death. The majority of patients with an AE required hospital admission (n = 102, 69.6%). Mean total costs of systemic therapy per cohort year of the patients who received a new systemic therapy within 90 days before death were 2.3%-2.8% of the total costs spent on melanoma therapies.

CONCLUSION The minority of Dutch patients with metastatic melanoma started a new systemic therapy in the last phase of life. However, the percentages varied between Dutch melanoma centers. Financial impact of these therapies in the last phase of life is relatively small.

JCO Oncol Pract 00. © 2022 by American Society of Clinical Oncology

INTRODUCTION

Clinical outcomes of patients with advanced melanoma have improved in recent years because of the introduction of new systemic therapies such as anti-programmed cell death protein-1 (PD-1) antibodies,^{1,2} BRAF/MEK inhibitors,^{3,4} anti-cytotoxic T-cell lymphocyte-4 inhibitors (ipilimumab),⁵ and combination therapy of ipilimumab plus nivolumab.⁶ However, a large proportion of these patients still do not achieve long-term remission and disease control. In addition, severe adverse events (AEs) caused by these systemic therapies are common.

Grade 3/4 AEs occur in $\pm 10\%$ of patients receiving monotherapy anti-PD-1 antibodies,^{1,2} $\pm 30\%$ of patients receiving ipilimumab,^{2,5} $\pm 60\%$ in patients receiving combination therapy of ipilimumab plus nivolumab,⁶ and 48%-50% of patients receiving BRAF/MEK

inhibitors.^{3,4} Not only are these AEs common, but they can also influence the quality of life of patients.⁷ Quality of life is especially important at the end of life where newly initiated therapies should ideally not cause any harm. The randomized clinical trials investigating these new therapies for advanced melanoma have measured quality of life as part of the marketing authorization process. However, these trials included relatively healthy patients,⁸ leading to a gap in knowledge in using these systemic therapies at the end of life. Another downside of these systemic therapies is the financial burden for the health care system. The costs of the systemic therapies are high, and the initiation of a new treatment at the end of life with uncertainty about the effect should be well considered.

Clinicians treating patients with advanced melanoma deal with a complex disease with uncertainty about the

Author affiliations and support information (if applicable) appear at the end of this article.

Accepted on June 17, 2022 and published at ascopubs.org/journal/op on August 9, 2022; DOI <https://doi.org/10.1200/OP.22.00061>

prognosis, progression, side effects, and possible outcomes.^{9,10} Treatment with BRAF/MEK inhibitors can provide patients with symptom relief in the last phase beyond progression as palliative therapy. A recent study shows that the use of systemic therapies in the last 30 days of life declined compared with the period before introduction of immune checkpoint inhibitors.¹¹ Little is known on the use of systemic therapies at the end of life in patients with advanced melanoma in specific patient subgroups. Retrospective evaluation of its effect might minimize unnecessary exposure to possible AEs and overtreatment at the end of life; this is important, also with high costs of therapy in mind. This study aimed to describe (1) the use of systemic therapies at the end of life of Dutch patients with advanced melanoma, (2) grade 3/4 AEs in the last phase of life, and (3) assess the costs of patients treated in the last phase of life.

METHODS

Patients

The study population consisted of patients with unresectable stage IIIc or stage IV melanoma (≥ 18 years) treated with systemic therapy who died between 2013 and 2019 (data set cutoff date was December 5, 2020). Only patients with a known date of death were included. Patients participating in clinical trials were also included.

Data Sources

This study used data from the Dutch Melanoma Treatment Registry (DMTR). Since July 1, 2013, all patients with unresectable stage IIIc or stage IV melanoma seen in one of the 14 melanoma centers have been registered in the DMTR. Trained data managers follow patients every 3 months during the first year and register changes in disease status, patients' characteristics, or treatment characteristics. In the following years, if a patient is clinically stable, follow-up is performed every 6 months. In case of progression, follow-up is performed every 4 months. A detailed description of the DMTR setup has been published by Jochems et al.¹² In compliance with Dutch regulations, the DMTR was approved by a medical ethical committee (METC Leiden University Medical Center, 2013) and is not considered subject to the Medical Research Involving Human Subjects Act.

Primary and Secondary Outcomes

The primary outcome was the probability of starting a new systemic therapy within 45 and 90 days before the date of death. If patients started multiple systemic therapies, we selected the most recent systemic therapy and corresponding patient and tumor characteristics. Systemic therapies included anti-PD-1, BRAF/MEK inhibitors, ipilimumab, combination therapy of ipilimumab plus nivolumab, or chemotherapy as monotherapy. Secondary outcomes were \geq grade 3 AEs according to Common Terminology Criteria for Adverse Events in the last 45 and

90 days before death, the number of treatment lines before last therapy, the number of individual doses given before death, and how frequently reintroduction of a previously used systemic therapy occurred. The number of doses is only registered for ipilimumab, anti-PD-1 antibodies, and combination therapy of ipilimumab plus nivolumab. We estimated the total number of days under treatment with BRAF/MEK inhibitors using the registered start and stop date. Unit costs were used to calculate total costs and total costs per patient. To estimate costs of dabrafenib plus trametinib, a schedule of dabrafenib twice daily plus trametinib once daily was used. For vemurafenib plus cobimetinib, a schedule of vemurafenib twice daily plus cobimetinib once daily was used with a break of seven days for cobimetinib after 21 days. For encorafenib plus binimetinib, a schedule of encorafenib once daily plus binimetinib twice daily was used. Costs of patients with dose reductions of BRAF/MEK inhibitors were adjusted on the basis of the date of the reduction and the dosage given after reduction. The costs represented in this study may not reflect actual costs as confidential financial agreements are made between the melanoma center and the pharmaceutical company and agreements between the pharmaceutical companies and the Ministry of Health.

Statistical Analysis

Baseline patient and disease characteristics of patients during the initiation of their last systemic treatment were analyzed using descriptive statistics. These characteristics included age, sex, lactate dehydrogenase (normal, 250-500 U/L and > 500 U/L), stage (unresectable IIIc, IV-M1a, M1b, and IV-M1c), Eastern Cooperative Oncology Group Performance Score (ECOG PS; 0-1 and ≥ 2), distant metastases (< 3 organ sites and ≥ 3 organ sites involved), liver metastases (yes/no), brain metastases (none, asymptomatic, and symptomatic) and *BRAF*^{V600} mutation status (wild-type or mutant). Baseline patient and disease characteristics of the centers with the lowest and highest percentages of new systemic therapies were compared.

The percentage of patients started with a new systemic therapy within ≤ 45 and ≤ 90 days before death is presented in funnel plots using 95% and 99% confidence limits that vary with the volume of patients per hospital.^{13,14}

Data handling and statistical analyses were performed using the R software system for statistical computing (version 4.0.2.; packages tidyverse,¹⁵ survival,¹⁶ and survminer¹⁷).

RESULTS

Use of Systemic Therapies at the End of Life

A total of 503 (13.2%) patients with advanced melanoma, who died between 2013 and 2019, received a new systemic therapy ≤ 45 days before death (Fig 1). Twenty-nine percent of all patients started a new systemic therapy within the last 90 days before death (N = 1,120). In total, 78




FIG 1. Study flowchart depicting the patients who died between 2013 and 2019 ($N = 3,797$) included for the analyses. Most analyses were on the basis of patients who received new systemic therapy ≤ 45 days ($n = 503$) or 46-90 days ($n = 617$) before death. DMTR, Dutch Melanoma Treatment Registry.

patients started two systemic therapies within 90 days of death. Three patients started three new systemic therapies within 90 days of death. Overall, the percentage of patients treated with a new systemic therapy ≤ 45 and ≤ 90 days before death did not change over the years (Fig 2). Patient and tumor characteristics at the initiation of the last systemic treatment are described in Table 1. Overall, at the start of a new systemic therapy, 30.0% of patients had an ECOG PS ≥ 2 , 95.2% had stage IV-M1c disease, and 72.0% had metastases in ≥ 3 organ sites. The main cause of death was melanoma-related in the majority of the patients (73%-82%), independent of whether they started a systemic therapy within 45 or 90 days before death (Table 1).

Variation Between Melanoma Centers

Between the 14 melanoma centers, the percentage of patients for whom new systemic therapies were initiated ≤ 45 and ≤ 90 days before death varied between 6%-23% and 20%-46%, respectively (Fig 3). The funnel plot shows that one center had a significantly higher percentage and two centers had a significantly lower percentage of patients who started with a new systemic therapy within 45 days of death using the 95% CI compared with the mean. Patient and tumor characteristics of patients treated in the three melanoma centers with the highest and lowest percentage of systemic therapy initiation at the end

of life are shown in the Data Supplement (online only). Overall, no significant differences in patient and tumor characteristics were observed between the centers with the lowest versus the highest percentage of newly initiated systemic therapies. Centers with the lowest percentage had less patients with symptomatic brain metastases, although this difference was not significant.

Type of Systemic Therapy

The type of systemic therapy initiated ≤ 45 and ≤ 90 days before death is shown in Table 2. In 216 of the 503 (42.9%) patients who died within 45 days, systemic therapy was initiated in the first line. First-line systemic treatment consisted mainly of anti-PD-1 antibodies (20.4%), combination therapy of ipilimumab plus nivolumab (17.1%), BRAF-inhibitor monotherapy (24.5%), and combination therapy of BRAF/MEK inhibitors (13.4%). Second- and third-line systemic therapy was initiated in, respectively, 35.1% and 13.7% of the patients. In 41 patients (8.2%), fourth-line or higher was initiated within 45 days of death. BRAF monotherapy and combination therapy of BRAF/MEK inhibitors were predominantly started in patients with brain metastases (54.1%). In total, 24.1% ($n = 121$) and 26.8% ($n = 300$) of patients received radiotherapy within the same episode as their last systemic therapy. Of these 300 patients, 259 (86%) received palliative radiotherapy, 29 patients (10%) received




FIG 2. Percentage of patients who started a new systemic therapy ≤ 45 and ≤ 90 days before death from the total number of patients who died over all years of death.

stereotactic radiation, and the remaining 11 (4%) patients received unspecified forms of radiotherapy.

Adverse Events and Reintroduction

Of all patients dying within 90 days after starting a new systemic therapy, 146 (13.0%) patients developed a total of 185 grade 3/4 AEs in this last period before their death. The most common grade 3/4 AEs were skin toxicities, colitis, hepatitis, and endocrine AEs affecting 18 (12.3%), 32 (21.9%), 18 (12.4%), and 20 (13.7%) patients, respectively (Data Supplement). Nineteen patients developed grade 3-5 AEs while receiving anti-PD-1 inhibitors (1.7%), 46 during BRAF inhibitors (4.1%), 29 during ipilimumab (2.6%), and 35 (3.1%) during ipilimumab plus nivolumab. Reintroduction of a systemic therapy used earlier occurred in 37 patients (12.1%). The reason for discontinuing the reintroduced systemic therapy in the earlier treatment line was progression in 20 patients (32.3%), toxicity in 15 patients (24.2%), planned in 13 patients (21.0%), and for other reasons in 24 patients (38.7%). For patients with an AE, hospital admission was necessary for 102 (69.9%) patients, one patient was admitted to the intensive care unit (0.7%), and one death occurred as a result of grade 5 toxicity (0.7%). In total, 9.1% of patients who died within 90 days of starting a new systemic therapy were admitted to the hospital.

Number of Doses and Costs

In total, 614 ipilimumab doses and 732 anti-PD-1 antibody courses were registered in the DMTR in patients who received a new systemic therapy within 90 days of death. Of the patients who started a new systemic therapy within 45 days of death, 207 ipilimumab courses and 222 anti-

PD-1 antibody courses were registered. On the basis of list prices, the total costs of the courses in the last 45 and 90 days before death are €3,537,934 and €10,688,871, respectively. On the basis of list prices, the total costs of BRAF/MEK inhibitors are €849,623 and €3,699,462 for the courses administered to patients who received a new systemic therapy within 45 and 90 days of death, respectively. Total costs of immune-checkpoint inhibitors and BRAF/MEK inhibitors combined were €4,387,557 and €14,388,333 for patients who received a new systemic therapy within 45 and 90 days of death, respectively. Mean costs per year were €626,793 and €2,055,476 for all patients starting a new systemic therapy within 45 and 90 days of death, respectively. Mean costs per patient were €8,722 and €12,846, respectively.

DISCUSSION

To our knowledge, this is the first study that evaluates the use of systemic therapies in the end of life of patients with advanced melanoma. Our study of a nationwide cohort of patients with advanced melanoma who died between 2013 and 2019 shows that 503 (13.2%) patients with advanced melanoma started a new systemic treatment within 45 days before death, and 1,120 patients (29.5%) had started a new systemic therapy within 90 days of death. Thirteen percent of patients experienced treatment-related severe AEs during these 90 days. However, a large proportion of patients with a grade 3/4 AE required hospital admission (69.9%). Mean total costs per year of the last treatment lines within 45 and 90 days before death were €626,793 and €2,055,476, respectively. We observed a significant difference in the percentage of new systemic therapies

TABLE 1. Characteristics at the Last Episode of the Patients Who Died Within 45 and 46-90 Days of Death and Patients Who Died But Did Not Start a Systemic Therapy Within 90 Days of Death

Baseline Variable	Started ≤ 45 Days (n = 503)	Started 46-90 Days (n = 617)	Died But Not Started ≤ 90 Days (n = 2,677)
Age, years, median (range)	63 (19-93)	63 (21-90)	67 (20-97)
Sex, No. (%)			
Male	330 (65.6)	360 (58.3)	1,553 (58.0)
Female	173 (34.4)	257 (41.7)	1,123 (42.0)
ECOG performance status, No. (%)			
0-1	254 (50.5)	396 (64.2)	1,674 (62.5)
≥ 2	151 (30.0)	138 (22.4)	524 (19.6)
Unknown	98 (19.5)	83 (13.5)	479 (17.9)
Stage (seventh edition AJCC), No. (%)			
Unresectable IIIc	5 (1.0)	9 (1.5)	88 (3.3)
IV-M1a	6 (1.2)	9 (1.5)	74 (2.8)
IV-M1b	8 (1.6)	17 (2.8)	151 (5.6)
IV-M1c	479 (95.2)	578 (93.7)	2,321 (86.7)
Unknown	5 (1.0)	4 (0.6)	43 (1.6)
LDH, No. (%)			
Normal	162 (32.2)	227 (37.8)	1,287 (52.4)
250-500 U/L	152 (30.2)	201 (33.5)	682 (27.8)
> 500 U/L	177 (35.2)	172 (28.7)	485 (19.8)
Unknown/not determined	12 (2.4)	17 (2.8)	223 (8.3)
Brain metastases, No. (%)			
No	227 (45.1)	328 (53.2)	1,604 (59.9)
Yes, asymptomatic	83 (16.5)	105 (17.0)	318 (11.9)
Yes, symptomatic	170 (33.8)	149 (24.1)	655 (24.5)
Unknown	23 (4.6)	35 (5.7)	100 (3.7)
Liver metastases, No. (%)			
No	234 (46.5)	305 (50.7)	1,567 (59.8)
Yes	257 (51.1)	296 (49.3)	1,054 (40.2)
Unknown	12 (2.4)	16 (2.6)	56 (2.1)
Organ sites, No. (%)			
0-2	141 (28.0)	195 (31.6)	1,225 (45.8)
≥ 3	362 (72.0)	422 (68.4)	1,452 (54.2)
<i>BRAF</i> ^{V600} -mutation, No. (%)			
Wild-type	198 (39.4)	263 (42.6)	1,444 (53.9)
Mutant	305 (60.6)	354 (57.4)	1,233 (46.1)

(continued on following page)

started between some of the 14 melanoma centers in the Netherlands.

We observed that the type of systemic therapy initiated ≤ 45 days before death varied between melanoma centers. All types of systemic therapies were started (BRAF monotherapy, BRAF/MEK inhibitors, ipilimumab, anti-PD-1, and combination therapy of ipilimumab plus nivolumab). The majority of systemic therapies were started as first-line treatment (42.9%). Overall, of all patients who died within 90 days of death of starting a systemic therapy, 30% started with BRAF monotherapy or combination therapy of BRAF/MEK inhibitors, which is a relative over-representation as only 60.6% of patients harbor a *BRAF*-mutation and are eligible for BRAF/MEK inhibitors. These therapies can provide patients with major symptom relief or palliation in the last phase of life.

Only a small proportion of the study population that died within 90 days of starting a new systemic therapy developed grade 3/4 AEs (13.0%). This percentage is lower than the randomized clinical trials, where the percentage of patients experiencing grade 3/4 AEs ranged from 20% to 59%.¹⁻⁶ The real-world patients included in this study were older, with a higher ECOG PS, and with more brain metastases compared with these randomized clinical trials. However, in the Checkmate-67, the median time until onset of these AEs ranged from 3.7 to 12.2 months.⁶ The selected patients in this study received a maximum of 1.5 months of systemic therapy. A possible explanation for the low percentage of AEs is that the included patients could not develop toxicity in the short time they received systemic therapy. Previous research has shown that the percentage of real-world patients experiencing grade 3/4 AEs in the DMTR is similar to randomized clinical trials,¹⁸ and more advanced disease is associated with a lower risk of grade 3/4 AEs.¹⁹

Although starting systemic therapy in the last phase of life can be meaningful, it is important to have insight into its financial impact. It is estimated that the mean total costs of Dutch patients with advanced melanoma measured from diagnosis are €89,240.²⁰ The current study estimates that mean total costs per cohort year in 2013-2019 for patients who died within 90 days of starting a new systemic therapy were €2,055,476. A previous study has shown that the total budget impact per cohort year of patients with advanced melanoma ranged between 73 and 90 million Euros.²¹ This budget impact may increase with emerging treatments for melanoma such as LAG-3,²² tumor-infiltrating lymphocytes,²³ and tebentafusp.²⁴ The costs of new systemic therapies in the last 45/90 days before death are a relatively small proportion (2.3%-2.8%). Patients starting a new systemic therapy only receive a small amount of courses, which could explain the small proportion of total costs. Starting a new systemic therapy should still be considered with caution in patients with poor condition who have doubts about starting a new systemic therapy. However, costs could be limited when patients die early after the onset of treatment. Unfortunately, no model

TABLE 1. Characteristics at the Last Episode of the Patients Who Died Within 45 and 46-90 Days of Death and Patients Who Died But Did Not Start a Systemic Therapy Within 90 Days of Death (continued)

Baseline Variable	Started ≤ 45 Days (n = 503)	Started 46-90 Days (n = 617)	Died But Not Started ≤ 90 Days (n = 2,677)
Cause of death, No. (%)			
Melanoma-related	414 (82.3)	498 (80.7)	1,952 (72.)
Toxicity because of treatment	7 (1.4)	8 (1.3)	6 (0.2)
Comorbidity	9 (1.8)	4 (0.6)	31 (1.2)
Other	17 (3.4)	19 (3.1)	65 (2.4)
Unknown	56 (11.1)	88 (14.2)	575 (21.5)

NOTE. In the Dutch Melanoma Treatment Registry, an episode is created when patients start a new systemic therapy.

Abbreviations: AJCC, American Joint Committee on Cancer; ECOG, Eastern Cooperative Oncology Group; LDH, lactate dehydrogenase.

exists to predict individual response to immunotherapy or targeted therapies.

During the total study period, differences exist between melanoma centers in the percentage of patients receiving a new systemic therapy in the Netherlands. This information

is important for both clinicians and patients. Clinicians might be unaware of the percentage of patients they are treating in the last phase of life with a new systemic therapy and how this percentage compares to other centers. By providing this information on differences between centers, the variation in the percentage of patients who are treated with systemic therapy in the last phase of life might become smaller. The results of this study have been discussed with the Dutch melanoma centers. To better understand the rationale of starting a new systemic therapies, comparisons between individual centers should take place. The current study did not measure quality of life in patients who received new systemic therapies ≤ 45 days before death. It would be interesting to investigate the quality of life in these patients, especially those who receive systemic therapy for symptom relief. In the current population, severe AEs in the last phase of life seem rare. If the patient understands that treatment may have a positive effect on the course of disease, and agrees with the risk of potentially severe AEs, there is no reason to not start a new systemic therapy.

This study has several limitations. First, we had to estimate the costs of BRAF/MEK inhibitors on the basis of the start and stop data. We argue that the estimated total and mean costs are a near-approximation of the true costs of the studied population. Second, this study only investigated Dutch patients with advanced melanoma. Although




FIG 3. Variation between melanoma centers in the percentage of patients who started a new systemic therapy ≤ 45 (left) and ≤ 90 (right) days before death among all deceased patients. The dotted line represents the average percentage of patients receiving a new systemic therapy ≤ 45 and ≤ 90 days before death. Hospitals falling outside the control limits deviate significantly from the national average. CL, control limit.

TABLE 2. Type of Systemic Therapy Started \leq 45 (top) and \leq 90 Days (bottom) Before Date of Death, Stratified for Line of Treatment

Line of Systemic Therapy	First-Line	Second-Line	Third-Line	\geq Fourth-Line	Total
\leq 45 days					
Total No. of patients, %	216 (42.9)	177 (35.2)	69 (13.7)	41 (8.2)	503 (100.0)
Anti-PD-1 antibodies	44 (20.4)	31 (17.5)	6 (8.7)	6 (14.6)	87 (17.3)
BRAF monotherapy	53 (24.5)	15 (8.5)	13 (18.8)	5 (12.2)	86 (17.1)
BRAF/MEK inhibitors	29 (13.4)	17 (9.6)	19 (27.5)	14 (34.1)	80 (15.7)
Chemotherapy	23 (10.6)	8 (4.5)	4 (5.8)	2 (4.9)	37 (7.4)
Ipilimumab	22 (10.2)	43 (24.3)	10 (14.5)	1 (2.4)	76 (15.1)
Ipilimumab plus nivolumab	37 (17.1)	43 (24.3)	7 (10.1)	4 (9.8)	94 (18.1)
Other systemic therapy	7 (3.2)	19 (10.7)	10 (14.5)	9 (22.0)	45 (8.9)
Unknown	1 (0.5)	1 (0.6)	0 (0.0)	0 (0.0)	2 (0.4)
\leq 90 days					
Total No. of patients, %	514 (45.9)	350 (31.3)	166 (14.8)	90 (8.0)	1,120 (100.0)
Anti-PD-1 antibodies	117 (22.8)	71 (20.3)	24 (14.5)	16 (17.8)	228 (20.4)
BRAF monotherapy	122 (23.7)	31 (8.9)	23 (13.9)	10 (11.1)	186 (16.6)
BRAF/MEK inhibitors	60 (11.7)	29 (8.3)	47 (28.3)	31 (34.4)	167 (14.9)
Chemotherapy	51 (9.9)	19 (5.4)	12 (7.2)	5 (5.6)	87 (7.8)
Ipilimumab	67 (13.0)	93 (26.6)	20 (12.0)	3 (3.3)	183 (16.3)
Ipilimumab plus nivolumab	71 (13.8)	72 (20.6)	14 (8.4)	7 (7.8)	164 (14.6)
Other systemic therapy	23 (4.5)	33 (9.4)	26 (15.7)	17 (18.9)	99 (8.8)
Unknown	1 (0.2)	1 (0.3)	0 (0.0)	0 (0.0)	2 (0.2)

NOTE. Only the most recent therapies are shown in this table. Patients who started with BRAF monotherapy who received BRAF/MEK combination therapy in the previous line of treatment were not seen as a start of new therapy.

Abbreviation: PD-1, programmed cell death protein-1.

treatment options for advanced melanoma are comparable across Western countries, it is unknown how our results compare with other countries. Third, we included patients who died between 2013 and 2019. Many of these patients who died between 2013 and 2014 received BRAF monotherapy or ipilimumab monotherapy as a last systemic therapy. This does not represent the current treatment landscape. Fourth, we could not show the characteristics of patients who stopped or did not start systemic therapy $>$ 45 days before death. Only characteristics at the start of systemic therapy are registered in the DMTR. Fifth, only costs of the systemic therapies itself are included in this study. Costs of infusion and other care are not included. Finally, this study does not provide insights on which patients are unlikely to benefit. A previous study of patients with advanced melanoma has

shown that patients with elevated lactate dehydrogenase, brain metastases, and a poor ECOG PS had lower OS.⁸ Future studies could focus on developing a prediction model for treatment outcomes when starting a systemic therapy in different phases of the disease.

In conclusion, this nationwide study shows the percentage of newly initiated systemic therapies within 45 and 90 days before death. Only a small percentage of these patients experience grade 3/4 AEs, but a large proportion of patients with a grade 3/4 AE had to be admitted to the hospital. The costs of new systemic therapies in the last 45/90 days before death are a relatively small proportion of the total costs of patients with advanced melanoma (2.3%-2.8%). Future studies should be aimed at identifying prognostic factors that predict a short-term poor outcome.

AFFILIATIONS

¹Dutch Institute for Clinical Auditing, Leiden, the Netherlands

²Department of Medical Oncology, Amsterdam UMC, VU University Medical Center, Cancer Center Amsterdam, Amsterdam, the Netherlands

³Division of Pharmacoepidemiology and Clinical Pharmacology, Utrecht Institute for Pharmaceutical Sciences, Utrecht, the Netherlands

⁴Department of Surgical Oncology, Netherlands Cancer Institute, Amsterdam, the Netherlands

⁵Department of Biomedical Data Sciences, Leiden University Medical Center, Leiden, the Netherlands

⁶Department of Pharmacy, Rode Kruis Ziekenhuis, Beverwijk, the Netherlands

⁷Department of Medical Oncology and Immunology, Netherlands Cancer Institute, Amsterdam, the Netherlands

⁸Division of Molecular Oncology and Immunology, Netherlands Cancer Institute, Amsterdam, the Netherlands

⁹Department of Medical Oncology, Leiden University Medical Center, Leiden, the Netherlands

¹⁰Department of Medical Oncology, GROW School for Oncology and Developmental Biology, Maastricht University Medical Center, Maastricht, the Netherlands

¹¹Department of Medical Oncology, Zuyderland Medical Center Sittard, Sittard-Geleen, the Netherlands

¹²Isala Oncology Center, Isala, Zwolle, the Netherlands

¹³Department of Medical Oncology, University Medical Center Groningen, University of Groningen, Groningen, the Netherlands

¹⁴Department of Internal Medicine, Medisch Spectrum Twente, Enschede, the Netherlands

¹⁵Department of Internal Medicine, Medical Center Leeuwarden, Leeuwarden, the Netherlands

¹⁶Department of Internal Medicine, Amphia Hospital, Breda, the Netherlands

¹⁷Department of Medical Oncology, Erasmus Medical Center, Rotterdam, the Netherlands

¹⁸Department of Internal Medicine, Maxima Medical Center, Eindhoven, the Netherlands

¹⁹Department of Medical Oncology, Radboud University Medical Center, Nijmegen, the Netherlands

²⁰Department of Medical Oncology, University Medical Center Utrecht, Utrecht, the Netherlands

CORRESPONDING AUTHOR

Jesper van Breeschoten, PharmD, Rijnsburgerweg 10, 2333AA Leiden, the Netherlands; e-mail: j.vanbreeschoten@dica.nl.

DISCLAIMER

This is a US Government work. There are no restrictions on its use. The funders had no role in the writing of this article or decision to submit it for publication.

SUPPORT

For the Dutch Melanoma Treatment Registry (DMTR), the Dutch Institute for Clinical Auditing foundation received a start-up grant from governmental organization The Netherlands Organization for Health

REFERENCES

- Weber JS, D'Angelo SP, Minor D, et al: Nivolumab versus chemotherapy in patients with advanced melanoma who progressed after anti-CTLA-4 treatment (CheckMate 037): A randomised, controlled, open-label, phase 3 trial. *Lancet Oncol* 16:375-384, 2015
- Robert C, Schachter J, Long GV, et al: Pembrolizumab versus ipilimumab in advanced melanoma. *N Engl J Med* 372:2521-2532, 2015
- Ascierto PA, McArthur GA, Dréno B, et al: Cobimetinib combined with vemurafenib in advanced BRAFV600-mutant melanoma (coBRIM): Updated efficacy results from a randomised, double-blind, phase 3 trial. *Lancet Oncol* 17:1248-1260, 2016
- Robert C, Karaszewska B, Schachter J, et al: Improved overall survival in melanoma with combined dabrafenib and trametinib. *N Engl J Med* 372:30-39, 2015
- Hodi FS, O'Day SJ, McDermott DF, et al: Improved survival with ipilimumab in patients with metastatic melanoma. *N Engl J Med* 363:711-723, 2010
- Larkin J, Chiarion-Sileni V, Gonzalez R, et al: Five-year survival with combined nivolumab and ipilimumab in advanced melanoma. *N Engl J Med* 381:1535-1546, 2019
- O'Reilly A, Hughes P, Mann J, et al: An immunotherapy survivor population: Health-related quality of life and toxicity in patients with metastatic melanoma treated with immune checkpoint inhibitors. *Support Care Cancer* 28:561-570, 2020
- van Zeijl MCT, Ismail RK, de Wreede LC, et al: Real-world outcomes of advanced melanoma patients not represented in phase III trials. *Int J Cancer* 147:3461-3470, 2020
- Fox JA, Langbecker D, Rosenberg J, et al: Uncertain diagnosis and prognosis in advanced melanoma: A qualitative study of the experiences of bereaved carers in a time of immune and targeted therapies. *Br J Dermatol* 180:1368-1376, 2019
- Temel JS, Shaw AT, Greer JA: Challenge of prognostic uncertainty in the modern era of cancer therapeutics. *J Clin Oncol* 34:3605-3608, 2016
- Khaki AR, Chennupati S, Fedorenko C, et al: Utilization of systemic therapy in patients with cancer near the end of life in the pre- versus postimmune checkpoint inhibitor eras. *JCO Oncol Pract* 17:e1728-e1737, 2021
- Jochems A, Schouwenburg MG, Leeneman B, et al: Dutch Melanoma Treatment Registry: Quality assurance in the care of patients with metastatic melanoma in the Netherlands. *Eur J Cancer* 72:156-165, 2017
- Rakow T, Wright RJ, Spiegelhalter DJ, et al: The pros and cons of funnel plots as an aid to risk communication and patient decision making. *Br J Psychol* 106:327-348, 2015
- Spiegelhalter DJ: Funnel plots for comparing institutional performance. *Stat Med* 24:1185-1202, 2005

Research and Development (ZonMW, Grant No. 836002002). The DMTR is structurally funded by Bristol Myers Squibb, Merck Sharpe & Dohme, Novartis, and Roche Pharma. Roche Pharma stopped and Pierre Fabre started funding of the DMTR in 2019.

AUTHORS' DISCLOSURES OF POTENTIAL CONFLICTS OF INTEREST

Disclosures provided by the authors are available with this article at DOI <https://doi.org/10.1200/OP.22.00061>.

AUTHOR CONTRIBUTIONS

Conception and design: Jesper van Breeschoten, Rawa K. Ismail, Michel W.J.M. Wouters, Doranne L. Hilarius, Karijn P.M. Suijkerbuijk, Alfons J.M. van den Eertwegh

Administrative support: Doranne L. Hilarius, Rawa K. Ismail, Michel W.J.M. Wouters, Doranne L. Hilarius, Alfons J.M. van den Eertwegh

Provision of study materials or patients: Michel W.J.M. Wouters, John B. Haanen, Christian U. Blank, Franchette W.P.J. van den Berkmortel, Jan Willem B. de Groot, Geke A.P. Hospers, Ellen Kapiteijn, Djura Piersma, Rozemarijn S. van Rijn, Marion A. Stevensen-den Boer, Astrid A.M. van der Veldt, Gerard Vreugdenhil, Marye J. Boers-Sonderen, Karijn P.M. Suijkerbuijk, Alfons J.M. van den Eertwegh

Collection and assembly of data: Jesper van Breeschoten, Michel W.J.M. Wouters, John B. Haanen, Christian U. Blank, Franchette W.P.J. van den Berkmortel, Jan Willem B. de Groot, Geke A.P. Hospers, Ellen Kapiteijn, Djura Piersma, Rozemarijn S. van Rijn, Marion A. Stevensen-den Boer, Astrid A.M. van der Veldt, Gerard Vreugdenhil, Marye J. Boers-Sonderen, Karijn P.M. Suijkerbuijk, Alfons J.M. van den Eertwegh

Data analysis and interpretation: All authors

Manuscript writing: All authors

Final approval of manuscript: All authors

Accountable for all aspects of the work: All authors

ACKNOWLEDGMENT

The authors thank all physicians and data managers who registered the patient data in the Dutch Melanoma Treatment Registry.

15. Wickham H, Averick M, Bryan J, et al: Welcome to the {tidyverse}. *J Open Source Softw* 4:1686, 2019
16. Thernau TM: A Package for Survival Analysis in R. 2020. <https://CRAN.R-project.org/package=survival>
17. Kassambara A, Kosinski M, Biecek P: Drawing Survival Curves Using "ggplot2". 2020. <https://CRAN.R-project.org/package=survminer>
18. Van Zeijl MCT, Haanen JBAG, Wouters MWJM, et al: Real-world outcomes of first-line anti-PD-1 therapy for advanced melanoma: A nationwide population-based study. *J Immunother* 43:256-264, 2020
19. Verheijden RJ, May AM, Blank CU, et al: Lower risk of severe checkpoint inhibitor toxicity in more advanced disease. *ESMO Open* 5:e000945, 2020
20. Leeneman B, Uyl-De Groot CA, Aarts MJB, et al: Healthcare costs of metastatic cutaneous melanoma in the era of immunotherapeutic and targeted drugs. *Cancers (Basel)* 12:1-11, 2020
21. Franken MG, Leeneman B, Aarts MJB, et al: Trends in survival and costs in metastatic melanoma in the era of novel targeted and immunotherapeutic drugs. *ESMO Open* 6:100320, 2021
22. Lipson EJ, Tawbi HA, Schadendorf D, et al: Relatlimab (RELA) plus nivolumab (NIVO) versus NIVO in first-line advanced melanoma: Primary phase III results from RELATIVITY-047 (CA224-047). *J Clin Oncol* 39, 2021 (15 suppl; abstr 9503)
23. Van Den Berg JH, Heemskerck B, Van Rooij N, et al: Tumor infiltrating lymphocytes (TIL) therapy in metastatic melanoma: Boosting of neoantigen-specific T cell reactivity and long-term follow-up. *J Immunother Cancer* 8:1-11, 2020
24. Nathan P, Hassel JC, Rutkowski P, et al: Overall survival benefit with tebentafusp in metastatic uveal melanoma. *N Engl J Med* 385:1196-1206, 2021



AUTHORS' DISCLOSURES OF POTENTIAL CONFLICTS OF INTEREST**End-of-Life Use of Systemic Therapy in Patients With Advanced Melanoma: A Nationwide Cohort Study**

The following represents disclosure information provided by authors of this manuscript. All relationships are considered compensated unless otherwise noted. Relationships are self-held unless noted. I = Immediate Family Member, Inst = My Institution. Relationships may not relate to the subject matter of this manuscript. For more information about ASCO's conflict of interest policy, please refer to www.asco.org/rwc or ascopubs.org/op/authors/author-center.

Open Payments is a public database containing information reported by companies about payments made to US-licensed physicians (Open Payments).

Rawa K. Ismail

Employment: MSD

John B. Haanen

Stock and Other Ownership Interests: Neogene Therapeutics

Consulting or Advisory Role: MSD Oncology (Inst), Pfizer (Inst), Bristol Myers Squibb (Inst), Novartis (Inst), Roche/Genentech (Inst), Ipsen (Inst), Achilles Therapeutics (Inst), Immunocore (Inst), Sanofi (Inst), Third Rock Ventures (Inst), Neogene Therapeutics, Molecular Partners (Inst), bioNTech (Inst), T-Knife (Inst), PokeAcell (Inst), Instil Bio (Inst), Iovance Biotherapeutics (Inst)

Research Funding: MSD (Inst), Bristol Myers Squibb (Inst), Novartis (Inst), Neon Therapeutics (Inst), Amgen (Inst), BioNTech (Inst), Asher Biotherapeutics (Inst)

Christian U. Blank

Stock and Other Ownership Interests: Immagene

Consulting or Advisory Role: Roche/Genentech (Inst), MSD Oncology (Inst), Bristol Myers Squibb (Inst), Novartis (Inst), GlaxoSmithKline (Inst), Pfizer (Inst), AstraZeneca (Inst), Lilly (Inst), Pierre Fabre (Inst), GenMab (Inst), Third Rock Ventures

Research Funding: Bristol Myers Squibb (Inst), Novartis (Inst), NanoString Technologies (Inst), 4SC (Inst)

Patents, Royalties, Other Intellectual Property: WO 2021/177822 A1

Expert Testimony: Freshfields

Travel, Accommodations, Expenses: Bristol Myers Squibb

Maureen J.B. Aarts

Research Funding: Pfizer (Inst)

Jan Willem B. de Groot

Consulting or Advisory Role: Bristol Myers Squibb, Pierre Fabre, Servier

Geke A.P. Hospers

Consulting or Advisory Role: Roche (Inst), MSD (Inst), Amgen (Inst), Bristol Myers Squibb (Inst), Novartis (Inst), Pierre Fabre (Inst), Sanofi (Inst)

Research Funding: Bristol Myers Squibb (Inst), Seerave Foundation (Inst)

Ellen Kapiteijn

Consulting or Advisory Role: Pierre Fabre (Inst), Bristol Myers Squibb (Inst), Novartis (Inst), Lilly (Inst), Ipsen (Inst), Delcath Systems (Inst)

Research Funding: BMS (Inst), Pierre Fabre (Inst)

Djura Piersma

Consulting or Advisory Role: Pierre Fabre (Inst), Novartis (Inst)

Astrid A.M. van der Veldt

Consulting or Advisory Role: Ipsen (Inst), Roche (Inst), Bristol Myers Squibb (Inst), Pfizer (Inst), MSD Oncology (Inst), Sanofi (Inst), Pierre Fabre (Inst), Novartis (Inst), Eisai (Inst), Merck (Inst)

Research Funding: Bayer (Inst)

Travel, Accommodations, Expenses: Roche, MSD Oncology, Novartis, Bayer

Karijn P.M. Suijkerbuijk

Consulting or Advisory Role: Bristol Myers Squibb (Inst), Novartis (Inst), MSD (Inst), Pierre Fabre (Inst), AbbVie (Inst)

Speakers' Bureau: Roche (Inst), Novartis (Inst)

Research Funding: Bristol Myers Squibb Foundation (Inst), TigaTx (Inst), Philips Research (Inst)

Alfonso J.M. van den Eertwegh

Honoraria: Bristol Myers Squibb

Consulting or Advisory Role: Bristol Myers Squibb, MSD Oncology, Amgen, Roche, Novartis, Sanofi, Pfizer, Ipsen, Merck, Pierre Fabre

Research Funding: Roche, Sanofi, BMS

No other potential conflicts of interest were reported.