

Left ventricle unloading in extracorporeal life support

Citation for published version (APA):

Meani, P. (2022). *Left ventricle unloading in extracorporeal life support*. [Doctoral Thesis, Maastricht University]. Maastricht University. <https://doi.org/10.26481/dis.20221222pm>

Document status and date:

Published: 01/01/2022

DOI:

[10.26481/dis.20221222pm](https://doi.org/10.26481/dis.20221222pm)

Document Version:

Publisher's PDF, also known as Version of record

Please check the document version of this publication:

- A submitted manuscript is the version of the article upon submission and before peer-review. There can be important differences between the submitted version and the official published version of record. People interested in the research are advised to contact the author for the final version of the publication, or visit the DOI to the publisher's website.
- The final author version and the galley proof are versions of the publication after peer review.
- The final published version features the final layout of the paper including the volume, issue and page numbers.

[Link to publication](#)

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal.

If the publication is distributed under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license above, please follow below link for the End User Agreement:

www.umlib.nl/taverne-license

Take down policy

If you believe that this document breaches copyright please contact us at:

repository@maastrichtuniversity.nl

providing details and we will investigate your claim.

Chapter 10. Impact

Abbreviations

CA, cardiac arrest

CS, Cardiogenic shock

IABP, Intra-aortic Balloon Pump

LV, Left Ventricle

V-A ECLS, veno-arterial extracorporeal life support

Cardiogenic shock (CS) and cardiac arrest (CA) are among the most lethal manifestation of acute cardiovascular disease, with high in-hospital mortality rates.

CA in North American and Europe approximates 50 to 100 cases per 100,000 [1] and the 30-day survival rate of out-of-hospital cardiac arrest patients who received cardiopulmonary resuscitation is only 10.7% worldwide [2]. Likewise, intensive care unit admission with CS has doubled from 4% to 8% over the past 15 years [3] and it is associated with a high mortality rate ranging from 30% to 40% [4-7].

Extracorporeal life support is increasingly used either in adults or children with acutely impaired cardiac function refractory to conventional medical management, mainly in profound cardiogenic shock and refractory cardiac arrest. In the United States, veno-arterial extracorporeal life support (V-A ECLS) use has raised from 1.613 per million in the year 2001 to 3.597 per million in 2011[8], and from 5.4 per million in the year 2000 to 44.3 per million in 2009 [9]. Despite this therapeutic advancement, mortality from CS and CA remain high [10-12]. Therefore, the efforts in improving the ECLS shortcomings have been increasing over the past decades. One of the most important concerns in V-A ECLS is the increased left ventricular (LV) pressure attributable to retrograde aortic perfusion. This could slow myocardial recovery or damage the myocardium and negatively affect survival [13]. This thesis provides a comprehensive treatise of LV unloading during V-A ECLS support, moving from a picture of the available strategies and the need of standardize the indications and treatments, through solid physiological data based on pre-clinical studies and ending to a clinical experience in a unique setting.

Scientific impact

Our work provided two main scientific improvements. First, our findings were based on a meticulous scientific methodology which firstly recognized the urgent need of experimental data supporting the use of these techniques in the daily clinical practice. This led to provide a detailed insight of the LV hemodynamics and workload investigated in a swine CS model, as shown in **chapter 4** and **chapter 6**. Additionally, a comprehensive scientific treatise on the Impella use was reported in **chapter 5**.

Second, this thesis was certainly one of the first attempt to align all the available indications and approaches. In **chapter 2**, we firstly recognized the need of a common definition of LV overload. In addition, a multiparametric approach was proposed and, nowadays, it has been one of the few clear indications available in the literature.

Clinical impact

Chapter 2 proposed, for the first time, a detailed literature review on this topic. We analyzed in depth the available literature and systematically treated each technique with its advantages and disadvantages. Furthermore, our conclusions strongly remarked a lack of knowledge. On one hand,

approximately 60% of the selected papers evaluated in this study accounted for case reports or case series. On the other hand, grounded data regarding the hemodynamic and physiological changes related to each method were absent. The scientific interest on LV unloading has been sharply raised over the following five years.

One of the controversies, mentioned in our review, was the role of intra-aortic balloon pump (IABP) as an effective unloading technique. In **chapter 3**, despite the limited number of patients, IABP implant will restore aortic valve opening in patients supported with V-A ECLS. Although the aortic valve dysfunction is considered only one of the LV overloading signs, the positive IABP impact confirmed its effectiveness in at least reversing this adverse event. This was a promising insight which drove the role IABP as a tool mainly when early signs of LV overload occurred. In **chapter 4** and **chapter 6** were the first studies which directly compared the effects of 3 different modalities of LV unloading during peripheral V-A ECLS. In this large animal model with profound cardiogenic shock (CS) supported with V-A ECLS, the Impella, pulmonary artery cannula and Atrial septostomy provided effective LV unloading maintaining adequate end-organ perfusion. Nevertheless, the grade of unloading significantly differed as demonstrated in our CS animal model. Besides the unique experimental data and the consequence better understanding of this phenomenon, these studies firstly provided solid proofs which may guide clinicians in their decision-making process. First, Impella was the most effective unloading technique in our experimental studies. Then, its effectiveness was further confirmed in large multicenter experiences and its use became widespread [14,15]. Second, although the role of pulmonary artery cannula as a dynamic and biventricular support was established [16], little was known regarding the effective impact on LV. Our experiment clarified and enhanced its effect as indirect LV unloading. Third, the use of atrial septostomy was supported by solid preclinical data which confirmed its effectiveness. With specific reference to the latter technique, **chapter 7** confirmed its benefits in a unique population of congenital heart disease patients supported V-A ECLS. To best authors' knowledge, this was the first and largest experience which demonstrated the impact of LV unloading on in-hospital survival in such population. Finally, **chapter 8** advanced a comprehensive algorithm as a diagnostic and therapeutic tool for properly unloading the LV during V-A ECLS. To summarize, this thesis investigated in depth the mechanisms and the hemodynamic impact of the most promising LV techniques. At the same time, we provided important insight which may support the clinicians in the decision-making process and patient selection. Finally, we opened a new research perspective regarding patients who might benefit LV unloading the most.

References

1. Wong, C.X.; Brown, A.; Lau, D.H.; Chugh, S.S.; Albert, C.M.; Kalman, J.M.; Sanders, P. Epidemiology of Sudden Cardiac Death: Global and Regional Perspectives. *Heart Lung Circ.* 2019, 28, 6–14.
2. Yan, S.; Gan, Y.; Jiang, N.; Wang, R.; Chen, Y.; Luo, Z.; Zong, Q.; Chen, S.; Lv, C. The global survival rate among adult out-of-hospital cardiac arrest patients who received cardiopulmonary resuscitation: A systematic review and meta-analysis. *Crit Care* 2020, 24, 61.
3. Puymirat E, Fagon JY, Aegerter P, et al. Cardiogenic shock in intensive care units: evolution of prevalence, patient profile, management and outcomes, 1997-2012. *Eur J Heart Fail* 2017;19:192–200.
4. Berg DD, Bohula EA, van Diepen S, et al. Epidemiology of shock in contemporary cardiac intensive care units. *Circ Cardiovasc Qual Outcomes.* 2019;12:e005618.
5. Hochman JS. Cardiogenic shock complicating acute myocardial infarction: expanding the paradigm. *Circulation.* 2003;107:2998–3002.
6. Hochman JS, Sleeper LA, Webb JG, et al. Early revascularization in acute myocardial infarction complicated by cardiogenic shock. SHOCK Investigators. Should we emergently revascularize occluded coronaries for cardiogenic shock. *N Engl J Med.* 1999;341:625–634.
7. Lawler PR, Bohula EA, Park J-G, Katz J, Diepen SV, Morrow DA. Clinical staging in cardiogenic shock discriminates survival: data from the
8. Yarmohammadi H, Duval S, Jain R. Trends and outcomes in the use of extracorporeal membrane oxygenation for adults in the United States. *Proc Am Heart Assoc Annu Meeting* 2013;128:A18565.
9. Nasr DM, Rabistein AA. Neurologic complications of extracorporeal membrane oxygenation. *J Clin Neurol* 2015;11: 383–9.
10. Ponikowski P, Voors AA, Anker SD, Bueno H, Cleland JGF, Coats AJS, Falk V, González-Juanatey JR, Harjola VP, Jankowska EA, et al; ESC Scientific Document Group. 2016 ESC guidelines for the diagnosis and treatment of acute and chronic heart failure: the task force for the diagnosis and treatment of acute and chronic heart failure of the European Society of Cardiology (ESC): developed with the special contribution of the Heart Failure Association (HFA) of the ESC. *Eur Heart J.* 2016;37:2129–2200.
11. Becher PM, Schrage B, Sinning CR, Schmack B, Fluschnik N, Schwarzl M, Waldeyer C, Lindner D, Seiffert M, Neumann JT, et al. Venoarterial extracorporeal membrane oxygenation for cardiopulmonary support. *Circulation.* 2018;138:2298–2300.
12. Chen, Y.S.; Lin, J.W.; Yu, H.Y.; Ko, W.J.; Jerng, J.S.; Chang, W.T.; Chen, W.J.; Huang, S.C.; Chi, N.H.; Wang, C.H.; et al. Cardiopulmonary resuscitation with assisted extracorporeal life-support versus conventional cardiopulmonary resuscitation in adults with in-hospital cardiac arrest: An observational study and propensity analysis. *Lancet* 2008, 372, 554–561
13. Ostadal P, Mlcek M, Kruger A, Hala P, Lacko S, Mates M, et al. Increasing venoarterial extracorporeal membrane oxygenation flow negatively affects left ventricular performance in a porcine model of cardiogenic shock. *Journal of translational medicine.* 2015;13:266.
14. Schrage B, Burkhoff D, RübSamen N, Becher PM, Schwarzl M, Bernhardt A, Grahn H, Lubos

Chapter 10. Impact

- E, Söffker G, Clemmensen P, Reichenspurner H, Blankenberg S, Westermann D. Unloading of the Left Ventricle During Venoarterial Extracorporeal Membrane Oxygenation Therapy in Cardiogenic Shock. *JACC Heart Fail.* 2018 Dec;6(12):1035-1043.
15. Schrage B, Becher PM, Bernhardt A, Bezerra H, Blankenberg S, Brunner S, Colson P, Cudemus Deseda G, Dabboura S, Eckner D, Eden M, Eitel I, Frank D, Frey N, Funamoto M, Goßling A, Graf T, Hagl C, Kirchhof P, Kupka D, Landmesser U, Lipinski J, Lopes M, Majunke N, Maniuc O, McGrath D, Möbius-Winkler S, Morrow DA, Mourad M, Noel C, Nordbeck P, Orban M, Pappalardo F, Patel SM, Pauschinger M, Pazzanese V, Reichenspurner H, Sandri M, Schulze PC, H G Schwinger R, Sinning JM, Aksoy A, Skurk C, Szczanowicz L, Thiele H, Tietz F, Varshney A, Wechsler L, Westermann D. Left Ventricular Unloading Is Associated With Lower Mortality in Patients With Cardiogenic Shock Treated With Venoarterial Extracorporeal Membrane Oxygenation: Results From an International, Multicenter Cohort Study. *Circulation.* 2020 Dec;142(22):2095-2106.
 16. Lorusso R, Raffa GM, Heuts S, Lo Coco V, Meani P, Natour E, Bidar E, Delnoij T, Loforte A. Pulmonary artery cannulation to enhance extracorporeal membrane oxygenation management in acute cardiac failure. *Interact Cardiovasc Thorac Surg.* 2020 Feb 1;30(2):215-222.