

# Fructose Intake From Fruit Juice and Sugar-Sweetened Beverages Is Associated With Higher Intrahepatic Lipid Content

Citation for published version (APA):

Buziau, A. M., Eussen, S. J. P. M., Kooi, M. E., van der Kallen, C. J. H., van Dongen, M. C. J. M., Schaper, N. C., Henry, R. M. A., Schram, M. T., Dagnelie, P. C., van Greevenbroek, M. M. J., Wesselius, A., Bekers, O., Meex, S. J. R., Schalkwijk, C. G., Stehouwer, C. D. A., & Brouwers, M. C. G. J. (2022). Fructose Intake From Fruit Juice and Sugar-Sweetened Beverages Is Associated With Higher Intrahepatic Lipid Content: The Maastricht Study. *Diabetes Care*, *45*(5), 1116-1123. https://doi.org/10.2337/dc21-2123

**Document status and date:** Published: 01/05/2022

DOI: 10.2337/dc21-2123

**Document Version:** Publisher's PDF, also known as Version of record

**Document license:** Taverne

### Please check the document version of this publication:

• A submitted manuscript is the version of the article upon submission and before peer-review. There can be important differences between the submitted version and the official published version of record. People interested in the research are advised to contact the author for the final version of the publication, or visit the DOI to the publisher's website.

• The final author version and the galley proof are versions of the publication after peer review.

• The final published version features the final layout of the paper including the volume, issue and page numbers.

Link to publication

#### General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

• Users may download and print one copy of any publication from the public portal for the purpose of private study or research.

• You may not further distribute the material or use it for any profit-making activity or commercial gain

• You may freely distribute the URL identifying the publication in the public portal.

If the publication is distributed under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license above, please follow below link for the End User Agreement:

www.umlib.nl/taverne-license

### Take down policy

If you believe that this document breaches copyright please contact us at:

repository@maastrichtuniversity.nl

providing details and we will investigate your claim.



Fructose Intake From Fruit Juice and Sugar-Sweetened Beverages Is Associated With Higher Intrahepatic Lipid Content: The Maastricht Study

Diabetes Care 2022;45:1116-1123 | https://doi.org/10.2337/dc21-2123



Amée M. Buziau,<sup>1,2,3</sup> Simone J.P.M. Eussen,<sup>2,4,5</sup> M. Eline Kooi,<sup>2,6</sup> Carla J.H. van der Kallen,<sup>2,7</sup> Martien C.J.M. van Dongen,<sup>4,5</sup> Nicolaas C. Schaper,<sup>1,2,5</sup> Ronald M.A. Henry,<sup>2,7,8</sup> Miranda T. Schram,<sup>2,7,8</sup> Pieter C. Dagnelie,<sup>2,7</sup> Marleen M.J. van Greevenbroek,<sup>2,3</sup> Anke Wesselius,<sup>9,10</sup> Otto Bekers,<sup>2,11</sup> Steven J.R. Meex,<sup>2,11</sup> Casper G. Schalkwijk,<sup>2,3</sup> Coen D.A. Stehouwer,<sup>2,7</sup> and Martijn C.G.J. Brouwers<sup>1,2</sup>

<sup>1</sup>Division of Endocrinology and Metabolic Disease, Department of Internal Medicine, Maastricht University Medical Center+, Maastricht, the Netherlands

<sup>2</sup>School for Cardiovascular Diseases (CARIM), Maastricht University, Maastricht, the Netherlands <sup>3</sup>Laboratory for Metabolism and Vascular Medicine, Division of General Internal Medicine, Department of Internal Medicine, Maastricht University Medical Center+, Maastricht, the Netherlands

<sup>4</sup>Department of Epidemiology, Maastricht University, Maastricht, the Netherlands

<sup>5</sup>Care and Public Health Research Institute (CAPHRI), Maastricht University, Maastricht, the Netherlands

<sup>6</sup>Department of Radiology and Nuclear Medicine, Maastricht University Medical Center+, Maastricht, the Netherlands

<sup>7</sup>Department of Internal Medicine, Maastricht University Medical Center+, Maastricht, the Netherlands

<sup>8</sup>Heart & Vascular Centre, Maastricht University Medical Center+, Maastricht, the Netherlands

<sup>9</sup>School for Nutrition and Translational Research in Metabolism (NUTRIM), Maastricht University, Maastricht, the Netherlands

<sup>10</sup>Department of Complex Genetics and Epidemiology, Maastricht University, Maastricht, the Netherlands

<sup>11</sup>Central Diagnostic Laboratory, Department of Clinical Chemistry, Maastricht University Medical Center+, Maastricht, the Netherlands

Corresponding author: Martijn C.G.J. Brouwers, mcgj.brouwers@mumc.nl

Received 12 October 2021 and accepted 1 January 2022

This article contains supplementary material online at https://doi.org/10.2337/figshare.18032555.

© 2022 by the American Diabetes Association. Readers may use this article as long as the work is properly cited, the use is educational and not for profit, and the work is not altered. More information is available at https://www. diabetesjournals.org/journals/pages/license.

See accompanying article, p. 1032.

EPIDEMIOLOGY/HEALTH SERVICES RESEARCH

# OBJECTIVE

Epidemiological evidence regarding the relationship between fructose intake and intrahepatic lipid (IHL) content is inconclusive. We, therefore, assessed the relationship between different sources of fructose and IHL at the population level.

# **RESEARCH DESIGN AND METHODS**

We used cross-sectional data from The Maastricht Study, a population-based cohort study (n = 3,981; mean ± SD age: 60 ± 9 years; 50% women). We assessed the relationship between fructose intake (assessed with a food-frequency questionnaire)—total and derived from fruit, fruit juice, and sugar-sweetened beverages (SSB)—and IHL (quantified with 3T Dixon MRI) with adjustment for age, sex, type 2 diabetes, education, smoking status, physical activity, and intakes of total energy, alcohol, saturated fat, protein, vitamin E, and dietary fiber.

# RESULTS

Energy-adjusted total fructose intake and energy-adjusted fructose from fruit were not associated with IHL in the fully adjusted models (P = 0.647 and P = 0.767). In contrast, energy-adjusted intake of fructose from fruit juice and SSB was associated with higher IHL in the fully adjusted models (P = 0.019 and P = 0.009). Individuals in the highest tertile of energy-adjusted intake of fructose from fruit juice and SSB had a 1.04-fold (95% Cl 0.99; 1.11) and 1.09-fold (95% Cl 1.03; 1.16) higher IHL, respectively, in comparison with the lowest tertile in the fully adjusted models. Finally, the association for fructose from fruit juice was stronger in individuals with type 2 diabetes (P for interaction = 0.071).

# CONCLUSIONS

Fructose from fruit juice and SSB is independently associated with higher IHL. These cross-sectional findings contribute to current knowledge in support of measures to reduce the intake of fructose-containing beverages as a means to prevent nonalcoholic fatty liver disease at the population level.

Nonalcoholic fatty liver disease (NAFLD) is highly prevalent among people with type 2 diabetes and emerging as the principal cause of liver transplantation in Western society (1,2). Furthermore, epidemiological evidence is accumulating that

NAFLD per se is a risk factor for type 2 diabetes (2–4). Currently, a myriad of pharmacological agents that target NAFLD have entered phase II–III clinical trials (5). However, given the high global prevalence of NAFLD ( $\sim$ 25% [6]), it is also desirable to use nonpharmacological measures to reduce the burden of NAFLD and its sequela at the population level.

There has been a long debate on whether dietary fructose is a modifiable risk factor of NAFLD. Despite convincing evidence derived from animal studies (7), there have been inconsistent experimental data in humans (8,9). Furthermore, the findings from observational studies that addressed the relationship between fructose intake and intrahepatic lipid (IHL) accumulation, the first stage of NAFLD, vary from positive (10,11) to inverse (12) and divergent (13) associations. Of note, some of these studies have been conducted in selected (pediatric) groups (10), did not use histology or imaging to quantify IHL (12,13), or did not sufficiently adjust for potential confounders (10-13). In addition, only one of these studies made a distinction between multiple sources of dietary fructose (13).

The aim of the current study was, therefore, to assess the independent relationship between habitual fructose intake—total and derived from fruit, fruit juice, and sugar-sweetened beverages (SSB)—and IHL, quantified with use of 3T Dixon MRI, in The Maastricht Study, an extensively phenotyped population-based cohort (14).

# **RESEARCH DESIGN AND METHODS**

#### Study Population

The Maastricht Study is a populationbased cohort study with an oversampling of individuals with type 2 diabetes (14). In brief, the focus of The Maastricht Study is on the etiology, pathophysiology, complications, and comorbidities of type 2 diabetes. All individuals between 40 and 75 years old who lived in the southern part of the Netherlands were eligible for participation.

The current study includes cross-sectional data from 7,689 participants who completed the baseline measurements from November 2010 until December 2017. MRI measurements of the liver were implemented from December 2013 onward (available for n = 5,180). Participants with invalid MRI measurements, missing data on dietary intake, implausible energy intake, and missing data on covariates were excluded from all analyses, resulting in a study population of 3,981 participants (Supplementary Fig. 1).

The Maastricht Study has been approved by the institutional medical ethics committee (NL31329.068.10) and the Minister of Health, Welfare and Sports of the Netherlands (permit 131088-105234-PG). All participants gave written informed consent.

### Assessment of Dietary Intake

Assessment of dietary intake has previously been reported in detail (15). In brief, habitual dietary intake over the past 12 months was estimated with use of a tailor-made and validated foodfrequency questionnaire, with assessment of the frequency of consumption and the amount of consumed food and nutrients. Intakes of total energy and individual mono- and disaccharides were calculated with use of the Dutch Food Composition Database (and, in the case of missing values in this composition database, this information was complemented with values obtained from other relevant food composition databases).

For the current study, fructose intake (grams per day) was defined as the sum of 50% sucrose intake plus free fructose intake. Further, fructose-containing food items were categorized as follows: 1) total fructose (grams per day), 2) fructose from fruit (fresh and dried fruit [grams per day]), 3) fructose from fruit juice (grams per day), and 4) fructose from SSB, including sugar-containing fruit drinks and syrups (grams per day).

# Assessment of Intrahepatic Lipid Content

IHL was assessed through Dixon MRI using a 3.0 Tesla MRI system (MAGNE-TOM Prismafit; Siemens Healthineers, Erlangen, Germany) with body matrix and supine radiofrequency coils. After a scout scan, transversal two-dimensional T2-weighted true fast imaging with steady-state free precession (T2w TRUFI) images were acquired through the liver with the following parameters: voxel size  $1.2 \times 1.2 \times 5.0 \text{ mm}^3$ , repetition time (TR) 422 ms, echo time (TE) 1.65 ms, flip angle 60°, number of signal averages = 1, and parallel imaging (GeneRalized Autocalibrating Partial Parallel Acquisition [GRAPPA]) factor 2. Next, transversal two-dimensional turbo spin-echo Dixon magnetic resonance images were acquired through the liver during a breathhold with the following parameters: voxel size 2.0  $\times$  $2.0 \times 6.0 \text{ mm}^3$ , number of slices = 4, TR 500 ms, TE 31 ms, turbo factor 5, number of signal averages = 1, and parallel imaging (GRAPPA) factor 3. Three regions of interest were drawn in the liver by trained observers on the T2w TRUFI images. Subsequently, these regions of interest were copied to the water and fat Dixon magnetic resonance images for calculation of the IHL fraction.

This method was validated and calibrated against proton MRS (<sup>1</sup>H-MRS), the gold standard to noninvasively quantify IHL, in 36 participants. After calibration, the intraclass correlation coefficient between Dixon MRI and <sup>1</sup>H-MRS was 0.989 (95% CI 0.979; 0.994). IHL was expressed as the ratio of CH<sub>2</sub> to H<sub>2</sub>O (\*100%).

#### **Measurement of Covariates**

All participants completed questionnaires regarding age, sex, educational level (low, medium, high), smoking status (never, former, current smoker), and history of cardiovascular disease (CVD) (14). Use of medication was assessed during medication interviews. Weight, height, waist circumference, and office systolic and diastolic blood pressure were measured during a physical examination. Fasting levels of glucose, HbA<sub>1c</sub>, and lipid profile were measured in venous blood samples. Daily physical activity levels were measured during eight consecutive days with the activPAL3 physical activity monitor (PAL Technologies, Glasgow, U.K.) (16).

Alanine aminotransferase (ALT) was measured enzymatically on a cobas 8000 modular analyzer (Roche Diagnostics, Basel, Switzerland).

All participants underwent a standardized 2-h 75-g oral glucose tolerance test (OGTT) after an overnight fast (14). Use of insulin or fasting capillary glucose levels >11.0 mmol/L were considered as contraindications for an OGTT. Participants fulfilling either of these criteria were automatically classified as having diabetes. Glucose metabolism status, i.e., normal glucose metabolism, prediabetes (i.e., impaired fasting glucose or impaired glucose tolerance), and diabetes, was categorized based on venous plasma glucose levels obtained during an OGTT according to the World Health Organization 2006 criteria in all other participants.

The Matsuda index  $(10,000/\sqrt{G0 \times I0 \times mean G \times mean I}$ , where G0 = fasting glucose, I0 = fasting insulin, mean G = mean glucose during OGTT, and mean I = mean serum insulin levels during an OGTT) was used as a measure of insulin sensitivity (14).

#### **Statistical Analyses**

Continuous data are presented as mean  $\pm$  SD or as median (interquartile range) in case of nonnormal distribution. Categorical data are presented as *n* (%).

All nutrient variables were adjusted for total energy intake with the residual method (17).

Multivariable linear regression models were constructed for studying the associations between the energy-adjusted intake of 1) total fructose, 2) fructose from fruit, 3) fructose from fruit juice, and 4) fructose from SSB and IHL, independent of confounders. Energyadjusted fructose intake was entered as either a continuous variable (for deriving a P for trend) or a category. Based on the distribution of energy-adjusted intake of fructose from fruit, fruit juice, and SSB, with a high number of low consumers in the latter two groups (Supplementary Fig. 2), we decided to categorize the participants according to tertiles of energy-adjusted fructose intake to obtain discriminative categories of intake. The tertiles of energy-adjusted fructose intake were entered in the models as independent variables (with the lowest tertile as a reference). The following regression models were used: model 1, crude; model 2, with adjustment for age, sex, and type 2 diabetes, the latter because of the oversampling of type 2 diabetes in The Maastricht Study; model 3, with additional adjustment for (proxies of) lifestyle, i.e., educational level, smoking status, physical activity, and total energy intake; model 4, with additional adjustment for nutritional factors that have been associated with IHL in randomized controlled trials, i.e., energy-adjusted intakes of alcohol, saturated fat, protein, and vitamin E (18-21); and model 5, with additional adjustment for energy-adjusted dietary fiber, which has

been associated with IHL in observational studies (22).

IHL was <sup>10</sup>log transformed to fulfill the assumption of normality for linear regression. To obtain interpretable results we back transformed the regression coefficients, which should be interpreted as the fold change (and not the additive change) in IHL that is associated with the difference between the tertile of fructose intake and the reference group (i.e., lowest tertile of energy-adjusted fructose intake), as can be deduced from the following:

$$Log(y) = \beta_0 + \beta_1 x$$
$$Exp(log(y)) = exp(\beta_0 + \beta_1 x)$$
$$y = exp(\beta_0)exp(\beta_1 x)$$

For instance, a regression coefficient of 0.019 implies that for every unit increase in fructose intake, (<sup>10</sup>log) IHL increases with 0.019. After backtransformation, the interpretation should be that for every unit increase in fructose intake, IHL increases 1.04-fold (=  $10^{0.019}$ ), i.e., by 4%.

Additional analyses were performed to test for the effect of an interaction between energy-adjusted fructose intake and type 2 diabetes or sex on IHL in the fully adjusted model.

Several sensitivity analyses were performed. First, multivariable logistic regression analyses were conducted with hepatic steatosis as a dependent, dichotomous variable, defined as IHL  $\geq$ 5.56% (23). This cutoff value, originally expressed as  $(CH_2 / (H_2O + CH_2))$ (23), corresponds to 5.89% (= 0.0556/ (1 - 0.0556)) when IHL is expressed as CH<sub>2</sub>/H<sub>2</sub>O, as was done in the current study. Second, the original analyses were repeated with replacement of 1) the covariate type 2 diabetes in models 2-5 by the Matsuda index (available for n = 1,415) for exploration of the role of insulin sensitivity in the relationship between fructose intake and IHL and 2) IHL by (<sup>10</sup>log transformed) serum ALT levels (available for n = 1,602).

Statistical analyses were performed with the use of SPSS (version 25.0; IBM, Chicago, IL). A *P* value of <0.05 was considered statistically significant in all analyses, except for interaction tests where a less stringent significance threshold of P < 0.10 was applied.

# Data and Resource Availability

The data of this study derive from The Maastricht Study, but restrictions apply to the availability of these data, which were used under license for the current study. Data are, however, available from the authors on reasonable request and with permission of The Maastricht Study management team.

# RESULTS

# **Study Population**

Table 1 shows the characteristics of the overall population with stratification according to IHL tertiles. Mean ± SD age of the study population was  $60 \pm 9$ years, 50% were female, 20% were diagnosed with type 2 diabetes, and the median IHL was 3.2% (interquartile range 2.0-6.1). Participants in the highest IHL tertile more often were men, older, had a lower educational level, were less physically active, and had a higher BMI compared with those in the lowest IHL tertile. Compared with participants in the lowest IHL tertile, those in the highest were metabolically unhealthy, as reflected by lower HDL cholesterol and higher serum triglycerides, HbA<sub>1c</sub>, prevalence of prediabetes, and systolic and diastolic blood pressure. Further, the prevalence of CVD and the use of medication (including lipid-modifying, glucose-lowering, and antihypertensive medication) were higher in the highest IHL tertile. Finally, intakes of total fructose, fructose from fruit, and dietary fiber were lower, while intakes of total energy and saturated fat were greater, in the highest IHL tertile.

# Relationship Between Fructose Intake and Intrahepatic Lipid Content

Total fructose intake was associated with lower IHL (P < 0.001) (Table 2, models 1–3), but this association was lost after adjustment for nutritional factors that are associated with IHL (P = 0.903) (Table 2, model 4).

When fructose intake was categorized according to different sources of fructose, a similar association was observed between intake of fructose from fruit and lower IHL (P < 0.001) (Table 2, models 1–3). Again, the strength of association was attenuated after adjustment for nutritional factors that are associated with IHL (P = 0.044) (Table 2, model 4) and was completely lost after additional

	Total (n = 3,981)	First tertile (n = 1,327)	Second tertile (n = 1,327)	Third tertile ( <i>n</i> = 1,327)
IHL, %	3.2 (2.0–6.1)	1.7 (1.3–2.0)	3.2 (2.7–3.9)	8.3 (6.1–12.6)
ALT, units/L	26.0 (21.0–34.0)	22.0 (19.0–28.0)	26.0 (21.0–33.0)	31.0 (24.0–42.0)
Age, years	60 ± 9	57 ± 9	60 ± 8	61 ± 8
Women, %	50	62	46	40
Education, % low/medium/high	32/28/40	26/30/44	32/27/41	38/27/35
Smoking, % never/former/current	40/49/12	44/44/12	39/49/12	35/53/12
Physical activity, min/day	51.4 (36.6–69.6)	56.0 (40.6–73.6)	52.7 (38.6–72.0)	45.1 (31.9–62.1)
BMI, kg/m <sup>2</sup>	26.5 ± 4.1	24.3 ± 3.0	26.3 ± 3.6	28.9 ± 4.1
Waist circumference, cm	93.7 ± 12.6	85.6 ± 9.6	93.6 ± 11.0	101.8 ± 11.4
Total cholesterol, mmol/L	5.3 ± 1.1	5.3 ± 1.0	5.3 ± 1.1	5.2 ± 1.2
HDL cholesterol, mmol/L	1.6 ± 0.5	1.8 ± 0.5	1.6 ± 0.5	$1.4 \pm 0.4$
LDL cholesterol, mmol/L	3.1 ± 1.0	3.1 ± 0.9	3.1 ± 1.0	3.0 ± 1.1
Triglycerides, mmol/L	1.2 (0.9–1.7)	1.0 (0.8–1.2)	1.2 (0.9–1.6)	1.5 (1.1–2.1)
Lipid-modifying medication, %	28	17	28	39
HbA <sub>1c</sub> , % and mmol/mol	5.5 (5.2–5.9)/37	5.4 (5.1–5.6)/35	5.5 (5.3–5.8)/36	5.7 (5.4–6.4)/39
Matsuda index of insulin sensitivity	3.56 (2.14–5.34)	4.87 (3.63–6.72)	3.81 (2.60–5.42)	2.31 (1.52–3.40)
GMS, % NGM/prediabetes/type 2 diabetes/other type of diabetes	65/15/20/1	82/9/8/1	71/13/15/1	41/21/38/0
Glucose-lowering medication, %	15	7	12	27
Office SBP, mmHg	133 ± 17	128 ± 17	133 ± 17	138 ± 16
Office DBP, mmHg	75 ± 10	73 ± 10	76 ± 9	78 ± 9
Antihypertensive medication, %	33	20	31	48
History of CVD, %	13	10	13	16
Total fructose, g/day	35.9 (26.2–47.5)	36.7 (27.0–47.6)	36.3 (26.3–47.6)	34.8 (25.1–47.2)
Fructose from fruit, g/day	9.1 (4.7–14.8)	9.6 (5.3–15.5)	9.1 (5.0–15.6)	8.2 (3.9–13.4)
Fructose from fruit juice, g/day	0.9 (0.1–3.8)	1.0 (1.2–3.8)	0.9 (0.1–3.8)	0.9 (0.1–3.9)
Fructose from SSB, g/day	0.4 (0.0–2.8)	0.3 (0.0–2.1)	0.3 (0.0–2.4)	0.6 (0.0–3.8)
Total energy, kcal/day	2,074 (1,721–2,486)	2,027 (1,699–2,446)	2,087 (1,748–2,515)	2,105 (1,703–2,500)
Alcohol, g/day	8.6 (1.8–18.7)	7.8 (1.5–15.7)	9.6 (2.5–19.5)	8.3 (1.6–21.0)
Carbohydrates, g/day	222 (179–272)	223 (179–273)	224 (181–273)	219 (178–269)
Saturated fat, g/day	27.3 (20.5–35.3)	26.2 (20.2–34.3)	27.6 (20.4–35.8)	28.1 (20.8–36.1)
Protein, g/day	82.0 (68.9–96.9)	80.8 (68.0–94.8)	82.7 (69.5–98.1)	82.5 (69.3–97.6)
Vitamin E, mg/day	12.5 (9.7–16.0)	12.5 (9.8–16.0)	12.5 (9.8–16.2)	12.5 (9.6–15.8)
Dietary fiber, g/day	26.1 (21.3–31.8)	26.4 (21.4–32.0)	26.6 (21.7–32.2)	25.5 (21.1-30.9)

Table 1—Characteristics of the overall population with stratification according to IHL tertiles

Data are means  $\pm$  SD, median (interquartile range), median (interquartile range)/median, or n (%) unless otherwise indicated. Nutrient variables represent absolute intake values. DBP, diastolic blood pressure; GMS, glucose metabolism status; NGM, normal glucose metabolism; SBP, systolic blood pressure.

adjustment for dietary fiber (P = 0.767) (Table 2, model 5).

In contrast, intake of fructose from fruit juice was associated with higher IHL, also after full adjustment for potential confounders (P = 0.019) (Table 2, model 5). Individuals in the highest tertile of energy-

adjusted intake of fructose from fruit juice had a 1.04-fold (95% CI 0.99; 1.11) (Table 2, model 5) higher IHL in comparison with the lowest tertile in the fully adjusted model.

Similarly, intake of fructose from SSB was associated with higher IHL in the

fully adjusted model (P = 0.009) (Table 2, model 5). Individuals in the highest tertile of energy-adjusted intake of fructose from SSB had a 1.09-fold (95% Cl 1.03; 1.16) (Table 2, model 5) higher IHL in comparison with the lowest tertile in the fully adjusted model.

	Energy-adjusted fructose intake tertiles			
	T1	T2	Т3	P <sub>trend</sub>
Total fructose, median g/day*	24.4	35.1	47.6	
Model 1	1	0.89 (0.84; 0.95)	0.83 (0.78; 0.88)	<0.001
Model 2	1	0.94 (0.89; 0.99)	0.90 (0.85; 0.96)	<0.001
Model 3	1	0.95 (0.90; 1.00)	0.91 (0.86; 0.97)	<0.001
Model 4	1	1.01 (0.95; 1.07)	1.01 (0.95; 1.08)	0.903
Model 5	1	1.01 (0.95; 1.07)	1.02 (0.95; 1.09)	0.647
Fructose from fruit, median g/day*	3.1	9.0	17.8	
Model 1	1	0.88 (0.83; 0.94)	0.82 (0.77; 0.87)	<0.001
Model 2	1	0.90 (0.85; 0.95)	0.84 (0.80; 0.90)	<0.001
Model 3	1	0.91 (0.86; 0.97)	0.87 (0.82; 0.92)	<0.001
Model 4	1	0.94 (0.89; 1.00)	0.91 (0.86; 0.97)	0.044
Model 5	1	0.96 (0.90; 1.01)	0.95 (0.89; 1.02)	0.767
Fructose from fruit juice, median g/day*	0.1	0.9	5.3	
Model 1	1	0.95 (0.90; 1.02)	0.96 (0.90; 1.02)	0.512
Model 2	1	1.01 (0.96; 1.08)	1.02 (0.96; 1.08)	0.078
Model 3	1	1.02 (0.96; 1.08)	1.03 (0.97; 1.09)	0.082
Model 4	1	1.03 (0.97; 1.09)	1.05 (1.00; 1.12)	0.008
Model 5	1	1.02 (0.96; 1.08)	1.04 (0.99; 1.11)	0.019
Fructose from SSB, median g/day*	0.0	0.5	4.5	
Model 1	1	1.00 (0.94; 1.07)	1.09 (1.02; 1.16)	<0.001
Model 2	1	1.02 (0.96; 1.08)	1.11 (1.04; 1.17)	0.001
Model 3	1	1.02 (0.96; 1.08)	1.08 (1.02; 1.14)	0.024
Model 4	1	1.03 (0.97; 1.09)	1.12 (1.06; 1.19)	<0.001
Model 5	1	1.02 (0.96; 1.08)	1.09 (1.03; 1.16)	0.009

# Table 2—Multivariable-adjusted associations of energy-adjusted fructose intake and IHL (n = 3,981)

Regression coefficients should be interpreted as the fold change in IHL that is associated with the difference between the tertile of fructose intake and the reference group (see *Research Design and Methods*). Data in parentheses are 95% Cl.  $P_{trend}$  values were obtained from linear regression with fructose as continuous variables (see *Research Design and Methods*). Model 1: energy-adjusted intake of fructose. Model 2: additional adjustment for age, sex, and type 2 diabetes. Model 3: additional adjustment for educational level, smoking status, physical activity, and intake of total energy. Model 4: additional adjustment for energy-adjusted intakes of alcohol, saturated fat, protein, and vitamin E. Model 5: additional adjustment for energy-adjusted intake of dietary fiber. T, tertile. \*Energy-adjusted fructose by means of the residual method.

There was a statistically significant interaction between type 2 diabetes and total fructose, fructose from fruit, and fructose from fruit juice in the effect on IHL (P for interaction = 0.089, 0.058, and 0.071, respectively), and the associations were more pronounced among individuals with type 2 diabetes (Supplementary Table 1). Furthermore, individuals with type 2 diabetes in the second and third tertile of intake of fructose from SSB had a statistically significantly higher IHL in comparison with individuals without type 2 diabetes (P = 0.001 and P = 0.020) (Supplementary Table 1). Of note, the strength of these associations did not differ between individuals with newly diagnosed type 2 diabetes (based on an OGTT) and individuals with prior diagnosed type 2 diabetes (data not shown).

Sex did not modify the association between fructose intake and IHL (P for interaction >0.10; data not shown).

# Sensitivity Analyses

First, when hepatic steatosis, defined as  $IHL \ge 5.56\%$  (CH<sub>2</sub> / (H<sub>2</sub>O + CH<sub>2</sub>)), was considered as a dichotomous variable, associations were generally similar (Fig. 1). In the fully adjusted model, individuals in the highest tertile of intake of fructose from SSB were more likely to have hepatic steatosis in comparison with the lowest tertile (odds ratio [OR] 1.37 [95% CI 1.12; 1.68]) (Fig. 1). Again, individuals with type 2 diabetes in the highest tertile of intake of fructose from fruit juice had a numerically higher risk of hepatic steatosis in comparison with individuals without type 2 diabetes (OR 1.33 [95% CI 0.93; 1.90] vs. OR 1.06 [95% CI 0.85; 1.34], respectively; *P* for interaction = 0.097) (Fig. 1).

Second, replacement of the covariate type 2 diabetes by the Matsuda index in the fully adjusted model showed a robust, positive association between fructose from SSB and IHL, whereas the association between fructose from fruit juice and IHL was attenuated toward the null (Supplementary Table 2).

Last, repeated analyses with ALT as an outcome variable resulted in null associations (Supplementary Table 3).

# CONCLUSIONS

In the current study, we found that the intake of fructose from fruit juice and SSB is independently associated with higher IHL in a large, extensively phenotyped, population-based cohort. The strength of the association between fructose from fruit juice and IHL appeared to be stronger among individuals with type 2 diabetes.

Only a few studies have addressed the relationship between dietary fructose and IHL at the population level. Kanerva et al. (12) previously reported a surprisingly inverse association between total fructose intake and prevalence of NAFLD in a Finnish population-based



**Figure 1**—Association between energy-adjusted fructose intake (highest vs. lowest tertile) and hepatic steatosis among the overall population (n = 3,981) ( $\bigcirc$ ), individuals without type 2 diabetes (n = 3,171) ( $\blacksquare$ ), and individuals with type 2 diabetes (n = 810) ( $\blacktriangle$ ). Data are presented for the fully adjusted model. T, tertile.

cohort. However, this study was limited by the use of surrogate outcome measures, i.e., the fatty liver index and the NAFLD liver fat score, and, in particular, incomplete adjustment for potential confounders. Indeed, although we observed a similar crude, inverse association between total fructose intake and IHL, quantified by MRI, this association was completely abrogated after additional adjustment for nutritional factors that previously have been reported to be associated with IHL (18–21).

Further, we were able to differentiate between sources of dietary fructose in relation to IHL. For fructose from fruit, we found that the crude, inverse association with IHL was attenuated toward the null after additional adjustments including dietary fiber. It is possible that overadjustment has occurred, since fruits are rich in dietary fiber. In agreement, in a previous study in Chinese adults investigators found an inverse association between fruit intake and the presence of NAFLD but did not adjust for dietary fiber (and other relevant confounders) (24).

In contrast to the findings for fructose from fruit, we observed an association between fructose from fruit juice and SSB and higher IHL, even after adjustment for nutritional factors that are associated with IHL. The role of SSB in the development of NAFLD and type 2 diabetes has extensively been studied, however, with inconsistent results for NAFLD (25,26). Of note, when this relationship was examined in a large cohort (n = 2,634) and IHL was accurately assessed (by computed tomography), a positive association was observed, even after adjustment for confounders (27). To date, only one study addressed the association of fructose from fruit juice and a surrogate marker of NAFLD, in a relatively small cohort of healthy individuals and individuals with type 2 diabetes, and did not find an association (13). Of note, in a recent meta-analysis  $(n \approx 35,000)$  investigators did find a positive association between fruit juice consumption and incident type 2 diabetes (26).

The divergent associations of fructose from fruit and fructose from fruit juice and SSB with IHL may be explained by the food matrix, i.e., "the physical domain that contains and/or interacts with specific constituents of a food (e.g., a nutrient) providing functionalities and behaviors which are different from those exhibited by the components in isolation or a free state" (28). For instance, the presence of fiber, vitamins, flavonoids, and antioxidants might counteract the deleterious effects of fructose (29,30). The abrogation of the inverse association of fructose from fruit with IHL after adjustment for dietary fiber supports the concept of the food matrix. Alternatively, consumption of fruit could be a proxy of a healthy lifestyle

(and vice versa for fruit juice and SSB). Although we extensively corrected for lifestyle variables, residual confounding may still be present and (partly) account for the current observations.

We generally observed stronger associations for individuals with type 2 diabetes, which warrant further investigation. One potential biological explanation could be a gene-environment interaction. Geneenvironment interactions have been reported for NAFLD susceptibility genes that also predispose to type 2 diabetes (31,32). Alternatively, the observed interactions may be methodologically flawed due to underreporting bias by specific subgroups, e.g., with higher BMI or type 2 diabetes, which is a limitation inherent to nutritional epidemiology (33). Adjustment for total energy intake can overcome this source of bias, except when there is differential bias in the reporting of macronutrient intake (33). However, we did not find differences in the strength of the associations between individuals with prior diagnosed type 2 diabetes and those with newly diagnosed type 2 diabetes (i.e., who were unaware of the diagnosis), which further reduces the likelihood of underreporting bias and suggests that the associations are truly stronger among individuals with type 2 diabetes.

Previous studies have shown that advanced liver fibrosis is particularly prevalent in type 2 diabetes (34,35). In the current study we did not find any association between fructose intake and ALT, used as a marker of hepatocyte damage. This may be explained by a lack of power (serum ALT was available for n = 1,602) and/or the fairly normal ALT levels in this population. Future studies are, therefore, warranted for further investigation of (different sources of) dietary fructose in relation to liver damage and fibrosis.

The current study has several strengths and limitations. We used a large population-based cohort, enriched with individuals with type 2 diabetes, extensively phenotyped with state-of-the-art methods (e.g., 3T Dixon MRI of the liver and physical activity monitoring by an accelerometer). This allowed for an accurate estimation of the dependent variable and the adjustment for a wide range of potential confounders. Our study also has specific limitations. First, dietary intake was assessed by means of a food-frequency questionnaire, which has been validated against 24-h dietary recalls for intakes of mono- and disaccharides and fruit but not for fruit juice and SSB (15). Further, we could not differentiate between intakes of fresh fruit juice and packed fruit juice, which warrants further study. Second, although the self-reported intake of total fructose in our cohort was comparable with that of the general Dutch population (36), self-reported intakes of fructose from fruit juice and particularly from SSB were low (consistent with reduced intakes of fruit juice and SSB with increasing age in the Dutch population [36]). Our results may, therefore, not be extrapolated to populations with high fructose consumption, such as the U.S. (25), although the effects of fructose restriction, in the case of any difference, are expected to be even greater in such populations. Third, similar to a previous study (13), we only calculated intakes of fructose from homogeneous and relatively easily quantifiable food products, such as fruit, fruit juice, and SSB. We did not specifically assess the association of other dietary sources of fructose, such as vegetables and processed foods (which are more difficult to quantify), with IHL. Fourth, this is a cross-sectional study, which, by design, does not allow inference of causality. We do, however, believe that reverse causality, i.e., high IHL leads to more intake of fructose from fruit juice and SSB, is less likely. Finally, we adjusted for type 2 diabetes in the regression models because of the oversampling of type 2 diabetes in The Maastricht Study. It is likely that overadjustment has occurred, since type 2 diabetes is believed to be a consequence of IHL (3). We, therefore, performed stratified analyses and generally observed stronger associations for individuals with type 2 diabetes. The role of insulin resistance is even more complicated, as it may be the consequence of both fructose intake (= exposure) and IHL (= outcome) (37). Adjustment for the Matsuda index may, therefore, have introduced collider bias and should be interpreted cautiously (38).

In view of implications for public health, we found that individuals in the highest tertile of intake of SSB may reduce their risk of hepatic steatosis by 37% by lowering their fructose intake to the lowest tertile of intake (i.e., a reduction of  $\sim$ 4.5 g fructose from SSB/day).

The corresponding absolute reduction of 0.3 percentage points in IHL (9%-fold change [Table 2, model 5] multiplied by the population median of 3.2% [Table 1]), is small, yet in line with our recently conducted double-blind randomized controlled trial showing that fructose restriction per se resulted in a reduction of 0.7 percentage points in IHL (9). Moreover, this seemingly small reduction in IHL should be viewed in the context of the global epidemic of NAFLD. It has been estimated that onequarter of the worldwide adult population (approximately five billion people) is affected by NAFLD. Moreover, NAFLD is more frequently observed in type 2 diabetes and, in fact, is currently viewed as a risk factor of type 2 diabetes (2-4). It is, therefore, expected that a relatively easily implementable change in lifestyle, i.e., reduction of fruit juice and SSB intake, will have major beneficial health effects at the population level. This finding is of particular interest, since there is growing evidence that an excise tax on SSB—as already implemented in U.K. and U.S. cities including Berkeley (California)—has a beneficial, reducing effect on SSB consumption (39,40). Of note, fruit juice (without added sugar) is currently exempted from all these levies (39.40).

In conclusion, our population-based cohort study shows that fructose from fruit juice and SSB is associated with higher IHL, independent of confounders. These cross-sectional findings contribute to current knowledge in support of measures to reduce the intake of fructose-containing beverages as a means to prevent hepatic steatosis at the population level.

Funding and Duality of Interest. This study was supported by the Dutch Diabetes Research Foundation (personal grant 2017.82.004 to M.C.G.J.B.). The Maastricht Study was supported by the European Regional Development Fund via OP-Zuid, the Province of Limburg, the Dutch Ministry of Economic Affairs (grant 310.041), Stichting De Weijerhorst (Maastricht, the Netherlands), the Pearl String Initiative Diabetes (Amsterdam, the Netherlands), School for Cardiovascular Diseases (CARIM), School for Public Health and Primary Care (CAPHRI) (Maastricht, the Netherlands), School for Nutrition and Translational Research in Metabolism (NUTRIM) (Maastricht, the Netherlands), Stichting Annadal (Maastricht, the Netherlands), and Health Foundation Limburg (Maastricht, the Netherlands) and by unrestricted grants from Janssen-Cilag B.V. (Tilburg, the Netherlands), Novo Nordisk Farma B.V. (Alphen aan den Rijn, the Netherlands), Sanofi Netherlands B.V. (Gouda, the Netherlands), and Medtronic (Tolochenaz, Switzerland). No other potential conflicts of interest relevant to this article were reported.

Author Contributions. A.M.B., S.J.P.M.E., C.G.S., C.D.A.S., and M.C.G.J.B. contributed to the conception and design of the study. A.M.B. and M.C.G.J.B. contributed to the generation, collection, assembly, and analysis or interpretation of data and drafting the manuscript. All authors contributed to the intellectual content of the manuscript and review of the manuscript and approved the final version for submission. M.C.G.J.B. is the guarantor of this work and, as such, had full access to all the data in the study and takes responsibility for the integrity of the data and the accuracy of the data analysis.

**Prior Presentation**. Parts of this study were presented in abstract form at the virtual 57th Annual Meeting of the European Association for the Study of Diabetes, 28 September–1 October 2021.

### References

1. Charlton MR, Burns JM, Pedersen RA, Watt KD, Heimbach JK, Dierkhising RA. Frequency and outcomes of liver transplantation for nonalcoholic steatohepatitis in the United States. Gastroenterology 2011;141:1249–1253

 Targher G, Corey KE, Byrne CD, Roden M. The complex link between NAFLD and type 2 diabetes mellitus - mechanisms and treatments. Nat Rev Gastroenterol Hepatol 2021;18:599–612

3. Mantovani A, Byrne CD, Bonora E, Targher G. Nonalcoholic fatty liver disease and risk of incident type 2 diabetes: a meta-analysis. Diabetes Care 2018;41:372–382

4. Brouwers MCGJ, de Graaf J, Simons N, et al. Incidence of type 2 diabetes in familial combined hyperlipidemia. BMJ Open Diabetes Res Care 2020;8:e001107

5. Francque S, Vonghia L. Pharmacological treatment for non-alcoholic fatty liver disease. Adv Ther 2019;36:1052–1074

6. Younossi ZM, Koenig AB, Abdelatif D, Fazel Y, Henry L, Wymer M. Global epidemiology of nonalcoholic fatty liver disease-meta-analytic assessment of prevalence, incidence, and outcomes. Hepatology 2016;64:73–84

7. Im YR, Hunter H, de Gracia Hahn D, et al. A systematic review of animal models of NAFLD finds high-fat, high-fructose diets most closely resemble human NAFLD. Hepatology 2021;74: 1884–1901

8. Chiu S, Sievenpiper JL, de Souza RJ, et al. Effect of fructose on markers of non-alcoholic fatty liver disease (NAFLD): a systematic review and meta-analysis of controlled feeding trials. Eur J Clin Nutr 2014;68:416–423

9. Simons N, Veeraiah P, Simons PIHG, et al. Effects of fructose restriction on liver steatosis (FRUITLESS); a double-blind randomized controlled trial. Am J Clin Nutr 2021;113:391–400

10. DiStefano JK, Shaibi GQ. The relationship between excessive dietary fructose consumption and paediatric fatty liver disease. Pediatr Obes 2021;16:e12759

11. OSullivan TA, Oddy WH, Bremner AP, et al. Lower fructose intake may help protect against

development of nonalcoholic fatty liver in adolescents with obesity. J Pediatr Gastroenterol Nutr 2014;58:624–631

12. Kanerva N, Sandboge S, Kaartinen NE, Männistö S, Eriksson JG. Higher fructose intake is inversely associated with risk of nonalcoholic fatty liver disease in older Finnish adults. Am J Clin Nutr 2014;100:1133–1138

13. Weber KS, Simon MC, Strassburger K, et al.; GDS Group. Habitual fructose intake relates to insulin sensitivity and fatty liver index in recent-onset type 2 diabetes patients and individuals without diabetes. Nutrients 2018;10:E774

14. Schram MT, Sep SJ, van der Kallen CJ, et al. The Maastricht Study: an extensive phenotyping study on determinants of type 2 diabetes, its complications and its comorbidities. Eur J Epidemiol 2014;29:439–451

15. van Dongen MC, Wijckmans-Duysens NEG, den Biggelaar LJ, et al. The Maastricht FFQ: development and validation of a comprehensive food frequency questionnaire for the Maastricht study. Nutrition 2019;62:39–46

16. Vandercappellen EJ, Henry RMA, Savelberg HHCM, et al. Association of the amount and pattern of physical activity with arterial stiffness: the Maastricht Study. J Am Heart Assoc 2020; 9:e017502

17. Willett W, Stampfer MJ. Total energy intake: implications for epidemiologic analyses. Am J Epidemiol 1986;124:17–27

18. Mouzaki M, Allard JP. The role of nutrients in the development, progression, and treatment of nonalcoholic fatty liver disease. J Clin Gastroenterol 2012;46:457–467

19. Luukkonen PK, Sädevirta S, Zhou Y, et al. Saturated fat is more metabolically harmful for the human liver than unsaturated fat or simple sugars. Diabetes Care 2018;41:1732–1739

20. Markova M, Pivovarova O, Hornemann S, et al. Isocaloric diets high in animal or plant protein reduce liver fat and inflammation in individuals with type 2 diabetes. Gastroenter-ology 2017;152:571–585.e8

21. Perumpail BJ, Li AA, John N, et al. The role of vitamin E in the treatment of NAFLD. Diseases 2018;6:E86

22. Rietman A, Sluik D, Feskens EJM, Kok FJ, Mensink M. Associations between dietary factors and markers of NAFLD in a general Dutch adult population. Eur J Clin Nutr 2018;72:117–123

23. Szczepaniak LS, Nurenberg P, Leonard D, et al. Magnetic resonance spectroscopy to measure hepatic triglyceride content: prevalence of hepatic steatosis in the general population. Am J Physiol Endocrinol Metab 2005;288:E462–E468

24. Chan R, Wong VW, Chu WC, et al. Dietquality scores and prevalence of nonalcoholic fatty liver disease: a population study using proton-magnetic resonance spectroscopy. PLoS One 2015;10:e0139310

25. Chen H, Wang J, Li Z, et al. Consumption of sugar-sweetened beverages has a dose-dependent effect on the risk of non-alcoholic fatty liver disease: an updated systematic review and dose-response meta-analysis. Int J Environ Res Public Health 2019;16:E2192

26. Imamura F, O'Connor L, Ye Z, et al. Consumption of sugar sweetened beverages, artificially sweetened beverages, and fruit juice and incidence of type 2 diabetes: systematic review, meta-analysis, and estimation of population attributable fraction. BMJ 2015;351:h3576

27. Ma J, Fox CS, Jacques PF, et al. Sugarsweetened beverage, diet soda, and fatty liver disease in the Framingham Heart Study cohorts. J Hepatol 2015;63:462–469

28. Aguilera JM. The food matrix: implications in processing, nutrition and health. Crit Rev Food Sci Nutr 2019;59:3612–3629

29. Vasdev S, Gill V, Parai S, Longerich L, Gadag V. Dietary vitamin E and C supplementation prevents fructose induced hypertension in rats. Mol Cell Biochem 2002;241:107–114

30. Parnell JA, Raman M, Rioux KP, Reimer RA. The potential role of prebiotic fibre for treatment and management of non-alcoholic fatty liver disease and associated obesity and insulin resistance. Liver Int 2012;32:701–711

31. Stender S, Kozlitina J, Nordestgaard BG, Tybjærg-Hansen A, Hobbs HH, Cohen JC. Adiposity amplifies the genetic risk of fatty liver disease conferred by multiple loci. Nat Genet 2017;49:842–847

32. Liu DJ, Peloso GM, Yu H, et al.; Charge Diabetes Working Group; EPIC-InterAct Consortium; EPIC-CVD Consortium; GOLD Consortium; VA Million Veteran Program. Exome-wide association study of plasma lipids in >300,000 individuals. Nat Genet 2017;49:1758–1766

33. Beaton GH, Burema J, Ritenbaugh C. Errors in the interpretation of dietary assessments. Am J Clin Nutr 1997;65(Suppl. ):1100S–1107S

34. Lomonaco R, Godinez Leiva E, Bril F, et al. Advanced liver fibrosis is common in patients with type 2 diabetes followed in the outpatient setting: the need for systematic screening. Diabetes Care 2021;44:399–406

35. Ciardullo S, Monti T, Perseghin G. High prevalence of advanced liver fibrosis assessed by transient elastography among U.S. adults with type 2 diabetes. Diabetes Care 2021; 44:519–525

36. Sluik D, Engelen AI, Feskens EJ. Fructose consumption in the Netherlands: the Dutch National Food Consumption Survey 2007-2010. Eur J Clin Nutr 2015;69:475–481

37. Softic S, Stanhope KL, Boucher J, et al. Fructose and hepatic insulin resistance. Crit Rev Clin Lab Sci 2020;57:308–322

38. Rothman K. Modern Epidemiology. Philadelphia, Wolters Kluwer Health/Lippincott Williams Wilkins, 2008

39. Pell D, Mytton O, Penney TL, et al. Changes in soft drinks purchased by British households associated with the UK soft drinks industry levy: controlled interrupted time series analysis. BMJ 2021;372:n254

40. Falbe J, Thompson HR, Becker CM, Rojas N, McCulloch CE, Madsen KA. Impact of the Berkeley excise tax on sugar-sweetened beverage consumption. Am J Public Health 2016; 106:1865–1871