

Chronic elevation of plasma fibroblast growth factor 19 in long-term farnesoid X receptor agonist therapy, a happy marriage or cause for oncological concern?

Citation for published version (APA):

Schaap, F. G., Jansen, P. L. M., & Damink, S. W. M. O. (2018). Chronic elevation of plasma fibroblast growth factor 19 in long-term farnesoid X receptor agonist therapy, a happy marriage or cause for oncological concern? *Hepatology*, 67(2), 782-784. <https://doi.org/10.1002/hep.29511>

Document status and date:

Published: 01/02/2018

DOI:

[10.1002/hep.29511](https://doi.org/10.1002/hep.29511)

Document Version:

Publisher's PDF, also known as Version of record

Document license:

Taverne

Please check the document version of this publication:

- A submitted manuscript is the version of the article upon submission and before peer-review. There can be important differences between the submitted version and the official published version of record. People interested in the research are advised to contact the author for the final version of the publication, or visit the DOI to the publisher's website.
- The final author version and the galley proof are versions of the publication after peer review.
- The final published version features the final layout of the paper including the volume, issue and page numbers.

[Link to publication](#)

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal.

If the publication is distributed under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license above, please follow below link for the End User Agreement:

www.umlib.nl/taverne-license

Take down policy

If you believe that this document breaches copyright please contact us at:

repository@maastrichtuniversity.nl

providing details and we will investigate your claim.

Download date: 03 May. 2024

Chronic elevation of plasma FGF19 in long-term FXR agonist therapy, a happy marriage or cause for oncologic concern?

Frank G. Schaap^{1,2} , Peter L.M. Jansen¹, Steven W.M. Olde Damink^{1,2}

¹Department of Surgery, Maastricht University, NUTRIM School of Nutrition and Translational Research in Metabolism, Maastricht University, Maastricht, The Netherlands.

²Department of General, Visceral and Transplantation Surgery, RWTH University Hospital Aachen, Aachen, Germany.

Steven Olde Damink steven.oldedamink@maastrichtuniversity.nl

Peter Jansen plm.jansen@maastrichtuniversity.nl

Frank Schaap frank.schaap@maastrichtuniversity.nl

Disclosures:

FGS receives research support from NGM Biopharmaceuticals and Intercept Pharmaceuticals. PLMJ serves on the data management board of NGM Biopharmaceuticals, and serves on the advisory board of Intercept Pharmaceuticals.

Corresponding author:

Frank G. Schaap, PhD
Maastricht University
Department of Surgery
NUTRIM School of Nutrition and Translational Research in Metabolism
PO BOX 616
6200 MD Maastricht
The Netherlands
Phone: +31 43 388 4502
Fax: +31 43 388 4154
Email: frank.schaap@maastrichtuniversity.nl

This article has been accepted for publication and undergone full peer review but has not been through the copyediting, typesetting, pagination and proofreading process which may lead to differences between this version and the Version of Record. Please cite this article as doi: 10.1002/hep.29511

An estimated 782,000 cases of primary liver cancer were diagnosed globally in 2012, making it the fifth and ninth most frequent neoplasm in males and females, respectively. Liver cancer is the second leading cause of cancer-related mortality worldwide, and estimated to account for 746,000 deaths in 2012¹. Hepatocellular carcinoma (HCC) represents the majority of liver cancer cases, and typically develops on a background of chronic liver disease. The main risk factor for development of HCC is cirrhosis due to alcohol abuse or chronic infection with HBV or HCV. Non-alcoholic fatty liver disease (NAFLD) is already the most common chronic liver disease in developed countries and its incidence is increasing. The progressive form non-alcoholic steatohepatitis (NASH) is considered an important cause of cryptogenic cirrhosis. NAFLD-associated HCC can however develop in the absence of cirrhosis, and is expected to become more frequent as a consequence of the global obesity epidemic².

At present, there is no approved pharmaceutical therapy for NAFLD, although new treatment options are in different phases of (pre)clinical testing³. One such approach, currently evaluated in phase II/III trials, targets the metabolic and fibro-inflammatory components in NASH and comprises agonistic activation of the Farnesoid-X Receptor (FXR)⁴. FXR is a ligand-activated transcription factor that -along with its target Fibroblast Growth Factor 19 (FGF19)- mediates signaling activities of bile salts. FXR is expressed in the intestines and liver, and is activated in the postprandial state by bile salts in enterohepatic circulation. FXR and ileum-derived FGF19 regulate a number of metabolic processes in the liver (*e.g.* bile salt homeostasis, gluconeogenesis, lipogenesis), and exert an anti-inflammatory action. FXR agonists are being developed by several pharmaceutical companies. The FXR agonist obeticholic acid improved liver histology (reduction of NAFLD activity score) and halted fibrotic progression during a 72-weeks intervention study in patients with biopsy-proven NASH⁴. An initial report indicates that daily subcutaneous administration of an FGF19-based biologic (NGM282) led to a rapid reduction -and in some cases complete normalization- of hepatic fat content in patients with NASH⁵.

Successful therapy for NAFLD is expected to have several health benefits including the prevention of NAFLD-associated HCC. However, at present it is unclear if chronic elevation of circulating FGF19 in patients treated for long periods with an FXR agonist, poses an oncological risk. The tumorigenic potential of FGF19 was first uncovered in 2002, when it was shown that overexpression of FGF19 in mice results in HCC, with mice of either gender developing one or multiple liver tumors after 10 months of age. FGF19-induced hepatocarcinogenesis appears to be accelerated on a background of fatty or cholestatic liver disease⁶. Of note, FGF19 does not occur naturally in mice, and overexpression of Fgf15 (the endogenous murine equivalent of FGF19) does

not induce tumor formation in normal or diseased liver⁶. In fact, *Fgf15* is held responsible for the prevention of liver tumor formation in mice that lack *Fxr*⁷. In the absence of *Fxr*, deranged bile salt homeostasis leads to a chronic inflammatory environment in which liver tumors develop after >15 months. In this background of complete genetic *Fxr* deficiency, re-expression of constitutively active *Fxr* in the intestines, which results in upregulation of ileal *Fgf15*, is sufficient to prevent hepatocarcinogenesis. It is currently unknown whether intestinal *Fxr*/*Fgf15* signaling is protective in other experimental models of HCC. Hepatic *Fxr* regulates expression of multiple tumor suppressor genes, and agonistic activation of *Fxr* reduces growth of liver tumors in xenografted nude mice⁸. Potential beneficial actions of FXR agonists on human tumor biology are unexplored.

The mechanism behind the tumor-inducing effect of FGF19 has remained obscure for a long period of time, with little more known than a requirement for FGF Receptor 4 (FGFR4). Using an elegant series of experiments, Zhou and colleagues fill this knowledge gap, and address the relevance of the unraveled pathway for human HCC⁹. Zhou *et al.* observed that FGF19-induced hepatocarcinogenesis depends on the presence of Stat3 in hepatocytes. FGF19 induces phosphorylation of this transcription factor, resulting in its activation and enhanced transcription of genes involved in proliferation and cell survival. Intriguingly, this is not a direct effect but depends on the presence of non-parenchymal liver cells. Screening for a number of cytokines and growth factors known to cause Stat3 activation, led to the identification of Interleukin-6 (Il6) as the mediator of FGF19-driven HCC formation. This is supported by lack of FGF19-induced tumorigenesis in mice nullizygous for *Il6*, and lack of tumors in (wild-type) mice treated with anti-Il6 antibodies. Further delineation of the cellular/molecular events led Zhou *et al.* to propose the following model for FGF19-driven hepatocarcinogenesis. FGF19 induces *Il6* expression in innate immune cells of the liver. Released Il6 activates the gp130 receptor on the cell surface of hepatocytes, leading to recruitment of the effector kinase Jak that in turn phosphorylates and activates Stat3. The importance of the Il6/Jak/Stat3 axis in FGF19-mediated tumorigenesis was studied in clinically relevant models of chronic liver disease. *db/db* mice served as model for NAFLD, and *Mdr2* knock-out mice as model of primary sclerosing cholangitis (PSC). The pan-JAK inhibitor tofacitinib (approved for treatment of rheumatoid arthritis) prevents FGF19-driven HCC in *db/db* mice, and anti-Il6 antibodies block tumor formation in mice lacking *Mdr2*, with additional improvement of cholestatic liver tests and liver histology in the PSC model.

The genomic and mutational landscape in human HCC is heterogeneous. Amplification of a gene cluster (*CCND1-ORAOV1-FGF19-FGF4-FGF3*) on chromosome 11q13.3 that includes *FGF19* is observed in up to 14% of HCC cases, as well as in a variety of other human cancers including

cholangiocarcinoma^{9,10}. *FGF19* gene amplification appears mutually exclusive with inactivating mutations in the tumor suppressor gene *TP53*⁹. In HCC, *FGF19* gene amplification is accompanied by increased *FGF19* expression^{9,10}, and *in vitro* studies indicate that *FGF19* is an important driver gene in HCC¹⁰. *FGF19*-positive HCC is associated with a more aggressive tumor type and poor prognosis. Although these observations do not prove a causal relation between *FGF19* and HCC formation in humans, it is conceivable that autocrine or paracrine *FGF19* signaling via the *FGFR4*/ β Klotho receptor complex provides a mitogenic signal in a subset of human HCCs. Zhou *et al.* found no evidence for a direct mitogenic effect of *FGF19* in isolated human hepatocytes, but noted elevated expression of *STAT3* target genes in *FGF19*-positive human HCCs in comparison with *FGF19*-negative HCCs or control livers⁹. This human data is in support of the concept of Zhou and colleagues, that *FGF19* promotes HCC in a non-cell autonomous manner.

The observation that exogenous human *FGF19* induces HCC in mice while endogenous murine *Fgf15* is devoid of such activity, raises questions on the relevance of *FGF19* in human hepatocarcinogenesis. This doubt was further fostered by the supraphysiological levels (>20 ng/mL) of *FGF19* in the systemic circulation that were attained in mice with transgenic or adenoviral overexpression of *FGF19*. However, in a recent study it was demonstrated that *FGF19* levels as low as 0.6 ng/mL induce liver tumors in diabetic *db/db* mice⁹. Although a direct comparison is not substantiated, these levels are in the range that is observed in patients with cholestasis and in patients treated with FXR agonist. Hence, the study of Zhou *et al.* asks for close monitoring of patients receiving long-term therapy with agents that cause systemic elevation of *FGF19*, such as FXR agonists. Surveillance screening for HCC by frequent ultrasonography, as already implemented in the management of at-risk patients¹¹, may be considered in future clinical trials of FXR agonists. A further merit of the study of Zhou *et al.* is that it provides a rationale for personalized treatment of *FGF19*-positive liver tumors with compounds that target the *IL6/IL6R/JAK* axis.

References

1. Ferlay J, Soerjomataram I, Ervik M, Dikshit R, Eser S, Mathers C et al. GLOBOCAN 2012 v1.0, Cancer Incidence and Mortality Worldwide: IARC CancerBase No. 11 [Internet]. Lyon, France: International Agency for Research on Cancer; 2013. Available from: <http://globocan.iarc.fr>. Accessed July 2017.

2. Piscaglia F, Svegliati-Baroni G, Barchetti A, Pecorelli A, Marinelli S, Tiribelli C, et al. Clinical patterns of hepatocellular carcinoma in nonalcoholic fatty liver disease: A multicenter prospective study. *Hepatology* 2016;63:827-838.
3. Townsend SA, Newsome PN. Review article: new treatments in non-alcoholic fatty liver disease. *Aliment Pharmacol Ther* 2017; 46:494-507.
4. Neuschwander-Tetri BA, Loomba R, Sanyal AJ, Lavine JE, Van Natta ML, Abdelmalek MF, et al. Farnesoid X nuclear receptor ligand obeticholic acid for non-cirrhotic, non-alcoholic steatohepatitis (FLINT): a multicentre, randomised, placebo-controlled trial. *Lancet* 2015;385:956-965.
5. Harrison SA, Abdelmalek MF, Trotter JF, Paredes AH, Arnold HL, Kugelmas M, et al. NGM282, a novel variant of FGF19, significantly reduces hepatic steatosis and key biomarkers of NASH: results of a Phase 2, multicenter, randomized, double-blinded, placebo controlled trial in biopsy-confirmed NASH patients. *J Hepatol* 2017;66:S92-S93. Abstract.
6. Zhou M, Luo J, Chen M, Yang H, Learned RM, DePaoli AM, et al. Mouse species-specific control of hepatocarcinogenesis and metabolism by FGF19/FGF15. *J Hepatol* 2017;66:1182-1192.
7. Degirolamo C, Modica S, Vacca M, Di Tullio G, Morgano A, D'Orazio A, et al. Prevention of spontaneous hepatocarcinogenesis in farnesoid X receptor-null mice by intestinal-specific farnesoid X receptor reactivation. *Hepatology* 2015;61:161-170.
8. Huang XF, Zhao WY, Huang WD. FXR and liver carcinogenesis. *Acta Pharmacol Sin* 2015;36:37-43.
9. Zhou M, Yang H, Learned RM, Tian H, Ling L. Non-cell-autonomous activation of IL-6/STAT3 signaling mediates FGF19-driven hepatocarcinogenesis. *Nat Commun* 2017;8:15433.
10. Sawey ET, Chanrion M, Cai C, Wu G, Zhang J, Zender L, et al. Identification of a therapeutic strategy targeting amplified FGF19 in liver cancer by Oncogenomic screening. *Cancer Cell* 2011;19:347-358.
11. Bruix J, Reig M, Sherman M. Evidence-based diagnosis, staging, and treatment of patients with hepatocellular carcinoma. *Gastroenterology* 2016;150:835-853.