

Letter regarding article by Thijssen et al "Temporal and spatial variations in structural protein expression during the progression from stunned to hibernating myocardium" o Response

Citation for published version (APA):

Thijssen, V. L. J. L., Borgers, M., Lenders, M. H. J. H., Ramaekers, F. C. S., Suzuki, G., Palka, B., Fallavollita, J. A., Thomas, S. A., & Cantry jr., J. M. (2005). Letter regarding article by Thijssen et al "Temporal and spatial variations in structural protein expression during the progression from stunned to hibernating myocardium" o Response. *Circulation*, *111*, e378-e379. <https://doi.org/10.1161/CIRCULATIONAHA.104.529107>

Document status and date:

Published: 01/01/2005

DOI:

[10.1161/CIRCULATIONAHA.104.529107](https://doi.org/10.1161/CIRCULATIONAHA.104.529107)

Document Version:

Publisher's PDF, also known as Version of record

Please check the document version of this publication:

- A submitted manuscript is the version of the article upon submission and before peer-review. There can be important differences between the submitted version and the official published version of record. People interested in the research are advised to contact the author for the final version of the publication, or visit the DOI to the publisher's website.
- The final author version and the galley proof are versions of the publication after peer review.
- The final published version features the final layout of the paper including the volume, issue and page numbers.

[Link to publication](#)

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal.

If the publication is distributed under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license above, please follow below link for the End User Agreement:

www.umlib.nl/taverne-license

Take down policy

If you believe that this document breaches copyright please contact us at:

repository@maastrichtuniversity.nl

providing details and we will investigate your claim.

Download date: 20 Apr. 2024

Letter Regarding Article by Thijssen et al, "Temporal and Spatial Variations in Structural Protein Expression During the Progression From Stunned to Hibernating Myocardium"

To the Editor:

It has been stated that glycogen accumulation and myofibrillar loss are specific pathognomonic histological abnormalities of hibernating myocardium.¹ We therefore read with great interest the study of Thijssen et al,² who reported that in a porcine model of hibernating myocardium histological findings of myofibrillar changes are global not regional, and therefore seem unlikely to be caused by chronic myocardial ischemia. These findings extend the experimental findings on glycogen content reported previously by the same group.³ Thijssen et al² note that nearly all previous biopsy studies in humans have not used control samples from regions with normal contractile function within the same patient. The conclusions drawn in these reports may therefore be biased by differences in patient characteristics. Apart from the weaknesses in design as discussed by Thijssen et al, the conclusions on glycogen content in previous studies have limitations that weaken the conclusions on this other "hallmark" of hibernating myocardium. The semiquantitative periodic acid-Schiff staining method used in previous studies also stains nonglycogenic material. Furthermore, a considerable part of the glycogen pool may not be stainable by this method because it may be dissolved in the cytoplasm,⁴ the increase in periodic acid-Schiff staining thus representing changes of the composition of the glycogen pool rather than an increase in the total glycogen pool. In a previous study of patients with heart failure, we quantified biopsy content of acid-extractable and protein-bound glycogen and related it to the content of noncollagen protein in hibernating regions. We observed that these contents were similar to control regions within the same patient.⁵ These findings in humans and the experimental findings of Thijssen et al² and Thomas et al³ question the generally accepted hypothesis of glycogen accumulation and myofibrillar loss as markers of metabolic derangement.¹ It seems that the prevailing conception that glycogen accumulation and myofibrillar loss are pathognomonic of hibernating myocardium¹ needs to be revised.

Henrik Wiggers, MD, PhD
Hans Erik Bøtker, MD, PhD, DMSc
Department of Cardiology
Skejby Hospital
Aarhus University Hospital Brendstrupgaardsvej
Aarhus N, Denmark
henrikwiggers@dadlnet.dk

1. Camici PG, Wijns W, Borgers M, de Silva R, Ferrari R, Knuuti J, Lammertsma AA, Liedtke AJ, Paternostro G, Vatner SF. Pathophysiological mechanisms of chronic reversible left ventricular dysfunction due to coronary artery disease (hibernating myocardium). *Circulation*. 1997; 96:3205-3214.
2. Thijssen VL, Borgers M, Lenders MH, Ramaekers FC, Suzuki G, Palka B, Fallavollita JA, Thomas SA, Cauty JM Jr. Temporal and spatial variations in structural protein expression during the progression from stunned to hibernating myocardium. *Circulation*. 2004;110:3313-3321.
3. Thomas SA, Fallavollita JA, Suzuki G, Borgers M, Cauty JM Jr. Dissociation of regional adaptations to ischemia and global myolysis in an accelerated swine model of chronic hibernating myocardium. *Circ Res*. 2002;91:970-977.
4. Botker HE, Randsbaek F, Hansen SB, Thomassen A, Nielsen TT. Superiority of acid extractable glycogen for detection of metabolic changes during myocardial ischaemia. *J Mol Cell Cardiol*. 1995;27:1325-1332.
5. Wiggers H, Noreng M, Paulsen PK, Bottcher M, Egeblad H, Nielsen TT, Botker HE. Energy stores and metabolites in chronic reversibly and irreversibly dysfunctional myocardium in humans. *J Am Coll Cardiol*. 2001;37:100-108.

Response

We thank Drs Wiggers and Bøtker for their interest in our study demonstrating that structural remodeling in cardiomyocytes occurs independently of chronic ischemia and most likely reflects a global response to chronic elevations in left ventricular preload. Wiggers and associates have previously described a similar observation with regard to metabolite storage and glycogen levels in patients with heart failure: As referred to in their letter, they observed equal levels of total glycogen content in normal and chronic reversibly dysfunctional myocardium.

The observations from both studies indicate a discrepancy between some of the previously defined characteristics of hibernation and chronic ischemia as being the supposed underlying trigger. In fact, besides our own observations and those of Wiggers et al, an increasing number of studies have reported a dissociation between cardiomyocyte remodeling and ischemia.¹⁻³ Thus, we support the view of Wiggers and Bøtker that the hibernating myocyte phenotype cannot be directly attributed to an adaptive response arising from regional alterations in myocardial perfusion, and that glycogen accumulation and myofibrillar loss are probably not pathognomonic of hibernating myocardium.

Myolysis, glycogen storage, and structural remodeling have been demonstrated at the structural and ultrastructural levels in a variety of different cardiac pathologies. It appears that this adaptive phenotype is a global response of the cardiomyocytes to cardiac stress, independent of regional flow reductions or ischemia. It remains unknown which triggers underlie this response, but we have shown that elevated preload may be involved.⁴ This could also account for the "hibernating" phenotype of atrial cardiomyocytes during chronic atrial fibrillation.⁵ The ultimate reversibility of this phenotype has not been clearly established other than in atrial fibrillation, in which it does not completely normalize.

Many new insights have emerged about the pathophysiology of viable chronically dysfunctional myocardium during the last several years, including the global nature of structural remodeling, transmural variations, reduced flow being a result rather than a cause, metabolic adaptation, and the importance of coronary flow reserve in the progression from stunning to hibernation. Insights from human studies are now complemented by chronic animal models that recapitulate most, if not all, of the features of human pathophysiology. We would applaud a critical revision and review of the criteria distinguishing the multiple pathophysiological entities underlying reversible contractile dysfunction with the recognition that the "hibernating cardiomyocyte" is a general adaptive cardiac phenotype to stress.

V.L.J.L. Thijssen, PhD
Department of Pathology
Research Institute for Growth and Development
Maastricht University
Maastricht, The Netherlands
v.thijssen@path.unimaas.nl

M. Borgers, PhD
M.-H. Lenders, BSc
F.C.S. Ramaekers, PhD
Department of Molecular Cell Biology
Cardiovascular Research Institute Maastricht
Maastricht University
Maastricht, The Netherlands

G. Suzuki, MD, PhD
B. Palka, BS
J.A. Fallavollita, MD
S.A. Thomas, PhD
J.M. Canty, Jr, MD

*Veterans Affairs Western New York Health Care System
Departments of Medicine and Physiology and Biophysics
State University of New York at Buffalo
Buffalo, NY*

1. Sherman AJ, Klocke FJ, Decker RS, Decker ML, Kozlowski KA, Harris KR, Hedjbeli S, Yaroshenko Y, Nakamura S, Parker MA, Checchia PA, Evans DB. Myofibrillar disruption in hypocontractile myocardium showing perfusion- contraction matches and mismatches. *Am J Physiol Heart Circ Physiol.* 2000;278:H1320–H1334.
2. Gunning MG, Kaprielian RR, Pepper J, Pennell DJ, Sheppard MN, Severs NJ, Fox KM, Underwood SR. The histology of viable and hibernating myocardium in relation to imaging characteristics. *J Am Coll Cardiol.* 2002;39:428–435.
3. Thomas SA, Fallavollita JA, Suzuki G, Borgers M, Canty JM Jr. Dissociation of regional adaptations to ischemia and global myolysis in an accelerated swine model of chronic hibernating myocardium. *Circ Res.* 2002;91:970–977.
4. Thijssen VL, Borgers M, Lenders MH, Ramaekers FC, Suzuki G, Palka B, Fallavollita JA, Thomas SA, Canty JM Jr. Temporal and spatial variations in structural protein expression during the progression from stunned to hibernating myocardium. *Circulation.* 2004;110:3313–3321.
5. Thijssen VL, Ausma J, Borgers M. Structural remodelling during chronic atrial fibrillation: act of programmed cell survival. *Cardiovasc Res.* 2001; 52:14–24.