

Conveying temporal information to the auditory system via transcranial current stimulation

Citation for published version (APA):

Riecke, L., & Zoefel, B. (2018). Conveying temporal information to the auditory system via transcranial current stimulation. *Acta Acustica united with Acustica*, 104(5), 883-886.
<https://doi.org/10.3813/AAA.919235>

Document status and date:

Published: 01/01/2018

DOI:

[10.3813/AAA.919235](https://doi.org/10.3813/AAA.919235)

Document Version:

Publisher's PDF, also known as Version of record

Document license:

Taverne

Please check the document version of this publication:

- A submitted manuscript is the version of the article upon submission and before peer-review. There can be important differences between the submitted version and the official published version of record. People interested in the research are advised to contact the author for the final version of the publication, or visit the DOI to the publisher's website.
- The final author version and the galley proof are versions of the publication after peer review.
- The final published version features the final layout of the paper including the volume, issue and page numbers.

[Link to publication](#)

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal.

If the publication is distributed under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license above, please follow below link for the End User Agreement:

www.umlib.nl/taverne-license

Take down policy

If you believe that this document breaches copyright please contact us at:

repository@maastrichtuniversity.nl

providing details and we will investigate your claim.

Conveying Temporal Information to the Auditory System via Transcranial Current Stimulation

Lars Riecke¹⁾, Benedikt Zoefel²⁾

¹⁾ Department of Cognitive Neuroscience, Faculty of Psychology and Neuroscience, Maastricht University, 6229 EV Maastricht, The Netherlands. L.Riecke@MaastrichtUniversity.nl

²⁾ MRC Cognition and Brain Sciences Unit, University of Cambridge, Cambridge CB2 7EF, United Kingdom

Summary

This review paper investigates whether non-invasive application of electric current to the human scalp can be utilized to convey perceptually relevant temporal information to the auditory system. Recent studies have corroborated this notion by demonstrating that transcranial current stimulation (TCS) with temporally structured (sinusoidal and/or sound envelope-shaped) current biases neural processing and auditory perception toward the temporal pattern of the applied current. However, the perceptual benefits achieved with TCS so far are fairly modest. In sum, the temporally specific modulatory ability of TCS makes it a useful scientific tool for identifying temporal mechanisms for auditory perception. Practical or clinical applications (e.g., to enhance or restore auditory functions in normal or hearing-impaired populations) are currently still premature and require further optimization of stimulation parameters.

© 2018 The Author(s). Published by S. Hirzel Verlag · EAA. This is an open access article under the terms of the Creative Commons Attribution (CC BY 4.0) license (<https://creativecommons.org/licenses/by/4.0/>).

PACS no. 43.64.Me, 43.64.Ri, 43.64.Sj, 43.66.Ba, 43.66.Mk, 43.66.Ts, 43.71.Qr

1. Introduction

TCS is a non-invasive and soundless brain stimulation technique that is becoming increasingly popular in auditory research, as reflected by a rapidly growing body of published studies over the last years [1, 2]. TCS involves the non-invasive and harmless application of low-intensity current (typically within ± 2 mA) via scalp electrodes. Current-flow simulations and intracranial recordings in humans [3, 4] have shown that the induced current spreads mainly along the highly conductive skin and to a smaller amount into the cranium, where it propagates widely across the brain. Extracellular electric fields applied to pyramidal neurons in vitro almost instantaneously elevate (or reduce, depending on the relative field orientation) spontaneous neural firing (e.g., [5]). Thus, intracranial stimulation with temporally structured (alternating or complex-shaped) current can entrain neural excitability to the temporal pattern of that current [6]. Application of the current at the scalp (TCS) is thought to induce corresponding excitability changes at the neural population level [7, 8]. Neural oscillations in neocortex align their high-excitability phases to expected relevant sensory events, thereby providing a mechanism for temporal filtering of sensory input [9]. If TCS with temporally-structured cur-

rent was also able to control the timing of neural excitability, then this technique could be utilized (in analogy to neural oscillations) to provide the brain with temporal information that is critical for extracting specific events from the sensory environment. The presumed ability of TCS to entrain neural oscillations [10] would be evidenced by demonstrating that neural excitability—and therewith the neural processing of sensory stimuli—covaries with the temporal pattern of TCS. Similarly, the presumed ability of TCS to convey perceptually relevant temporal information would require showing that perception of sensory stimuli covaries with the temporal pattern of TCS. Such evidence would have interesting consequences for both scientific research and clinical/practical application: It would confirm that TCS can effectively manipulate specific temporal activity patterns (including neural oscillations), which would enable to experimentally identify causal roles of these patterns in auditory functions, such as hearing, auditory scene analysis, and speech comprehension. Moreover, it would suggest that TCS can be utilized to enhance or restore these functions in normal and hearing-impaired populations [11].

The goal of this review paper is to assess whether TCS can be utilized to provide the human auditory system with perceptually relevant temporal information. The focus is on auditory studies that have applied TCS with temporally structured currents. Auditory studies using TCS variants that seem less suited for conveying temporal information (direct currents, random noise currents, or alternating cur-

Received 28 February 2018,
accepted 17 July 2018.

rents bearing no temporal relationship to acoustic input) have been reviewed elsewhere [1, 2].

2. TCS can entrain neural oscillations

Electric stimulation with alternating current can entrain spontaneous neural firing (measured intracranially) in animal neocortex, even when the alternating current is applied transcranially (TACS) [12]. Studies attempting to translate these animal findings to humans found that TACS elevates power at the TACS frequency in subsequent scalp magneto-/electroencephalographic recordings (M/EEG) [13]. However, these aftereffects more likely reflect TACS-induced synaptic plasticity than oscillatory phase entrainment [14]. More convincing evidence of TACS-induced entrainment would show that oscillations follow the applied current waveform *during* the stimulation. Using sophisticated approaches to reduce TACS-induced artefacts in simultaneous M/EEG recordings (e.g., [15, 16]), several studies have indeed shown an *online* increase in both power at the TACS frequency [17, 18] and phase-locking between TACS waveform and M/EEG oscillation [16, 17, 19]. While the minimum current intensity required to reliably entrain human neocortical oscillations is still being debated [3, 4, 20, 21], the aforementioned studies strongly suggest that TACS can entrain neural oscillations and thereby convey temporal information to sensory systems. In future studies, it needs to be shown whether TCS with more complex- shaped (aperiodically fluctuating) currents resembling naturally occurring sensory signals (e.g., speech) can align endogenous neural activity as well [6].

3. TCS can modulate auditory perception in a temporally specific manner

Several studies have shown that TCS applied above temporal cortex can modulate auditory perception in a temporally specific manner. Neuling *et al.* [22] applied direct current modulated at 10 Hz. They reported that the threshold for detecting tone pips in a simultaneous noise masker depends on the relative phase of the simultaneously applied 10-Hz electric stimulation. The size of this *phase-effect* on simultaneous-masked threshold was ~ 0.3 dB. Using a similar paradigm, Riecke *et al.* [23] found that detection of 4-Hz click trains in silence depends on the relative phase of the click train and simultaneously applied 4-Hz TACS (size of phase effect on click-detection accuracy: $\sim 2\%$). Consistent with these click-detection results, Riecke *et al.* [24] found that the perceptual buildup of 4-Hz-modulated complex tones in informational maskers (i.e., the time required for listeners to perceptually segregate a rhythmic target stream from background noise) depends on the relative phase of the target stream and 4-Hz TACS (size of phase effect on buildup time: ~ 40 ms). Because the applied current carried temporal cues regarding the occurrence of the target stream constituents, this observation

supports the notion that TCS can provide perceptually relevant temporal information to the auditory system.

Temporal information might be of particular importance for speech processing, as a rapid sequence of densely structured information has to be precisely deciphered. The amplitude envelope of the speech signal is critical for this, as its disruption strongly impairs speech comprehension (e.g. [25]). Indeed, several recent studies show that speech processing can be manipulated by applying external stimulation carrying speech-envelope information. E.g., Riecke *et al.* [26] investigated speech comprehension under 4-Hz TACS in a cocktail party-like two-talker situation. Speech envelopes were artificially fixed at 4 Hz and enhanced, and the two simultaneous speech signals were mixed so that their envelopes alternated. Consistent with results on click- and tone-detection, speech-recognition accuracy was found to depend on the relative phase of the target-speech envelope and TACS (size of phase-effect on speech-recognition accuracy: $\sim 3\%$). Zoefel *et al.* [27] combined TACS at 3.125 Hz with functional magnetic resonance imaging (fMRI) while participants listened to rhythmic sentences (spoken at the TACS frequency) that were modified with vocoders to resemble either intelligible speech or only broadband noise fluctuating at the speech rhythm. Speech-evoked fMRI responses in superior temporal gyrus were found to depend on the relative phase of the auditory speech rhythm and TACS. Strikingly, this effect was only observed for intelligible speech, but not fluctuating noise. Moreover, certain relative phases resulted in suppression of speech-evoked responses (compared with sham stimulation). Importantly, these results provide a potential neural mechanism underlying the aforementioned TACS-phase effects on speech recognition [26].

Two recent studies also applied TCS at different time lags relative to speech sounds; however, current waveforms matched the envelope of the natural speech stimuli. Wilsch *et al.* [28] observed that speech-in-noise recognition at listeners' individual "best" time lag (the lag revealing the smallest threshold) is better than under sham stimulation, with a threshold difference between individual best lag and sham of ~ 0.4 dB. In a second experiment, Riecke *et al.* [26] removed the low-frequency envelope from auditory speech stimuli and applied it via TCS to restore this "aurally missing" temporal information in the brain. Speech comprehension was found to depend on the lag between envelope-reduced speech and envelope-shaped TCS. The maximal benefit was observed when TCS led the acoustic input by 375 ms, with a difference in word-recognition accuracy of $\sim 4\%$ (average best lag vs. average worst lag). In sum, these studies show consistently that TCS applied above temporal cortex can convey perceptually relevant temporal information to the auditory system.

4. TACS frequency modulates perceptual sampling frequency

It is conceivable that TACS transmits temporal information via not only its relative phase but also its *frequency* – in form of a sampling rhythm that affects the temporal resolution of perception. Based on the observation that gap-detection thresholds correlate negatively with listeners' gamma resonance frequency in auditory cortex (defined as the modulation frequency within 20–70 Hz that evokes the strongest auditory steady-state EEG response), Baltus *et al.* used TACS near individual gamma-resonance frequency to manipulate the temporal resolution of the auditory system [29, 30]. They found that TACS with a frequency slightly above (vs. below) resonance frequency reduced gap-detection threshold. This indicates that TACS can modulate the rate at which the human auditory system samples acoustic input.

5. Clinical utility of TCS and current limitations

There currently exists only little evidence to support an effective clinical/practical utilization of rhythmic auditory TCS. TCS effects observed so far are rather modest (see section 3) and their actual size and direction (benefit vs disruption) are difficult to estimate. E.g., most studies correct for inter-individual differences in the best time lag (presumed to originate from anatomical variations [19]) to improve the power of group-level analyses. However, this approach requires sacrificing the listeners' maximum-performance data, which inevitably results in an underestimation of both average effect sizes and potential benefits. Indeed, beyond the effects of TCS timing, only few studies could validly detect significant differences between TCS and sham stimulation. Thus, it remains to be shown whether TCS can reliably provide sufficiently strong auditory perceptual benefits, as required for a clinical or practical application.

Potential benefits may be strengthened by systematically improving TCS parameters to promote the specific spatiotemporal brain-activity patterns that underlie normal auditory functions [31]. While the TCS studies reviewed here have primarily considered basic temporal parameters, future research may focus on complementary (spatial and spectral) parameters, e.g., by comparing the benefits of different electrode shapes, locations, and numbers. Clinical TCS utilization also requires more basic research on the mechanisms underlying the observed TCS effects, which are still being debated [20, 21, 32]. While it seems safe to conclude that TACS with sufficient intensity can entrain endogenous neural activity and modulate auditory perception (sections 2 and 3), it is still unclear how this entrainment originates: directly in the cortex, indirectly via rhythmic peripheral responses elicited by the current, or both. An exclusive mediating role of peripheral responses seems unlikely because TACS applied above the mastoids

does not entrain otoacoustic emissions [33] and TACS-phase effects on auditory cortical speech processing have been observed exclusively for auditory stimuli that can be identified as speech [27]. Nevertheless, even though the TCS studies reviewed here attempted to reduce potential tactile or visual sensations [34], the applied currents might have induced subliminal tactile or visual temporal cues that potentially facilitated (or disrupted) neural activity. Future studies need to rule out this possibility by applying TCS also to non-auditory control locations (e.g., near the eyes or on the hand).

6. Conclusions

Our review shows that the timing of TCS modulates endogenous neural activity, auditory cortical speech processing, and various perceptual auditory functions (hearing in silence or noise, auditory streaming, and speech comprehension). This converging evidence indicates that TCS can convey temporal information to the auditory system (by entraining slow endogenous neural activity) and therewith alter processing and perception of acoustic input. This makes TCS a highly useful scientific tool for identifying causal dynamic mechanisms underlying perceptual auditory functions. However, perceptual benefits achieved with current stimulation settings are still modest, indicating that any clinical or practical utilization of TCS (e.g., in assisted hearing) still requires substantial methodological improvement.

References

- [1] K. Heimrath, M. Fiene, K. S. Rufener, T. Zaehle: Modulating human auditory processing by transcranial electrical stimulation. *Front Cell Neurosci* **10** (2016) 53.
- [2] B. Zoefel, M. H. Davis: Transcranial electric stimulation for the investigation of speech perception and comprehension. *Lang Cogn Neurosci* **32** (2017) 910–923.
- [3] A. Opitz, A. Falchier, C. G. Yan, E. M. Yeagle, G. S. Linn, P. Megevand, A. Thielscher, A. R. Deborah, M. P. Milham, A. D. Mehta, et al.: Spatiotemporal structure of intracranial electric fields induced by transcranial electric stimulation in humans and nonhuman primates. *Sci Rep* **6** (2016) 31236.
- [4] Y. Huang, A. A. Liu, B. Lafon, D. Friedman, M. Dayan, X. Wang, M. Bikson, W. K. Doyle, O. Devinsky, L. C. Parra: Measurements and models of electric fields in the in vivo human brain during transcranial electric stimulation. *Elife* **6** (2017).
- [5] M. Bikson, M. Inoue, H. Akiyama, J. K. Deans, J. E. Fox, H. Miyakawa, Jefferys, J. G.: Effects of uniform extracellular dc electric fields on excitability in rat hippocampal slices in vitro. *J Physiol* **557** (2004) 175–190.
- [6] F. Frohlich, D. A. McCormick: Endogenous electric fields may guide neocortical network activity. *Neuron* **67** (2010) 129–143.
- [7] A. Priori: Brain polarization in humans: a reappraisal of an old tool for prolonged non-invasive modulation of brain excitability. *Clin Neurophysiol* **114** (2003) 589–595.
- [8] W. Paulus: Transcranial electrical stimulation (tES - tDCS; tRNS, tACS) methods. *Neuropsychol Rehabil* **21** (2011) 602–617.

- [9] C. E. Schroeder, P. Lakatos: Low-frequency neuronal oscillations as instruments of sensory selection. *Trends Neurosci* **32** (2009) 9–18.
- [10] C. S. Herrmann, S. Rach, T. Neuling, D. Struber: Transcranial alternating current stimulation: a review of the underlying mechanisms and modulation of cognitive processes. *Front Hum Neurosci* **7** (2013) 279.
- [11] D. J. Schutter, M. Wischniewski: A meta-analytic study of exogenous oscillatory electric potentials in neuroenhancement. *Neuropsychologia* **86** (2016) 110–118.
- [12] S. Ozen, A. Sirota, M. A. Belluscio, C. A. Anastassiou, E. Stark, C. Koch, G. Buzsaki: Transcranial electric stimulation entrains cortical neuronal populations in rats. *J Neurosci* **30** (2010) 11476–11485.
- [13] D. Veniero, A. Vossen, J. Gross, G. Thut: Lasting EEG/MEG aftereffects of rhythmic transcranial brain stimulation: Level of control over oscillatory network activity. *Front Cell Neurosci* **9** (2015) 477.
- [14] A. Vossen, J. Gross, G. Thut: Alpha power increase after transcranial alternating current stimulation at alpha frequency (alpha-tACS) reflects plastic changes rather than entrainment. *Brain Stimul* **8** (2015) 499–508.
- [15] N. Noury, M. Siegel: Analyzing EEG and MEG signals recorded during tES, a reply. *Neuroimage* **167** (2017) 53–61.
- [16] M. Witkowski, E. Garcia-Cossio, B. S. Chander, C. Braun, N. Birbaumer, S. E. Robinson, S. R. Soekadar: Mapping entrained brain oscillations during transcranial alternating current stimulation (tACS). *Neuroimage* **140** (2016) 89–98.
- [17] R. F. Helfrich, T. R. Schneider, S. Rach, S. A. Trautmann-Lengsfeld, A. K. Engel, C. S. Herrmann: Entrainment of brain oscillations by transcranial alternating current stimulation. *Curr Biol* **24** (2014) 333–339.
- [18] T. Neuling, P. Ruhnau, M. Fusca, G. Demarchi, C. S. Herrmann, N. Weisz: Friends, not foes: Magnetoencephalography as a tool to uncover brain dynamics during transcranial alternating current stimulation. *Neuroimage* **118** (2015) 406–413.
- [19] P. Ruhnau, T. Neuling, M. Fusca, C. S. Herrmann, G. Demarchi, N. Weisz: Eyes wide shut: Transcranial alternating current stimulation drives alpha rhythm in a state dependent manner. *Sci Rep* **6** (2016) 27138.
- [20] B. Lafon, S. Henin, Y. Huang, D. Friedman, L. Melloni, T. Thesen, W. Doyle, G. Buzsaki, O. Devinsky, L. C. Parra, et al.: Low frequency transcranial electrical stimulation does not entrain sleep rhythms measured by human intracranial recordings. *Nat Commun* **8** (2017) 1199.
- [21] M. Voroslakos, Y. Takeuchi, K. Brinyiczki, T. Zombori, A. Oliva, A. Fernandez-Ruiz, G. Kozak, Z. T. Kincses, B. Ivanyi, G. Buzsaki, et al.: Direct effects of transcranial electric stimulation on brain circuits in rats and humans. *Nat Commun* **9** (2018) 483.
- [22] T. Neuling, S. Rach, S. Wagner, C. H. Wolters, Herrmann, C. S.: Good vibrations: oscillatory phase shapes perception. *Neuroimage* **63** (2012) 771–778.
- [23] L. Riecke, E. Formisano, C. S. Herrmann, A. T. Sack: 4-Hz transcranial alternating current stimulation phase modulates hearing. *Brain Stimul* **8** (2015) 777–783.
- [24] L. Riecke, A. T. Sack, C. E. Schroeder: Endogenous delta/theta sound-brain phase entrainment accelerates the build-up of auditory streaming. *Curr Biol* **25** (2015) 3196–3201.
- [25] R. Drullman, J. M. Festen, R. Plomp: Effect of temporal envelope smearing on speech reception. *J Acoust Soc Am* **95** (1994) 1053–1064.
- [26] L. Riecke, E. Formisano, B. Sorger, D. Baskent, E. Gaudrain: Neural entrainment to speech modulates speech intelligibility. *Curr Biol* **28** (2018) 161–169 e165.
- [27] B. Zoefel, A. Archer-Boyd, M. H. Davis: Phase entrainment of brain oscillations causally modulates neural responses to intelligible speech. *Curr Biol* **28** (2018) 401–408 e405.
- [28] A. Wilsch, T. Neuling, J. Obleser, C. S. Herrmann: Transcranial alternating current stimulation with speech envelopes modulates speech comprehension. *Neuroimage* (2018).
- [29] A. Baltus, C. S. Herrmann: Auditory temporal resolution is linked to resonance frequency of the auditory cortex. *Int J Psychophysiol* **98** (2015) 1–7.
- [30] A. Baltus, S. Wagner, C. H. Wolters, C. S. Herrmann: Optimized auditory transcranial alternating current stimulation improves individual auditory temporal resolution. *Brain Stimul* **11** (2018) 118–124.
- [31] V. Romei, G. Thut, J. Silvanto: Information-based approaches of noninvasive transcranial brain stimulation. *Trends Neurosci* **39** (2016) 782–795.
- [32] P. Ruhnau, K. S. Rufener, H. J. Heinze, T. Zaehle: Sailing in a sea of disbelief: In vivo measurements of transcranial electric stimulation in human subcortical structures. *Brain Stimul* **11** (2018) 241–243.
- [33] M. A. Ueberfuhr, A. Braun, L. Wiegrebe, B. Grothe, M. Drexler: Modulation of auditory percepts by transcutaneous electrical stimulation. *Hear Res* **350** (2017) 235–243.
- [34] Z. Turi, G. G. Ambrus, K. Janacsek, K. Emmert, L. Hahn, W. Paulus, A. Antal: Both the cutaneous sensation and phosphene perception are modulated in a frequency-specific manner during transcranial alternating current stimulation. *Restor Neurol Neurosci* **31** (2013) 275–285.