

Alcohol consumption in relation to cardiovascular risk and mortality

Citation for published version (APA):

Costanzo, S. (2012). *Alcohol consumption in relation to cardiovascular risk and mortality*. [Doctoral Thesis, Maastricht University]. Maastricht University. <https://doi.org/10.26481/dis.20120126sc>

Document status and date:

Published: 01/01/2012

DOI:

[10.26481/dis.20120126sc](https://doi.org/10.26481/dis.20120126sc)

Document Version:

Publisher's PDF, also known as Version of record

Please check the document version of this publication:

- A submitted manuscript is the version of the article upon submission and before peer-review. There can be important differences between the submitted version and the official published version of record. People interested in the research are advised to contact the author for the final version of the publication, or visit the DOI to the publisher's website.
- The final author version and the galley proof are versions of the publication after peer review.
- The final published version features the final layout of the paper including the volume, issue and page numbers.

[Link to publication](#)

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal.

If the publication is distributed under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license above, please follow below link for the End User Agreement:

www.umlib.nl/taverne-license

Take down policy

If you believe that this document breaches copyright please contact us at:

repository@maastrichtuniversity.nl

providing details and we will investigate your claim.

5. Summary

A consistent, significant protective effect of regular and moderate alcohol consumption in coronary heart disease and ischaemic stroke, already shown in previous epidemiological studies, has been clearly confirmed and established by the work assembled in the present thesis. The association between alcohol consumption and CHD mortality was described indeed through a J-shaped relationship, where teetotalers and heavy drinkers were at the highest risk whereas light-moderate drinkers were at the lowest risk.

Thus, if low alcohol intake is inversely related to CHD, the other side of the coin shows an increased risk for certain cancers, cirrhosis and death from accidents mainly associated with increasing alcohol consumption.

Total mortality too was significantly reduced in moderate drinkers; however, excess drinking was definitely harmful also when this hard global end-point was considered. Thus, data assembled in this thesis and previous epidemiologic data—though only derived at present from a large number of prospective observational studies — confirm the hazards of excess drinking, but also indicate the existence of potential windows of alcohol intake which may confer a net beneficial effect of drinking, in terms of survival, both in men and in women.

Regarding cardiovascular patients, if not contraindicated, regular alcohol consumers should not exceed one drink/day for women or up to two drinks/day for men as a component of a balanced cardio-protective dietary pattern, with appropriate energy intake levels. At present, according to a prudent approach, a cardiovascular patient who is teetotaler should neither be recommended, nor "prescribed" to start drinking for health gain.

In spite of a large number of experimental studies suggesting a protective role of wine-derived polyphenols on cardiovascular risk, epidemiological evidence of a similar effect of wine and beer has been now established and it seems likely that the ethanol component of different beverages is a major factor that reduces the cardiovascular risk.

The present thesis provides reasonable evidence that regular and moderate alcohol intake is significantly associated with a reduction in the incidence of cardiovascular and total mortality, in apparently healthy people and in patients at high

cardiovascular risk. However, the fact that regular moderate alcohol consumption is not easily attainable or maintainable in all parts of the world must be taken into serious consideration. There are certainly appreciable differences between Mediterranean and Northern European or Russian habits of alcohol consumption. Therefore, in some low-income populations and poor countries, even if the net effect on CVD might be beneficial, the effect of alcohol on the overall burden of disease might be detrimental because of more frequent uncontrolled alcohol-use disorders, cancer, liver cirrhosis, and injury.