

# Respectful Caring for the Agitated Elderly (ReCAGE)

Citation for published version (APA):

Mendes, A., Bergh, S., Cesana, B. M., Handels, R., Ciccone, A., Cognat, E., Fabbo, A., Fascendini, S., Frisoni, G. B., Froelich, L., Jori, M. C., Mecocci, P., Merlo, P., Peters, O., Tsolaki, M., & Defanti, C. A. (2023). Respectful Caring for the Agitated Elderly (ReCAGE): A Multicentre, Prospective, Observational Study to Evaluate the Effectiveness of Special Care Units for People with Dementia. *Journal of Alzheimer's Disease*, 96(3), 1083-1096. Article JAD-230708. <https://doi.org/10.3233/JAD-230708>

## Document status and date:

Published: 21/11/2023

## DOI:

[10.3233/JAD-230708](https://doi.org/10.3233/JAD-230708)

## Document Version:

Publisher's PDF, also known as Version of record

## Document license:

Taverne

## Please check the document version of this publication:

- A submitted manuscript is the version of the article upon submission and before peer-review. There can be important differences between the submitted version and the official published version of record. People interested in the research are advised to contact the author for the final version of the publication, or visit the DOI to the publisher's website.
- The final author version and the galley proof are versions of the publication after peer review.
- The final published version features the final layout of the paper including the volume, issue and page numbers.

[Link to publication](#)

## General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal.

If the publication is distributed under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license above, please follow below link for the End User Agreement:

[www.umlib.nl/taverne-license](http://www.umlib.nl/taverne-license)

## Take down policy

If you believe that this document breaches copyright please contact us at:

[repository@maastrichtuniversity.nl](mailto:repository@maastrichtuniversity.nl)

providing details and we will investigate your claim.

# Respectful Caring for the Agitated Elderly (ReCAGE): A Multicentre, Prospective, Observational Study to Evaluate the Effectiveness of Special Care Units for People with Dementia

Aline Mendes<sup>a,\*</sup>, Sverre Bergh<sup>b</sup>, Bruno Mario Cesana<sup>c</sup>, Ron Handels<sup>d</sup>, Alfonso Ciccone<sup>e</sup>, Emmanuel Cognat<sup>f</sup>, Andrea Fabbo<sup>g</sup>, Sara Fascendini<sup>h</sup>, Giovanni B. Frisoni<sup>i</sup>, Lutz Froelich<sup>j</sup>, Maria Cristina Jori<sup>k</sup>, Patrizia Mecocci<sup>l,m</sup>, Paola Merlo<sup>n</sup>, Oliver Peters<sup>o</sup>, Magdalini Tsolaki<sup>p</sup> and Carlo Alberto Defanti<sup>h</sup>

<sup>a</sup>*Division of Geriatrics and Rehabilitation, University Hospitals of Geneva and University of Geneva, Geneva, Switzerland*

<sup>b</sup>*Research Center for Age-Related Functional Decline and Disease, Innlandet Hospital Trust, Ottestad, Norway*

<sup>c</sup>*Department of Clinical Sciences and Community Health, Unit of Medical Statistics, Biometry and Bioinformatics “Giulio A. Maccacaro” Faculty of Medicine and Surgery, University of Milan, Milan, Italy*

<sup>d</sup>*Faculty of Health Medicine and Life Sciences, Department of Psychiatry and Neuropsychology School for Mental Health and Neuroscience Alzheimer Centre Limburg Maastricht University Medical Centre Maastricht the Netherlands*

<sup>e</sup>*Department of Neurology with Neurosurgical Activity “Carlo Poma” Hospital, ASST di Mantova, Mantua, Italy*

<sup>f</sup>*Cognitive Neurology Centre, Lariboisière-Fernand Widal Hospital GHU AP-HP Nord, Paris, France*

<sup>g</sup>*Department of Primary Care, Geriatric Service-Cognitive Disorders and Dementia, Local Health Authority of Modena (AUSL), Modena, Italy*

<sup>h</sup>*FERB Alzheimer Centre, Gazzaniga, Italy*

<sup>i</sup>*Memory Centre, Division of Geriatrics and Rehabilitation, University Hospitals of Geneva and University of Geneva, Geneva, Switzerland*

<sup>j</sup>*Department of Geriatric Psychiatry, Central Institute of Mental Health, Medical Faculty Mannheim, Heidelberg University, Mannheim, Germany*

<sup>k</sup>*Mediolanum Cardio Research, Milano, Italy*

<sup>l</sup>*Department of Medicine and Surgery, Institute of Gerontology and Geriatrics, University of Perugia, Italy*

<sup>m</sup>*Division of Clinical Geriatrics; NVS Department, Karolinska Institutet Stockholm, Sweden*

<sup>n</sup>*Neurological Unit (PM), U.V.A. Centre, Humanitas Gavazzeni, Bergamo, Italy*

<sup>o</sup>*Department of Psychiatry, Charité-Universitätsmedizin Berlin, Campus Benjamin Franklin, Berlin, Germany*

<sup>p</sup>*Greek Association of Alzheimer's Disease and Related Disorders (GAADR), Thessaloniki, Makedonia, Hellas. 1st Department of Neurology, School of Medicine, Aristotle University of Thessaloniki (AUTH), Macedonia, Hellas*

Accepted 12 September 2023

Pre-press 1 November 2023

---

\*Correspondence to: Aline Mendes, Division of Geriatrics and Rehabilitation, Chemin du Pont-Bochet 3, 1226 Thônex, Switzerland. Tel.: +41 223056111; E-mail: aline.mendes@hcuge.ch.

**Abstract.**

**Background:** Behavioral and psychological symptoms of dementia (BPSD) bring complexity in the clinical management of people with dementia; therefore, it is important to evaluate different models of care, such as Special Care Units (SCU-B).

**Objective:** To evaluate the SCU-B effectiveness toward alleviating BPSD and improving the quality of life (QoL) of patients and their caregivers.

**Methods:** ReCAGE was a multicenter, controlled, longitudinal study where 508 patients with BPSD were enrolled in two cohorts: 262 patients from centers endowed with a SCU-B, and 246 from centers without SCU-B. Statistical analyses included factorial ANCOVA for comparison among centers. The primary endpoint was effectiveness of the SCU-B, measured through the Neuropsychiatric Inventory (NPI) changes. Secondary endpoints were change in QoL of patients and caregivers, and the tertiary endpoint was time to nursing home admission.

**Results:** The NPI scores decreased in both arms, with a statistically significant difference from baseline to 36 months ( $p < 0.0001$ ) in both cohorts. Over time, NPI decreased more steeply during the first year in the SCU-B arm, but in the following two years the slope was clearly in favor of the control arm. This different pattern of the two cohorts reached statistical significance at the interaction “cohort by time” ( $p < 0.0001$ ). Conflicting results were found regarding the outcomes of quality of life, while there were no differences in time to institutionalization in both cohorts.

**Conclusions:** The ReCage study did not confirm the long-term superiority of the pathway comprising a SCU-B. A *post-hoc* analysis revealed data supporting their acute effectiveness during behavioral crises.

Keywords: Alzheimer’s disease, behavioral and psychological symptoms of dementia, caregiver burden, dementia, Neuropsychiatric Inventory, special care unit

## INTRODUCTION

Currently, the number of people with dementia is 55 million around the world, with estimates predicting a striking increase to 139 million by 2050, showing the importance of adapted health models to meet the needs of patients and caregivers [1]. If dementia is characterized by cognitive impairment with significant repercussions on the performance of daily activities, the concomitant presence of non-cognitive symptoms is also a constant feature in the trajectory of persons with dementia (PwD) [2].

The behavioral and psychological symptoms of dementia (BPSD) represent a heterogeneous group of non-cognitive symptoms occurring at some point in the disease trajectory in almost all PwD. They constitute a major component of the dementia syndrome irrespective of the underlying etiology and are not less clinically relevant than cognitive symptoms, as they strongly correlate with the degree of functional impairment, caregiver burden, inappropriate hospital admissions, and institutionalization [3–5]. Worldwide guidelines of good clinical practices recommend a comprehensive assessment of BPSD followed by multicomponent non-pharmacological approaches to be the first line of treatment, but there is lacking evidence of the exact standards of implementation among different settings [6].

However, the use of different classes of drugs such as anticholinesterase inhibitors, memantine, antidepressants, sedatives, and antipsychotics is frequent in the management of BPSD [7]. Up to 60% of persons with moderate or severe dementia PwD receive antipsychotics, although such medications are associated with a poor prognosis, increasing mortality, the risk of accelerated cognitive deterioration, and of stroke [8–10]. This panorama reveals the great challenge that is the management of BPSD in clinical practice [11].

To prevent these undesirable outcomes, Special Care Units (SCU-B) have in some countries been created in both acute and rehabilitation settings, with other special units integrating long-term care and nursing homes [12]. Despite the lack of a common definition of their standards, in this study we defined such units as those present in hospital wards where patients with BPSD are temporarily admitted when their behavioral symptoms are not amenable to being treated at home. The specific role of the SCU-B is to mitigate the challenging symptoms and to allow patients to get back home whenever possible [13].

Our group performed a scoping review of the literature that identified 33 studies that dealt with SCU-B (Pecoraro et al., unpublished). Nine studies provided only descriptive information about the SCU-B structure and organization. Among the studies that

evaluated the impact of SCU-B on patient-centered outcomes, only one was a randomized study and 23 were uncontrolled case series. As to clinical effectiveness, the only randomized clinical trial did not show statistically significant differences versus standard care as regards the primary outcomes, but patients and families were more satisfied [14]. From the uncontrolled studies there is some evidence that a short stay in SCU-Bs can improve BPSD, at least temporarily, and allow return home in approximately 50% of cases. As regards the possible long-term effectiveness, evidence is limited to short-term (1-year) and/or limited cost or health outcomes. In summary, the literature about SCU-B is scant, and the retrieved studies do not allow us to draw confident conclusions about the effectiveness of SCU-B on patient-centered outcomes in the long term.

We hypothesized that patients from regions where SCU-B is available had favorable short- and long-term outcomes in BPSD mitigation than those where such units did not exist. In this context, the main objective of the RECAGE (REspectful Caring for agitated Elderly) study is to evaluate the effectiveness of BPSD management of PwD in centers with and without SCU-B.

## METHODS

### *Design, setting, and population*

RECAGE was a multicenter, prospective observational study composed of two cohorts of patients with BPSD recruited in centers endowed with SCU-B and in centers lacking this facility. Of the 11 centers that participated in the study, five (Italy, Germany, France, Switzerland, and Norway) and six centers (Italy, Greece, France, and Germany), respectively, composed the two cohorts where there SCU-B were available or not.

Inclusion criteria comprised patients of any age, with a diagnosis of dementia of any etiology according to the DSM-IV, with a Mini-Mental State Examination (MMSE) less than or equal to 24 [15]. In addition, patients presented BPSD, with a Neuropsychiatric Inventory (NPI) score greater than or equal to 32 [16]. The presence of a family member or patient's caregiver during the study committed to the proposed follow-up was mandatory. Exclusion criteria comprised: 1) presence of uncontrolled physical diseases potentially contributing to the cognitive decline and BPSD, 2) concomitant psychiatric

disorders or chronic alcoholism, and 3) concomitant diseases severe enough to reduce life expectancy.

The sample size was calculated on the comparison between the two cohorts only at the final time point. So, a difference given by an effect size of about 0.25 at a Student's *t* test for unpaired data with a power of 0.80 and a significance level of 0.05 (two-sided) could be demonstrated by a sample size of 250 patients in each cohort.

All centers had memory clinics that were the entry point for enrollment in the study. It is important to emphasize that hospitalization in an SCU-B was not mandatory for participation in the study in the centers where it was available. The centers endowed with a SCU-B facility contributed with 266 patients, while the centers without a SCU-B, contributed with 252 patients, for a total of 518 included patients. Ten patients did not perform any assessment after baseline and were therefore excluded from the modified intention to treat population (Fig. 1). The study was conducted according to the ethical principles of the Declaration of Helsinki. All patients, representatives, and caregivers received oral and written information about the study and provided written consent to participate. This study was approved by the local research ethics committees (Comitato Etico di Bergamo, REG.SPERIM. 25/18, 90.02.2018).

### *Study procedures*

Detailed information on the different procedures and the study protocol was the subject of a previous publication [17]. In brief, from the inclusion visit and during the 3-year follow-up of the study, patients and caregivers were extensively assessed by the project team, allowing the collection of clinical data from different domains. The dataset included demographic information, vital signs, active comorbidities and relevant medical history, dementia characteristics (age at diagnosis, etiology, severity of cognitive impairment), clinical examination, functional status, pharmacological treatment, BPSD assessment as well as the quality-of-life parameters of the patient and the caregiver. In addition, we also collected information on the medical-economic dimension, whose in-depth analysis will be the subject of another publication.

### *Follow-up*

The patients and their caregivers attended follow-up visits scheduled every 6 months. This follow-up




Fig. 1. Flowchart of enrollment and follow-up. SCU-B, special medical care unit for people with behavioral and psychological symptoms of dementia.

duration was chosen to encompass the different short- and long-term outcomes defined in this study. Unscheduled visits were organized if requested by the patient or the caregiver.

### Study outcomes

The primary outcome referred to the effectiveness in the clinical management of BPSD in centers where SCU-B were available. Such effectiveness was measured through the changes in the total NPI scores from the first visit to the end of the study in both cohorts.

The patient's and caregiver's quality of life throughout the follow-up was established as a secondary outcome. We applied different clinical scales that concern the patient such as the Quality of Life in Alzheimer's Disease questionnaire (QoL-AD; self-rated and proxy-rated) [18] and caregivers such as Adult Carer Quality of life questionnaire (AC-QoL) [19], the Caregiver Burden Inventory (CBI) [20], and the Dementia Attitude Scale (DAS) [21]. It is important to emphasize that in chronic diseases of

a progressive nature, as in the case of dementia, the measure of quality of life and caregiver burden will tend to worsen with evolution, even if appropriate interventions are implemented. Although this secondary outcome is not likely to improve in the long term, the differences in the trajectory of the two cohorts were of interest for this analysis. Moreover, the delay in admission to a nursing home was defined as a tertiary outcome.

Several safety-related outcomes were documented throughout the study. They included intercurrent adverse events, such as admission to an emergency room, hospitalization, and mortality. Other serious events anticipated were falls, injury due to falls, accelerated cognitive decline, accelerated functional decline, and parkinsonism. As the COVID-19 pandemic happened during the follow-up of patients and caregivers, the potential repercussions in the study results were also considered in the analysis of adverse events. We highlight the fact that one of the Italian centers with SCU-B, Gazzaniga, was in the Italian region most severely affected by the pandemic.

### Statistical analysis

Descriptive statistics (mean with standard deviation, minimum and maximum, and 95% confidence interval for the variables with assumed Gaussian distribution; median with the first and third quartile, and interquartile range for the corresponding variables assumed not Gaussian distributed) have been calculated for continuous variables. For categorical variables, patient counts and percentages are provided. The baseline characteristics of the cohorts have been compared for qualitative variables by means of the chi-squared test. Student's *t*-test for unpaired data or its alternative non-parametric Wilcoxon rank sum test has been used for quantitative variables.

To compare the primary outcome, changes over time between the two cohorts and the propensity score was calculated by means of the SAS PROC PSMATCH with the cohort (SCU-B or non SCU-B) as the dependent variable and all the baseline patients' characteristics as the independent variables. Then, the pattern of the different scales over the time has been compared between the two cohorts by means of repeated measures of the ANCOVA model with the propensity score as a covariate. Particularly, the considered models have "Cohorts" at two levels (SCU-B and non-SCU-B) as a fixed "between subjects" factor, "Time" at seven levels (baseline or V0, V6, V12, V18, V24, V30, and V36) as a fixed factor "within subjects", and their interaction "Cohort by time". In addition, "subjects" is a random factor nested into the fixed factor "Cohorts".

The *post hoc* multiple comparison analysis has been carried out by adjusting the significant level according to Šidák [22]. These analyses have been carried out by means of SAS PROC MIXED according to several models (general linear model, heterogeneous general linear model, random coefficients linear model in order to consider the trend for each subject, and heterogeneous random coefficients linear model in considering the trend for each subject and a different pattern of the variance-covariance matrix of the two cohorts). Among the several considered models and patterns of variance-covariance matrices the heterogeneous general linear model with the unstructured matrix pattern has been selected according to a lower value of the Akaike's information criterion for the NPI analysis [23]. Pragmatically, a similar approach was used for the secondary outcomes with the total scores from the QoL-AD self-rated and proxy-rated, AC-QoL, CBI, and DAS scales. SAS PROC MIXED allows to deal with the

missing data under the assumption that they are at least "missing at random".

Time to admission to a nursing home has been analyzed by means of the Kaplan-Meier estimate of the cumulative probability of placement to a nursing home and compared between the two cohorts by means of the log-rank test. Also, Cox's proportional hazard regression model to predict time to institutionalization was carried out in order to estimate the hazard ratio between the two cohorts. The statistical significance has been put at  $p=0.05$ . All statistical analyses were performed using the SAS® (Statistical Analysis System) version 9.4.

## RESULTS

### *Characteristics of the study population*


Overall, in both cohorts, BPSD severity significantly decreased at follow-up (Fig. 2). However, this decrease was more significant in the cohort without specialized units compared to centers with SCU-B (Fig. 3). These results contradict our initial hypothesis.

Of a total of 508 PwD included in the study, 54.9% were female, with a mean age of  $78 \pm 7.93$  years, and  $8.93 \pm 4.53$  years of formal education. Caregivers were mainly the spouse or the child, 70.2% were female, with a mean age of  $61.89 \pm 12.75$  years. The most frequent etiology of dementia was Alzheimer's disease (58.4%), followed by dementia with multiple etiologies (16.4%). The mean MMSE was  $15.45 \pm 6.25$  points at baseline. Multimorbidity was frequent in this population (90.6%), with hypertension (52.2%), dyslipidemia (17.5%), diabetes (17.1%), and depression (11.4%) being the most prevalent comorbidities documented. The mean NPI score was of 52.44 points, with the domains of apathy/indifference (85.4%), agitation/aggression (83.1%), depression/dysphoria (81.7%), and irritability/lability (80.9%) being the BPSD most frequently described. The AC-QoL scores of  $71.28 \pm 18.09$  indicated a mid-range quality of life of caregivers at baseline. Furthermore, a high burden of caregivers was present with 47.4% presenting a CBI >36 points.

When comparing the characteristics of the centers with and without SCU-B, we highlight a lower proportion of women in those with SCU-B (50.4% versus 59.8%;  $p=0.0338$ ). Furthermore, caregivers from centers with SCU-B had higher quality of life in baseline according to the AC-QoL ( $73.94 \pm 18.44$  versus  $68.49 \pm 17.31$ ;  $p=0.0007$ ). Similarly, there was a

**A. Centers with SCU-B**

p<0.0001



**B. Centers without SCU-B**

p<0.0001




Fig. 2. Mean values of NPI domains at baseline and at the end of the study.

higher burden of caregivers in centers without SCU-B ( $32.99 \pm 17.88$  versus  $38.07 \pm 16.91$ ;  $p = 0.0006$ ). Table 1 summarizes the different variables evaluated in both cohorts that finally showed a similar profile in the other parameters of the study.

*Effectiveness*

The Least Squares Means estimates of the total NPI score decreased in both cohorts according to a different pattern (interaction “cohort by time”




Fig. 3. Least square means of the Neuropsychiatric Inventory in baseline and during follow-up visits in centers with and without SCU-B. *p* value corresponds to the interaction “cohort by time”.

was statistically significant:  $p < 0.0001$ ) (Fig. 3). In the centers with SCU-B, the total NPI score decreased from 52.68 (baseline) by 16.21% (at 6 months), 22.21% (at 12 months), 24.0% (at 18 months), 23.95% (at 24 months), 20.73% (at 30 months), and finally, by 19.51% to a score of 42.30 at the 36-month visit. In the centers without SCU-B, the NPI decreased from a baseline of 52.12 by 9.27% (at 6 months), 16.82% (at 12 months), 24.12% (at 18 months), 34.69% (at 24 months), 40.81% (at 30 months), and finally by 41.40% to a score of 30.54 at the 36-month visit. In both cohorts the difference of NPI scores between baseline to the visit performed at the end of study was statistically significant, showing decreased levels of BPSD ( $p < 0.0001$ ) (Fig. 2).

This difference favored the centers without SCU-B, which presented an increased pattern of

Table 1  
Baseline characteristics of patients and caregivers included in the study

Characteristics	Centers with SCU-B N = 262	Centers without SCU-B N = 246	<i>p</i>	Total N = 508
<b>Demographics</b>				
Age, y	78.14 (8.42)	78.04 (7.40)	0.8909	78.09 (7.93)
Female sex	132 (50.4%)	147 (59.8%)	0.0338	279 (54.9%)
Educational level, y	9.11 (4.41)	8.73 (4.65)	0.1737	8.93 (4.53)
Primary caregiver			0.8627	
Spouse or child	245 (93.8%)	230 (93.5%)		475 (93.7%)
Other	17 (6.2%)	16 (6.5%)		33 (6.3%)
Caregiver age, y	62.85 (12.55)	60.87 (12.90)	0.0797	61.89 (12.75)
Caregiver Female sex	185 (70.9%)	171 (69.5%)	0.7362	356 (70.2%)
Caregiver educational level, y	11.87 (4.06)	12.69 (3.89)	0.0269	12.24 (4.00)
<b>Vital signs<sup>a</sup></b>				
Height, cm	166.62 (8.88)	163.14 (9.89)		164.97 (9.52)
Weight, kg	69.78 (12.81)	70.62 (16.50)		70.17 (14.64)
BMI, kg/m <sup>2</sup>	25.10 (3.56)	26.39 (4.97)		25.71 (4.33)
Systolic blood pressure, mmHg	126.77 (15.70)	130.79 (17.73)		128.75 (16.83)
Diastolic blood pressure, mmHg	77.06 (9.56)	76.92 (10.14)		76.99 (9.84)
Heart rate, pulse/min	72.65 (11.04)	70.50 (11.65)		71.63 (11.37)
<b>Disease characteristics</b>				
Age at diagnosis	75.35 (8.80)	75.15 (7.79)	0.7646	75.25 (8.32)
Etiology <sup>b</sup>			0.0828	
Alzheimer's disease	162 (62.1%)	134 (54.5%)		296 (58.4%)
Vascular dementia	28 (10.7%)	3 (1.2%)		31 (6.1%)
Lewy body dementia	13 (5.0%)	10 (4.1%)		23 (4.5%)
Parkinson's disease	1 (0.4%)	5 (2.0%)		6 (1.2%)
Frontotemporal dementia	19 (7.3%)	19 (7.7%)		38 (7.5%)
Multiple etiologies	34 (13.0%)	49 (19.9%)		83 (16.4%)
Not specified	5 (1.5%)	26 (10.6%)		31 (5.9%)
<b>Cognitive impairment</b>				
Memory	252 (96.2%)	240 (97.6%)	0.1357	492 (96.9%)
Executive functioning	211 (80.5%)	200 (81.3%)	0.0553	411 (80.9%)
Language	105 (40.1%)	114 (46.3%)	0.0939	219 (43.1%)
Visuospatial	136 (51.9%)	127 (51.6%)	0.3436	263 (51.8%)
Frontal	111 (42.4%)	131 (53.3%)	0.0007	242 (47.6%)
Apraxia	103 (39.3%)	125 (50.8%)	0.0014	228 (44.9%)

(Continued)



Table 1  
(Continued)

Characteristics	Centers with SCU-B N = 262	Centers without SCU-B N = 246	<i>p</i>	Total N = 508
<b>Patient status</b>				
MMSE (0–30)	15.88 (6.08)	14.99 (6.41)	0.1102	15.45 (6.25)
ADCS-ADL (0–78)	34.59 (18.47)	35.93 (17.65)	0.4058	35.24 (18.07)
Basic activities	14.77 (5.94)	13.72 (5.72)	0.0434	14.26 (5.85)
Instrumental activities	19.82 (13.61)	22.21 (12.85)	0.0425	20.98 (13.29)
NPI total score (0–144) <sup>c</sup>	52.87 (16.45)	51.98 (21.27)	0.5963	52.44 (18.92)
Delusions	178 (67.9%)	145 (58.9%)	0.0352	323 (63.6%)
Hallucinations	100 (38.2%)	80 (32.5%)	0.1835	180 (35.4%)
Agitation / aggression	225 (85.9%)	197 (80.1%)	0.0817	422 (83.1%)
Depression / dysphoria	221 (84.4%)	194 (78.9%)	0.1099	415 (81.7%)
Anxiety	211 (80.5%)	193 (78.5%)	0.5617	404 (79.5%)
Elation / euphoria	52 (19.8%)	56 (22.8%)	0.4220	108 (21.3%)
Apathy / indifference	226 (86.3%)	208 (84.6%)	0.5858	434 (85.4%)
Disinhibition	101 (38.5%)	142 (57.7%)	<0.0001	243 (47.8%)
Irritability / lability	220 (84.0%)	191 (77.6%)	0.0698	411 (80.9%)
Aberrant motor behavior	183 (69.8%)	162 (65.9%)	0.3352	345 (67.9%)
Sleep disturbances	165 (63.0%)	149 (60.6%)	0.5767	314 (61.8%)
Appetite and eating disorders	133 (50.8%)	163 (66.3%)	0.0004	296 (58.3%)
<b>Patient and caregiver quality of life</b>				
Patient QoL-AD; proxy-rated	28.33 (6.35)	28.52 (6.39)	0.7317	28.42 (6.36)
Patient QoL-AD; self-rated	34.67 (5.52)	34.24 (6.62)	0.4743	34.45 (6.09)
AC-QoL	73.94 (18.44)	68.49 (17.31)	0.0007	71.28 (18.09)
CBI			0.0004	
0–24	92 (35.5%)	51 (20.8%)		143 (28.4%)
25–36	63 (24.3%)	59 (24.1%)		122 (24.2%)
>36	104 (40.2%)	135 (55.1%)		239 (47.4%)
CBI	32.99 (17.88)	38.07 ± 16.91	0.0006	35.57 (17.67)
DAS	95.24 (15.80)	95.80 (16.91)	0.7064	95.52 (16.34)

SCU-B, special medical care unit for people with behavioral and psychological symptoms of dementia; BMI, body mass index; MMSE, Mini-Mental State Examination; ADCS-ADL, Alzheimer's Disease Cooperative Study Activities of Daily Life Scale; NPI, Neuropsychiatric Inventory; QoL-AD, Quality of Life-Alzheimer's Disease; AC-QoL, Adult Carer Quality of Life Questionnaire; CBI, Caregiver's Burden Inventory; DAS, Dementia Attitude Scale. <sup>a</sup>No statistical tests have been carried out, as differences can be statistically significant owing to the sample numbers, without clinical relevance. <sup>b</sup>*p* value refers to the comparison of Alzheimer's disease versus all other etiologies. <sup>c</sup>NPI sub-domains have been calculated only for patients with symptom. *p* values refer to the comparison of the proportion of patients with symptoms versus without symptoms.

Table 2  
Effectiveness end-point analysis as NPI least-square means at baseline and during follow-up in centers with and without SCU-B

NPI	Centers with SCU-B		Centers without SCU-B		<i>p</i>
	LSM (SE)	Δ	LSM (SE)	Δ	
Baseline	52.6823 (1.0698)		52.1232 (1.3804)		1
6 months	44.1381 (1.1554)	–8.544	47.2894 (1.4358)	–4.834	0.9999
12 months	40.9805 (1.1991)	–3.158	43.3579 (1.4551)	–3.932	1
18 months	40.0342 (1.4325)	–0.946	39.5505 (1.3916)	–3.807	1
24 months	40.0601 (1.5482)	0.026	34.0424 (1.1632)	–5.508	0.1981
30 months	41.7625 (1.7699)	1.702	30.8522 (1.3118)	–3.19	0.0001
36 months	42.4031 (1.8834)	0.641	30.5436 (1.5145)	–0.309	0.0001

NPI, Neuropsychiatric Inventory; SCU-B, Special medical care unit for people with behavioral and psychological symptoms of dementia; LSM, Least Square Means; SE, Standard Error. *p*-value is the adjusted significance value for the comparison between the two cohorts at each. Δ, Changes are calculated as the difference between the value at the visit and the previous value. A negative change indicates an improvement.

improvement of the NPI total score at the different assessment points, especially after the visit at the 18-month follow-up (Table 2).

#### Quality of life

Although there was no statistically significant difference between the two cohorts at each time point




Fig. 4. Least square means of the quality-of-life scales in baseline and during follow-up visits in centers with and without SCU-B. QoL-AD, Quality of Life-Alzheimer’s Disease; AC-QoL, Adult Carer Quality of Life Questionnaire; CBI, Caregiver’s Burden Inventory; DAS, Dementia Attitude Scale. *p* values correspond to the interaction “cohort by time”.

(Supplementary Material), we found a different qualitative pattern between the two cohorts in the AC-QoL scores: caregivers’ quality of life of increased in the non-SCU-B centers according to this questionnaire ( $p=0.0169$ ) (Fig. 4C). On the opposite, DAS total scores significantly improved in centers with SCU-B compared to centers without SCU-B at the end of the study ( $p=0.0203$ ) (Fig. 4E).

Owing to the statistical significance of the interaction “cohort by time”, we specifically considered the differences among the time points within each cohort for both questionnaires. Regarding the AC-QoL, we observed a statistically significant difference between month 24 and month 36 ( $p=0.0094$ ), as well as between month 30 and month 36 ( $p=0.0177$ ) of follow-up in centers with SCU-B, both in the direction of a decreased quality of life. No statistically significant difference between any two visits was found in centers without SCU-B for the AC-QoL. All Least squares mean values in both cohorts with their respective differences along follow-up are shown in detail in the Supplementary Material (Tables 1 and 2).

Regarding the DAS, the difference was statistically significant between baseline and month 12 ( $p=0.0032$ ), month 18 ( $p=0.0067$ ), month 24 ( $p<0.0001$ ), month 30 ( $p<0.0001$ ), and month 36

( $p=0.0010$ ) of follow-up, as well as between month 6 and month 30 ( $p=0.0319$ ). In the centers without SCU-B, the difference between two visits was never statistically significant except for the comparison between baseline and month 12 ( $p=0.0490$ ).

Neither the QoL-AD nor the CBI scores disclosed a statistically different qualitative/quantitative pattern between the two cohorts, therefore there was no statistically significant interaction (Fig. 4A, B, D). For this reason, we considered only a pooled analysis of both cohorts together. It revealed a statistically significant difference from baseline to the end of the study in the two versions of QoL-AD, patient-rated and proxy-rated ( $p<0.0001$ ) and in the CBI ( $p=0.0165$ ). These differences showed a decreased quality of life of PwD and an increased caregiver burden by the end of the study (Supplementary Material).

*Time to institutionalization*

The evolution of the rate of admission to a nursing home during the 36 months of follow-up is shown in Fig. 5. There was no significant difference between centers with and without SCU-B ( $p=0.3552$ ). The last probability values in the two curves were: SCU-B: 0.42 (95%CI 0.34; 0.50); non-SCU-B: 0.52 (95% CI 0.45; 0.59). The Cox’s proportional-hazard model



follow-up of the study. We analyzed a sample of 39 patients with available NPI and MMSE values at admission and at discharge from a SCU-B. They had a mean NPI and a mean MMSE of  $62.2 \pm 19.02$  and of  $14.9 \pm 7.86$ , respectively. NPI decreased to  $21.4 \pm 12.52$  at discharge, while MMSE slightly increased to  $15.08 \pm 7.55$ . The decrease between before and after (admission and hospital discharge) was statistically significant for NPI ( $p < 0.0001$ ), whilst the increase of the MMSE ( $p = 0.0803$ ) was not.

## DISCUSSION

When we elaborated the concept of this study, the initial hypothesis was that the care pathways of patients with BPSD in centers with a SCU-B (available for admission in case of behavioral crisis, but not mandatory) were superior to the pathways lacking this facility. This was based on an assumption of better quality of care in centers with SCU-B, translated as efficient mitigation of behavioral crisis in SCU-B, lower prescription of psychotropic drugs, increased delay to nursing home admission, as well as better confidence, knowledge, and support of caregivers. However, the findings of this study do not confirm the initial hypothesis.

As regards the primary outcome of the study, the results did not confirm the research hypothesis of the long-term clinical superiority of the SCU-B arm over the control one. In fact, while a gradual decrease of the score over time in both arms was observed, a trajectory expected from previous evidence [24], the slope was steeper in the SCU-B cohort only until 6 months, but afterward the score continued to decrease only in the control arm, reaching statistical significance at 30 and 36 months of follow-up in favor of the latter. Therefore, in the long term, the outcome improved more in those patients whose behavioral crises were managed without the possibility of being admitted to a SCU-B. Our results do not provide any likely explanation for this (unexpected) difference in favor of the non-SCU-B cohort during the second part of the follow-up, despite several exploratory analyses performed to identify potential confounding/explanatory factors. We carried out several sensitivity analyses by means of ANCOVA, restricting the population to patients with a baseline MMSE within 10 and 20 and including only the first 10 items of NPI. Furthermore, the analysis was repeated excluding centers with the lowest number of enrolled patients; on the patients

with all seven visits performed; on the patients with at least the 30-month visit performed; and finally, on the patients with at least the 24-month visit performed. The analyses did not substantially change the results. Other hypotheses could not be tested on the basis of our data: we do not know in depth how the acute crises of patients with BPSD were managed in centers without SCU-B and possibly there were differences not measured by the study variables and responsible, at least in part, for the observed finding. Moreover, differences in primary health care including day care centers, availability of home care, and costs of such interventions could significantly impact on our primary outcome.

Another point to underline is that a much higher number of admissions to SCU-B was expected during the follow-up by the clinicians adhering to the REcage Consortium. The relatively small number of admissions over three years (one admission of 45 pts/508, 56 total number of admissions) was possibly due, at least to some extent, to the pandemic. This means that only 45 pts were exposed, to the SCU-B.

As for the secondary endpoints, a similar result in favor of the centers without SCU-B cohort was found for the quality of life of caregivers (AC-QoL), whereas the caregivers' attitude toward dementia (DAS) improved in the SCU-B cohort, especially concerning the knowledge factor.

Finally, no significant difference between the cohorts was found on time of institutionalization.

To sum up, we must emphasize that REcage was carried out in the European context, where large differences exist as regards the clinical management of patients with BPSD across the participating countries. Outpatient care in centers is probably rather similar across countries, although not superposable, due to the context they operate in, whereas the diffusion of SCU-Bs is very unequal: for instance, Italy does not have a developed network of SCU-B, but only a few, whereas the outpatient clinics for PwD are widespread. In Germany, the treatment of BPSD in severe dementia is provided by the fully sectorized care of the psychiatric hospitals in collaboration with geriatric internal medicine specialists. In Norway, psychiatric wards in all hospital trusts do admit people with dementia, but in their acute ward. Innlandet Hospital trust (Ottestad), which participated in the present study, is the only hospital trust with a SCU-B ward. In France there have for many years been a well-developed network of SCU-Bs called "Unités Cognitivo-Comportementales" and are widespread all over the country. In Switzerland the situation is

variable from a canton to another. In Greece, there are no SCU-B.

Regarding safety-related outcomes, we would like to discuss our findings on psychotropic medication use. Although the prescription was significantly lower in centers with SCU-B, in general, we noticed a high prevalence of psychotropic medications in both cohorts. Although psychotropic medication is not the first-line treatment recommendation, they remain widely prescribed. Several studies showed that a high rate of patients with moderate-severe dementia were under at least one psychotropic medication, including antipsychotics [25]. The main factors associated with the use of pharmacological interventions include persistent neuropsychiatric symptoms, living in nursing homes and prior psychiatric diagnosis [26].

Regarding special units for people with dementia, previous studies showed mixed results, with a few reports of increased psychotropic medication use in people admitted to a SCU-B [27–29]. One explanation is the selection of patients admitted to those units based on the severity of symptoms. They often present criteria of acute crisis that could not be mitigated outside the hospital or symptoms refractory to previous non-pharmacological interventions implemented by the team. Therefore, in certain circumstances, despite the limited efficacy, clinical management in specialized units may culminate in an increase (or absence of decrease) in drug prescription.

#### *Limitations of the study*

The outbreak of the COVID-19 pandemic, six months after the end of the recruitment period, created many difficulties, especially hindering regular “in presence” visits and compelling the memory clinics to do only or mostly phone visits (NPI is not validated for phone visits), with the obvious consequent loss of data, easing withdrawals from the study, and delayed visits. Moreover, it is important to remark that COVID-19 was specially devastating in Italy, where two SCU-Bs (Gazzaniga and Modena) with a high recruitment rate (150/262) are located, with ensuing mortality (total of 15 COVID-19 related deaths with 14 in SCU-B and only 1 in non-SCU-B arm). Finally, the pandemic may have had other indirect impacts on the study, such as social isolation due to the repeated and prolonged lockdowns (whose effects on cognition and behavior are well known).

The possible COVID-19 impact was evaluated to verify whether the administration of the scales over the phone had any methodological influence and

whether there was a country effect due to the different (spatial and temporal) incidence of the pandemic. NPI analysis excluding assessments performed as phone visits and NPI analysis by comparing pattern of scores between non-SCU-B and SCU-B located in Italy versus non-SCU-B and SCU-B in the other countries were carried out. The difference between the SCU-B and non-SCU-B arm persisted after excluding the Gazzaniga and Modena units, the most COVID-stricken centers (and their 150/262 pts). However, all the above analyses confirmed a statistically significant interaction “time by cohorts” in favor of the non-SCU-B.

A variety of factors other than COVID-19 interfered with the trial and possibly led to biased results. Among them is observed discrepancies at baseline, as consequence of the non-randomized study design, which may be relevant despite the use of the propensity score analysis. Most important, the social context of the trial was different across the countries participating in the RECaGe Consortium.

Many of these units could not carry out usual admissions according to predefined criteria, while other units had to absorb patients with COVID-19 on a temporary basis. In addition, many of the outpatient interventions, such as consultations and therapies, were also suspended or modified to meet the isolation measures imposed in each region and country. Therefore, potential biases regarding the comparison of centers with and without SCU-B were introduced. Supplementary measurements that could help solve this problem would be to accurately incorporate the different interventions and services available at each center, as well as which ones were implemented for each person with dementia and caregiver during the entire follow-up of the study. Such measures would have helped in a deeper understanding of the factors that influenced our results, including during the pandemic period.

#### *Implications for further research*

Due to the many limitations of the trial, its negative results must be interpreted with caution.

In fact, since literature shows evidence for short-term effect of the SCU-B, albeit mainly from non-controlled studies as well as from our *post-hoc* analysis of the patients admitted to the unit, the RECaGe results do not invalidate the “acute” clinical effectiveness of the admission to the SCU-B to solve behavioral crises [30–32]. Therefore we think that, where SCU-Bs are available, they should

be supported, at least unless there is an available operational alternative [33]. Despite the potential effectiveness of mitigating behavioral crises, the overall evidence from previous studies shows mixed results, which was also observed in special care units specifically located in long-term care facilities [12].

As to future research, the lack of long-term results puts the discussion of ethical approval for randomization on the table. A randomized clinical trial could be ethically justified, comparing SCU-B admission with another planned alternative, for instance mobile BPSD teams. If studied in a multicenter setting, the participant centers should be stratified for social context of the country or region to minimize biases.

Finally, the definition of a standardized panel of outcomes and measurements to guide future studies is essential, taking into consideration meaningful outcomes for patients, families, staff, and stakeholders.

## ACKNOWLEDGMENTS

The authors thank all the members of the RECage study group who, throughout the study, contributed to the accomplishment of this work. Also, we wish to thank all the patients, family members and caregivers taking part in this study.

## FUNDING

This project has received funding from the European Union's Horizon 2020 Research and Innovation program under grant agreement No 779237.

## CONFLICT OF INTEREST

The authors have no conflict of interest to report.

## DATA AVAILABILITY

The data supporting the findings of this study are available on request from the corresponding author. The data are not publicly available due to privacy and ethical restrictions.

## SUPPLEMENTARY MATERIAL

The supplementary material is available in the electronic version of this article: <https://dx.doi.org/10.3233/JAD-230708>.

## REFERENCES

- [1] (2022) World Alzheimer Report 2022: Life after diagnosis: Navigating treatment, care and support.
- [2] Rabins PV, Schwartz S, Black BS, Corcoran C, Fauth E, Mielke M, Christensen J, Lyketsos C, Tschanz J (2013) Predictors of progression to severe Alzheimer's disease in an incidence sample. *Alzheimers Dement* **9**, 204-207.
- [3] Gitlin LN, Kales HC, Lyketsos CG (2012) Nonpharmacologic management of behavioral symptoms in dementia. *JAMA* **308**, 2020-2029.
- [4] Herrmann N, Lanctôt KL, Sambrook R, Lesnikova N, Hébert R, McCracken P, Robillard A, Nguyen E (2006) The contribution of neuropsychiatric symptoms to the cost of dementia care. *Int J Geriatr Psychiatry* **21**, 972-976.
- [5] Piercy KW, Fauth EB, Norton MC, Pfister R, Corcoran CD, Rabins PV, Lyketsos C, Tschanz JT (2013) Predictors of dementia caregiver depressive symptoms in a population: The Cache County dementia progression study. *J Gerontol B Psychol Sci Soc Sci* **68**, 921-926.
- [6] Reus VI, Fochtmann LJ, Eyster AE, Hilty DM, Horvitz-Lennon M, Jibson MD, Lopez OL, Mahoney J, Pasic J, Tan ZS, Wills CD, Rhoads R, Yager J (2016) The American Psychiatric Association Practice Guideline on the use of antipsychotics to treat agitation or psychosis in patients with dementia. *Am J Psychiatry* **173**, 543-546.
- [7] Ijaopo EO (2017) Dementia-related agitation: A review of non-pharmacological interventions and analysis of risks and benefits of pharmacotherapy. *Transl Psychiatry* **7**, e1250.
- [8] Steinberg M, Lyketsos CG (2012) Atypical antipsychotic use in patients with dementia: Managing safety concerns. *Am J Psychiatry* **169**, 900-906.
- [9] Russ TC, Batty GD, Starr JM (2012) Cognitive and behavioural predictors of survival in Alzheimer disease: Results from a sample of treated patients in a tertiary-referral memory clinic. *Int J Geriatr Psychiatry* **27**, 844-853.
- [10] Ma H, Huang Y, Cong Z, Wang Y, Jiang W, Gao S, Zhu G (2014) The efficacy and safety of atypical antipsychotics for the treatment of dementia: A meta-analysis of randomized placebo-controlled trials. *J Alzheimers Dis* **42**, 915-937.
- [11] Backhouse T, Camino J, Mioshi E (2018) What do we know about behavioral crises in dementia? A systematic review. *J Alzheimers Dis* **62**, 99-113.
- [12] Orth J, Flowers D, Betz G, Cagle JG (2023) A systematic review of specialty dementia care units in long-term care settings. *Int J Geriatr Psychiatry* **38**, e5907.
- [13] Guazzarini AG, Casanova G, Buchholz F, Kozori M, Lavalpe S, Lichtwarck B, Margioli E, Mendes A, Montandon M-L, Murasecco I, Myhre J, Poptsi E, Reda V, Ulshöfer DE, Fascendini S (2022) The Special Care Unit for People with Behavioral and Psychological Symptoms of Dementia (SCU-B) in the context of the project "RECage-Respectful Caring for Agitated Elderly": A qualitative study. *Int J Environ Res Public Health* **19**, 16913.
- [14] Goldberg SE, Bradshaw LE, Kearney FC, Russell C, Whitmore KH, Foster PER, Mamza J, Gladman JRF, Jones RG, Lewis SA, Porock D, Harwood RH, Medical Crises in Older People Study Group (2013) Care in specialist medical and mental health unit compared with standard care for older people with cognitive impairment admitted to general hospital: Randomised controlled trial (NIHR TEAM trial). *BMJ* **347**, f4132.
- [15] Folstein MF, Folstein SE, McHugh PR (1975) "Mini-mental state". A practical method for grading the cognitive state of patients for the clinician. *J Psychiatr Res* **12**, 189-198.

- [16] Cummings JL (1997) The Neuropsychiatric Inventory: Assessing psychopathology in dementia patients. *Neurology* **48**, S10-S16.
- [17] Poptsi E, Tsolaki M, Bergh S, Cesana BM, Ciccone A, Fabbo A, Frisoni GB, Frölich L, Lavolpe S, Guazzarini AG, Hugon J, Fascendini S, Defanti CA (2021) Rationale, design, and methodology of a prospective cohort study for coping with behavioral and psychological symptoms of dementia: The RECage Project. *J Alzheimers Dis* **80**, 1613-1627.
- [18] Torisson G, Stavenow L, Minthon L, Londos E (2016) Reliability, validity and clinical correlates of the Quality of Life in Alzheimer's disease (QoL-AD) scale in medical inpatients. *Health Qual Life Outcomes* **14**, 90.
- [19] Elwick H, Joseph S, Becker S, Becker F (2010) *Manual for the Adult Carer Quality of Life Questionnaire (AC-QoL)*. The Princess Royal Trust for Carers, London.
- [20] Novak M, Guest C (1989) Application of a multidimensional caregiver burden inventory. *Gerontologist* **29**, 798-803.
- [21] O'Connor ML, McFadden SH (2010) Development and psychometric validation of the Dementia Attitudes Scale. *In J Alzheimers Dis* **2010**, e454218.
- [22] Šidák Z (1967) Rectangular confidence regions for the means of multivariate normal distributions. *J Am Stat Assoc* **62**, 626-633.
- [23] Akaike H (1974) A new look at the statistical model identification. *IEEE Trans Automat Contr* **19**, 716-723.
- [24] Aalten P, de Vugt ME, Jaspers N, Jolles J, Verhey FRJ (2005) The course of neuropsychiatric symptoms in dementia. Part II: Relationships among behavioural sub-syndromes and the influence of clinical variables. *Int J Geriatr Psychiatry* **20**, 531-536.
- [25] Selbæk G, Kirkevold Ø, Engedal K (2008) The course of psychiatric and behavioral symptoms and the use of psychotropic medication in patients with dementia in Norwegian nursing homes—a 12-month follow-up study. *Am J Geriatr Psychiatry* **16**, 528-536.
- [26] Nørgaard A, Jensen-Dahm C, Gasse C, Hansen ES, Walde-mar G (2017) Psychotropic polypharmacy in patients with dementia: Prevalence and predictors. *J Alzheimers Dis* **56**, 707-716.
- [27] Koskas P, Bourdonnec L, Kohler S, Romdhani M, Drunat O (2021) Factors influencing length of hospital stay in an acute psychogeriatric unit. *Psychogeriatrics* **21**, 478-482.
- [28] Djekovic K, Clowes K, Spalding N, Waite L (2022) A retrospective study of the behavioural and psychological symptoms of dementia in patients admitted to a Specialist Dementia Care Unit. *Australas J Ageing* **41**, e397-e405.
- [29] Ghianda D, Cornali C, Franzoni, S, Di Fazio I, Rozzini L, Padovani A, Trabucchi M (2005) L'Istituto di Riabilitazione Geriatrica (IDRG): Un modello clinico-assistenziale per le demenze? *G Gerontol* **53**, 87-96.
- [30] Koskas P, Feugeas MCH, Saad S, Belqadi S, Daraux J, Drunat O (2011) Management of behavioral and psychological symptoms of dementia in a dedicated psychogeriatric unit: A pilot experience. *Alzheimer Dis Assoc Disord* **25**, 184-186.
- [31] Pongan E, Dorey J-M, Krolak-Salmon P, Federico D, Sellier C, Auguste N, Fabre F, Laurent B, Trombert-Paviot B, Rouch I (2017) Predictors of discharge destinations and three-month evolution of patients initially hospitalized in a cognitive behavioral unit. *J Alzheimers Dis* **60**, 1259-1266.
- [32] Soto ME, Andrieu S, Villars H, Secher M, Gardette V, Coley N, Nourhashemi F, Vellas B (2012) Improving care of older adults with dementia: Description of 6299 hospitalizations over 11 years in a special acute care unit. *J Am Med Dir Assoc* **13**, 486.e1-6.
- [33] Tay FHE, Thompson CL, Nieh CM, Nieh CC, Koh HM, Tan JJC, Yap PLK (2018) Person-centered care for older people with dementia in the acute hospital. *Alzheimers Dement (N Y)* **4**, 19-27.