

Pragmatic Trials in Long-Term Care

Citation for published version (APA):

Resnick, B., Zimmerman, S., Gaugler, J., Ouslander, J., Abrahamson, K., Brandt, N., Colón-Emeric, C., Galik, E., Gravenstein, S., Mody, L., Sloane, P. D., Unroe, K., & Verbeek, H. (2022). Pragmatic Trials in Long-Term Care: Research Challenges and Potential Solutions in Relation to Key Areas of Care. *Journal of the American Medical Directors Association*, 23(3), 330-338.
<https://doi.org/10.1016/j.jamda.2021.12.011>

Document status and date:

Published: 01/03/2022

DOI:

[10.1016/j.jamda.2021.12.011](https://doi.org/10.1016/j.jamda.2021.12.011)

Document Version:

Publisher's PDF, also known as Version of record

Document license:

Taverne

Please check the document version of this publication:

- A submitted manuscript is the version of the article upon submission and before peer-review. There can be important differences between the submitted version and the official published version of record. People interested in the research are advised to contact the author for the final version of the publication, or visit the DOI to the publisher's website.
- The final author version and the galley proof are versions of the publication after peer review.
- The final published version features the final layout of the paper including the volume, issue and page numbers.

[Link to publication](#)

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal.

If the publication is distributed under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license above, please follow below link for the End User Agreement:

www.umlib.nl/taverne-license

Take down policy

If you believe that this document breaches copyright please contact us at:

repository@maastrichtuniversity.nl

providing details and we will investigate your claim.



JAMDA

journal homepage: www.jamda.com

Special Article

Pragmatic Trials in Long-Term Care: Research Challenges and Potential Solutions in Relation to Key Areas of Care



Barbara Resnick PhD, CRNP^{a,*}, Sheryl Zimmerman PhD^b, Joseph Gaugler PhD^c, Joseph Ouslander MD^d, Kathleen Abrahamson PhD, RN^e, Nicole Brandt PharmD, MBA^f, Cathleen Colón-Emeric MD, MHSc^g, Elizabeth Galik PhD, CRNP^a, Stefan Gravenstein MD, MPH^h, Lona Mody MD, MSCⁱ, Philip D. Sloane MD, MPH^b, Kathleen Unroe MD, MHA, MS^j, Hilde Verbeek PhD^k

^a University of Maryland School of Nursing, Baltimore, MD, USA

^b Cecil G. Sheps Center for Health Services Research and Schools of Social Work and Public Health, University of North Carolina at Chapel Hill, Chapel Hill, NC, USA

^c Division of Health Policy & Management, School of Public Health, University of Minnesota, Minneapolis, MN, USA

^d Department of Integrated Medical Sciences, Charles E. Schmidt College of Medicine, Florida Atlantic University, Boca Raton, FL, USA

^e Purdue University, West Lafayette, IN, USA

^f University of Maryland School of Pharmacy, Baltimore, MD, USA

^g Duke University School of Medicine and the Durham VA GRECC, Durham, NC, USA

^h Brown University and Providence Veterans Administration Medical Center, Providence, RI, USA

ⁱ University of Michigan and Veterans Affairs, Ann Arbor Healthcare System, Ann Arbor, MI, USA

^j Indiana University School of Medicine and Regenstrief Institute, Inc, Center for Aging Research, Indianapolis, IN, USA

^k Maastricht University and Living Lab in Ageing and Long-Term Care, Maastricht, the Netherlands

ABSTRACT

Keywords:
Pragmatic trials

As a method of research, pragmatic trials are recommended so as to generate results that are applicable to real-world care. This intent is especially important for the millions of older adults who receive long-term care in thousands of nursing homes and assisted living communities across the country—and many millions more around the globe. This article presents key points raised by experts participating in a conference funded by the National Institute of Aging held at the 2021 conference of the Society for Post-Acute and Long-term Care Medicine. The purpose of the conference was to convene leading clinicians, researchers, and industry partners to address special considerations of pragmatic trials in long-term care. Cross-cutting and unique challenges and solutions to conducting pragmatic trials were discussed focusing on 3 areas of clinical relevance to long-term care: (1) functional care and outcomes, (2) psychosocial care and quality of life, and (3) medical care and outcomes, with a special focus on persons with dementia. Challenges and innovative solutions were organized across the 9 domains of the revised Pragmatic-Explanatory Continuum Indicator Summary (PRECIS) Tool, and future research recommendations for pragmatic trials in long-term care were identified.

© 2021 AMDA – The Society for Post-Acute and Long-Term Care Medicine.

This work was supported by the National Institute on Aging (NIA) of the National Institutes of Health, Award R13 AG067681. Sheryl Zimmerman and Joseph Gaugler were also supported by the NIA under Award Number U54AG063546, which funds the NIA Imbedded Pragmatic Alzheimer's Disease and AD-Related Dementias Clinical Trials Collaboratory (NIA IMPACT Collaboratory). The content is solely the responsibility of the authors and does not necessarily represent the official views of the National Institutes of Health.

The authors declare no conflicts of interest.

* Address correspondence to Barbara Resnick, PhD, CRNP, University of Maryland School of Nursing, 655 W Lombard St, Baltimore, MD 21201, USA.

E-mail address: Resnick@umaryland.edu (B. Resnick).

There have been calls to improve the quality of long-term care for more than 30 years,¹ but changing behavior is challenging and implementing new approaches to care is slow. When new care practices are developed, they are typically evaluated in well-controlled clinical trials, meaning that the extent to which they are suited for practical real-world implementation is unclear. On the other hand, pragmatic trials examine real-world efficacy of an intervention in relevant settings and provide the opportunity to adapt it to address implementation challenges. Pragmatic trials are needed to better disseminate and implement effective approaches to care and deimplement ineffective approaches. They also encourage clinicians and

others working in care settings to partner with researchers in the development of innovative approaches to care.² The National Institute on Aging provides guidance in designing research to effect change through their Stage Model, which defines the stages of research from basic science (Stage 0) to dissemination and implementation (Stage V); the conduct of pragmatic trials fits in Stage IV, effectiveness.³

Nursing homes and assisted living communities are complex adaptive systems, composed of numerous individuals interacting in ways that are not always predictable and always having to adapt.^{4,5} Thus, optimal strategies to conduct pragmatic trials in long-term care are not straightforward. In response to the need to address special considerations of pragmatic trials in long-term care, especially for persons living with dementia, the National Institute of Aging funded a conference to bring together leaders to discuss related research challenges and potential solutions. To create focus, the organizers identified 3 areas of clinical relevance to long-term care: (1) functional care and outcomes, (2) psychosocial care and quality of life, and (3) medical care and outcomes. Clinical trials have demonstrated effective approaches to care in these areas, including to increase physical activity,^{6,7} decrease falls,^{8,9} improve management of behavioral expressions,^{10,11} prevent transfers to acute care settings and address goals of care,¹² improve medication management,¹³ and enhance infection/viral control.^{14,15} However, evidence of efficacy based on randomized controlled trials does not generally change care practices in real world settings. Further, some care practices persist despite questionable effectiveness such as obtaining computed tomography of the head and neurologic checks following all falls¹⁶ and use of bed and chair alarms to prevent falls.¹⁷

Conducting Pragmatic Trials in Long-Term Care

Investigators must understand the unique characteristics of people who live in long-term care settings when conducting pragmatic trials. The majority of residents have cognitive impairment among other comorbidities.^{18–21} They frequently exhibit behavioral and psychological symptoms of dementia (now recognized as behavioral expressions)²² and experience undertreated pain, functional decline, and limited physical activity.²³ In addition, 25% of hospitalizations are considered potentially avoidable.²⁴ Thus, there is a need to provide long-term care for persons with dementia and other geriatric syndromes that is not only efficacious but also scalable and practical for these complex care environments. Appreciation of other key aspects of these environments that influence intervention design and implementation include such things as staffing models, reimbursement, and regulatory constraints.

In contrast to the development of new knowledge, funders and investigators have traditionally been less focused on dissemination or implementation of research findings into real world settings.²⁵ Evidence-based research protocols are frequently too complex for implementation³ or not developed for actual users,²⁶ and clinical settings lack management support, organizational policies and practices, financial resource availability, organizational readiness for change, and measurement that is acceptable, compatible, practical, and useful.^{27,28} To gain a better understanding of how to best integrate optimal care practices into long-term care settings, there has been an increased focus on conducting pragmatic trials.²⁵

Differences Between Explanatory Trials vs Pragmatic Trials

Explanatory trials are used to demonstrate the initial efficacy of an intervention and are conducted to support or refute a clear clinical hypothesis. These trials are critically important to determine whether an intervention will work under optimal conditions. In explanatory trials, interventions are tested under ideal and controlled situations and generally focus on individuals who are most likely to receive the greatest benefit; also, resources and staff are often provided beyond what is available in usual care environments. Explanatory trials strive

Table 1
Description of the Domains Within PRECIS-2³⁰

Domain	Description
Eligibility	Who is eligible to be in the trial?
Recruitment	How are participants recruited into the trial?
Setting	Where is the trial being conducted?
Organization	What expertise and resources are needed from the organization to deliver the intervention?
Flexibility in delivery	How should the intervention be delivered?
Flexibility in adherence	What is being done to ensure adherence to the intervention?
Follow-up	How closely are the participants monitored or followed?
Primary outcome	How relevant is the outcome to participants?
Primary analysis	Are all data included regardless of each individual's level of participation?

to maximize the internal validity of results so that investigators have confidence that findings are due primarily to the effects of an intervention and not confounding factors. These trials create an evidence base for efficacy. Conversely, results from pragmatic trials are used to inform clinical or policy decisions by providing evidence that the intervention demonstrates effectiveness in any relevant setting.²⁹ There are often, however, varying degrees to which a trial is considered “pragmatic” vs “explanatory” based on the type of intervention and the participants involved. To help determine the extent to which a trial is pragmatic, the Pragmatic-Explanatory Continuum Indicator Summary (PRECIS-2) Tool³⁰ was developed. The purpose of PRECIS-2 is to help researchers design studies that are truly pragmatic.

Description of PRECIS-2

PRECIS-2 includes the following 9 domains (Table 1): Eligibility, Recruitment, Setting, Organization, Flexibility in Delivery, Flexibility in Adherence, Follow-up, Primary Outcome, and Primary Analysis. Each of the domains is evaluated by the individuals designing the study to determine to what degree the trial is pragmatic. Scoring on each domain ranges from very explanatory (1), rather explanatory (2), equally pragmatic and explanatory (3), rather pragmatic (4), or very pragmatic (5). Use of PRECIS-2 can help assure that truly practical trials are being conducted in long-term care and that they will be relevant for those living and working in these communities.

Pragmatic Trials in Long-Term Care: Challenges and Solutions

The NIA-funded Pragmatic Trials in Long-Term Care conference included presentations by 10 experts (see acknowledgments) who discussed research challenges and solutions in 10 areas related to the 3 clinical foci delineated above: falls prevention, activities of daily living, function-focused care (related to functional care and outcomes); nonpharmacologic practices, aging in place, advance care planning (related to psychosocial care and quality of life); and medication use, prevention, infections, and health service use (related to medical care and outcomes). Using the PRECIS-2, they commented on up to 4 domains constituting particular challenges in pragmatic trials in long-term care in their area. Table 2 summarizes key challenges for the 9 PRECIS-2 domains, all of which are described in detail below. Although Table 2 is not a comprehensive list of all challenges, it demonstrates that challenges can occur across multiple domains of the PRECIS-2 Tool.

Eligibility of Participants

Highly pragmatic trials include broad eligibility criteria for participants so as to include diverse settings of care, residents, and/or staff. Even broad criteria require parameters, though, in that some interventions may be more useful for select residents and settings;

Table 2
Challenges Identified in Pragmatic Trials in Long-Term Care Based on Categories in the PRECIS-2

Topic	Challenges*								
	Participants	Recruitment	Settings	Organization	Delivery	Adherence	Measurement Follow-up	Primary Outcome	Primary Analyses
Falls prevention*	Heterogeneity among residents; resident goals may conflict with adherence			Heterogeneity in staff training and background; appropriate level of randomization may vary across facilities	Heterogeneity in staffing models; turnover	Multifactorial interventions require multiple fidelity or process measures		Unreliable facility data; Hawthorne effect; surrogate outcomes not important to residents; process outcomes difficult to document	
Activities of daily living*	Variability of length of stay; timing of assessment and intervention		Variability of settings; contamination risk if not randomized by facility				Lack of sensitivity of assessments; attrition in target group	Bias in proxy reports of function	
Function-focused care*		Challenges to assenting and consenting residents with dementia			Identifying invested champion(s); securing oversight at the facility level; anticipating and managing staff turnover; allowing flexibility in delivery	Competition with new initiatives; lack of accountability; need to educate new staff	Lack of sensitivity in proxy reports; timing of routine data collection may not capture results		
Nonpharmacologic practices*				Intervention must be relevant for the setting and have low initiation and maintenance costs	Intervention must not require initiation by resident and should not exceed routine care activities	Adherence must be monitored over time		Outcome measure must take into account that the week is 168 h	
Aging in place*	Need to consider cognition and function and presence of informal caregiver; should include persons in remote areas and diverse populations (eg, those with difficult family, mental illness)		Remote or underserved settings are difficult to access; communities may change over time (eg, in food availability, walkability, gentrification, transportation)		Ensuring a variety of preferences and individualized needs may result in low intervention specificity; requiring an infusion of resources may be impractical			Outcomes are broad (eg, relate to location, health status, mortality, social roles, activities)	

Advance care planning (ACP)*	Vulnerable population with complex consent issues; trials require assessment of capacity for participation		Poor data integration between settings; longitudinal data require robust plan for input from diverse settings			Intervention fidelity data should be collected to understand outcomes		ACP research lacks consensus on key outcome measures; there are few validated measures of the quality of communication and goal-concordant care; hospitalization and utilization occur for a subset, limiting power	
Medication use*	Obtaining consent from residents who are decisionally impaired vs waiving consent			Differences in care models across countries; financing of long-term care and implications; provider expertise and resource availability	Study fidelity; stakeholder engagement; organizational support and readiness		Completeness of regulatory data and completeness and accuracy of existing data		
Prevention*		Recruit facilities, not residents	Infection control regulations vary by state	Practice change is slow absent formal recommendation			Challenges to maintaining access and completing entries for primary data, and data matching between data sets (provider and resident IDs)	Evolving coding practices	Adjusting for facility-level factors
Infections*	Targeted vs inclusive approach requires consideration	Short-stay residents may be discharged before recruitment, introducing selection bias	Facilities are often diverse in their focus, services, case mix, infection control infrastructure, availability and engagement of providers, and strength of connections with local hospitals	Focus on QI and research engagement can vary across facilities	Time constraints, staff turnover, distance from academic centers, competing priorities	Staff turnover, prior staff training, leadership engagement, priorities of the facility and their parent corporations	Outcome measures that require frequent follow-up	Outcome definition cannot be complex (ie, definitions such as for UTI are often very specific); expectations from academic journals require rigorous assessments	Need to measure at the specimen, pathogen, visit, or patient level depending on the research question
Health services use*				Scarce resources, staff resistance, competing demands, instability of leadership	Interventions are complex; may need evidence to justify more intensive interventions	Fidelity and fidelity monitoring need attention			
Total number	6	3	5	6	7	6	5	7	2

EHR, electronic health record; QI, quality improvement; UTI, urinary tract infection.

*Each presenter was asked to identify the 3 or 4 most pressing challenges to highlight; therefore, this list is not complete.

Table 3
Future Areas of Suggested Research

Functional care and outcomes
<ul style="list-style-type: none"> • Evaluation of the environment as an active intervention ingredient (eg, Green Care Farms) • Use of Ecological Momentary Assessment and assessment of resident participation in activities of daily living • Strategies to motivate staff to engage residents in physical activity • Interventions to increased administrative support of Function Focused Care • Evaluation of phenotypes of residents who fall • Use of wearable devices to measure falls
Psychosocial care and quality of life
<ul style="list-style-type: none"> • Evaluation of what organizational assessment tools inform successful implementation • Impact of the environment on stimulation of persons with dementia • How activities of daily living can result in positive experiences between persons with dementia and their caregivers • Whether evidence of outcomes related to psychosocial care and quality of life is transferable to rural settings • Cognitive capacity and frailty as related to completion of advance care plans • Advance care planning outside the nursing home setting and in diverse populations
Medical care and outcomes
<ul style="list-style-type: none"> • Interventions to improve the quality of prescribing, including overprescribing • Impact of immunization on functional loss, cardiovascular events, and outbreak prevention, and comparisons of enhanced vaccines • Engaging residents, family, and visitors in infection prevention • The role of in-room surfaces in the transfer of bacteria and viruses • Development of systematic solutions to decrease transmission of pathogens, and implementation of interventions with proven evidence-based infection prevention • Evaluation of the potential of telehealth to provide services to residents • Connecting electronic health records across systems to improve patient transfers • Regaining a focus on person-centered care and the match of patient/family goals with treatment

such parameters need to be articulated and justified. One consideration for determining eligibility is ensuring that residents' personal goals and preferences do not conflict with the intervention, such as a resident who prefers autonomy over falls-risk reduction when the setting is focused on falls prevention. Temporal considerations also influence eligibility considerations as for some studies it may be more appropriate to target new admissions regardless of the potential merit for all residents. Again, such decisions must be carefully weighed and justified if the trial is to be fully pragmatic.

Further, different interventions may be necessary in pragmatic trials to recognize the heterogeneity of residents, which may affect eligibility or the intervention itself. Implementation of interventions for advance care planning, for example, requires inclusion of surrogate decision makers; the absence of such decision makers may affect eligibility. Likewise, health literacy can influence the appropriateness of the intervention for certain groups of participants, suggesting a need to attend to modify the intervention lest it not be optimally pragmatic.

Potential solutions

To overcome some of the challenges associated with the heterogeneity of potential participants, it may be necessary to utilize administrative data and identify key subgroups that will most likely benefit from the intervention or that reduce confounding. Conducting secondary or post hoc analyses on original efficacy trial data to obtain a better understanding of which subgroups responded positively or with the least amount of variance to the intervention can help target specific participants in the pragmatic trial phase. In this manner, the intervention could deliberately address different subgroups within a setting and tailor interventions for each of those groups. Consideration of advanced trial designs, such as sequential, multiple

assignment, randomized trials (SMARTs) where participants are randomized at multiple stages during an intervention to allow for more adaptive approaches, may also be warranted to ensure effective tailoring of an intervention in long-term care.^{31,32} Shared decision-making tools are another way to match individual goals with the intervention being implemented.¹²

Recruitment of Participants

When conducting pragmatic trials, the goal is to include all eligible participants to best reflect standard practice. Therefore, recruitment must avoid strategies for recruitment that may bias participation. Challenges to recruitment in pragmatic trials may include the need to consent individuals if gathering identifiable data, making them de facto less pragmatic. Cognitive impairment impacts both consent and data collection. The inability to access caregivers or guardians and the lack of assent among the very residents who might benefit the most from the intervention (eg, those with behavioral expressions) are major challenges to recruitment to all trials in long-term care.³³

Potential solutions

Various approaches to avoid the need for or to facilitate consent include such things as consent waivers, broadcast notification, integrated consent, and targeted consent.³⁴ Consent waivers allow researchers to gather deidentified data already collected. Broadcast notification involves placing notices in prominent locations that inform potential patients of the ability to participate in research related to their care. Integrated consent integrates clinical and research consent into a single clinical encounter (often at the time of admission), whereas targeted consent involves a brief consent process followed by an information sheet for participants informing them that they are helping researchers explore a specific topic.

One overall approach that ensures a trial is more pragmatic is to reduce reliance on primary data collection and instead use deidentified data extracted from health records; this approach reduces the need for informed consent but also limits the nature of the data available for study. It is also helpful to use a modified consent procedure and obtain verbal informed consent to reduce the burden of signed consent for a *low-risk* intervention. Limiting the complexity of an intervention will increase willingness to participate as will highlighting the relevance of the effort for potential participants, particularly when reaching out to those from underrepresented racial and ethnic groups.

Inclusion of Settings

Pragmatic trials should strive to include all types of long-term care settings as appropriate for their target population, but inherent differences in settings (such as nursing homes vs assisted living) may require separate trials for each setting type. In addition, pragmatic trials often rely on cluster randomization to avoid contamination between intervention and control within a given setting, thereby requiring a large sample of randomization units such as nursing homes to ensure adequate statistical power (eg, because individuals are nested within settings) and address variation. It may be challenging to ensure that settings randomized to the intervention vs usual care are in fact similar, particularly when a relatively small number of settings are included. It is also challenging for research teams to include settings that may be more rural and therefore more difficult to access. In addition, whether or not settings are part of a larger entity (eg, health system or corporation) may affect the resources available for an intervention; a pragmatic trial is one that is suitable for all relevant settings. Other considerations for pragmatic trials in long-term care include the services that are available (eg, separate dementia care units, electronic health records that connect

with other sites such as hospitals), and also variations in relevant regulations due to state or regional differences.

Potential solutions

To best address or overcome these challenges, consideration should be given to the match between components of the intervention and care settings. For example, an intervention focused on optimizing function and physical activity would not match with a setting that focused on managing pain by (erroneously) decreasing mobility in an effort to lessen pain. Instead, a better match would be with a setting focused on falls prevention that supported the philosophy that physical activity can prevent falls. If it is feasible to include a large number of settings, it may be useful to account for variation in settings by creating subgroups (eg, within and without memory care units) and proceed with stratified random sampling. In addition, because imbalances between treatment and control group settings may occur when using cluster randomization, it may be advisable to use multiple-stage constrained randomization techniques to limit imbalance.³⁵ Further to the point of matching the intervention with the setting, some have suggested assessing organizational readiness to determine the setting's readiness and capability to carry out the various elements of the pragmatic trial.³⁶ However, there is no uniform endorsement to limit intervention opportunities to those that are the most ready to enact them because doing so may disadvantage the settings that are most needy.³⁷ Finally, because some administrators may not prioritize a given intervention, it may be more useful to focus on settings with a particular interest in the intervention being offered; once effectiveness and pragmatism are established, hesitant administrators may be persuaded.

Organization of Care Delivery

Because pragmatic trials conduct interventions in “real-world” practice rather than highly controlled research settings, the organization of care delivery in long-term care is highly consequential. In this regard, challenges include heterogeneity in staff training and staffing ratios; limited and variable organizational resources; few or no policy requirements, recommendations, or accreditation standards related to the intervention content; and hands-off involvement of administrative staff, to name but a few. It is usually the case that time is limited to educate staff regarding a new intervention (especially if it is impractical and complex), that there are competing demands on staff time (even if time is available for training), that high industry-wide turnover rates affect sustainability, and that the lack of stable leadership or “champions” impedes adoption.

Potential solutions

To overcome infrastructure challenges, it may be helpful to assess the culture of the organization using culture assessment tools³⁸ prior to starting the trial as well as organizational readiness as described earlier, perhaps with an eye toward promoting the culture and readiness of those less able. A principal objective when doing so is to match the intervention and study outcomes to the culture and priorities of care of the partner organization. Use of electronic health records and other tools embedded within the healthcare system can facilitate intervention delivery processes and should be fully incorporated into the data collection of a proposed pragmatic trial as appropriate. Other implementation approaches that may be helpful include using outside resources to provide education or training of staff (or relying on remote or asynchronous methods to do so)³⁹ and making sure that initiation and maintenance costs are low. Overall, it is best that the intervention fit the existing infrastructure, such as capitalizing on electronic learning management systems where modules can be uploaded, or regularly scheduled staff in-service

sessions that can be leveraged to deliver intervention-related trainings.

Flexible Delivery of the Intervention

Pragmatic trials typically ensure that the delivery of the intervention is flexible so that it can be individualized for participant needs and fit better into real-world settings. A key concern in this regard is that it is not often clear which components of the intervention are modifiable if the outcomes shown to be efficacious in an explanatory trial are to be achieved. Further, the components that are modified by the staff may in fact be those that are central to achieving the intended outcome. For example, one study of a multicomponent intervention conducted in long-term care found that the combination of components (in this case, families attending a workshop, a care plan being developed, and being followed) was more related to outcomes than was any single component.⁴⁰ In this case, it was critical that all components be included to achieve the desired outcomes. Had the workshop not been offered, the intervention would have been less effective; it may have been possible to modify the workshop, but not to omit it completely.

Potential solutions

To optimize delivery, it is critical to ensure that the intervention is clearly defined and fits within the current workflow of the setting; doing so may be facilitated by offering a menu of potential modification strategies from which staff can choose to minimize the practice change required. When doing a pragmatic trial, the use of hybrid effectiveness designs are recommended because outcomes are considered alongside key measures of implementation (eg, appropriateness, feasibility, acceptability). This design can help ensure that the intervention aligns with the capacity and resources of the organization and can be implemented as intended.⁴¹ For a pragmatic trial to be successful, researchers must work with the care organization to optimize delivery of all components as intended and to identify the core elements of an intervention that must remain as well as the flexible elements that can be adapted to better fit the preferences and needs of residents, families, and staff. Doing so requires that the pragmatic trial include pragmatic measures of fidelity.

Flexibility of Adherence

Beyond delivering the intervention with suitable flexibility, it must be adhered to and sustained. Challenges to adherence include competing initiatives, resource-intensive interventions, and those perceived as useless to the staff or residents. There is widespread recognition among pragmatic trialists that fidelity must be assessed in an ongoing manner, must be done as unobtrusively as possible, and in a flexible manner that fits with daily workflow. Asking staff to monitor adherence can be burdensome, as found in a recent trial in which charting new care practices was more often missing than not.⁴² Moreover, it is not clear when to measure adherence and whether the focus is relevant for short- (less than 6 months) vs long-term adherence (greater than 6 months). Also challenging is when supervisors do not require accountability for changed care, which relates to the earlier pragmatic consideration regarding the setting itself.

Potential solutions

Much has been written about the benefits of identifying and involving a committed champion and stakeholder team to promote adoption and fidelity. Toward this end, embedding pragmatic trials in a model of quality improvement is likely to achieve the most buy-in and will by design include follow-up monitoring. Alternate modes of monitoring might also be considered, especially those that do not impose on staff. Technology is one such option, such as when the

intervention itself provides counts or an observable means regarding use; in fact, it has been noted there is need to better use technology in care for persons with dementia.⁴³ For example, the use of actigraphy with devices such as the Motionwatch8 is a useful way to capture data on physical activity in a pragmatic trial. Another option for monitoring may be including resident and family reports—if such can be done in a pragmatic way.

Measurement of Follow-Up

There are numerous challenges related to follow-up measurement of residents in long-term care. When care is intended to benefit a resident with dementia, the person may not be able to self-report, or self-report may be limited to responses with limited sensitivity to change, such as yes/no or simple Likert-type scales. Family may be called on as respondents, but proxy reports are known to be biased, often in a negative direction.⁴⁴ Also, both of these options may create undue burden, contrary to the intent of pragmatic trials. Observational measures may be resource intensive and similarly not pragmatic unless they are technology-based. Further, use of secondary data may not coincide with optimal timing to detect change, and items such as from Medicare claims data or the Minimum Data Set may not be optimally relevant or sensitive to change.

Potential solutions

To overcome measurement challenges, it is helpful to capitalize on existing data. In so doing, it may be advisable to triangulate those data with other data sources to determine reliability (eg, physical activity data from Minimum Data Set data and actigraphy). If leadership is supportive of the trial, it may be possible to collaborate to incorporate new structured data elements within paper or electronic templates. Indeed, focusing on outcomes that are included in the clinical record, such as basing them on documented notes or goal attainment⁴⁵ is a promising option to facilitate the availability of data if the records are sufficiently detailed. Regardless the strategy, it is critical that data elements be clearly defined (eg, definition of pneumonia or a urinary tract infection) and that if others are asked to provide the data, that they find it to be important.

Relevance of Primary Outcomes

Outcomes should be relevant to residents and other stakeholders in long-term care if the trial is to be pragmatic. In this regard, a key challenge is reconciling outcomes that are important to residents and stakeholders with available data (eg, recorded hospitalizations) and validated outcome measures. For example, researchers tend to consider hospitalizations as something to be avoided, whereas in some cases residents and their family may be reassured if there is a hospitalization. Relatedly, it may be that the wrong component of an outcome is being measured—such as measuring the completion of an advance directive when what truly mattered was having the related conversation. Process outcomes, such as whether or not the activity associated with the intervention was done, may be particularly relevant for pragmatic trial results, but difficult to document and capture.

Potential solutions

Pilot studies and review of existing literature may be the best strategy to ensure that the outcomes being collected are relevant to stakeholders. Toward this end, collaborating with long-term care settings during the development of the trial is indicated, and has been noted as a critical gap.³⁷ If residents, families, and staff endorse the importance of key outcomes, they may be more likely to support the collection of other data they find less central, including fidelity data noted earlier.

Nature of Primary Analyses

Pragmatic trials use an intention to treat analysis so as to ascertain effects of an intervention in real-world settings—as opposed to per-protocol analyses that examine effects when the intervention is provided as intended. As such, data from all participants are relevant, but in long-term care, attrition and dropout are common given resident death and staff turnover. Not only might their outcome data be unavailable but the resulting data may be biased given the selective nature of attrition.

Potential solutions

Analytically, adjusting for covariates that relate to outcomes may allow a more valid indication of the treatment effect. To do so, it is necessary that those covariates be known in advance and collected in advance, presumably from residents' medical records (even if they are not fully sufficient for adjustment). Further, prior research may suggest setting specific factors known to influence outcomes and attrition that can be included in analytic models. Use of theory or conceptual models that consider the multilevel effects of interventions in long-term care may help inform setting-level characteristics to address in pragmatic trials.

Successful Pragmatic Trials and Approaches Used for Sustainability of Interventions

Across multiple areas, experts reported that effective and sustainable interventions were those that were feasible and allowed for flexibility in implementation. Feasible interventions are those that are integrated into the care plan or part of ongoing care processes,^{7,46} aligned with policy, or important to the setting based on relevant organizational, patient-centered, and/or regional/national goals.⁴⁷ Active participation of a champion and stakeholder team was consistently mentioned as essential to successful implementation and sustainability.^{46,48,49} Care initiatives that are perceived as beneficial to staff and residents are also more likely to be sustained by staff providing the care.^{50–52}

The use of technology was repeatedly encouraged to facilitate scalability of interventions and approaches.^{53,54} Technology included wearable devices, use of electronic health records, and the development and use of apps to facilitate intervention delivery and staff training. There was strong consensus that successful implementation required that the intervention be adaptable or adjustable based on feedback from organizational partners and that consideration be given to the business of health care and cost/benefit of the new approach when compared to usual care. Lastly, deimplementation of ineffective interventions was recognized as an important component to successful implementation of new care practices.⁵⁵ Deimplementation involves the removal or the replacement of care interventions that are known to be ineffective, harmful, or not beneficial.⁵⁶

Conclusions and Implications

Challenges to designing and implementing pragmatic trials in long-term care were noted across all 9 domains of PRECIS-2. Although not a comprehensive list, the most commonly noted challenges included identifying and obtaining primary outcomes that are relevant to residents and families as well as practical to collect, adherence to the intervention, allowing flexibility in delivery, including heterogeneous participants while simultaneously ensuring sufficient control to determine intervention effectiveness, and the variable nature of the organization.

In long-term care, flexibility in delivery is critical and it may be beneficial to help facilities become ready to initiate the intervention.⁵⁷ In so doing, it is possible to ensure that more disadvantaged facilities are able to participate successfully. If a trial is to be optimally

pragmatic, exclusions should be limited to settings in which the intervention lacks relevance. Recognizing setting-specific differences, there may be need to revise training or intervention materials so that they are culturally appropriate and relevant for staff and participants.⁵⁸ Technology and use of existing data are critical going forward, yet both need further development and evaluation.

In a pragmatic trial, recruitment is done by providers and is offered to all who may benefit. Future work is required to establish strategies to ethically include residents who may not have the capability to assent or consent and may not have an identified proxy.^{59–61} Waiving consent is allowable in situations in which the intervention is low risk, does not impact the welfare or rights of the participants, and when comparing 2 different but equally effective care approaches.⁶² The waiver of consent, however, impacts the type of outcomes that can be obtained.

Table 3 provides an overview of the areas of pragmatic trial research suggested by the expert participants in the 2021 conference. Some topics relate to methodology such as using new measurement approaches; others address learning more about strategies to ensure successful implementation by establishing factors within organizations that inform success, or translating interventions for new settings. The majority of the suggested areas of research involve using pragmatic trials to learn more about factors that influence functional care and outcomes (eg, falls), psychosocial care and quality of life (eg, advance care planning), and medical care and outcomes (eg, medication prescribing, infection prevention). The ideas promoted by the group at this meeting are not likely to be comprehensive of all possible challenges and solutions. For example, there may be a benefit to engaging with resident or family councils to vet potential outcome measures of importance to them or to seek the help of an ombudsman to facilitate recruitment of a participant.

Pragmatic trials are critically important to our ability to disseminate and implement useful and sustainable interventions in long-term care. These types of trials ensure that the intervention is not only effective but can be implemented in real-world settings and organizations among a range of residents and staff. Although there are some challenges to designing and implementing truly pragmatic trials based on PRECIS-2 domains, challenges can be overcome using innovative approaches. In doing so, it is possible to build from the explanatory to the pragmatic and help ensure that dissemination and implementation of new interventions will be successful and sustainable in real-world settings.

Acknowledgment

Conference presenters and moderators included the authors and also Laura Hanson, MD, MPH, University of North Carolina at Chapel Hill.

References

- Institute of Medicine (US) Committee on Nursing Home Regulation. Improving the Quality of Care in Nursing Homes. National Academies Press; 1986. <https://pubmed.ncbi.nlm.nih.gov/25032432/>. Accessed November, 2021.
- Voorberg W, Bekkers V, Tummers L. A systematic review of co-creation and co-production: Embarking on the social innovation journey. *Public Manage Rev* 2015;17:1333–1357.
- Onken L, Carroll K, Shoham V, Cuthbert B, Riddle M. Reenvisioning clinical science: unifying the discipline to improve the public health. *Clin Psychol Sci* 2014;2:22–34.
- Anderson R, Issel L, McDaniel RJ. Nursing homes as complex adaptive systems: relationship between management practice and resident outcomes. *Nurs Res* 2003;52:12–21.
- Zimmerman S, Mitchell C, Beeber A, et al. Strategies to reduce potentially inappropriate antibiotic prescribing in assisted living and nursing homes and inform other quality improvement efforts. *Am J Med Res* 2015;2:41–52.
- Galik E, Resnick B, Lerner N, Hammersla M, Gruber-Baldini A. Function focused care for assisted living residents with dementia. *Gerontologist* 2015;55(Suppl 1):S13–S26.
- Resnick B, Boltz M, Galik E, et al. Testing the implementation of function focused care in assisted living settings. *J Am Med Dir Assoc* 2021;22:1706–1713.
- Resnick B, Boltz M, Galik E, et al. The impact of function focused care and physical activity on falls in assisted living residents. *Can J Nurs Res* 2020;52:45–53.
- Cameron I, Dyer S, Panagoda C, et al. Interventions for preventing falls in older people in care facilities and hospitals. *Cochrane Database Syst Rev* 2018;9:CD005465.
- Cohen-Mansfield J. Nonpharmacologic interventions for inappropriate behaviors in dementia: a review, summary, and critique. *Am J Geriatr Psychiatry* 2001;9:361–381.
- Kolanowski AM, Litaker M, Buettner L, Moeller J, Costa P. A randomized clinical trial of theory-based activities for the behavioral symptoms of dementia in nursing home residents. *J Am Geriatr Soc* 2011;59:1032–1041.
- Hanson L, Zimmerman S, Song M, et al. Effect of the goals of care intervention for advanced dementia: a randomized clinical trial. *JAMA Intern Med* 2017;177:24–31.
- Garland C, Guénette L, Kröger E, Carmichael P, Rouleau R, Sirois C. A new care model reduces polypharmacy and potentially inappropriate medications in long-term care. *J Am Med Dir Assoc* 2021;22:141–147.
- Mody L, Krein S, Saint S, et al. A targeted infection prevention intervention in nursing home residents with indwelling devices: a randomized clinical trial. *JAMA Intern Med* 2015;175:714–723.
- Gravenstein S, Dahal R, Gozalo P, et al. A cluster randomized controlled trial comparing relative effectiveness of two licensed influenza vaccines in US nursing homes: design and rationale. *Clin Trials* 2016;13:364–374.
- Pages P, Boncoeur-Martela M, Dalmay F, et al. Relevance of emergency head CT scan for fall in the elderly person. *J Neurol* 2020;47:54–58.
- Shorr R, Chandler A, Mion L, et al. Effects of an intervention to increase bed alarm use to prevent falls in hospitalized patients: a cluster randomized trial. *Ann Intern Med* 2012;157:692–699.
- Hyde J, Perez R, Forester B. Dementia and assisted living. *Gerontologist* 2007;47:51–67.
- Leon J, Cheng C, Neumann P. Alzheimer's disease care: costs and potential savings. *Health Aff* 1998;17:206–216.
- Zimmerman S, Sloane P, Reed D. Dementia prevalence and care in assisted living. *Health Aff* 2014;33:658–666.
- Harris-Kojetin L, Sengupta M, Park-Lee E, et al. Long-term care providers and services users in the United States: data from the National Study of Long-Term Care Providers, 2013–2014. *Vital Health Stat* 2016;38:x–xii; 1–105.
- Steinberg M, Shao H, Zandi P, Lyketsos CG, Welsh-Bohmer K, Norton M; Cache County Investigators. Point and 5-year period prevalence of neuropsychiatric symptoms in dementia: the Cache County study. *Int J Geriatr Psychiatry* 2018;23:170.
- Shiells K, Pivodic L, Holmerová I, Van den Block L. Self-reported needs and experiences of people with dementia living in nursing homes: a scoping review. *Aging Ment Health* 2019;1:1–16.
- Healthy People 2020. Dementias, including Alzheimer's disease. <http://www.healthypeople.gov/2020/topics-objectives/topic/dementias-including-alzheimers-disease/national-snapshot>. Accessed November, 2021.
- Glasgow R, Vinson C, Chambers D, et al. National Institutes of Health approaches to dissemination and implementation science: current and future directions. *Am J Public Health* 2012;102:1274–1281.
- Miller H, Ward K, Zimmerman S. Implementation effectiveness of psychosocial and environmental care practices in assisted living. *Geriatr Nurs* 2021;42:295–302.
- Powell B, Stanick C, Halko H, et al. Toward criteria for pragmatic measurement in implementation research and practice: a stakeholder-driven approach using concept mapping. *Implement Sci* 2017;12:118.
- Weiner B. A theory of organizational readiness for change. *Implement Sci* 2009;4:67.
- Schwartz D, Lellouch J. Explanatory and pragmatic attitudes in therapeutical trials. *J Chron Dis* 1967;20:637–648.
- Loudon K, Treweek S, Sullivan F, Donnan P, Thorpe K, Zwarenstein M. The PRECIS-2 tool: Designing trials that are a fit for purpose. *BMJ* 2015;350:1–11.
- The Penn State Methodology Center. Why Use a SMART Design to Build an Adaptive Intervention? The Pennsylvania State University College of Education and Human Development. 2020. <https://www.methodology.psu.edu/ra/adap-inter/research/>. Accessed November, 2021.
- Almirall D, Compton S, Gunlicks-Stoessel M, Duan N, Murphy S. Designing a pilot sequential multiple assignment randomized trial for developing an adaptive treatment strategy. *Stat Med* 2012;31:1887–1902.
- Resnick B, Kolanowski A, Van Haitsma K, et al. Current psychotropic medication use and contributing factors among nursing home residents with cognitive impairment. *Clin Nurs Res* 2021;30:59–69.
- Carpenter J, Connie Ulrich C, Nancy Hodgson N, Hanson L, Ersek M. Alternative consent models in pragmatic palliative care. *J Pain Symptom Manag* 2021;62:183–190.
- Roth D, Huang J, Gitlin L, Gaugler J. Application of randomization techniques for balancing site covariates in the adult day service plus pragmatic cluster-randomized trial. *Contemp Clin Trials Commun* 2020;19:100628.
- Miake-Lye I, Delevan D, Ganz D, Mittman D, Finley E. Unpacking organizational readiness for change: an updated systematic review and content analysis of assessments. *BMC Health Serv Res* 2020;20:106.
- Levy C, Zimmerman S, Mor V, et al. Pragmatic trials in long-term care: Implementation and dissemination challenges and opportunities. *J Am Med Dir Assoc* 2022;23:339–344.

38. American Nurses Credentialing Center. Self-assessment of organizational culture (for long term care). <https://www.nursingworld.org/globalassets/organizational-programs/self-assessment-of-organizational-culture-for-long-term-care.pdf>. Accessed November, 2021.
39. Gitlin L, Marx K, Scerpellac D, et al. Embedding caregiver support in community-based services for older adults: a multi-site randomized trial to test the Adult Day Service Plus Program (ADS Plus). *Contemp Clin Trials* 2019;83:97–108.
40. Washington T, Zimmerman S, Cagle J, et al. Fidelity decision making in social and behavioral research: alternative measures of dose and other considerations. *Soc Work Res* 2014;38:154–162.
41. Curran G, Bauer M, Mittman B, Pyne J, Stetler C. Effectiveness-implementation hybrid designs: combining elements of clinical effectiveness and implementation research to enhance public health impact. *Med Care* 2012;50:217–226.
42. Zimmerman S, Wretman C, Ward K, Tandan M, Sloane P, Preisser J. Fidelity and sustainability of Mouth Care Without a Battle and lessons for other innovations in care. *Geriatr Nurs* 2020;41:878–884.
43. Kolanowski A, Fortinsky R, Calkins M, et al. Advancing research on care needs and supportive approaches for persons with dementia: recommendations and rationale. *J Am Med Dir Assoc* 2018;19:1047–1053.
44. Magaziner J, Zimmerman S, Gruber-Baldini A, Hebel JR, Fox KM. Proxy reporting in five areas of functional status. Comparison with self-reports and observations of performance. *Am J Epidemiol* 1997;146:418–428.
45. Gordon JE, Powell C, Rockwood K. Goal attainment scaling as a measure of clinically important change in nursing home patients. *Age Ageing* 1999;28:275–281.
46. Palmer J, Parker V, Barre L, et al. Understanding implementation fidelity in a pragmatic randomized clinical trial in the nursing home setting: a mixed-methods examination. *Trials* 2019;20:2–10.
47. Iida K, Ryan A, Hasson F, Payne S, McIlfratrick S. Palliative and end-of-life educational interventions for staff working in long-term care facilities: An integrative review of the literature. *Int J Older People Nurs* 2021;16:1–14.
48. Papaioannou A, Loannidis G, McArthur C, et al. Preventing fractures in long-term care: translating recommendations to clinical practice. *J Am Med Dir Assoc* 2021;22:36–42.
49. Seshadri S, Felsen C, Sellers C, Dumyati G. "There is no one to pick up the pieces": sustainability of antibiotic stewardship programs in nursing homes. *Infect Control Hosp Epidemiol* 2021;42:440–447.
50. Resnick B, Aboff-Pretzer I, Galik E, et al. Barriers and benefits to implementing a restorative care intervention in nursing homes. *J Am Med Dir Assoc* 2008;2:103–108.
51. Sloane PD, Hoeffler B, Mitchell CM, et al. Effect of person-centered showering and the towel bath on bathing-associated aggression, agitation, and discomfort in nursing home residents with dementia: a randomized, controlled trial. *J Am Geriatr Soc* 2004;52:1795–1804.
52. Calleson D, Sloane P, Cohen L. Effectiveness of mailing 'Bathing Without a Battle' to all US nursing homes. *Gerontol Geriatr Educ* 2006;27:67–79.
53. Germain A, Markwald R, King E, et al. Enhancing behavioral sleep care with digital technology: study protocol for a hybrid type 3 implementation-effectiveness randomized trial. *Trials* 2021;22:1–14.
54. Ben-Zeev D, Buck B, Meller S, Hudenko W, Hallgren K. Augmenting evidence-based care with a texting mobile interventionist: a pilot randomized controlled trial. *Psychiatr Serv* 2020;71:1218–1224.
55. Shelton R, Brotzman L, Johnson D, Erwin D. Trust and mistrust in shaping adaptation and de-implementation in the context of changing screening guidelines. *Ethn Dis* 2021;31:119–132.
56. Norton W, Chambers D. Unpacking the complexities of de-implementing inappropriate health interventions. *Implement Sci* 2020;15:2.
57. Ojo T, Kabasele L, Boyd B, et al. The role of implementation science in advancing resource generation for health interventions in low- and middle-income countries. *Health Services Insights* 2021;3:1–12.
58. Green HA. community-based evaluation of Screening, Brief Intervention, and Referral to Treatment (SBIRT) for the Black community. *Qual Health Res* 2018;28:418–432.
59. Shepherd V. How nurses can support the inclusion in research of older people who lack capacity to consent. *Nurs Older People* 2021;33:26–31.
60. O'Reilly M, Fetherstonhaugh D, McMaster M, Moyle W, Fielding E, Beattie E. Supporting autonomy of nursing home residents with dementia in the informed consent process. *Dementia* 2019;18:2821–2835.
61. Gustavson A, Drake C, Lakin A, et al. Conducting clinical research in post-acute and long-term nursing home care settings: regulatory challenges. *J Am Med Dir Assoc* 2019;20:798–803.
62. United States Department of Human Health and Services. Assuring compliance with this policy research conducted or supported by any Federal department or agency. 45 CFR. 2009. <https://www.hhs.gov/ohrp/sites/default/files/ohrp/policy/ohrpregulations.pdf>. Accessed November, 2021.