

medBERT.de

Citation for published version (APA):

Bressem, K. K., Papaioannou, J. M., Grundmann, P., Borchert, F., Adams, L. C., Liu, L., Busch, F., Xu, L., Loya, J. P., Niehues, S. M., Augustin, M., Grosser, L., Makowski, M. R., Aerts, H. J. W. L., & Löser, A. (2023). medBERT.de: A comprehensive German BERT model for the medical domain. *Expert Systems with Applications*, 237, Article 121598. <https://doi.org/10.1016/j.eswa.2023.121598>

Document status and date:

Published: 14/09/2023

DOI:

[10.1016/j.eswa.2023.121598](https://doi.org/10.1016/j.eswa.2023.121598)

Document Version:

Publisher's PDF, also known as Version of record

Document license:

Taverne

Please check the document version of this publication:

- A submitted manuscript is the version of the article upon submission and before peer-review. There can be important differences between the submitted version and the official published version of record. People interested in the research are advised to contact the author for the final version of the publication, or visit the DOI to the publisher's website.
- The final author version and the galley proof are versions of the publication after peer review.
- The final published version features the final layout of the paper including the volume, issue and page numbers.

[Link to publication](#)

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal.

If the publication is distributed under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license above, please follow below link for the End User Agreement:


www.umlib.nl/taverne-license

Take down policy

If you believe that this document breaches copyright please contact us at:

repository@maastrichtuniversity.nl

providing details and we will investigate your claim.



medBERT.de: A comprehensive German BERT model for the medical domain

Keno K. Bressen^{a,b,i,1,*}, Jens-Michalis Papaioannou^{c,1}, Paul Grundmann^{c,1}, Florian Borchert^d, Lisa C. Adams^e, Leonhard Liu^c, Felix Busch^a, Lina Xu^a, Jan P. Løyen^a, Stefan M. Niehues^a, Moritz Augustin^f, Lennart Grosser^f, Marcus R. Makowski^e, Hugo J.W.L. Aerts^{g,h,i}, Alexander Löser^c

^a Charité – Universitätsmedizin Berlin, Freie Universität Berlin and Humboldt-Universität zu Berlin, Department of Radiology, Hindenburgdamm 30, 12203 Berlin, Germany

^b Institute of Health at Charité – Universitätsmedizin Berlin, Charitéplatz 1, 10117 Berlin, Germany

^c Berlin University of Applied Sciences and Technology (BHT), Luxemburger Str. 10, 13353 Berlin, Germany

^d Hasso-Plattner Institute for Digital Engineering, Digital Health Center, University of Potsdam, Prof.-Dr.-Helmert-Straße 2-3, 14482 Potsdam, Germany

^e Technical University of Munich, Department of Diagnostic and Interventional Radiology, School of Medicine and Klinikum Rechts der Isar, Ismaninger Str. 22, 81675 Munich, Germany

^f Tiplu GmbH, Karnapp 25, 21079 Hamburg, Germany

^g Artificial Intelligence in Medicine (AIM) Program, Mass General Brigham, Harvard Medical School, 221 Longwood Avenue, Boston, MA 02115, USA

^h Departments of Radiation Oncology and Radiology, Dana-Farber Cancer Institute and Brigham and Women's Hospital, 221 Longwood Avenue, Boston, MA 02115, USA

ⁱ Radiology and Nuclear Medicine, CARIM & GROW, Maastricht University, Minderbroedersberg 4-6, 6211 LK Maastricht, Netherlands

ARTICLE INFO

Keywords:

BERT
Natural language processing
Medicine

ABSTRACT

This paper presents [medBERT.de](https://medbert.de), a pre-trained German BERT model specifically designed for the German medical domain. The model has been trained on a large corpus of 4.7 Million German medical documents and has been shown to achieve new state-of-the-art performance on eight different medical benchmarks covering a wide range of disciplines and medical document types. In addition to evaluating the overall performance of the model, this paper also conducts a more in-depth analysis of its capabilities. We investigate the impact of data deduplication on the model's performance, as well as the potential benefits of using more efficient tokenization methods. Our results indicate that domain-specific models such as [medBERT.de](https://medbert.de) are particularly useful for longer texts, and that deduplication of training data does not necessarily lead to improved performance. Furthermore, we found that efficient tokenization plays only a minor role in improving model performance, and attribute most of the improved performance to the large amount of training data. To encourage further research, the pre-trained model weights and new benchmarks based on radiological data are made publicly available for use by the scientific community.

1. Introduction

Self-supervised pre-training has become a popular approach in natural language processing (NLP) because it allows the creation of high-performance language models. By training a model on a large corpus

of text, the model can learn useful representations of the language. However, the effectiveness of these representations is closely related to the type of data used for pre-training.

When applied to a different type of text, such as a different language, the model may not perform as well, requiring the development of

* Corresponding author.

E-mail addresses: keno-kyrill.bressen@charite.de (K.K. Bressen), michalis.papaioannou@bht-berlin.de (J.-M. Papaioannou), pgrundmann@bht-berlin.de (P. Grundmann), florian.borchert@hpi.de (F. Borchert), lisa.adams@tum.de (L.C. Adams), leonhard.liu@bht-berlin.de (L. Liu), felix.busch@charite.de (F. Busch), lina.xu@charite.de (L. Xu), jan-philipp.loeyen@charite.de (J.P. Løyen), stefan.niehues@charite.de (S.M. Niehues), m.agustin@tiplu.de (M. Augustin), l.grosser@tiplu.de (L. Grosser), marcus.makowski@tum.de (M.R. Makowski), haerts@bwh.harvard.edu (H.J.W.L. Aerts), alooser@bht-berlin.de (A. Löser).

¹ Contributed equally.

specialized models. Consequently, language-specific models, such as those for German (Scheible et al., 2020), French (Martin et al., 2019), or Spanish (Canete et al., 2020; Martin et al., 2019), have been developed and have been successful in improving performance for these languages. However, even within a language, technical languages can be very different from spoken language, necessitating the development of domain-specific models (Gu et al., 2021). Medical terminology is a prime example.

Medical language models that are specifically trained to process and structure medical text have the potential to greatly improve the efficiency and accuracy of medical document analysis.

However, the challenge in training such models lies in the limited availability of relevant text data, especially for languages other than English. In addition, the sensitive nature of medical information often limits the generation of large medical text corpora.

Despite these challenges, the development of medical language models is highly desirable, as they could help process and structure the vast amounts of text generated daily in hospitals.

For English, specialized models for biomedical language processing have already been developed (Gu et al., 2021; Lee et al., 2020). Lee et al. proposed BioBERT (Lee et al., 2020), a version of the BERT (Bidirectional Encoder Representations from Transformers) model trained on English biomedical abstracts, which outperforms previous models on biomedical benchmarks while maintaining the performance of the original BERT model (Devlin et al., 2019). Med-BERT, proposed by (Rasmy et al., 2021), is the first model to be fully trained using hospital data, specifically semi-structured EHRs (electronic health records), resulting in improved performance for downstream prediction models. For non-English languages, the development of specialized models is more difficult due to less available data. Nevertheless, previous work on German medical models has demonstrated the potential of using specialized models for biomedical language processing (Bressem et al., 2020; Frei & Kramer, 2022; Lentzen et al., 11 2022). For example, BioGottBERT (Lentzen et al., 11 2022), which is trained on open medical German texts from Wikipedia and scientific abstracts, could outperform its generalized counterpart (Scheible et al., 2020) on medical tasks. However, these models often suffer from limitations such as limited training data, narrow focus (i.e., focused on only one medical subspecialty), or unrepresentative benchmarks, which limit their comparability.

To address these challenges, our goal was to build a comprehensive German clinical language model - *medBERT.de* - that is trained on a diverse set of medical texts, including scientific texts, medical books, and hospital data from various medical domains. Furthermore, by providing openly available benchmarks using realistic hospital data, we aimed to enable the reproducibility of our results and facilitate comparison with other models.

2. Materials and methods

This study was approved by the local ethics committees of *Charité – Universitätsmedizin Berlin* (EA2/078/22). In accordance with local laws and regulations, written informed consent was not required due to the retrospective design. A total of 4.7 million documents from 11 different sources were included, representing 10 GB of raw data. For details, see Table 1 or Appendix 1.

2.1. Data annotation and benchmarking

2.1.1. Radiology benchmarks

Three medical benchmarks, each based on 2000 radiology reports, were created from radiology reports. Patients that undergo radiological examinations often have a variety of different conditions. For this reason, radiology reports capture a greater diversity of information than, e.g., electronic health records that stem from a single medical field. In this study, we focused on three tasks. The first task is a classification task

Table 1
Datasources.

Source	No. Documents	No. Sentences	No. Words	Size (MB)
DocCheck Flexikon	63,84	720,404	12,299,257	92
GGPOnc 1.0 (Borchert et al., 2022)	4,369	66,256	1,194,345	10
Webcrawl (Shrestha, 2021)	11,322	635,806	9,323,774	65
PubMed abstracts	12,139	108,936	1,983,752	16
Radiology reports	3,657,801	60,839,123	520,717,615	4,195
Spinger Nature	257,999	14,183,396	259,284,884	1,986
Electronic health records (Schmidt et al., 2021)	373,421	4,603,461	69,639,020	440
Doctoral theses	7,486	4,665,850	90,380,880	648
Thieme Publishing Group	330,994	10,445,580	186,200,935	2,898
Wikipedia	3,639	161,714	2,799,787	22
Summary	4,723,010	96,430,526	1,153,824,249	10,372

based on short text reports from chest X-rays. The second task, also a classification task, is based on longer reports of CT (computed tomography) chest examinations. The third task is a named entity recognition task (NER), that is based on medium-sized reports of CT/X-ray examinations of the wrist. All of these reports were obtained from a large, level 1 hospital in Germany and cover a wide range of bone, lung, heart, and vascular diseases.

For the first medical benchmark, a multi-label text classification task, three board-certified radiologists (KKB, LCA, SMN) manually labeled 2000 reports for the global presence/absence of four pathologies and four different types of therapy devices. The second benchmark, a text classification task, contains 2000 reports of computed tomography (CT) scans. All reports had to include the chest but could include additional body parts (head, neck, abdomen). All studies were labeled by a final-year medical student (LX) for the presence/absence of 23 different chest pathologies and were then reviewed by two board-certified radiologists (LCA, KKB).

For the third benchmark, a named entity recognition task, 2000 reports were labeled by a final-year medical student (JPL) for the presence/absence of 42 labels. The labels were then verified by two board-certified radiologists (LCA, KKB). Details of each label are provided in Appendix 3.

Benchmarks and training code can be found at: <https://github.com/DATEXIS/medBERT.de>.

2.1.2. Open medical benchmarks

As an additional openly available benchmark task, we considered the prediction of named entities in GGPOnc 2.0, a corpus of German clinical practice guidelines in oncology and the largest freely distributable data set of semantically annotated German medical texts (Borchert et al., 2022). Seven medical students (all passed their first medical exam) annotated more than 200 K mentions of clinically named entities. For the benchmark, we considered the most challenging setting with 8 fine-grained semantic classes and long entity spans. We used the same training/development/test splits as in the original baseline NER experiments. We also considered GraSCCo, a synthetic corpus of 62 clinical case reports, as a benchmark (Modersohn et al., 2022). Since the initial release of GraSCCo does not include semantic annotations, a single annotator from the GGPOnc annotation team created named entity annotations using the original GGPOnc annotation scheme and instructions. This resulted in 5.8 K entity annotations in the long/fine setting.

2.1.3. Private medical benchmarks

A significant proportion of radiology reports in the training corpus can lead to overfitting, hindering the model's ability to generalize to new clinical scenarios. Therefore, the benchmarks based on radiological texts alone may not be sufficient to assess how well the models would perform on new types of clinical data. To mitigate this, we developed benchmarks using new, unseen clinical records, namely surgical reports and discharge summaries. Both types of clinical records were not previously included in the training data. The benchmarks were multi-label classification tasks, where the model had to predict multiple diagnoses (ICD-10) or procedure (OPS) codes from the text, allowing a thorough evaluation of its performance in real-world situations.

Discharge notes and operative reports are more challenging benchmarks than radiology reports because they tend to be longer and more complex. In addition, the information contained in discharge summaries and operative reports differs from that found in radiology reports. While radiology reports focus primarily on describing anatomy, pathological findings, and formulating a diagnosis, discharge notes and operative reports provide a broader context and include information about the patient's condition, treatment, medications, and follow-up plans.

Discharge summaries, surgery reports, as well as all ICD-10 and OPS codes were extracted from the hospital information system of the same level I hospital as before. For these tasks, no manual labeling was performed. Instead, we assigned to the surgery reports as labels all OPS codes of the same patient that matched the date of the text document. Furthermore, we restricted the codes to the surgery chapter of the OPS system.² For the discharge summaries, we assigned all codes (in one task diagnoses as ICD-10, in another task procedures as OPS) of the patient as labels. For each of these tasks, we included as labels the most frequent codes, such that the test set consisted of at least 10 examples for each label.

Since all three tasks involve automatic and, particularly for the discharge summaries tasks, only approximately accurate labels (e.g., codes are present but the information could be described elsewhere in the respective EHR), performance even for perfect pattern recognition is not expected to be very high - by construction. Thus, the results of these tasks should not be interpreted in an absolute manner but rather relatively by comparing the performance between models.

The OPS code 5-984 ("Microsurgical technique") is the most common label used for surgery reports, 8-930 ("Monitoring of respiration, heart, and circulation without measurement of pulmonary artery pressure and central venous pressure") for OPS-labeled discharge summaries, and the ICD code Z11 ("Special procedures for testing for infectious and parasitic diseases") for ICD-10-labeled discharge summaries.

2.1.4. General domain benchmark

Ultimately, we included GermEval18 (Risch et al., 2018), a widely adopted standard benchmark in the general domain for evaluating German language models. GermEval18 serves as a classification task specifically designed to assess the performance of models on offensive text messages extracted from social media platforms. This dataset is commonly utilized to evaluate the proficiency of German language models.

2.2. Data anonymization

Radiology reports extracted from our hospital database typically do not contain identifiable information such as name or date of birth. However, in rare cases, this information may have been added by the radiologist. Therefore, we used a named entity recognition model for the German language to identify all patient names in the (Akbik et al., 2018)

data set. Identified names were then manually verified by two radiologists (KKB, LCA) and subsequently removed from the text. In addition, dates were removed from the text and replaced with wildcards. For benchmarks, each document was reviewed at least three times by the authors (KKB, LCA, JMP, PG) to ensure that no identifiable information was present.

2.3. Architectures/models

To evaluate the performance of our proposed model, we compared six different models based on the BERT or RoBERTa architecture. We justify our model choice given the amount of training data available (Kaplan et al. 2020). Training a larger model would exceed the available domain specific pre-training data. Therefore, we settled on a BERT base sized model and compared against similar parametrized models, one general German language model, one multilingual model that includes German training data, two German medical models, and two versions of our pre-trained BERT model:

- GottBERT, which is the current state-of-the-art RoBERTa-based model for German text (Liu et al., 2019; Scheible et al., 2020). GottBERT was pre-trained on the German portion of the OSCAR dataset (Suárez et al. 2020), resulting in 145 GB of text containing approximately 21.5 billion words.
- A multilingual BERT-based model (Devlin et al., 2019)
- BioGottBERT, a version of GottBERT fine-tuned to medical texts and German Wikipedia (Lentzen et al., 11 2022)
- German-MedBERT, a version of the German BERT (Chan et al., 2020) fine-tuned on a crawl of German medical websites.
- *medBERT.de*, our model pre-trained on 4.7 Million German medical texts
- *medBERT.de_{dedup}*, a variant of *medBERT.de* trained on a slightly smaller corpus where duplicated radiology reports had been removed

2.4. Deduplication

Radiology reports are often written in a semi-structured form with very similar sentences. Because of this repetition, the information content of many documents is lower in terms of semantic concepts than other data sources used.

Language models tend to quickly overfit due to these data-inherent properties. A common strategy to counteract this behavior is to deduplicate the pre-training data (K. Lee et al., 2021). Therefore, we measured the cosine distance between all reports by encoding them as bag of word representations. We only kept documents for which there is no other document with a similarity greater than 0.75. Due to computational restriction, we had to limit this approach to short reports only. Still, using this approach, we reduced the number of radiology reports for pre-training from 4,504,167 to 3,657,801 reports. To evaluate the impact of deduplication, we report the performance of the two BERT models trained with and without deduplicated data.

2.5. Hyperparameters/pre-training details

We pre-trained the model using the Lamb optimizer (You et al., 2019). As usual for BERT-based models, we trained the model in two phases, the first with a maximum sequence length of 128 and the second with a sequence length of 512. To pre-train our model from scratch, we used the hyperparameters given by (You et al., 2019). We initialized our model using the default Kaiming uniform distribution in PyTorch. We used a normal distribution with a standard deviation of 0.02 for the embedding layer of the model. In the first phase, we used a learning rate of $6e^{-3}$ and a batch size of 65,536 with 2,000 warm-up steps and a polynomial decaying learning rate for a total of 7,038 steps. In the second phase, we trained with a batch size of 32,768 and a maximum

² <https://www.dimdi.de/static/de/klassifikationen/ops/kode-suche/opshtm12022/chapter-5.htm>.

learning rate of $4e^{-3}$, 200 warm up steps and a total of 1,563 steps. We removed very rare Unicode characters that appear less than three times from our pre-training data. This allows the tokenizer vocabulary to contain more specific sub-words and removes unnecessary tokens from the vocabulary that have an impact on the memory footprint. In addition, we set the number of occurrences required for a word to be included in the vocabulary to 20 to avoid including patient names in the vocabulary that may have been missed during anonymization.

3. Experimental design

We performed a hyperparameter optimization on all downstream tasks with median pruning on either the area under the receiver operating characteristic curve (AUROC, classification tasks) or token F1 (NER tasks) with a threshold of 0.5. We did not perform any calibration for any of the models for the calculation of Precision and Recall. For each model-task combination, we performed 100 runs and tune the learning rate, the batch size, and the number of warm-up steps. We tune the learning rate from $1e^{-5}$ to $1e^{-4}$ with logarithmic sampling. Also, we optimize the number of warmup steps where we sample from 0 to 10,000. Lastly, we tune the batch size, selecting one of three options: 8, 16, or 32. Finally, we evaluated the best performing models for which the hyperparameters are reported in [Appendix 4](#).

4. Results

4.1. Radiology benchmarks

In the chest x-ray task, we found that the two best performing models were our own pre-trained BERT models (s. [Table 2](#) and [Fig. 1](#)). Our model trained on the corpus with duplicates removed (medBERT.de_{dedup}) achieves a slightly better performance with an average AUROC of 83.65 compared to 83.42 of the model trained on the whole corpus (medBERT.de). The third-best performance was achieved by GottBERT with a mean AUROC of 83.48, which also outperformed other medical models.

In the chest CT task, our model trained on the whole corpus performed best with an AUROC of 96.69, closely followed by medBERT.de_{dedup} (AUROC 96.39). Other models showed a significantly lower performance of 19%. GottBERT and BioGottBERT both perform similarly with mostly sub-percentage differences in all metrics except precision.

For the NER task, medBERT.de_{dedup} showed the best performance in all metrics except for global Recall. However, the scores of all models were in a similar range, with mostly 1–3% differences between the best and worst-performing models.

The results suggest that the advantage of domain-specific models is more pronounced for longer texts. On the X-ray (98 ± 27 words) and NER (108 ± 41 words) tasks, which consist of short reports of only a few sentences, the difference between the models is not as pronounced as on the CT reports, which are considerably longer (258 ± 100 words).

[Tables 2 and 3](#) provide an overview of all tasks and metrics. Detailed metrics for each class on the radiology benchmarks can be found in [Appendix 2](#).

4.2. Open medical benchmarks

The GGPOnc benchmark ([Borchert et al., 2022](#)), a German corpus based on clinical practice guidelines for oncology, and the GraSCCo benchmark ([Modersohn et al., 2022](#)), a corpus consisting of artificially generated electronic health records for various diseases, were used for this comparison.

On GGPOnc, our models achieved higher AUROC, precision (global and token-level) and token-level recall than the other models, while BioGottBERT achieved the highest macro F1 score and recall. Deduplication of training data did not seem to have a positive impact on model

Table 2

Classification Tasks.

Model	AUROC	Macro F1	Micro F1	Precision	Recall
Chest CT					
GottBERT	92.48	69.06	83.98	76.55	65.92
BioGottBERT	92.71	69.42	83.41	80.67	65.52
Multilingual BERT	91.90	66.31	80.86	68.37	65.82
German-MedBERT	92.48	66.40	81.41	72.77	62.37
medBERT.de	96.69	81.46	89.39	87.88	78.77
medBERT.de _{dedup}	96.39	78.77	89.24	84.29	76.01
Chest X-Ray					
GottBERT	83.18	64.86	74.18	59.67	78.87
BioGottBERT	83.48	64.18	74.87	59.04	78.90
Multilingual BERT	82.43	63.23	73.92	56.67	75.33
German-MedBERT	83.22	63.13	75.39	55.66	78.03
medBERT.de	84.65	67.06	76.20	60.44	83.08
medBERT.de _{dedup}	84.42	66.92	76.26	60.31	82.99
ICD-10 code classification on discharge notes					
GottBERT	77.23	18.32	51.23	38.30	14.27
BioGottBERT	78.01	17.96	50.56	35.97	13.95
Multilingual BERT	76.64	19.48	51.19	38.39	15.60
German-MedBERT	75.44	23.41	53.63	41.39	18.94
medBERT.de	80.78	23.41	53.84	41.42	18.75
medBERT.de _{dedup}	80.84	21.44	52.46	40.45	17.04
OPS code classification on discharge notes					
GottBERT	71.37	16.46	39.54	29.63	13.02
BioGottBERT	69.90	15.97	38.60	35.06	12.23
Multilingual BERT	70.39	15.53	39.16	27.94	12.35
German-MedBERT	71.79	15.90	38.22	29.30	12.76
medBERT.de	76.83	20.48	44.96	33.82	16.57
medBERT.de _{dedup}	77.07	21.33	46.33	39.72	16.83
OPS code classification on surgery reports					
GottBERT	89.17	52.58	64.25	61.42	48.08
BioGottBERT	92.88	54.82	66.27	61.38	50.98
Multilingual BERT	91.28	65.42	72.92	71.74	61.78
German-MedBERT	92.50	55.21	66.32	62.37	51.24
medBERT.de	94.36	66.39	73.95	69.67	64.25
medBERT.de _{dedup}	93.44	65.28	73.87	73.19	61.09
GermEval-18					
GottBERT	86.07	79.19	81.11	80.10	78.58
BioGottBERT	82.36	76.57	78.95	77.72	75.88
Multilingual BERT	83.65	76.76	78.63	77.09	76.49
German-MedBERT	84.72	77.85	79.60	77.69	78.02
medBERT.de	81.45	74.84	77.65	76.77	73.97
medBERT.de _{dedup}	82.05	74.56	77.40	76.22	73.75

performance, as medBERT.de consistently achieved higher metrics than medBERT.de_{dedup}.

On GraSCCo, GottBERT and medBERT.de showed the best performance. medBERT.de had the highest AUROC (85.14), AUROC_{tok} (75.17), and Recall_{tok} (75.17), while GottBERT had the best performance on all other metrics. However, the overall difference between the models is small.

4.3. Private medical benchmarks

Three medical benchmarks were constructed using discharge letters and surgical reports, which are not publicly available for privacy reasons. The discharge summaries and surgical reports included were anonymized texts randomly sampled from a larger, single hospital, database containing records from an array of medical and surgical subspecialties (e.g., cardiology, nephrology, ophthalmology, orthopedic surgery or rheumatology). Given the anonymization prior to sampling, there was no control over the distribution of different specialties in each benchmark. Each benchmark consisted of 2000 texts, which were stratified into a training set of 1000 texts, a validation set of 500 texts, and a test set of 500 texts.

The first task is to predict 65 different ICD-10 codes from discharge summaries. Both of our models showed superior performance on this task, outperforming all other models. The best model is medBERT.de, although the difference to medBERT.de_{dedup} was in the sub-percentage range.




Fig. 1. AUROC performance on classification tasks. Fig. 1 compares the areas under the receiver operating characteristics curve (AUROC) achieved by the different evaluated models on our clinical text classification benchmarks. It shows, that both [medBERT.de](#) and [medBERT.de dedup](#) consistently achieve a higher AUROC than the other models. However, there seems to be now advantage in deduplication of the training data, as the difference between [medBERT.de](#) and [medBERT.de dedup](#) is small. CT = Computed Tomography, ICD-10 = International Classification of Diseases 10th edition, OPS = German procedure classification (Operationen- und Prozedurenschlüssel).

Table 3
NER Tasks.

Model	AUROC	AUROC _{tok}	F1 _{mac}	F1 _{tok}	Prec	Prec _{tok}	Rec	Rec _{tok}
WristNER								
GottBERT	86.24	86.31	55.22	52.94	62.26	60.45	56.18	54.17
BioGottBERT	87.40	87.48	54.89	52.53	61.39	59.37	57.15	54.74
Multilingual BERT	87.29	87.24	54.63	53.12	62.95	61.53	55.09	53.43
German MedBERT	88.25	88.31	53.40	51.26	61.33	59.77	54.54	52.50
medBERT.de	86.84	86.88	59.11	58.68	65.66	65.39	60.59	60.06
medBERT.de dedup	87.45	87.40	58.15	57.03	62.88	61.87	60.26	59.11
GraSCCo								
GottBERT	84.83	84.81	76.00	75.75	76.72	76.50	75.42	75.06
BioGottBERT	84.82	84.56	75.52	74.95	76.37	75.96	74.84	74.08
Multilingual BERT	83.90	83.73	72.54	71.98	72.93	72.56	72.22	71.44
German-MedBERT	84.03	83.95	73.14	72.85	73.05	72.86	73.41	73.00
medBERT.de	85.14	85.07	75.60	75.46	75.91	75.78	75.33	75.17
medBERT.de dedup	84.89	84.83	75.20	75.25	75.85	75.97	74.72	74.66
GGPOnc								
GottBERT	98.01	98.07	75.18	73.43	76.40	74.96	74.43	72.42
BioGottBERT	97.96	98.05	76.07	74.62	77.06	75.45	75.60	74.18
Multilingual BERT	97.81	97.95	73.75	72.20	75.47	73.93	72.95	71.11
German-MedBERT	97.55	97.66	74.48	72.45	75.59	73.73	73.76	71.57
medBERT.de	98.20	98.35	75.93	75.12	77.37	76.33	75.15	74.45
medBERT.de dedup	98.10	98.22	75.57	74.92	76.74	75.98	75.05	74.43

The second task is to predict 49 OPS codes from discharge summaries. Again, [medBERT.de](#) and [medBERT.de dedup](#) performed better than all other models. However, the overall performance of all models was below average, with particularly low scores for the F1 measure and recall.

In the third task, the classification of 10 OPS codes from surgical reports, our domain-specific models again showed the best performance. Compared to the other two tasks, the overall scores were also higher. We attribute this to the fact that this task is less complex since surgical reports are generally shorter and have less variability than discharge

summaries. In addition, the number of labels was reduced, which further contributed to the reduced difficulty.

4.4. General domain benchmarks

In addition to the medical and radiological benchmark tasks, we also evaluated performance on the GermEval18 ([Risch et al., 2018](#)). We found that GottBERT outperformed all domain-adapted models in all evaluated metrics.

We also observed that the [medBERT.de](#) models were outperformed

by the BioGottBERT model. This can be attributed to the fact that the pre-training corpus for GottBERT contains a greater amount of general domain knowledge and language, giving the model an advantage in general domain tasks. In addition, BioGottBERT, which is based on GottBERT, may have the same advantage but may be affected by catastrophic forgetting because it was only further pre-trained on corpora from the medical domain.

4.5. Tokenizer fertility

Rust et al. suggested that a tokenizer that produces fewer sub-words per word may improve performance due to better-developed embeddings (Rust et al., 2021). Therefore, we measured tokenizer fertility, which measures the average number of sub-words per tokenized word for all of our evaluated models. For the evaluation, we used the text data from our chest CT, chest x-ray classification, and wrist NER tasks.

As expected, we measured the lowest fertility for the tokenizer of the medBERT.de model, which was trained on data following a similar distribution.

We found that tokenizer fertility did not necessarily correlate with improved performance. For example, our medBERT.de models both had the lowest fertility score of 1.18 (see Table 4) and were the best-performing models on the chest CT and x-ray as well as the wrist NER task. However, it was closely followed by GottBERT, which had a much higher fertility score of 1.75, but performed similarly well. Furthermore, we observed that deduplication of the training data had almost no effect on the fertility of the medBERT.de model.

5. Discussion

In this study, we trained a domain-specific German BERT model on a large data set of German medical texts, including articles, papers, and electronic medical records. We then fine-tuned the model on various medical benchmarks and found that it outperformed both general domain language models and other medical domain models, demonstrating the model's ability to capture the unique characteristics and terminology of German medical language more effectively than general German models. Our results highlight the advantages of using domain-specific language models for the German medical language but also the importance of a large training corpus. In line with previous research (Pérez-Mayos et al., 2021), we attribute the superior performance of our model on medical benchmarks to the larger amount of data used in training compared to German-MedBERT or BioGottBERT. However, our results also suggest that data for pre-training or fine-tuning domain-specific models should not consist solely of specialized language, as this may negatively affect the model's performance on general tasks. This was demonstrated by the Germeval18 task, where GottBERT outperformed BioGottBERT, even though the two models had identical architectures and BioGottBERT was initialized with GottBERT's weights. This performance difference can be attributed to the fact that BioGottBERT was trained on the entire German Wikipedia, which contains general domain language.

In addition, we observed a drastically improved performance on the OPS and ICD code classification tasks compared to all other domain-specific and general domain models. This indicates that our models

are able to generalize to different subdomains that are not included in the pre-training data. The results, therefore, suggest that the models are able to capture relevant clinical and medical concepts, thus demonstrating their ability to adapt to the different clinical language used in discharge notes.

In contrast to (Rust et al., 2021), we did not observe a direct correlation between the fertility of the tokenizer and the performance of the model on downstream tasks. This suggests that fertility alone is not a predictive measure of a model's performance on specialized downstream tasks. Nevertheless, it is likely that the fertility of the tokenizer played a role in the model's performance on tasks involving longer texts, particularly on clinical benchmarks based on discharge notes and surgical reports. Since the texts for these benchmarks were truncated to fit into 512 tokens, some information may have been lost in the process. A more efficient tokenizer may be able to encode more information, potentially improving the model's performance on these tasks.

In our study, we found a mixed impact of data deduplication. While earlier research suggested benefits from deduplication (Lee et al., 2021), we did not see a consistent improvement in performance with our model (medBERT.de) compared to the deduplicated version (medBERT.de_{dedup}). While on certain benchmarks, our medBERT.de performed better than the deduplicated version, on others it performed worse. This discrepancy could be due to the fact that our deduplication process was not as extensive, as it was only applied to short reports. Furthermore, we have not yet applied deduplication to non-radiological text, which might contain duplicates.

6. Limitations

A limitation of our study is that about 40% of the data consists of radiology reports, which may differ in style from other types of electronic health records. In addition, certain medical specialties, such as ophthalmology and pathology, are underrepresented in our sample due to their limited use of radiological imaging. On the other hand, other specialties, such as psychiatry, may be underrepresented because the conditions they treat are not typically seen on imaging. It is also worth noting that the texts were collected from a single university hospital, and it is possible that the performance of our model on new data may be affected by differences in reporting styles between institutions. We suspect that our training data does not sufficiently capture the semantic information needed to improve performance compared to a language model trained on a general domain corpus. This can be partly explained by the repetitive nature of radiology reports.

7. Conclusion and outlook

In conclusion, this study has shown the benefits of using a domain-specific German BERT model, trained on a large data set of German medical texts, for tasks related to the German medical language. The model achieved superior performance compared to the general domain and other medical domain models, underlining the value of using domain-specific models. However, to further improve performance, a future German clinical language model should be trained on a more diverse data set, e.g., including discharge summaries from a broad range of medical specialties. Nevertheless, this model represents a new state-of-the-art for the German clinical language, outperforming GottBERT.

CRediT authorship contribution statement

Keno K. Bressem: Conceptualization, Methodology, Software, Validation, Validation, Investigation, Resources, Data curation, Writing – original draft, Writing – review & editing, Project administration. **Jens-Michalis Papaioannou:** Methodology, Software, Validation, Formal analysis, Investigation, Writing – original draft, Writing – review & editing. **Paul Grundmann:** Methodology, Software, Validation, Formal analysis, Investigation, Writing – original draft, Writing – review

Table 4
Fertility Table.

Model	Fertility
GottBERT	1.75
BioGottBERT	1.75
Multilingual BERT	1.97
German-MedBERT	1.86
medBERT.de	1.18
medBERT.de _{dedup}	1.18

& editing. **Florian Borchert**: Data curation, Writing – review & editing. **Lisa C. Adams**: Data curation, Writing – original draft, Writing – review & editing. **Leonhard Liu**: Software, Writing – review & editing. **Felix Busch**: Data curation, Writing – review & editing. **Lina Xu**: Data curation, Writing – review & editing. **Jan P. Loyen**: Data curation, Writing – review & editing. **Stefan M. Niehues**: Resources, Data curation, Writing – review & editing. **Moritz Augustin**: Validation, Writing – review & editing. **Lennart Grosser**: Validation, Writing – review & editing. **Marcus R. Makowski**: Resources, Writing – review & editing. **Hugo J. W.L. Aerts**: Resources, Writing – review & editing. **Alexander Löser**: Conceptualization, Resources, Writing – review & editing, Supervision, Project administration.

Declaration of Competing Interest

The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

Appendix

Appendix 1. – Details on data sources

DocCheck Flexikon

The DocCheck Flexikon (<https://flexikon.doccheck.com/>) is an open wiki dealing with medical topics. It contains overview articles about diseases, diagnostic procedures, or treatments in all areas of medicine. This study includes all articles of the Flexikon that have been published until January 1st, 2022. In addition, entries from the DocCheck forum and product descriptions from the medical store were included. This resulted in 63,884 documents (92 MB of raw text).

GGPOnc

GGPOnc is a freely available German language corpus based on clinical practice guidelines for oncology with expert annotations. It is available at (Borchert et al., 2020, 2022).

Webcrawl

A webcrawl of several German medical forums was performed, as described in (Shrestha, 2021). The webcrawl consisted of 11,322 documents (65 MB of raw text).

Pubmed abstracts

We crawled PubMed (<https://pubmed.ncbi.nlm.nih.gov>) for German publications with openly available German abstracts published by September 1st 2022. This resulted in 12,139 documents, representing 16 MB of raw text.

Radiology reports

All radiology reports created between January 1st, 2009, and December 31st, 2021, were extracted from the Radiology Information System of Charité - Universitätsmedizin Berlin. After removing texts with less than 100 characters, we performed a similarity analysis to exclude duplicate reports. All remaining documents were then anonymized. This resulted in 3.66 million radiology reports that were included in the training corpus (4,195 MB of raw text).

Springer nature corpus

Using the Springer Nature API, we identified German-language, open-access publications from the Springer Nature group. Abstracts and full text of the publications were extracted and added to the training corpus. In total, this involved 257,999 documents and 1,986 MB of data.

Thieme publishing group corpus

With the permission of the Thieme Publishing Group, medical textbooks, licensed by the Charité, and journals for continuing medical education (including the Up2Date series, available at <https://www.thieme.de/de/aerzte-in-weiterbildung/up2date-fachzeitschriften-20280.htm>) were included in the training corpus. After cleaning the texts by removing figures and tables, this resulted in 330,994 documents and 2,898 MB of raw text.

Electronic health records

We included 373,421 electronic health records (EHR) from the TBase database from the Department of Nephrology and the Center for Kidney Transplantation at Charité - Universitätsmedizin Berlin (Schmidt et al., 2021). These included physician letters, microbiology reports, pathology reports, and diagnostic procedure reports (440 MB of raw text).

PhD theses

In this study, we used a data set of 7,481 open-access German medical dissertations and postdoctoral theses from the Charité - Universitätsmedizin

Data availability

Radiology benchmarks are openly available. Pretraining data cannot be shared because of privacy concerns

Acknowledgements

We would like to thank DocCheck AG and Thieme Medical Publishers for their assistance with data collection. We would also like to thank Manjil Shrestha and Rolf Becker for providing additional training data from their German MedBERT model. We also thank the Scientific Computing of the IT Division at the Charité – Universitätsmedizin Berlin for providing computational resources that contributed to the research results reported in this paper. KKB is grateful for his participation in the BIH Charité Digital Clinician Scientist Program funded by Charité-Universitätsmedizin Berlin and the Berlin Institute of Health.

Berlin available at <https://refubium.fu-berlin.de/handle/fub188/13>. We cleaned the data by removing sentences that did not contain German stop words and excluded theses with a length of fewer than 15 pages. This process ensured that only relevant, high-quality information was included in our analysis. In total, 648 MB of data was added to the training corpus.

Wikipedia

Entries from the German Wikipedia dealing with medical topics were extracted and added to the training corpus. There were 3,639 texts, corresponding to 22 MB.

Appendix 2. – Detailed metrics per class for benchmarks based on radiology reports

In evaluating the performance of our models on radiology report benchmarks, we analyzed detailed metrics per class to provide a deeper understanding of the model’s performance on specific classes within the data set. The metrics included precision, recall, F1 score, and AUROC, which were calculated for each class individually, allowing for a more nuanced evaluation of the model’s performance and potential shortcomings.

Per Class Metrics for X-Ray Report Classification Task

Class	AUROC	F1	Precision	Recall	AUROC	F1	Precision	Recall
	GottBERT				BioGottBERT			
Congestion	65.85	36.23	27.12	54.55	66.49	35.63	27.67	50.00
Effusion	76.30	60.20	50.43	74.68	77.11	61.27	51.05	76.58
Opacity	66.20	45.36	36.07	61.11	68.75	47.60	36.40	68.75
Pneumothorax	84.85	25.12	14.53	92.86	84.01	24.06	13.83	92.86
Gastric tube	89.92	76.57	70.53	83.75	91.06	81.10	72.20	92.50
Thoracic drain	94.00	85.87	82.56	89.44	93.46	84.46	79.13	90.56
Tracheal tube	89.19	84.58	82.63	86.64	87.56	84.93	82.20	87.85
Venous catheter	92.79	91.57	91.30	91.84	91.59	91.91	91.12	92.71
Misplaced	89.52	78.26	81.82	75.00	91.26	66.67	77.78	58.33
	Multilingual BERT				German MedBERT			
Congestion	66.31	37.61	30.14	50.00	66.18	36.63	27.03	56.82
Effusion	77.96	61.58	52.70	74.05	76.11	61.73	51.71	76.58
Opacity	64.80	41.90	35.05	52.08	67.57	49.23	39.02	66.67
Pneumothorax	82.14	23.45	14.53	60.71	82.65	22.89	13.77	67.86
Gastric tube	88.31	79.89	71.43	90.62	91.39	80.65	71.50	92.50
Thoracic drain	90.93	84.78	82.98	86.67	93.05	85.57	79.81	92.22
Tracheal tube	88.54	84.89	80.43	89.88	87.75	85.60	82.40	89.07
Venous catheter	91.16	90.41	90.14	90.67	92.07	92.53	91.22	93.88
Misplaced	91.71	64.52	52.63	83.33	92.21	53.33	44.44	66.67
	medBERT.de				medBERT.de _{dedup}			
Congestion	65.02	37.04	27.47	56.82	68.90	38.55	28.34	60.23
Effusion	78.65	61.58	51.49	76.58	78.54	61.78	52.68	74.68
Opacity	70.88	49.25	38.58	68.06	67.00	48.47	38.31	65.97
Pneumothorax	86.33	24.64	14.21	92.86	84.90	25.84	14.92	96.43
Gastric tube	90.88	80.55	71.71	91.87	91.07	80.11	72.59	89.38
Thoracic drain	94.49	88.08	82.52	94.44	94.10	88.31	82.93	94.44
Tracheal tube	90.19	86.55	83.46	89.88	90.30	86.33	83.40	89.47
Venous catheter	93.45	92.53	91.22	93.88	93.05	92.87	92.73	93.00
Misplaced	91.93	83.33	83.33	83.33	91.93	80.00	76.92	83.33

Per Class Metrics for CT Report Classification Task

Class	AUROC	F1	Precision	Recall	AUROC	F1	Precision	Recall
	GottBERT				BioGottBERT			
Aortic dissection	88.71	0	0	0	98.99	0	0	0
Bone tumors	75.26	0	0	0	76.80	14.81	40.00	9.09
Bronchial abnorm.	93.18	82.24	86.27	78.57	98.04	81.36	77.42	85.71
Congestion	93.55	76.92	88.24	68.18	87.18	41.38	85.71	27.27
Effusion	99.10	97.16	96.86	97.47	99.28	97.79	97.48	98.10
Emphysema								
Lung	96.67	92.68	92.68	92.68	95.42	92.02	92.59	91.46
Soft tissue	84.08	57.89	64.71	52.38	89.48	45.71	57.14	38.10
Enlarged Heart	96.65	91.30	91.30	91.30	93.50	80.99	94.23	71.01
Fibrosis	93.53	86.75	94.74	80.00	92.03	82.35	87.50	77.78
Fractures	92.38	73.77	80.36	68.18	93.08	77.17	80.33	74.24
Hiatal hernia	99.67	77.78	87.50	70.00	98.44	80.00	80.00	80.00
Lung tumor	94.69	88.37	90.05	86.76	95.12	88.04	87.05	89.04
Lymphadenopathy								
All	95.69	92.24	89.54	95.11	93.03	90.64	86.94	94.67
Malignant	88.88	68.12	83.93	57.32	87.23	66.67	84.91	54.88
Mediastinal tumor	78.50	0	0	0	81.03	27.59	57.14	18.18
Pericard. effusion	97.96	88.89	90.32	87.50	97.41	93.55	96.67	90.62
Pleural abnorm.	72.27	25.00	100.00	14.29	79.55	44.44	100.00	28.57
Pneumonia	94.10	84.25	81.56	87.12	95.51	86.14	85.19	87.12
Pneumothorax	97.23	87.50	93.33	82.35	97.03	81.36	96.00	70.59
Pulm. embolism	99.48	90.62	90.62	90.62	99.48	84.06	78.38	90.62
Therapy devices	97.47	92.04	95.68	88.67	95.33	90.72	93.62	88.00

(continued on next page)

(continued)

Class	AUROC	F1	Precision	Recall	AUROC	F1	Precision	Recall
Ventilation	96.50	94.02	94.02	94.02	96.21	94.05	93.55	94.57
Others	93.02	29.63	57.14	20.00	90.03	32.00	80.00	20.00
Multilingual BERT				German-MedBERT				
Aortic dissection	99.40	0	0	0	97.08	0	0	0
Bone tumors	67.18	0	0	0	73.90	0	0	0
Bronchial abnorm.	94.26	80.39	89.13	73.21	95.52	80.00	81.48	78.57
Congestion	87.23	45.71	61.54	36.36	96.58	66.67	85.71	54.55
Effusion	98.63	94.86	90.75	99.37	98.97	96.53	96.23	96.84
Emphysema								
Lung	95.74	89.16	88.10	90.24	96.12	92.59	93.75	91.46
Soft tissue	84.70	57.78	54.17	61.90	86.48	52.94	69.23	42.86
Enlarged Heart	97.44	86.11	82.67	89.86	91.71	70.09	85.42	59.42
Fibrosis	96.44	77.42	75.00	80.00	96.36	82.50	94.29	73.33
Fractures	90.30	70.15	69.12	71.21	91.48	72.57	87.23	62.12
Hiatal hernia	98.87	70.00	70.00	70.00	99.92	90.00	90.00	90.00
Lung tumor	94.13	86.88	82.11	92.24	93.67	86.71	88.57	84.93
Lymphadenopathy								
All	93.72	90.83	87.30	94.67	95.10	91.70	90.13	93.33
Malignant	88.40	67.11	74.63	60.98	88.40	59.50	92.31	43.90
Mediastinal tumor	73.64	25.00	40.00	18.18	76.49	0	0	0
Pericard. effusion	97.34	86.15	84.85	87.50	93.64	85.71	100.00	75.00
Pleural abnorm.	80.87	16.67	20.00	14.29	70.51	0	0	0
Pneumonia	92.44	77.52	68.00	90.15	95.18	83.97	84.62	83.33
Pneumothorax	98.92	88.89	96.55	82.35	97.31	85.71	93.10	79.41
Pulm. embolism	99.22	77.61	74.29	81.25	99.01	84.06	78.38	90.62
Therapy devices	95.74	87.37	89.51	85.33	95.69	86.52	92.42	81.33
Ventilation	93.84	89.30	87.89	90.76	96.93	94.28	94.54	94.02
Others	88.13	51.61	72.73	40.00	95.06	51.61	72.73	40.00
medBERT.de				medBERT.de _{dedup}				
Aortic dissection	80.65	66.67	100.00	50.00	67.44	0	0	0
Bone tumors	94.10	64.86	80.00	54.55	97.19	64.86	80.00	54.55
Bronchial abnorm.	96.70	91.74	94.34	89.29	98.16	85.98	90.20	82.14
Congestion	98.60	72.22	92.86	59.09	95.02	72.22	92.86	59.09
Effusion	99.62	97.50	96.30	98.73	99.17	97.14	97.45	96.84
Emphysema								
Lung	97.56	95.06	96.25	93.90	97.96	93.83	95.00	92.68
Soft tissue	91.66	54.55	75.00	42.86	92.12	63.16	70.59	57.14
Enlarged heart	99.44	93.53	92.86	94.20	99.86	95.04	93.06	97.10
Fibrosis	96.33	82.98	79.59	86.67	97.94	87.06	92.50	82.22
Fractures	97.71	89.55	88.24	90.91	98.09	92.31	93.75	90.91
Hiatal hernia	99.69	73.68	77.78	70.00	99.73	73.68	77.78	70.00
Lung tumor	97.68	93.24	92.00	94.52	98.42	93.75	91.70	95.89
Lymphadenopathy								
All	97.06	95.03	92.44	97.78	97.01	94.02	90.53	97.78
Malignant	97.68	85.00	87.18	82.93	97.89	84.21	91.43	78.05
Mediastinal tumor	86.78	16.00	66.67	9.09	93.40	8.33	50.00	4.55
Pericard. effusion	97.96	90.00	96.43	84.38	94.20	88.14	96.30	81.25
Pleural abnorm.	97.36	57.14	57.14	57.14	97.12	69.23	75.00	64.29
Pneumonia	97.43	87.86	83.11	93.18	97.46	87.54	82.55	93.18
Pneumothorax	99.79	92.31	96.77	88.24	99.85	92.31	96.77	88.24
Pulm. embolism	99.83	90.62	90.62	90.62	99.94	92.06	93.55	90.62
Therapy devices	99.00	96.00	96.00	96.00	98.86	95.39	94.16	96.67
Ventilation	97.96	94.43	92.23	96.74	97.49	95.16	94.15	96.20
Others	98.28	66.67	84.62	55.00	98.34	70.59	85.71	60.00

Per Class, Token Level Metrics for NER Task on CT and X-rays of the Upper Extremity

Class	AUROC	F1	Precision	Recall	AUROC	F1	Precision	Recall
GottBERT				BioGottBERT				
Amput.	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
Carpal Fx	96.87	82.05	92.31	73.85	98.83	83.19	97.92	72.31
Comb. Forearm Fx	99.93	65.12	48.28	100.0	100.0	100.0	100.0	100.0
DISI	99.6	46.67	53.85	41.18	98.89	51.85	70.0	41.18
Disloc.	99.97	86.49	76.19	100.0	99.98	85.71	78.95	93.75
Dist. Rad. Fx	98.73	40.0	42.86	37.5	99.97	55.56	50.0	62.5
Finger Fx	82.47	0.0	0.0	0.0	91.12	0.0	0.0	0.0
Fi ss.	99.29	79.16	75.38	83.33	98.63	86.02	83.33	88.89
Hum. Fx	99.01	81.24	84.91	77.87	98.95	85.95	88.66	83.4
Incorr. Mat. Pos.	95.98	78.49	100.0	64.6	98.89	70.21	88.0	58.41
Joint Involv.	78.21	0.0	0.0	0.0	75.43	0.0	0.0	0.0
Lux.	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
Mat. Break	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0

(continued on next page)

(continued)

Class	AUROC	F1	Precision	Recall	AUROC	F1	Precision	Recall
	GottBERT				BioGottBERT			
Mat. Intact	100.0	80.0	66.67	100.0	100.0	83.33	71.43	100.0
Midhand Fx	99.99	87.5	77.78	100.0	99.99	87.5	77.78	100.0
No Disloc.	99.39	75.86	68.75	84.62	99.71	78.05	74.42	82.05
No Disloc.	98.66	81.1	88.06	75.16	97.98	80.66	83.11	78.34
No Fx	98.65	89.6	89.24	89.96	99.07	85.69	82.14	89.56
No Joint Involv.	99.63	75.0	75.51	74.5	99.23	77.85	77.85	77.85
No Scapholun. Diss.	99.21	78.81	75.86	81.99	97.27	78.86	76.23	81.68
No Ulna Fx	100.0	73.33	100.0	57.89	99.42	59.26	100.0	42.11
Old Fx	99.87	20.0	100.0	11.11	94.17	13.79	100.0	7.41
PISI	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
Pseudoarth.	100.0	90.0	85.71	94.74	99.99	89.47	89.47	89.47
Rad. Fx	99.85	59.34	45.0	87.1	99.94	56.25	41.54	87.1
STT OA	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
Scap. Fx	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
Scaphoid Fx	98.85	76.01	73.29	78.93	98.01	74.61	77.14	72.24
Scapholun. Diss.	99.18	81.02	77.72	84.62	99.34	75.81	69.46	83.43
Sublux.	100.0	98.18	96.43	100.0	100.0	100.0	100.0	100.0
Thumb OA	99.68	0.0	0.0	0.0	99.94	28.0	16.28	100.0
Ulna Fx	98.0	97.19	97.08	97.31	98.02	97.05	97.15	96.96
Ulna+	95.44	31.82	55.26	22.34	93.68	37.04	60.98	26.6
Ulna-	98.1	71.26	68.0	74.84	98.65	65.79	56.56	78.62
Wrist OA	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0

Class	AUROC	F1	Precision	Recall	AUROC	F1	Precision	Recall
	Multilingual BERT				German-MedBERT			
Amput.	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
Carpal Fx	97.52	68.52	84.09	57.81	95.52	64.22	77.78	54.69
Comb. Forearm Fx	96.71	0.0	0.0	0.0	98.61	60.0	100.0	42.86
DISI	96.9	12.5	100.0	6.67	99.24	38.1	25.0	80.0
Disloc.	99.97	88.89	80.0	100.0	99.98	88.89	80.0	100.0
Dist. Rad. Fx	98.4	46.15	60.0	37.5	99.35	42.86	50.0	37.5
Finger Fx	94.58	0.0	0.0	0.0	88.48	0.0	0.0	0.0
Fiss.	97.29	81.42	79.68	83.24	99.33	74.16	64.85	86.59
Hum. Fx	98.84	82.05	82.54	81.57	97.36	79.68	80.97	78.43
Incorr. Mat. Pos.	98.88	68.97	98.36	53.1	99.8	74.89	74.56	75.22
Joint Involv.	77.39	0.0	0.0	0.0	72.16	0.0	0.0	0.0
Lux.	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
Mat. Break	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
Mat. Intact	100.0	83.33	71.43	100.0	100.0	100.0	100.0	100.0
Midhand Fx	99.99	87.5	77.78	100.0	99.99	87.5	77.78	100.0
No Disloc.	99.55	64.15	50.75	87.18	97.73	70.0	68.29	71.79
No Disloc.	89.9	78.45	88.1	70.7	97.5	77.12	75.93	78.34
No Fx	96.91	87.04	85.77	88.35	99.14	83.27	79.06	87.95
No Joint Involv.	99.34	74.68	71.88	77.7	96.89	78.32	81.16	75.68
No Scapholun. Diss.	98.38	76.83	77.56	76.1	97.74	76.48	82.78	71.07
No Ulna Fx	99.38	0.0	0.0	0.0	92.92	0.0	0.0	0.0
Old Fx	100.0	100.0	100.0	100.0	99.53	20.69	100.0	11.54
PISI	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
Pseudoarth.	100.0	90.0	85.71	94.74	100.0	87.8	81.82	94.74
Rad. Fx	98.96	74.67	65.12	87.5	99.82	62.79	50.0	84.38
STT OA	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
Scap. Fx	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
Scaphoid Fx	98.15	77.03	76.51	77.55	98.06	74.96	78.44	71.77
Scapholun. Diss.	98.05	73.87	75.0	72.78	98.24	73.72	87.1	63.91
Sublux.	99.63	89.8	100.0	81.48	99.98	68.29	100.0	51.85
Thumb OA	96.86	57.14	42.86	85.71	99.69	92.31	100.0	85.71
Ulna Fx	97.2	96.26	95.66	96.86	97.21	96.02	95.64	96.41
Ulna+	94.63	31.15	63.33	20.65	90.81	24.24	40.0	17.39
Ulna-	95.84	61.35	59.88	62.89	98.09	72.73	68.13	77.99
Wrist OA	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0

Class	AUROC	F1	Precision	Recall	AUROC	F1	Precision	Recall
	medBERT.de				medBERT.de _{dedup}			
Amput.	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
Carpal Fx	97.1	69.23	90.0	56.25	98.2	83.05	90.74	76.56
Comb. Forearm Fx	100.0	100.0	100.0	100.0	95.36	0.0	0.0	0.0
DISI	93.81	44.44	50.0	40.0	97.97	63.64	100.0	46.67
Disloc.	99.96	72.73	57.14	100.0	99.98	80.0	66.67	100.0
Dist. Rad. Fx	99.98	66.67	53.85	87.5	99.99	72.73	57.14	100.0
Finger Fx	48.2	0.0	0.0	0.0	84.76	0.0	0.0	0.0
Fiss.	99.2	81.35	75.85	87.71	99.01	85.71	82.81	88.83

(continued on next page)

(continued)

Class	AUROC	F1	Precision	Recall	AUROC	F1	Precision	Recall
Hum. Fx	98.97	84.35	82.16	86.67	97.97	83.64	86.25	81.18
Incorr. Mat. Pos.	98.97	63.16	93.1	47.79	99.35	75.14	100.0	60.18
Joint Involv.	70.89	0.0	0.0	0.0	73.7	0.0	0.0	0.0
Lux.	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
Mat. Break	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
Mat. Intact	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Midhand Fx	100.0	87.5	77.78	100.0	99.99	87.5	77.78	100.0
No Disloc.	92.52	80.28	89.76	72.61	93.75	78.41	81.94	75.16
No Disloc.	99.57	65.12	59.57	71.79	99.75	69.05	64.44	74.36
No Fx	98.42	87.14	83.46	91.16	97.9	87.35	84.72	90.16
No Joint Involv.	98.78	72.03	68.71	75.68	99.05	77.15	68.78	87.84
No Scapholun. Diss.	98.07	81.63	81.5	81.76	98.14	78.34	76.58	80.19
No Ulna Fx	99.87	77.42	100.0	63.16	100.0	78.79	92.86	68.42
Old Fx	99.97	70.0	100.0	53.85	100.0	96.0	100.0	92.31
PISI	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
Pseudoarth.	99.99	95.0	90.48	100.0	100.0	95.0	90.48	100.0
Rad. Fx	98.86	63.16	54.55	75.0	96.29	72.73	70.59	75.0
STT OA	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
Scap. Fx	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0
Scaphoid Fx	98.19	78.32	77.41	79.25	97.9	76.45	72.7	80.61
Scapholun. Diss.	98.9	67.48	69.38	65.68	98.77	78.31	79.75	76.92
Sublux.	99.99	82.61	100.0	70.37	98.89	89.8	100.0	81.48
Thumb OA	95.96	80.0	75.0	85.71	100.0	82.35	70.0	100.0
Ulna Fx	97.74	96.52	96.48	96.55	97.22	96.54	96.47	96.62
Ulna+	95.64	26.09	65.22	16.3	92.93	36.64	61.54	26.09
Ulna-	97.95	74.01	67.18	82.39	90.87	71.52	71.97	71.07
Wrist OA	100.0	100.0	100.0	100.0	100.0	100.0	100.0	100.0

Fx: Fracture, OA: Osteoarthritis, Hum.: Humerus, Scapholun.: Scapholunate, Diss.: Dissociation, Disloc.: Dislocation, Rad.: Radius, Amput.: Amputation, Involv.: Involvement, STT: Scaphotrapeziotrapezoid, Ulna: Ulna Minus, Scap.: Scapula, Mat.: Material, Comb.: Combined, Pseudoarth.: Pseudarthrosis, Dist.: Distal, Incorr.: Incorrect, Pos.: Position, Lux.: Luxation, Sublux.: Subluxation.

Appendix 3. – distribution of classes for benchmarks based on radiology reports

For the benchmarks based on radiology reports, the distribution of classes is an important factor to consider when evaluating the performance of our models. As some diagnoses are rarer than others, the class distribution is highly skewed, especially for the CT and NER tasks.

Class Distribution for X-ray Classification Task

Class	Train	Valid	Test
Congestion	273	89	88
Opacity	584	120	144
Effusion	460	214	158
Pneumothorax	74	19	28
Thoracic drain	190	80	180
Venous catheter	579	283	343
Gastric tube	244	117	160
Tracheal tube	397	230	247
Misplaced	49	20	12

Class Distribution for CT Classification Task

Class	Train	Valid	Test
Aortic dissection	1	2	2
Bone tumors	25	19	22
Bronchial abnormalities	98	38	56
Congestion	40	19	22
Effusion	251	138	158
Lung	179	93	82
Soft tissue	42	24	21
Enlarged heart	128	62	69
Fibrosis	91	45	45
Fractures	118	77	66
Hiatal hernia	26	10	10
Lung tumor	427	224	219
All	396	223	225
Malignant	154	83	82
Mediastinal tumor	33	21	22
Pericardial effusion	69	32	32
Pleural abnormalities	26	15	14
Pneumonia	268	119	132
Pneumothorax	49	33	34
Pulmonary embolism	49	33	32

(continued on next page)

(continued)

Class	Train	Valid	Test
Therapy devices	243	135	150
Ventilation	306	165	184
Others	41	17	20

Class Distribution for NER Task on X-ray/CT of the Wrist

Class	Train	Valid	Test
Amput.	1	0	1
Carpal Fx	175	41	30
Comb. Forearm Fx	9	2	4
DISI	17	2	1
Disloc.	444	73	75
Dist. Rad. Fx	410	48	53
Finger Fx	7	2	3
Fiss.	5	2	3
Hum. Fx	42	5	8
Joint Involv.	504	70	75
Lux.	14	4	2
Mat. Intact	25	5	2
Midhand Fx	117	16	26
No Disloc.	613	113	88
No Fx	751	106	91
No Joint Involv.	42	5	13
No Scapholun. Diss.	110	21	7
Old Fx	75	10	13
Pseudoarth.	9	2	2
Rad. Fx	112	14	13
STT OA	18	4	2
Scaphoid Fx	90	19	17
Scapholun. Diss.	35	3	5
Sublux.	35	0	3
Thumb OA	47	7	12
Ulna Fx	220	29	25
Ulna-	22	2	3
Wrist OA	6	0	2

Appendix 4. – model selection for radiology tasks

For the benchmarks based on radiology reports, the hyperparameter optimization resulted in the following parameters:

Parameters of the Best Models for CT Classification Task

Model	Batch Size	Learning Rate	Warmup Steps
GottBERT	16	3.42e-05	23
BioGottBERT	16	6.69e-05	78
Multilingual BERT	8	2.96e-05	648
German-MedBERT	16	3.96e-05	146
medBERT.de	8	2.41e-05	1
medBERT.de _{dedup}	8	1.9e-05	49

Parameters of the Best Models for X-Ray Classification Task

Model	Batch Size	Learning Rate	Warmup Steps
GottBERT	8	2.04e-05	454
BioGottBERT	16	3.81e-05	293
Multilingual BERT	16	7.77e-05	995
German-MedBERT	8	4.20e-05	249
medBERT.de	8	1.18e-05	140
medBERT.de _{dedup}	16	3.80e-05	2

Parameters of the Best Models for NER Task on CT and X-rays of the Upper Extremity

Model	Batch Size	Learning Rate	Warmup Steps
GottBERT	8	2.18e-05	957
BioGottBERT	8	3.81e-05	243
Multilingual BERT	16	6.92e-05	882
German-MedBERT	8	3.32e-05	232
medBERT.de	8	4.79e-05	466
medBERT.de _{dedup}	8	2.22e-05	40

References

- Akbik, A., Blythe, D., & Vollgraf, R. (2018). Contextual String Embeddings for Sequence Labeling. *COLING 2018, 27th International Conference on Computational Linguistics*, 1638–1649.
- Borchert, F., Lohr, C., Modersohn, L., Langer, T., Follmann, M., Sachs, J. P., Hahn, U., & Schapranow, M.-P. (2020). GGPONC: A Corpus of German Medical Text with Rich Metadata Based on Clinical Practice Guidelines. *Proceedings of the 11th International Workshop on Health Text Mining and Information Analysis*, 38–48.
- Borchert, F., Lohr, C., Modersohn, L., Witt, J., Langer, T., Follmann, M., Gietzelt, M., Arnrich, B., Hahn, U., & Schapranow, M.-P. (2022). GGPONC 2.0-the German clinical guideline corpus for oncology: Curation workflow, annotation policy, baseline NER taggers. *Proceedings of the Thirteenth Language Resources and Evaluation Conference*, 3650–3660.
- Bressemer, K. K., Adams, L. C., Gaudin, R. A., Tröltzsch, D., Hamm, B., Makowski, M. R., ... Niehues, S. M. (2020). Highly accurate classification of chest radiographic reports using a deep learning natural language model pre-trained on 3.8 million text reports. *Bioinformatics*, 36(21), 5255–5261.
- Canete, J., Chaperon, G., Fuentes, R., Ho, J.-H., Kang, H., & Pérez, J. (2020). Spanish pre-trained bert model and evaluation data. *Pml4dc at Iclr*, 2020, 1–10.
- Chan, B., Schweter, S., & Möller, T. (2020). German's Next Language Model. *Proceedings of the 28th International Conference on Computational Linguistics*, 6788–6796.
- Devlin, J., Chang, M.-W., Lee, K., & Toutanova, K. (2019). BERT: Pre-training of Deep Bidirectional Transformers for Language Understanding. In J. Burstein, C. Doran, & T. Solorio (Eds.), *Proceedings of the 2019 Conference of the North American Chapter of the Association for Computational Linguistics: Human Language Technologies, NAACL-HLT 2019, Minneapolis, MN, USA, June 2-7, 2019, Volume 1 (Long and Short Papers)* (pp. 4171–4186). Association for Computational Linguistics.
- Frei, J., & Kramer, F. (2022). GERNERMED: An open German medical NER model. *Software Impacts*, 11, Article 100212.
- Gu, Y., Tinn, R., Cheng, H., Lucas, M., Usuyama, N., Liu, X., ... Poon, H. (2021). Domain-specific language model pretraining for biomedical natural language processing. *ACM Transactions on Computing for Healthcare (HEALTH)*, 3(1), 1–23.
- J. Kaplan, S. McCandlish, T. Henighan, T. B. Brown, B. Chess, R. Child, S. Gray, A. Radford, J. Wu, and D. Amodei (2020) Scaling laws for neural language models. CoRR, abs/2001.08361.
- Lee, J., Yoon, W., Kim, S., Kim, D., Kim, S., So, C. H., & Kang, J. (2020). BioBERT: A pre-trained biomedical language representation model for biomedical text mining. *Bioinformatics*, 36(4), 1234–1240.
- Lee, K., Ippolito, D., Nystrom, A., Zhang, C., Eck, D., Callison-Burch, C., & Carlini, N. (2021). Deduplicating training data makes language models better. *arXiv Preprint arXiv:2107.06499*.
- Lentzen, M., Madan, S., Lage-Rupprecht, V., Kühnel, L., Fluck, J., Jacobs, M., Mittermaier, M., Witzelrath, M., Brunecker, P., Hofmann-Apitius, M., Weber, J., & Fröhlich, H. (11 2022). Critical assessment of transformer-based AI models for German clinical notes. *JAMIA Open*, 5(4). <https://doi.org/>.
- Liu, Y., Ott, M., Goyal, N., Du, J., Joshi, M., Chen, D., Levy, O., Lewis, M., Zettlemoyer, L., & Stoyanov, V. (2019). Roberta: A robustly optimized bert pretraining approach. *arXiv Preprint arXiv:1907.11692*.
- Martin, L., Muller, B., Suárez, P. J. O., Dupont, Y., Romary, L., de La Clergerie, É. V., Seddah, D., & Sagot, B. (2019). CamemBERT: a tasty French language model. *arXiv Preprint arXiv:1911.03894*.
- Modersohn, L., Schulz, S., Lohr, C., & Hahn, U. (2022). GRASCO-The First Publicly Shareable, Multiply-Alienated German Clinical Text Corpus. *Studies in Health Technology and Informatics*, 296, 66–72.
- Pedro Javier Ortiz Suárez, Laurent Romary, and Benoît Sagot. (2020). A Monolingual Approach to Contextualized Word Embeddings for Mid-Resource Languages. In *Proceedings of the 58th Annual Meeting of the Association for Computational Linguistics*, pages 1703–1714, Online. Association for Computational Linguistics.
- Pérez-Mayos, L., Ballesteros, M., & Wanner, L. (2021). How much pretraining data do language models need to learn syntax? In M.-F. Moens, X. Huang, L. Specia, & S. W.-T. Yih (Eds.), *Proceedings of the 2021 Conference on Empirical Methods in Natural Language Processing, EMNLP 2021, Virtual Event / Punta Cana, Dominican Republic, 7-11 November, 2021* (pp. 1571–1582). Association for Computational Linguistics.
- Rasmy, L., Xiang, Y., Xie, Z., Tao, C., & Zhi, D. (2021). Med-BERT: Pretrained contextualized embeddings on large-scale structured electronic health records for disease prediction. *NPJ Digital Medicine*, 4(1), 1–13.
- Risch, J., Krebs, E., Löser, A., Riese, A., & Krestel, R. (2018). Fine-Grained Classification of Offensive Language. *Proceedings of GermEval 2018 (co-located with KONVENS)*, 38–44.
- Rust, P., Pfeiffer, J., Vulic, I., Ruder, S., & Gurevych, I. (2021). How Good is Your Tokenizer? On the Monolingual Performance of Multilingual Language Models. In C. Zong, F. Xia, W. Li, & R. Navigli (Eds.), *Proceedings of the 59th Annual Meeting of the Association for Computational Linguistics and the 11th International Joint Conference on Natural Language Processing, ACL/IJCNLP 2021, (Volume 1: Long Papers), Virtual Event, August 1-6, 2021* (pp. 3118–3135). Association for Computational Linguistics.
- Scheible, R., Thomczyk, F., Tippmann, P., Jaravine, V., & Boeker, M. (2020). GottBERT: a pure German language model. *arXiv Preprint arXiv:2012.02110*.
- Schmidt, D., Osmanodja, B., Pfefferkorn, M., Graf, V., Raschke, D., Duettmann, W., et al. (2021). TBase-an integrated electronic health record and research database for kidney transplant recipients. *JoVE (Journal of Visualized Experiments)*, 170(1), e61971.
- Shrestha, M. (2021). *Development of a Language Model for Medical Domain*. Hochschule Rhein-Waal.
- You, Y., Li, J., Reddi, S., Hseu, J., Kumar, S., Bhojanapalli, S., Song, X., Demmel, J., Keutzer, K., & Hsieh, C.-J. (2019). Large batch optimization for deep learning: Training bert in 76 minutes. *arXiv Preprint arXiv:1904.00962*.