

The Kauffman bracket skein module of the lens spaces via unoriented braids

Citation for published version (APA):

Diamantis, I. (2024). The Kauffman bracket skein module of the lens spaces via unoriented braids. Communications in Contemporary Mathematics, 26(02), Article 22500766. https://doi.org/10.1142/S0219199722500766

Document status and date: Published: 01/01/2024

DOI: 10.1142/S0219199722500766

Document Version: Publisher's PDF, also known as Version of record

Document license: Taverne

Please check the document version of this publication:

 A submitted manuscript is the version of the article upon submission and before peer-review. There can be important differences between the submitted version and the official published version of record. People interested in the research are advised to contact the author for the final version of the publication, or visit the DOI to the publisher's website.

• The final author version and the galley proof are versions of the publication after peer review.

 The final published version features the final layout of the paper including the volume, issue and page numbers.

Link to publication

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these riahts.

Users may download and print one copy of any publication from the public portal for the purpose of private study or research.

You may not further distribute the material or use it for any profit-making activity or commercial gain
 You may freely distribute the URL identifying the publication in the public portal.

If the publication is distributed under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license above, please follow below link for the End User Agreement:

www.umlib.nl/taverne-license

Take down policy

If you believe that this document breaches copyright please contact us at:

repository@maastrichtuniversity.nl

providing details and we will investigate your claim.





The Kauffman bracket skein module of the lens spaces via unoriented braids

Ioannis Diamantis

Department of Data Analytics and Digitalisation School of Business and Economics, Maastricht University P. O. Box 616, 6200 MD Maastricht, The Netherlands i.diamantis@maastrichtuniversity.nl

> Received 23 June 2022 Revised 3 October 2022 Accepted 19 October 2022 Published 19 December 2022

In this paper, we develop a braid theoretic approach for computing the Kauffman bracket skein module of the lens spaces L(p,q), KBSM(L(p,q)), for $q \neq 0$. For doing this, we introduce a new concept, that of an *unoriented braid*. Unoriented braids are obtained from standard braids by ignoring the natural top-to-bottom orientation of the strands. We first define the generalized Temperley-Lieb algebra of type B, $TL_{1,n}$, which is related to the knot theory of the solid torus ST, and we obtain the universal Kauffman brackettype invariant, V, for knots and links in ST, via a unique Markov trace constructed on $TL_{1,n}$. The universal invariant V is equivalent to the KBSM(ST). For passing now to the KBSM(L(p,q)), we impose on V relations coming from the band moves (or slide moves), that is, moves that reflect isotopy in L(p,q) but not in ST, and which reflect the surgery description of L(p,q), obtaining thus, an infinite system of equations. By construction, solving this infinite system of equations is equivalent to computing KBSM(L(p,q)). We first present the solution for the case q = 1, which corresponds to obtaining a new basis, \mathcal{B}_p , for $\operatorname{KBSM}(L(p,1))$ with $(\lfloor p/2 \rfloor + 1)$ elements. We note that the basis \mathcal{B}_p is different from the one obtained by Hoste and Przytycki. For dealing with the complexity of the infinite system for the case q > 1, we first show how the new basis \mathcal{B}_p of KBSM(L(p, 1)) can be obtained using a diagrammatic approach based on unoriented braids, and we finally extend our result to the case q > 1. The advantage of the braid theoretic approach that we propose for computing skein modules of c.c.o. 3-manifolds, is that the use of braids provides more control on the isotopies of knots and links in the manifolds, and much of the diagrammatic complexity is absorbed into the proofs of the algebraic statements.

Keywords: Skein module; Kauffman bracket; unoriented braids; solid torus; lens spaces; mixed links; mixed braids; braid group of type B; generalized Hecke algebra of type B; generalized Temperley–Lieb algebra of type B.

Mathematics Subject Classification 2020: 57K31, 57K14, 20F36, 20F38, 57K10, 57K12, 57K45, 57K35, 57K99, 20C08

0. Introduction

Skein modules were introduced independently by Przytycki [31] and Turaev [32] as generalizations of knot polynomials in S^3 to knot polynomials in arbitrary 3-manifolds. They are quotients of free modules over isotopy classes of links in 3-manifolds by properly chosen local (skein) relations. A skein module of a 3-manifold yields all possible isotopy invariants of knots which satisfy a particular skein relation. Skein modules based on the Kauffman bracket skein relation

$$L_+ - AL_0 - A^{-1}L_\infty,$$

where L_{∞} and L_0 are represented schematically by the illustrations in Fig. 1, are called *Kauffman bracket skein modules* (KBSM).

For example, the Kauffman bracket skein module of S^3 , KBSM(S^3), is freely generated by the unknot and it is equivalent to the Kauffman bracket for framed links in S^3 up to regular isotopy. It is ambient isotopy counterpart is the wellknown Jones polynomial. Recall that the algebraic counterpart construction of the Kauffman bracket is the Temperley–Lieb algebra together with a unique Markov trace constructed by Jones (see [26] and references therein). It is worth pointing out the importance of the Temperley–Lieb algebras. Through the pioneering work of Jones [26, 25], these algebras related knot theory to statistical mechanics, topological quantum field theories and the construction of quantum invariants for 3-manifolds (works of Witten, Reshetikhin–Turaev, Lickorish, etc).

The precise definition of KBSM is as follows.

Definition 1. Let M be an oriented 3-manifold and \mathcal{L}_{fr} be the set of isotopy classes of unoriented framed links in M. Let $R = \mathbb{Z}[A^{\pm 1}]$ be the Laurent polynomials in Aand let $R\mathcal{L}_{fr}$ be the free R-module generated by \mathcal{L}_{fr} . Let S be the ideal generated by the skein expressions $L_{+} - AL_{0} - A^{-1}L_{\infty}$ and $L \bigsqcup O - (-A^{2} - A^{-2})L$. Note that blackboard framing is assumed and that $L \bigsqcup O$ stands for the union of a link L and the trivially framed unknot in a ball disjoint from L.

Then the Kauffman bracket skein module of M, KBSM(M), is defined to be

$$\operatorname{KBSM}(M) = \frac{R\mathcal{L}_{\operatorname{fr}}}{S}.$$

Skein modules of 3-manifolds have become very important algebraic tools in the study of 3-manifolds, since their properties renders topological information about



Fig. 1. The links L, L_0 and L_∞ locally.

Commun. Contemp. Math. Downloaded from www.worldscientific.com by Ioannis Diamantis on 03/03/23. Re-use and distribution is strictly not permitted, except for Open Access articles.

the 3-manifolds. In this paper, we are interested in the Kauffman bracket skein module of the lens spaces L(p,q), for $p \neq 0$ using braid theoretical tools. We consider the lens spaces L(p,q) obtained from S^3 by rational surgery along the unknot with coefficient p/q. Surgery along the unknot is realized by considering the complementary solid torus and attaching to it a solid torus according to a (p,q)-homeomorphism on the boundary. Hence, the Kauffman bracket skein module of the solid torus, KBSM(ST), is our starting point. KBSM(ST) is essential in the study of Kauffman bracket skein modules of arbitrary c.c.o. 3-manifolds, since every c.c.o. 3-manifold can be obtained by surgery along a framed link in S^3 with unknotted components. The family of the lens spaces, L(p,q), comprises the simplest example, since, as mentioned before, they are obtained by rational surgery on the unknot. For a survey on skein modules, see [31].

From the above it is clear that the knot theory of ST is of fundamental importance for studying knot theory in other c.c.o. 3-manifolds. A basis for KBSM(ST) is presented in [22] using diagrammatic methods (see also [32]). More precisely:

Theorem 1 ([22]). The Kauffman bracket skein module of ST, KBSM(ST), is freely generated by an infinite set of generators $\{x^n\}_{n=0}^{\infty}$, where x^n denotes n parallel copies of a longitude of ST and x^0 is the affine unknot (see Fig. 2).

We develop a braid theoretic approach to the Kauffman bracket skein module of L(p,q), for $p \neq 0$. Note that the KBSM(L(p,q)) has been computed via a diagrammatic approach in [23], where it is shown that:

Theorem 2 ([23, Theorem 4]). The Kauffman bracket skein module of the lens spaces L(p,q) for $p \ge 1$ is freely generated by $\{x^i\}_{i=0}^{\lfloor p/2 \rfloor}$, where $\lfloor p/2 \rfloor$ denotes the integer part of p/2.

We propose a new and more efficient method for computing Kauffman bracket skein modules via the notion of *unoriented braids*, which we define along the way. The algebraic techniques developed in [29, 30, 27, 28, 12, 13, 17, 15, 1–4] are of great importance. More precisely, in [28] it is shown that the generalized Hecke algebra of type B, $H_{1,n}$, is related to the knot theory of the solid torus and the Artin group of the Coxeter group of type B, $B_{1,n}$. Moreover, the most generic analog of the HOMFLYPT polynomial, X, for links in the solid torus ST has been derived from



Fig. 2. The standard basis of KBSM(ST).

 $H_{1,n}$ via a unique Markov trace constructed on it. The invariant X recovers the HOMFLYPT skein module of ST (see [28, 13, 14]) and as shown in [17, 15], it can be extended to an invariant for knots in L(p, 1) by considering the effect of the *braid band moves*, i.e. the analog of band moves on the braid level (see [29, 30, 12]) on elements in some basis of the HOMFLYPT skein module of ST. As shown in [31], this is also true for the case of the Kauffman bracket skein module of $L(p,q), p \ge 1$. That is, in order to compute KBSM(L(p,q)), we consider elements in some basis of the universal invariant for knots in ST, relations coming from the band moves. That is,

$$\text{KBSM}(L(p,q)) = \frac{\text{KBSM}(\text{ST})}{\langle a - \text{bbm}(a) \rangle}, \text{ where } a \text{ basis element of KBSM}(\text{ST}).$$
(1)

In this paper, we give the complete analog of KBSM(L(p,q)) via braids. We first define the generalized Temperley–Lieb algebra of type B, $\text{TL}_{1,n}$, as the quotient of $H_{1,n}$ over the ideal generated by expressions of the form

$$\sigma_{i,i+1} := 1 + u(\sigma_i + \sigma_{i+1}) + u^2(\sigma_i \sigma_{i+1} + \sigma_{i+1} \sigma_i) + u^3 \sigma_i \sigma_{i+1} \sigma_i.$$
(2)

That is, $\operatorname{TL}_{1,n} := \frac{H_{1,n}}{\langle \sigma_{i,i+1} \rangle}$. We then show that the trace function constructed in [28] on $H_{1,n}$ factors through $\operatorname{TL}_{1,n}$ and, by normalizing it, we obtain the universal Kauffman bracket, V, for knots and links in ST. We then extend this invariant to knots and links in L(p,q) for $q \neq 0$, by solving the infinite system of equations resulting from the band moves. Namely, we force

$$V_{\widehat{a}} = V_{\widehat{\mathrm{bm}(a)}},\tag{3}$$

for all a in the basis of KBSM(ST), where bm(a) denotes the result of a band move on a.

Equation (3) has particularly simple formulations with the use of a new basis, $B_{\rm ST}$, for the Kauffman bracket skein module of ST, that is presented first in Sec. ?? (see Eq. (8) in this paper), in terms of mixed braids (that is, classical braids with the first strand identically fixed). We prove that $B_{\rm ST}$ is a basic set of KBSM(ST) by relating the braided form of the standard basis, $B'_{\rm ST}$, to $B_{\rm ST}$ via a lower triangular matrix with invertible elements in the diagonal. For an illustration of elements in the basis $B_{\rm ST}$, see the bottom of Fig. 8. Note that the same basis is presented in [1] via the Temperley–Lieb algebra of type B and in [21] via a diagrammatic approach based on arrow diagrams. We then solve the infinite system of equations (3) for the case of L(p, 1) and we arrive at the main result of this paper. In particular, we have the following result.

Theorem 3. The Kauffman bracket skein module of the lens spaces $L(p, 1), q \neq 0$, is freely generated by elements in the set \mathcal{B}_p , consisting of element in the form $\{t^i\}_{i=0}^{\lfloor p/2 \rfloor}$. For an illustration of elements in \mathcal{B}_p , see Fig. 8.

Note that the basis \mathcal{B}_p is different from the basis presented by Hoste and Przytycki in [23] for KBSM(L(p, 1)). After we establish the algebraic method for the case q = 1, we present a diagrammatic method based on braids and we use it to derive the basis \mathcal{B}_p for KBSM(L(p, 1)). We then extend this diagrammatic method for the case of L(p,q), $q \neq 0$, and in particular, we show that the set \mathcal{B}_p forms a basis for KBSM(L(p,q)). We finally discuss the algebraic method for the case of KBSM(L(p,q)).

This paper is organized as follows: In Sec. 1, we recall the setting and the essential techniques and results from [27–30, 12]. More precisely, we present isotopy moves for knots and links in L(p,q) and we then describe braid equivalence for knots and links in L(p,q). In Sec. 2, we present results from [28, 18] and we extend these results in order to construct the universal invariant for knots in ST of the Kauffman bracket type. More precisely, we start by recalling results on the generalized Hecke algebra of type B, $H_{1,n}$, and we present the universal invariant of the HOMFLYPT type for knots in ST via a unique Markov trace, tr, defined on the algebra. This invariant captures the HOMFLYPT skein module of ST. We then pass to the generalized Temperley–Lieb algebra of type B, defined as a quotient algebra from $H_{1,n}$ over the ideal generated by relations (2), and we find necessary and sufficient conditions so that the trace tr factors through $TL_{1,n}$. Using the trace, we define the universal invariant V for knots in ST of the Kauffman bracket type. In Sec. 3, we present results on KBSM(ST) via braids and we present a new basis, $B_{\rm ST}$, for KBSM(ST) via an ordering relation defined in [13]. This basis is also presented in [1] using the Temperley–Lieb algebra of type B, which, roughly speaking, is related to an invariant for knots in ST, but not the most generic one that captures KBSM(ST). Moreover, the results presented for obtaining $B_{\rm ST}$ are essential for the computation of the Kauffman bracket skein module of the lens spaces L(p, 1)that we compute in Sec. 4 using algebraic techniques. This is done with the use of the newly introduced concept of unoriented braids defined in Sec. 3.1. These new objects seem promising for studying knots in c.c.o. 3-manifolds (and c.c.o. 3manifolds in general) via the proposed technique. Finally, in Sec. 5 we present the basis \mathcal{B}_p for L(p,1) using a diagrammatic approach based on unoriented braids and in Sec. 5.2 we show how these results can be generalized for the case $L(p,q), q \neq 0$.

It is worth mentioning that the importance of our approach lies in the fact that it can shed light to the problem of computing (various) skein modules of arbitrary c.c.o. 3-manifolds (see [4] for the case of the Kauffman bracket skein module of the complement of (2, 2p + 1)-torus knots). The main difficulty of the problem lies in solving the infinite system of equations (3).

1. Preliminaries

1.1. Mixed links and isotopy in L(p,q)

In this section, we recall results from [29, 30, 12]. We shall consider ST to be the complement of a solid torus in S^3 . Then, an oriented link L in ST can be represented by an oriented *mixed link* in S^3 , that is, a link in S^3 consisting of the unknotted fixed part \hat{I} representing the complementary solid torus in S^3 , and the moving part



Fig. 3. A mixed link.

L that links with \widehat{I} . A mixed link diagram is a diagram $\widehat{I} \cup \widetilde{L}$ of $\widehat{I} \cup L$ on the plane of \widehat{I} , where this plane is equipped with the top-to-bottom direction of I (for an illustration, see Fig. 3).

The lens spaces L(p,q) can be obtained from S^3 by surgery on the unknot with surgery coefficient $p \in \mathbb{Z}$. Surgery along the unknot can be realized by considering first the complementary solid torus and then attaching to it a solid torus according to some homeomorphism h on the boundary. For L(p, 1), the homeomorphism hon the boundaries of the solid tori maps the meridian, m_1 , of one solid torus to a (p, 1)-curve on the other, that is, a $(p \cdot l_2 + 1 \cdot m_2)$ -curve, where l_2 denotes the longitude and m_2 denotes the meridian of the second solid torus:

$$h: \partial \operatorname{ST}_1 \to \partial \operatorname{ST}_2$$
$$m_1 \mapsto p \cdot l_2 + 1 \cdot m_2$$

Thus, isotopy in L(p,q) can be viewed as isotopy in ST together with the *band* moves in S^3 , which reflect the surgery description of the manifold. In Fig. 4, the two types of band moves for the case of L(p, 1) are illustrated, according to the orientation of the component of the knot and of the surgery curve. In the α -type the orientation of the arc is opposite to the orientation of the surgery curve, but after the performance of the move their orientations agree. In the β -type the orientations initially agree, but disagree after the performance of the move. For the case of L(p,q), see Fig. 17. In [12], it is shown that in order to describe isotopy for knots and links in a c.c.o. 3-manifold, it suffices to consider only one type of band moves



Fig. 4. The two types of band moves for q = 1.

(cf. [12, Theorem 6]) and thus, isotopy between oriented links in L(p,q) is reflected in S^3 by means of the following theorem.

Theorem 4. Two oriented links in L(p,q) are isotopic if and only if two corresponding mixed link diagrams of theirs differ by isotopy in ST together with a finite sequence of one type of band moves.

1.2. Mixed braids and braid equivalence for knots and links in L(p,q)

In this section, we introduce the notion of mixed braids, the mixed braid group $B_{1,n}$ and braid equivalence in L(p,q). Note that a mixed braid in S^3 is a braid where, without loss of generality, its first strand represents \hat{I} , the fixed part, and the other strands, β , represent the moving part L. We shall call the subbraid β the moving part of $I \cup \beta$ (see bottom left-hand side of Fig. 4). We now recall the analog of the Alexander theorem for knots in ST (cf. [28, Theorem 1]).

Theorem 5 (The analog of the Alexander theorem for ST). A mixed link diagram $\hat{I} \cup \tilde{L}$ of $\hat{I} \cup L$ may be turned into a mixed braid $I \cup \beta$ with isotopic closure.

In order to translate isotopy for links in L(p,q) into braid equivalence, we first perform the technique of *standard parting* introduced in [30] in order to separate the moving strands from the fixed strand that represents the lens spaces L(p,q). This can be realized by pulling each pair of corresponding moving strands to the right and *over* or *under* the fixed strand that lies on their right (for more details the reader is referred [30]). Then, we define a *braid band move* to be a move between mixed braids, which is a α -type band move between their closures. It starts with a little band oriented downward, which, before sliding along a surgery strand, gets one twist *positive* or *negative* (see bottom-part of Fig. 5 for q = 1 and Fig. 6 for q > 1). For now we shall only consider the α -type band moves, since the result of an α -type band move remain braided, while the result of a β -band move does not. The β -band moves will become important later, when we compute KBSM(L(p,q)) diagrammatically. Note also that there are two different types of braid band moves; the positive and the negative braid band move, depending on the kind of the twist the moving strand gets before sliding along the fixed strand.

The sets of braids related to ST form groups, which are in fact the Artin braid groups of type B, denoted $B_{1,n}$, with presentation:

$$B_{1,n} = \left(t, \sigma_1, \dots, \sigma_{n-1} \middle| \begin{array}{c} \sigma_1 t \sigma_1 t = t \sigma_1 t \sigma_1 \\ t \sigma_i = \sigma_i t, \quad i > 1 \\ \sigma_i \sigma_{i+1} \sigma_i = \sigma_{i+1} \sigma_i \sigma_{i+1}, \quad 1 \le i \le n-2 \\ \sigma_i \sigma_j = \sigma_j \sigma_i, \quad |i-j| > 1 \end{array} \right),$$

where the generators σ_i and t are illustrated in Fig. 7(i).



Fig. 5. Isotopy in L(p, 1) and the two types of braid band moves on mixed braids.



Fig. 6. The braid band move for L(p,q), q > 1.



Fig. 7. The generators of $B_{1,n}$ and the "looping" elements t'_i and t_i .

2250076-8

Let now \mathcal{L} denote the set of oriented knots and links in ST. Isotopy in L(p,q) is then translated on the level of mixed braids by means of the following theorem.

Theorem 6 ([30, Theorem 5]). Let L_1, L_2 be two oriented links in L(p, 1) and let $I \cup \beta_1$, $I \cup \beta_2$ be two corresponding mixed braids in S^3 . Then L_1 is isotopic to L_2 in L(p, 1) if and only if $I \cup \beta_1$ is equivalent to $I \cup \beta_2$ in \mathcal{B} by the following moves:

 $\begin{array}{ll} \text{(i)} & Conjugation: & \alpha \sim \beta^{-1} \alpha \beta & \text{if } \alpha, \beta \in B_{1,n}; \\ \text{(ii)} & Stabilization moves: } \alpha \sim \alpha \sigma_n^{\pm 1} \in B_{1,n+1} & \text{if } \alpha \in B_{1,n}; \\ \text{(iii)} & Loop \ conjugation: & \alpha \sim t^{\pm 1} \alpha t^{\mp 1} & \text{if } \alpha \in B_{1,n}; \\ \text{(iv)} & Braid \ band \ moves: & \alpha \sim t^p \alpha_+ \sigma_1^{\pm 1}, & a_+ \in B_{1,n+1} \ for \ L(p,1), \\ & \beta \sim [(\sigma_{q-1} \ \dots \ \sigma_1) \ t]^p \ \beta + \sigma_q^{\pm 1}, \ a_+ \in B_{1,n+q} \ for \ L(p,q), \end{array}$

where α_+ is the word α with all indices shifted by +1 and β_+ is the word β with all indices shifted by q. Note that moves (i), (ii) and (iii) correspond to link isotopy in ST.

Notation 1. We denote a braid band move by bbm and, specifically, the result of a positive or negative braid band move performed on a mixed braid β by $bbm_{\pm}(\beta)$.

Note also that in [30] it was shown that the choice of the position of connecting the two components after the performance of a bbm is arbitrary.

2. Knot Algebras and Invariants of Knots in ST

In this section, we present the most generic invariant, V, for knots and links in ST that captures the Kauffman bracket skein module of ST. Therefore, V is the most appropriate invariant to be extended to a Kauffman bracket invariant for knots and links in L(p,q). This invariant is derived via a unique Markov trace, tr, on the generalized Temperley–Lieb algebra of type B, $\text{TL}_{1,n}$. This algebra is defined in Sec. 2.2 as a quotient of the generalized Hecke algebra of type B, $H_{1,n}$, and tr is the unique Markov trace constructed on $H_{1,n}$ which factors through $\text{TL}_{1,n}$. We first recall some important results on the generalized Hecke algebra of type B.

2.1. The generalized Hecke algebra of type B

In [28], the most generic analog of the HOMFLYPT polynomial, X, for links in the solid torus ST has been derived from the generalized Hecke algebras of type B, $H_{1,n}$, via a unique Markov trace constructed on them. This algebra was defined by Lambropoulou as the quotient of $\mathbb{C}[q^{\pm 1}]B_{1,n}$ over the quadratic relations $g_i^2 = (q-1)g_i + q$. Namely,

$$H_{1,n}(q) = \frac{\mathbb{C}[q^{\pm 1}]B_{1,n}}{\langle \sigma_i^2 - (q-1)\sigma_i - q \rangle}.$$

In [28], it is also shown that the following sets form linear bases for $H_{1,n}(q)$ [28, Proposition 1 and Theorem 1]):

(i)
$$\Sigma_n = \{ t_{i_1}^{k_1} \dots t_{i_r}^{k_r} \cdot \sigma \}, \text{ where } 0 \le i_1 < \dots < i_r \le n-1;$$

(ii) $\Sigma'_n = \{ t_{i_1}'^{k_1} \dots t_{i_r}'^{k_r} \cdot \sigma \}, \text{ where } 0 \le i_1 < \dots < i_r \le n-1,$
(4)

where $k_1, \ldots, k_r \in \mathbb{Z}$, $t'_0 = t_0 := t$, $t'_i = g_i \ldots g_1 t g_1^{-1} \ldots g_i^{-1}$ and $t_i = g_i \ldots g_1 t g_1 \ldots g_i$ are the "looping elements" in $H_{1,n}(q)$ (see Fig. 7(ii)) and σ a basic element in the Iwahori–Hecke algebra of type A, $H_n(q)$, for example in the form of the elements in the set [26]:

$$S_n = \{ (g_{i_1}g_{i_1-1} \dots g_{i_1-k_1})(g_{i_2}g_{i_2-1} \dots g_{i_2-k_2}) \dots (g_{i_p}g_{i_p-1} \dots g_{i_p-k_p}) \},\$$

for $1 \leq i_1 < \cdots < i_p \leq n-1$. In [28], the bases Σ'_n are used for constructing a Markov trace on $\mathcal{H} := \bigcup_{n=1}^{\infty} H_{1,n}$, and using this trace, a universal HOMFLYPT-type invariant for oriented links in ST is constructed.

Theorem 7 ([28, Theorem 6 and Definition 1]). Given z, s_k with $k \in \mathbb{Z}$ specified elements in $R = \mathbb{C}[q^{\pm 1}]$, there exists a unique linear Markov trace function on \mathcal{H}

$$\operatorname{tr}: \mathcal{H} \to R(z, s_k), \quad k \in \mathbb{Z}$$

determined by the rules

- (1) $\operatorname{tr}(ab) = \operatorname{tr}(ba)$ for $a, b \in H_{1,n}(q)$;
- (2) tr(1) = 1 for all $H_{1,n}(q)$;
- (3) $\operatorname{tr}(ag_n) = z\operatorname{tr}(a)$ for $a \in H_{1,n}(q)$;
- (4) $\operatorname{tr}(at'_n{}^k) = s_k \operatorname{tr}(a) \text{ for } a \in H_{1,n}(q), \ k \in \mathbb{Z}.$

Then, the function $X : \mathcal{L} \to R(z, s_k)$

$$X_{\widehat{\alpha}} = \Delta^{n-1} \cdot (\sqrt{\lambda})^e \operatorname{tr}(\pi(\alpha))$$

is an invariant of oriented links in ST, where $\Delta := -\frac{1-\lambda q}{\sqrt{\lambda}(1-q)}, \lambda := \frac{z+1-q}{qz}, \alpha \in B_{1,n}$ is a word in the σ_i 's and t'_i 's, $\widehat{\alpha}$ is the closure of α , e is the exponent sum of the σ_i 's in α , π the canonical map of $B_{1,n}$ to $H_{1,n}(q)$, such that $t \mapsto t$ and $\sigma_i \mapsto g_i$.

Remark 1. As shown in [28, 13] the invariant X recovers the HOMFLYPT skein module of ST. For a survey on the HOMFLYPT skein module of the lens spaces L(p, 1) via braids, the reader is referred to [14, 11].

2.2. The generalized Temperley-Lieb algebra of type B

We now introduce the analog of the (normalized) Kauffman bracket polynomial, V, for links in the solid torus via the generalized Temperley–Lieb algebra of type B.

Definition 2. The generalized Temperley-Lieb algebra of type B, $\text{TL}_{1,n}$, is defined as the quotient of the generalized Hecke algebra of type B, $H_{1,n}(q)$, over the ideal generated by the elements $\sigma_{i,i+1}$, $i \in \mathbb{N} \setminus \{0\}$ (recall Eq. (2)).

Remark 2. In [18], the analog of the normalized Kauffman bracket polynomial, V, for links in the solid torus ST has been derived from the Temperley–Lieb algebra of type B, TL_n^{B} . This algebra defined as a quotient of the Hecke algebra of type B, $H_{1,n}(q, Q)$, over the ideal generated by the $\sigma_{i,i+1}$ elements and

$$h_B := 1 + u \,\sigma_1 + v \,t + uv \,(\sigma_1 t + t\sigma_1) + u^2 v \,\sigma_1 t\sigma_1 + uv^2 \,t\sigma_1 t + (uv)^2 \,\sigma_1 t\sigma_1 t.$$

It is also worth mentioning that in [18] a different presentation for $H_{1,n}$ is used, that involves the parameters u, v and the quadratic relations

$$\sigma_i^2 = (u - u^{-1})\sigma_i + 1.$$
(5)

One can switch from one presentation to the other by a taking $\sigma_i = u\sigma_i$, t = vtand $q = u^2$. We will adapt this setting from now on.

Since the generalized Temperley–Lieb algebra of type B is a quotient of the generalized Hecke algebra of type B, we look for necessary and sufficient conditions so that the Markov trace defined in $H_{1,n}$ factors through to $TL_{1,n}$. We have the following result.

Theorem 8. The trace defined in $H_{1,n}$ factors through to $TL_{1,n}$ if and only if the trace parameters take the following values:

$$z = -\frac{1}{u(1+u^2)}.$$
 (6)

Proof. In order to evaluate the values of z so that the trace function factors through $TL_{1,n}$, we solve the equation $tr(g_{i,i+1}) = 0$ and we have

$$u^{3} \operatorname{tr}(g_{i} g_{i+1} g_{i}) = -1 - 2u z - 2u^{2} z^{2} \Leftrightarrow$$
$$u^{3} z \operatorname{tr}(g_{i}^{2}) = -1 - 2u z - 2u^{2} z^{2} \Leftrightarrow$$
$$u^{3} z^{2} (u - u^{-1}) + u^{3} z = -1 - 2u z - 2u^{2} z^{2} \Leftrightarrow$$
$$(uz + 1) (u^{3}z + uz + 1) = 0 \Leftrightarrow$$
$$z = -\frac{1}{u} \quad \text{or} \quad z = -\frac{1}{u (1 + u^{2})}.$$

As explained in [18], only the values in (6) are of topological interest and the proof is now concluded. $\hfill \Box$

Note now that for $z = -\frac{1}{u(1+u^2)}$, one deduces $\lambda = u^4$. Following [18], we obtain the following result.

Theorem 9. The following invariant is the universal invariant of the Kauffman bracket type, for knots and links in ST:

$$V_{\widehat{\alpha}}(u,v) := \left(-\frac{1+u^2}{u}\right)^{n-1} (u)^{2e} \operatorname{tr}(\overline{\pi}(\alpha)),$$

where $\alpha \in B_{1,n}$ is a word in the σ_i 's and t'_i ', s $\widehat{\alpha}$ is the closure of α , e is the exponent sum of the σ_i 's in α , $\overline{\pi}$ the canonical map of $B_{1,n}$ to $\mathrm{TL}_{1,n}$, such that $t \mapsto t$ and $\sigma_i \mapsto g_i$.

3. The Kauffman Bracket Skein Module of ST via Braids

In the braid setting, the elements of the standard basis of KBSM(ST) correspond bijectively to the elements of the following set:

$$B'_{\rm ST} = \{tt'_1 \dots t'_n, \ n \in \mathbb{N}\}.$$
(7)

The set B'_{ST} forms a basis of KBSM(ST) in terms of braids (for an illustration, see Fig. 8). Note that B'_{ST} is a subset of \mathcal{H} and, in particular, B'_{ST} is a subset of $\Sigma' = \bigcup_n \Sigma'_n$. Note also that in contrast to elements in Σ' , the elements in B'_{ST} have no gaps in the indices, the exponents are all equal to one and there are no "braiding tails".

The invariant V defined in Theorem 9 recovers KBSM(ST). Indeed, it gives distinct values to distinct elements of B'_{ST} , since $\operatorname{tr}(tt'_1 \dots t'_n) = s_1^n$. Hence, in order to compute the Kauffman bracket skein module of the lens spaces L(p,q), it suffices to extend the (most generic) invariant V for knots in L(p,q) following the ideas in [13–15]. That is, it suffices to solve the infinite system of equations (3), $V_{\widehat{\alpha}} = V_{\widehat{\text{bbm}_{\pm}(\alpha)}}$, for all α in a basis of KBSM(ST). These equations have particularly simple formulations with the use of the new basis, B_{ST} , for the Kauffman bracket skein module of ST first presented in [1].

We now present a different basis $B_{\rm ST}$ for the Kauffman bracket skein module of the solid torus, which is crucial toward the computation of KBSM(L(p, 1)) and which is described in Eq. (8) in open braid form. For an illustration, see Fig. 8. In particular, we have the following.

Theorem 10. The following set is a basis for KBSM(ST):

$$B_{\rm ST} = \{t^n, \ n \in \mathbb{N}\}.\tag{8}$$

The importance of the new basis $B_{\rm ST}$ of KBSM(ST) lies in the simplicity of the algebraic expression of braid band moves, which extend the link isotopy in ST to link isotopy in L(p,q) (recall Theorem 1(iv)). This was our motivation for establishing this new basis $B_{\rm ST}$. Note also that in [1] the same basis was derived using the Temperley–Lieb algebra of type B. In this section, we present a new and

 $The {\it Kauffman \ bracket \ skein \ module}$



Fig. 8. Elements in the different bases of KBSM(ST).

different proof for Theorem 10. The results presented here will also be used for the computation of KBSM(L(p, 1)) in Sec. 4. The method for proving Theorem 10 is the following:

- We first recall the total ordering defined in [13] for elements in the sets Σ' and Σ .
- We then pass from the set B'_{ST} consisting of monomials of the form $tt'_1t'_2...$ $t'_n, n \in \mathbb{N}$, to an augmented set B_{ST}^{aug} , consisting of monomials of the form t^n , where $n \in \mathbb{Z}$.
- We express elements in $B_{\rm ST}^{\rm aug} \setminus B_{\rm ST}$ to sums of elements in $B_{\rm ST}$.
- The two (ordered) sets $B'_{\rm ST}$ and $B_{\rm ST}$ are then related via a lower triangular infinite matrix with invertible elements on the diagonal.
- We conclude that the set $B_{\rm ST}$ forms a basis of KBSM(ST).

Before we proceed with the proof of Theorem 10, we introduce the notion of unoriented braids, which is crucial for working with unoriented knots on the "braid" level.

3.1. Unoriented braids

The Kauffman bracket polynomial is defined for unoriented knots, but when working on the braid level, there is a natural top-to-bottom orientation. Applying the Kauffman bracket skein relation on a crossing from standard braid results in merging strands with opposite orientations (see Fig. 9).

For this reason, we introduce the notion of *unoriented braids* as follows.

Definition 3. An *unoriented braid* is defined as a classical braid by ignoring the natural top-to-bottom orientation.

Remark 3. The concept of unoriented braids is new and it seems promising in serving as a tool for studying knots in c.c.o. 3-manifolds and for deriving invariants for 3-manifolds in general. For a thorough study of unoriented braids, see [16].

The significance of unoriented braids for the Kauffman bracket skein module of the solid torus (and for Kauffman bracket skein modules in general), is illustrated in Fig. 10, where the looping generators t and t^{-1} are shown to be "equivalent" in KBSM(ST) (compare also with Fig. 14).

From now on, by braids we shall mean unoriented braids unless otherwise stated.



Fig. 9. The skein relation on the braid level.



Fig. 10. Unoriented braids and KBSM(ST).

3.2. An ordering relation

We now present an ordering relation defined on Σ'_n , B'_{ST} and Σ_n . We first introduce the notion of the index of a word in B'_{ST} , in Σ'_n and in Σ_n .

Definition 4. The *index* of a word τ in B'_{ST} (or in Σ'_n or in Σ_n), denoted $ind(\tau)$, is defined to be the highest index of the t'_i 's (respectively, of the t_i 's), in τ . Similarly, the *index* of an element in Σ'_n or in Σ_n is defined in the same way by ignoring possible gaps in the indices of the looping generators and by ignoring the braiding part in H_n . Moreover, the index of a monomial in H_n and in $TL_{1,n}$ is equal to 0.

For example, $\operatorname{ind}(t'^{k_0}t'_1^{k_1}\dots t'_n^{k_n}) = \operatorname{ind}(t^{u_0}\dots t^{u_n}_n) = n.$

We now proceed with presenting an ordering relation in the sets Σ and Σ' , which passes to their respective subsets $B_{\rm ST}$ and $B'_{\rm ST}$.

Definition 5 ([13, Definition 2]). Let $w = t'_{i_1}^{k_1} \dots t'_{i_{\mu}}^{k_{\mu}} \cdot \beta_1$ and $u = t'_{j_1}^{\lambda_1} \dots t'_{j_{\nu}}^{\lambda_{\nu}} \cdot \beta_2$ in Σ' , where $k_t, \lambda_s \in \mathbb{Z}$ for all t, s and β_1, β_2 are standard braid words. Then, we define the following ordering in Σ' :

(a) If $\sum_{i=0}^{\mu} k_i < \sum_{i=0}^{\nu} \lambda_i$, then w < u. (b) If $\sum_{i=0}^{\mu} k_i = \sum_{i=0}^{\nu} \lambda_i$, then: (i) if ind(w) < ind(u), then w < u; (ii) if ind(w) = ind(u), then: (α) if $i_1 = j_1, \dots, i_{s-1} = j_{s-1}, i_s < j_s$, then w > u; (β) if $i_t = j_t$ for all t and $k_{\mu} = \lambda_{\mu}, k_{\mu-1} = \lambda_{\mu-1}, \dots, k_{i+1} = \lambda_{i+1}, |k_i| < |\lambda_i|$, then w < u;

(γ) if $i_t = j_t$ for all t and $k_\mu = \lambda_\mu, k_{\mu-1} = \lambda_{\mu-1}, \dots, k_{i+1} = \lambda_{i+1}, |k_i| = |\lambda_i|$ and $k_i > \lambda_i$, then w < u; (δ) if $i_t = j_t \forall t$ and $k_i = \lambda_i, \forall i$, then w = u.

The ordering in the set Σ is defined as in Σ' , where t'_i 's are replaced by t_i 's.

Definition 6. We define the augmented sets $B_{\rm ST}^{\rm aug}$ and $B_{\rm ST}'^{\rm aug}$ as follows:

$$B_{\rm ST}^{\rm aug} := \{t^n, n \in \mathbb{Z}\},\$$
$$B'_{\rm ST}^{\rm aug} := \{t^{k_0} t'_1^{k_1} \dots t'_m^{k_m}, k_i \in \mathbb{Z}, \forall i\}.$$

Proposition 1. The sets $B_{\rm ST}$, $B_{\rm ST}$, $B_{\rm ST}$ and $B'_{\rm ST}$ and $B'_{\rm ST}$ equipped with the ordering relation of Definition 5, are totally ordered sets. Moreover, the sets $B_{\rm ST}$ and $B'_{\rm ST}$ are well-ordered sets.

Proof. In [13, Proposition 1], it is shown that the sets Σ_n' and Σ_n equipped with the ordering relation of Definition 5, are totally ordered sets. Since $B'_{\rm ST}$ and $B'_{\rm ST}$ are subsets of Σ_n' , and since $B_{\rm ST}$ and $B_{\rm ST}^{\rm aug}$ are subsets of Σ_n , the sets $B_{\rm ST}$, $B_{\rm ST}^{\rm aug}$, $B'_{\rm ST}$ and $B'_{\rm ST}^{\rm aug}$ inherit the property of being totally ordered sets from Σ_n (cor. Σ'_n).

Moreover, t^0 is the minimum element of both B_{ST} and B'_{ST} and thus, these sets are well-ordered sets.

3.3. Useful lemmas

In this section, we prove a series of results in order to convert elements in B'_{ST} to elements in B_{ST} . We will use the symbol $\widehat{\cong}$ when conjugation and stabilization



Fig. 11. Conjugation on the t'_i 's.

moves are performed, and \cong_{skein} when both conjugation, stabilization moves and the Kauffman bracket skein relation is performed. Note that most of the times and in order to simplify our results, we will omit the coefficients involved. Finally, it is worth mentioning that the t'_i 's are conjugates, i.e. $t'_i t'_j \cong t'_j t'_i$ for all $i \neq j$. A general case is illustrated in Fig. 11, for $w \in B'_{ST}$ ^{aug}.

In order to simplify the algebraic expressions obtained throughout this procedure and throughout the paper in general, we first introduce the following notation.

Notation 2. We set $\tau'_{i,i+m}^{k_{i,i+m}} := t'^{k_i}_i \dots t'^{k_{i+m}}_{i+m} \in B'_{\mathrm{ST}}^{\mathrm{aug}}$, for $m \in \mathbb{N}$, $k_j \neq 0$ for all j.

Consider now for example elements of the form $t^k t'_1^m$, such that $k, m \in \mathbb{N}$ and $k \geq m > 0$. As illustrated in Fig. 12, applying the skein relations and conjugation, we may express these elements in terms of elements in the basis B_{ST} , together with elements of the form $t^{k-i} t'_1^{m-i}$, i > 0. The process is repeated until we are left with elements in B_{ST} .



$$= -A^{2}t^{k+m} + t^{k+m-2} + A^{2}t^{k-1}t'_{1}^{m-1}$$

Fig. 12. $t^{k}t'_{2}^{m} \cong -A^{2}t^{k+m} + t^{k+m-2} + A^{2}t^{k-1}t'_{2}^{m-1}$.

Applying induction on the exponents k and m and using the ordering relation defined in Definition 5, we obtain the following result.

Lemma 1. For $k, m \in \mathbb{N}$ and $k \ge m > 0$, the following relation hold in $\mathrm{TL}_{1,n}$:

$$t^{k} t_{1}^{\prime m} \widehat{\cong} - A^{2} t^{k+m} + \sum_{i=1}^{\lfloor m/2 \rfloor} A^{2(i-1)} (1 - A^{4}) t^{k+m-2i} + A^{2m} t^{k-m}.$$

Remark 4. In Lemma 1, we considered elements of the form $t^k t'_1^m$, where $k \ge m > 0$. If k < m, then since the t'_i 's are conjugates, we order the exponents on the closure of the mixed braids and we braid the resulting loops again (recall Fig. 11). Namely, $t^k t'_1^m \cong t^m t'_1^k$. Thus, from now on we will be using Lemma 1 without the restriction on the exponents.

Lemma 1 and Remark 4 lead to the following result.

Corollary 1. Let $w \in B'_{ST}$ such that ind(w) = n - 1. Then, for $k \ge m$ in \mathbb{Z} the following relations hold in KBSM(ST):

$$w t'_n^{k} t'_{n+1}^{m} \widehat{\cong} \sum_{\text{skein}}^m \sum_{i=0}^m a_i \cdot w t'_n^{k+m-2i},$$

where a_i are coefficients for all i.

Note that when applying Corollary 1 on an element in $B'_{\rm ST}$, we obtain a sum of elements in $B'_{\rm ST}$ ^{aug} of lower index than the initial element in $B'_{\rm ST}$. We now convert elements in $B'_{\rm ST}$ ^{aug} to sums of elements in $B'_{\rm ST}$. We have the following result.

Lemma 2. The following relations hold in KBSM(ST):

$$\tau'^{k_{0,n}}_{0,n} \widehat{\cong}_{\text{skein}} \sum_{i < k} a_i t^i,$$

where $k = \sum_{i=0}^{n} k_i$, $k_i \in \mathbb{N}$ and a_i coefficients for all *i*.

Proof. We prove Lemma 2 by strong induction on the order of $\tau'_{0,n}^{k_{0,n}}$.

The base of the induction is the monomial $tt'_1 \in B'_{ST}$ of index 1, and it is illustrated in Fig. 13.



Fig. 13. The base of induction.

2250076-18

Assume now that Lemma 2 holds for all monomials that are of lower order than $\tau'_{0,n}^{k_{0,n}}$. Then, we have

$$\tau_{0,n}^{\prime k_{0,n}} = \tau_{0,n-2}^{\prime k_{0,n-2}} \underline{t_{n-1}^{\prime k_{n-1}} t_{n}^{\prime k_{n}}} \stackrel{\text{Cor. 1}}{\cong} \frac{1}{\sum_{i=0}^{k_{n}}} \tau_{0,n-2}^{\prime k_{0,n-2}} t_{n-1}^{\prime k_{n-1}+k_{n}-2i}$$

According to the ordering relation, on the right-hand side of this equation we have monomials of index $n - 1 < n = \operatorname{ind}(\tau'_{0,n}^{k_{0,n}})$. Thus, these monomials are of lower order than $\tau'_{0,n}^{k_{0,n}}$. By the induction hypothesis the proof is concluded.

Remark 5. Figure 12 suggests that Lemma 2 is true for $k_i \in \mathbb{Z}$, that is,

$$\tau'_{0,n}^{k_{0,n}} \widehat{\cong}_{\substack{s \in in \\ s \notin in}} \sum_{i \le k} a_i t^i, \tag{9}$$

where $k = \sum_{i=0}^{n} |k_i|, k_i \in \mathbb{Z}$ and a_i coefficients for all i.

Applying now Lemma 9 (together with Eq. (9) when needed) on elements in B'_{ST} leads to the following result.

Corollary 2. Elements in B'_{ST} can be written as sums of elements in B_{ST}^{aug} . In particular,

$$\tau_{0,n}' \stackrel{\widehat{\cong}}{\underset{i \le n+1}{\cong}} \sum_{i \le n+1} a_i t^i, \tag{10}$$

where a_i coefficients for all *i*. Equivalently, the set B_{ST}^{aug} spans KBSM(ST).

In the following example, we apply Lemma 9 and Eq. (9) to some elements in the standard basis $B'_{\rm ST}$.

Example 1.

$$\begin{split} t\,t_1'\, & \stackrel{\cong}{\underset{\text{skein}}{\cong}} \,1+t^2, \\ t\,t_1'\,t_2'\, & \stackrel{\cong}{\underset{\text{skein}}{\cong}} \,t\,t_1'^2 + t\,\stackrel{\cong}{\underset{\text{skein}}{\cong}} t^{-1} + t + t^3, \\ t\,t_1'\,t_2'\,t_3'\, & \stackrel{\cong}{\underset{\text{skein}}{\cong}} \,t\,t_1'\,t_2'^2 + t\,t_1'\,\stackrel{\cong}{\underset{\text{skein}}{\cong}} t\,t_1'^{-1} + t\,t_1'\stackrel{\cong}{\underset{\text{skein}}{\cong}} t^{-2} + 1 + t^2 + t^4, \\ t\,t_1'\,t_2'\,t_3'\,t_4'\, & \stackrel{\cong}{\underset{\text{skein}}{\cong}} \,t\,t_1'\,t_2'\,t_3'^2 + t\,t_1'\,t_2'\stackrel{\cong}{\underset{\text{skein}}{\cong}} t\,t_1'\,t_2'^3 + tt_1'\,t_2'^{-1} + tt_1't_2' \\ & \stackrel{\cong}{\underset{\text{skein}}{\cong}} \,t\,t_1'^4 + t\,t_1'^{-2} + t + t\,t_1'^2 + t\,t_1'\,t_2' \\ & \stackrel{\cong}{\underset{\text{skein}}{\cong}} t^{-3} + t^{-1} + t + t^3 + t^5. \end{split}$$

Following the calculations in Example 1, one may notice that there are "closed formulas" for Eq. (10), by ignoring the coefficients. More precisely, simple calculations lead to the following result.

Lemma 3. For $n \in \mathbb{N}$, the following relations hold in KBSM(ST):

$$\tau_{0,n}' \cong \begin{cases} \sum_{i=0}^{(n+1)/2} a_i t^{2i} + \sum_{i=1}^{(n-1)/2} b_i t^{-2i} & \text{for } n \text{ odd,} \\ \sum_{i=0}^{n/2} c_i t^{2i+1} + \sum_{i=1}^{n/2} d_i t^{-2i+1} & \text{for } n \text{ even,} \end{cases}$$

where a_i, b_i, c_i, d_i are coefficients.

3.4. Dealing with negative exponents

We now deal with monomials in $B_{\rm ST}^{\rm aug}$ with negative exponents, i.e. elements in $B_{\rm ST}^{\rm aug} \setminus B_{\rm ST}$, and express them to sums of elements in $B_{\rm ST}$. As mentioned before, due to the fact that the Kauffman bracket does not take orientation into consideration, we use unoriented braids. In the figures that follow, we start with standard braids and we ignore the orientation when a Kauffman bracket skein relation is performed, resulting in unoriented braids. Note that we will be using the same notation for standard braids and unoriented braids. In Fig. 14, the cases of t^{-1} and t^{-2} are illustrated and it is important to note that the resulting monomial tt'_1 , that



Fig. 14. Converting t^{-1} and t^{-2} to elements in B'_{ST} .

appears when treating t^{-2} , is an unoriented braid. Finally, it is worth mentioning that in order to demonstrate the importance of the new proposed method for computing Kauffman bracket skein modules, we will follow a more algebraic procedure for converting elements in $B_{\rm ST}^{\rm aug} \setminus B_{\rm ST}$ to elements in $B_{\rm ST}$ using the Kauffman bracket skein relation, instead of straightforward using the fact that $t^{-n} \sim t^n$, for all $n \in \mathbb{N}$ on the unoriented braid level.

Note that in the case of t^{-2} , when we apply the skein relation for the first time, a closed looping \hat{t} is "isolated". We then apply again the skein relation by ignoring \hat{t} (readers may find useful to think of it as being shrunk). Recall also that $tt'_1 \cong 1 + t^2$ (recall Fig. 13). Thus, $t^{-2} \cong 1 + t^2$ in KBSM(ST).

Let us consider now the general case for $n \in \mathbb{N}$. We first perform an RI move on the top of the looping generator creating a kink. We then apply the Kauffman bracket skein relation on the resulting crossing and the result is $t^{-n+2} + t^{-n+1} t'_1$. We apply the same procedure on the resulting monomials. Note that in the monomial $t^{-n+1} t'_1$, the looping t'_1 will not interfere with the above-mentioned process on the closure $t^{-n+1} t'_1$. We continue until all negative exponents become either zero or one. Obviously, this depends on n being odd or even. Hence, we have the following result.

Lemma 4. For $n \in \mathbb{N}$, the following relations hold in KBSM(ST):

$$t^{-n} \widehat{\cong}_{\text{skein}} \begin{cases} \sum_{i=0}^{\frac{n-1}{2}} a_i \tau'_{0,2i} & \text{for } n \text{ odd,} \\ \\ \sum_{i=0}^{\frac{n}{2}} a'_i \tau'_{0,2i} & \text{for } n \text{ even,} \end{cases}$$

where a_i, a'_i coefficients.

It is crucial now to note that when we apply Lemma 3 on a monomial τ' in $B'_{\rm ST}$, we obtain monomials in $B_{\rm ST}$ and in $B_{\rm ST}^{\rm aug} \backslash B_{\rm ST}$. We then apply Lemma 4 on the resulting monomials in $B_{\rm ST}^{\rm aug} \backslash B_{\rm ST}$ and we obtain elements in $B'_{\rm ST}$ again, but of lower order than the initial monomial τ' according to the ordering relation defined in Definition 5. Thus, if we continue this way, we will eventually reach at elements in $B_{\rm ST}$. Indeed, we have the following result.

Proposition 2. For $n \in \mathbb{N}$, the following hold in KBSM(ST):

$$t^{-n} \widehat{\cong} \sum_{i=0}^{n} a_i t^i,$$

where a_i are coefficients for all *i*.

Proof. Let n be an odd natural number. The case that n is even follows similarly. We have that

$$t^{-n} \overset{\text{Lemma } 4}{\sim} \frac{1}{\sum_{i=0}^{n-1}} a_i \tau'_{0,2i} \overset{\text{Lemma } 3}{\sim} \frac{1}{\sum_{i=0}^{n-1}} a_i \left(\sum_{j=0}^{i} a'_i t^{2j+1} + \sum_{j=1}^{i} a''_i t^{-2j+1} \right)$$
$$= \left(\underbrace{\sum_{i=0}^{n-1} \sum_{j=0}^{i} b_i t^{2j+1}}_{A} \right) + \left(\underbrace{\sum_{i=0}^{n-1} \sum_{j=1}^{i} b'_i t^{-2j+1}}_{B} \right),$$

where a_i, a'_i, a''_i, b_i and b'_i are coefficients for all *i*.

We now deal with terms in B that consist of elements in $B_{\rm ST}^{\rm aug} \setminus B_{\rm ST}$, since all terms in A are in $B_{\rm ST}$. The highest exponent in the terms in B is (-n+3>-n) and thus, we may continue applying Lemmas 3 and 4 until we eliminate all negative exponents. All resulting monomials will eventually belong to $B_{\rm ST}$ and the proof is concluded.

Remark 6. It is possible to find a "closed formula" for t^{-n} , $n \in \mathbb{N}$ in KBSM(ST). It is very challenging to keep track of the coefficients, but simple calculations lead to the following result:

$$t^{-n} \widehat{\cong}_{\text{skein}} \begin{cases} \sum_{i=0}^{n/2} a_i t^{2i} & \text{ for } n \text{ even,} \\ \frac{n-1}{2} \\ \sum_{i=0}^{n-1} a'_i t^{2i+1} & \text{ for } n \text{ odd,} \end{cases}$$

where a_i, a'_i coefficients, for all *i*.

3.5. The new basis of KBSM(ST)

We are now in position to prove the main result of this section. More precisely

Theorem 11. The set B_{ST} forms a basis for the Kauffman bracket skein module of the solid torus.

Proof. Consider $\tau'_{0,n} \in B'_{ST}$ of index *n*. We apply Corollary 2 and obtain a sum of elements in B_{ST}^{aug} with (n+1) being the highest exponent appearing in the sum. More precisely, for *n* odd we have

$$B_{\mathrm{ST}}' \ni \tau_{0,n}' \underset{\mathrm{skein}}{\cong} \underbrace{\sum_{i=0}^{n+1}}_{\in B_{\mathrm{ST}}, \forall i} a_i t^{2i} + \underbrace{\sum_{i=1}^{n-1}}_{\in B_{\mathrm{ST}}^{\mathrm{aug}}} \underbrace{\widehat{a}_i t^{-2i}}_{\mathrm{skein}} \underbrace{\widehat{\sum_{i=0}^{n+1}}}_{\in B_{\mathrm{ST}}, \forall i} a_i t^{2i} + \underbrace{\sum_{i=1}^{n-1}}_{\in B_{\mathrm{ST}}} \underbrace{\sum_{i=0}^{n+1}}_{\in B_{\mathrm{ST}}} a_i t^{2i} + \underbrace{\sum_{i=1}^{n-1}}_{\in B_{\mathrm{ST}}} \underbrace{\sum_{i=0}^{n+1}}_{\in B_{\mathrm{ST}}} a_i t^{2i} + \underbrace{\sum_{i=1}^{n-1}}_{\in B_{\mathrm{ST}}} a_i t^{2i} + \underbrace{\sum_{i=1}^{n-1}}_{E_{\mathrm{ST}}} a_i t^{2i} + \underbrace{\sum_{i=1}^{n-1}}_{E_{\mathrm{ST}}} a_i t^{2i} + \underbrace{\sum_{i=1}^{n-1}}_{E$$

icles.

The monomial of highest order in the sum $\sum_{i=1}^{\frac{n-1}{2}} (\sum_{j=0}^{i} a'_{i} \tau'_{0,2j})$ is $\tau'_{0,n-1}$, which according to the ordering relation of Definition 5, is of lower order than the initial monomial $\tau'_{0,n}$. Hence, if we continue applying the same procedure on the resulting monomials, we will eventually reach at a sum of elements in $B_{\rm ST}$, with (n+1) being the highest exponent.

Thus, we have proved that any monomial τ' in B'_{ST} with $\operatorname{ind}(\tau') = n$, can be written as a sum of elements in B_{ST} and in particular, $\tau' \cong \sum_{\substack{s \in in \\ s \neq i = 0}}^{n+1} a_i t^i$, for some coefficients a_i . Moreover, t^{n+1} is the monomial of the highest order appearing when converting the monomial $\tau' \in B'_{ST}$ of index n to elements in B_{ST} . Hence, the sets B'_{ST} and B_{ST} are related via an infinite lower triangular matrix with invertible elements in the diagonal. Thus, the set B_{ST} forms a basis of KBSM(ST).

4. A Basis for the Kauffman Bracket Skein Module of L(p, 1) via Braids

In this section, we solve the infinite system of equations (3) for q = 1, which is equivalent to computing the Kauffman bracket skein module of the lens spaces L(p, 1). Recall that this infinite system of equations is obtained from elements in the basis of KBSM(ST), $B_{\rm ST}$, by performing braid band moves and then by imposing to the generic invariant V for knots and links in ST relations of the form $V_{\hat{t}^n} = V_{\widehat{\text{bbm}(t^n)}}$, for all $n \in \mathbb{N}$, where $\text{bbm}(t^n) = t^p t_1^n \sigma_1^{\pm 1}$ (recall Theorem 6). Recall also that the unknowns in the system are the s_i 's, coming from the fourth rule of the trace function in Theorem 7, that is, $\text{tr}(t^n) = s_n$, for all $n \in \mathbb{Z}$.

4.1. The infinite system

In this section, we present a series of results toward the solution of the infinite system. Our main task is the evaluation of $\operatorname{tr}(t^p t_1^n \sigma_1^{\pm 1})$ for $n \in \mathbb{N}$. Recall now that the trace function is defined using the t'_i 's and not the t_i 's that we obtain when performing a bbm on elements in B_{ST} (recall the fourth rule of Theorem 7). Thus, our first task is to express the monomials $t^p t_1^n \sigma_1^{\pm 1} \in \Sigma$ to sums of monomials in Σ' (recall Eq. (4)). In [13], a method for expressing monomials in Σ' to sums of monomials in Σ (and vice-versa) is presented on the generalized Hecke algebra of type B, $H_{1,n}$, level. More precisely, it is shown that a monomial $\tau \in \Sigma$ can be written as a sum of elements in Σ' , such that the homologous word of τ in Σ' , i.e. the monomial τ where all t_i 's are replaced by t'_i 's, is the monomial of the highest order appearing in the resulting sum. All other monomials appearing in the sum have smaller index, or equal index but different exponents (recall the ordering relation of Definition 5). Finally, the braiding "tails", i.e. the σ_i 's that come after the t_i 's in monomials in Σ , are eliminated in this process.

Since now the generalized Temperley–Lieb algebra of type B, $TL_{1,n}$, is a quotient of $H_{1,n}$ over the ideal generated by elements in Eq. (2) that only involves the σ 's, and since the only braiding generator in the monomials $bbm(t^n)$ is $\sigma_1^{\pm 1}$, it follows that the results presented in [13] are also true in $TL_{1,n}$ for the monomials $bbm(t^n) = t^p t_1^n \sigma_1^{\pm 1}$. Hence, we obtain the following results.

Lemma 5. For $n, p \in \mathbb{N} \setminus \{0\}$, the following holds in $\mathrm{TL}_{1,n}$:

$$t^p t_1^n \sigma_1 \cong \sum_{i=0}^n a_i t^{p+i} t_1'^{n-i}$$
 and $t^p t_1^n \sigma_1^{-1} \cong \sum_{i=1}^n b_i t^{p+i} t_1'^{n-i}$,

where a_i, b_i coefficients for all *i*.

Note that in the monomial of highest order in $\sum_{i=0}^{n} a_i t^{p+i} t'_1^{n-i}$ is the homologous monomial of $t^p t_1^n$, $t^p t'_1^n$. In order now to evaluate $\operatorname{tr}(\sum_{i=0}^{n} a_i t^{p+i} t'_1^{n-i})$, we first perform Lemma 1 on the monomials in the sum and we obtain the following:

$$t^{p} t_{1}^{\prime n} \widehat{\cong} \sum_{i=0}^{\lfloor n/2 \rfloor} a_{i} t^{p+n-2i}, \qquad (11)$$

where a_i are coefficients for all i.

Hence, we have that $\operatorname{tr}(t^p t_1^n \sigma_1) = \sum_{i=0}^{\lfloor n/2 \rfloor} a_i s_{p+n-2i}$ for some coefficients a_i , since

$$\operatorname{tr}(t^{p} t_{1}^{n} \sigma_{1}) = \operatorname{tr}\left(\sum_{i=0}^{n} a_{i} t^{p+i} t_{1}^{\prime n-i}\right) = \operatorname{tr}\left(\sum_{i=0}^{n} \sum_{j=0}^{\lfloor \frac{n-i}{2} \rfloor} (a_{i}^{\prime} t^{p+n-2j})\right).$$

Remark 7. Consider now a negative braid band move applied on elements in $B_{\rm ST}$, i.e. $t^n \to t^p t_1^n \sigma_1^{-1}$. Observe that $t^p t_1^n \sigma_1^{-1} = t^p t_1^{n-1} \sigma_1 t \cong t^{p+1} t_1^{n-1} \sigma_1$. Thus, applying Lemma 5 and Eq. (11) on $t^{p+1} t_1^{n-1} \sigma_1$ we obtain

$$t^{p+1} t_1^{n-1} \sigma_1 \stackrel{\cong}{\cong} \sum_{i=0}^{n-1} a_i t^{p+1+i} t_1'^{n-1-i} \stackrel{\cong}{\cong} \sum_{i=0}^{n-1} \sum_{j=0}^{\lfloor \frac{n-1-i}{2} \rfloor} \left(a_i' t^{p+n-2j} \right) + \frac{1}{2} \left(a_$$

for some coefficients a'_i . Evaluating now the trace of these elements, we have that

$$\operatorname{tr}(t^p t_1^n \sigma_1^{-1}) = \sum_{i=0}^{n-1} \sum_{j=0}^{\lfloor \frac{n-1-i}{2} \rfloor} (a'_i s_{p+n-2j}).$$

Less number of unknowns of the infinite system appear in the equation $V_{t\hat{n}} = V_{\overline{\text{bbm}_{-}(t^n)}}$, compared to the equation $V_{t\hat{n}} = V_{\overline{\text{bbm}_{+}(t^n)}}$, and thus, we consider only the positive braid band moves for obtaining the equations of the infinite system for $n \in \mathbb{N}$. Another reason for that is the relation between a positive (respectively, a negative) braid band move and a band move of type β . More precisely, consider $bbm_{+}(t^n) = t^p t_1^n \sigma_1$, and apply the skein relation on the braiding generator σ_1 . The result is a sum of two terms, one of which is the result of the performance of a β band move on the closure. Recall now that one type of band moves suffices to translate isotopy in L(p, q) and the result follows.

4.2. Dealing with indices greater than p

We are now in position to prove the first fundamental result toward the solution of the infinite system. We have the following.

Proposition 3. For $n \in \mathbb{N}$ the following relations hold in KBSM(L(p, 1)) for $n \in \mathbb{N}$:

$$s_{p+n} = \begin{cases} \sum_{i=0}^{n/2} (c_i \, s_{2i} \, + \, d_i \, s_{p-2i}) & \text{for } n \text{ even,} \\ \\ \sum_{i=0}^{(n-1)/2} (c_i' \, s_{2i-1} + d_i' \, s_{p-2i-1}) & \text{for } n \text{ odd,} \end{cases}$$

where c_i, d_i, c'_i, d'_i are coefficients.

Proof. The following diagram describes how the equations of the infinite system are obtained:



Hence, $s_n = \left(-\frac{1+u^2}{u}\right) u^{2e} \cdot \sum_{i=0}^n s_{p-n+2i}$, for all $n \in \mathbb{N}$. We now consider the following cases:

• For $n = 2k, k \in \mathbb{N}$, we have

÷

For
$$n = 0$$
: $s_0 = s_p \Leftrightarrow s_p = s_0 := 1$,
For $n = 2$: $s_2 = s_{p-2} + s_p + s_{p+2} \Leftrightarrow s_{p+2} = s_0 + s_2 + s_{p-2}$,
For $n = 4$: $s_4 = s_{p-4} + s_{p-2} + s_p + s_{p+2} + s_{p+4} \Leftrightarrow s_{p+4} = s_0 + s_2 + s_4 + s_{p-4} + s_{p-2}$,

• For $n = 2k + 1, k \in \mathbb{N}$, we have

÷

For
$$n = 1$$
: $s_1 = s_{p-1} + s_{p+1} \Leftrightarrow s_{p+1} = s_1 + s_{p-1}$,
For $n = 3$: $s_3 = s_{p-3} + s_{p-1} + s_{p+1} + s_{p+3} \Leftrightarrow s_{p+3} = s_1 + s_3 + s_{p-3} + s_{p-1}$

For
$$n = 2k + 1 : \Leftrightarrow s_{p+2k+1} = \sum_{i=0}^{k} (s_{2i+1} + s_{p-2i-1}).$$

The proof is concluded.

An immediate result of Proposition 3 is that every unknown of the infinite system with index greater than or equal to p, can be written as a sum of the unknowns with index less than p. That is

Corollary 3. For $k \in \mathbb{N}$, the following hold in KBSM(L(p, 1)):

$$s_{p+k} = \sum_{i < p} q_i s_i$$
, where $i \in \mathbb{Z}$ and q_i coefficients.

Remark 8. Note that since $tr(t^n) = s_n$, we have proved that the loop elements $t^{p+i} \in \text{KBSM}(\text{ST}), i \in \mathbb{N}$, can be expressed as a sum of elements in the form $t^{p-j}, j \in \mathbb{N}$. Roughly speaking, the exponents of the *t*'s are mod*p* in KBSM (L(p, 1)).

It is important to observe now that depending on p, the exponents on some monomials of the resulting sum may be negative. We shall deal with negative indices in the following section.

4.3. Dealing with negative indices

We now deal with the unknowns of the infinite system that have negative indices and express them in terms of s_i 's where $i \in \mathbb{N}$. Recall that $s_{-n} = \operatorname{tr}(t^{-n})$ and that $t^{-n} \stackrel{\sim}{\cong} \sum_{i=0}^{n} a_i t^i$ (Proposition 2). Thus, we have that

$$s_{-n} = \operatorname{tr}(t^{-n}) = \operatorname{tr}\left(\sum_{i=0}^{n} a_i t^i\right) = \sum_{i=0}^{n} a_i s_i.$$

If instead of Proposition 2 we use relations from Remark 6, we obtain the following.

 $s_{-n} = \begin{cases} \sum_{i=0}^{n/2} s_{2i} & \text{for } n \text{ even,} \\ \\ \sum_{i=0}^{(n-1)/2} s_{2i+1} & \text{for } n \text{ odd.} \end{cases}$

for some coefficients a_i, b_i . Omitting the coefficients we have that

$$s_{-2} = s_0 + s_2, \quad s_{-4} = s_0 + s_2 + s_4, \dots, s_{-j} = s_0 + s_2 + \dots + s_j$$
 for j even and
 $s_{-1} = s_1, \quad s_{-3} = s_1 + s_3, \dots, s_{-j} = s_1 + s_3 + \dots + s_j$ for j odd.

The result follows.

4.4. The solution of the infinite system

We are now in position to present the solution of the infinite system that is equivalent to computing the Kauffman bracket skein module of the lens spaces L(p, 1). We first generalize Proposition 3 and in particular we show that the unknowns with indices greater than or equal to p can be written as sums of unknowns with non-negative indices and also that these indices are mod p. Indeed, we have the following.

Proposition 4. The following relations hold in KBSM(L(p, 1)) for all $n \in \mathbb{N}$:

$$s_{p+n} = \sum_{i=0}^{p} a_i s_i.$$

Proof. For $n \leq p$ the results follows from Proposition 3. Let now n > p and consider the case where n is even. The case n being odd follows similarly.

Let $j \in \mathbb{N} \setminus \{0\}$ such that n = p + j. Omitting the coefficients, we have

$$s_{p+n} \stackrel{\text{Proposition } 3}{=} \sum_{i=0}^{n/2} (s_{2i} + s_{p-2i})$$

$$= \underbrace{[(s_0 + s_p) + (s_2 + s_{p-2}) + \dots + (s_p + s_0)]}_A$$

$$= \underbrace{[(s_{p+2} + s_{-2}) + (s_{p+4} + s_{-4}) + \dots + (s_{p+j} + s_{-j})]}_B$$

Note now that all indices in terms in A are in $\{0, 1, \ldots, p\}$ and thus, we focus on terms in B, i.e. terms of the form s_{-k} and s_{p+k} for $k \in \{1, 2, \ldots, j\}$.

We distinguish the following cases:

- **Case I.** We deal first with terms of the form s_{-k} for $k \in \{1, 2, ..., j\}$. We apply Lemma 6 on these terms and we obtain sums of unknowns with all indices being less than or equal to j. Thus, if j < p the result follows. If j > p, we repeat the same procedure until we reduce all indices mod p.
- **Case II.** We now deal with terms of the form s_{p+k} for $k \in \{1, 2, ..., j\}$. We apply Proposition 3 and obtain elements in s_i 's such that i is either in $\mathbb{Z}\setminus\mathbb{N}$ (see Case I), or in \mathbb{N} such that i < p. Continuing that way, we eventually obtain a sum of elements of the form $\sum_{i=0}^{p} s_i$.

We shall now reduce the indices in the unknowns further. Let

$$I = \{0, 1, \dots, \lfloor p/2 \rfloor\},\$$

where $\lfloor k \rfloor$ denotes the greatest integer less than or equal to k. We have

Theorem 12. For all $n \in \mathbb{N} \setminus I$, we have that $s_n = \sum_{i \in I} s_i$.

Proof. From Proposition 4, it follows that it suffices to show that $s_n = \sum_{i \in I} s_i$ for $n \in \{\lfloor p/2 \rfloor + 1, \ldots, p\}$.

Consider elements in $B_{\rm ST}^{\rm aug} \backslash B_{\rm ST}$ of the form t^{-n} for $n \in I$. We perform bbm's on these elements in order to obtain equations for the infinite system. Using Lemma 6 and Proposition 3, we may convert the unknowns with negative indices to sums of unknowns with indices in I and thus, we conclude that the unknowns with indices I suffice to generate all unknowns s_k , where $k \notin I$. Equivalently, the elements t^k , where $k \notin I$, can be expressed as sums of elements of the form t^n , where $n \in I$. More precisely, we have

 $\begin{array}{cccc} \bullet t^{0} \stackrel{\text{bbm}}{\to} t^{p} \sigma_{1}^{\pm 1} & \stackrel{V}{\Rightarrow} 1 := s_{0} = s_{p} & \Rightarrow & s_{p} = s_{0}, \\ \bullet t^{-1} \stackrel{\text{bbm}}{\to} t^{p} t_{1}^{-1} \sigma_{1}^{\pm 1} \stackrel{V}{\Rightarrow} s_{-1} = s_{p-1} & \stackrel{\text{Lemma } 6}{\underset{s_{-1} = s_{1}}{\Rightarrow}} & s_{p-1} = s_{1}, \\ \bullet t^{-2} \stackrel{\text{bbm}}{\to} t^{p} t_{1}^{-2} \sigma_{1}^{\pm 1} \stackrel{V}{\Rightarrow} s_{-2} = s_{p-2} + s_{p} + s_{p+2} & \stackrel{\text{Lemma } 6 \text{ and Proposition } 4}{\underset{s_{-2} = s_{0} + s_{2}}{\Rightarrow}} & s_{p-2} = \sum_{i=0}^{p-3} s_{i}, \\ \begin{cases} s_{-2} = s_{0} + s_{2} \\ s_{p+2} = \sum_{p}^{i=0} s_{i} \end{cases} \end{cases} \\ \vdots \end{array}$

•
$$t^{-\lfloor p/2 \rfloor + 1} \xrightarrow{\text{bbm}} t^p t_1^{-\lfloor p/2 \rfloor + 1} \sigma_1^{\pm 1} \xrightarrow{V} s_{-\lfloor p/2 \rfloor + 1} = s_{p-\lfloor p/2 \rfloor - 1} + \dots + s_{p+\lfloor p/2 \rfloor + 1}$$

 $\Rightarrow s_{p-\lfloor p/2 \rfloor + 1} = \sum_{i=0}^{p-\lfloor p/2 \rfloor} s_i.$

Note that in all computations above, we omit the coefficients in order to simplify the algebraic expressions. The proof is now concluded.

Corollary 4. The set $\mathcal{B}_p := \{t^n, n \in I\}$ spans the Kauffman bracket skein module of the lens spaces L(p, 1).

We now show that the set \mathcal{B}_p is linearly independent and thus, we prove the main theorem of the paper, Theorem 3. An easy and straightforward way to prove that the set \mathcal{B}_p forms a basis for KBSM(L(p, 1)), is by relating this set to the basis presented in [23] via an invertible lower triangular matrix (see [13] for the case of the HOMFLYPT skein module of the Solid Torus). To make this paper more self-contained, we present here a proof based on the results presented before. Before doing so, however, a final note is necessary. The freeness of the module comes from the fact that the solution of the infinite system is unique, i.e. there are no contradictions. In other words, the equations obtained from elements in the set \mathcal{B}_p by performing negative braid band moves.

Proof. From the discussion and the results presented above, it follows that it suffices to show that $t^{\lfloor p/2 \rfloor}$ cannot be written as a sum of elements in $B_p \setminus \{t^{\lfloor p/2 \rfloor}\}$, or equivalently, that the unknown $s_{\lfloor p/2 \rfloor}$ cannot be written as a sum of unknowns of indices in $I \setminus \{\lfloor p/2 \rfloor\}$. We consider the most interesting case, that of p being even. The case that p is odd follows immediately from the proof of Theorem 12.

Note that since $t^{p/2}$ is in the basis of KBSM(ST), and since KBSM(L(p, 1)) = KBSM(ST)/ \langle band moves \rangle , it suffices to show that the braid band moves do not affect $t^{\lfloor p/2 \rfloor}$. Recall first that isotopy in L(p, 1) can be viewed as isotopy in ST together with one of the two different types of band moves (Theorem 6). So far,

we have been using the α -type band moves that are immediately translated on the level of braids by the braid band moves. In order to prove that $t^{p/2}$ for p even, cannot be expressed as a sum of looping generators of exponents in $I \setminus \lfloor p/2 \rfloor$, we use the β -type band moves. This simplifies the calculations involved a lot.

Indeed, consider $t^{p/2} \in B_{ST}$ and perform the β -type band move as illustrated in Fig. 15. Note that the result of the performance of a type β band move on a looping generator is not a mixed braid. But as shown in Fig. 15, isotopy in ST results in the looping generator $\widehat{t^{p-p/2}} = \widehat{t^{p/2}}$, that we braid and obtain $t^{p/2}$ again. Hence, $t^{p/2}$ cannot be written as sum of elements in $\mathcal{B}_p \setminus \{t^{p/2}\}$ and the proof is now concluded.

Remark 9. It is possible to prove that $t^{p/2}$ cannot be written as a sum of elements in $\mathcal{B}_p \setminus \{t^{p/2}\}$ using the braid band moves, i.e. the α -type band moves. The difficulty lies in keeping track of all coefficients involved, since following the same procedure as in the proof of Theorem 12 for $t^{-p/2} \xrightarrow{\text{bbm}} t^p t_1^{-p/2} \sigma_1^{\pm 1}$ (applying Lemma 6 and Proposition 3), results in an equation of the form $\sum_{i=0}^{p/2} a_i s_i = \sum_{i=0}^{p/2} b_i s_i$, where a_i, b_i coefficients for all *i*. We would then have to prove that $a_i = b_i$ for all *i*, which is very complicated and technical.

Remark 10. The case $L(0,1) \cong S^1 \times S^2$ is of special interest and in [10], the algebraic approach is applied in order to compute $\text{KBSM}(S^1 \times S^2)$. As shown in [24], $\text{KBSM}(S^1 \times S^2)$ contains torsion, and in [10] we show how torsion is detected by solving the corresponding infinite system of equations (3). This emphasizes on the importance of the algebraic approach based on unoriented braids that we present in this paper.

5. The Diagrammatic Approach for Computing KBSM(L(p,q))

In this section, we compute the Kauffman bracket skein module of the lens spaces L(p,q) using a diagrammatic method that is different to the diagrammatic method used in [23]. We first demonstrate this method for the q = 1 case and we then extend it for q > 1. It is worth mentioning that in [4], a new basis for the Kauffman bracket skein module of the complement of (2, 2p + 1)-torus knots is presented via this new method, which is based on unoriented braids.

5.1. The L(p, 1) case

In this section, we describe how the basis \mathcal{B}_p can be obtained using a diagrammatic approach based on unoriented braids. Instead of the α -band move, we will be using the β -type band move throughout this section (recall Fig. 4).

Let $t^n \in B_{ST}$. Apply a β -type band move as shown in the first two illustrations of Fig. 15. The red strand corresponds to the strand that appears after the band move is performed, which wraps around the surgery curve I, p times. The orange strands



Fig. 15. The β -type band move on an element in B_{ST} and its result in KBSM(L(p, 1)).

in the middle illustration indicate the places where isotopy in ST is performed in order to unwrap the moving strand from the surgery strand.

As shown in Fig. 15, using the β = band move and isotopy in ST, we may reduce t^n to t^{n-p} for all $n \in \mathbb{Z}$. In particular, for $n \in \mathbb{N}$, we have that

$$\begin{array}{rrrr} n=0 & : & t^0=t^p, & n=1:t^1=t^{p-1}, \\ n=2 & : & t^2=t^{p-2}, & n=3:t^3=t^{p-3}, \\ \vdots & & \vdots \\ n=\lfloor p/2 \rfloor -1:t^{\lfloor p/2 \rfloor -1}=t^{p-\lfloor p/2 \rfloor +1} \end{array}$$

and we conclude that the t^n 's for $n \in \{\lfloor p/2 \rfloor + 1, \ldots, p\}$ are not in the basis of KBSM(L(p, 1)).

Similarly, for $n \in \mathbb{Z} \setminus \mathbb{N}$, we have

$$n = -1: t^{-1} = t^{p+1}, \ n = -2: t^{-2} = t^{p+2},$$

$$n = -3: t^{-3} = t^{p+3}, \ n = -4: t^{-4} = s_{p+4},$$

$$\vdots \qquad \vdots$$

and we conclude that $t^{p+k} \sim t^{-k}$, for $k \in \mathbb{N}$. Recall now that $t^{-k} \cong \sum_{\text{skein}}^{k} t^i$ (Proposition 2). Thus, $t^{p+1} \sim t^{-1} = t$, etc. We conclude again that the t^n 's for n > p are not in the basis of KBSM(L(p, 1)). Equivalently, we have that the set \mathcal{B}_p spans KBSM(L(p, 1)). Linear independence follows then from the proof of Theorem 3.

5.2. The $L(p,q), q \notin \{0,1\}$ case

In this section, we adjust the proof of Theorem 3 for the case of L(p,q), $q \notin \{0,1\}$, and in particular we show that the diagrammatic approach using the unoriented braids that we presented in Sec. 5.1, also works for all $q \notin \{0,1\}$. Hence, we shall conclude that the set \mathcal{B}_p forms a basis for KBSM(L(p,q)).

We first note that the band moves are more complicated for $q \neq 1$, in the sense that when a moving strand of a mixed link approaches the surgery curve \hat{I} , q new



Fig. 16. The β -type band move for L(3, 2).



Fig. 17. The band move for L(p,q), q > 1.

strands will appear wrapping around \hat{I} , *p*-times. For an illustration, see Fig. 16, for p = 3 and q = 2 and where band move is shortened to bm.

On the unoriented braid level, a generic (p, q)-band move is illustrated in Fig. 17. The shaded area depicts the braided (p, q)-torus knots that appears after the performance of the band move, coming from the fact that this torus knot bounds a disk in L(p, q). Note also that the standard closure for mixed braids on the resulted "braid" is a β -band move on the mixed link level.

Our starting point is the basis $B_{\rm ST}$ of the Kauffman bracket skein module of the solid torus. The idea is the same as in the q = 1 case. More precisely, we consider elements $t^n \in B_{\rm ST}$ and perform the β -type band moves. Then, using isotopy in ST we may reduce t^n to t^{n-p} for all n. In Fig. 18, we demonstrate this process for L(3,2) and in particular we show how to reduce t^2 to t in L(3,2).

Thus, for $k \notin I$, we have that $t^k \cong t^n$, for some $n \in I$. The same arguments that we used in the proof of Theorem 3 for the q = 1 case, in order to prove that the set \mathcal{B}_p is linearly independent, also apply for the q > 1 case, and we conclude that the set \mathcal{B}_p forms a basis for KBSM(L(p,q)).

Remark 11. It is possible to use the algebraic method presented in Sec. 4 to derive the basis \mathcal{B}_p for KBSM(L(p,q)), q > 1. Only the calculations involved are more complicated due to the complexity of the (p,q)-braid band move: $\text{bbm}(t^n) =$



Fig. 18. $t^2 \sim t$ in L(3, 2).

 $[(\sigma_{q-1} \ldots \sigma_1) t]^p t_q^n \sigma_q^{\pm 1}$. If we consider $bbm(t^n)$ in KBSM(ST), we may express it as a sum of elements in B_{ST} . Using the Kauffman bracket skein relations together with Lemma 6 and Proposition 3, one may show that $bbm(t^n) = \sum_{i=0}^{p+n} a_i t^i$, for some coefficients a_i and for $n \in \mathbb{N}$. The result then follows similarly to the L(p, 1) case.

6. Conclusions

In this paper, we introduce the concept of unoriented braids, that play an important role for computing Kauffman bracket skein modules of c.c.o. 3-manifolds, and that seems promising in serving as a tool for studying c.c.o. 3-manifold invariants in general. Then, we introduce the generalized Temperley–Lieb algebra of type B, $TL_{1,n}$, and Jones' original idea [26], we present the most generic invariant, V, for knots and links in the solid torus ST, of the Kauffman bracket type. We then extend V to an invariant for knots and links in the lens spaces by imposing on V relations coming from the (braid) band moves. Our starting point is the Kauffman bracket

skein module of ST, KBSM(ST), since

$$\operatorname{KBSM}(L(p,q)) = \frac{\operatorname{KBSM}(\operatorname{ST})}{\langle \text{band moves} \rangle}.$$

We first present a new basis for KBSM(ST), $B_{\rm ST}$, using the concept of unoriented braids and we then solve the infinite system of equations $V_{\hat{a}} = V_{\widehat{\text{bbm}(a)}}$, where bbm(a) denotes the result of a (p, 1)-braid band move on an element a in the basis of KBSM(ST). This is equivalent to computing the Kauffman bracket skein module of L(p, 1) and it is the first time that a skein module is fully computed via this algebraic approach based on braids.

For the case of the HOMFLYPT skein module of the lens spaces L(p, 1), the reader is referred to [14, 15, 17, 3] for an algebraic approach based on braids and the generalized Hecke algebra of type B, and [20] for a diagrammatic approach via arrow diagrams (see also [19, 11]). We then present a new diagrammatic approach for computing KBSM(L(p, 1)) and we finally extend this approach for computing KBSM(L(p,q)), $q \neq 0$. Finally, it is worth mentioning that in [5, 6, 8, 7, 9], skein modules are also discussed for different families of "knotted objects" in 3-manifolds, such as tied links, pseudo and singular links and knotoids.

Acknowledgment

I would like to acknowledge several discussions with Prof. Sofia Lambropoulou, who provided insight and expertise that greatly assisted this research.

References

- I. Diamantis, An alternative basis for the Kauffman bracket skein module of the solid torus via braids, in *Knots, Low-Dimensional Topology and Applications (KNOTS16* 2016), eds. C. Adams et al., Springer Proceedings in Mathematics and Statistics, Vol. 284 (Springer, Cham, 2019), pp. 329–345, doi:10.1007/978-3-030-16031-9_16.
- [2] I. Diamantis, The Kauffman bracket skein module of the handlebody of genus 2 via braids, J. Knot Theory Ramifications 28(13) (2019) 1940020, doi:10.1142/S0218216519400200.
- [3] I. Diamantis, HOMFLYPT skein sub-modules of the lens spaces L(p, 1) via braids, *Topology Appl.* **301** (2021) 107500, doi:10.1016/j.topol.2020.107500, arXiv:2005.00737.
- [4] I. Diamantis, The Kauffman bracket skein module of the complement of (2, 2p + 1)-torus knots via braids, preprint (2021), arXiv:2106.04965 [math.GT].
- [5] I. Diamantis, Tied pseudo links & pseudo knotoids, Mediterr. J. Math. 18 (2021) 201, doi:10.1007/s00009-021-01842-1.
- [6] I. Diamantis, Tied links in various topological settings, J. Knot Theory Ramifications 30(07) (2021) 2150046, doi:10.1142/S0218216521500462.
- [7] I. Diamantis, Knotoids, pseudo knotoids, Braidoids and pseudo braidoids on the Torus, *Commun. Korean Math. Soc.* **37**(4) (2022) 1221–1248, doi:10.4134/ CKMS.c210169, arXiv:2103.16433 [math.GT].
- [8] I. Diamantis, Pseudo links and singular links in the Solid Torus, Communications in Mathematics 31(1) (2023), doi:10.46298/cm.10438, arXiv:2101.03538v1 [math.GT].

- [9] I. Diamantis, Pseudo links and singular links in the Solid Torus, Bull. Hellenic Math. Soc. 65 (2021) 17–34.
- [10] I. Diamantis, The Kauffman bracket skein module of $S^1 \times S^2$ via unoriented braids, manuscript in preparation.
- [11] I. Diamantis, B. Gabrovšek, S. Lambropoulou and M. Mroczkowski, Knot Theory of Lens Spaces, De Gruyter Expositions in Mathematics (De Gruyter, 2022).
- [12] I. Diamantis and S. Lambropoulou, Braid equivalences in 3-manifolds with rational surgery description, *Topology Appl.* **194** (2015) 269–295, doi:10.1016/ j.topol.2015.08.009.
- [13] I. Diamantis and S. Lambropoulou, A new basis for the HOMFLYPT skein module of the solid torus, J. Pure Appl. Algebra 220(2) (2016) 577–605, doi:10.1016/j.jpaa.2015.06.014.
- [14] I. Diamantis and S. Lambropoulou, The braid approach to the HOMFLYPT skein module of the lens spaces L(p, 1), in Algebraic Modeling of Topological and Computational Structures and Application, Springer Proceedings in Mathematics and Statistics (PROMS) (Springer, Cham, 2017), pp. 143–176, doi:10.1007/978-3-319-68103-0_7.
- [15] I. Diamantis and S. Lambropoulou, An important step for the computation of the HOMFLYPT skein module of the lens spaces L(p, 1) via braids, J. Knot Theory Ramifications **28**(11) (2019) 1940007, doi:10.1142/S0218216519400078.
- [16] I. Diamantis and S. Lambropoulou, The theory of unoriented braids, in preparation.
- [17] I. Diamantis, S. Lambropoulou and J. H. Przytycki, Topological steps on the HOM-FLYPT skein module of the lens spaces L(p, 1) via braids, J. Knot Theory Ramifications **25**(14) (2016) 1650084.
- [18] M. Flores and D. Goundaroulis, Framization of a Temperley–Lieb algebra of type B, J. Pure Appl. Algebra 224(6) (2020) 106273.
- B. Gabrovsek, Tabulation of prime knots in lens spaces, Mediterr. J. Math. 14 (2017) 88.
- [20] B. Gabrovšek and M. Mroczkowski, The HOMFLYPT skein module of the lens spaces L(p, 1), Topology and its Appl. 175 (2014) 72–80.
- [21] B. Gabrovšek and M. Mroczkowski, Link diagrams and applications to skein modules, in Algebraic Modeling of Topological and Computational Structures and Applications, Springer Proceedings in Mathematics & Statistics (Springer, 2017), pp. 117–141.
- [22] J. Hoste and M. Kidwell, Dichromatic link invariants, Trans. Amer. Math. Soc. 321(1) (1990) 197–229.
- [23] J. Hoste and J. Przytycki, The $(2, \infty)$ -skein module of lens spaces: A generalization of the Jones polynomial, J. Knot Theory Ramifications **02**(03) (1993) 321–333.
- [24] J. Hoste and J. Przytycki, The Kauffman bracket skein module of $S^1 \times S^2$, Math. Z. **220**(1) (1995) 63–73.
- [25] V. F. R. Jones, A polynomial invariant for links via von-Neumann algebras, Bull. Amer. Math. Soc. 12 (1985) 103–111.
- [26] V. F. R. Jones, Hecke algebra representations of braid groups and link polynomials, Ann. Math. 126 (1987) 335–388.
- [27] S. Lambropoulou, Solid torus links and Hecke algebras of B-type, in *Quantum Topology*, ed. D. N. Yetter (World Scientific Press, 1994), pp. 225–245.
- [28] S. Lambropoulou, Knot theory related to generalized and cyclotomic Hecke algebras of type B, J. Knot Theory Ramifications 8(5) (1999) 621–658.
- [29] S. Lambropoulou and C. P. Rourke, Markov's theorem in 3-manifolds, *Topology Appl.* 78 (1997) 95–122.

- [30] S. Lambropoulou and C. P. Rourke, Algebraic Markov equivalence for links in 3-manifolds, *Compos. Math.* 142 (2006) 1039–1062.
- [31] J. Przytycki, Skein modules of 3-manifolds, Bull. Polish Acad. Sci. Math. 39(1-2) (1991) 91-100.
- [32] V. G. Turaev, The Conway and Kauffman modules of the solid torus, Zap. Nauchn. Sem. LOMI 167 (1988) 79–89 [English translation: J. Soviet Math. (1990) 2799– 2805].