

The anabolic properties of plant-derived proteins

Citation for published version (APA):

Pinckaers, P. J. M. (2024). *The anabolic properties of plant-derived proteins*. [Doctoral Thesis, Maastricht University]. Maastricht University. <https://doi.org/10.26481/dis.20240410pp>

Document status and date:

Published: 01/01/2024

DOI:

[10.26481/dis.20240410pp](https://doi.org/10.26481/dis.20240410pp)

Document Version:

Publisher's PDF, also known as Version of record

Please check the document version of this publication:

- A submitted manuscript is the version of the article upon submission and before peer-review. There can be important differences between the submitted version and the official published version of record. People interested in the research are advised to contact the author for the final version of the publication, or visit the DOI to the publisher's website.
- The final author version and the galley proof are versions of the publication after peer review.
- The final published version features the final layout of the paper including the volume, issue and page numbers.

[Link to publication](#)

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal.

If the publication is distributed under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license above, please follow below link for the End User Agreement:

www.umlib.nl/taverne-license

Take down policy

If you believe that this document breaches copyright please contact us at:

repository@maastrichtuniversity.nl

providing details and we will investigate your claim.

IMPACT

The main aim of this dissertation was to investigate the anabolic properties of plant-derived proteins, and to evaluate how they compare with animal-derived proteins in their capacity to stimulate muscle protein synthesis. This thesis extends on previous work reporting on the inferior anabolic properties of soy protein isolate ingestion when compared with animal-derived protein in humans [1-3]. This dissertation evaluated the plasma amino acid profiles and muscle protein synthesis rates following ingestion of wheat, corn, pea, and potato derived protein in comparison to animal-derived milk protein in healthy, young men. The amino acid profiles following ingestion of 30 g of the selected plant-derived proteins showed lower plasma essential amino acid availability, when compared to the ingestion of the same amount of milk-derived protein, especially for leucine, lysine, and methionine. Despite a lower essential amino acid provision, muscle protein synthesis rates were strongly increased following ingestion of the selected plant-derived protein sources. Importantly, this muscle protein synthetic response did not differ when compared to the ingestion of the same amount of milk protein. Similar results were observed when potato protein isolate was ingested during recovery from exercise. Furthermore, we presented data on the anabolic properties of protein blends, in which different plant-derived proteins are combined, with or without animal-derived protein. We showed that despite lower plasma essential amino acid availability, the ingestion of the selected protein blends increased muscle protein synthesis rates, which was not different from the increase observed following ingestion of milk protein. Lastly, we reported that ingestion of a lysine enriched plant-based meat substitute, consisting of wheat- and chickpea-derived protein, increases muscle protein synthesis rates to an extent not different from chicken breast filet, when both provide 40 g protein. Collectively, we can conclude, that ingestion of an ample amount of plant-derived protein (i.e. ≥ 30 g) stimulates muscle protein synthesis to an extent not different from ingesting the same amount of animal-derived protein in healthy, young men.

Stimulating muscle protein synthesis is fundamental for maintaining muscle health and support the exercise induced muscle reconditioning response [4]. When protein is ingested in close proximity to physical activity or exercise, the increased muscle protein synthetic response is greater and prolonged, when compared with the stimulation observed following exercise or protein ingestion only [5-8]. However, it is important to note that not all proteins are the same and do not have the same potential to stimulate muscle protein synthesis rates [9, 10]. Therefore, this thesis provides valuable information outlining the anabolic properties of various plant-derived proteins, their protein blends, and a plant-based meat substitute. The impact of this dissertation, as well as the implications for translation into practice will be discussed below.

Scientific relevance

This thesis broadens our understanding of the anabolic properties of plant-derived proteins. We have made substantial contributions to the scientific field, to better understand the important factors impacting the anabolic response following protein ingestion. Early studies on protein quality identified three main factors that would determine protein quality, and the subsequent anabolic response: 1) essential amino acid composition, 2) leucine content,

and 3) protein digestion and amino acid absorption of the specific protein or protein source [10]. Given that on all these factors plant-derived proteins are subpar when compared to animal-derived proteins, plant-derived proteins have traditionally been considered to exhibit a lower capacity to increase muscle protein synthesis rates [11]. This assumption was primarily based on earlier studies evaluating the anabolic properties following ingestion of soy protein isolates [1-3, 12]. Several of these studies showed that soy protein ingestion stimulates muscle protein synthesis to a lower extent when compared to the ingestion of dairy-derived proteins, both at rest and during post-exercise recovery [1-3]. There are, however, many different plant protein sources with each holding their own unique properties [9, 10]. This thesis expands our knowledge by investigating the anabolic properties of wheat, corn, pea, and potato protein, protein blends, and a plant-based meat substitute *in vivo* in humans. The current work clearly shows that low essential amino acid contents of plant-derived proteins do not necessarily compromise the acute anabolic response when a sufficient amount of protein (i.e. ≥ 30 g) is ingested in healthy, young men. Therefore, neither protein characteristics, such as essential amino acid composition, nor the availability of essential amino acids in the circulation following ingestion, predict the muscle protein synthetic response following protein ingestion. Hence, statements on the anabolic properties of a specific protein can only be sustained when muscle protein synthesis rates have been assessed directly following the ingestion of that specific protein *in vivo* in humans.

With the work presented in this thesis we have substantially contributed to the current knowledge on the anabolic properties of plant-derived proteins and have provided many new directions for future research. It would be of interest to evaluate the muscle protein synthetic response of the selected proteins in older individuals. Given that older individuals suffer from anabolic resistance [13], it is of substantial interest to determine how plant-derived proteins compare with animal-derived proteins in stimulating muscle protein synthesis rates in this population. Secondly, the amount of protein ingested is of key importance to consider when evaluating plant-derived protein sources, particularly in older adults [14, 15]. A key question that should be addressed is whether older individuals need to ingest more plant-derived protein to elicit a similar anabolic response when compared with the ingestion of animal-derived protein. Furthermore, as discussed in chapter 9, there is a need to transition towards assessing the anabolic response following ingestion of whole foods and more complex meals, to take into account all interactions between foods and nutrients on protein digestion, amino acid absorption, and anabolic signaling originating from all other food components besides protein.

Target groups and societal relevance

There is a global transition towards the consumption of more plant-derived proteins, which will become more important to attain future global protein needs [16-18]. Plant-derived proteins are considered to be a more sustainable protein source as production requires less water, land, and energy resources when compared to animal-derived proteins [19]. Given the essential role of dietary protein in the maintenance of muscle health, it is key to understand the anabolic properties of plant-derived proteins.

The work in this thesis shows that wheat, corn, pea, and potato-derived protein isolates may potentially be used as effective alternative protein sources when transitioning towards more plant-based protein consumption in young, healthy individuals. In addition, we show that plant-derived protein blends, and plant-based meat substitutes, may also be considered as alternative protein sources when transitioning towards the consumption of more plant-derived proteins. This provides direction and opportunities for industry for the development of plant-derived protein rich products and meat substitutes, i.e. the type and amount of protein to be used to ensure an anabolic response similar to the ingestion of an equivalent amount of animal-derived products. We identified potato protein concentrate as an effective protein source to stimulate muscle protein synthesis rates at rest and during recovery from exercise. This is of special interest, since potato protein concentrate is produced from the waste product of potato starch extraction. Hence, this not only shows that plant-derived proteins can support the transition towards a more plant-based diet, but also supports the current drive towards a more circular economy. The studies presented in this thesis have been performed within a public-private partnership, in which a collaboration is found between the Dutch government, university and industry to perform pre-competitive research. Beside the financial support to perform independent scientific research studies, such a public-private partnership facilitates the translation of scientific discoveries into practical applications and products. Private (industry) partners generally have expertise in commercialization and manufacturing, and distribution which is key to bridge the gap between research findings and real-world implementation. This dissertation capitalizes on this collaboration by direct utilization of the results and products used by the private partners in the development of their consumer products.

In the athletic community there is an increasing interest towards specific plant-derived sports nutrition products to support athletes adhering to a vegan and vegetarian diet [20]. More than 80% of the sports nutrition market sales originates from protein products, including powders, bars, and drinks [21]. Given the combined interest towards plant-derived proteins and the continuously growing sports nutrition market, the application of plant-derived proteins in sports nutrition will substantially increase in the years to come. This dissertation, provides key knowledge in this transition by showing that potato-derived protein can effectively support post-exercise muscle protein synthesis rates, and thereby support muscle recovery and remodeling following exercise. Like in athletes, skeletal muscle recovery and/or reconditioning is of equal importance in many clinical situations. The loss of skeletal muscle mass and function is a well-known phenomenon during hospitalization [22, 23]. Apart from the negative effects of physical inactivity, the loss of muscle mass observed during hospitalization may be attributed to an insufficient protein intake as a direct consequence of the lower energy intake [24, 25]. Within the field of clinical nutrition there is equal interest in the use of more sustainable protein sources [26]. In addition, there will be a need to accommodate future patient choices to consume a more plant-based diet. Therefore, it is key to evaluate the anabolic responses of plant-derived protein administration in frail and older individuals, to support optimal recovery with the use of both animal as well as plant-derived protein.

Translation into practice

In today's society, there is a lot of information available and accessible to every individual, and people share their thoughts and opinions with the world by just a few clicks on social media. On the flipside, there is a large spread of misinformation, hype, and very narrow perspectives without scientific rigor. Especially when considering food supplements and diets excluding certain foods or food groups, it is important to elucidate the potential health consequences in an evidence-based manner. Throughout the completion of this dissertation we have not only provided scientific reports and presentations at scientific conferences, but also actively incorporated results in lectures in nutrition and exercise physiology educational courses. Outside the academic world we have, and still see an increasing interest in the transition towards the consumption of more plant-derived proteins, in which we are being approached to translate our knowledge to be applied by e.g. sports nutrition companies and the military.

The transition towards incorporation of more plant-derived protein in our diet, seems inevitable from both a consumers' choice as well as a sustainability perspective, to meet future global protein demands. Therefore, it is important to not only address the potential shortcomings in the anabolic response of plant-derived proteins, but to also give directions towards potential solutions. The scientific field should not be blinded by the initial studies showing a lower anabolic response following soy vs dairy protein ingestion [2, 3], but should rather support this transition by providing directions towards potential solutions. In chapter 2 we have given potential solutions to the lower quality of plant-derived proteins, i.e. 1) providing protein isolates, 2) fortifying the protein source with its limiting amino acids, 3) providing protein blends, and 4) consuming more of a lower quality protein. Future research should focus on translating these findings even further, by investigating the clinical relevance of the potential differential anabolic response between plant- and animal-derived proteins. Lastly, the knowledge obtained from plant-derived protein isolates and meat substitutes, requires further translation towards whole meals in the everyday kitchen.

REFERENCES

1. Tang JE, Moore DR, Kujbida GW, Tarnopolsky MA, Phillips SM. Ingestion of whey hydrolysate, casein, or soy protein isolate: Effects on mixed muscle protein synthesis at rest and following resistance exercise in young men. *Journal of applied physiology (Bethesda, Md : 1985)*. 2009;107(3):987-92; <https://doi.org/10.1152/jappphysiol.00076.2009>.
2. Wilkinson SB, Tarnopolsky MA, Macdonald MJ, Macdonald JR, Armstrong D, Phillips SM. Consumption of fluid skim milk promotes greater muscle protein accretion after resistance exercise than does consumption of an isonitrogenous and isoenergetic soy-protein beverage. *Am J Clin Nutr*. 2007;85(4):1031-40; <https://doi.org/10.1093/ajcn/85.4.1031>.
3. Yang Y, Churchward-Venne TA, Burd NA, Breen L, Tarnopolsky MA, Phillips SM. Myofibrillar protein synthesis following ingestion of soy protein isolate at rest and after resistance exercise in elderly men. *Nutr Metab (Lond)*. 2012;9(1):57; <https://doi.org/10.1186/1743-7075-9-57>.
4. Wilkinson SB, Phillips SM, Atherton PJ, Patel R, Yarasheski KE, Tarnopolsky MA, et al. Differential effects of resistance and endurance exercise in the fed state on signalling molecule phosphorylation and protein synthesis in human muscle. *J Physiol*. 2008;586(15):3701-17; <https://doi.org/10.1113/jphysiol.2008.153916>.
5. Tipton KD, Ferrando AA, Phillips SM, Doyle D, Jr., Wolfe RR. Postexercise net protein synthesis in human muscle from orally administered amino acids. *Am J Physiol*. 1999;276(4 Pt 1):E628-34; <https://doi.org/10.1152/ajpendo.1999.276.4.E628>.
6. Biolo G, Tipton KD, Klein S, Wolfe RR. An abundant supply of amino acids enhances the metabolic effect of exercise on muscle protein. *American Journal of Physiology-Endocrinology And Metabolism*. 1997;273(1):E122-E9; <https://doi.org/10.1152/ajpendo.1997.273.1.E122>.
7. Howarth KR, Moreau NA, Phillips SM, Gibala MJ. Coingestion of protein with carbohydrate during recovery from endurance exercise stimulates skeletal muscle protein synthesis in humans. *J Appl Physiol*. 2009;106(4):1394-402; <https://doi.org/10.1152/jappphysiol.90333.2008>.
8. Breen L, Philp A, Witard OC, Jackman SR, Selby A, Smith K, et al. The influence of carbohydrate-protein co-ingestion following endurance exercise on myofibrillar and mitochondrial protein synthesis. *The Journal of Physiology*. 2011;589(16):4011-25; <https://doi.org/10.1113/jphysiol.2011.211888>.
9. Gorissen SH, Crombag JJ, Senden JM, Waterval WH, Bierau J, Verdijk LB, et al. Protein content and amino acid composition of commercially available plant-based protein isolates. *Amino Acids*. 2018;50(12):1685-95; <https://doi.org/10.1007/s00726-018-2640-5>.
10. van Vliet S, Burd NA, van Loon LJ. The skeletal muscle anabolic response to plant- versus animal-based protein consumption. *J Nutr*. 2015;145(9):1981-91; <https://doi.org/10.3945/jn.114.204305>.
11. Lim MT, Pan BJ, Toh DWK, Sutanto CN, Kim JE. Animal protein versus plant protein in supporting lean mass and muscle strength: A systematic review and meta-analysis of randomized controlled trials. *Nutrients*. 2021;13(2):661; <https://doi.org/10.3390/nu13020661>.
12. Churchward-Venne TA, Pinckaers PJM, Smeets JSJ, Peeters WM, Zorenc AH, Schierbeek H, et al. Myofibrillar and mitochondrial protein synthesis rates do not differ in young men following the ingestion of carbohydrate with whey, soy, or leucine-enriched soy protein after concurrent resistance- and endurance-type exercise. *J Nutr*. 2019;149(2):210-20; <https://doi.org/10.1093/jn/nxy251>.
13. Wall BT, Gorissen SH, Pennings B, Koopman R, Groen BBL, Verdijk LB, et al. Aging is accompanied by a blunted muscle protein synthetic response to protein ingestion. *PLoS One*. 2015;10(11):e0140903; <https://doi.org/10.1371/journal.pone.0140903>.
14. Witard OC, Jackman SR, Breen L, Smith K, Selby A, Tipton KD. Myofibrillar muscle protein synthesis rates subsequent to a meal in response to increasing doses of whey protein at rest and after resistance exercise. *Am J Clin Nutr*. 2014;99(1):86-95; <https://doi.org/10.3945/ajcn.112.055517>.

15. Moore DR, Churchward-Venne TA, Witard O, Breen L, Burd NA, Tipton KD, et al. Protein ingestion to stimulate myofibrillar protein synthesis requires greater relative protein intakes in healthy older versus younger men. *Journals of Gerontology Series A: Biomedical Sciences and Medical Sciences*. 2015;70(1):57-62; <https://doi.org/10.1093/gerona/glu103>.
16. Brink E, Van Rossum C, Postma-Smeets A, Stafleu A, Wolvers D, Van Dooren C, et al. Development of healthy and sustainable food-based dietary guidelines for the netherlands. *Public Health Nutr*. 2019;22(13):2419-35; <https://doi.org/10.1017/s1368980019001435>.
17. Aiking H. Protein production: Planet, profit, plus people? *Am J Clin Nutr*. 2014;100 Suppl 1:483S-9S; <https://doi.org/10.3945/ajcn.113.071209>.
18. Aiking H, de Boer J. The next protein transition. *Trends in Food Science & Technology*. 2020;105:515-22; <https://doi.org/10.1016/j.tifs.2018.07.008>.
19. Pimentel D, Pimentel M. Sustainability of meat-based and plant-based diets and the environment. *Am J Clin Nutr*. 2003;78(3 Suppl):660S-3S; <https://doi.org/10.1093/ajcn/78.3.660S>.
20. López-Martínez MI, Miguel M, Garcés-Rimón M. Protein and sport: Alternative sources and strategies for bioactive and sustainable sports nutrition. *Front Nutr*. 2022;9:926043; <https://doi.org/10.3389/fnut.2022.926043>.
21. Arenas-Jal M, Suñé-Negre JM, Pérez-Lozano P, García-Montoya E. Trends in the food and sports nutrition industry: A review. *Crit Rev Food Sci Nutr*. 2020;60(14):2405-21; <https://doi.org/10.1080/10408398.2019.1643287>.
22. Kouw IWK, Groen BBL, Smeets JSJ, Kramer IF, Van Kranenburg JMX, Nilwik R, et al. One week of hospitalization following elective hip surgery induces substantial muscle atrophy in older patients. *J Am Med Dir Assoc*. 2019;20(1):35-42; <https://doi.org/10.1016/j.jamda.2018.06.018>.
23. Van Ancum JM, Scheerman K, Jonkman NH, Smeenk HE, Kruizinga RC, Meskers CGM, et al. Change in muscle strength and muscle mass in older hospitalized patients: A systematic review and meta-analysis. *Exp Gerontol*. 2017;92:34-41; <https://doi.org/10.1016/j.exger.2017.03.006>.
24. Weijzen MEG, Kouw IWK, Geerlings P, Verdijk LB, Van Loon LJC. During hospitalization, older patients at risk for malnutrition consume <0.65 grams of protein per kilogram body weight per day. *Nutr Clin Pract*. 2020;35(4):655-63; <https://doi.org/10.1002/ncp.10542>.
25. Weijzen MEG, Kouw IWK, Verschuren AAJ, Muyters R, Geurts JA, Emans PJ, et al. Protein intake falls below 0.6 g·kg⁻¹·d⁻¹ in healthy, older patients admitted for elective hip or knee arthroplasty. *J Nutr Health Aging*. 2019;23(3):299-305; <https://doi.org/10.1007/s12603-019-1157-2>.
26. Saldivar B, Al-Turk B, Brown M, Aggarwal M. Successful incorporation of a plant-based menu into a large academic hospital. *Am J Lifestyle Med*. 2022;16(3):311-7; <https://doi.org/10.1177/15598276211048788>.