

# Intracochlear electrical stimulation to suppress tinnitus

Citation for published version (APA):

Arts, R. A. G. J. (2016). *Intracochlear electrical stimulation to suppress tinnitus*. [Doctoral Thesis, Maastricht University]. Maastricht University. <https://doi.org/10.26481/dis.20160115ra>

## Document status and date:

Published: 01/01/2016

## DOI:

[10.26481/dis.20160115ra](https://doi.org/10.26481/dis.20160115ra)

## Document Version:

Publisher's PDF, also known as Version of record

## Please check the document version of this publication:

- A submitted manuscript is the version of the article upon submission and before peer-review. There can be important differences between the submitted version and the official published version of record. People interested in the research are advised to contact the author for the final version of the publication, or visit the DOI to the publisher's website.
- The final author version and the galley proof are versions of the publication after peer review.
- The final published version features the final layout of the paper including the volume, issue and page numbers.

[Link to publication](#)

## General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal.

If the publication is distributed under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license above, please follow below link for the End User Agreement:

[www.umlib.nl/taverne-license](http://www.umlib.nl/taverne-license)

## Take down policy

If you believe that this document breaches copyright please contact us at:

[repository@maastrichtuniversity.nl](mailto:repository@maastrichtuniversity.nl)

providing details and we will investigate your claim.

## Valorization addendum

Although the results obtained in these feasibility studies are promising, one could doubt whether patients would accept such an invasive and expensive procedure to treat their tinnitus.

### Treatment acceptance

To reduce tinnitus completely, 38% of the 197 tinnitus sufferers who attended a meeting of the Australian Tinnitus Society would accept (i.e., they indicated an acceptance of 91-100%) a CI as treatment option while 25% would accept a CI in case of reduced tinnitus loudness and annoyance by half. This acceptance rate is comparable to noninvasive wearable devices (42% and 30%, respectively) [Tyler, 2012]. In contrast, based on a Dutch online survey on 415 tinnitus sufferers and member of the Dutch society for the hearing impaired, 29% of the population express a large interest in CI (i.e., they scored their acceptance 8-10 on a visual analogue scale from 0 to 10) in case there is a 100% chance of full tinnitus suppression while 22% is very interested in case there is a 50% chance of full tinnitus suppression. A decrease in treatment acceptability was observed compared to the noninvasive conventional hearing aids, with an acceptance rate of 53% and 51% respectively (unpublished data). The difference in CI-acceptance in relation to noninvasive wearable devices between both studies can probably be explained by selection bias. Nevertheless, both studies have shown that the acceptance of cochlear implantation as a treatment option for tinnitus is considerable which is consistent with the results obtained in an American internet survey with 439 responders. Here, almost three-fourths of the patients would be willing to have a device implanted in their body for a therapy that could eliminate or reduce their tinnitus perception by half [Engineer et al., 2013].

### Economic considerations

As mentioned in the introduction of this thesis, tinnitus affects about 50 million people in the United States and an estimated 70 million in the European Union [Cederroth et al., 2013] which is about 10-15% of the general population [Langguth et al., 2013]. The estimated tinnitus prevalence in the Netherlands is 2 million of whom about 60.000 severely suffer from their tinnitus [Cima et al. 2009]. Furthermore, Maes and colleagues [2013] concluded that the economic burden of tinnitus to society is substantial with an on average annual tinnitus related societal cost per patient of €5,315 in the Netherlands. Most of these costs were associated with production losses; on average €3,702. Overall, the annual costs of tinnitus in the Netherlands examined from a societal perspective is about €6.8 billion (95% confidence interval: €3.9 billion - €10.8 billion). The annual health care costs were €1.9 billion (95% confidence interval: €1.4 billion - €2.5 billion) which amounts to 2.3% of the total Dutch health care expenditure [Maes et al., 2013].

Based on these results it is striking that within the National Institute on Deafness and other Communication Disorders, tinnitus funding accounts for only 2% (about \$5 million per year from 2009 to 2011) of total hearing research funds [Cederroth et al., 2013]. Private donors like for example the founder of the Tinnitus Research Initiative; Matteo de Nora, the fundraising efforts of for example the American Tinnitus Association and investments from the industry are therefore essential for the knowledge about and treatment options for tinnitus.

Moreover, Tyler [2012] reported about the willingness to pay for tinnitus treatments and asked 197 tinnitus sufferers to indicate how much money they would be willing to pay for effective treatments. Most of these patients indicated to pay at least \$5000 and 20.3% were willing to pay as much as \$25,000 for complete tinnitus reduction. Furthermore, Engineer and colleagues [2013] reported about an online survey with 439 responders of which 94% reported that they had health insurance. Here, almost 40% had

already spent between \$500 and \$10,000 on tinnitus therapies. Therefore, from an economical point of view, the tinnitus population is an interesting market for the industry with in particular the industry of medical devices as novel drugs take a long time and hundreds of millions of dollars to develop. For purposes of illustration, Auris Medical recently invested \$51 million for a phase 3 trial in tinnitus for the pharmacotherapy using AM-101 [Cederroth et al., 2013]. In contrast, the average implantable medical device takes about \$10 million and five years to develop [Engineer et al., 2013].

This research project is funded by one of our CI industry partners. They are interested in the long-term effects of electrical stimulation independent of environmental sounds on tinnitus as they see new opportunities to expand their market. Although the highly sophisticated CI shows promising results concerning the tinnitus suppression, the considerable costs (about €25,000 for the device [Broersen, 2010]) have hampered the clinical application. With the promising long-term effects of electrical stimulation independent of environmental sounds on tinnitus, a "Tinnitus Implant" (TI) should be viable. This relatively simple neurostimulator could be produced considerably cheaper because it does not require a strategy to encode environmental sounds. This thesis includes feasibility studies (Chapter 3-7) which open new possibilities for the development of this treatment option.

### Possible effects on the standard clinical CI

The results obtained on tinnitus suppression may indirectly lead to future research in order to expand the inclusion criteria for CI, especially regarding residual hearing. To date, there is a considerable risk of damaged residual hearing due to the implantation itself, which is the main reason why only subjects who were deaf in the ear to be implanted were included. Changes in both the device and the surgical procedure could reduce this risk. If this risk can be neglected, this treatment option could possibly be suitable for the tinnitus population with significant residual hearing. This thesis can contribute to the interest to develop an atraumatic procedure for electrode insertion. Although Chapter 2 of this thesis seems to suggest that deterioration of residual hearing as a result of the implantation itself does not cause postoperative chronic tinnitus, hearing loss as a side effect of the electrode insertion for tinnitus treatment is undesirable.

For safety as well as methodological reasons the long-term effects of intracochlear electrical stimulation independent of environmental sounds were investigated using the standard clinical CI which was reprogrammed to function as a tinnitus specific neurostimulator. This gave us the opportunity to additionally investigate the audiological effects of the standard CI in the SSD population (Chapter 7). The positive effects of CI obtained can contribute to the broadening of the indication for the standard clinical CI. For example, in Belgium and Germany the indication for CI is expanded already to SSD associated with tinnitus (the CI system is CE-Market for SSD in both children and adults), because in these countries the value of CI in SSD has been proven [Van de Heyning et al., 2008; Kleinjung et al., 2009; Vermeire et al., 2009; Arndt et al., 2010; Buechner et al., 2010; Arndt et al., 2011; Jacob et al., 2011; Stelzig et al., 2011; Hassepas et al., 2012; Mertens et al., 2013]. It is plausible that the Dutch policy makers (College voor Zorgverzekering) will follow after research in the Netherlands has reproduced these findings.

### Knowledge valorization

The findings and ideas have been communicated to and shared with fellow researchers around the world at (inter)national congresses and in scientific journals. Furthermore, a huge appreciation for our work is observed in the tinnitus population. The Maastricht UMC+ organizes for example every six months a tinnitus symposium in collaboration with the Dutch society for the hearing impaired (Nederlandse Vereniging Voor Slechthorenden). These symposia were always fully booked. Here, background

information was given about tinnitus and hearing loss and recent findings from research projects were shared. Frequent interest of national media resulted in a generally better understanding of tinnitus. As tinnitus remains refractory to current medical treatment, sufferers are regularly disappointed after consulting a physician and do often not feel taken seriously. This clinical research contributes to the patient's awareness of global developments in this field of research.

## References

- Arndt S, Aschendorff A, Laszig R, Beck R, Schild C, Kroeger S, Ihorst G, Wesarg T: Comparison of pseudobinaural hearing to real binaural hearing rehabilitation after cochlear implantation in patients with unilateral deafness and tinnitus. *Otol Neurotol* 2010;32:39-47.
- Arndt S, Laszig R, Aschendorff A, Beck R, Schild C, Hassepaß F, Ihorst G, Kroeger S, Kirchem P, Wesarg T: Einseitige Taubheit und Cochleaimplantat-Versorgung. *HNO* 2011;59:437-446.
- Bartels H, Staal MJ, Holm AF, Mooij JJ, Alberts FW: Long-term evaluation of treatment of chronic, therapeutically refractory tinnitus by neurostimulation. *Stereotact Funct Neurosurg* 2007;85(4):150-7.
- Broersen SFM: Cochleair implantaat opent de wereld. *Medisch Contact* 2010;10:528-531.
- Buechner A, Brendel M, Lesinski-Schiedat A, Wenzel G, Frohne-Buechner C, Jaeger B, Lenarz T: Cochlear implantation in unilateral deaf subjects associated with ipsilateral tinnitus. *Otol Neurotol* 2010;31:1381-1385.
- Cederroth CR, Canlon B, Langguth: Hearing loss and tinnitus – are funders and industry listening? *Nature Biotechnology* 2013;31(11):972-4.
- Cima RF, Maes IH, Joore MA, Scheyen DJ, El Refaie A, Baguley DM, Anteunis LJ, van Breukelen GJ, Vlaeyen JW: Specialised treatment based on cognitive behaviour therapy versus usual care for tinnitus: a randomized controlled trial. *Lancet* 2012;379(9830):1951-9.
- De Martino F, Moerel M, van de Moortele PF, Ugurbil K, Goebel R, Yacoub E, Formisano E: Spatial organization of frequency preference and selectivity in the human inferior colliculus. *Nat Commun* 2013;4:1386.
- Di Nardo W, Cianfrone F, Scorpecci A, Cantone I, Giannantonio S, Paludetti G: Transtympanic electrical stimulation for immediate and long-term tinnitus suppression. *Int Tinnitus J* 2009;15(1):100-6.
- Dobie RA: A review of randomized clinical trials in tinnitus. *Laryngoscope* 1999;109:1202-1211.
- Engelhardt J, Dauman R, Arné P, Allard M, Dauman N, Branchard O, Perez P, Germain C, Caire F, Bonnard D, Cuny E: Effect of chronic cortical stimulation on chronic severe tinnitus: a prospective randomized double-blind cross-over trial and long-term follow up. *Brain Stimul* 2014;7(5):694-700.
- Engineer ND, Rosellini WM, Tyler RS: Willingness to accept and pay for implantable tinnitus treatments: a survey. *Neuromodulation* 2013;16(2):154-62.
- Gazibegovic D, Dees G, van Hoof M, Stokroos R: The assessment of the scalar position of the HiFocus Mis Scala electrode based on MRI and Cone Beam CT fusion. 8<sup>th</sup> International symposium on objective measures in auditory implants 2014; <http://www.eposteronline.com/omai2014/node/202>
- Gilles A, Van Hal G, De Ridder D, Wouters K, Van de Heyning P: Epidemiology of noise-induced tinnitus and the attitudes and beliefs towards noise and hearing protection in adolescence. *PLoS ONE* DOI: 10.1371/journal.pone.0070297.

- Greenwood DD: A cochlear frequency-position function for several species – 29 years later. *J Acoust Soc Am* 1990;87(6):2592-2605.
- Grimes DA, Schulz KF. Bias and causal associations in observational research. *Lancet* 2002;359:248-52.
- Hassepass F, Aschendorff A, Wesarg T, Kröger S, Laszig R, Beck RL, Schild C, Arndt S: Unilateral deafness in children: audiologic and subjective assessment of hearing ability after cochlear implantation. *Otol Neurotol* 2012;34:53-60.
- Hallam RS, Jakes JC, Hinchcliffe R: Cognitive variables in tinnitus annoyance. *Br J Clin Psychol* 1988;27:213-222.
- Hoare DJ, Edmondson-Jones M, Sereda M, Akeroyd MA, Hall D: Amplification with hearing aids for patients with tinnitus and co-existing hearing loss. *Cochrane Database of Systematic Reviews* DOI: 10.1002/14651858.CD010151.pub2.
- Jacob R, Stelzig Y, Nopp P, Schleich P: Audiologische Ergebnisse mit Cochleaimplantat bei einseitiger Taubheit. *HNO* 2011;59:453-460.
- Jastreboff PJ: Tinnitus retraining therapy. *Br J Audiol* 1999;33(1):68-70.
- Kleinjung T, Steffens T, Strutz J, Langguth B: Curing tinnitus with a cochlear implant in a patient with unilateral sudden deafness: a case report. *Cases J* 2009;2:7462.
- Landsberger DM, Svrakic M, Roland JT, Svirsky M: The relationship between insertion angles, default frequency allocations, and spiral ganglion place pitch in cochlear implants. *Ear Hear* 2015;[Epub ahead of print].
- Langguth B, Kreuzer PM, Kleinjung T, De Ridder D: Tinnitus: causes and clinical management. *Lancet Neurol* 2013;12:920-30.
- Maes IHL, Cima RFF, Vlaeyen JW, Anteunis LJC, Joore MA: Tinnitus: a cost study. *Ear Hear* 2013;34(4):508-514.
- Mazurek B, Szczepek AJ, Hebert S: Stress and tinnitus. *HNO* 2015;63:258-265.
- Meeus O, Blaivie C, Van de Heyning P: Validation of the Dutch and the French version of the Tinnitus Questionnaire. *B-ENT* 2007;3(Suppl 7):11-17.
- Meikle MB, Henry JA, Griest SE, Stewart BJ, Abrams HB, McArdle R, Myers PJ, Newman CW, Sandridge S, Turk DC, Folmer RL, Frederick EJ, House JW, Jacobson GP, Kinney SE, Martin WH, Nagler SM, Reich GE, Searchfield G, Sweetow R, Vernon JA: The Tinnitus Functional Index: development of a new clinical measure for chronic, intrusive tinnitus. *Ear Hear* 2012;33(2):153-76.
- Mertens G, Kleine Punte A, De Ridder D, Van de Heyning P: Tinnitus in a single-sided deaf ear reduces speech reception in the nontinnitus ear. *Otol Neurotol* 2013;34:662-666.
- Moerel M, De Martino F, Formisano E: An anatomical and functional topography of human auditory cortical areas. *Front Neurosci* 2014;8:225.
- Newman CW, Jacobson GP, Spitzer JB: Development of the Tinnitus Handicap Inventory. *Arch Otolaryngol Head Neck Surg* 1996;122:143-148.
- Smit JV, Janssen ML, Schulze H, Jahanshahi A, Van Overbeeke JJ, Temel Y, Stokroos RJ: Deep brain stimulation in tinnitus: current and future perspectives. *Brain Res* 2015;1608:51-65.
- Stakhovskaya O, Sridhar D, Bonham BH, Leake P: Frequency map for the human cochlear spiral ganglion: implications for cochlear implants. *JARO* 2007;8:220-233.
- Stelzig Y, Jacob R, Mueller J: Preliminary speech recognition results after cochlear implantation in patients with unilateral hearing loss: a case series. *Journal of Medical Case Reports* 2011;5:343.
- Tass PA, Adamchic I, Freund H-J, von Stackelberg T, Hauptmann C: Counteracting tinnitus by acoustic coordinated reset neuromodulation. *Restor Neurol Neurosci* 2012;30:137-159.
- Tyler RS: Patients preferences and willingness to pay for tinnitus treatments. *J Am Acad Audiol* 2012;23:115-125.
- Tyler RS, Ji H, Perreau A, Witt S, Noble W, Coelho C: Development and validation of the tinnitus primary function questionnaire. *Am J Audiol* 2014;23(3):260-72.
- Tyler RS, Keiner AJ, Walker K, Deshpande AK, Witt S, Killian M, Ji H, Patrick J, Dillier N, van Dijk P, Lai WK, Hansen MR, Gantz B: A series of case studies of tinnitus suppression with mixed background stimuli in a cochlear implant. *Am J Audiol* 2015;[Epub ahead of print].
- Van Abel KM, Dunn CC, Sladen DP, Oleson JJ, Beatty CW, Neff BA, Hansen M, Gantz BJ, Driscoll CL: Hearing preservation among patients undergoing cochlear implantation. *Otol Neurotol* 2015;36(3):416-21.
- Van de Heyning P, Vermeire K, Diebl M, Nopp P, Anderson I, De Ridder D: Incapacitating unilateral tinnitus in single-sided deafness treated by cochlear implantation. *Ann Oto Rhinol Laryn* 2008;117(9):645-652.
- Vermeire K, Van de Heyning P: Binaural hearing after cochlear implantation in subjects with unilateral sensorineural deafness and tinnitus. *Audiol Neurotol* 2009;14:163-171.
- Zeman F, Koller M, Schecklmann M, Langguth B, Landgrebe M: Tinnitus assessment by means of standardized self-report questionnaires: Psychometric properties of the Tinnitus Questionnaire (TQ), the Tinnitus Handicap Inventory (THI), and their short versions in an international and multi-lingual sample. *Health and Quality of Life Outcomes* DOI: 10.1186/1477-7525-10-128.
- Zimmerman E, Timboe A: Tinnitus: steps to take, drugs to avoid. *J Fam Pract* 2014;63(2):82-8.