

# Kinetic aspects of the interaction of blood clotting enzymes

Citation for published version (APA):

Hemker, H. C., Kop, J., & Willems, G. M. (1979). Kinetic aspects of the interaction of blood clotting enzymes: VIII. The relation between clotting time and clotting velocity. *Thrombosis and Haemostasis*, 41(2), 309-313. <https://doi.org/10.1055/s-0038-1646650>

## Document status and date:

Published: 01/01/1979

## DOI:

[10.1055/s-0038-1646650](https://doi.org/10.1055/s-0038-1646650)

## Document Version:

Other version

## Please check the document version of this publication:

- A submitted manuscript is the version of the article upon submission and before peer-review. There can be important differences between the submitted version and the official published version of record. People interested in the research are advised to contact the author for the final version of the publication, or visit the DOI to the publisher's website.
- The final author version and the galley proof are versions of the publication after peer review.
- The final published version features the final layout of the paper including the volume, issue and page numbers.

[Link to publication](#)

## General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal.

If the publication is distributed under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license above, please follow below link for the End User Agreement:

[www.umlib.nl/taverne-license](http://www.umlib.nl/taverne-license)

## Take down policy

If you believe that this document breaches copyright please contact us at:

[repository@maastrichtuniversity.nl](mailto:repository@maastrichtuniversity.nl)

providing details and we will investigate your claim.

# THROMBOSIS AND HAEMOSTASIS

(formerly Thrombosis et Diathesis Haemorrhagica)

Journal of the International Society on Thrombosis and Haemostasis

No. 2 · Vol. 41  
April 23, 1979

Editor:  
Dr. Rosemary Biggs,  
Oxford

Associate Editors: E. A. Beck, Bern · F. Duckert, Basel · G. I. C. Ingram, London Congress · V. J. Marder, Philadelphia Congress · C. R. Rizza, Oxford · H. R. Roberts, Chapel Hill · P. N. Walsh, Philadelphia

## Kinetic Aspects of the Interaction of Blood Clotting Enzymes

### VIII. The Relation between Clotting Time and Clotting Velocity

H. C. HEMKER, J. KOP, G. M. WILLEMS

*Department of Biochemistry, State University Limburg, Maastricht, The Netherlands.*

We have shown that the relation between coagulation time and prothrombin concentration in a one-stage prothrombin assay is

$$t_c = a/c + b \quad (1)$$

where  $t_c$  = clotting time,  $c$  = prothrombin concentration and  $a, b$  = constants. The same type of relation very probably holds when  $c$  is the concentration of any of the coagulation factors V, VII, or X. (Hemker et al. 1967, 77). In a reaction catalyzed by an enzyme according to the Michaelis-Menten model the reaction velocity is given by

$$1/v = a/c + b \quad (2)$$

Comparison of (1) and (2) immediately suggests that clotting time might be inversely proportional to the velocity of coagulation. This article reports experiments to verify this suggestion.  $c$  may apply to either a substrate or an enzyme under the conditions prevailing in coagulation.

#### Theoretical Considerations

Reaction velocities are additive, that is: the result of two or more reaction velocities ( $v_1, v_2$ , etc.) yielding the same product is product formation ( $v$ ), with a velocity equal to the sum of the constituent velocities,  $v = v_1 + v_2$ , etc., in general  $v = \sum_{i=1}^n v_i$ .

In the course of a reaction, the velocity is increased stepwise, then one can calculate the velocity which would have had the same effect as the discontinuous velocities occurred. In Fig. 1 an initial velocity  $v_1$  is increased stepwise to a final velocity  $v_2$ .



F. K. SCHATTAUER VERLAG · STUTTGART – NEW YORK

# Kinetic Aspects of the Interaction of Blood Clotting Enzymes

## VIII. The Relation between Clotting Time and Clotting Velocity

H. C. HEMKER, J. KOP, G. M. WILLEMS

*Department of Biochemistry, State University Limburg, Maastricht, The Netherlands.*

### Summary

It is shown that in a system where thrombin acts on fibrinogen, the clotting time can be used to assess coagulation velocity because clotting time and clotting velocity are inversely proportional.

### Introduction

We have shown that the relation between coagulation time and prothrombin concentration in a one-stage prothrombin assay is

$$t_c = a.1/c + b \quad (1)$$

where  $t_c$  = clotting time,  $c$  = prothrombin concentration and  $a$ ,  $b$  = constants. The same type of relation very probably holds when  $c$  is the concentration of any of the coagulation factors V, VII, or X. (Hemker et al. 1967, 77). In a reaction catalyzed by an enzyme according to the Michaelis Menten model the reaction velocity is given by

$$1/v = a.1/c + b \quad (2)$$

Comparison of (1) and (2) immediately suggests that clotting time might be inversely proportional to the velocity of coagulation. This article reports experiments to verify this suggestion.  $c$  may apply to either a substrate or an enzyme under the conditions prevailing in coagulation.

### Theoretical Considerations

Reaction velocities are additive, that is: the result of two or more reaction velocities ( $v_1$ ,  $v_2$  etc.) yielding the same product is product formation ( $v_r$ ), with a velocity equal to the sum of the constituent velocities.  $v_r = v_1 + v_2$  etc., in general  $v_r = \sum_{i=1}^n v_i$

When in the course of a reaction, the velocity is increased stepwise, then one can calculate the constant velocity which would have had the same effect as the discontinuous velocities that actually occurred. In Fig. 1 an initial velocity  $c$  is increased at moment  $a$  to the level  $d$ . Then at moment  $b$  the total effect of the two velocities is taken to be equivalent to one other




Fig. 1 Explanation in text.

velocity  $e$ . Because velocities are additive  $e(a + b) = c \times a + d \times b$   
 from which it follows that

$$e = \frac{db + ca}{a + b} \tag{3}$$

In coagulation we can execute this experiment by adding a given amount of thrombin ( $T_0$ ) at zero time, and after an interval ( $t_i$ ) before clotting occurs, add a second amount to obtain a higher concentration of thrombin ( $T_n$ ). Then one measures the clotting time  $t_{exp}$ . In an independent experiment one measures the clotting times that will be obtained by  $T_0$  or  $T_n$  added at zero time. We will call them  $t_0$  and  $t_n$ , respectively.

On the assumption that the clotting time is inversely proportional to clotting velocity we can translate this experiment in terms of Fig. 1, because ( $n$  is a proportionality constant)  $t_0 = n 1/c$ ;  $t_n = n 1/d$  and  $t_{exp} = a + b = n.1/e$ . Applying this to formula (3) gives:

$$1/t_{exp} = \frac{1/t_n (t_{exp} - t_i) + 1/t_0 t_i}{t_{exp}}$$

from which it follows that

$$t_{exp} = t_n + t_i - (t_n/t_0)t_i \tag{4}$$

$$t_{exp} = t_n + \frac{(t_0 - t_n)}{t_0} t_i$$

This formula can be put to test experimentally.



Fig. 2 Explanation in text.

## Materials and Methods

The experiments were carried out in a thermostated (37°C) reaction vessel of a Radiometer pH Stat placed on a magnetic stirrer. As a stirring magnet a 5 mm steel rod was used. The vessel contained 500  $\mu$ l Michaelis buffer pH 7.2 and 250  $\mu$ l fibrinogen solution (Kabi fibrinogen 100 mg/ml). At zero time 50  $\mu$ l of a thrombin solution were added (Thrombin Roche 50 U/ml or suitable dilution, kept at 0°C). At variable intervals before clotting occurred again 50  $\mu$ l of thrombin were added. The moment of clotting was assessed by observing the first white thread that winds around the stirring magnet (Fig. 2).

In control experiments it could be shown that the dilution of the fibrinogen brought about by the second addition of thrombin did not change the coagulation time to a significant degree. The dilution brought about by the addition therefore was neglected.

### Experimental

A representative experiment is shown in Table 1. In order to prevent systematic deviations each individual determination was assigned a random number and carried out in that order.

Table 1

Interval time ( $t_i$ ) (sec)	0	5	10	15	20	25	30	35	$\infty$
Coagulation times ( $t_{exp}$ ) (sec)	26.8	28.5	31.2	33.3	35.4	38.5	40.3	42.2	49.7
	26.0	29.0	31.5	34.1	34.7	37.6	39.8	42.7	49.0
	27.0	28.7	31.3	33.5	35.2	37.5	39.6	42.5	49.1
	26.3	29.2	31.2	33.2	35.0	37.6	39.6	42.3	48.4
	26.6	28.3	31.4	32.6	35.4	37.7	40.1	42.0	49.8
	26.1	28.6	31.3	33.0	34.8	37.8	40.1	42.2	48.8
	26.3	28.8	31.9	33.3	35.9	37.5	40.0	41.9	48.9
	26.4	28.0	31.4	32.7	36.5	37.5	39.7	42.0	48.9
	27.3	28.6	30.5	33.5	36.0	38.5	39.9	41.7	49.4
	26.5	28.6	30.7	34.0	35.8	37.6	40.1	42.6	49.0
Mean	26.5	28.6	31.2	33.3	35.5	37.8	39.9	42.2	49.1
Standard deviation	0.4	0.3	0.4	0.5	0.6	0.4	0.3	0.3	0.4

At zero time 50  $\mu$ l of a thrombin solution were added; after the interval time indicated the same amount was again added. Infinite interval times indicate no second addition. In terms of formula (4) we found  $t_n = 26.5$  and  $t_o = 49.1$ . When  $t_{exp}$  is plotted as a function of interval time, Fig. 3 is obtained. We see that  $t_{exp}$  is a linear function of  $t_1$ . We measure a slope of 0.46. According to formula (4) this slope should be

$$\frac{t_o - t_n}{t_o} = \frac{49.1 - 26.5}{49.1} = 0.46.$$

In repeated experiments of the same type the slopes calculated from equation (4) were found to be equal within the experimental error.

## Discussion

The evidence obtained does not counteract the derivations that are based on the assumption that clotting time is inversely proportional to clotting velocity. Any other type of




Fig.3 Moment of clotting as observed in the reaction vessel.

relation between clotting time and clotting velocity would have led to different formula. We therefore cannot falsify the original assumption. Still it may be possible that a type of relationship which resembles  $v = n.1/t_c$  very much (for instance  $v = n.1/t_c + a$  with small  $a$  etc.) would describe the situation better. As the experimental limits of our method are the same as in those where the relationship will be used in practice, i. e. the measuring of clotting times, this will be of little practical importance.

Résumé

Il est démontré que dans un système où la thrombine agit sur le fibrinogène, le temps de coagulation peut être utilisé pour établir la vitesse de coagulation car le temps de coagulation et la vitesse de coagulation sont inversement proportionnels.

Zusammenfassung

Es wird gezeigt, daß in einem System, in dem Thrombin Fibrinogen umsetzt, die Gerinnungszeit zur Bestimmung der Koagulationsgeschwindigkeit herangezogen werden kann, da die Gerinnungszeit und die Gerinnungsgeschwindigkeit invers proportional sind.

**References**

- HEMKER, H. C. and HEMKER, P. W. (1969): The kinetics of enzyme cascades. *Proceedings of the Royal Society B* 173, 411.
- HEMKER, H. C., SIEPEL, T., ALTMAN, R. and LOELIGER, E. A. (1967): Kinetic aspects of the interaction of blood clotting enzymes. II. The relation between clotting time and plasma concentration in prothrombin-time estimations. *Thrombosis et Diathesis Haemorrhagica* 17, 350.
- HEMKER, H. C., VERMEER, C. and GOVERS-RIEMSLAG, J. (1977): Kinetic aspects of the interaction of blood clotting enzymes. VII. The relation between clotting time and prothrombin concentration. *Thrombosis and Haemostasis* 37, 81.

Received 12th December, 1977 Accepted 15th March, 1978.

## Kinetic Aspects of the Interaction of Blood Clotting Enzymes

### VIII. The Relation between Clotting Time and Clotting Velocity

H. C. HEMKER, J. KOP, G. M. WILLEMSEN

*Department of Biochemistry, State University Limburg, Maastricht, The Netherlands.*

