

Linking surgical skills to postoperative outcomes

Citation for published version (APA):

Beulens, A. J. W., Brinkman, W. M., Van der Poel, H. G., Vis, A. N., van Basten, J. P., Meijer, R. P., Wijburg, C. J., Hendriks, A. J. M., van Merriënboer, J. J. G., & Wagner, C. (2019). Linking surgical skills to postoperative outcomes: a Delphi study on the robot-assisted radical prostatectomy. *Journal of Robotic Surgery*, 13(5), 675-687. <https://doi.org/10.1007/s11701-018-00916-9>

Document status and date:

Published: 01/10/2019

DOI:

[10.1007/s11701-018-00916-9](https://doi.org/10.1007/s11701-018-00916-9)

Document Version:

Publisher's PDF, also known as Version of record

Document license:

Taverne

Please check the document version of this publication:

- A submitted manuscript is the version of the article upon submission and before peer-review. There can be important differences between the submitted version and the official published version of record. People interested in the research are advised to contact the author for the final version of the publication, or visit the DOI to the publisher's website.
- The final author version and the galley proof are versions of the publication after peer review.
- The final published version features the final layout of the paper including the volume, issue and page numbers.

[Link to publication](#)

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal.

If the publication is distributed under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license above, please follow below link for the End User Agreement:

www.umlib.nl/taverne-license

Take down policy

If you believe that this document breaches copyright please contact us at:

repository@maastrichtuniversity.nl

providing details and we will investigate your claim.



Linking surgical skills to postoperative outcomes: a Delphi study on the robot-assisted radical prostatectomy

A. J. W. Beulens^{1,2,12} · W. M. Brinkman³ · H. G. Van der Poel⁴ · A. N. Vis⁵ · J. P. van Basten⁶ · R. P. Meijer⁷ · C. J. Wijburg⁸ · A. J. M. Hendriks⁹ · J. J. G. van Merriënboer¹⁰ · C. Wagner^{1,11}

Received: 9 October 2018 / Accepted: 18 December 2018 / Published online: 4 January 2019
© Springer-Verlag London Ltd., part of Springer Nature 2019

Abstract

Objective To develop an assessment instrument for the evaluation of surgical videos to elucidate the association between surgical skills and postoperative outcomes after a robot-assisted radical prostatectomy (RARP).

Design A Delphi study consisting of two consecutive online surveys and a consensus group meeting.

Setting Urology departments of general, teaching and university hospitals in the Netherlands.

Participants All Dutch urologists with a specialization in RARP.

Results Of 18 invited experts, 12 (67%) participated in the first online survey. In the second round, 9 of the 18 invited experts participated (50%). The Delphi meeting was attended by 5 of the 18 (27%) invited experts. The panel identified seven surgical steps with a possible association to postoperative outcomes. The experts also expected an association between adverse postoperative outcomes and the frequency of camera removals, the number of stitches placed, the amount of bleeding, and the extent of coagulation. These factors were incorporated into an assessment instrument.

Conclusions Experts in the field of RARP achieved consensus on 7 surgical steps and 4 aspects of the RARP procedure that may be related to adverse postoperative outcomes. The resulting assessment instrument will be tested in future research to determine its validity.

Keywords Delphi · RARP · Video analysis · Prostatectomy · Surgical skills · Postoperative outcomes

Introduction

In the Netherlands, approximately 2500 radical prostatectomies are performed annually of which 90% are performed using the surgical robot, i.e. robot-assisted radical

prostatectomy (RARP). The RARP is a complex but highly standardized operation to cure local prostate cancer. However, RARP is hampered by serious side-effects [1–3] such as urinary incontinence, which occurs in 4 to 26% of the

✉ A. J. W. Beulens
Alexander.beulens@catharinaziekenhuis.nl

¹ Netherlands Institute for Health Services Research (NIVEL), Utrecht, The Netherlands

² Department of Urology, Catharina Hospital, Eindhoven, The Netherlands

³ Department of Urology, Erasmus Medical Centre, Rotterdam, The Netherlands

⁴ Department of Urology, Dutch Cancer Institute-Antoni van Leeuwenhoek Hospital, Amsterdam, The Netherlands

⁵ Department of Urology, VU Medical Centre, Amsterdam, The Netherlands

⁶ Department of Urology, Canisius Wilhelmina Hospital, Nijmegen, The Netherlands

⁷ Department of Oncological Urology, University Medical Centre Utrecht, Utrecht, The Netherlands

⁸ Department of Urology, Rijnstate Hospital, Arnhem, The Netherlands

⁹ Department of Urology, Urologist NP, Catharina Hospital, Eindhoven, The Netherlands

¹⁰ School of Health Professions Education, Maastricht University, Maastricht, The Netherlands

¹¹ VU Medical Centre, Amsterdam Public Health Research Institute, Amsterdam, The Netherlands

¹² Catharina Hospital Eindhoven, P.O. Box 1350, 5602 ZA Eindhoven, The Netherlands

patients [4–6], and erectile dysfunction, which occurs in 14 to 90% of the patients [7, 8].

Previous research has shown that greater surgical experience is associated with better postoperative outcomes [9–11]. Therefore, the Dutch Society of Urology (NVU) increased the minimally required number of annual RARP per hospital from 50 to 100 procedures to improve functional results and reduce complications. However, at the moment there is no minimum annual number of procedures per surgeon.

Various authors suggested that systematic evaluation of skills, both technical (surgical) and non-technical (communication and teamwork) may be more effective in improving the surgeons' skills than a quota alone [12, 13]. Thorough analysis of surgical videos can possibly elucidate which steps or facets of surgery may be related to disadvantageous results such as postoperative complications (i.e. bleeding and leakage of the vesico-urethral anastomosis) and adverse functional outcomes (i.e. erectile dysfunction, incontinence) [12, 14].

To standardize video analysis, a detailed description of all the separate surgical steps is needed. In the past, different assessment instruments containing individual steps of the RARP have been defined [15–17], but these methods are mostly intended for providing feedback during training of new robotic surgeons or to evaluate the skills of more experienced robotic surgeons by means of video analysis. So far, no specific method has been developed to investigate how a surgeon's skills and surgical events as assessed on video are related to adverse postoperative outcomes of RARP.

The present Delphi study is designed to evaluate whether experts in the field of RARP can identify the surgical and non-surgical factors in RARP that are potentially associated with negative aspects of postoperative outcomes.

The following key questions were to be answered: which steps of the RARP and which peri-operative events (i.e. bleeding, usage of coagulation, usage of haemostatic clips and suturing) are most likely associated with postoperative complications (i.e. bleeding and leakage of the vesico-urethral anastomosis) and adverse functional outcomes (i.e. erectile dysfunction, incontinence)? How can these steps of the RARP and these peri-operative events be incorporated in an RARP assessment instrument?

Methods

During a focus group consisting of three Dutch urologists, one urologist in training, and one cognitive task analysis expert a list of statements was created, describing the surgical steps and possible peri-operative events of the RARP procedure as well as their possible association with (1) direct postoperative complications and (2) functional outcomes.

These statements were formulated in order to investigate which steps of the surgery and which peri-operative events should be included in an instrument for video analysis. This assessment instrument will form the basis for further research on the possible associations between surgical skills and adverse postoperative outcomes.

Expert panel

The expert panel for the Delphi study was selected based on recommendations of three separate independent urologists who are experts in the field of robotic surgery. Based on these recommendations, 18 experts in the field of robot-assisted radical prostatectomy were selected. In this group, multiple proctors and educators of different fellowships in robotic surgery were included since they have intricate knowledge of the possible origins of complications in surgeons with all levels of experience. The experts were invited by e-mail. If no response was given the experts were contacted by telephone to ask whether they were interested to participate in the Delphi study.

Consensus procedure

To achieve consensus, a two-step procedure was used (Appendix Fig. 1): the first step was an online two-round Delphi Survey involving Dutch urologists experienced in RARP. The second step was a consensus group meeting with the same Dutch urologists to discuss the results of the online survey and to identify the aspects of the surgery and the peri-operative events which might be associated with postoperative adverse outcomes. The steps of the Delphi process are based on protocols for consensus finding [18–21].

Online two-round Delphi Survey

The results of the initial focus group were used to define seven domains in which the statements could be categorized. The domains were organized as follows: (Tables 1, 2).

1. The relation of the statement to postoperative complications;
2. The relation of the statement to functional results;
3. Surgical steps associated with complications (i.e. bleeding and leakage of the vesico-urethral anastomosis);
4. Surgical steps associated with postoperative erectile dysfunction;
5. Surgical steps associated with postoperative urinary incontinence;
6. Factors that play a role in the origins of postoperative complications;
7. Elements that should certainly be included in the training of novice surgeons.

Table 1 Domains of statements used in the online Delphi Survey and consensus meeting

Domain	Statements in survey (n)	Consensus in survey (n)	Consensus survey (%)	Statements in Delphi panel (n)	Consensus Delphi panel (n)	Consensus Delphi panel (%)	Consensus combined Delphi Survey and panel (%)
The relation of the statement to postoperative complications	14	9	64	6	3	50	79
The relation of the statement to functional results	14	7	50	7	2	29	64
Steps of the surgery associated with complications	11	9	82	2	0	0	82
Steps of the surgery associated with postoperative erectile dysfunction	11	9	82	2	1	50	91
Steps of the surgery associated with postoperative urinary incontinence	11	8	73	3	2	66	91
Factors that play a role in the origins of postoperative complications	4	4	100	–	–	–	100
Elements that are essential for the training of novice surgeons	7	7	100	–	–	–	100

The statements were used to design a two-round online Delphi Survey to obtain consensus on identifying the relevant steps of the RARP procedure and their possible causal relation to postoperative complications and adverse functional outcomes.

The panel members were asked to rate the relevance of each statement using a 9-point Likert scale according to the discriminatory power of each surgical step to correspond with the specified postoperative complication. A rating of 1 was defined as “extremely disagree” and a rating of 9 was defined as “extremely agree”. As described in the RAND/UCLA Appropriateness Method [22], for each item, the median agreement score, lower limit interpercentile range (IPR), and upper limit IPR and Disagreement Index (DI) were calculated. A median agreement score of 1.0–3.0 was considered to be “disagree”, 3.1–6.9 as “uncertain”, and 7.0–9.0 as “agree”. A DI value above one (> 1) indicated a lack of consensus among the participants regarding the association between the statement and the postoperative complication.

In addition to the consensus statements, seven general questions (Table 3) were included in the first online survey to assess the experts opinions on the project and their willingness to cooperate in further research. In the first round, the participants were invited to suggest additional items that should be included in the second-round survey. The second survey consisted of the consensus statements. After each round, the scores for each item were anonymized to a mean

ranking score for the whole group and reported back to the participants.

Consensus group meeting

During a consensus panel meeting, the statements on which consensus had been reached in the two-round online survey were reviewed and statements on which no consensus had been reached were discussed and voted on. The meeting was chaired by a urologist of the Dutch Cancer Institute, Amsterdam (HvdP).

The statements from the online survey were presented to the panel, and participants were asked to motivate their opinions on each of the statements for which no consensus had been reached previously. The list of approved steps and aspects was then categorized to develop an initial RARP assessment instrument for evaluating the surgical procedure on video. This assessment instrument was subsequently judged on face validity by the 12 experts who participated in the Delphi process.

Informed consent

All procedures followed were in accordance with the ethical standards of the responsible committee on human experimentation (institutional and national) and with the Helsinki Declaration of 1975, as revised in 2013 [23]. Informed

Table 2 Results of the first and second online Delphi Survey and combined consensus results after Delphi meeting

Statement	MAS*	Consensus	MAS*	Consensus	Combined Consensus after Delphi meeting
	Round 1	Round 1	Round 2	Round 2	
The relation of the statement to postoperative complications					
Operating quickly results in better outcomes for the patient (speed is related to insight and therefore a good measure) in terms of complications	5	no	6	no	no
Shorter operating times result in fewer complications	6,5	no	7	agree	agree
The use of as few different instruments as possible lead to fewer complications	3	disagree	3	disagree	disagree
Not to using the 3rd arm of the robot leads to fewer complications.	1	disagree			disagree
Zooming in more on the operation field is better (closer gives better vision) as it leads to fewer complications.	5	no	5	no	no
More zooming out of the operating field is better (further away gives better overview and less dirt on the camera lens) as it leads to fewer complications.	3	disagree	3	disagree	disagree
The suture material used has an influence on the development of complications.	4,5	no	5	no	no
Fewer camera movements result in fewer complications.	5	no	3	disagree	disagree
Fewer instrument movements result in fewer complications	5	no	6	no	agree
A lower estimated blood loss results in fewer complications.	5	no	6	no	agree
A shorter duration of coagulation results in fewer complications	6	no	7	agree	agree
Placing fewer stitches results in fewer complications.	3	disagree	3	disagree	disagree
Placing fewer clips results in fewer complications.	3	disagree	4	disagree	disagree
Inspection of the abdomen leads to fewer complications	7	agree	8	agree	agree
The relation of the statement to functional results					
Operating quickly results in better functional outcomes for the patient (speed is related to insight and therefore a good measure)	4	no	5	no	no
Shorter operating times results in improved functional results.	4	no	5	no	no
The use of as few different instruments as possible leads to better functional results	3,5	no	3	disagree	disagree
Not to using the 3rd arm of the robot leads to better functional results.	2	disagree	2	disagree	disagree

Table 2 (continued)

Zooming in more on the operation field is better (closer gives better vision) as it leads to better functional results.	7	agree	7	agree	agree
More zooming out of the operating field is better (further away gives better overview and less dirty of the camera lens) as it leads to better functional results	4	no	2,5	disagree	disagree
The suture material used has an influence on the functional results	6	no	6	no	no
Fewer camera movements result in improved functional results	3	disagree	5,5	no	no
Fewer instrument movements result in improved functional results	5	no	6,5	no	agree
A lower estimated blood loss results in improved functional results.	4	no	6	no	no
A shorter duration of coagulation results in improved functional results	7	agree	7	agree	agree
Placing fewer stitches results in improved functional results.	4	no	4	no	disagree
Placing fewer clips results in improved functional results	5	no	3	disagree	disagree
Inspection of the abdomen results in improved functional results	3	disagree	2,5	disagree	disagree
Steps of the surgery associated with complications					
Abdominal cavity approach/port placement	5	no	6	no	no
Retropubic space approach/mobilisation of Retzius	3	disagree	2	disagree	disagree
Pelvic floor muscle exposure/opening of the endopelvic fascia	5	no	5,5	no	no
Bladder neck dissection	7	agree	7,5	agree	agree
Ligation of prostate pedicles	7,5	agree	8,5	agree	agree
Nerve preservation	8	agree	8	agree	agree
Management of prostate apex/urethra	8	agree	8,5	agree	agree
Prostate removal	3	disagree	2,5	disagree	disagree
Urethro-vesical anastomosis	8	agree	8	agree	agree
Lymph node dissection	7,5	agree	8	agree	agree
Wound closure and specimen removal	7,5	agree	7	agree	agree
Steps of the surgery associated with postoperative erectile dysfunction					
Abdominal cavity approach/port placement	1	disagree			disagree
Retropubic space approach/mobilisation of Retzius	1	disagree			disagree
Pelvic floor muscle exposure/opening of the endopelvic fascia	6	no	6,5	no	no
Bladder neck dissection	3	disagree	2,5	disagree	disagree
Ligation of prostate pedicles	8	agree	8	agree	agree
Nerve preservation	9	agree			agree

Table 2 (continued)

Management of prostate apex/urethra	9	agree			agree
Prostate removal	1	disagree			disagree
Urethro-vesical anastomosis	7	agree	7	agree	agree
Lymph node dissection	4,5	no	4	no	agree
Wound closure and specimen removal	1	disagree			disagree
Steps of the surgery associated with postoperative urinary incontinence					
Abdominal cavity approach/port placement	1	disagree			disagree
Retropubic space approach/mobilisation of Retzius	1	disagree			disagree
Pelvic floor muscle exposure/opening of the endopelvic fascia	7	agree	7	agree	agree
Bladder neck dissection	6,5	no	6,5	no	no
Ligation of prostate pedicles	5	no	4,5	no	disagree
Nerve preservation	7	agree	6,5	no	agree
Management of prostate apex/urethra	9	agree			agree
Prostate removal	1	disagree			disagree
Urethro-vesical anastomosis	8,5	agree	9	agree	agree
Lymph node dissection	2	disagree	3	disagree	disagree
Wound closure and specimen removal	1	disagree		disagree	disagree
Factors that play a role in the origins of postoperative complications					
Teamwork	9	agree			agree
Communication between the surgeon and the surgical team	9	agree			agree
Surgical skills of the surgeon	9	agree			agree
Patient factors (i.e. Age, BMI, tumour size)	8	agree	7,5	agree	agree
Elements that are essential for the training of novice surgeons					
Theoretical education	8	agree	8,5	agree	agree
Simulator training, practice on virtual reality simulators	8	agree	8,5	agree	agree
Wetlab practice, training on animals	8	agree	8	agree	agree
Cadaver training	7	agree	7	agree	agree
Drylab training, practice on models	7	agree	7	agree	agree
Supervised practice on real patients	9	agree			agree
Fellowship	9	agree			agree

*MAS Median Agreement Score

Table 3 The participants and their institute, occupation and voting status

Participant	Institute	Occupation	Voting status
H. van der Poel	Antoni van Leeuwenhoek Hospital, Amsterdam/Netherlands Cancer Institute, Amsterdam	Urologist	Voting
R. Meijer	University Medical Center Utrecht	Urologist	Voting
H. Beerlage	Jeroen Bosch Hospital, Den Bosch	Urologist	Voting
M. Busstra	Erasmus Medical Centre, Rotterdam	Urologist	Voting
C. Wijburg	Rijnstate Hospital, Arnhem	Urologist	Voting
A. Hendrikk	Catharina Hospital, Eindhoven	Urologist NP	Non-voting
J. van Merienboer	Maastricht University	Educational expert	Non-voting
W. Brinkman	Erasmus Medical Center, Rotterdam	Urologist in training	Non-voting
A. Beulens	Catharina Hospital, Eindhoven/Netherlands institute for health services research (NIVEL), Utrecht	PhD student	Non-voting

consent was obtained from all participants for being included in the study.

Results

Delphi Survey

The results of the preliminary focus group meeting were used to formulate 72 statements on surgical steps and possible perioperative events of the RARP procedure and their possible association with (1) direct postoperative complications and (2) functional outcomes. These statements were divided over seven domains and incorporated in an online survey (Table 1).

A total of 18 Dutch experts in robot-assisted radical prostatectomy (RARP) were identified and invited to participate in the two-round online Delphi Survey. In the first round, 12 of the 18 (67%) invited experts participated in the survey. Of these 12 participants, 10 experts responded to all statements, and two participants reported difficulties with the survey resulting in a partial response to the statements.

In the first round, participants did not propose any additional statements. Of the 72 statements reviewed in the first round, 18 statements on which a clear consensus had been reached (i.e. a median agreement score of 1 or 9) were excluded from the second round. The remaining 54 statements were incorporated in the second online survey round.

In the second round, nine of the 12 participants of the first round participated in the survey. Of these nine participants, eight completed the survey and one reported difficulties with the survey resulting in a partial response to the questionnaire.

General questions

Results of the general questionnaire (Table 3) show that 83.0% of the experts believe that patient outcome can be improved by analysis of critical surgical factors. According

to 75.0% and 83.3% of the experts who participated in this study, video assessment is suitable for predicting complications and functional patient outcomes, respectively. According to 92.0% of the experts, the use of video assessment could reduce the risk of complications. All experts were interested in participating in the analysis of surgical videos. Most experts had the means to record surgical videos (80.0%) and can link these videos to surgical data (90.1%).

Consensus group meeting

Of the 18 invited experts, five participated in the consensus group meeting. In total, this meeting was attended by nine participants, whose occupation and voting status are presented in Table 4.

Final consensus statements

Table 2 shows the statements on which consensus was reached, organized per domain. The results of the Delphi Survey and the consensus group meeting were used to develop the assessment instrument PROTEST (PROstatectomy video Observation to Evaluate and Score Technical skill) (Appendix Table 5). This instrument contains the seven steps of the RARP surgery and the perioperative measurements that are considered to be most likely to be related to complications and adverse postoperative outcomes.

The relation between the statements and postoperative complications

Consensus of ‘agreement’ was reached on three out of 14 statements (Table 2) regarding the relation of the statement to postoperative complications. Consensus of ‘disagreement’ was reached on six out of 14 statements, one of which received a unanimous ‘disagreement’ score (i.e. median score of 1 and disagreement index = 0). No consensus was

Table 4 Results of general questions about video analysis, registration of postoperative outcomes, and intention to participate in the analysis of RARP videos

Question	Percentage of respondents (n)
Do you believe that it is useful to analyse surgical factors in order to improve the outcomes of patients (several options possible)?	
Yes, because we can learn from mistakes made	83.33 (10)
Yes, because we can develop new and better surgical techniques	83.33 (10)
No, because there is a chance that the considerations of the surgeon and patient selection play a more important role than the actual operation	8.33 (1)
Is it possible in your view to predict possible perioperative and postoperative complications by means of video assessments?	
Yes	75 (9)
Is it possible in your view to predict postoperative (functional) outcomes by means of video assessments?	
Yes	83,33 (10)
Is it possible in your view to reduce the risk of complications by means of data obtained by video analysis?	
Yes	92% (11)
Are you prepared to participate in the analysis of surgical videos?	
Yes	100% (11)
Do you record the robot-assisted radical prostatectomy procedure on videos?	
Yes	80% (8)
Do you have the option to correlate outcome data such as complications and functional outcomes to surgical videos?	
Yes	90.1% (10)

reached on the five remaining statements. The panel agreed on the following statements: “Shorter operating times result in fewer complications” and “Shorter duration of coagulation results in fewer complications”. All participants disagreed with the statement “It is better not to use the third arm of the robot when looking at complications”.

The relation between the statements and functional results

The participants of the Delphi Survey reached consensus of ‘agreement’ on two out of 14 statements concerning functional results (Table 2). The panel reached a consensus of ‘disagreement’ on five out of 14 statements. None of the statements received a unanimous score. No consensus was reached on the seven remaining statements. The panel agreed on the following statements: “Zooming in more on the operation field provides better vision as it leads to better functional results.” and “A shorter duration of coagulation results in improved functional results.”

Steps of the surgery associated with complications

During the Delphi Survey, consensus of ‘agreement’ was reached on seven out of 11 statements regarding the steps of the surgery that might be associated with complications. The panel reached consensus of ‘disagreement’ on two out of 11 statements (Table 2). No statements received a unanimous score. No consensus was reached on two remaining statements.

The panel agreed that the following steps of the surgery might be associated with complications: “Bladder neck dissection”, “Ligation of prostate pedicles”, “Nerve preservation”, “Management of prostate apex/urethra”, “Vesico-urethral anastomosis”, “Lymph node dissection”, and “Wound closure and specimen removal”.

Steps of the surgery associated with postoperative erectile dysfunction

The Delphi Survey panel reached a consensus of ‘agreement’ on four out of 11 statements regarding the steps of the surgery that might be associated with postoperative erectile dysfunction (Table 2). The experts unanimously agreed that “Nerve preservation” and “Management of prostate apex/urethra” might be associated with the incidence of postoperative erectile dysfunction. A consensus of ‘disagreement’ was reached on five out of 11 statements, four of which received a unanimous ‘disagreement’ score. No consensus was reached on the two remaining statements.

Steps of the surgery associated with postoperative urinary incontinence

During the Delphi Survey, the panel reached a consensus of ‘agreement’ on two out of 11 statements (Table 2) regarding steps of the surgery that might be related to postoperative urinary incontinence. The panel reached a consensus of ‘disagreement’ on five out of 11 of these

statements. No consensus was reached on the four remaining statements. A unanimous consensus of ‘agreement’ was reached on one of these steps, and a unanimous consensus of disagreement was reached on three of these steps.

The panel agreed that the following steps of the surgery might be associated with postoperative urinary incontinence: “Pelvic floor muscle exposure/opening of the endopelvic fascia”, and “Urethrovesical anastomosis”.

Factors that play a role in the origins of postoperative complications

Both the Delphi Survey and the consensus meeting reached a consensus of ‘agreement’ that all (four out of four) the proposed factors (Table 2) could play a role in the origins of postoperative complications. The experts unanimously agreed on the relevance of the following factors: “Team-work”, “Communication between the surgeon and the surgical team”, and “Surgical skills of the surgeon”.

Elements that are essential for the training of novice surgeons

The participants of both the Delphi Survey and the consensus meeting reached a consensus of ‘agreement’ that all (seven out of seven) proposed elements of training (Table 2) are essential for the training of novice surgeons. There was unanimous agreement on the need to implement the following training assessment methods: “Supervised practice on real patients”, and “Fellowship”.

Discussion

The aim of this study was to develop an assessment instrument for the evaluation of surgical videos to elucidate the association between surgical skills and postoperative outcomes after a robot-assisted radical prostatectomy (RARP). To investigate this association, we invited all Dutch experts in RARP to participate in a standardized Delphi procedure in order to identify surgical and non-surgical factors in RARP that are potentially associated with an adverse postoperative course and to assess whether any of these parameters may be worth evaluating for the prediction of postoperative outcomes.

We found that the majority of Dutch urologists specialized in RARP are interested in an instrument for video analysis of their surgical skills in relation to the postoperative outcomes. These urologists indicated that they were interested to participate in the current study because they considered video

analysis to be useful for the improvement of surgical skills and the subsequent reduction of postoperative complications.

Consensus group meeting

During the consensus group meeting the panel members agreed that the duration of the surgical procedure and the duration of coagulation could be causally related to the rate of postoperative complications. They advised to investigate whether such a causal relation exists. The panel also agreed there could be a causal relation between the duration of coagulation and the level of adverse postoperative functional results.

The panel agreed that postoperative complications, postoperative erectile dysfunction and postoperative urinary incontinence could result from events during the following steps of the RARP procedure: “Pelvic floor muscle exposure/opening of the endopelvic fascia”, “Bladder neck dissection”, “Ligation of prostate pedicles”, “Nerve preservation”, “Management of prostate apex/urethra”, “Urethrovesical anastomosis”, “Lymph node dissection”, and “Wound closure and specimen removal”. The panel agreed that these steps should be incorporated in an assessment instrument to investigate whether they are related to adverse postoperative outcomes.

PROTEST assessment instrument

Based on the consensus reached during the Delphi Survey and the consensus meeting, the PROTEST assessment instrument was developed (Appendix 2). This instrument can be used to assess the skills of a surgeon through analysis of a video recording of the surgery.

This assessment instrument was developed with the input of the panel members and was reviewed by all panel members in order to give them the opportunity to refine and clarify the assessment instrument. One item, “Wound closure and specimen removal”, was not included in the PROTEST assessment instrument, because this step is not recorded on surgical videos.

When comparing the results of the current Delphi study to existing assessment instruments [15, 24, 25], the developed PROTEST assessment instrument shows a combination of subjective surgical skills assessment and objective metrics of procedural steps and events. The developed PROTEST assessment instrument is different from the GEARS assessment instrument where the focus lies solely on the subjective scoring of 5 technical domains of surgical skill, with no objective measurements [23]. The GERT assessment instrument comparable to the PROTEST assessment instrument as it focusses on different

features of the surgery (i.e. clipping, suturing, use of the retractors and use of suction), but it only gives feedback on possible errors whilst performing these features, there is no room to score subjective surgical skill [24]. The PACE assessment instrument is similar to the GERT assessment instrument as it evaluates specific steps of the RARP procedure, similar to the PROTEST assessment instrument, but only gives feedback based on errors whilst performing these steps and there is no room for subjective surgical skill analysis [15].

Implications of study findings for clinical practice and research

This Delphi procedure resulted in an overview of possible origins of complications after RARP and in a new assessment instrument that can be used to objectively assess a surgeon's skills.

The PROTEST assessment instrument gives detailed insight into the proficiency of the surgeon on each of the individual surgical steps of the RARP. It combines the answers to two general subjective questions with multiple objective measurements in order to provide detailed feedback to the surgeon.

Future studies should explore whether the factors identified in this Delphi process are indeed causally related to postoperative complications and whether video assessments by means of the PROTEST instrument can help in the training of novice surgeons and improving the skills of RARP surgeons.

Limitations

A limitation of this study is that we consulted all the urologists specialized in RARP who are registered in the Netherlands. Future studies with larger panels and international

participants might add other factors that could contribute to complications after RARP. A second limitation is that the answers to the general questions might be subject to participation bias and hence they cannot be generalized to the total group of Dutch urologists. Finally, the responses of panel members could have been influenced by the fact that the consensus meeting was not led by an independent chair.

Conclusion

Dutch experts in the field of RARP have reached consensus on seven surgical steps and four aspects of the RARP procedure that may be related to postoperative complications after RARP. The resulting assessment instrument, PROTEST, can be used to assess surgical skill. The resulting assessment instrument will be tested in future research to determine its validity for assessing the relationship between surgical skills and adverse postoperative outcomes after RARP.

Acknowledgements The authors thank H.P. Beerlage, A.E. Boeken Kruger, J.B.W. Rietbergen, J.P.M. Sedelaar, P.C. Weijerman, M.B. Busstra, O.S. Klaver, and B.C. Knipscheer for their participation during the Delphi process.

Compliance with ethical standards

Conflict of interest Drs. Beulens, Dr. Brinkman, Dr. van der Poel, Dr. Vis, Dr. van Basten, Dr. Meijer, Dr. Wijburg, Dr. Hendrikx, Prof. van Merriënboer, and Prof. Wagner have no conflicts of interest or financial ties to disclose.

Appendix

See Table 5 and Fig. 1.

Table 5 PROTEST assessment instrument

PROTEST assessment instrument						
Pelvic floor muscle exposure	Surgical skill	1 = uncoordinated	2	3	4	5 = perfect coordination
		1 = inaccurate	2	3	4	5 = perfectly accurate
	Total time step (s)					
	Time bleeding (s)					
	Time coagulating (s)					
	Time suturing (s)					
	Number of times camera removal (<i>n</i>)					
	Comments					
	Events					
	Bladder neck dissection	Surgical skill	1 = uncoordinated	2	3	4
		1 = inaccurate	2	3	4	5 = perfectly accurate
Total time step (s)						
Time bleeding (s)						
Time coagulating (s)						
Time suturing (s)						
Number of times camera removal (<i>n</i>)						
Comments						
Events						
Ligation of prostatic pedicles		Surgical skill	1 = uncoordinated	2	3	4
		1 = inaccurate	2	3	4	5 = perfectly accurate
	Total time step (s)					
	Time bleeding (s)					
	Time coagulating (s)					
	Time suturing (s)					
	Number of times camera removal (<i>n</i>)					
	Comments					
	Events					
	Nerve preservation	Surgical skill	1 = uncoordinated	2	3	4
		1 = inaccurate	2	3	4	5 = perfectly accurate
Total time step (s)						
Time bleeding (s)						
Time coagulating (s)						
Time suturing (s)						
Number of times camera removal (<i>n</i>)						
Comments						
Events						
Management of prostatic apex/urethra		Surgical skill	1 = uncoordinated	2	3	4
		1 = inaccurate	2	3	4	5 = perfectly accurate
	Total time step (s)					
	Time bleeding (s)					
	Time coagulating (s)					
	Time suturing (s)					
	Number of times camera removal (<i>n</i>)					
	Comments					
	Events					

Table 5 (continued)

PROTEST assessment instrument

Urethrovesical anastomosis	Surgical skill	1 = uncoordinated	2	3	4	5 = perfect coordination
		1 = inaccurate	2	3	4	5 = perfectly accurate
	Total time step (s)					
	Time bleeding (s)					
	Time coagulating (s)					
	Time suturing (s)					
	Number of times camera removal (<i>n</i>)					
	Comments					
	Events					
	Lymph node dissection (if applicable)	Surgical skill	1 = uncoordinated	2	3	4
		1 = inaccurate	2	3	4	5 = perfectly accurate
Total time step (s)						
Time bleeding (s)						
Time coagulating (s)						
Time suturing (s)						
Number of times camera removal (<i>n</i>)						
Comments						
Events						
Was bladder neck preservation attempted (Y/N)?						
Were both ureteral orifices in sight during preparation of the bladder neck? (Y/N)						
Was the capsula damaged during nerve sparing? (Y/N)						
Was there a tear in the vesiculae during preparation? (Y/N)						
Was the diathermia used during transection of the plexus of Santorini? (Y/N)						
Was the diathermia used during transection of the urethra? (Y/N)						
Was the colliculus in sight during transection of the urethra? (Y/N)						
Was a bladder neck reconstruction performed? (Y/N)						
Was the Rocco stitch (median fibrous raphe) reconstruction used? (Y/N)						
Was a barbed suture used for the bladder/urethra anastomosis? (Y/N)						
How many stitch throws were used in the anastomosis? (<i>n</i>)						
Total time surgery (s)						
Total time bleeding (s)						
Total time coagulation (s)						
Total time suturing (s)						
Total number of camera removals (<i>n</i>)						
Total number of events (<i>n</i>)						
Average score surgical skills						
2d/3d images						
Nerve sparing						
BMI						
Date of surgery						
Age of patient						
Tumour stage						
Prostate size						




Fig. 1 Visual representation of the Delphi Survey

References

- Barocas DA, Alvarez J, Resnick MJ, Koyama T, Hoffman KE, Tyson MD, Conwill R, McCollum D, Cooperberg MR, Goodman M, Greenfield S, Hamilton AS, Hashibe M, Kaplan SH, Paddock LE, Stroup AM, Wu X-C, Penson DF (2017) Association between radiation therapy, surgery, or observation for localized prostate cancer and patient-reported outcomes after 3 years. *JAMA* 317(11):1126
- Donovan JL, Hamdy FC, Lane JA, Mason M, Metcalfe C, Walsh E, Blazeby JM, Peters TJ, Holding P, Bonnington S, Lennon T, Bradshaw L, Cooper D, Herbert P, Howson J, Jones A, Lyons N, Salter E, Thompson P et al (2016) Patient-reported outcomes after monitoring, surgery, or radiotherapy for prostate cancer. *N Engl J Med* 375(15):1425–1437
- Lardas M, Liew M, van den Bergh RC, De Santis M, Bellmunt J, Van den Broeck T, Cornford P, Cumberbatch MG, Fossati N, Gross T, Henry AM, Bolla M, Briers E, Joniau S, Lam TB, Mason MD, Mottet N, van der Poel HG, Rouvière O et al (2017) Quality of life outcomes after primary treatment for clinically localised prostate cancer: a systematic review. *Eur Urol* 72(6):869–885
- Maike HJ, Cathelijne MP, Ziedses des Plantjes DM, Somford JA, van Erkelens RG, de Vries KH, Jozette JC, Bart P (2011) Wijsman MBB en J van L. Hoe vaak incontinentie na radicale prostatectomie? *Ned Tijdschr Geneesk* 162:D2294
- Wilson LC, Gilling PJ. Post-prostatectomy urinary incontinence: a review of surgical treatment options. *BJU Int* 107:7–10
- Goluboff ET, Saidi JA, Mazer S, Bagiella E, Heitjan DF, Benson MC, Olsson CA (1998) Urinary continence after radical prostatectomy: the Columbia experience. *J Urol* 159(4):1276–1280
- Tollefson MK, Gettman MT, Karnes RJ, Frank I (2011) Administrative data sets are inaccurate for assessing functional outcomes after radical prostatectomy. *J Urol* 185(5):1686–1690
- Salonia A, Burnett AL, Graefen M, Hatzimouratidis K, Montorsi F, Mulhall JP, Stief C (2012) Prevention and management of postprostatectomy sexual dysfunctions. Part 1: choosing the right patient at the right time for the right surgery. *Eur Urol* 62(2):261–272
- Vickers A, Savage C, Bianco F, Mulhall J, Guillonneau B, Cronin A, Scardino P (2011) Cancer control and functional outcomes after radical prostatectomy as markers of surgical quality: analysis of heterogeneity between surgeons at a single cancer center. *Eur Urol* 59(3):317–322
- Cathcart P, Sridhara A, Ramachandran N, Briggs T, Nathan S, Kelly J (2015) Achieving quality assurance of prostate cancer surgery during reorganisation of cancer services. *Eur Urol* 68(1):22–29
- Gershman B, Meier SK, Jeffery MM, Moreira DM, Tollefson MK, Kim SP, Karnes RJ, Shah ND (2017) Redefining and contextualizing the hospital volume-outcome relationship for robot-assisted radical prostatectomy: implications for centralization of care. *J Urol* 198(1):92–99
- Birkmeyer JD, Finks JF, O'reilly A, Oerline M, Carlin AM, Nunn AR, Dimick J, Banerjee M, Birkmeyer NJO (2013) Surgical skill and complication rates after bariatric surgery for the Michigan Bariatric Surgery Collaborative. *N Engl J Med* [Internet] 369:1434–1442
- Goldenberg MG, Lee JY, Kwong JC, Grantcharov TP, Costello A (2018) Implementing assessments of robotic-assisted technical skill in urologic education: a systematic review and synthesis of the validity evidence. *BJU Int*
- Goldenberg MG, Goldenberg L, Grantcharov TP (2017) Surgeon performance predicts early continence after robot-assisted radical prostatectomy. *J Endourol* 31(9):858–863
- Hussein AA, Ghani KR, Peabody J, Sarle R, Abaza R, Eun D, Hu J, Fumo M, Lane B, Montgomery JS, Hinata N, Rooney D, Comstock B, Chan HK, Mane SS, Mohler JL, Wilding G, Miller D, Guru KA (2017) Development and validation of an objective scoring tool for robot-assisted radical prostatectomy: prostatectomy assessment and competency evaluation. *J Urol* 197(5):1237–1244
- Morris C, Hoogenes J, Shayegan B, Matsumoto ED (2017) Towards development and validation of an intraoperative assessment tool for robot-assisted radical prostatectomy training: results of a Delphi study. *Int Braz J Urol* 43(4):661–670
- Stolzenburg J-U, Schwaibold H, Bhanot SM, Rabenalt R, Do M, Truss M, Ho K, Anderson C (2005) Modular surgical training for endoscopic extraperitoneal radical prostatectomy. *BJU Int* 96(7):1022–1027
- Gillissen S, Attard G, Beer TM, Beltran H, Bossi A, Bristow R, Carver B, Castellano D, Chung BH, Clarke N, Daugaard G, Davis ID, de Bono J, Borges dos Reis R, Drake CG, Eeles R, Efstathiou E, Evans CP, Fanti S et al (2018) Management of patients with advanced prostate cancer: the report of the advanced prostate cancer consensus conference APCCC 2017. *Eur Urol* 73(2):178–211
- van den Bos W, Muller BG, Ahmed H, Bangma CH, Barret E, Crouzet S, Eggenger SE, Gill IS, Joniau S, Kovacs G, Pahernik S, de la Rosette JJ, Rouvière O, Salomon G, Ward JF, Scardino PT (2014) Focal therapy in prostate cancer: international multidisciplinary consensus on trial design. *Eur Urol* 65(6):1078–1083
- Lutkenhaus LJ, van Os RM, Bel A, Hulshof MCCM (2016) Clinical results of conformal versus intensity-modulated radiotherapy using a focal simultaneous boost for muscle-invasive bladder cancer in elderly or medically unfit patients. *Radiat Oncol* 11(1):45
- MacLennan S, Bekema HJ, Williamson PR, Campbell MK, Stewart F, MacLennan SJ, N'Dow JM, Lam TB (2015) A core outcome set for localised prostate cancer effectiveness trials: protocol for a systematic review of the literature and stakeholder involvement through interviews and a Delphi survey. *Trials* 16(1):76
- Fitch K, Bernstein SJJ, Aguilar MDD, Burnand B, LaCalle JRR, Lazaro P, Van Het Loo M, McDonnell J, Vader JPP, Kahan JPP, Loo M (2001) The RAND/UCLA appropriateness Method User's Manual [Internet]. Transformation
- Kong H, West S (2013) WMA declaration of Helsinki—ethical principles for scientific requirements and research protocols 29–32
- Goh AC, Goldfarb DW, Sander JC, Miles BJ, Dunkin BJ (2012) Global evaluative assessment of robotic skills: validation of a clinical assessment tool to measure robotic surgical skills. *J Urol* 187(1):247–252
- Husslein H, Shirreff L, Shore EM, Lefebvre GG, Grantcharov TP (2015) The generic error rating tool: a novel approach to assessment of performance and surgical education in gynecologic laparoscopy. *J Surg Educ* 72(6):1259–1265