

Associations between energy demands, physical activity, and body composition in adult humans between 18 and 96 y of age.

Citation for published version (APA):

Speakman, J. R., & Westerterp, K. R. (2010). Associations between energy demands, physical activity, and body composition in adult humans between 18 and 96 y of age. *American Journal of Clinical Nutrition*, 92(4), 826-834. <https://doi.org/10.3945/ajcn.2009.28540>

Document status and date:

Published: 01/01/2010

DOI:

[10.3945/ajcn.2009.28540](https://doi.org/10.3945/ajcn.2009.28540)

Document Version:

Publisher's PDF, also known as Version of record

Document license:

Taverne

Please check the document version of this publication:

- A submitted manuscript is the version of the article upon submission and before peer-review. There can be important differences between the submitted version and the official published version of record. People interested in the research are advised to contact the author for the final version of the publication, or visit the DOI to the publisher's website.
- The final author version and the galley proof are versions of the publication after peer review.
- The final published version features the final layout of the paper including the volume, issue and page numbers.

[Link to publication](#)

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal.

If the publication is distributed under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license above, please follow below link for the End User Agreement:

www.umlib.nl/taverne-license

Take down policy

If you believe that this document breaches copyright please contact us at:

repository@maastrichtuniversity.nl

providing details and we will investigate your claim.

Associations between energy demands, physical activity, and body composition in adult humans between 18 and 96 y of age^{1–3}

John R Speakman and Klaas R Westerterp

ABSTRACT

Background: Associations between body composition and the energy expended on basal metabolism and activity are complex and age dependent.

Objective: The objective was to examine associations between body composition and daily (DEE), basal (BEE), and activity energy expenditure (AEE) throughout the adult life span.

Design: A cross-sectional study was conducted in 529 adults aged 18–96 y. DEE was measured by using doubly labeled water, BEE by using respirometry, and body composition by isotope dilution. AEE was calculated as DEE – BEE, and physical activity level (PAL) was calculated as DEE/BEE.

Results: Up to age 52 y, fat-free mass (FFM) and fat mass (FM) were positively associated with age in men, but no significant effect was observed in women. No effects of age on DEE and AEE were observed. The average DEE in men (14.1 MJ/d) was 27% greater than that in women (10.7 MJ/d). PAL averaged 1.84 in men and 1.75 in women. Above and including the age of 52 y, FFM, FM, DEE, BEE, and AEE were all negatively associated with greater age. The effect of age on AEE was greater than on BEE; consequently, PAL by the age of 95 y was only 1.36. PAL and AEE were both unrelated to FFM (both age adjusted).

Conclusions: PAL and AEE were not associated with age in subjects aged <52 y. AEE, BEE, and PAL were all negatively associated with age in subjects aged ≥52 y. An absence of a relation between age-adjusted PAL and FFM suggested that greater physical activity was not associated with higher FFM in the elderly. *Am J Clin Nutr* 2010;92:826–34.

INTRODUCTION

The interrelations between energy expenditure and body composition during adulthood are complex and appear to vary with life stage and with sex. During the first half of adulthood, individuals gradually accumulate body fat (1–4), but the extent of accumulation differs widely. Low energy expenditure, due either to low basal energy demands (5–7) or to low physical activity energy expenditure (AEE) (8, 9), have both been suggested to contribute to the energy imbalance that influences the extent of this increasing fat deposition. However, other studies have disputed these associations with basal (10) and AEE (5, 11, 12). Basal energy expenditure (BEE), however, clearly depends on the amount of metabolizing tissue (13), with independent effects of fat-free mass (FFM) and fat mass (FM) observed in many studies (14–23). Age has also been suggested to have a direct effect on tissue basal metabolic rate (15, 24–28), and consequently, the

actual change in BEE with age is a balance between this direct negative effect and an increase due to the expanding quantity of respiring tissue.

In contrast, in older individuals, total body mass tends to decline (29). In particular, the skeletal muscle component of FFM may be progressively lost (sarcopenia), notably in males (30). In later adulthood this decline in skeletal muscle mass is one of the key factors leading to immobility (31). Combined with the age-related decline in basal tissue metabolism, the loss of FFM leads to a large decline in BEE as individuals age. Physical activity level (PALs) also decline as individuals get older (32). However, whether this decline in activity is driven by the progression of sarcopenia or whether physical activity can reduce the rate of skeletal tissue loss is unknown. The causes of sarcopenia are complex and multifactorial (33). Strong evidence indicates that resistance training can stem the extent of skeletal muscle loss (34–37), but the role of general physical activity as a protective measure is controversial (38–42). Our aim in the present study, therefore, was to explore associations between PALs (by using the gold standard doubly labeled water method to assess energy expended on physical activity) and body composition. More generally, we aimed to define energy requirements across the life span. Energy is a key nutrient, and understanding patterns of energy requirements with age is of fundamental importance for nutritional professionals.

We used a large cross-sectional sample of >500 adults aged 18–96 y to explore the relations between energy demands and body composition across adulthood. Daily energy expenditure (DEE) was measured by using the doubly labeled water method, BEE by indirect calorimetry, and body composition by isotope dilution. Activity energy demands were captured directly from the difference between total and basal expenditures.

¹ From the Aberdeen Centre for Energy Regulation and Obesity, Institute of Biological and Environmental Sciences, University of Aberdeen, Aberdeen, Scotland (JRS), and the Department of Human Biology, Maastricht University, Maastricht, Netherlands (KRW).

² The original studies that contributed data to this analysis were funded by various sources acknowledged in the original manuscript. No additional funding was required to perform this work.

³ Address correspondence to JR Speakman, Institute of Biological and Environmental Sciences, University of Aberdeen, Aberdeen, Scotland, United Kingdom. E-mail: j.speakman@abdn.ac.uk.

Received August 14, 2009. Accepted for publication July 23, 2010.

First published online September 1, 2010; doi: 10.3945/ajcn.2009.28540.

SUBJECTS AND METHODS

Between 1983 and 2005, 529 subjects (289 men and 240 women) underwent measurements in Maastricht, Netherlands (first publication 1988), with both the DLW method and direct estimates of their BEE. This sample excluded individuals aged <18 y, undergoing interventions involving energy intake, undergoing physical activity including athletic performance, and those who were pregnant, lactating, or diseased (Table 1). For studies involving dietary or exercise interventions, only the preintervention data or those for control groups were used. Because data were collected over a prolonged time period during which there has been a profound change in obesity levels there is the potential for the date of the study to act as a confounding factor. However, during the time period in question, obesity levels increased modestly in the Netherlands (from 5% to 9%), and the age of the subjects was not biased in relation to the dates of the studies. Moreover, we previously showed no effect of the year of study on DEE (11). The population of the Netherlands has a slightly higher level of physical activity (measured as the PAL, *see* below) than other Western societies, such as the United States (11), but otherwise is a representative population.

Estimates of DEE, BEE, AEE, PAL, and body composition were derived from doubly labeled water studies. Doubly labeled water is an isotope-based technique in which subjects are dosed with labels of heavy oxygen and heavy hydrogen, and their differential elimination is a measure of their DEE (43). The method has a long history of application to the study of human energy demands (44). BEE was measured with indirect calorimetry. DEE was measured over 2-wk periods according to the Maastricht protocol (45). In brief, isotopes were administered orally as a mixture of $^2\text{H}_2\text{O}$ (99.9 atom%) and H_2^{18}O (10 atom%), which resulted in an initial excess body water enrichment of ≈ 150 ppm for deuterium and 300 ppm for oxygen-18. This dose leaves a sufficient excess enrichment at the end of the 14-d observation period. The dose volume was typically 80–160 mL,

depending on body mass. The isotopes were administered as the last consumption before bedtime. Subjects provided a background body water sample (blood or urine) immediately before isotope consumption (46; method D). Equilibration took place overnight, and the first postdosing sample was collected in the early morning, in case of urine from the second voiding. Additional duplicate samples (second daily void) were collected on days 8 and 14 after dosing.

PAL was derived by expressing DEE as a multiple of BEE (PAL = DEE/BEE). Alternatively, AEE was calculated as $(0.9 \times \text{DEE}) - \text{BEE}$, assuming that diet-induced energy expenditure is one-tenth of DEE in subjects in whom intake matches expenditure (47). FFM was derived from isotope dilution, assuming a 73% hydration of FFM (48), where dilution spaces were corrected for exchange of the label with nonaqueous substances in the body by dividing the dilution space of ^{18}O and ^2H by 1.01 and 1.04, respectively (49).

All analyses were performed separately for women ($n = 228$) and men ($n = 275$). Differences in FFM between subjects were adjusted for differences in FM by calculating the residual of the regression of FM on FFM, where fat mass was calculated as body mass minus FFM. To answer the question of whether age-related differences in FFM were related to PAL, the residual of the regression of PAL on age was analyzed as a function of the residual of FFM on fat mass and age. Data were tested for consistency to normality before analysis (Shapiro-Wilk test). When data were not normally distributed, they were transformed by log conversion to normalize the data distributions. Data analysis refers to transformed data, but the original data values are shown in the figures for clarity. When data were transformed, the nature of the transformation is mentioned at the appropriate location in the results. Data were considered significant at $P < 0.05$.

Visual inspection of the data showed that, in all cases, there was a transition point at the age of ≈ 50 y. Before 50 y, traits tended to be constant or higher with increasing age, whereas

TABLE 1
Subject characteristics by age decade¹

Age	No. of subjects	BMI	FFM	FM	FM	BEE	AEE	DEE	PAL
Men		<i>kg/m²</i>	<i>kg</i>	<i>kg</i>	<i>%</i>	<i>MJ/d</i>	<i>MJ/d</i>	<i>MJ/d</i>	
18–29 y	62	22.4 ± 3.4	60.0 ± 7.7	13.6 ± 7.2	17.9 ± 6.0	7.5 ± 1.0	5.1 ± 1.8	14.0 ± 2.5	1.88 ± 0.24
30–39 y	71	27.1 ± 7.0	63.7 ± 9.6	24.4 ± 17.6	25.6 ± 8.9	7.8 ± 1.4	4.7 ± 1.9	14.0 ± 2.8	1.79 ± 0.25
40–49 y	58	28.1 ± 5.7	63.2 ± 7.0	24.8 ± 12.9	26.7 ± 8.0	7.6 ± 1.2	5.2 ± 1.8	14.2 ± 2.6	1.88 ± 0.25
50–59 y	23	29.6 ± 4.5	65.7 ± 7.8	28.8 ± 11.1	29.6 ± 6.0	7.8 ± 0.8	5.3 ± 2.3	14.5 ± 2.7	1.88 ± 0.34
60–69 y	23	26.1 ± 4.7	56.3 ± 7.7	22.6 ± 9.4	27.7 ± 6.3	7.0 ± 0.9	3.6 ± 1.7	11.7 ± 2.2	1.69 ± 0.30
70–79 y	39	25.9 ± 2.6	53.3 ± 5.6	24.4 ± 6.0	31.1 ± 5.4	6.9 ± 1.1	2.8 ± 1.3	10.7 ± 1.7	1.57 ± 0.23
80–89 y	39	26.2 ± 2.8	49.9 ± 4.2	24.5 ± 6.0	32.6 ± 4.8	6.5 ± 0.6	1.6 ± 1.0	9.0 ± 1.7	1.37 ± 0.14
90–99 y	7	23.6 ± 3.4	46.3 ± 2.6	21.2 ± 8.8	30.5 ± 8.8	6.0 ± 0.8	1.3 ± 0.8	8.0 ± 0.6	1.36 ± 0.17
Women									
18–29 y	83	23.5 ± 5.6	46.1 ± 5.7	20.7 ± 12.5	29.1 ± 9.3	6.1 ± 0.8	3.5 ± 1.1	10.7 ± 1.7	1.75 ± 0.20
30–39 y	51	27.3 ± 6.9	46.6 ± 6.8	27.9 ± 13.7	35.2 ± 10.2	6.2 ± 1.1	3.1 ± 1.0	10.4 ± 1.8	1.68 ± 0.19
40–49 y	32	29.2 ± 8.2	50.9 ± 8.8	30.5 ± 17.4	35.0 ± 10.8	6.4 ± 1.2	3.6 ± 1.2	11.2 ± 2.3	1.71 ± 0.16
50–59 y	19	28.5 ± 7.4	45.2 ± 5.3	32.8 ± 14.3	40.7 ± 6.8	5.9 ± 0.7	2.9 ± 0.7	9.8 ± 1.3	1.65 ± 0.11
60–69 y	24	25.7 ± 2.5	42.4 ± 4.7	24.7 ± 5.6	36.5 ± 5.0	5.6 ± 0.7	2.7 ± 0.9	9.2 ± 0.9	1.67 ± 0.24
70–79 y	6	23.6 ± 3.4	39.0 ± 3.3	22.5 ± 6.5	36.2 ± 7.9	5.2 ± 0.3	2.4 ± 1.0	8.4 ± 1.2	1.63 ± 0.21
80–89 y	9	22.1 ± 3.2	36.8 ± 4.7	18.5 ± 3.8	33.3 ± 5.1	5.0 ± 0.5	0.6 ± 0.4	6.2 ± 0.8	1.23 ± 0.08
90–99 y	10	23.3 ± 4.4	37.0 ± 5.1	21.0 ± 9.7	34.6 ± 9.9	4.9 ± 0.6	1.1 ± 0.7	6.7 ± 1.2	1.36 ± 0.13

¹ All values are means ± SDs. FFM, fat-free mass; FM, fat mass; BEE, basal energy expenditure; AEE, activity energy expenditure; DEE, daily energy expenditure; PAL, physical activity level (ie, DEE/BEE).

they tended to be lower above the approximate age of 50 y with increasing age. The exact location of the transition appeared to vary with the trait under consideration and between the sexes. We explored the position of this transition using segmented regression (50) and the publically available computer program SEGREG (www.waterlog.info/segreg.htm). This program identified the location and significance of the breakpoint in the data relative to the fit of a simple linear regression model. We subsequently split the data at the appropriate breakpoints and performed the analyses for each trait separately on younger and older cohorts. All procedures described in this manuscript were reviewed by the University of Maastricht Ethical review board prior to their use.

RESULTS

Body-composition and energy expenditure characteristics of the study population stratified by age and sex are presented in Table 1. There was a strong difference in the effect of age on these variables at different times of life. Visual examination indicated that there appeared to be transitions in the data for all traits around the age of 40–60 y. We identified the exact position of these transitional breakpoints in the data by using segmented regression analysis (Table 2) on the data separated by sex. In all cases, the incorporation of a breakpoint in the analysis significantly improved the fit of the data above a simple linear regression with age as the predictor (Table 2). However, the position of the breakpoints varied significantly between traits and also between the sexes. With respect to body composition, the breakpoint for FFM in men (57.8 y; SE = 0.7) was significantly later than for women (46.8 y; SE = 0.45), whereas the breakpoint for FM did not differ significantly between the sexes (women: 47.8 y, SE = 3.55; men: 54.2 y, SE = 5.3). With respect to the BEE and DEE, the breakpoints were also at significantly greater ages in men than in women (Table 2). For AEE and PAL, no significant sex effect was observed on the position of the breakpoint. We performed regression analyses of the effect of

subject age on the various variables, splitting each trait into younger and older cohorts for both men and women (Table 3) using the optimal breakpoints from the segmented regression analysis to identify where to split the data. This analysis showed that all the traits were either constant or greater with subject age up to the breakpoints, except PAL in women, which was significantly lower with increasing subject age; however, for subjects above the breakpoint, age was negatively associated with all the traits except FM in men.

The breakpoints had an average age of 54.6 y for men and 48.9 y for women (paired $t_5 = 1.44$, $P = 0.21$). To facilitate comparisons between the sexes we took the average of these breakpoints (52 y), and the analyses were performed separately for a younger cohort of subjects aged 18–52 y ($n = 166$ women and 185 men) and for an older cohort of subjects aged ≥ 52.1 y ($n = 74$ women and 104 men).

Effects in subjects aged <52 y

Up to the age of 52 y, a large sex-related difference in body composition and energy expenditure was observed. On average, women younger than 52 y had a mean (\pm SD) FFM of 47.2 ± 7.0 kg; in men, FFM was 30% greater and averaged 62.5 ± 8.4 kg ($P < 0.001$). Both FFM and FM were significantly positively associated with subject age between 18 and 52 y in women (least-squares regression: \log_e converted data to normalize: $F_{1, 164} = 10.42$, $P = 0.002$ and $F_{1, 164} = 17.59$, $P < 0.001$, respectively) and in men (least-squares regression: \log_e converted data to normalize: $F_{1, 183} = 11.3$, $P < 0.001$ and $F_{1, 183} = 47.8$, $P < 0.001$, respectively, Figure 1). Because both FFM and FM were higher with increasing age, body mass index (in kg/m^2) was also positively related to subject age (Table 1). At age 45 y, almost half the subjects were obese (body mass index > 30). It is well established that greater FM is associated with elevated FFM. When FM was adjusted for the levels of FFM, there was no longer any trend with subject age between 18 and 52 y in women ($F_{1, 164} = 1.79$, $P = 0.18$), but there was

TABLE 2
Characteristics of segmented regression analysis on all traits¹

Trait	BP	BPse	F	P	r ² (lin)	r ² (seg)
Men						
FFM	57.8	0.68	71.9	<0.001	0.133	0.306
FM	54.2	5.3	28.4	<0.001	0.038	0.125
BEE	57.8	0.98	27.1	<0.001	0.069	0.139
DEE	53.1	0.99	209.7	<0.001	0.229	0.319
AEE	52.3	1.1	24.4	<0.001	0.214	0.275
PAL	52.3	1.27	15.5	<0.001	0.173	0.214
Women						
FFM	46.8	0.45	31.6	<0.001	0.082	0.189
FM	47.8	3.55	32.3	<0.001	0.023	0.123
BEE	39.8	2.14	19.6	<0.001	0.099	0.168
DEE	43.7	1.57	18.2	<0.001	0.272	0.324
AEE	54.6	1.44	8.91	<0.01	0.269	0.296
PAL	60.9	3.7	5.67	<0.02	0.214	0.231

¹ BP, optimal breakpoint age (in y); BPse, SE of optimal breakpoint; F, test statistic for significance, including breakpoint into regression analysis (df = 1, 287 for men and 1, 238 for women); r²(lin), coefficient of determination for the simple linear regression; r²(seg), coefficient of determination for the segmented regression; FFM, fat-free mass; FM, fat mass; BEE, basal energy expenditure; AEE, activity energy expenditure; DEE, daily energy expenditure; PAL, physical activity level.

TABLE 3

Parameters of regression fits for various traits in relation to age (y) below and above the optimal breakpoint (< or >BP) identified by segmented regression analysis¹

	Regression equation	df	F	P
Men				
FFM (kg)				
<BP	FFM = 54.99 + 0.222Age	1, 198	16.0	<0.001
>BP	FFM = 78.83 - 0.349Age	1, 87	28.1	<0.001
FM (kg)				
<BP	FM = 1.58 + 0.587Age	1, 186	27.8	<0.001
>BP	FM = 28.62 - 0.07Age	1, 79	0.94	0.335
BEE (MJ/d)				
<BP	BEE = 7.108 + 0.016Age	1, 198	4.04	0.046
>BP	BEE = 9.463 - 0.036Age	1, 87	11.9	<0.001
DEE MJ/d				
<BP	DEE = 13.33 + 0.022Age	1, 185	1.08	0.299
>BP	DEE = 22.89 - 0.165Age	1, 100	87.5	<0.001
AEE (MJ/d)				
<BP	AEE = 4.85 + 0.005Age	1, 183	0.12	0.727
>BP	AEE = 10.37 - 0.102Age	1, 102	55.9	<0.001
PAL				
<BP	PAL = 1.86 - 0.0004Age	1, 183	0.04	0.84
>BP	PAL = 2.53 - 0.013Age	1, 102	29.3	<0.001
Women				
FFM (kg)				
<BP	FFM = 40.07 + 0.236Age	1, 160	13.1	<0.001
>BP	FFM = 51.52 - 0.139Age	1, 76	8.1	0.006
FM (kg)				
<BP	FM = 6.68 + 0.589Age	1, 161	19.8	<0.001
>BP	FM = 42.44 - 0.251Age	1, 75	10.8	0.002
BEE (MJ/d)				
<BP	BEE = 5.505 + 0.022Age	1, 132	3.16	0.078
>BP	BEE = 7.607 - 0.029Age	1, 104	38.1	<0.001
DEE (MJ/d)				
<BP	DEE = 10.45 + 0.009Age	1, 150	0.20	0.655
>BP	DEE = 14.56 - 0.086Age	1, 86	95.75	<0.001
AEE (MJ/d)				
<BP	AEE = 3.94 - 0.015Age	1, 168	2.13	0.146
>BP	AEE = 6.13 - 0.056Age	1, 68	62.82	<0.001
PAL				
<BP	PAL = 1.86 - 0.004Age	1, 187	9.05	0.003
>BP	PAL = 2.43 - 0.012Age	1, 49	28.1	<0.001

¹ The actual BPs are shown in Table 2. FFM, fat-free mass; FM, fat mass; BEE, basal energy expenditure; AEE, activity energy expenditure; DEE, daily energy expenditure; PAL, physical activity level.

a significant positive relation in men ($F_{1, 183} = 8.04, P = 0.005$) (Figure 2). Percentage body fat in the youngest age group (18–29) averaged 17.9% in men and 29.1% in women and in the oldest cohort under 52 y (40–52) averaged 26.8% and 34.7% in men and women, respectively.

The mean (\pm SD) total DEE in men younger than 52 y was 14.1 ± 2.6 MJ/d, whereas DEE was significantly lower in women by 27% (10.7 ± 1.9 MJ/d). This difference of 3.4 MJ/d in DEE was contributed to by significantly greater basal and AEE in the men. Hence, BEE was 6.2 ± 1.0 MJ/d in the women and 7.6 ± 1.2 MJ/d in the men (analysis on log-converted data: $t = 13.2, P < 0.001$)—an average difference of 1.4 MJ/d (40% of the overall difference in DEE). The difference in AEE between men and women was even greater. Hence, women expended an average of 3.4 ± 1.1 MJ/d in AEE compared with 5.0 ± 1.8 MJ/d in men (comparison on log-converted data: $t_{321} = 9.46, P < 0.001$) (Figure 3). This difference contributed to 60% of the overall difference in DEE between the sexes.

At the individual level, BEE was significantly positively related to FFM (general linear model with \log_e FFM, \log_e FM and age entered as covariates to predict \log_e converted BEE: $F_{1, 348} = 1015, P < 0.001$), independently to \log_e FM ($F_{1, 348} = 87.4, P < 0.001$), and negatively related to subject age ($F_{1, 348} = 20.7, P < 0.01$). The least-squares fit multiple regression explained 80.1% of the total variation in \log_e BEE. Once these effects had been accounted for, there was still a small residual effect of sex ($F_{1, 350} = 8.7, P = 0.003$); women had a slightly lower BEE than anticipated from body composition. Because the oldest individuals in the cohort aged <52 y (ages 40–52 y) were fatter and heavier than the younger members of this cohort, the net effect of subject age on BEE was a marginally significant positive effect in women ($F_{1, 164} = 4.12, P = 0.044$) but no significant effect in men ($F_{1, 183} = 2.99, P = 0.085$; Figure 4). In contrast, AEE was independent of age (least-squares fit linear regression of \log_e AEE against age: $F_{1, 350} = 0.50, P = 0.54$; Figure 3). Consequently, there was no significant trend in PAL




FIGURE 1. Fat-free mass (FFM; A) and fat mass (FM; B) as a function of age in women ($n = 240$; gray squares) and men ($n = 289$; black circles). Lines are fitted regressions for each sex above and below an average break point of 52 y.

with age in men ($F_{1, 183} = 0.04$, $P = 0.84$). The distribution of PAL was consistent with normality by the Shapiro-Wilk test ($P > 0.15$) and was not transformed, but there was a slight but significantly lower PAL with age in women ($F_{1, 164} = 5.58$, $P = 0.19$; **Figure 5**). There was no significant relation between FFM (logged data adjusted for \log_e FM and age) and PAL (adjusted




FIGURE 2. The residual (res) of the regression of fat-free mass (ffm) on fat mass (FM) as a function of age for women ($n = 240$; gray squares) and men ($n = 289$; black circles). Lines are fitted regressions for each sex above and below an average break point of 52 y.




FIGURE 3. Activity energy expenditure (AEE) as a function of age in women ($n = 240$; gray squares) and men ($n = 289$; black circles). Lines are fitted regressions for each sex above and below an average break point of 52 y.

for age) in men aged 18–52 y ($F_{1, 183} = 3.47$, $P = 0.064$), but in women aged 18–52 y there was a statistically significant but very weak positive relation ($F_{1, 164} = 3.9$, $P = 0.05$, $r^2 = 0.023$) (**Figure 6A**). Individuals aged 18–52 y with greater FFM (independent of FM) were generally not more physically active.

Effects in subjects aged ≥ 52 y

Above and including the age of 52 y in both men and women, greater subject age was associated with lower FFM and FM (ie, decadal averages from 50 to 59 y onward show progressively lower values; Table 1, Figure 1). Moreover, the lower FFM in older subjects remained when it was adjusted for differences in \log_e FM (FFM distribution was consistent with normality, but FM was not and had to be log transformed: $F_{1, 175} = 25.1$, $P < 0.001$). The effect of age on adjusted FFM was different between sexes, being lower in women than in men. On average, FFM in the women aged ≥ 52 y was lower by 0.13 kg/y (95% CI: 0.05, -0.20 kg/y), whereas in men it was lower by an average of 0.42 kg/y (95% CI: -0.32 , -0.53 kg/y). In the very oldest individuals aged 90–100 y, the average sex difference in FFM was




FIGURE 4. Basal energy expenditure (BEE) as a function of age in women ($n = 240$; gray squares) and men ($n = 289$; black circles). Lines are fitted regressions for each sex above and below an average break point of 52 y.




FIGURE 5. Physical activity level (PAL) as a function of age in women ($n = 240$; gray squares) and men ($n = 289$; black circles). Lines are fitted regressions for each sex above and below an average break point of 52 y.

<10 kg, compared with a 15-kg difference in subjects aged 18–52 y (Table 1).

Above and including the age of 52 y, all distributions of the measures of energy expenditure (BEE, AEE, DEE, and PAL) were consistent with normality, except for AEE and PAL in men, which were log transformed to normalize them. All of these traits were age dependent and were lower in the older subjects (Figures 3–6). The lower BEE with increasing subject age in subjects aged ≥ 52 y (Figure 4) was similar for women ($-22 \text{ kJ} \cdot \text{d}^{-1} \cdot \text{y}^{-1}$; 95% CI: $-15, -38 \text{ kJ} \cdot \text{d}^{-1} \cdot \text{y}^{-1}$) and men ($-45 \text{ kJ} \cdot \text{d}^{-1} \cdot \text{y}^{-1}$; 95% CI: $-26, -60 \text{ kJ} \cdot \text{d}^{-1} \cdot \text{y}^{-1}$). As established for individuals aged < 52 y, BEE was significantly related to FFM and \log_e FM, both of which were negatively related to subject age in subjects aged ≥ 52 y (Table 1 and Figure 1B). When we included FFM and \log_e FM as predictors of BEE in the multiple regression model, the effect of subject age disappeared (FFM: $F_{1, 173} = 368, P < 0.001$; \log_e FM: $F_{1, 173} = 7.28, P = 0.008$; age: $F_{1, 173} = 0.12, P = 0.735$). The lower BEE in older subjects in this age group, therefore, appeared solely due to older subjects having a lower FM and FFM.

AEE was also significantly lower in older subjects (Figure 3). AEE was not normally distributed and was therefore was log converted to normalize it before analysis. There was a strong effect of a subject’s age on \log_e AEE ($F_{1, 173} = 124, P < 0.001$) but no effect of sex ($F_{1, 173} = 0.12, P = 0.75$) and no sex-by-age interaction ($F_{1, 173} = 0.21, P = 0.64$) (Figure 4); hence, the effect of age was not significantly different between men and women. Because the lower AEE in older subjects was much greater than the effect of subject age on BEE, PAL was strongly dependent on age for those subjects aged ≥ 52 y (Figure 5). On average the PAL between ages 18 and 52 y averaged 1.84 in men and 1.75 in women. In the subjects aged 90–100 y, PAL averaged only 1.36 in both sexes (Table 1). The sex difference in the effect of subject age on PAL after age 52 y for women (-0.010 ; 95% CI: $-0.007, -0.014$) and men (-0.013 ; 95% CI: $-0.008, -0.017$) was not significant.

Above and including the age of 52 y, both FFM (adjusted for \log_e FM) and PAL were generally lower in older individuals, although an age effect was less noticeable early in this period, particularly in men. It was necessary, therefore, to statistically remove this age effect before seeking a relation between adjusted FFM and PAL. In effect, this analysis asked whether there





FIGURE 6. Residual of the regression of physical activity level (PAL) on age as a function of the residual of the regression of fat-free mass (FFM) on age and fat mass in women (gray squares) and men (black circles). The upper plot is for adults aged < 52 y, and the lower plot is for adults aged ≥ 52 y.

is a relation between adjusted FFM and PAL, independent of the overall effect of age on these variables. There was no significant relation between age-adjusted FFM (also adjusted for FM) and age-adjusted PAL (Figure 6B) in men ($F_{1, 102} = 1.06, P = 0.306$) or in women ($F_{1, 72} = 0.01, P = 0.901$). Consequently, individuals in this older age cohort with greater FFM for their age and body fatness were not more physically active. Alternatively greater physical activity at any particular age was not associated with higher FFM.

DISCUSSION

Transition points

All the traits measured showed transition points around the middle of the age range. Before the transitions, the traits tended to be independent of subject age or increased in older subjects; after

the transition, all the traits tended to be reduced in individuals that were older. These trends were consistent with many previous studies (references for individual traits follow). The age of the breakpoints did not differ significantly between men and women and occurred at a mean (\pm SE) age of 52 ± 1.8 y ($n = 12$). In cross-sectional data, such as in the current study, little significance can be drawn from the different breakpoints for the different traits.

Effects of age on body-composition differences

Up to the age of 52 y in this cohort, greater subject age was associated with elevated FM and FFM in men but not in women. The elevation of FM was not just associated with elevated FFM, because there was a significant independent effect of age on the adjusted FM. Our data are consistent with many previous studies that have found age-related increases in FM in younger men (1–3, 51, 52), although other studies have reported no change (53). Most previous studies have also established similar age-related effects in women, and the absence of an effect in our cohort was therefore unexpected. Subjects who were older than 60 y tended to have lower FFM and FM. This pattern was also observed previously (1–3), but, again was not universal—some studies reporting progressive accumulation of FM beyond the age of 50 y (54). In this cohort, the effect of subject age was much greater in men than in women. A greater prevalence of sarcopenia in older men was also previously indicated (30, 55).

Effect of age on BEE

Up to the age of 52 y, there was a significant positive effect of FFM and an independent contribution of FM, which, combined with a small negative effect of age, explained 81% of the total variation in BEE. This pattern is consistent with many other studies that have explored the effect of body composition and age on BEE (12, 13, 15–19, 21–23), although some studies have failed to find an independent effect of FM beyond that of FFM (56–60). Several previous studies have suggested that the large sex effect on BEE is only due to the large difference in size and body composition between women and men (12, 22, 61, 62), whereas our data were consistent with other data that have suggested a persistent sex effect after body composition is accounted for (63, 64). The reasons for the differences between these studies are not immediately apparent. The lower BEE of older subjects in the age range 50–100 y has been reported many times (15, 25, 26, 65–67) although contradictory studies showing no age effect have also been published (68). There has been much dispute over the contribution of changes in body composition to this effect (69), with some studies consistent with the current study, suggesting that body-composition changes can explain the reduced BEE or lack of change, but others suggesting it cannot (66). The present data set suggested no additional age effect beyond that caused by subject differences in FFM and FM. Differences between studies may reflect variation in the causes of differences in FFM between studies. It is well established that FFM is heterogeneous in composition and comprises tissues with widely differing metabolic rates (70, 71) and that this variability may contribute to wider scaling effects of metabolic rate (72–74). Recent studies attempting to account for this heterogeneity have suggested that there may be an additional physiologic component to the age-related differences in BEE (19, 75–77).

Effects of subject age on AEE and PAL

There were no effects of subject age on either AEE or PAL, up to the age of 52 y. Previous studies have not reported significant age effects on these variables in this age group, which suggests that it is generally the case that physical activity is sustained until subjects are in their 50s. Thereafter, there was a profound negative effect of subject age on both AEE and PAL. This pattern is consistent with that previously observed (24, 24, 78, 79). Blanc et al (62) noted that women in their 80s had a much lower AEE than did men but that PAL was not different, consistent with our observations in the older subjects.

The data presented here are cross-sectional. Some longitudinal studies have been performed in older adults, and their results generally contrast with those found in the current study. For example, Rothenberg et al (67) found no changes in AEE and PAL in a longitudinal comparison between ages 73 and 78 y. Moreover, Reilly et al (80) also found no differences in PAL over time in a cohort studied between the ages of 65 and 72 y. However, in both of these cases the sample sizes were small and the duration of follow-up short. Studies of larger cohorts with longer follow-up, however, are much harder to perform, hence their absence in the literature. It seems likely that the cross-sectional nature of our sample would actually tend to minimize the effect of age on PAL, because in all age classes we selected participants who were free-living and able-bodied. As subjects age they become increasingly likely to suffer from a lack of mobility, and such subjects would have a low PAL and a low probability of being included in these studies.

The lower levels of physical activity in older subjects who also had a lower FFM has led to the suggestion that these phenomena are causally linked (39, 81–85), either because elevated physical activity protects against age-related FFM loss or because preserved FFM loss allows greater physical activity. The data presented here do not support this interpretation because, once the confounding influence of age had been removed, we found no association between FFM and PAL (Figure 6B). This analysis is consistent with previous studies based on a smaller sample size (66) and studies measuring physical activity by accelerometry (42) or by questionnaire rather than directly as expended energy on exercise (38, 40, 41). Consistent with our correlational findings, other studies have shown that physical activity interventions may alter muscle function and increase fat loss, but do not halt the progress of muscle mass loss in the elderly (86, 87). Interestingly, Hunter et al (36) found that resistance training could reverse the decline in FFM in the elderly, and that the increase in FFM was associated with an increase in DEE. This was due in part to an increase in BEE because of the greater FFM. However, there was also a suggestion of an increase in AEE, but this was not significant ($P = 0.06$). Whether this was a type II error because of the low sample size (8 men and 7 women) remains uncertain. Generally, however, it is now regarded that lower physical activity is due more to changes in muscle composition and physiology than to quantity (88, 89) and that sarcopenia is caused by a complex interaction of nutritional factors, such as a protein intake, oxidative stress, inflammatory changes, and hormonal effects (84, 90–95).

In summary, our data show that up to the age of 52 y, fat and FFM were positively associated with subject age, whereas BEE, DEE, AEE, and hence PALs, were broadly independent of subject

age. After the age of 52 y, older subjects tended to have lower values for all these variables. This effect of subject age was most noticeable in FFM and also in AEE and PAL. Older subjects aged > 52 y had a lower FFM and lower levels of energy expenditure on physical activity. For a person at any given age (≥ 52 y), however, there was no association between the amount of energy expended on physical activity and FFM, which suggests that routine physical activity does not protect against loss of FFM, and that loss of FFM is not associated with lower routine PALs.

Four referees made valuable contributions to improving the manuscript.

The authors' responsibilities were as follows—KRW: involved in all of the studies that generated the original data, and JRS and KRW: analyzed the data and cowrote the manuscript. Neither author had any conflicts of interest to declare.

REFERENCES

- Ito H, Ohshima A, Ohto N, et al. Relation between body composition and age in healthy Japanese subjects. *Eur J Clin Nutr* 2001;55:462–70.
- Kyle UG, Genton L, Hans D, Karsgaard L, Slosman DO, Pichard C. Age-related differences in fat-free mass, skeletal muscle, body cell mass and fat mass between 18 and 94 years. *Eur J Clin Nutr* 2001;55:663–72.
- Kyle UG, Genton L, Slosman DO, Pichard C. Fat-free and fat mass percentiles in 5225 healthy subjects aged 15 to 98 years. *Nutrition* 2001;17:534–41.
- Larsson I, Forslund HB, Lindroos AK, et al. Body composition in the SOS (Swedish obese subjects) reference study. *Int J Obes Relat Metab Disord* 2004;28:1317–24.
- Tataranni PA, Harper IT, Snitker S, et al. Body weight gain in free-living Pima Indians: effect of energy intake vs expenditure. *Int J Obes Relat Metab Disord* 2003;27:1578–83.
- Ravussin E. Low resting metabolic rate as a risk factor for weight gain: role of the sympathetic nervous system. *Int J Obes Relat Metab Disord* 1995;19:S8–9.
- Ravussin E, Lillioja S, Knowler WC, et al. Reduced rate of energy expenditure as a risk factor for body-weight gain. *N Engl J Med* 1988;318:467–72.
- Esparza J, Fox C, Harper IT, et al. Daily energy expenditure in Mexican and USA Pima Indians: low physical activity as a possible cause of obesity. *Int J Obes Relat Metab Disord* 2000;24:55–9.
- Levine JA. Non-exercise, activity thermogenesis (NEAT). *Nutr Rev* 2004;62:S82–97.
- Flatt JP. Differences in basal energy expenditure and obesity. *Obesity (Silver Spring)* 2007;15:2546–8.
- Westertep KR, Speakman JR. Physical activity energy expenditure has not declined since the 1980s and matches energy expenditures of wild mammals. *Int J Obes Relat Metab Disord* 2008;32:1256–63.
- Klausen B, Toubro S, Astrup A. Age and sex effects on energy expenditure. *Am J Clin Nutr* 1997;65:895–907.
- Weinsier RL, Schutz Y, Bracco D. Reexamination of the relationship of resting metabolic rate to fat-free mass and to the metabolically active components of fat-free mass in humans. *Am J Clin Nutr* 1992;55:790–4.
- Cunningham JJ. A reanalysis of the factors influencing basal metabolic rate in normal adults. *Am J Clin Nutr* 1980;33:2372–4.
- Fukagawa NK, Bandini LG, Young JB. Effect of age on body composition and resting metabolic rate. *Am J Physiol* 1990;259:E233–8.
- Cunningham JJ. Body composition as a determinant of energy expenditure—a synthetic review and a proposed general prediction equation. *Am J Clin Nutr* 1991;54:963–9.
- Nelson KM, Weinsier RL, Long CL, Schutz Y. Prediction of resting energy expenditure from fat-free mass and fat mass. *Am J Clin Nutr* 1992;56:848–56.
- Karhunen L, Franssilakallunki A, Rissanen A, Kervinen K, Kesaniemi YA, Uusitupa M. Determinants of resting energy expenditure in obese non-diabetic Caucasian women. *Int J Obes Relat Metab Disord* 1997;21:197–202.
- Wang Z, Heshka S, Gallagher D, Boozer CN, Kotler DP, Heymsfield SB. Resting energy expenditure-fat-free mass relationship: new insights provided by body composition modeling. *Am J Physiol Endocrinol Metab* 2000;279:E539–45.
- Muller MJ, Bosy-Westphal A. Assessment of energy expenditure in children and adolescents. *Curr Opin Clin Nutr Metab Care* 2003;6:519–30.
- Krems C, Luhrmann PM, Strassburg A, Hartmann B, Neuhauser-Berthold M. Lower resting metabolic rate in the elderly may not be entirely due to changes in body composition. *Eur J Clin Nutr* 2005;59:255–62.
- Johnstone AM, Murison SD, Duncan JS, Rance KA, Speakman JR. Factors influencing variation in basal metabolic rate include fat-free mass, fat mass, age, and circulating thyroxine but not sex, circulating leptin, or triiodothyronine. *Am J Clin Nutr* 2005;82:941–8.
- Ganpule AA, Tanaka S, Ishikawa-Takata K, Tabata I. Interindividual variability in sleeping metabolic rate in Japanese subjects. *Eur J Clin Nutr* 2007;61:1256–61.
- Roberts SB, Dallal GE. Effects of age on energy balance. *Am J Clin Nutr* 1998;68:975S–9S.
- Vaughan L, Zurlo F, Ravussin E. Aging and energy expenditure. *Am J Clin Nutr* 1991;53:821–5.
- Piers LS, Soares MJ, McCormack LM, O'Dea K. Is there evidence for an age-related reduction in metabolic rate? *J Appl Physiol* 1998;85:2196–204.
- Johnstone AM, Rance KA, Murison SD, Duncan JS, Speakman JR. Additional anthropometric measures may improve the predictability of basal metabolic rate in adult subjects. *Eur J Clin Nutr* 2006;60:1437–44.
- Rothenberg EM, Bosaeus IG, Westertep KR, Steen BC. Resting energy expenditure, activity energy expenditure and total energy expenditure at age 91–96 years. *Br J Nutr* 2000;84:319–24.
- Newman AB, Lee JS, Visser M, et al. Weight change and the conservation of lean mass in old age: the Health, Aging and Body Composition Study. *Am J Clin Nutr* 2005;82:872–8.
- Iannuzzi-Sucich M, Prestwood KM, Kenny AM. Prevalence of sarcopenia and predictors of skeletal muscle mass in healthy, older men and women. *J Gerontol A Biol Sci Med Sci* 2002;57:M772–7.
- Fantin F, Di Francesco V, Fontana G, et al. Longitudinal body composition changes in old men and women: interrelationships with worsening disability. *J Gerontol A Biol Sci Med Sci* 2007;62:1375–81.
- Harris AM, Lanningham-Foster LM, McCrady SK, Levine JA. Non-exercise movement in elderly compared with young people. *Am J Physiol Endocrinol Metab* 2007;292:E1207–12.
- Lee SK, Choi HJ, Kim YC, et al. Basal metabolic rate is related to lumbar bone mineral density and body composition in pre- and post-menopausal women at Korea. *J Bone Miner Res* 2004;19:S306.
- Adamo ML, Farrar RP. Resistance training, and IGF involvement in the maintenance of muscle mass during the aging process. *Ageing Res Rev* 2006;5:310–31.
- Bamman MM, Clarke MSF, Feeback DL, Lieberman SA, Stevens BR, Greenisen MG. Resistance exercise prevents sarcopenia during bed rest unloading. *FASEB J* 1997;11:326.
- Hunter GR, Wetzstein CJ, Fields DA, Brown A, Bamman MM. Resistance training increases total energy expenditure and free-living physical activity in older adults. *J Appl Physiol* 2000;89:977–84.
- Hunter GR, McCarthy JP, Bamman MM. Effects of resistance training on older adults. *Sports Med* 2004;34:329–48.
- Hughes VA, Frontera WR, Roubenoff R, Evans WJ, Singh MAF. Longitudinal changes in body composition in older men and women: role of body weight change and physical activity. *Am J Clin Nutr* 2002;76:473–81.
- Roubenoff R, Hughes VA, Dallal GE, et al. The effect of gender and body composition method on the apparent decline in lean mass-adjusted resting metabolic rate with age. *J Gerontol A Biol Sci Med Sci* 2000;55:M757–60.
- Castillo EM, Goodman-Gruen D, Kritiz-Silverstein D, Morton DJ, Wingard DL, Barrett-Connor E. Sarcopenia in elderly men and women—the Rancho Bernardo Study. *Am J Prev Med* 2003;25:226–31.
- Raguso CA, Kyle U, Kossovsky MP, et al. A 3-year longitudinal study on body composition changes in the elderly: role of physical exercise. *Clin Nutr* 2006;25:573–80.
- Starling RD. Energy expenditure and aging: effects of physical activity. *Int J Sport Nutr Exerc Metab* 2001;11:S208–17.
- Speakman JR. Doubly-labelled water: theory and practice. New York, NY: Kluwer Academic Publishers, 1997.
- Speakman JR. The history and theory of the doubly labeled water technique. *Am J Clin Nutr* 1998;68:932S–8S.

45. Westerterp KR, Wouters L, Lichtenbelt WDV. The Maastricht protocol for the measurement of body-composition and energy expenditure with labeled water. *Obes Res* 1995;3:49–57.
46. Speakman JR, Racey PA. The equilibrium concentration of O-18 in body-water—implications for the accuracy of the doubly-labeled water technique and a potential new method of measuring RQ in free-living animals. *J Theor Biol* 1987;127:79–95.
47. Westerterp KR, Plasqui G. Physical activity and human energy expenditure. *Curr Opin Clin Nutr Metab Care* 2004;7:607–13.
48. Pace N, Rathbun EN. Studies on body composition, III: The body water and chemically combined nitrogen content in relation to fat content. *J Biol Chem* 1945;158:685–91.
49. Schoeller DA, Vansanten E. Measurement of energy expenditure in humans by doubly labeled water method. *J Appl Physiol* 1982;53:955–9.
50. Oosterbaan RJ, Sharma DP, Singh KN, Rao KVGK. Crop production and soil salinity: evaluation of field data from India by segmented linear regression with breakpoint. Proceedings of the symposium on land drainage for salinity control in arid and semi-arid regions. 1990;3:373–83.
51. Jankowska EA, Rogucka E, Medras M, Welon Z. Relationships between age-related changes of sex steroids, obesity and body fat distribution among healthy Polish males. *Med Sci Monit* 2000;6:1159–64.
52. Heitmann BL. Body fat in the adult Danish population aged 35-65 years: an epidemiologic study. *Int J Obes* 1991;15:535–45.
53. Atlantis E, Martin SA, Haren MT, Taylor AW, Wittert GA. Lifestyle factors associated with age-related differences in body composition: the Florey Adelaide Male Aging Study. *Am J Clin Nutr* 2008;88:95–104.
54. Coin A, Sergi G, Miniucuci N, et al. Fat-free mass and fat mass reference values by dual-energy X-ray absorptiometry (DEXA) in a 20-80 year-old Italian population. *Clin Nutr* 2008;27:87–94.
55. Bunout D, de la Maza MP, Barrera G, Leiva L, Gattas V, Hirsch S. Assessment of sarcopenia: longitudinal versus cross sectional body composition data. *Aging Clin Exp Res* 2007;19:295–9.
56. Luke A, Rotimi CN, Adeyemo AA, et al. Comparability of resting energy expenditure in Nigerians and US blacks. *Obes Res* 2000;8:351–9.
57. Dugas LR, Ebersole K, Schoeller D, et al. Very low levels of energy expenditure among pre-adolescent Mexican-American girls. *Int J Pediatr Obes* 2008;3:123–6.
58. Bogardus C, Lillioja S, Ravussin E, et al. Familial dependence of the resting metabolic rate. *N Engl J Med* 1986;315:96–100.
59. Segal KR, Pi-Sunyer FX, Nyman AM, Brehm BA, Gutin B. Energy expenditure of lean and obese men of similar body weight. *Fed Proc* 1985;44:1282.
60. Ferraro R, Ravussin E. Fat mass in predicting resting metabolic rate. *Am J Clin Nutr* 1992;56:460.
61. Cunningham JJ. Body composition and resting metabolic-rate: the myth of feminine metabolism. *Am J Clin Nutr* 1982;36:721–6.
62. Blanc S, Schoeller DA, Bauer D, et al. Energy requirements in the eighth decade of life. *Am J Clin Nutr* 2004;79:303–10.
63. Ferraro R, Lillioja S, Fontvieille AM, Rising R, Bogardus C, Ravussin E. Lower sedentary metabolic rate in women compared with men. *J Clin Invest* 1992;90:780–4.
64. Muller MJ, Bosity-Westphal A, Klaus S, et al. World Health Organization equations have shortcomings for predicting resting energy expenditure in persons from a modern, affluent population: generation of a new reference standard from a retrospective analysis of a German database of resting energy expenditure. *Am J Clin Nutr* 2004;80:1379–90.
65. Roberts SB, Fuss P, Heyman MB, Young VR. Influence of age on energy requirements. *Am J Clin Nutr* 1995;62:S1053–8.
66. Pannemans DLE, Westerterp KR. Energy expenditure, physical activity and basal metabolic rate of elderly subjects. *Br J Nutr* 1995;73:571–81.
67. Rothenberg EM, Bosaeus IG, Steen BC. Energy expenditure at age 73 and 78—a five year follow-up. *Acta Diabetol* 2003;40:S134–8.
68. Murray LA, Reilly JJ, Choudhry M, Durnin JVGA. A longitudinal study of changes in body composition and basal metabolism in physically active elderly men. *Eur J Appl Physiol Occup Physiol* 1996;72:215–8.
69. Poehlman ET. Energy expenditure and requirements in aging humans. *J Nutr* 1992;122:2057–65.
70. Elia M. Organ and tissue contribution to metabolic rate. In: Kinney JM, Tucker HN, eds. *Energy metabolism: tissue determinants and cellular corollaries*. New York, NY: Raven, 1992:61–80.
71. Krebs HA. Body size and tissue respiration. *Biochim Biophys Acta* 1950;4:249–69.
72. Heymsfield SB, Gallagher D, Kotler DP, Wang ZM, Allison DB, Heshka S. Body-size dependence of resting energy expenditure can be attributed to nonenergetic homogeneity of fat-free mass. *Am J Physiol Endocrinol Metab* 2002;282:E132–8.
73. Heymsfield SB, Gallagher D, Wang ZM. Body composition modeling—application to exploration of the resting energy expenditure fat-free mass relationship. 2000.
74. Pietrobello A, Heymsfield SB, Gallagher D, et al. Body size dependence of resting energy expenditure can be attributed to non-energetic homogeneity of fat-free mass. *Obes Res* 2001;9:79S.
75. Gallagher D, He Q, Albu J, et al. Loss of high metabolic rate organ mass does not explain resting metabolic rate decrease with age. *Obes Res* 2005;13:A6.
76. Gallagher D, Allen A, Wang ZM, Heymsfield SB, Krasnow N. Smaller organ tissue mass in the elderly fails to explain lower resting metabolic rate. *Ann NY Acad Sci* 2000;904:449–55.
77. Gallagher D, Belmonte D, Deurenberg P, et al. Organ-tissue mass measurement allows modeling of REE and metabolically active tissue mass. *Am J Physiol* 1998;275:E249–58.
78. Roberts SB, Rosenberg I. Nutrition and aging: changes in the regulation of energy metabolism with aging. *Physiol Rev* 2006;86:651–67.
79. Roberts SB, Dallal GE. Energy requirements and aging. *Public Health Nutr* 2005;8:1028–36.
80. Reilly JJ, Lord A, Bunker VW, et al. Energy balance in healthy elderly women. *Br J Nutr* 1993;69:21–7.
81. Kamel HK. Sarcopenia and aging. *Nutr Rev* 2003;61:157–67.
82. Kamel HK, Maas D, Duthie EH. Role of hormones in the pathogenesis and management of sarcopenia. *Drugs Aging* 2002;19:865–77.
83. Marcell TJ. Sarcopenia: causes, consequences, and preventions. *J Gerontol A Biol Sci Med Sci* 2003;58:M911–6.
84. Roubenoff R, Hughes VA. Sarcopenia: current concepts. *J Gerontol A Biol Sci Med Sci* 2000;55:M716–24.
85. Baumgartner RN, Waters DL, Gallagher D, Morley JE, Garry PJ. Predictors of skeletal muscle mass in elderly men and women. *Mech Ageing Dev* 1999;107:123–36.
86. Goodpaster BH, Chomentowski P, Ward BK, et al. Effects of physical activity on strength and skeletal muscle fat infiltration in older adults: a randomized controlled trial. *J Appl Physiol* 2008;105:1498–503.
87. Puggaard L, Larsen JB, Ebbesen E, Jeune B. Body composition in 85 year-old women: effects of increased physical activity. *Aging (Milano)* 1999;11:307–15.
88. Hollmann W, Strueder HK, Tagarakis CVM, King G. Physical activity and the elderly. *Eur J Cardiovasc Prev Rehabil* 2007;14:730–9.
89. Stenholm S, Harris TB, Rantanen T, Visser M, Kritchevsky SB, Ferrucci L. Sarcopenic obesity: definition, cause and consequences. *Curr Opin Clin Nutr Metab Care* 2008;11:693–700.
90. Lee CE, McArdle A, Griffiths RD. The role of hormones, cytokines and heat shock proteins during age-related muscle loss. *Clin Nutr* 2007;26:524–34.
91. Chaput JP, Lord C, Cloutier M, et al. Relationship between antioxidant intakes and class I sarcopenia in elderly men and women. *J Nutr Health Aging* 2007;11:363–9.
92. Di Iorio A, Abate M, Di Renzo D, et al. Sarcopenia: age-related skeletal muscle changes from determinants to physical disability. *Int J Immunopathol Pharmacol* 2006;19:703–19.
93. Stookey JD, Adair LS, Popkin BM. Do protein and energy intakes explain long-term changes in body composition? *J Nutr Health Aging* 2005;9:5–17.
94. Evans WJ, Campbell WW. Sarcopenia and age-related-changes in body composition and functional capacity. *J Nutr* 1993;123:465–8.
95. Lynch GS, Ryall JG. Role of beta-adrenoceptor signaling in skeletal muscle: implications for muscle wasting and disease. *Physiol Rev* 2008;88:729–67.