

# Bring the Noise!

Citation for published version (APA):

Pereira Da Cruz Gomes Lourenco, M. (2021). *Bring the Noise! Fear, assessment and treatment of tinnitus*. [Doctoral Thesis, Maastricht University, KU Leuven]. Ridderprint. <https://doi.org/10.26481/dis.20211029mp>

**Document status and date:**

Published: 01/01/2021

**DOI:**

[10.26481/dis.20211029mp](https://doi.org/10.26481/dis.20211029mp)

**Document Version:**

Publisher's PDF, also known as Version of record

**Please check the document version of this publication:**

- A submitted manuscript is the version of the article upon submission and before peer-review. There can be important differences between the submitted version and the official published version of record. People interested in the research are advised to contact the author for the final version of the publication, or visit the DOI to the publisher's website.
- The final author version and the galley proof are versions of the publication after peer review.
- The final published version features the final layout of the paper including the volume, issue and page numbers.

[Link to publication](#)

**General rights**

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal.

If the publication is distributed under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license above, please follow below link for the End User Agreement:

[www.umlib.nl/taverne-license](http://www.umlib.nl/taverne-license)


**Take down policy**

If you believe that this document breaches copyright please contact us at:

[repository@maastrichtuniversity.nl](mailto:repository@maastrichtuniversity.nl)

providing details and we will investigate your claim.

**B  
R  
I  
N  
G  
  
T  
H  
E  
  
N  
O  
I  
S  
E**



Fear, assessment and treatment of tinnitus  
by **Matheus P. C. G. Lourenco**

# **BRING THE NOISE!**

Fear, assessment and treatment of tinnitus

Matheus Pereira da Cruz Gomes Lourenco

ISBN: 978-94-6416-838-9  
Cover design: Ruth Poppe  
Lay-out: Publiss | [www.publiss.nl](http://www.publiss.nl)  
Print: Ridderprint | [www.ridderprint.nl](http://www.ridderprint.nl)

© Copyright 2021: Matheus Pereira da Cruz Gomes Lourenco

All rights reserved. No part of this publication may be reproduced, stored in a retrieval system, or transmitted in any form or by any means, electronic, mechanical, by photocopying, recording, or otherwise, without the prior written permission of the author.

BRING THE NOISE!

Fear, assessment and treatment of tinnitus

DISSERTATION

to obtain the joint degree of Doctor at the Maastricht University and Doctor of Psychology at KU Leuven, on the authority of the Rector Magnificus, Prof. dr. Rianne M. Letschert and the Rector Prof.dr. Luc Sels in accordance with the decision of the Board of Deans, to be defended in public on October 29th, 2021

by

Matheus Pereira da Cruz Gomes Lourenco

*Supervisor(s):*

Prof. dr. J. W. S. Vlaeyen

Dr. R. F. F. Cima

Assessment Committee:

Prof. dr. M. Peters (chair)

Prof. dr. I. van Diest, KU Leuven, Belgium

Dr. D. Hoare, Nottingham University, United Kingdom

Dr. W. Schlee, University of Regensburg, Germany

This project has received funding from the European Union's Horizon 2020 research and innovation programme under the Marie Skłodowska-Curie grant agreement number 722046.

À Geni





## *Table of contents*

<b>Chapter 1</b>	General introduction	3
<b>Chapter 2</b>	Fear in the ear	31
<b>Chapter 3</b>	Effects of EMA-induced monitoring on tinnitus experience	51
<b>Chapter 4</b>	The daily experience of subjective tinnitus	75
<b>Chapter 5</b>	Better together	95
<b>Chapter 6</b>	General discussion	123
<b>Impact statement</b>		138
<b>Acknowledgments</b>		145



# CHAPTER 1

---

General introduction

## From trees to tinnitus

*“If a tree were to fall on an island where there were no human beings would there be any sound?”* – The Chautauquan (1883)

The answer provided to the infamous question postulated by The Chautauquan was *no*. Sound is the transformation of vibrations that travel through a medium (e.g. air) into coherent neurological signals, and consequently cannot exist outside an interpreter of sound. The corollary that may follow – *“If a human being were to hear a tree fall, would there be any tree?”* – is at the heart of this dissertation as we seek to better understand the experience of phantom sounds, specifically tinnitus.

### *What is tinnitus?*

Current tinnitus definitions (e.g. Baguley et al., 2013; Cima, 2018; Pawel J. Jastreboff et al., 1994; Langguth, 2011; Table 1) converge in specifying a lack of an acoustic source while sound is perceived. In other words, authors agree that *tinnitus* is the perception of a sound without a corresponding acoustic origin. Different from the experience of other phantom acoustic perceptions such as voices or music, tinnitus is limited to sounds without explicit semantic meaning (e.g. tones, hissing, or chirping). Transient tinnitus experiences may subside within seconds, minutes or days (i.e. acute tinnitus), though for some its experience, whether continuous or intermittent, becomes chronic. Persistent tinnitus that does not spontaneously remit over a significant period of time (considered to be at least 3 months) may be classified as chronic tinnitus (Fuller, 2021). Chronic tinnitus may take a toll in the life of those who experience it. Sleep disturbance, concentration difficulties, suicidal thoughts, depressive symptomatology, anxiety, anger, avoidance environments perceived as too silent or loud, and decreased quality of life are some of the reported effects of tinnitus (Hall et al., 2018). Such chronic tinnitus that is associated with emotional reactivity and related disability can be further classified as Chronic Disabling Tinnitus (Fuller, 2021). The current work then utilizes the definition of Chronic Disabling Tinnitus as the unremitted perception of a sound (for at least 3 months) that is without semantic meaning nor corresponding acoustic origin, which produces significant emotional reactivity and related disability.

**Table 1:** Example of tinnitus definitions

Definition	Author
<i>“An auditory phantom perception, and therefore cannot be associated with any sound measurement”</i> (p. 216)	Jastreboff et al. (1994)
<i>“A common and distressing condition that is typically characterized by the perceived sensation of sound in the absence of an external stimulus”</i> (p. 1635)	Langguth (2011)
<i>“The conscious perception of an auditory sensation in the absence of a corresponding external stimulus”</i> (p. 1600)	Baguley et al. (2013)
<i>“The symptom itself, tinnitus aurium, can be defined as the phantom perception of continuous sound or noise in the absence of an external (or adequate) source”</i> (p. 369)	Cima (2018)
<i>“Tinnitus is the conscious awareness of a tonal or composite noise for which there is no identifiable corresponding external acoustic source, which becomes Tinnitus Disorder ‘when associated with emotional distress, cognitive dysfunction, and/or autonomic arousal, leading to behavioural changes and functional disability”</i> (p. 1)	De Ridder et al. (2021)

## *Epidemiology*

A review of 35 tinnitus prevalence studies found that estimates for tinnitus vary widely, between 5.1 to 42.7% (McCormack et al., 2016). Variability in estimates were in part attributed to differences in the geographical location of the study, population demographics, tinnitus assessment and heterogeneous reporting (Biswas & Hall, 2020; McCormack et al., 2016). Recently, a standardized tinnitus assessment was created to be used across Europe in an attempt to provide a more accurate picture of tinnitus epidemiology. Biswas et al. (2020) included and 11 427 participants representing 11 different languages across Europe. The prevalence was estimated to be 14.7%, ranging from 8.7% (Ireland) to 28.3% (Bulgaria), without differences between genders (Biswas et al., 2020). Higher tinnitus prevalence was confirmed with increased age and worsening hearing. The authors further investigated tinnitus severity (i.e. self-reported level of annoying, worrisome or bothersome tinnitus), finding it prevalent in 1% of the participants, ranging from 0.6% (Ireland) to 1.4% (Romania), with a difference between women (1.4%) and men (1%). Moreover, the role of lifestyle risk factors (e.g. alcohol consumption, smoking, obesity) have contradictory findings and are yet to be fully understood (Biswas & Hall, 2020). As with previous reports, current epidemiological findings in the tinnitus field are restricted to the geographical location in which the study is conducted (mostly in developed countries) as well as the assessment methods of tinnitus.

## *Assessing tinnitus*

The subjective tinnitus experience cannot be directly observed or measured, thus mostly relying on patient self-report assessments. As audiological features of the tinnitus percept (e.g. loudness, pitch, location) do not adequately explain tinnitus severity (Andersson, 2003), assessments methods evolved to include and reflect tinnitus related distress, disability, coping, attention and beliefs (Table 2).

**Table 2:** Tinnitus related self-report assessments

<b>Assessment</b>	<b>Author</b>	<b>Purpose</b>	<b>Sub-scales</b>	<b>Recall Timeframe</b>
<b>Tinnitus Questionnaire (TQ)</b>	Hallam (1988)	Measure psychological aspects of tinnitus complaints and distress	Emotional distress, cognitive distress, intrusiveness, auditory perceptual difficulties, sleep disturbance, somatic complaint.	One week.
<b>Tinnitus Handicap Questionnaire (THQ)</b>	Kuk et al. (1990)	Measure level of perceived tinnitus related handicap.	Physical, health, emotional status, social consequences hearing and communication, personal viewpoint.	No timeframe specified.
<b>Tinnitus Handicap Inventory (THI)</b>	Newman et al. (1996)	Measure level of perceived tinnitus severity.	Functional, emotional and catastrophic responses.	No timeframe specified.
<b>Tinnitus Coping Style Questionnaire (TCSQ)</b>	Budd and Pugh (1996)	Measure tinnitus related coping strategies and style.	Maladaptive coping, effective coping, and passive coping.	No time frame specified.
<b>Tinnitus Vigilance and Awareness Questionnaire (TVAQ)</b>	Cima et al. (2011)	Measure heightened attention towards tinnitus		No time frame specified.
<b>Fear of Tinnitus Questionnaire (FTQ)</b>	Cima et al. (2011)	Measure tinnitus related fear		No time frame specified.
<b>Tinnitus Functional Index (TFI)</b>	Meikle et al. (2012)	Measure tinnitus severity and treatment related change	Tinnitus intrusiveness, sense of control, quality of life, sleep, hearing, concentration, relaxation, and emotion.	One week.

Such a list of assessments contributes towards a comprehensive picture of those who experience tinnitus and shed light into the high heterogeneity of experience (e.g. Henry et al., 2012; Schlee et al., 2016). However, self-report assessments have limitations, which

hinder the accurate portrayal of tinnitus. High levels of reading difficulty in the tools have been reported to potentially affect tinnitus assessment (Atcherson et al., 2011). Furthermore, effects of the psychological state during assessment have been documented to influence responses (Belli et al., 2008; Brüggemann et al., 2016; Das et al., 2012; Langguth et al., 2011). More importantly, bias associated with memory reconstruction (i.e. recall bias) can be influenced by the setting of the assessment (e.g. hospital), the recency of the experience, and the averaging of experiences within longer (or unspecified) time frames (Shields et al., 2016; Stone & Shiffman, 1994).

### *Underlying cognitive and behavioral mechanisms of Chronic Disabling Tinnitus*

Despite the limitations, a comprehensive assessment of the tinnitus experience allows for insights into the underlying cognitive and behavioral mechanisms which may explain tinnitus disability. Whereas tinnitus triggers (e.g. excess earwax, increased stress, ototoxicity) and audiological features of the tinnitus percept (e.g. loudness, pitch, location) do not adequately predict disability (Andersson, 2003; Wallhäusser-Franke et al., 2017), the role of fear, attention and avoidance have been postulated to play a major role in the development and maintenance of Chronic Disabling Tinnitus (Table 3). A brief description of each model follows.

**Table 3:** Overview of tinnitus models

<b>Model</b>	<b>Fundamental assumptions</b>	<b>Core prediction</b>
<b>Habituation model (Hallam et al., 1984)</b>	Decrease response to the tinnitus perception (i.e. habituation) can be achieved through repeated exposure.	Habituation is disrupted due to heightened attention in moments of high physiological arousal (e.g. due to daily stressors).
<b>Neurophysiological model (Pawel J. Jastreboff et al., 1988; Pawel J. Jastreboff, 1990)</b>	Through classical conditioning the tinnitus perception is associated with aversive emotional responses.	Weakening the association between tinnitus perception and emotional responses is essential for habituation. Thus, extinction, the presentation of the conditioned stimulus without the unconditioned stimulus, results in decrease tinnitus disability and distress.
<b>Cognitive Behavioural Model (McKenna et al., 2014)</b>	Automatic evaluations of the tinnitus percept are at the core of dysfunctional responses.	Improvements in tinnitus distress and disability are achieved through altering the negative automatic thoughts on the tinnitus meaning and controllability.
<b>Fear Avoidance model of tinnitus (Cima, Crombez, et al., 2011; Kleinstäuber et al., 2013)</b>	Associative learning processes and catastrophic misinterpretations of the tinnitus percept are responsible for a downward spiral of emotional and behavioural reactions. Avoidance plays a pivotal role in the development and maintenance of distress and disability. Opposed to the downward spiral, a neutral or positive interpretation of the tinnitus percept is expected elicit a functional response.	Improvements are expected with the use of exposure therapy in order to facilitate extinction of the learned associations between fear and tinnitus.


### *The Habituation model*

The model specifies that crucial to Chronic Disabling Tinnitus is the habituation process (Hallam et al., 1984), more specifically, the failure to habituate to the tinnitus percept. Habituation is defined as the decrease in responses (e.g. physiological arousal) to a repeated or continuous exposure to the stimulus (e.g. tinnitus). Thus, habituation may be hindered when attention is continuously directed towards the percept due to affective learning associations. The model assumes that tinnitus features (e.g. pitch, intensity) are constant, and thus can be classified as a repeated or continuous stimulus. In reality, tinnitus features fluctuate (Cederroth et al., 2019) and may be experienced as new stimulus whenever changes are perceived, and hence not a repeated or continuous stimulus.



Avoidant behaviours were later incorporated into the model to help explain disability and decreased quality of life (e.g. avoidance of environments perceived as too silent or loud; Figure 1; Kröner-Herwig et al., 2003). The model further elaborates on the avoidant behaviours as the result of operant learning, which refers to the change in frequency of behaviour through positive or negative reinforcement or punishment (Skinner, 1938). In the context of the Habituation model, an individual may avoid situations where heightened physiological arousal (e.g. fear) is expected or experienced. The avoidance of these experiences (i.e. removal of negative stimulus) provides immediate relief (i.e. negative reinforcement). According to the model, reduction of physiological arousal before entering such environments is a necessary part in reducing tinnitus distress by weakening the negative association (Mckenna, 2004). It should be noted that strategies to decrease physiological arousal (e.g. relaxation techniques) may act as an avoidance strategy in itself, hindering the habituation process by not challenging the originally created association between tinnitus and heightened physiological arousal (i.e. fear).

**Figure 1:** The Habituation model




Note: adapted from Kröner-Herwig et al. (2003).

### *Neurophysiological model*

Building upon the Habituation model, the Neurophysiological model (Figure 2) further assumes that the habituation process is hindered due to an aversive emotional state being associated (i.e. conditioned) with the tinnitus percept (Pawel J. Jastreboff et al., 1988; Pawel J. Jastreboff, 1990). Central to the model is classical (i.e. Pavlovian) conditioning, where two previously unrelated stimuli are paired (Pavlov, 1927). More specifically, a neutral stimulus (Conditioned Stimulus; CS) is paired with a biologically relevant one (Unconditioned Stimulus; US) which elicits a response (Unconditioned Response; UR). Successful pairing results in responses (Conditioned Response; CR) to the CS in the absence of the US. Despite the model's reliance on classical conditioning paradigms, there is a lack of clarity on the model's proposed associations (Baguley et al., 2013). Moreover, the research which provides the theoretical underpinnings is based on animal models and thus focused on tinnitus perception generation and limited in the explanation of tinnitus disability and distress.

**Figure 2:** The Neurophysiological model




Note: adapted from Jastreboff (2011; p. 579).

Nonetheless, treatment based on the Neurophysiological model is aimed at appeasing the patients' emotional reactions through psycho-education while utilizing sound therapy to reduce tinnitus perception (Tinnitus Retraining Therapy; Pawel J. Jastreboff & Hazell, 2004). The use of sound therapy relies on the generation of sounds that mask the characteristics of the tinnitus percept while psycho-education is used to deconstruct any tinnitus-related fears. Limited evidence for the efficacy of this particular treatment exists (Cima et al., 2019). More specifically, masking of tinnitus perception may provide short-term relief, thus avoiding the feared stimulus and working as negative reinforcement. Furthermore, distress and disability potentially increase as patients may become dependent on sound generation to avoid tinnitus perception (McKenna & Irwin, 2008).

### *Cognitive Behavioural model*

Contrary to the Neurophysiological model, the Cognitive Behavioural (CB) model of tinnitus aims at explaining tinnitus distress through the cognitive process of appraisal (Figure 3; McKenna et al., 2014). According to Lazarus (1991) appraisal, can be divided into primary – appraisal of the causal attributions (e.g. what causes tinnitus) – and secondary – appraisal of controllability (e.g. what can it be done to reduce tinnitus). The dual appraisal model may then explain the negative tinnitus evaluation (e.g. tinnitus as a result of irreversible hearing damage), which in turn increases physiological arousal leading to active monitoring of and selective attention towards the tinnitus percept. Safety behaviours are the direct result of such appraisals. Beliefs that loud environments may further increase tinnitus severity may lead to the avoidance of such environments or to other coping strategies (e.g. ear plugs).

Consequently, the CB model justifies treatments that focus on tinnitus appraisal (e.g. through CBT) and reducing physiological arousal (e.g. relaxation, mindfulness). Some indirect support for the model may be inferred from the efficacy of treatments, such as mindfulness based cognitive therapy (McKenna et al., 2017) in reducing tinnitus related distress. However, studies to test the model itself are lacking with only limited evidence for the separate components (McKenna et al., 2014).

**Figure 3:** The Cognitive Behavioural model

Note: adapted from McKenna (2014).

### *Fear Avoidance model*

As the name suggests, fear – through catastrophic misinterpretations of tinnitus (e.g. indication of brain damage) – is the hinge which pivots patients into a pathological spiral instead of a path to recovery (Cima, Crombez, et al., 2011; Kleinstäuber et al., 2013). The role of fear in the development and maintenance of chronic disability has been supported in the field of chronic pain (e.g. Meulders, 2020), from where the model originates (Lethem et al., 1983; Vlaeyen & Linton, 2000, 2012). Parallels between pain and tinnitus have been stipulated for decades (e.g. Møller, 1997). A starting point between similarities stems from the observation that the chronic experiences from both fields are characterized by the lack of identifiable source and (further) physical harm (D. De Ridder et al., 2011; Møller, 1997, 2007). Furthermore, severity of chronic pain and tinnitus are marked by oversensitivity to specific stimuli, such as light touch and soft sounds respectively (Møller, 1997). More interestingly, chronic experiences of pain and tinnitus are not necessarily

associated with distress and decreased quality of life. Fear, attention and avoidance contribute to the pathological expression of these experiences. Successful treatment for both chronic experiences are similar, relying on Cognitive Behavioural Therapy (CBT) with a highlighted role of exposure during treatment (e.g. Fuller et al., 2020; Vlaeyen et al., 2012). The model is further elaborated on in the next.


## From fear to freedom

*Or in the night, imagining some fear,*

*How easy is a bush supposed a bear!* – Shakespeare (1605, 5.1.1)

The FA model postulates that the pathological cycle is triggered through negative misinterpretations of tinnitus. As in the name of the model, *fear* is thought to play a pivotal role in the development of Chronic Disabling Tinnitus. Tinnitus-related fear, increased attention and relentless monitoring follow. The result is defensive mechanisms, such as the avoidance of stimuli that are deemed threatening, which in turn increases disability, distress, and decreased quality of life (Figure 4). Avoidance (i.e. an adaptive behavioural response to fear; Watson & Rayner, 1920), also highlighted in previous models, takes a fundamental role in the development of disability as individuals may avoid common situations in daily life. Environments perceived to have higher levels of sound (e.g. restaurants, bars, social gatherings, movie theatres) are avoided due to the perceived threat of potential tinnitus worsening. On the other hand, moments in which the tinnitus perception may be more salient due to silence (e.g. evening and night times), are also avoided and maladaptive strategies may be employed (e.g. masking of the tinnitus percept), affecting not only the one who perceives tinnitus, but those close to them (Hall et al., 2018).

Alternatively, tinnitus may be perceived as a relatively harmless experience. In this case, no detrimental cycle is initiated, and the individual may continue with a normal, unaltered daily life. The crucial moment of tinnitus experience in which the path to recovery or pathological cycle diverge is postulated to be dependent on the learned association between tinnitus and fear.

**Figure 4:** Fear Avoidance model (replicated from Cima et al., 2018)

Fear is an emotional response resulting from the perception of threat to health or safety. Fear serves as an adaptive response by motivating animals to avoid harm. Examples are easily observed from mice to men: fear of heights, predators, or the dark. While some fears may be innate (e.g. fear of snakes), others may be learned (e.g. fear of flying). Learned fear holds evolutionary benefits as a harmful experience (e.g. pain resulting from a snake bite), produces anticipatory defensive behaviours (e.g. avoiding snakes). Some events may not be learned fast enough from a first-hand experience (e.g. a poisonous snake bite may end one's life). As such, indirect learning without undergoing the experience (e.g. through observation or verbal communication) provides an added benefit in the evolutionary process.

The process of creating (or re-creating) associations through observation (i.e. social learning; Bandura et al., 1967) adds yet another layer to the process of fear acquisition. Overall, humans may acquire fear from a variety of sources, such as personal experiences (i.e. classical or operant conditioning), observation and verbal instructions derived from cultural norms. Beliefs (e.g. tinnitus may signal a brain tumour vs. tinnitus as benign) can then be passed on and may become ingrained in a population. Such a rich learning environment and innate capacity to learn fear occasionally leads to maladaptive fear responses to unharmed and unthreatening events or experiences (e.g. fear of flying). Sustained maladaptive fear associations and responses may lead to the development of pathological cycles that reverberate through one's life (as illustrated in the FA model).

Biologically salient negative experiences, such as pain, are powerful motivators to seek safety and trigger avoidance behaviours. More specifically, pain may quickly create fear of stimuli which then are avoided due to potential harm. Pain is a naturally feared stimulus, and biologically imperative in the evolutionary process. The universality of pain led to comprehensive development of associate learning paradigms which robustly support the FA model. While the evidence on the relationship between fear and pain is rich in the field of chronic pain (e.g. Vlaeyen & Crombez, 2020), the development of fundamental research to support the FA model in the tinnitus field is in its infancy. Thus, the fundamental research of the FA model was based on fear of pain. Currently there is no evidence that tinnitus is as naturally as feared as pain, hampering the adaptation of the FA model. The origins of Chronic Disabling Tinnitus fear are not yet clear, though it has been suggested by the different models to rely on learned associations processes (i.e. classical conditioning, operant conditioning, social learning).

As in the Neurophysiological model, classical conditioning is postulated to be one integral mechanism in the development of Chronic Disabling Tinnitus. More specifically, fear conditioning, which relies on the pairing between a neutral (CS) and a biologically salient (US) stimulus which elicits a fear response (UR). A powerful US elicits an innate UR. In a successful association, the contingent pairings of CS and US will result in a motivational change of the CS, which activates the representation of the US, and hence starts eliciting an anticipatory conditioned response (CR) in the absence of the US. An example may be drawn from the field of chronic pain where a normally innocuous movement (proprioceptive CS; e.g. bending over) has occurred with a pain (US), leading to fear of the bending over. This type of conditioning can be and is reliability re-created in lab setting with healthy participants, where fear of movement is established, and avoidance of the movement is measured (for a review on fear conditioning in the context of pain see Meulders, 2020). Moreover, the experience of the feared CS (e.g. bending over) without the expected negative outcome (e.g. pain), provides the opportunity for the formation of new CS – no-pain-US associations, thereby challenging dysfunctional expectations and weakening the previously made association (i.e. extinction).

An unanswered question is whether we can simply substitute *tinnitus* for *pain* and establish learned fear of tinnitus sounds in a similar way. Currently, paradigms in tinnitus experimental studies are limited to animal models (e.g. Brozoski et al., 2012; Pawel J. Jastreboff et al., 1988). Animal studies have supported the correlations between hearing damage (i.e. loss of outer hair functioning) and tinnitus, as it may trigger

related hyperactivity (i.e. in the dorsal cochlear nucleus; Kaltenbach et al., 2002). Such paradigms, however, are severely limited in the understanding of tinnitus distress and disability. Moreover, even establishing that animals have acquired tinnitus perception is challenging and restricted by behaviour experiments. The clear role of tinnitus in classical conditioning models (e.g. US, CR) remains unclear and it is not known if the experience of tinnitus is as naturally feared as pain (and thus a possible US). Furthermore, the creation of tinnitus in humans within lab conditions remains a challenge as it is not known if the experience of a constant and even loud tone may lead to reliable fear conditioning.

### *Treating tinnitus*

Currently, a common cure for tinnitus perception does not exist, and since tinnitus does not automatically translate into suffering (as the majority of individuals with chronic tinnitus do not suffer from Chronic Disabling Tinnitus), a cure, albeit desirable, may not be necessary. However, Cognitive Behaviour Therapy (CBT) is highly recommended for the treatment of tinnitus (Cima et al., 2019), with a recent Cochrane review reporting potential benefits of CBT on reducing the negative impact of tinnitus on quality of life (Fuller et al., 2020).

With a long tradition, CBT aims at the complex relationship between behaviour, cognition and emotion (Beck, 1979; Ellis & Grieger, 1977). CBT is an umbrella term that includes therapeutic approaches stemming from both behaviour, and cognitive therapies. Learned associations are challenged through behavioural techniques (e.g. exposure), while cognitive techniques modify the relationship between thought (e.g. catastrophizing) and emotion (e.g. fear). By combining cognitive and behavioural methods, CBT encompasses various psychological intervention techniques that, in the case of tinnitus, are aimed at reducing the impact of tinnitus rather than altering tinnitus features (i.e. loudness). Altering cognitions and responses by targeting different steps of the pathological cycle (i.e. catastrophizing, fear, avoidance) can be achieved through the delivery of different techniques (e.g. psychoeducation, exposure) making CBT a flexible and adaptive treatment option. The variety in techniques allows for tailored treatment approaches, resulting in unique combinations of techniques in function of the patient, therapist and setting differences. On the other hand, increased variability decreases replicability in research and treatment, with CBT treatment protocols potentially differing significantly. Thus, the role that each potential variation within treatment delivery (e.g. techniques, setting, length) plays is yet to be fully understood.



## *From bench to bedside to bench*

Given the current state of tinnitus research and treatment we have identified three points of interest to further develop the field. They fall within the topics of (1) testing of a tinnitus fear conditioning paradigm in humans; (2) improving tinnitus assessment; and (3) the investigation of CBT components in order to better understand what works best for whom. We will subsequently expand on each point and draw the objectives of this dissertation.

### *1. Developing a tinnitus fear conditioning paradigm for human participants*

In order to develop a comprehensive understanding of tinnitus development, maintenance and treatment, the systematic investigation of the role of tinnitus-related fear seems promising. Fundamental knowledge on the underlying learning mechanisms of tinnitus-related fear is still in its infancy. The main obstacle when adapting the FA model from pain to tinnitus is that the stimuli (pain and tinnitus) may translate differently into fundamental paradigms. Pain is considered a naturally aversive and negative experience whereas the experience of tinnitus may not as easily or universally trigger immediate alarm as pain might. Pain, when considered a representation of a serious (i.e. threatening) bodily harm (e.g. nerve damage), may act as a US, therefore creating a functional CS – US association. The implication for tinnitus research is that tinnitus itself may not be perceived as threatening nor used to consistently trigger an aversive response (US). Like pain, tinnitus threat must be associated with a perception of serious (potential) harm (e.g. damaged hearing, brain tumour). Therefore, a fear learning paradigm for tinnitus may be adapted as follows: tinnitus, conceptualized as a perception of a neutral sound (CS), acquires fear responses (CR) experienced due its association with an aversive (threatening) event (US). In other words, a neutral sound becomes a signal for a threat to well-being.

Considering fear associations in tinnitus research, an ecologically relevant US is necessary. Due to (1) the report of reduced sound tolerance on those suffering from Chronic Disabling Tinnitus (Baguley, 2003), (2) the avoidance of loud sounds due to the fear of triggering or worsening tinnitus perception (Kleinstäuber et al., 2013), and (3) the common reporting of tinnitus after experiencing of loud sounds (Gilles et al., 2013), loud sounds may be considered threatening (e.g. causing hearing damage and increased tinnitus) and thus provide a viable US. Consequently, a potentially replicable fear learning paradigm may be achieved in the tinnitus field.

*Objective:*

Adapt/create an associative learning model for the field of tinnitus in order to establish a replicable fear learning paradigm with healthy human participants.

*2. (Re)Assessing tinnitus*

Assessing tinnitus has proven a challenge, with audiological measures of tinnitus pitch, type, or intensity not translating into tinnitus distress (Andersson, 2003; Figueiredo et al., 2010; Henry, 2016; Rabau et al., 2015). Such measures do not provide insight into the toll that Chronic Disabling Tinnitus may take on one's life (e.g. social isolation, annoyance, fear). Instead, patient self-report must not be seen as the *next best thing*, but the most adequate assessment of Chronic Disabling Tinnitus. In other words, it is not specific tinnitus *features* that define the pathology but the entirety of the tinnitus *experience*, following a patient centered, rather than disease centered, approach to research and treatment.

Research in the field of chronic pain, in which a lack of direct measure has also been debated, has produced substantial arguments for the use of self-report tools as a valid technique for pain assessment (Robinson et al., 2013). We are then left with properly providing the patient with an accurate method to measure each individual's unique tinnitus experience. Hence the use of standardized self-report questionnaires to operationalize tinnitus. Many have been developed and improved upon, such as the Tinnitus Questionnaire (TQ; Hallam et al., 1988), Tinnitus Functional Inventory (TFI; Meikle et al., 2012) the Tinnitus Handicap Inventory (THI; Newman et al., 1996, 1998) and the Tinnitus Disability Index (TDI; Cima, Vlaeyen, et al., 2011). Independently from each conceptualization and objective, these tools are susceptible to the same issues of commonly used self-report questionnaires: recall bias, reading difficulty, and current psychological state.

Novel methodological approaches, namely Ecological Momentary Assessments (EMA) and End-of-Day Diaries (EDD), increase ecological validity and reduce the influence of recall bias by: (1) shortening the time between experience and assessment and (2) eliminating the artificial settings of self-report assessments (e.g. hospitals, laboratories, clinics). While EDD relies on once-a-day administration of questionnaires, EMA aims to reach information that is still in working memory by prompting participants with short and simple questions during their daily life. Previously, such methodologies were

implemented at painstakingly efforts and costs, but the development of affordable technology to a wider population (i.e. smartphones) made these methodologies more accessible for researchers and clinicians alike, providing an increasingly used alternative in the tinnitus field (e.g. Gerull et al., 2019; Goldberg et al., 2017; Schlee et al., 2016; Timmer et al., 2018).

EMA is an attractive tool in investigating tinnitus, with the added benefit to understand fluctuations during the day and further decreasing the risk of recall bias when compared to EDD. These potential benefits come at a cost since it relies on participants to remain in possession of their smartphone at all times. Participants must allow themselves to be interrupted in the midst of activities and research on the detrimental value of increased screen time is not lacking. Higher levels of stress, anxiety, depression (Elhai et al., 2017; Vahedi & Saipho, 2018) and lower wellbeing (Horwood & Anglim, 2019) have been associated with smartphone use. Furthermore, simply answering to an EMA prompt may elicit a cascade of events when reminders of unanswered emails, messages or tasks are readily available and highlighted by smartphones. Beyond that, increased awareness to negative experiences (i.e. tinnitus for patients suffering from Chronic Disabling Tinnitus) has been thought to negatively impact well-being as well. Conversely, EDD's minimize the burden of screen time while also potentially sacrificing recall bias and ecological validity when compared to EMA (Schneider & Stone, 2016). EDD's have been an established methodology for decades (e.g. Verbrugge, 1980), though its use in tinnitus research is scarce and a direct comparison between the tools in the field of tinnitus is lacking.

The development of technology and increased availability of smartphones has made novel assessment methods (EDD and EMA) viable alternatives to questionnaire and interview methods of measuring tinnitus. Nevertheless, these methodologies are yet to be fully tested within the field of tinnitus and must be better understood before being applied to wider use. We identified two main areas of focus in regard to this area. First, following one of the main principals of healthcare – *Primum non nocere* (*first, do no harm*) – we seek to explore the possible negative effects that EMA may have on well-being and tinnitus experience by filling the gap of previous studies on the matter. Second, we seek to compare EMA and EDD methodologies in the hope of making a substantiated recommendation for the use of either methodology.

*Objectives:*

Investigate whether EMA negatively affects tinnitus experience.

&

Investigate whether EMA provides a more accurate measure of tinnitus experience when compared to diary assessments.

*3. CBT for tinnitus: what works for whom?*

Finding a cure for tinnitus has challenged the field, with a lack of consensus on tinnitus definition, treatment and resource allocation (McFerran et al., 2019). Despite lack of consensus, the literature has repeatedly shown that, even without a common cure, meaningful improvement in tinnitus distress and quality of life can be achieved through the use of CBT (Fuller et al., 2020). In tinnitus, CBT aims at breaking, altering or recreating learned associations between tinnitus and maladaptive responses (e.g. avoidance) as well as identifying and modifying cognitive interpretations in the tinnitus experience (e.g. fear).

The variety of CBT techniques and combinations is an issue that does not only pertain to the tinnitus field. In an attempt to increase reproducibility of interventions, Michie et al. (2013) established a taxonomy that expands to 93 possible behavioural change techniques. Even with such detailed framework, authors of the recent Cochrane review (Fuller et al., 2020) point to the lack of availability of detailed protocols used in CBT interventions. Furthermore, mode of delivery (e.g. face-to-face, videocall), length and frequency of sessions, setting (e.g. hospital, clinic), delivery agent (e.g. therapist, chat-robot) and unit of delivery (i.e. group or individual) create an infinite number of possible CBT protocols that preclude the consideration of equivalent treatment. Thus, whereas CBT has been shown to be an added value in tinnitus treatment, lack of replicable protocols hinders the capacity for research to understand what works best and for whom, preventing more tailored approaches to be designed.

Tailoring of treatments can be accomplished in different degrees, such as the stepped care in the CBT approach used by Cima et al. (2012), which delivers care as needed (by steps) according to severity (increasing in treatment complexity). More tailored approaches can be reached through stratification or matching (Linton et al., 2018). Stratification, different from stepped care, classifies patients according to severity levels from the first visit. This approach curbs the assumption made by stepped care, in which most patients

will recover with little to no treatment (i.e. filtering out the less severe cases at each step). Further tailoring can be achieved through matched care. Expanding on the severity-based stratification, matched care includes individual risk factors which are then the focus of treatment.

It must be clarified that *tailored* treatment does not necessarily translate to *individual* treatment. Individual treatment simply refers to receiving a treatment in an individual setting (i.e. alone). Therefore, an individual treatment is not, by definition, tailored and vice versa, with tailored treatment sometimes being delivered in group settings. Beyond potentially reducing the costs of treatment delivery, group treatment may allow for social learning processes to be used as a powerful agent of change, particularly within CBT. Group-based CBT has the added benefit of changing behaviour, cognitions and emotions through increased availability of social learning cues (Bandura et al., 1967). These cues, such as observing another patient being exposed to the feared stimulus (e.g. loud noise, tinnitus) without the expected negative outcome (e.g. anxiety, loss of control) challenges previously learned pathological associations. Moreover, observing someone else expose themselves to a feared stimulus (i.e. vicarious extinction of fear; Rachman, 1977) may lead to superior fear reduction when compared to the standard extinction of undergoing the experience themselves (Golkar et al., 2013, 2016). Hence, beyond the behaviour and cognitive techniques employed in CBT, the additional layer of yet another learning mechanism, may prove beneficial.

Group-based CBT has had some positive results in the tinnitus literature (e.g. Cima et al., 2012), but it is not yet known if the group-based treatment is a contributing part of CBT for tinnitus or for whom it may be more beneficial. Group and individual CBT methods applied to tinnitus were previously compared by Fuller et al. (2020) in the context of a meta-analysis. Both individual and group-based CBT were more effective than wait list control or “active comparison” conditions and no difference between individual or group-based treatment was found. These comparisons were made on the results from separate studies, using either individual or group-based CBT and no study has compared one specific treatment protocol under each condition.

*Objective:*

Contrast and compare individual to group-based CBT in the treatment of tinnitus

## Summary

With a brief account of the current state of the tinnitus field we have identified three areas of interest in which a novel contribution to the field would more likely result in direct progress in tinnitus research and treatment. From each area of interest specific objectives were derived as follows:

The first objective of the current work relies on the adaptation/creation of an associative learning model for the field of tinnitus in order to establish a replicable fear learning paradigm with healthy human participants (Chapter 2).

Two objectives were drawn from the area of tinnitus assessment. They were to (1) investigate whether EMA negatively affects tinnitus experience and to (2) investigate whether EMA provides a more accurate measure of tinnitus experience when compared to EDD assessments. These objectives are respectively reflected in the studies reported on Chapters 3 and 4 of the current PhD dissertation.

In the area of tinnitus treatment, the current CBT approach and desire to tailor treatment inspired the investigation of contrasting and comparing individual to group-based CBT for tinnitus (Chapter 5).

With the results reported in each study, Chapter 6 offers a synthesis and general discussion on the overall findings of the current body of work. Future research perspectives are drawn and a statement on the impact of this dissertation is made.

## References

- Andersson, G. (2003). Tinnitus loudness matchings in relation to annoyance and grading of severity. *Auris Nasus Larynx*, *30*(2), 129–133. [https://doi.org/10.1016/S0385-8146\(03\)00008-7](https://doi.org/10.1016/S0385-8146(03)00008-7)
- Atcherson, S. R., Zraick, R. I., & Brasseur, R. E. (2011). Readability of Patient-Reported Outcome Questionnaires For Use With Persons With Tinnitus. *Ear and Hearing*, *32*(5), 671–673. <https://doi.org/10.1097/AUD.0b013e3182134654>
- Baguley, D. (2003). Hyperacusis. *JRSM*, *96*(12), 582–585. <https://doi.org/10.1258/jrsm.96.12.582>
- Baguley, D., Andersson, G., McFerran, D., & McKenna, L. (2013). Psychological Models of Tinnitus. In *Tinnitus: A Multidisciplinary Approach* (pp. 102–109). John Wiley & Sons, Ltd.
- Bandura, A., Grusec, J. E., & Menlove, F. L. (1967). Vicarious extinction of avoidance behavior. *Journal of Personality and Social Psychology*, *5*(1), 16–23. <https://doi.org/10.1037/h0024182>
- Beck, A. (1979). *Cognitive Therapy and the Emotional Disorders*. International Universities Press.
- Belli, S., Belli, H., Bahcebasi, T., Ozcetin, A., Alpay, E., & Ertem, U. (2008). Assessment of psychopathological aspects and psychiatric comorbidities in patients affected by tinnitus. *European Archives of Oto-Rhino-Laryngology*, *265*(3), 279–285. <https://doi.org/10.1007/s00405-007-0440-8>
- Biswas, R., & Hall, D. A. (2020). Epidemiology of Tinnitus. In *Tinnitus* (pp. 23–45). [https://doi.org/10.1007/7854\\_2020\\_154](https://doi.org/10.1007/7854_2020_154)
- Biswas, R., Lugo, A., Hall, D., Akeroyd, M., Liu, X., Schlee, W., & Gallus, S. (2020). A study of the pan-European prevalence of tinnitus and hearing difficulty using a standardized set of questions. *Poster Presented at Association of Research in Otolaryngology 43rd Annual MidWinter Meeting*. [https://aro.org/wp-content/uploads/2020/02/2020-Abstracts\\_1-21-20-Web.pdf](https://aro.org/wp-content/uploads/2020/02/2020-Abstracts_1-21-20-Web.pdf)
- Brozoski, T. J., Wisner, K. W., Sybert, L. T., & Bauer, C. A. (2012). Bilateral Dorsal Cochlear Nucleus Lesions Prevent Acoustic-Trauma Induced Tinnitus in an Animal Model. *Journal of the Association for Research in Otolaryngology*, *13*(1), 55–66. <https://doi.org/10.1007/s10162-011-0290-3>
- Brüggemann, P., Szczepek, A. J., Rose, M., McKenna, L., Olze, H., & Mazurek, B. (2016). Impact of Multiple Factors on the Degree of Tinnitus Distress. *Frontiers in Human Neuroscience*, *10*. <https://doi.org/10.3389/fnhum.2016.00341>
- Budd, R. J., & Pugh, R. (1996). The relationship between coping style, tinnitus severity and emotional distress in a group of tinnitus sufferers. *British Journal of Health Psychology*, *1*(3), 219–229. <https://doi.org/10.1111/j.2044-8287.1996.tb00504.x>
- Cederroth, C. R., Gallus, S., Hall, D. A., Kleinjung, T., Langguth, B., Maruotti, A., Meyer, M., Norena, A., Probst, T., Pryss, R., Searchfield, G., Shekhawat, G., Spiliopoulou, M., Vanneste, S., & Schlee, W. (2019). Editorial: Towards an Understanding of Tinnitus Heterogeneity. *Frontiers in Aging Neuroscience*, *11*. <https://doi.org/10.3389/fnagi.2019.00053>
- Cima, R. F. F. (2018). Bothersome tinnitus: Cognitive behavioral perspectives. *Hno*, *66*(5), 369–374. <https://doi.org/10.1007/s00106-018-0502-9>
- Cima, R. F. F., Crombez, G., & Vlaeyen, J. W. S. (2011). Catastrophizing and Fear of Tinnitus Predict Quality of Life in Patients With Chronic Tinnitus. *Ear and Hearing*, *32*(5), 634–641. <https://doi.org/10.1097/AUD.0b013e31821106dd>

- Cima, R. F. F., Maes, I. H., Joore, M. A., Scheyen, D. J. W. M., El Refaie, A., Baguley, D. M., Anteunis, L. J. C., van Breukelen, G. J. P., & Vlaeyen, J. W. S. (2012). Specialised treatment based on cognitive behaviour therapy versus usual care for tinnitus: a randomised controlled trial. *The Lancet*, *379*(9830), 1951–1959. [https://doi.org/10.1016/S0140-6736\(12\)60469-3](https://doi.org/10.1016/S0140-6736(12)60469-3)
- Cima, R. F. F., Mazurek, B., Haider, H., Kikidis, D., Lapira, A., Noreña, A., & Hoare, D. J. (2019). A multidisciplinary European guideline for tinnitus: diagnostics, assessment, and treatment. *HNO, February 2018*. <https://doi.org/10.1007/s00106-019-0633-7>
- Cima, R. F. F., van Breukelen, G., & Vlaeyen, J. W. S. (2018). Tinnitus-related fear: Mediating the effects of a cognitive behavioural specialised tinnitus treatment. *Hearing Research*, *358*, 86–97. <https://doi.org/10.1016/j.heares.2017.10.003>
- Cima, R. F. F., Vlaeyen, J. W. S., Maes, I. H. L., Joore, M. A., & Anteunis, L. J. C. (2011). Tinnitus Interferes With Daily Life Activities: A Psychometric Examination of the Tinnitus Disability Index. *Ear and Hearing*, *32*(5), 623–633. <https://doi.org/10.1097/AUD.0b013e31820dd411>
- Das, S. K., Wineland, A., Kallogjeri, D., & Piccirillo, J. F. (2012). Cognitive speed as an objective measure of tinnitus. *The Laryngoscope*, *122*(11), 2533–2538. <https://doi.org/10.1002/lary.23555>
- De Ridder, D., Elgoyhen, A. B., Romo, R., & Langguth, B. (2011). Phantom percepts: Tinnitus and pain as persisting aversive memory networks. *Proceedings of the National Academy of Sciences*, *108*(20), 8075–8080. <https://doi.org/10.1073/pnas.1018466108>
- De Ridder, Dirk, Schlee, W., Vanneste, S., Londero, A., Weisz, N., Kleinjung, T., Shekhawat, G. S., Elgoyhen, A. B., Song, J.-J., Andersson, G., Adhia, D., de Azevedo, A. A., Baguley, D. M., Biesinger, E., Binetti, A. C., Del Bo, L., Cederroth, C. R., Cima, R., Eggermont, J. J., ... Langguth, B. (2021). *Tinnitus and tinnitus disorder: Theoretical and operational definitions (an international multidisciplinary proposal)* (pp. 1–25). <https://doi.org/10.1016/bs.pbr.2020.12.002>
- Elhai, J. D., Dvorak, R. D., Levine, J. C., & Hall, B. J. (2017). Problematic smartphone use: A conceptual overview and systematic review of relations with anxiety and depression psychopathology. *Journal of Affective Disorders*, *207*, 251–259. <https://doi.org/10.1016/j.jad.2016.08.030>
- Ellis, A., & Grieger, R. (1977). *Handbook of Rational-Emotive Therapy* (Vol. 1.). Springer.
- Figueiredo, R. R., Rates, M. A., Azevedo, A. A. de, Oliveira, P. M. de, & Navarro, P. B. A. de. (2010). Correlation analysis of hearing thresholds, validated questionnaires and psychoacoustic measurements in tinnitus patients. *Brazilian Journal of Otorhinolaryngology*, *76*(4), 522–526. <http://www.ncbi.nlm.nih.gov/pubmed/20835541>
- Fuller, T. E. (2021). *Background noise: Clinical guidelines, assessment & CBT for disabling tinnitus*.
- Fuller, T. E., Cima, R., Langguth, B., Mazurek, B., Vlaeyen, J. W. S., & Hoare, D. J. (2020). Cognitive behavioural therapy for tinnitus. *Cochrane Database of Systematic Reviews*, *4*. <https://doi.org/10.1002/14651858.CD012614.pub2>
- Gerull, K. M., Kallogjeri, D., Piccirillo, M. L., Rodebaugh, T. L., Lenze, E. J., & Piccirillo, J. F. (2019). Feasibility of Intensive Ecological Sampling of Tinnitus in Intervention Research. *Otolaryngology–Head and Neck Surgery*, 019459981984496. <https://doi.org/10.1177/0194599819844968>
- Gilles, A., Van Hal, G., De Ridder, D., Wouters, K., & Van de Heyning, P. (2013). Epidemiology of Noise-Induced Tinnitus and the Attitudes and Beliefs towards Noise and Hearing Protection in Adolescents. *PLoS ONE*, *8*(7), e70297. <https://doi.org/10.1371/journal.pone.0070297>



- Goldberg, R. L., Piccirillo, M. L., Nicklaus, J., Skillington, A., Lenze, E., Rodebaugh, T. L., Kallogjeri, D., & Piccirillo, J. F. (2017). Evaluation of ecological momentary assessment for tinnitus severity. *JAMA Otolaryngology - Head and Neck Surgery*, *143*(7), 700–706. <https://doi.org/10.1001/jamaoto.2017.0020>
- Hall, D. A., Fackrell, K., Li, A. B., Thavayogan, R., Smith, S., Kennedy, V., Tinoco, C., Rodrigues, E. D., Campelo, P., Martins, T. D., Lourenço, V. M., Ribeiro, D., & Haider, H. F. (2018). A narrative synthesis of research evidence for tinnitus-related complaints as reported by patients and their significant others. *Health and Quality of Life Outcomes*, *16*(1), 1–15. <https://doi.org/10.1186/s12955-018-0888-9>
- Hallam, R. S., Jakes, S. C., & Hinchcliffe, R. (1988). Cognitive variables in tinnitus annoyance. *The British Journal of Clinical Psychology*, *27* ( Pt 3)(pt 3), 213–222. <http://www.ncbi.nlm.nih.gov/pubmed/3191301>
- Hallam, R. S., Rachman, S., & Hinchcliffe, R. (1984). Psychological aspects of tinnitus. *Contributions to Medical Psychology*, *3*, 31–53.
- Henry, J. A. (2016). “Measurement” of Tinnitus. *Otology & Neurotology*, *37*(8), e276–e285. <https://doi.org/10.1097/MAO.0000000000001070>
- Henry, J. A., Galvez, G., Turbin, M. B., Thielman, E. J., Mcmillan, G. P., & Istvan, J. A. (2012). *Pilot Study to Evaluate Ecological Momentary Assessment of Tinnitus*. 279–290.
- Horwood, S., & Anglim, J. (2019). Problematic smartphone usage and subjective and psychological well-being. *Computers in Human Behavior*, *97*, 44–50. <https://doi.org/10.1016/j.chb.2019.02.028>
- Jastreboff, P. J. (2011). Tinnitus Retraining Therapy. In A. R. Møller, B. Langguth, D. D. De Ridder, & T. Kleinjung (Eds.), *Textbook of Tinnitus* (pp. 575–596). Springer New York.
- Jastreboff, Pawel J. (1990). Phantom auditory perception (tinnitus): mechanisms of generation and perception. *Neuroscience Research*, *8*(4), 221–254. [https://doi.org/10.1016/0168-0102\(90\)90031-9](https://doi.org/10.1016/0168-0102(90)90031-9)
- Jastreboff, Pawel J., Brennan, J. F., Coleman, J. K., & Sasaki, C. T. (1988). Phantom auditory sensation in rats: An animal model for tinnitus. *Behavioral Neuroscience*, *102*(6), 811–822. <https://doi.org/10.1037/0735-7044.102.6.811>
- Jastreboff, Pawel J., & Hazell, J. W. P. (2004). *Tinnitus Retraining Therapy*. Cambridge University Press. <https://doi.org/10.1017/CBO9780511544989>
- Jastreboff, Pawel J., Hazell, J. W. P., & Graham, R. L. (1994). Neurophysiological model of tinnitus: Dependence of the minimal masking level on treatment outcome. *Hearing Research*, *80*(2), 216–232. [https://doi.org/10.1016/0378-5955\(94\)90113-9](https://doi.org/10.1016/0378-5955(94)90113-9)
- Kaltenbach, J. A., Rachel, J. D., Mathog, T. A., Zhang, J., Falzarano, P. R., & Lewandowski, M. (2002). Cisplatin-Induced Hyperactivity in the Dorsal Cochlear Nucleus and Its Relation to Outer Hair Cell Loss: Relevance to Tinnitus. *Journal of Neurophysiology*, *88*(2), 699–714. <https://doi.org/10.1152/jn.2002.88.2.699>
- Kleinstäuber, M., Jasper, K., Schweda, I., Hiller, W., Andersson, G., & Weise, C. (2013). The Role of Fear-Avoidance Cognitions and Behaviors in Patients with Chronic Tinnitus. *Cognitive Behaviour Therapy*, *42*(2), 84–99. <https://doi.org/10.1080/16506073.2012.717301>
- Kröner-Herwig, B., Frenzel, A., Fritsche, G., Schilkowsky, G., & Esser, G. (2003). The management of chronic tinnitus. *Journal of Psychosomatic Research*, *54*(4), 381–389. [https://doi.org/10.1016/S0022-3999\(02\)00400-2](https://doi.org/10.1016/S0022-3999(02)00400-2)

- Kuk, F. K., Tyler, R. S., Russell, D., & Jordan, H. (1990). The Psychometric Properties of a Tinnitus Handicap Questionnaire. *Ear and Hearing, 11*(6), 434–445. <https://doi.org/10.1097/00003446-199012000-00005>
- Langguth, B. (2011). A review of tinnitus symptoms beyond ‘ringing in the ears’: a call to action. *Current Medical Research and Opinion, 27*(8), 1635–1643. <https://doi.org/10.1185/03007995.2011.595781>
- Langguth, B., Landgrebe, M., Kleinjung, T., Sand, G. P., & Hajak, G. (2011). Tinnitus and depression. *The World Journal of Biological Psychiatry, 12*(7), 489–500. <https://doi.org/10.3109/15622975.2011.575178>
- Lazarus, R. S. (1991). Progress on a cognitive-motivational-relational theory of emotion. *American Psychologist, 46*(8), 819–834. <https://doi.org/10.1037/0003-066X.46.8.819>
- Lethem, J., Slade, P. D., Troup, J. D., & Bentley, G. (1983). Outline of a Fear-Avoidance Model of exaggerated pain perception--I. *Behaviour Research and Therapy, 21*(4), 401–408. <http://www.ncbi.nlm.nih.gov/pubmed/6626110>
- Linton, S. J., Nicholas, M., & Shaw, W. (2018). Why wait to address high-risk cases of acute low back pain? A comparison of stepped, stratified, and matched care. *Pain, 159*(12), 2437–2441. <https://doi.org/10.1097/j.pain.0000000000001308>
- McCormack, A., Edmondson-Jones, M., Somerset, S., & Hall, D. A. (2016). A systematic review of the reporting of tinnitus prevalence and severity. *Hearing Research, 337*, 70–79. <https://doi.org/10.1016/j.heares.2016.05.009>
- McFerran, D. J., Stockdale, D., Holme, R., Large, C. H., & Baguley, D. M. (2019). Why Is There No Cure for Tinnitus? *Frontiers in Neuroscience, 13*. <https://doi.org/10.3389/fnins.2019.00802>
- Mckenna, L. (2004). Models of tinnitus suffering and treatment compared and contrasted. *Audiological Medicine, 2*(1), 41–53. <https://doi.org/10.1080/16513860410028735>
- McKenna, L., Handscomb, L., Hoare, D. J., & Hall, D. A. (2014). A Scientific Cognitive-Behavioral Model of Tinnitus: Novel Conceptualizations of Tinnitus Distress. *Frontiers in Neurology, 5*(OCT), 1–15. <https://doi.org/10.3389/fneur.2014.00196>
- Mckenna, L., & Irwin, R. (2008). Sound therapy for tinnitus – sacred cow or idol worship?: An investigation of the evidence. *Audiological Medicine, 6*(1), 16–24. <https://doi.org/10.1080/16513860801899389>
- McKenna, L., Marks, E. M., Hallsworth, C. A., & Schiette, R. (2017). Mindfulness-Based Cognitive Therapy as a Treatment for Chronic Tinnitus: A Randomized Controlled Trial. *Psychotherapy and Psychosomatics, 86*(6), 351–361. <https://doi.org/10.1159/000478267>
- Meikle, M. B., Henry, J. A., Griest, S. E., Stewart, B. J., Abrams, H. B., McArdle, R., Myers, P. J., Newman, C. W., Sandridge, S., Turk, D. C., Folmer, R. L., Frederick, E. J., House, J. W., Jacobson, G. P., Kinney, S. E., Martin, W. H., Nagler, S. M., Reich, G. E., Searchfield, G., ... Vernon, J. A. (2012). The Tinnitus Functional Index. *Ear and Hearing, 33*(2), 153–176. <https://doi.org/10.1097/AUD.0b013e31822f67c0>
- Meulders, A. (2020). Fear in the context of pain: Lessons learned from 100 years of fear conditioning research. *Behaviour Research and Therapy, 131*(October 2019), 103635. <https://doi.org/10.1016/j.brat.2020.103635>
- Michie, S., Richardson, M., Johnston, M., Abraham, C., Francis, J., Hardeman, W., Eccles, M. P., Cane, J., & Wood, C. E. (2013). The Behavior Change Technique Taxonomy (v1) of 93 Hierarchically Clustered Techniques: Building an International Consensus for the Reporting of Behavior Change Interventions. *Annals of Behavioral Medicine, 46*(1), 81–95. <https://doi.org/10.1007/s12160-013-9486-6>

- Møller, A. R. (1997). Similarities between chronic pain and tinnitus. *The American Journal of Otolaryngology*, 18(5), 577–585. <http://www.ncbi.nlm.nih.gov/pubmed/9303153>
- Møller, A. R. (2007). *Tinnitus and pain* (pp. 47–53). [https://doi.org/10.1016/S0079-6123\(07\)66004-X](https://doi.org/10.1016/S0079-6123(07)66004-X)
- Newman, C. W., Jacobson, G. P., & Spitzer, J. B. (1996). Development of the Tinnitus Handicap Inventory. *Archives of Otolaryngology - Head and Neck Surgery*, 122(2), 143–148. <https://doi.org/10.1001/archotol.1996.01890140029007>
- Newman, C. W., Sandridge, S. A., & Jacobson, G. P. (1998). Psychometric adequacy of the Tinnitus Handicap Inventory (THI) for evaluating treatment outcome. *Journal of the American Academy of Audiology*, 9(2), 153–160. <http://www.ncbi.nlm.nih.gov/pubmed/9564679>
- Organ of the Chautauqua Literary and Scientific Circle. (1883). *The Chautauquan*, 3(9), 543.
- Pavlov, I. P. (1927). *Conditioned reflexes: an investigation of the physiological activity of the cerebral cortex*. Oxford Univ. Press.
- Rabau, S., Cox, T., Punte, A. K., Waelkens, B., Gilles, A., Wouters, K., de Varebeke, S. J., & Van de Heyning, P. (2015). Changes over time of psychoacoustic outcome measurements are not a substitute for subjective outcome measurements in acute tinnitus. *European Archives of Oto-Rhino-Laryngology*, 272(3), 573–581. <https://doi.org/10.1007/s00405-013-2876-3>
- Rachman, S. (1977). The conditioning theory of fear-acquisition: A critical examination. *Behavior Research and Therapy*, 15, 375–387.
- Robinson, M. E., Staud, R., & Price, D. D. (2013). Pain Measurement and Brain Activity: Will Neuroimages Replace Pain Ratings? *The Journal of Pain*, 14(4), 323–327. <https://doi.org/10.1016/j.jpain.2012.05.007>
- Schlee, W., Pryss, R. C., Probst, T., Schobel, J., Bachmeier, A., Reichert, M., & Langguth, B. (2016). Measuring the moment-to-moment variability of Tinnitus: The TrackYourTinnitus smart phone app. *Frontiers in Aging Neuroscience*, 8(DEC), 1–8. <https://doi.org/10.3389/fnagi.2016.00294>
- Schneider, S., & Stone, A. A. (2016). Ambulatory and diary methods can facilitate the measurement of patient-reported outcomes. *Quality of Life Research*, 25(3), 497–506. <https://doi.org/10.1007/s11136-015-1054-z>
- Shakespeare, W. (1605). *A midsummer night's dream*.
- Shields, A. L., Shiffman, S., & Stone, A. (2016). Recall bias: Understanding and reducing bias in PRO data collection. In *EPro: Electronic solutions for patient-reported data* (pp. 5–21).
- Skinner, B. F. (1938). *The behavior of organisms: an experimental analysis*. Appleton-Century.
- Stone, A. A., & Shiffman, S. (1994). Ecological Momentary Assessment (Ema) in Behavioral Medicine. *Annals of Behavioral Medicine*, 16(3), 199–202. <https://doi.org/10.1093/abm/16.3.199>
- Timmer, B. H. B., Hickson, L., & Launer, S. (2018). The use of ecological momentary assessment in hearing research and future clinical applications. *Hearing Research*, 1–5. <https://doi.org/10.1016/j.heares.2018.06.012>
- Vahedi, Z., & Saiphoo, A. (2018). The association between smartphone use, stress, and anxiety: A meta-analytic review. *Stress and Health*, 34(3), 347–358. <https://doi.org/10.1002/smi.2805>
- Verbrugge, L. M. (1980). Health Diaries. *Medical Care*, 18(1), 73–95. <https://doi.org/10.1097/00005650-198001000-00006>

- Vlaeyen, J. W. S., & Crombez, G. (2020). Behavioral Conceptualization and Treatment of Chronic Pain. *Annual Review of Clinical Psychology, 16*, 187–212. <https://doi.org/10.1146/annurev-clinpsy-050718-095744>
- Vlaeyen, J. W. S., & Linton, S. J. (2000). Fear-avoidance and its consequences in chronic musculoskeletal pain: a state of the art. *Pain, 85*(3), 317–332. <http://www.ncbi.nlm.nih.gov/pubmed/10781906>
- Vlaeyen, J. W. S., & Linton, S. J. (2012). Fear-avoidance model of chronic musculoskeletal pain: 12 years on. *Pain, 153*(6), 1144–1147. <https://doi.org/10.1016/j.pain.2011.12.009>
- Vlaeyen, J. W. S., Morley, S., Linton, S. J., Boersma, K., & de Jong, J. (2012). *Pain-related fear: Exposure-based treatment of chronic pain*. IASP Press.
- Wallhäuser-Franke, E., D'Amelio, R., Glauner, A., Delb, W., Servais, J. J., Hörmann, K., & Repik, I. (2017). Transition from Acute to Chronic Tinnitus: Predictors for the Development of Chronic Distressing Tinnitus. *Frontiers in Neurology, 8*(NOV), 1–13. <https://doi.org/10.3389/fneur.2017.00605>
- Watson, J. B., & Rayner, R. (1920). Conditioned emotional reactions. *Journal of Experimental Psychology, 3*(1), 1–34. <https://doi.org/10.1037/0003-066X.55.3.313>





## CHAPTER 2

---

Fear in the ear:  
the role of fear conditioning in the  
acquisition and extinction of fear of sounds

## Abstract


**Introduction** Fear associations to a previously neutral sound may explain Chronic tinnitus experience. Replicable fundamental research of the associative learning models at the core of development of chronic tinnitus is yet to be created for human participants. This study aims at introducing a novel tinnitus relevant differential conditioning paradigm with healthy participants. **Method** Two acoustically equivalent probes presented for 5s at 50 dB were presented to participants where one probe predicted a second aversive acoustic probe at 98 dB for 1s. Expectancy ratings, heart rate acceleration and Skin Conductance Response (SCR) were collected from 62 participants (82% females,  $M_{\text{age}} = 20.65$ ). Audiological measures, Acoustic Stapedial Reflex Thresholds (ASRT) and Loudness Discomfort Levels (LDL) changes were explored within the paradigm. **Results** Expectancy ratings indicated differential fear acquisition and indiscriminate extinction and reinstatement to the two acoustic probes. Heart rate acceleration and SCR measures did not converge with expectancy ratings. LDL increased during the experiment while ASRT remained unchanged or decreased. **Conclusion** Support for the role of fear in the development and maintenance of tinnitus is found. The paradigm sets the foundations for future research on the underlying mechanisms of chronic tinnitus to be investigated within healthy human participants.



## 1. Introduction

Tinnitus, the perception of sound (e.g. ringing, hissing) without corresponding acoustic source, affects between 11.9-30.3% of individuals (McCormack et al., 2016). Despite being a relative common and harmless experience, 1-6% of individuals suffer from it (Davis and Refaie, 2000; Gallus et al., 2015; Kim et al., 2015), with a number of physical (e.g. sleep difficulties) and psychological (e.g. fear, distress) complaints (Hall et al., 2018). Characterizations of tinnitus (e.g. loudness, lateralization, chronicity, somatic modulation) have failed to explain tinnitus distress and disability (e.g. Ward et al., 2015). An explanation for the development and maintenance of chronic tinnitus may be found in the Fear Avoidance (FA) model of tinnitus (Figure 1; Cima et al., 2011; Kleinstäuber et al., 2013) – adapted from pain (Lethem et al., 1983; Vlaeyen and Linton, 2012, 2000). While there are other perspectives, such as the Habituation model (Hallam et al., 1984), Neurophysiological model (Jastreboff, 1990; Jastreboff et al., 1988), and Cognitive Behaviour model (McKenna et al., 2014), it is FA model that holds the biggest promise by being robustly supported in parallel fields (i.e. chronic pain). Vital to the FA model is the role of specific tinnitus-related fear, as opposed to audiologic characteristics, in triggering the dysfunctional cycle. The model further predicts that those who do not hold catastrophic (mis)interpretations of the tinnitus percept and associated fear more easily recover. Thus, the onset of chronic disabling tinnitus may lie in the development of learned fear responses to the tinnitus percept (i.e. fear conditioning).

**Figure 1:** Fear Avoidance model (replicated from Cima et al., 2018)



Associative learning, specifically fear conditioning, a particular type of classical (Pavlovian) conditioning, relies on the contingent pairing of a neutral (conditioned stimulus = CS), like a bell, and a biologically salient (unconditioned = US) stimulus, like shock or pain, which elicits a fear response (unconditioned response UR). After repeated pairings, the CS in itself will elicit fear responses (conditioned response = CR). In other words, it is learned that if the CS occurs, the US will or might follow, and fear responses ensue, even in the absence of the US. In the case of chronic pain for example, a normally innocuous movement (CS) (e.g. bending over), has co-occurred with a painful event (US). The individual learns that bending signals pain and fear for this movement (CR) ensues, even in the absence of such intense pain. Such a direct translation between CS, CR and US, to movement, fear and pain, has made a fear conditioning paradigm a particularly fruitful tool in the field of chronic pain research, which helped in providing robust foundations for the FA model (for a review on fear conditioning in the context of pain see Meulders, 2020). Moreover, the tenets of Cognitive Behavioural Therapy (CBT) can be investigated in fear learning paradigms. Exposure techniques, for example, are commonly used in CBT to diminish fear responses and entail the experience of the feared stimulus (e.g. bell) without the expected negative outcome (e.g. pain). This can be replicated in fear learning paradigms through extinction phases, where the CS+ is repeatedly presented without the US, challenging the previously learned association. Furthermore, a single presentation of the US without CS after extinction adds another layer to test for the strength of the fear association made during acquisition.

Considering hypothesized parallels between chronic pain and chronic tinnitus disability (e.g. Cima et al., 2011; Kleinstäuber et al., 2013), an associative learning paradigm for tinnitus might be adapted as follows: tinnitus, conceptualized as a perception of a neutral sound (CS), acquires fear responses (UR) experienced due its association with an aversive event (US). In translating fear associations to tinnitus research, a biologically relevant US is necessary. The choice of a loud sound as an US can be based on three reasons: (1) the overlap of hyperacusis (i.e. reduced sound tolerance) and chronic tinnitus (Baguley, 2003), (2) the avoidance of loud sounds due to perceived triggering or worsening of tinnitus (Kleinstäuber et al., 2013), and (3) the perception of tinnitus after the experience of loud sounds (Gilles et al., 2013). The focus of this study is to test such a novel fear-learning paradigm.

The investigation of associative learning models (i.e. fear conditioning) often employ differential conditioning paradigms (for a review on methodological considerations

for fear conditioning paradigms see Lonsdorf et al., 2017). These paradigms employ a stimulus that is paired with a US (the CS+) and a stimulus which is not (the CS-), such that only the CS+ signals the occurrence of the US. Conditioned fear is then measured by the differences between responses elicited by CS+ and CS-, removing the need for a control group and controlling for inter-individual differences. Assessments of fear are achieved through self-reported expectancy ratings (i.e. expectancy of US after CS+/-) and/or physiological responses (i.e. fear potentiated startle, heart-rate acceleration, skin conductance response). While expectancy ratings are often used and regarded as an accurate measure of fear (Boddez et al., 2013), self-report biases and contingency awareness may influence responses. Physiological responses are resistant to self-report bias. Skin Conductance Response (SCR) and Heart Rate (HR) acceleration are capable of measuring the difference between CS+ and CS- in fear conditioning paradigms. Nevertheless, SCR and HR acceleration measurements may not converge as they reflect different dimensions of fear (e.g. expectancy and affective components). Finally, Fear Potentiated Startles (FPS), defensive reflexes that may be probed through sudden sensory input, may increase during threatening circumstances (e.g. Aslaksen et al., 2016) providing yet another complementary measure of fear.

In the current study, we hypothesize that after repeated presentation of a neutral sound (CS+) followed with a short delay by the same sound at higher intensity (US), the CS+ will elicit fear responses (as measure by self-reported expectancy ratings and physiological responses) in healthy participants even in absence of the US. A different auditory stimulus that is not paired with the US (CS-) will not elicit these fear responses. At a second phase we expect a reduction (i.e. extinction) of fear, that is, a decreased fear differentiation between CS+ and CS-, after repeated presentation of CS- and CS+ without US. Reinstatement of fear is tested by one single presentation of US after extinction followed by repeated presentations of both CS+ and CS-, for which we expect an increased difference in fear, which is to diminish over time. Furthermore, FPS in the form of Acoustic Stapedial Reflex Thresholds (ASRT) are explored within the paradigm as well as changes in sound tolerance.

## 2. Method

### 2.1. Participants

Participants were recruited through advertisement flyers and posters placed in the university buildings of KU Leuven, as well as online (e.g. social media). Healthy adult

volunteers (between 18-55 years old) were eligible to participate if they reported (1) absence of significant hearing loss measured by pure tone audiometry, (2) absence of acute or chronic tinnitus as defined by the perception of sound(s) in your head or ears (such as ringing or buzzing) in the absence of any corresponding external source, (3) absence of a cardiovascular disease, mental/or neurological disorder, (4) not being asked by their doctor to avoid stressful situations, and (5) being proficient in the English language. Written informed consent was obtained prior to the start of the study and compensation was either course credit or five euros (to be chosen by the participant). The study was approved by the KU Leuven's Social and Societal Ethics Committee (SMEC; G-2019 07 1693), and the hypotheses and main analysis were pre-registered (<https://aspredicted.org/8fp3y.pdf>).

## *2.2. Hearing thresholds*

Formal audiometry with hearing thresholds and audiogram were conducted for both ears. A pure tone average (PTA) was calculated at 500, 1000, 2000, 4000, and 8000 Hz for each ear. Participants were excluded if the PTA in one or both ears exceeded 20 dB HL. A more sensitive averaged PTA of higher frequencies (1000, 2000 and 4000; PTA<sub>high</sub>) was further calculated if normal hearing thresholds were established by all participants.

## *2.3. Sound tolerance*

Sound tolerance was measured through Loudness Discomfort Level (LDL) test. The test utilizes the presentation of a 1 s white noise at 40 dB with stepped increases of 5 dB until participant indicated to stop or maximal level was reached (never exceeding 100 dB) for each ear. LDL was measured for 125, 250, 500, 1000, 4000, 6000 and 8000 Hz.

## *2.4. Expectancy measures*

Expectancy ratings were recorded on a Visual Analog Scale (VAS), with anchors set at 0, 50 and 100%. The question – “*How much do you expect a loud sound to follow?*” – was presented during each CS.

## *2.5. Physiological fear measures*

Skin Conductance Response (SCR) was measured continuously at 1000 Hz. Based on Sjouwerman & Lonsdorf (2018), skin conductance data was resampled to 100 Hz offline.

SCR was scored trough-to-peak in a response window of 1-5 s after stimulus onset. Responses smaller than 0.02  $\mu$ S are scored as an amplitude of 0  $\mu$ S. Log transformations of SCR were performed when distributions violated normality assumptions of the statistical analyses. Based on recommendations of Lonsdorf et al. (2019), non-responders were defined as those who held a response of 0 for at least three out of the five US during the acquisition phase.


Heart Rate (HR) responses were captured by assessing the interpolated HR in the first 5 s after CS onset and segmenting that signal into 0.5 s epochs. HR acceleration was measured by subtracting the mean HR of the second prior to CS onset from every 0.5 s epoch (if data for the second prior to CS was missing, the first available epoch was used instead). To assess the conditioned preparation for defensive action, we measured the maximum HR acceleration within a time window of 1-5 s after CS onset (Burger et al., 2016; Van Diest et al., 2009).

Fear potentiated startles were adapted to the paradigm by exploring Acoustic Stapedial Reflex Threshold (ASRT). While eye-blink startles have been previously used, an acoustically specific reflex response would raise ecological validity of the paradigm. Located in the middle ear, the stapedius muscle contraction may be triggered by loud noises, making it an objectively measurable reflex that may further clarify the role of fear in auditory perceptions. ASRT, was measured with 1 s pure tone (contralateral) and 1s pulsed tone (ipsilateral) of 1000 Hz. Stimulus level was increased by 5 dB from 80 dB with a 2 s break between probes (never exceeding 100 dB). Once the ASRT was exceeded (deviation of 0,02 ml), the stimulus level presentation was repeated and then recorded. To our knowledge, the ASRT are yet to be utilized in the context of fear learning. As such, ASRT responsiveness is explored in the current study.

## ***2.6. Procedure***

Audiometry, LDL and ASRT measurements were recorded at baseline (T0), after acquisition (T1) and at the end of the experiment (T2) in a different audiometric soundproof room from the experimental paradigm set-up (Figure 2).

**Figure 2:** Sequence of events in the experimental design. Audiometry, Loudness Discomfort Levels (LDL) and Acoustic Stapedial Reflex Thresholds (ASRT) performed at a separate room from the experimental set-up.



Two acoustically equivalent probes served as conditioned stimuli (CS). The probes were a warble (4 kHz, modulated by 100 Hz, moderation frequency of 20 Hz) and a pure (4 kHz) tone counterbalanced as CS+ and CS-. Both CS+ and CS- were presented for 5 s at 50 dB. The US was an acoustic probe equal to the CS+ however presented at almost twice the intensity (98 dB) for 1 s. A black screen with a fixation cross was presented for each 8 s intertrial interval (ITI).

Acquisition relied on a delayed differential conditioned paradigm of 20 trials. Ten CS+ and CS- were counterbalanced and delivered at a pseudo-random sequence (i.e. never exceeding two successive presentations). A 100% reinforcement schedule was used where the US immediately followed CS+. Extinction comprised ten trials of CS+ and CS- at a pseudo-random order without US. Reinstatement was tested through a single non-contingent presentation of the US (8 s after the last CS of the extinction phase and before the first CS of the reinstatement phase) and followed by another ten trials of CS+ and CS- at a pseudo-random delivery order.

## 2.7. Apparatus

Acoustic stimuli were presented via an external soundcard (Fireface UCX; RME, Haimhausen, Germany), connected to DD45 transducers embedded in Peltor caps. The set-up was calibrated with a Brüel & Kjaer Sound (Nærum, Denmark) level meter 2260 and a Brüel & Kjaer artificial ear 4153 using the flat plate (mono calibration of right

side). Due to binaural summation, stimuli were calibrated at 3 dB lower. CS stimuli were calculated from 80 dB. In order to correct for environmental low band noise in the echoic room, “A peaks” (dB A) were averaged from original sounds (Warble = 46.6 dB A equivalent average; Pure = 47.3 dB A equivalent average). US stimuli were calibrated to approximately 95dB A peak (Warble = 94.6 dB A peak; Pure = 94.7 dB A peak).

Formal audiometry and Loudness Discomfort Level (LDL) tests were carried-out through the Audiometer Madson Astera by Otometrics (Natus Medical Incorporated, California, USA) type 1066 with TDH-39 headphones. ASRT tests utilized the Madsen Zodiac Diagnostic & Clinical Stand-Alone by Otometrics (Natus Medical Incorporated, California, USA) type 1096 SA, TDH-39P headphones (Telephonics, New York, USA) and diagnostic probe (with shoulder strap).

SCR was recorded with a Coulbourn Instruments Isolated Skin Conductance Coupler S71-23 (Coulbourn Instruments, Pennsylvania, USA), through isotonic disposable electrodes (EL507; Biopac Systems Inc., California, USA) connected at the hypothenar eminence of the non-dominant hand. HR responses were recorded through gelled disposable electrodes (Kendall H66LG; Covidien, Dublin, Ireland) attached to the chest in a triangle, with one underneath each collarbone and one under the participants lowermost left rib. The signal was recorded at 1000 Hz, using the Isolated Bioamplifier Model V75-04 (Coulbourn Instruments, Pennsylvania, USA) with a band pass filter of 1000 Hz.

Randomization and data acquisition were controlled by custom-made stimulus presentation software Affect5 (Spruyt et al., 2010) through a Microsoft Windows 10 based PC (Intel® Core™ i7; 8GB ram; Radeon RT250) connected to a Dell P2014H 20” WXGA++ monitor.

## *2.8. Analysis*

Three main hypotheses are tested in this study: during acquisition we (1) hypothesize that after repeated presentation of CS+ followed by the US, the CS+ will elicit fear responses (as measured by expectancy ratings, SCR, and HR) while the CS-, never paired with the US, will not elicit these fear responses; (2) extinction of fear responses to the CS+ are expected after repeated presentation of CS- and CS+ but this time without the presentation of the US; and (3) reinstatement of fear is expected after a single presentation of US after extinction.

First, visual inspection of raw physiological data was used in order to exclude artefacts and carried out through PhysioData Toolbox software (Sjak-Shie, 2017) running on

MATLAB Release 2016a (The MathWorks, Inc., Natick, MA, USA). Next, multilevel mixed model analyses were used to assess each hypothesis by testing whether the conditioning procedure resulted in successful differential fear- and extinction learning as measured by expectancy ratings, SCR and HR responses. All multilevel mixed models were created using maximum likelihood modeling. Intercepts were allowed to vary randomly across participants. Random slopes were added in a second step to see whether this improves model fit. An AR1 variance-covariance matrix was added to see whether this improves model fit. The independent variable Trial, signifying trial number within each session, was group mean centered. CS type was dummy-coded, using CS- trials as the reference category. To account for possible non-linear learning rates, linear and loglinear time curves were fitted to all models and either of these variables were removed if this resulted in a better model fit according to BIC estimates. Alterations to the basic model (e.g. random slopes, loglinear time curve, AR1 matrix) are stated in the analysis when it resulted in an improved model.

Potential changes in ASRT responsiveness and sound tolerance levels (as measured by the LDL) were analysed through multilevel modeling in order to explore changes in LDL and ASRT over time (i.e. T0, T1 and T2). All analyzes were conducted through SPSS 26.0.

### 3. Results


Sixty-two participants (51 females and 11 males;  $M_{\text{age}} = 20.65$ ) aged between 18 and 25 years participated in the study (Table 1). All participants showed normal hearing thresholds ( $PTA_{\text{high}}$ ). Inspection of expectancy ratings revealed that 19 participants recorded higher US expectations for CS- than CS+ on the final trial (non-learners). Eight participants' SCR recordings were eliminated from analysis due to excessive artefacts (4 in the warble as CS+ condition and 4 in the pure tone as CS+ condition). As a result of a programming error, no physiological data was recorded during sections of the last two CS- trials of 9 participant and the last CS+ trial of 11 participants during the reinstatement phase.

**Table 1:** Summary of demographic characteristics, State Trait, Skin Conductance Response (SCR) non-responders, and Pure Tone Average of high frequencies ( $PTA_{\text{high}}$ ) for all participants per group.

	CS+ tone type	
	Warble (n = 30)	Pure (n = 32)
Mean age (SD)	20.83 (9.95)	20.47 (2.02)
Female (%)	26 (86.7)	25 (78.1)
Non-learners (%)	13 (43.3)	6 (18.7)
SCR Non-responders (%)	12 (46.2)	10 (35.7)
$PTA_{\text{high}}$ (SD)	5.00 (4.28)	6.61 (5.02)



**Figure 3:** Mean US expectancy ratings, heart rate acceleration, and skin conductance per phase for CS+ and CS-. Error bars: 95% confidence interval.



### 3.1. Acquisition

Participants reported significantly higher expectancy ratings for CS+ trials compared to CS- trials (Figure 3), as reflected by the main effect of CS type,  $b = 11.07$ ,  $t(101.93) = 3.92$ ,  $p < .001$ . Over the course of the acquisition phase, there was a main effect of Trial ( $b = -5.66$ ,  $t(336.90) = -5.08$ ,  $p < .001$ ) and a significant CS-type\*Trial interaction ( $b = 6.61$ ,  $t(343.24) = 4.12$ ,  $p < .001$ ).

Participants' heart rate acceleration did not reflect differential fear learning. Participants did not display stronger heart rate acceleration for CS+ trials compared to CS- trials ( $b = .01$ ,  $t(124.67) = .03$ ,  $p = .979$ ). While heart rate acceleration in response to CS presentation increased over time (Figure 3), as reflected by the main effect of Trial,  $b = .40$ ,  $t(236.73) = 2.10$ ,  $p = .037$ , this increase did not differ between CS+ and CS- trials (CS type\*Trial),  $b = -.08$ ,  $t(236.73) = -.31$ ,  $p = .755$ .

Participants' SCR did not reflect differential fear learning. Figure 3 suggests that participants did not display stronger SCR for CS+ trials compared to CS- trials (CS-type main effect), ( $b < -.01$ ,  $t(164.72) = -.58$ ,  $p = .562$ ). Participants' SCRs displayed significant habituation over time, as displayed by significant reduction of SCRs over time,  $b = -.01$ ,  $t(164.72) = -4.47$ ,  $p < .001$ , irrespective of CS type,  $b = -.00$ ,  $t(84.50) = .792$ ,  $p = .430$ .

### 3.2. Extinction

A random slope was added to the extinction model for expectancy ratings as this improved the model fit. US expectancy ratings indicated successful extinction of fear. Specifically, a main effect of trial indicated a general decrease in expectancy ratings over time,  $b = -7.65$ ,  $t(129.35) = -6.10$ ,  $p < .001$ . A steeper extinction curve was observed for the CS+ trials compared to CS- trials, which is supported by the significant CS-type\*Trial interaction ( $b = -3.10$ ,  $t(272.32) = -2.14$ ,  $p = .034$ ). A non-significant CS-type effect,  $b = 4.77$ ,  $t(83.55) = 1.82$ ,  $p = .073$ , indicated a lack of general distinction of safety (CS-) and threatening (CS+) probes.

Heart rate acceleration did not differ between CS-types ( $b = .31$ ,  $t(134.36) = .63$ ,  $p = .531$ ). Moreover, there was no decrease in acceleration over time, as seen by the lack of a main effect of Trial,  $b = -.01$ ,  $t(264.56) = -.047$ ,  $p = .963$ , for either CS-type (CS-type\*trial),  $b = .33$ ,  $t(264.56) = .96$ ,  $p = .340$ .

However, SCR extinction was confirmed through the main effect of Trial,  $b = -.01$ ,  $t(116.49) = -5.22$ ,  $p < .001$ , though there was no difference in the decrease, as evidenced by the interaction CS-type\*Trial,  $b = .00$ ,  $t(116.49) = 1.15$ ,  $p = .251$ . No discrimination between CS-type,  $b = .00$ ,  $t(40.87) = -.21$ ,  $p = .832$ , was observable.

### 3.3. Reinstatement

While slight increases in CS+ and CS- are observed when comparing the last trial of extinction with the first trial of reinstatement, no difference between CSs is confirmed. The expectancy ratings model for the reinstatement phase included loglinear time curve for better fit. A decrease of expectancy ratings over time, as indicated by a main effect of Trial ( $b = -60.69$ ,  $t(343.36) = -3.86$ ,  $p < .001$ ), though no difference in the decrease is observed between CS+ and CS- trials as the CS-type\*trial interaction failed to reach significance ( $b = -2.99$ ,  $t(343.36) = -1.35$ ,  $p = .893$ ). Furthermore, indiscriminate expectancy between CS-types is revealed by the lack of main effect of CS-type ( $b = 2.30$ ,  $t(281.49) = .398$ ,  $p = .691$ ).


Heart rate acceleration did not support reinstatement as reflected by the lack of Trial main effect ( $b = -.26$ ,  $t(260.34) = -.83$ ,  $p = .408$ ), with CS+ trials and CS- trials holding no differences over the duration of the phase (CS-type\*trial interaction:  $b = .03$ ,  $t(262.00) = .067$ ,  $p = .947$ ). Participants did not have different heart rate acceleration for CS+ or CS- trials, as seen by the results of the main effect of CS-type ( $b = .17$ ,  $t(126.23) = .26$ ,  $p = .793$ ).

Similar to expectancy ratings, a slight increase from the last trial of extinction and the first trial of reinstatement phase depicted in Figure 3 are observed, though no difference between CS+ and CS- is confirmed. Loglinear time curve was added to the basic SCR model to improve fit. Extinction after reinstatement is confirmed through the decrease of SCR through time, as seen by the main effect of Trial ( $b = -.01$ ,  $t(144.84) = -2.02$ ,  $p = .045$ ), although there was no difference in the decrease of CS+ trials and CS- trials, as seen by the CS-type\*Trial interaction ( $b < -.01$ ,  $t(145.09) = -.43$ ,  $p = .666$ ). Furthermore, there was no difference between CS-type ( $b < .01$ ,  $t(57.44) = .524$ ,  $p = .602$ ).

### 3.4. Loudness Discomfort Levels (LDL) and Stapedial Acoustic Reflex Thresholds (ASRT)

Over 70% of participants reach the maximum probe available for LDL at 125 Hz (60 dB) at all testing timepoints and so this frequency was eliminated from analysis. Figure 4 shows mean LDL at each measurement point for each frequency. Those who exceeded the maximum threshold (18.1% of all tests) were recorded at the next theoretical level: 105 dB for all frequencies except at 250 Hz (90 dB) and at 8 kHz (85 dB). Left and right LDL were averaged per participant and tested through multilevel modeling for an effect of time.

**Figure 4:** Mean Loudness Discomfort Levels (LDL) and mean Acoustic Stapedial Reflex thresholds (ASRT) at baseline (T0), after acquisition (T1) and after reinstatement (T2).



Nineteen participants (32%) did not record ASRT at least once at maximum threshold (100 dB). One participant did not present ASRT at any point. When ASRT was not detected at maximum level (13% of all tests) data was recorded at the next theoretical test threshold (105 dB). Figure 4 depicts mean for ipsilateral and contralateral ASRT measures at each time point.

Thresholds of LDL at 4 kHz (equal to the frequency of CS and US) increased at each measurement point (main effect of time estimates are depicted in Table 2). Moreover, this habituation pattern is generalized to LDL at all frequencies with the exception of 1 kHz. Specifically, the threshold of the right ear (first test conducted).

**Table 2:** Summary of results from multilevel mixed models for Loudness Discomfort Levels (LDL) at each frequency tested. The regression estimate (*b*) refers to main effect of time.

Frequency (Hz)	Estimate ( <i>b</i> )	df	<i>t</i>	<i>p</i>
250	.93	65.43	3.14	.003
500	1.33	62.78	3.55	.00
1 k	-.02	67.83	-.06	.952
2 k	1.19	66.54	3.62	.001
4 k	1.01	133.65	2.70	.008
8 k	.84	68.90	2.66	.010

Ipsilateral ASRT did not respond to time,  $b = -.24$ ,  $t(59.06) = -1.22$ ,  $p = .228$ , while contralateral ASRT decreased through time,  $b = -.50$ ,  $t(62.63) = -2.27$ ,  $p = .026$ .

## 4. Discussion

The FA model, long established in pain research, has been adapted to tinnitus in order to provide a framework for a better understanding of the development and maintenance of chronic disabling tinnitus. While pain-related fear learning research has supported the FA model, its application in the area of tinnitus is in its infancy. The development of a tinnitus-relevant fear learning paradigm with healthy participants may allow for underlying mechanisms to be explored. The current study introduced a novel paradigm relying solely on acoustic probes to study differential fear learning that is ecologically valid to chronic tinnitus patients. We hypothesized that after repeated presentation of CS+ followed by the US, the CS+ will elicit fear responses while the CS-, which is never paired with the US, will not elicit these fear responses. While fear expectancy corroborated the hypothesis, physiological responses indicated quick habituation (SCR) and indiscriminate fear learning (heart rate acceleration). Fear reduction was also investigated through the extinction of fear responses to the CS+ after repeated presentation of CS- and CS+ without US. Interestingly, extinction was observed in expectancy ratings and SCR to both CS+ and CS-, while heart rate acceleration indicated no extinction. Lastly, reinstatement of fear was expected after a single presentation of US after extinction. Similarly, to the extinction phase, expectancy ratings and SCR indicated indiscriminate fear reinstatement while heart rate accelerations remained high. Additionally, noise tolerance levels (LDL) increased at each timepoint, opposing ASRT which remained constant or decreased.

We found support for differential fear learning through US expectancy ratings at acquisition. Physiological measures diverged as heart rate acceleration indicated indiscriminate fear learning while habituation patterns were observed for SCR. Furthermore, indiscriminate extinction and reinstatement of fear were also confirmed

through expectancy ratings and supported by SCR. Indiscriminate fear learning was also confirmed through heart rate acceleration during acquisition whereas patterns observed during extinction and reinstatement did not support the fear reduction observed in expectancy ratings and SCR. Despite the lack of differential fear learning indicated by the physiological measures, previous research on expectancy ratings indicates its strength as an accurate measure of fear, holding face validity, diagnostic validity, predictive validity and construct validity (Boddez et al., 2013). Moreover, recording of expectancy ratings are less subjected to measurement errors. Small movement artifacts, electrode placement, room temperature or sweatiness of a participant's hand may affect SCR measurement. Quick habituation patterns, as observed in the study, are a further limitation of SCR ("Publication recommendations for electrodermal measurements," 2012). Similarly, heart rate acceleration may have been influenced by external factors such as walking between the soundproof audiometric rooms or water intake beforehand (Heathers et al., 2018).

The change of context for the measurement of ASRT and LDL might have played a significant role in the absence of differentiation of US expectancy ratings between CS+ and CS- trials at the start of the extinction phase, despite the observed differential fear learning observed during acquisition. Beyond the interruption of the task, the repeated stimuli presented by the ASRT and LDL tests may have served as an extinction of fear due to its similarity with the US (in frequency and/or intensity). Participants who reached the 100 dB probe limit in both LDL and ASRT tests, potentially lessened the negative valence of the US in the experiment. Thus, the fear measured subsequently, at extinction and reinstatement phases, may have been associated to the context rather than the CS+. Moreover, the relatively small number of trials may have not been enough for full awareness of the contingencies between CSs and US. It is important to note that a higher number of trials also signifies a larger burden for the auditory system, more specifically repeated high intensity (> 75 dB) sounds can be harmful to the inner ear hair cells (Services USDoHH, 1990; Ward et al., 1981). Further limitations may include the aversiveness of the US, which may have failed to elicit adequate UR in all participants, and the assumption of the CSs as neutral stimuli.

Exploratory analysis of sound tolerance levels (LDL) indicated increased habituation to loud sounds at each subsequent measuring point – after acquisition (T1) and after the end of the experiment (T2). An exception to this pattern was found in the first test conducted (1 kHz, right ear at baseline), which indicated an overestimation by participants. A second measure of the 1 kHz frequency during the first battery of tests would be advisable for

future research and clinical practice. Despite this exception, the habituation pattern was generalized to all frequencies and not specific to 4 kHz (the frequency of CSs and US). Congruent with the physiological responses, which appear to indicate indiscriminate fear learning. This is discordant to the findings of ASRT, which did not increase with time. In fact, contralateral ASRT decreased, indicating sensitization. Interestingly, the ASRT, thought to be a possible objective measure of sound tolerance, contrasted with the subjective measure of sound tolerance (LDL). The 5 dB stepped increases used in the current study may not have been sensitive enough to achieve a better understanding of ASRT. Further research on the responsiveness of ASRT to threat is warranted.

Our findings on sound tolerance are limited due to the lack of consistent fear learning effects and the use of 5 dB step increases, reducing measurement sensitivity. Further research with multimodal paradigms and more precise operationalization of ASRT and LDL may test these hypotheses and further elaborate on the role of ASRT in sound tolerance. Nevertheless, these findings indicate the strength of this novel tinnitus relevant paradigm. Participants successfully acquired fear responses to previously neutral acoustic stimuli by the presentation of the same stimuli at a higher intensity. By building the basis for future research to expand and improve on the current paradigm, tinnitus research may develop its own fundamental research that fully supports the FA model.

## References

- Aslaksen, P.M., Åsli, O., Øvervoll, M., Bjørkedal, E., 2016. Nocebo hyperalgesia and the startle response. *Neuroscience* 339, 599–607. <https://doi.org/10.1016/j.neuroscience.2016.10.040>
- Baguley, D., 2003. Hyperacusis. *JRSM* 96, 582–585. <https://doi.org/10.1258/jrsm.96.12.582>
- Boddez, Y., Baeyens, F., Luyten, L., Vansteenwegen, D., Hermans, D., Beckers, T., 2013. Rating data are underrated: Validity of US expectancy in human fear conditioning. *J. Behav. Ther. Exp. Psychiatry* 44, 201–206. <https://doi.org/10.1016/j.jbtep.2012.08.003>
- Burger, A.M., Verkuil, B., Van Diest, I., Van der Does, W., Thayer, J.F., Brosschot, J.F., 2016. The effects of transcutaneous vagus nerve stimulation on conditioned fear extinction in humans. *Neurobiol. Learn. Mem.* 132, 49–56. <https://doi.org/10.1016/j.nlm.2016.05.007>
- Cima, R.F.F., Crombez, G., Vlaeyen, J.W.S., 2011. Catastrophizing and Fear of Tinnitus Predict Quality of Life in Patients With Chronic Tinnitus. *Ear Hear.* 32, 634–641. <https://doi.org/10.1097/AUD.0b013e31821106dd>
- Davis, A., Refaie, A.E., 2000. Epidemiology of Tinnitus, in: Tyler, R.S. (Ed.), *Handbook of Tinnitus*. Singular Thompson Learning, San Diego, pp. 1–24.
- Gallus, S., Lugo, A., Garavello, W., Bosetti, C., Santoro, E., Colombo, P., Perin, P., La Vecchia, C., Langguth, B., 2015. Prevalence and Determinants of Tinnitus in the Italian Adult Population. *Neuroepidemiology* 45, 12–19. <https://doi.org/10.1159/000431376>
- Gilles, A., Van Hal, G., De Ridder, D., Wouters, K., Van de Heyning, P., 2013. Epidemiology of Noise-Induced Tinnitus and the Attitudes and Beliefs towards Noise and Hearing Protection in Adolescents. *PLoS One* 8, e70297. <https://doi.org/10.1371/journal.pone.0070297>
- Hall, D.A., Fackrell, K., Li, A.B., Thavayogan, R., Smith, S., Kennedy, V., Tinoco, C., Rodrigues, E.D., Campelo, P., Martins, T.D., Lourenço, V.M., Ribeiro, D., Haider, H.F., 2018. A narrative synthesis of research evidence for tinnitus-related complaints as reported by patients and their significant others. *Health Qual. Life Outcomes* 16, 1–15. <https://doi.org/10.1186/s12955-018-0888-9>
- Hallam, R.S., Rachman, S., Hinchcliffe, R., 1984. Psychological aspects of tinnitus. *Contrib. to Med. Psychol.* 31–53.
- Heathers, J., Quintana, D.S., Angus, D., Krygier, J., Kemp, A., de Rosnay, M., 2018. Water consumption as a source of error in the measurement of heart rate variability. <https://doi.org/https://doi.org/10.31219/osf.io/83exy>
- Jastreboff, P.J., 1990. Phantom auditory perception (tinnitus): mechanisms of generation and perception. *Neurosci. Res.* 8, 221–254. [https://doi.org/10.1016/0168-0102\(90\)90031-9](https://doi.org/10.1016/0168-0102(90)90031-9)
- Jastreboff, P.J., Brennan, J.F., Coleman, J.K., Sasaki, C.T., 1988. Phantom auditory sensation in rats: An animal model for tinnitus. *Behav. Neurosci.* 102, 811–822. <https://doi.org/10.1037/0735-7044.102.6.811>
- Kim, H.-J., Lee, H.-J., An, S.-Y., Sim, S., Park, B., Kim, S.W., Lee, J.S., Hong, S.K., Choi, H.G., 2015. Analysis of the Prevalence and Associated Risk Factors of Tinnitus in Adults. *PLoS One* 10, e0127578. <https://doi.org/10.1371/journal.pone.0127578>
- Kleinstäuber, M., Jasper, K., Schweda, I., Hiller, W., Andersson, G., Weise, C., 2013. The Role of Fear-Avoidance Cognitions and Behaviors in Patients with Chronic Tinnitus. *Cogn. Behav. Ther.* 42, 84–99. <https://doi.org/10.1080/16506073.2012.717301>
- Lethem, J., Slade, P.D., Troup, J.D., Bentley, G., 1983. Outline of a Fear-Avoidance Model of exaggerated pain perception--I. *Behav. Res. Ther.* 21, 401–8.



- Lonsdorf, T.B., Klingelhöfer-Jens, M., Andreatta, M., Beckers, T., Chalkia, A., Gerlicher, A., Jentsch, V.L., Drexler, S.M., Mertens, G., Richter, J., Sjouwerman, R., Wendt, J., Merz, C.J., 2019. Navigating the garden of forking paths for data exclusions in fear conditioning research. *Elife* 8, 1–36. <https://doi.org/10.7554/eLife.52465>
- Lonsdorf, T.B., Menz, M.M., Andreatta, M., Fullana, M.A., Golkar, A., Haaker, J., Heitland, I., Hermann, A., Kuhn, M., Kruse, O., Meir Drexler, S., Meulders, A., Nees, F., Pittig, A., Richter, J., Römer, S., Shiban, Y., Schmitz, A., Straube, B., Vervliet, B., Wendt, J., Baas, J.M.P., Merz, C.J., 2017. Don't fear 'fear conditioning': Methodological considerations for the design and analysis of studies on human fear acquisition, extinction, and return of fear. *Neurosci. Biobehav. Rev.* 77, 247–285. <https://doi.org/10.1016/j.neubiorev.2017.02.026>
- McCormack, A., Edmondson-Jones, M., Somerset, S., Hall, D.A., 2016. A systematic review of the reporting of tinnitus prevalence and severity. *Hear. Res.* 337, 70–79. <https://doi.org/10.1016/j.heares.2016.05.009>
- McKenna, L., Handscomb, L., Hoare, D.J., Hall, D.A., 2014. A Scientific Cognitive-Behavioral Model of Tinnitus: Novel Conceptualizations of Tinnitus Distress. *Front. Neurol.* 5, 1–15. <https://doi.org/10.3389/fneur.2014.00196>
- Meulders, A., 2020. Fear in the context of pain: Lessons learned from 100 years of fear conditioning research. *Behav. Res. Ther.* 131, 103635. <https://doi.org/10.1016/j.brat.2020.103635>
- Publication recommendations for electrodermal measurements, 2012. *Psychophysiology* 49, 1017–1034. <https://doi.org/10.1111/j.1469-8986.2012.01384.x>
- Services USDoHH (Ed.), 1990. Noise and Hearing Loss. Consensus Development Conference Statement., in: National Institutes of Health. Bethesda, MB.
- Sjak-Shie, E.E., 2017. PsysioData Toolbox.
- Sjouwerman, R., Lonsdorf, T.B., 2018. Latency of skin conductance responses across stimulus modalities. *Psychophysiology* 56, 1–11. <https://doi.org/10.1111/psyp.13307>
- Spruyt, A., Clarysse, J., Vansteenwegen, D., Baeyens, F., Hermans, D., 2010. Affect 4.0: A free software package for implementing psychological and psychophysiological experiments. *Exp. Psychol.* 36-45.
- Van Diest, I., Bradley, M.M., Guerra, P., Van den Bergh, O., Lang, P.J., 2009. Fear-conditioned respiration and its association to cardiac reactivity. *Biol. Psychol.* 80, 212–217. <https://doi.org/10.1016/j.biopsycho.2008.09.006>
- Vlaeyen, J.W.S., Linton, S.J., 2012. Fear-avoidance model of chronic musculoskeletal pain: 12 years on. *Pain* 153, 1144–1147. <https://doi.org/10.1016/j.pain.2011.12.009>
- Vlaeyen, J.W.S., Linton, S.J., 2000. Fear-avoidance and its consequences in chronic musculoskeletal pain: a state of the art. *Pain* 85, 317–32.
- Ward, J., Vella, C., Hoare, D.J., Hall, D.A., 2015. Subtyping somatic tinnitus: A cross-sectional UK cohort study of demographic, clinical and audiological characteristics. *PLoS One* 10, 1–11. <https://doi.org/10.1371/journal.pone.0126254>
- Ward, W.D., Duvall, A.J., Santi, P.A., Turner, C.W., 1981. Total Energy and Critical Intensity Concepts in Noise Damage. *Ann. Otol. Rhinol. Laryngol.* 90, 584–590. <https://doi.org/10.1177/000348948109000615>



## CHAPTER 3

---

# Effects of Ecological Momentary Assessment (EMA) induced monitoring on tinnitus experience: a multiple-baseline single-case experiment

Published as:

Lourenco, M., Cima, R. F. F., & Vlaeyen, J. W. S. (2021).

Effects of Ecological Momentary Assessment (EMA) induced monitoring on tinnitus experience:  
A multiple-baseline single-case experiment. *Prog Brain Res*, 263, 153-170.  
<https://doi.org/10.1016/bs.pbr.2021.04.009>

## Abstract

**Introduction:** Ecological momentary assessment (EMA) is a method capable of assessing tinnitus experience throughout the day, enabling the exploration of daily dynamic changes of tinnitus expression. However, the effects on patients' tinnitus experience itself are still largely unknown. This study seeks to test the hypothesis that the use of EMA negatively influences tinnitus experience in participants with severe tinnitus. **Method:** A multiple-baseline single-case experimental design included four severely affected tinnitus volunteers who were recruited online and randomized into different phasing schedules. Baseline phase (A) ranged from 11-24 days, followed by an EMA phase (B) for the remainder of the 33-day schedule. End-of-day diary assessments of tinnitus experience (e.g. annoyance, intrusiveness, mood) were visually inspected, and complemented with inferential statistics (randomization tests and Tau-U). **Results:** End-of-day diary data revealed no change in broadened median between phases. Nevertheless, tinnitus experience scores improved as variability decreased and a significant improvement in stress was observed through weighted Tau-U statistics. **Conclusion:** Findings of this study corroborate that EMA assessment does not negatively affect tinnitus experience. On the contrary, participants may have improved. The underlying mechanism of improvements are still to be uncovered. Findings are limited to severely affected tinnitus sufferers at present.


## 1. Introduction

Tinnitus is defined as the perception of sound(s) (e.g. ringing or buzzing) in the ear or head without a detectable corresponding physical source. While nearly 20% of the adult population reports tinnitus, it is only a small subset (1-6%) who suffer from it, experiencing severe distress and disturbances in numerous aspects of daily life (Davis and Refaie, 2000; Cima, Crombez and Vlaeyen, 2011; Kim *et al.*, 2015; Bhatt, Lin and Bhattacharyya, 2016; McCormack *et al.*, 2016; Hall *et al.*, 2018). There is currently no cure for chronic tinnitus and, while treatment options are varied, Cognitive Behavioural Therapy (CBT) approaches are strongly recommended for treatment (Cima *et al.*, 2019) with a recent Cochrane review emphasizing the positive effect it has on quality of life (Fuller *et al.*, 2020).

The Fear Avoidance (FA) model of chronic pain (Lethem *et al.*, 1983; Vlaeyen and Linton, 2000, 2012) provides the underpinnings for CBT by predicting the development, maintenance and increase of chronic pain disability, and has successfully been applied to tinnitus as well (Cima, Crombez and Vlaeyen, 2011; Kleinstäuber *et al.*, 2013). The model states that a pathological cycle may start with catastrophic misinterpretations about the tinnitus perception that feed into tinnitus-specific fear, heightened attention and constant monitoring, avoidance of stimuli that are expected to increase tinnitus, which in turn increases disability, distress, and decreased quality of life (Figure 1). In an alternative to the detrimental spiral, tinnitus may be perceived as a common and harmless sensation, leading to continuation of valued activities, thereby confronting stimuli that may increase tinnitus (e.g. exposure) and eventual recovery. Research has supported the negative role of tinnitus catastrophizing, fear and attention in regards to quality of life (i.e. disability; Cima, Crombez and Vlaeyen, 2011). Beyond the tinnitus field, exposure techniques have been long supported in literature and are considered the golden standard for their effectiveness in reducing psychological distress associated with other chronic conditions, such as pain (Vlaeyen *et al.*, 2001; Woods and Asmundson, 2008; e.g. Hedman-Lagerlöf *et al.*, 2018) and anxiety disorders (Meuret *et al.*, 2012; e.g. Carpenter *et al.*, 2018). Exposure techniques may take different forms but rely on the repeated confrontation with the fear-provoking stimuli. Simply put, fear and catastrophic thoughts are reduced by confronting the patient with the distressing experience (e.g. tinnitus) without the anticipated negative outcome (e.g. loss of control), thus violating the expected prediction (Craske *et al.*, 2014; for a detailed review on underlying mechanism of inhibitory learning and exposure see Craske, Hermans and Vervliet, 2018). Despite

empirical support, mechanisms behind these exposure components of CBT and, more specifically, the role of attention to tinnitus are still debated, with competing techniques and therapies recommended in standard practice (e.g. masking, sound enrichment therapy, TRT). Controversies also exist about the rise of novel assessment methodologies (e.g. Ecological Momentary Assessment) and technologies (e.g. TrackYourTinnitus app, Schlee *et al.*, 2016) that aim to capture tinnitus fluctuations by repeatedly assessing participants throughout the day.

**Figure 1:** Fear Avoidance model (replicated from Cima, van Breukelen and Vlaeyen, 2018)




Ecological Momentary Assessment (EMA) has been under scrutiny for potentially increasing attention to tinnitus (e.g. Henry *et al.*, 2012; Schlee *et al.*, 2016). Worries exist that the repeated daily measurements will lead to increase in awareness of tinnitus with detrimental effects on patients and their disability due to tinnitus. EMA attempts to capture in-the-moment experiences in real-life situations by randomly delivering questions throughout the day, thereby avoiding common limitations of standardized self-report questionnaires such as possible fallacies of recalling and reconstructing experiences (Stone and Shiffman, 1994). EMA questionnaires are designed to be rapidly answered and items may focus on different dimensions of an individual's life, including well-being (e.g. happiness, stress, sleep), daily activities (e.g. current location, social contact, environmental noise), and tinnitus experience (e.g. annoyance, anger, distraction). Technological advances and the rising use of smartphones has enabled EMA to be

commonly delivered through purpose-built apps, increasing its research applications. Despite the growing use of EMA the effects of it on tinnitus experience are still widely unknown.

### *1.1. Previous research*

Henry et al. (2012), the first to look into the potential effects of EMA on tinnitus subjects, used personal digital assistants (PDAs) and 24 participants who underwent a 2-week long EMA phase. Reactivity to EMA was analysed through pre- and post-EMA group mean measurements of the Tinnitus Handicap Inventory (THI; Newman, Jacobson and Spitzer, 1996; Newman, Sandridge and Jacobson, 1998), where an observed worsening was not statistically significant. Moreover, the researchers evaluated the individual trajectories of participants' EMA responses by plotting and visually inspecting the daily scores over time. The authors observed a high degree of intra- and inter-individual differences and nearly half of the participants (54.2%) were classified as “consistent” where no clear trend could be visually fitted. Six (25.0%) participants were observed to have an “improving trend” while the remaining five participants (20.8%) were found to “worsen” during the EMA phase. The lack of statistical support prevents a more accurate trend fitting. Furthermore, the absence of a control condition (i.e. baseline phase) hinders the interpretability of the data. For instance, an “improving trend” may be deceiving, as baseline data could have revealed a steeper improvement trend before the introduction of EMA (Figure 2). Beyond these limitations, the interpretation of the individual data sheds light into the group level analysis. The stipulated improvement and worsening of individual participants within the group may cancel each other out when analysing group means, and true effects might therefore remain undetected. In other words, EMA may have significant negative or positive effect on a particular subset of participants.

**Figure 2:** Example of possible baseline trends (A) prior to intervention phase (B) with higher distress levels reflected through higher scores on y-axis. From top to bottom graphs respectively indicate worsening, improving and no change after intervention onset.



More recently, Schlee et al. (2016) investigated the potential effects of EMA on tinnitus loudness and distress. Data gathered online through an EMA delivery app for smartphones was analysed. From 857 volunteers who participated in data collection, linear regression was applied with participants who used the app for at least one month ( $n=66$ ). No significant association between EMA use and tinnitus loudness or distress was observed. The lack of reactivity was further supported by a different group-level analysis in which the average first and last five assessments were compared through a t-test. The same test was conducted for the remainder of participants who had completed the EMA from anywhere between 5 and 30 days. While no significant change was found using group level analysis, the authors observed high intra- and inter-individual differences, congruent with findings from Henry et al. (2012). While linear regression analysis is a methodological improvement to visual inspection of trend, the reliance on group level analysis continues to limit the interpretability of data as participants high variability in scores at the individual level may offset each other's when combining the data.



## 1.2. Single-Case Experimental Design (SCED)

As group studies fail to acknowledge the idiosyncratic nature of tinnitus, individualized research designs offer an alternative that thrives on such variability. Previous group-based research has supported lack of reactivity in scores of tinnitus handicap, distress and loudness to EMA, but it has also emphasized the high degree of divergence in tinnitus experience. Such individual variability remains hidden when interpreting results from group level analysis, possibly omitting significant individual effects, thereby limiting the generalizability of group-findings to a specific patient's profile. As such, the conclusion that tinnitus sufferers may lack EMA reactivity excludes subgroups (e.g. severe tinnitus sufferers) who could potentially benefit or worsen from its utilization.

In order to bridge the gap in knowledge about the effects of employing EMA, the current study uses a Single-Case Experimental Design (SCED) to investigate whether EMA influences tinnitus experience. Different from non-experimental designs (e.g. case-studies), SCED relies on repeated measurements of the dependent variable (e.g. tinnitus experience) over time during at least two different levels of the independent variable (e.g. with/without EMA) (Morley, 2018). Recent guidelines (i.e. Tate *et al.*, 2016) have emerged as the result of a growing interest in SCEDs. Alternatively to group-based designs and congruent with the push for individualized treatment and research (Schork, 2015), SCEDs can equally investigate causal relationships between variables (i.e. EMA and tinnitus experience). By repeatedly assessing individuals before (baseline phase: A) and after the introduction of an intervention or manipulation (phase: B), usually at a random starting point within a predefined time window, changes between phases' level, trend and variability are inspected for each participant and statistically analysed (i.e. randomization tests, Tau-U). Furthermore, the multiple observations at baseline create robust control conditions, despite the high intra- and inter-individual variance, which allows for addressing the limitations of previous research.

Building on previous findings, a specific subgroup is analysed in this study. The FA model illustrates the role of constant monitoring and unwarranted attention to the tinnitus percept and its pathological progression. As higher levels of attention towards tinnitus haven been associated with higher degrees of tinnitus severity (e.g. lower quality of life; Cima, Crombez and Vlaeyen, 2011), it is hypothesized that severe tinnitus sufferers respond negatively to the use of EMA (e.g. increased tinnitus annoyance, increased stress levels, decreased sleep quality).

## 2. Method

### 2.1. Study design

An AB multiple-baseline SCED where baseline phase (A) always precedes an experimental phase (B) was employed. Within the different SCED possibilities, the utilization of a multiple-baseline design across participants limits confounding factors by requiring different individuals to undergo an AB schedule with randomized B phase onset.

End-of-day diary assessments were continuously gathered throughout both phases, while during phase B, EMA was added, which included real-time assessment, equivalent to those assessed in the diary. Data was gathered through a purpose-built app (TinNotes) developed in-house by Maastricht University's Instrumentation Engineering department. TinNotes ran on iOS devices and notified participants daily (at 8pm) for the completion of the diary assessment and, during phase B, also delivered daily EMA at 7-random time points with at least 30-minutes in-between prompts. The participant was able to set a sleeping schedule and to snooze EMA prompts twice for 5-minutes each time, after which the assessment was cancelled. Data was sent directly to Maastricht University's servers when the device was connected to the internet. Six different AB schedules were randomly determined through a Single Case Data Analysis app (SCDA; <https://tamalkd.shinyapps.io/scda/>). Five consecutive observation points were considered minimally necessary per phase, and 23 potential phase B (EMA introduction) onset points were available, totaling 33 potential diary assessments per participant. According to the regulated randomization principle (Koehler and Levin, 1998), after the six randomized schedules were created, each participant was randomly assigned to a schedule and unaware of phase B onset day. The study was pre-registered (Lourenco, Cima and Vlaeyen, 2019) and approved by Maastricht University's Ethical Review Committee Psychology and Neuroscience (ERCPN-204\_23\_02\_2019). Reporting of the study follows the guidelines established at SCRIBE (Tate *et al.*, 2016).

### 2.2. Participants

Recruitment took place through an advertisement published in the newsletter of the Dutch national tinnitus patient association (Stichting Hoormij). The research was also announced at the annual symposium of the association. Compensation for participants who completed the study was 75€ (seventy-five euros). Inclusion criteria included: (1) own and use an iPhone; (2) self-reported tinnitus; (3) not currently undergoing treatment for tinnitus or psychological, psychiatric or any other kind of therapy

addressing psychological, social, emotional, and or behavioral problems; (4) absence of severe anxiety or depression; (5) able to read and write in Dutch. Participants were excluded if: (1) the criteria for severe tinnitus was not met, as measured by the Tinnitus Questionnaire (TQ > 59; Hallam, Jakes and Hinchcliffe, 1988), or (2) met the criteria for severe Anxiety or Depression, assessed through the Hospital Anxiety and Depression Scale (HADS-A > 14 or HADS-D >14; Zigmond and Snaith, 1983). Participants were blinded to the study's objective and were invited to test a new tinnitus tracking app. After data collection ended participants were debriefed to the true objective of the study.

Participants who completed the screening online and met inclusion criteria (n=175) were systematically contacted to participate. The first six participants who filled out the survey at T0 were randomly allocated to an AB schedule (Table 1). Participants were instructed to download the app and login via a unique code that allowed the app to deliver the specific AB schedule. Participants' codes were encrypted, and the key was maintained in a secure server.

**Table 1:** Six randomly created schedules where baseline (A) and EMA phase (B) required a minimum of five consecutive observations per phase. PP = participant.

	Observation																																
	1	...	10	11	12	13	14	15	16	17	18	19	20	21	22	23	24	...	33														
<b>PP 1</b>	A	...	A	A	A	A	A	A	A	A	A	A	A	A	A	A	B	...	B														
<b>PP 6</b>	A	...	A	A	A	A	A	A	A	A	A	A	A	B	B	B	B	...	B														
<b>PP 2</b>	A	...	A	A	A	A	A	A	A	A	A	B	B	B	B	B	B	...	B														
<b>PP 3</b>	A	...	A	A	A	A	A	A	A	A	B	B	B	B	B	B	B	...	B														
<b>PP 4</b>	A	...	A	A	A	A	B	B	B	B	B	B	B	B	B	B	B	...	B														
<b>PP 5</b>	A	...	A	B	B	B	B	B	B	B	B	B	B	B	B	B	B	...	B														

## 2.3. Measurements

### 2.3.1. Primary outcomes

The end-of-day diary assessments were the primary study endpoints gathered by the TinNotes app. Compiled of 15 questions on a 7-point Likert scale and one qualitative question, the diary items were extracted and adapted from existing questionnaires (e.g. TFI, TQ) by a group of four specialists which included researchers and therapists. The questions were delivered in a different order every day of the week. Each question assessed a different aspect of tinnitus experience, including avoidance, annoyance, intrusiveness, invasiveness, fear, sadness, pleasantness, distraction, masking and anger. It also included questions about overall well-being, assessing happiness, feeling stressed, sleep quality,

activity level, anxiousness and social interaction (qualitative). Treatment fidelity was assessed through the completion rate of EMA, with at least a 75% completion rate required.

### 2.3.2. Secondary outcomes

Standardized questionnaires were collected before baseline (T0) and after EMA (T1) phases.

Tinnitus Functional Inventory (TFI; Meikle et al., 2012): The TFI is a 25-item self-report measure of impairment in daily functioning. Respondents use 10-point Likert scales to indicate what tinnitus related experiences they have had over the previous week. The TFI has subscales on intrusiveness; reduced sense of control; cognitive interference; sleep disturbance; auditory difficulties attributed to tinnitus; interference with relaxation; reduced quality of life; and emotional distress. A total score, with a maximum of 100 can be calculated with higher scores reflecting greater levels of interference in daily activities. The TFI has excellent psychometric properties and was specifically designed to be used as an outcome measure in clinical trials with a 13 point change deemed as clinically significant (Meikle *et al.*, 2012). A Dutch version of the TFI was recently developed and validated for use with Dutch speaking patients (Rabau, Wouters and Van de Heyning, 2014).

Tinnitus Questionnaire (TQ; Hallam, Jakes and Hinchcliffe, 1988; Meeus, Blaivie and Van de Heyning, 2007): The TQ is a 52-item measure of tinnitus-severity. Each question is rated on a three-point scale and assesses psychological distress associated with tinnitus. The TQ is widely used in tinnitus research (Hall *et al.*, 2016) and has good psychometric properties (Fackrell *et al.*, 2014). A Dutch version has been developed and validated for use (Meeus, Blaivie and Van de Heyning, 2007).

Fear of Tinnitus Questionnaire (FTQ; Cima, Crombez and Vlaeyen, 2011): The FTQ is a 17-item self-report measure of a person's feared outcomes of living with subjective tinnitus. A higher total score on the FTQ is associated with higher levels of interference in daily activities of living. A recent analysis of the psychometric properties reported that it is a reliable and valid measure (Fuller *et al.*, 2019) and has demonstrated that it is responsive to clinical change (Cima *et al.*, 2012).

Tinnitus Catastrophizing Scale (TCS; Cima, Crombez and Vlaeyen, 2011): adapted from the 13-item Pain Catastrophizing Scale (Sullivan, Bishop and Pivik, 1995) for use with

tinnitus patients – the word “pain” being replaced by the word “tinnitus”. Participants are asked to respond on a five-point Likert-type scale (0 = Not at all; 1= to a small extent; 2 = to some extent; 3 = to a large extent; 4 = always) to statements describing thoughts and feelings that might be related to their tinnitus. Higher levels of catastrophizing as measured by the TCS has been found to be closely related to poorer quality of life (Cima, Crombez and Vlaeyen, 2011).

Hospital Anxiety and Depression Scale (HADS; Zigmond and Snaith, 1983): The HADS is a widely used scale to screen for depression and anxiety. There is a total of 14-items with a 4-point Likert scale. Each subscale has a possible score of 21, with higher scores indicating higher levels of anxiety (HADS-A) and depression (HADS-D). The scale was used for screening with scores over 14 for any of the subscales indicating the presence of severe depression and/or anxiety.

Demographic and tinnitus characteristics data were gathered through the ESIT-SQ (Genitsaridi *et al.*, 2019): developed by the Tinnitus Research Initiative (TRI) and the European School for Interdisciplinary Tinnitus research (ESIT), the Screening Questionnaire (ESIT-SQ) is a comprehensive self-report tool for healthy individuals and tinnitus patients that provide demographic and multidisciplinary information on tinnitus relevant variables. Translated into six languages, this tool offers a standardized assessment for tinnitus research across Europe.

### 2.3.3. Analysis

Each item of the end-of-day diary was analysed per participant. For the purpose of this study the end-of-day diary items on social interaction and activity level were removed from analysis as they are treatment specific goals of CBT and do not pertain to this study’s main hypothesis. Visual inspection of annoyance and stress items were selected to illustrate findings with the remaining individual visual analysis presented as supplemental material.

The end-of-day diary scores were plotted over time. Visual inspection of the data was carried out in order to determine changes in level and variability. The broadened median (Rosenberger and Gasko, 1983) was utilized due to its resistance to outliers and plotted for change in level inspection. Visual inspection of variability was aided by including range lines per phase. Randomization tests were calculated for each participant with a combined p-value calculated (Onghena and Edgington, 2005). Monte Carlo sampling

(1000) was used with the test statistic defined as  $\bar{A}-\bar{B}$  due to the dependent variables' projected increase (worsening) after EMA introduction, while inverted items' (happiness, tinnitus pleasantness and sleep quality) test statistics was defined as  $\bar{B}-\bar{A}$ . Visual analysis and randomization tests were carried out through a purpose-built web-application which provides an interface for the shiny SCDA software, which utilizes the R packages SCRT, SCVA and SCM (<https://tamalkd.shinyapps.io/scda/>; Bulté and Onghena, 2013). Tau-U, a more conservative analysis of non-overlap statistics, was utilized to detect and correct for baseline trend when comparing phases. Tau-U enables confidence intervals and p-values to be calculated by following the "S" sampling distribution (Parker *et al.*, 2011). The web-application 'Single Case Research: web based calculators for SCR analysis (Version 2.0)' (Vannest *et al.*, 2016) was utilized for Tau-U calculations.

### 3. Results

Data was retrieved from the server after the completion of the study. An unknown error caused the app to malfunction for two participants, from whom the data was not included for analysis. One participant completed less than half of the EMA delivered (44.8%) compromising treatment fidelity and thus the data was not included for analysis. Three participants completed the study with over 83.3% and 91.7% compliance rate on end-of-day diary and EMA respectively. Participants who completed the study had been allocated to schedules 1 (PP 1), 2 (PP 6) and 4 (PP 3). Participants' demographic and tinnitus characteristics are presented in Table 2.

**Table 2:** Demographic and tinnitus characteristics

	Age	Sex	HADS		Duration	Location	Tinnitus	
			A	D			Daily fluctuation	Type
<b>PP 1</b>	55	Female	10	8	5 months	Both ears, worse in left	Stable	High pitch tone
<b>PP 6</b>	49	Female	12	9	10 years	Both ears, worse in left	Sometimes fluctuates	High pitch tone
<b>PP 3</b>	66	Male	7	4	12 years	Right ear	Stable	High pitch tone

### 3.1. Diary assessments

Visual inspection of the end-of-day diary scores for items on tinnitus annoyance and stress levels are presented in figure 3. P-values for the combined randomization tests and weighted Tau-U analysis are presented in Table 3 (individual Tau-U tables are presented as supplemental material). No statistically significant worsening was found by randomization tests, while Tau-U analysis revealed a statistically significant improvement in stress levels.

**Table 3:** P-values for combined Randomization Tests (RT) and weighted Tau-U analysis per item.


Item	RT	Tau-U
Anger	.767	.588
Annoyance	.521	.677
Anxiety	.603	.297
Avoidance	.486	.262
Distraction	.541	.792
Fear	.496	.435
Happiness†	.228	.549
Interference	.640	.683
Invasiveness	.514	.533
Pleasantness†	.220	.282
Sadness	.534	.335
Sleep quality†	.147	.644
Stress	.112	< .001**

†inverted items (test statistic =  $\bar{B} - \bar{A}$ )

\*p-value < .05

\*\*p-value < .001

**Figure 3:** Visual analysis of multiple-baseline AB-design of end-of-day diary for participants PP 1, PP 6 and PP 3 respectively showing level of each phase using broadened median scores and variability of each phase using range lines. Range lines are omitted when concur with broadened median.





### 3.2. Standardized assessments

Change in standardized questionnaires were calculated for each participant (Table 4). No participant had clinically meaningful improvement or worsening according to the TFI. Analyses of the TFI sub-scales (Table 5) revealed that in participant 6 (PP 6) overall worsening was mainly associated with decreased sleep quality (SI), though visual inspection of diary data for sleep quality revealed no shift in level and variability.

Table 4: Scores before baseline (T0), after EMA phase (T1) and change ( $\Delta$ ) of the Tinnitus Questionnaire (TQ), Tinnitus Functional Index (TFI), Tinnitus Catastrophizing Scale (TCS), and the Fear of Tinnitus Questionnaire (FTQ).

	TQ			TFI			TCS			FTQ		
	T0	T1	$\Delta$	T0	T1	$\Delta$	T0	T1	$\Delta$	T0	T1	$\Delta$
PP 1	68	60	-8	48.8	40	-8.8	37	20	-8	12	10	-2
PP 6	78	72	-6	67.6	80	12.4	31	20	-11	14	11	-3
PP 3	66	71	5	58	63.6	5.6	37	42	5	10	10	0

Table 5: Score change ( $\Delta$ ) of the Tinnitus Functional Index (TFI) sub-scales: Intrusive (I), Sense of Control (Sc), Cognitive (C), Sleep (SI), Auditory (Au), Relaxation (R), Quality of Life (Q), and Emotional (E).

	TFI								
	$\Delta I$	$\Delta Sc$	$\Delta C$	$\Delta SI$	$\Delta Au$	$\Delta R$	$\Delta Q$	$\Delta E$	
PP 1	13.33	-10	-13.33	-20	10	-3.33	-25	-16.66	
PP 6	-10	-3.33	-6.66	60	20	16.66	17.5	3.33	
PP 3	33.33	16.66	-20	-10	10	-13.33	15	10	

## 4. Discussion

The aim of this study was to investigate whether tinnitus monitoring induced by the use of EMA negatively affects overall tinnitus experience. In order to mitigate previous research limitations (e.g. lack of control condition), the present study employed a SCED with participants suffering from severe tinnitus.

Change in tinnitus experience was primary assessed through visual inspection of end-of-day diary data, showing no meaningful and consistent shift in level in any variable. Visual inspection also revealed that while one participant's (PP 1) scores at baseline presented a floor effect on the item regarding tinnitus pleasantness, all other items and participants presented adequate variation of scores and patterns that did not show floor

and ceiling effects. This lack of negative reactivity of tinnitus experience to EMA was further confirmed through randomization tests, which rendered no significant change for each variable. While level remained similar between phases, improvement was observed through variability decrease in all variables for at least two participants with the exception of tinnitus invasiveness and pleasantness. Decrease in variability of answers has been previously reported in literature and it is not attributed to instrumental effects (Vachon *et al.*, 2016). Surprisingly, Tau-U analysis indicated a significant decreased stress-levels after EMA introduction. Overall, EMA may have had a positive effect on participants' experience by increasing awareness of their current state during the day and allowing them to rate it on a scale. Such a monitoring exercise may lead to a more accurate reflection of overall daily experiences, which is then reported on the end-of-day diary. Furthermore, in the parallel field of chronic pain, monitoring of pain sensations have previously demonstrated potential long-term benefits when compared to distraction techniques for patients (Nouwen *et al.*, 2006) and highly fearful individuals (Roelofs *et al.*, 2004). These findings also fit the clinical benefits of exposure, in which increased tinnitus awareness is evoked to change threat-expectancies, leading to decreased safety-seeking and fear-responding.

Closer analysis of Tau-U calculations revealed that one participant's (PP 3) stress improvement outweighed other participants' lack of changes. More specifically, the change found may have been heavily influenced by changes in level and variability across most items of that participant observed in the one week. Due to the number of observations and the robust baseline established, caution in interpretation of the effects is warranted in that improvements reported during observations 23-27 of PP 3 might be due to external influences.

Standardized outcomes (TQ and TFI) revealed no clinically significant changes. Moreover, one participant's (PP 6) TFI score increase was mainly attributed to worsened quality of sleep, though diary data showed that there was no change in level or variability when comparing phases. Consequently, sleep quality worsening is not likely due to EMA. The remaining assessments (TQ, TCS and FTQ) of the participant indicated marginal improvements, which may also be observed in most items of the end-of-day diary.

Despite the dissonant result on the TFI, special consideration must be given to its interpretation. The TFI was developed to assess clinical change and is thus favoured over the TQ (Jacquemin *et al.*, 2019). Nevertheless, as with other self-report measures of subjective experiences, the TFI is susceptible to fallacies such as memory recall and

reconstruction. Retrospective self-reports may be influenced by biases in reconstruction of events according to the individual's own beliefs, behaviours or knowledge acquired after the event, as well as the current emotional state and physical location at the time of assessment (Stone and Shiffman, 1994; Kahneman *et al.*, 2004). Highly variable experiences such as tinnitus, further increase the burden of assessment, challenging individuals to quantify (e.g. average) it over a longer time period. The lack of convergence between the TFI and other assessments – for PP 6 – highlights the possible discrepancies that can result from retrospective self-reports of highly variable and subjective experiences (i.e. tinnitus). Novel approaches were developed to tackle these limitations, and as such this study favours end-of-day diary use, which decreases the timeframe of recall, therefore, reducing the risk of bias.

Tinnitus fear and catastrophizing, as measured by the FTQ and TCS respectively, revealed no meaningful negative change. According to the FA model, increase at any point of the pathological cycle strengthens the negative experience of tinnitus. A hypothetical monitoring effect induced by EMA must not exceed the current monitoring level of severely affected tinnitus sufferers. Interestingly, two participants demonstrated a decrease in tinnitus fear and catastrophizing. Exposure techniques reduce fear and catastrophic thoughts by repeatedly confronting the individual with the distressing experience without the expected negative outcome. Although speculative, the EMA-induced monitoring may have increased the number of instances where violation of expectations occurred (i.e. exposure). While the reported changes in fear and catastrophic misinterpretations may not be considered meaningful at this point, findings may pave the way for future research on the underlying mechanism of potential EMA-induced improvements.

At the time of the study, the app was limited to iOS devices only. Research on the differences between Android and iOS users in a tinnitus population have been previously conducted. Pryss *et al.* (2018) found small but significant differences in age and tinnitus duration. Android users were found to be slightly older and perceived tinnitus for longer when compared to iOS users. However, meaningfulness of previous findings is based on group-level analysis and therefore limited for the current study. Although different operational systems for smartphones are not expected to have an effect on the findings, future research which includes Android users is warranted.

The multiple-baseline SCED employed in the current study made it possible to create control conditions (i.e. baseline), which was lacking in previous research. Moreover, the minimum number of participants recommended for a multiple-baseline AB design was

reached (Kratochwill *et al.*, 2013). Only a subset of tinnitus sufferers was selected, and the focus on severe tinnitus is considered a strength of this research, which aims at untangling previous findings in the field by following novel standards for individualized medicine and research (Schork, 2015).

## **5. Conclusion**

The present study corroborates and expands on previous findings regarding EMA reactivity in tinnitus sufferers (Henry *et al.*, 2012; Schlee *et al.*, 2016). Inter- and intra-individual tinnitus experience variability is narrowed in the present study by including only severe tinnitus sufferers using single-case methodology. These participants were not observed to have meaningful negative reactions to EMA utilization. Contrary to expectations, slight improvements after EMA onset were observed. The underlying mechanism of the EMA-induced improvements are still to be uncovered.

## References

- Bhatt, J. M., Lin, H. W. and Bhattacharyya, N. (2016) 'Prevalence, Severity, Exposures, and Treatment Patterns of Tinnitus in the United States', *JAMA Otolaryngology–Head & Neck Surgery*, 142(10), p. 959. doi: 10.1001/jamaoto.2016.1700.
- Bulté, I. and Onghena, P. (2013) 'The Single-Case Data Analysis Package: Analysing Single-Case Experiments with R Software', *Journal of Modern Applied Statistical Methods*, 12(2), pp. 450–478. doi: 10.22237/jmasm/1383280020.
- Carpenter, J. K. *et al.* (2018) 'Cognitive behavioral therapy for anxiety and related disorders: A meta-analysis of randomized placebo-controlled trials', *Depression and Anxiety*, 35(6), pp. 502–514. doi: 10.1002/da.22728.
- Cima, R. F. F. *et al.* (2012) 'Specialised treatment based on cognitive behaviour therapy versus usual care for tinnitus: a randomised controlled trial', *The Lancet*, 379(9830), pp. 1951–1959. doi: 10.1016/S0140-6736(12)60469-3.
- Cima, R. F. F. *et al.* (2019) 'A multidisciplinary European guideline for tinnitus: diagnostics, assessment, and treatment', *HNO*, (February 2018). doi: 10.1007/s00106-019-0633-7.
- Cima, R. F. F., Crombez, G. and Vlaeyen, J. W. S. (2011) 'Catastrophizing and Fear of Tinnitus Predict Quality of Life in Patients With Chronic Tinnitus', *Ear and Hearing*, 32(5), pp. 634–641. doi: 10.1097/AUD.0b013e31821106dd.
- Craske, M. G. *et al.* (2014) 'Maximizing exposure therapy: An inhibitory learning approach', *Behaviour Research and Therapy*, 58(310), pp. 10–23. doi: 10.1016/j.brat.2014.04.006.
- Craske, M. G., Hermans, D. and Vervliet, B. (2018) 'State-of-the-art and future directions for extinction as a translational model for fear and anxiety', *Philosophical Transactions of the Royal Society B: Biological Sciences*, 373(1742), p. 20170025. doi: 10.1098/rstb.2017.0025.
- Davis, A. and Refaie, A. E. (2000) 'Epidemiology of Tinnitus', in Tyler, R. S. (ed.) *Handbook of Tinnitus*. San Diego: Singular Thompson Learning, pp. 1–24.
- Fackrell, K. *et al.* (2014) 'Tools for tinnitus measurement: development and validity of questionnaires to assess handicap and treatment effects', in Signorelli, F. and Turjman, F. (eds) *Tinnitus: Causes, Treatment and Short & Long-term Health Effects*. New York: Nova Science Publishers Inc, pp. 13–60.
- Fuller, T. *et al.* (2020) 'Cognitive behavioural therapy for tinnitus', *Cochrane Database of Systematic Reviews*, (4). doi: 10.1002/14651858.CD012614.pub2.
- Fuller, T. E. *et al.* (2019) 'The Fear of Tinnitus Questionnaire', *Ear and Hearing*, 40(6), pp. 1467–1477. doi: 10.1097/AUD.0000000000000728.
- Genitsaridi, E. *et al.* (2019) 'Standardised profiling for tinnitus research: The European School for Interdisciplinary Tinnitus Research Screening Questionnaire (ESIT-SQ)', *Hearing Research*. Elsevier B.V, (xxxx), pp. 1–7. doi: 10.1016/j.heares.2019.02.017.
- Hall, D. A. *et al.* (2016) 'Systematic review of outcome domains and instruments used in clinical trials of tinnitus treatments in adults', *Trials*. *Trials*, 17(1), p. 270. doi: 10.1186/s13063-016-1399-9.
- Hall, D. A. *et al.* (2018) 'A narrative synthesis of research evidence for tinnitus-related complaints as reported by patients and their significant others', *Health and Quality of Life Outcomes*. *Health and Quality of Life Outcomes*, 16(1), pp. 1–15. doi: 10.1186/s12955-018-0888-9.

- Hallam, R. S., Jakes, S. C. and Hinchcliffe, R. (1988) 'Cognitive variables in tinnitus annoyance.', *The British journal of clinical psychology*, 27 ( Pt 3)(pt 3), pp. 213–22. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/3191301>.
- Hedman-Lagerlöf, M. *et al.* (2018) 'Internet-Delivered Exposure Therapy for Fibromyalgia A Randomized Controlled Trial', *Clinical Journal of Pain*, 34(6), pp. 532–542. doi: 10.1097/AJP.0000000000000566.
- Henry, J. A. *et al.* (2012) 'Pilot Study to Evaluate Ecological Momentary Assessment of Tinnitus (Henry et al, 2012)', pp. 279–290.
- Jacquemin, L. *et al.* (2019) 'Sensitivity to change and convergent validity of the Tinnitus Functional Index (TFI) and the Tinnitus Questionnaire (TQ): Clinical and research perspectives', *Hearing Research*. Elsevier B.V, 382, p. 107796. doi: 10.1016/j.heares.2019.107796.
- Kahneman, D. *et al.* (2004) 'A Survey Method for Characterizing Daily Life Experience: The Day Reconstruction Method', *Science*, 306(5702), pp. 1776–1780. doi: 10.1126/science.1103572.
- Kim, H.-J. *et al.* (2015) 'Analysis of the Prevalence and Associated Risk Factors of Tinnitus in Adults', *Plos One*, 10(5), p. e0127578. doi: 10.1371/journal.pone.0127578.
- Kleinstäuber, M. *et al.* (2013) 'The Role of Fear-Avoidance Cognitions and Behaviors in Patients with Chronic Tinnitus', *Cognitive Behaviour Therapy*, 42(2), pp. 84–99. doi: 10.1080/16506073.2012.717301.
- Koehler, M. J. and Levin, J. R. (1998) 'Regulated randomization: A potentially sharper analytical tool for the multiple-baseline design.', *Psychological Methods*, 3(2), pp. 206–217. doi: 10.1037/1082-989X.3.2.206.
- Kratochwill, T. R. *et al.* (2013) 'Single-Case Intervention Research Design Standards', *Remedial and Special Education*, 34(1), pp. 26–38. doi: 10.1177/0741932512452794.
- Lethem, J. *et al.* (1983) 'Outline of a Fear-Avoidance Model of exaggerated pain perception--I.', *Behaviour research and therapy*, 21(4), pp. 401–8. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/6626110>.
- Lourenco, M., Cima, R. and Vlaeyen, J. (2019) *Tinnotes: An usability study*, OSF. Available at: [osf.io/avhb2](http://osf.io/avhb2).
- McCormack, A. *et al.* (2016) 'A systematic review of the reporting of tinnitus prevalence and severity', *Hearing Research*. Elsevier B.V., 337, pp. 70–79. doi: 10.1016/j.heares.2016.05.009.
- Meeus, O., Blavie, C. and Van de Heyning, P. (2007) 'Validation of the Dutch and the French version of the Tinnitus Questionnaire.', *B-ENT*, 3 Suppl 7, pp. 11–7. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/18228680>.
- Meikle, M. B. *et al.* (2012) 'The Tinnitus Functional Index', *Ear and Hearing*, 33(2), pp. 153–176. doi: 10.1097/AUD.0b013e31822f67c0.
- Meuret, A. E. *et al.* (2012) 'Coping Skills and Exposure Therapy in Panic Disorder and Agoraphobia: Latest Advances and Future Directions', *Behavior Therapy*. Elsevier B.V., 43(2), pp. 271–284. doi: 10.1016/j.beth.2011.08.002.
- Morley, S. (2018) *Single-Case Methods in Clinical Psychology: A Practical Guide*. Kindle Edi. Edited by M. Ciara and C. J. Main. Taylor & Francis.
- Newman, C. W., Jacobson, G. P. and Spitzer, J. B. (1996) 'Development of the Tinnitus Handicap Inventory', *Archives of Otolaryngology - Head and Neck Surgery*, 122(2), pp. 143–148. doi: 10.1001/archotol.1996.01890140029007.

- Newman, C. W., Sandridge, S. A. and Jacobson, G. P. (1998) 'Psychometric adequacy of the Tinnitus Handicap Inventory (THI) for evaluating treatment outcome.', *Journal of the American Academy of Audiology*, 9(2), pp. 153–60. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/9564679>.
- Nouwen, A. *et al.* (2006) 'Effects of Focusing and Distraction on Cold Pressor-Induced Pain in Chronic Back Pain Patients and Control Subjects', *The Journal of Pain*, 7(1), pp. 62–71. doi: 10.1016/j.jpain.2005.08.004.
- Ongghena, P. and Edgington, E. S. (2005) 'Customization of Pain Treatments', *The Clinical Journal of Pain*, 21(1), pp. 56–68. doi: 10.1097/00002508-200501000-00007.
- Parker, R. I. *et al.* (2011) 'Combining Nonoverlap and Trend for Single-Case Research: Tau-U', *Behavior Therapy*. Elsevier B.V., 42(2), pp. 284–299. doi: 10.1016/j.beth.2010.08.006.
- Pryss, R. *et al.* (2018) 'Differences between Android and iOS Users of the TrackYourTinnitus Mobile Crowdsensing mHealth Platform', in *2018 IEEE 31st International Symposium on Computer-Based Medical Systems (CBMS)*. IEEE, pp. 411–416. doi: 10.1109/CBMS.2018.00078.
- Rabau, S., Wouters, K. and Van de Heyning, P. (2014) 'Validation and translation of the Dutch tinnitus functional index.', *B-ENT*, 10(4), pp. 251–8. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/25654947>.
- Roelofs, J. *et al.* (2004) 'Does fear of pain moderate the effects of sensory focusing and distraction on cold pressor pain in pain-free individuals?', *Journal of Pain*, 5(5), pp. 250–256. doi: 10.1016/j.jpain.2004.04.001.
- Rosenberger, J. L. and Gasko, M. (1983) 'Comparing location estimators: Trimmed means, medians and trimean', in Hoaglin, D., Mosteller, F., and Tukey, J. (eds) *Understanding robust and exploratory data analysis*, pp. 297–336.
- Schlee, W. *et al.* (2016) 'Measuring the moment-to-moment variability of Tinnitus: The TrackYourTinnitus smart phone app', *Frontiers in Aging Neuroscience*, 8(DEC), pp. 1–8. doi: 10.3389/fnagi.2016.00294.
- Schork, N. J. (2015) 'Personalized medicine: Time for one-person trials', *Nature*, 520(7549), pp. 609–611. doi: 10.1038/520609a.
- Stone, A. A. and Shiffman, S. (1994) 'Ecological Momentary Assessment (Ema) in Behavioral Medicine', *Annals of Behavioral Medicine*, 16(3), pp. 199–202. doi: 10.1093/abm/16.3.199.
- Sullivan, M. J. L., Bishop, S. R. and Pivik, J. (1995) 'The Pain Catastrophizing Scale: Development and validation.', *Psychological Assessment*, 7(4), pp. 524–532. doi: 10.1037/1040-3590.7.4.524.
- Tate, R. L. *et al.* (2016) 'The Single-Case Reporting Guideline In BEhavioural Interventions (SCRIBE) 2016 Statement', *Archives of Scientific Psychology*, 4(1), pp. 10–31. doi: 10.1037/arc0000027.
- Vachon, H. *et al.* (2016) 'Repeated self-evaluations may involve familiarization: An exploratory study related to Ecological Momentary Assessment designs in patients with major depressive disorder', *Psychiatry Research*. Elsevier, 245, pp. 99–104. doi: 10.1016/j.psychres.2016.08.034.
- Vannest, K. J. *et al.* (2016) 'Single Case Research: web based calculators for SCR analysis'. College Station, TX: Texas A&M University. Available at: <http://www.singlecaseresearch.org/calculators/tau-u>.

- Vlaeyen, J. W. S. *et al.* (2001) 'Graded exposure in vivo in the treatment of pain-related fear: a replicated single-case experimental design in four patients with chronic low back pain', *Behaviour Research and Therapy*. IEEE Comput. Soc, 39(2), pp. 151–166. doi: 10.1016/S0005-7967(99)00174-6.
- Vlaeyen, J. W. S. and Linton, S. J. (2000) 'Fear-avoidance and its consequences in chronic musculoskeletal pain: a state of the art.', *Pain*, 85(3), pp. 317–32. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/10781906>.
- Vlaeyen, J. W. S. and Linton, S. J. (2012) 'Fear-avoidance model of chronic musculoskeletal pain: 12 years on', *Pain*, 153(6), pp. 1144–1147. doi: 10.1016/j.pain.2011.12.009.
- Woods, M. P. and Asmundson, G. J. G. (2008) 'Evaluating the efficacy of graded in vivo exposure for the treatment of fear in patients with chronic back pain: A randomized controlled clinical trial', *Pain*, 136(3), pp. 271–280. doi: 10.1016/j.pain.2007.06.037.
- Zigmond, A. S. and Snaith, R. P. (1983) 'The Hospital Anxiety and Depression Scale', *Acta Psychiatrica Scandinavica*, 67(6), pp. 361–370. doi: 10.1111/j.1600-0447.1983.tb09716.x.







## CHAPTER 4

---

# The daily experience of subjective tinnitus: Ecological Momentary Assessment (EMA) vs End-of-Day Diary (EDD)

Published as:

*Lourenco, Matheus P. C. G.; Simoes, Jorge; Vlaeyen, Johan W. S.; & Cima, Rilana F. F.1,2,4 (2021)*  
*The Daily Experience of Subjective Tinnitus, Ear and Hearing:*  
*doi: 10.1097/AUD.0000000000001054*

**Abstract**

**Objective** Traditional methods of self-report assessments are susceptible to bias (i.e. memory, recall, recency). Ecological Momentary Assessment (EMA) may curb these biases by repeated momentary assessment of the participant throughout the day. High costs and participant burden may however impede the use of EMA. End-of-Day Diary (EDD) provides an attractive alternative to EMA, though no direct comparison has been carried-out in the tinnitus field. **Design** 4,732 data entries were collected from nine participants undergoing Cognitive Behavioural Treatment (CBT) for tinnitus. Eleven equivalent EMA and EDD items were collected for approximately 3-months. Tinnitus experience (i.e. anger, annoyance, avoidance, distraction, fear, invasiveness, pleasantness and sadness) and wellbeing (i.e. anxiety, happiness and stress) were correlated and means compared (t-tests). **Results** All variables presented adequate correlation ( $r > .68$ ) between the EMA and EDD counterparts. Small ( $< 3.9\%$ ) significant daily mean differences between EMA and EDD were found for six variables (tinnitus anger, invasiveness, pleasantness, sadness, as well as anxiety and stress) with worse results reported in EDD. **Conclusion** The small significant effects found may be attributed to the large number of data points. When EMA is not possible or recommended, EDD provides a viable alternative to assess tinnitus experience daily. Further research on the underlying mechanisms of tinnitus experience and recollection is warranted.

## 1. Introduction

The use of Ecological Momentary Assessment (EMA) has risen with the development of technology and growing availability of smartphones. The increased use of EMA has been reflected in a variety of research fields, including suicidal ideation (e.g. Kleiman et al., 2017), substance use (e.g. Jones et al., 2019) and chronic pain (e.g. May et al., 2018), to name but a few (for a comprehensive review on EMA we guide the interested reader to Shiffman et al., 2008). EMA aims at capturing experiences during real-life activities and situations by assessing individuals at several random times during the day. The advantages of these in-the-moment assessments are threefold: (1) reduced recall bias, (2) increased ecological validity, and (3) the exploration of symptom fluctuation (Schneider & Stone, 2016). Recall bias refers to any unwanted bias associated with the cognitive processes of memory reconstruction (e.g. mood, setting, recency) and summation (i.e. average) of these experiences (Shields et al., 2016; Stone & Shiffman, 1994). Reducing the time between events and assessment reduces recall bias and focuses on reaching information that can be accessed in working memory. Increased ecological validity is achieved by assessing the participant during real daily-life situations, and thus outside a setting that may unduly influence responses (e.g. hospital, clinic, lab). Fluctuation patterns of experiences (e.g. stress, tinnitus annoyance) during the individual's daily life may provide insights in the relationship of those variables with specific (e.g. social) or cyclical (e.g. sleep/awake) patterns. In order to capture such data, EMA is deployed several times during each day, requiring participants to remain in possession of their smartphone at all times, and allow interruption of activities in order to respond to the assessments. Such intrusiveness has been suggested to potentially produce negative outcomes in participants. Smartphone use has been associated with increased stress, anxiety, depression (Elhai et al., 2017; Vahedi & Saiphoo, 2018) and lower wellbeing (Horwood & Anglim, 2019). While symptom fluctuation during the day may be of importance to researchers and clinicians, daily average EMA scores can provide a broader daily picture.

Similarly, to EMA, End-of-Day Diary (EDD) minimizes the effects of recall bias by being deployed once a day. An established methodology for decades (e.g. Verbrugge, 1980), EDD has been used in a variety of fields, including chronic pain (e.g. Rost et al., 2016), eating behaviour (e.g. Debeuf et al., 2018), and emotionality during the COVID-19 pandemic (Morón & Biolik-Morón, 2021). EDD's benefit of reduced burden to the participant potentially sacrifices ecological validity when compared to EMA (Schneider & Stone, 2016). In order to make an informed choice between EMA and EDD a direct

comparison is warranted. Broderick et al. (2009) found little differences between both assessment methods after one week, however the findings are limited to the field of chronic pain and fatigue as well as to the period of one week. Moreover, results diverged according to the experience being assessed (i.e. pain, fatigue), specifically when comparing the daily equivalence between EMA and EDD. Research on different experiences (i.e. tinnitus) and over longer time periods that usually comprise existent intervention protocols are warranted.

The assessment of tinnitus, the experience of phantom sounds (e.g. high-pitched tone, chirping), relies on self-report only, and a precise evaluation of the experience is paramount for the development of research and symptom management. EMA use within tinnitus is in its infancy, with studies exploring possibilities and limitations of its use (e.g. Gerull et al., 2019; Lourenco et al., 2019; Pryss et al., 2018; Schlee et al., 2016). However, its superiority to retrospective self-reports has been confirmed (Goldberg et al., 2017). On the other hand, use of EDD, while common for decades, it is rarely utilized as an outcome measure within the tinnitus field. The current study aims at comparing results from EMA and EDD assessments in tinnitus patients undergoing treatment, in order to provide recommendations for future research. More specifically, EDD mean values are compared to EMA means. Moreover, EMA gathered close (late in the day) to the EDD completion are compared with earlier-in-the-day EMA and EDD. These analyses elucidate if EDD accurately reflects the overall daily picture, as illustrated by EMA.

## **2. Methodology**

### ***2.1. Participants***

As part of a larger project on the effects of Cognitive Behavioural Therapy (CBT) on chronic tinnitus, we collected data from two subsequent clinical studies (duration of 3-months each) in which both assessment methods were used: EMA and EDD. Studies within the project applied a Single-Case Experimental Design (SCED) approach. In such a design, a small number of participants are repeatedly and consistently assessed to establish an individual and unique control condition (baseline phase). Afterwards, each participant undergoes a manipulation phase (e.g. treatment onset), while maintaining the continuous assessment (for an in depth review of SCED we guide the interested reader to Kazdin, 2018; Morley, 2018). As such, these powerful designs rely on large amount of data from a small number of participants. Each study included six tinnitus patients undergoing specialised CBT for tinnitus which contained a variety of treatment

components (e.g. exposure, relaxation, psychoeducation) delivered twice a week in 2-hour treatment sessions for a total of 20 sessions (for detailed treatment protocol see Cima et al., 2012). Patients on the waiting list for CBT treatment from the Adelante Department of Audiology and Communication (Hoensbroek, The Netherlands) were sequentially invited to participate in the project. Exclusion criteria comprised: (1) undergoing other tinnitus-related or psychological treatment during the time of the study; (2) commenced the use of hearing aid within three months of the start of treatment; (3) commenced or ceased the use of antidepressants, antipsychotics, anxiolytics, Ritalin, hormone replacement therapy, or medication to lower high blood pressure within three months of treatment; (4) unable to read and write in Dutch; (5) disclosed current suicidal intent or (6) had more than 40dB of uncorrected hearing loss in one or both ears as measured by calculating a Pure Tone Average (on the frequencies of 500, 1000 and 2000 Hz).

Patients' tinnitus severity was measured at baseline by the validated Dutch version (Meeus et al., 2007) of the Tinnitus Questionnaire (TQ; Hallam et al., 1988), which utilizes 52 items on a three-point scale for a total score ranging from 0 (low severity) to 104 (high severity). Further characterization of the sample is provided through the Dutch (de Beurs et al., 2001) version of the Depression Anxiety and Stress Scale (DASS-21; Lovibond & Lovibond, 1995). Consistent of 21-items on a 4-point Likert scale, each sub-scale indicates levels of depression, anxiety and stress on a score from 0 (low) to 21 (high).

Each of the two studies included were conducted consecutively starting in May 2019 and registered at the Netherlands Trial Register (trial numbers NL7826 and NL8056). Ethical approval was obtained from the Medical Ethical Committee at Maxima Medical Centre, Veldhoven, The Netherlands (METC; NL63262.016.18).

## *2.2. Ecological Momentary Assessments (EMA) and End-of-Day Diary (EDD)*

EMA and EDD data were collected continuously throughout the duration of treatment (3 months). EMA and EDD were collected through purpose-built apps installed on participants' smartphones. One study utilized an in-house developed app (TinNotes) by Maastricht University's Instrumentation Engineering department, while the subsequent study utilized an equivalent third-party app (mEMA; ilumivu, Inc., Cambridge, MA, USA; www.ilumivu.com). EDD assessments were delivered at 8-pm with a 4-hour time limit for completion. EMA questions were prompted seven times during the day, at random points with at least 2-hours in between prompts. Participants had the option

to snooze the prompt twice for 5-mins each time, after which the questionnaire would not be available any longer and result in a missing EMA measure for that time-point. Individualized sleeping hours were set so that prompts would not be delivered during those hours. Participants had to complete at least 50% of EDD assessments to be included for analysis. Assessments comprised of 16 (EDD) and 17 (EMA) items, presented in random order, of which 12 had content-equivalence. Eleven of the equivalent items (Table 1) were rated on a 7-point Likert scale (0-6) and related to either tinnitus experience (i.e. anger, annoyance, avoidance, distraction, fear, invasiveness, pleasantness and sadness) or overall wellbeing (i.e. anxiety, happiness and stress). One item (Social Interaction; EMA - “*Who are you with?*”; Diary - “*Who did you spend time with today?*”) was descriptive and not included for analysis.

**Table 1:** Equivalent items of both assessment types: End-of-Day Diary (EDD) and Ecological Momentary Assessment (EMA).

EDD	EMA
How angry did your tinnitus make you today?	My tinnitus makes me angry
How annoying was your tinnitus today?	My tinnitus is annoying
How anxious were you today?	I feel anxious
How hard did you try to avoid your tinnitus today?	I try to avoid the tinnitus
How distracting was your tinnitus today?	My tinnitus is distracting
How afraid of hearing your tinnitus were you today?	I am afraid of hearing my tinnitus
How happy were you today?	I feel happy
How invasive was your tinnitus today?	My tinnitus is invasive
How pleasant was your tinnitus today?	How pleasant is your tinnitus?
How sad did your tinnitus make you today?	My tinnitus makes me sad
How stressful has your day been?	I feel stressed

### 2.3. Analysis

Pairwise comparisons using Spearman Rank Correlation between EMA and EDD were carried out between all equivalent items. EMA data of each item was plotted through time and a daily mean calculated. In order to compare EMA gathered proximally to EDD (delivered at 8-pm) and given the minimum 2-hour gap between EMA prompts, EMA delivered from 6-pm (2-hours before EDD delivery) was separated. Two new EMA means were calculated: (1) early EMA (before 6-pm), and (2) late EMA (after 6-pm). Paired t-tests between EMA means and EDD were conducted and corrected for multiple comparison (Holm, 1979). The Holm method controls for family-wise Type I error, with corrections decreasing the threshold of significance for each hypothesis tested. Following



convention, we considered  $p$ -values below 0.05 “statistically significant”. Pairwise deletions were used to account for missing values. Effect sizes were calculated through Cohen’s  $D$  (Cohen, 1988).


Statistical analyses were performed with R version 4.0.1 (R Core Team, 2020) with supporting packages (Grolemund & Wickham, 2011; Tiedemann, 2020; Tierney, 2017; Wei & Simko, 2017; Wickham, 2019; Wickham et al., 2018, 2019).

### 3. Results


Nine participants (88.9% Men; Mean age = 58.11, SD = 9.98) were included for analysis for a total of 4,732 data entries (Table 2). From the original pool of 12 participants, 1 participant dropped out due to personal reasons unrelated to the treatment. An unknown error with the TinNotes app deemed data for two other participants to be unreliable. Data for one participant, who had recently commenced the use hearing aid, was included for analysis as the use of the hearing aid was not continued during treatment.

EMA fluctuations (Figure 1) show the difference between experience variability according to the time of day (e.g. decrease of tinnitus fear after 7-pm). Strong correlations ( $r > .70$ ) were found for all but one (i.e. stress) EMA and EDD items (Figure 2). Paired  $t$ -tests (Table 3) indicated significant differences between EMA and EDD daily means on six variables (i.e. tinnitus anger, anxiety, tinnitus invasiveness, tinnitus pleasantness, tinnitus sadness, stress). EDD reports for these variables were significantly worse with the exception of tinnitus avoidance, which indicated no differences (Figure 3). Comparisons between EDD and early EMA (before 6-pm) indicate similar results of worse EDD scores for five variables (i.e. tinnitus anger, anxiety, tinnitus invasiveness, tinnitus sadness, stress). Moreover, EDD comparisons with late EMA (after 6-pm) indicated worse EDD scores for four variables (i.e. tinnitus anger, anxiety, tinnitus pleasantness, tinnitus sadness, stress) and improved scores for tinnitus avoidance.

**Figure 1:** Mean Ecological Momentary Assessment (EMA) recording per hour.



**Figure 2:** Correlation coefficient strengths.



**Table 2:** Demographic characteristics

P	Age	Gender	Hearing aid (time)	DASS-21			Compliance (%)			Tinnitus		
				D	A	S	EMA	EDD	Duration	Location	Type	Severity
1	67	Men	10 years	4	0	2	15.8	77.5	2 years	Both ears	High pitch	81
2	62	Men	> 30 years	8	4	6	51.3	73.6	12 years	Both sides of head	High pitch	48
3	59	Men	No hearing aid	2	0	6	70.6	98.8	9 months	Both ears	High pitch	61
4	65	Men	No hearing aid	0	0	0	89.8	83.8	15 years	In the head	Middle pitch	68
5	66	Men	1 week	12	6	16	64.8	87.5	12 years	Right ear	High pitch	57
6	57	Women	No hearing aid	6	10	11	72.9	90.9	35 years	Both ears	Buzzing	54
7	64	Men	No hearing aid	1	1	3	90.5	98.5	2 years	Right side on the back of the head	High pitch	38
8	43	Men	No hearing aid	13	9	11	76.8	100	11 months	Whole head	High pitch	80
9	40	Men	No hearing aid	2	0	6	70.6	89.4	6 years	Both ears	High and middle pitch	77

Note: Tinnitus severity measured by Tinnitus Questionnaire (TQ); Depression (D), Anxiety (A) and Stress (S) measured by the Depression Anxiety and Stress Scale (DASS-21).

**Table 3:** Paired sample t-tests with adjusted p-values for equivalent variables of Ecological Momentary Assessment (EMA) and End-of-Day Diary (EDD) on a Likert scale (0-6).


	<b>Mean EMA (SD)</b>	<b>Mean EDD (SD)</b>	<b>p</b>	<b>Cohen's D</b>
<b>Anger</b>	1.756 (1.723)	1.870 (1.748)	< .001**	.065
<b>Annoyance</b>	3.382 (1.931)	3.414 (1.932)	.156	.017
<b>Anxiety</b>	1.575 (1.42)	1.648 (1.44)	< .001**	.050
<b>Avoidance</b>	1.454 (1.477)	1.422 (1.451)	.156	-.022
<b>Distracting</b>	3.242 (1.891)	3.202 (1.855)	.110	.065
<b>Fear</b>	1.403 (1.399)	1.425 (1.431)	.170	.016
<b>Happy†</b>	2.503 (1.308)	2.507 (1.247)	.815	.003
<b>Invasiveness</b>	3.461 (1.895)	3.522 (1.805)	< .001**	.033
<b>Pleasant†</b>	3.649 (2.271)	3.587 (2.28)	.030*	-.027
<b>Sadness</b>	1.869 (1.679)	1.956 (1.697)	< .001**	.051
<b>Stress</b>	1.831 (1.576)	2.102 (1.587)	< .001**	10.172
	<b>Mean early EMA (SD)</b>	<b>Mean EDD (SD)</b>	<b>p</b>	<b>Cohen's D</b>
<b>Anger</b>	1.885 (1.777)	1.997 (1.791)	< .001**	.063
<b>Annoyance</b>	3.337 (1.954)	3.372 (1.939)	.348	.018
<b>Anxiety</b>	1.690 (1.447)	1.750 (1.452)	< .001**	.042
<b>Avoidance</b>	1.563 (1.514)	1.550 (1.485)	.974	-.009
<b>Distracting</b>	3.199 (1.912)	3.175 (1.858)	.920	.063
<b>Fear</b>	1.507 (1.43)	1.532 (1.463)	.580	.017
<b>Happy†</b>	2.52 (1.354)	2.527 (1.283)	.974	.006
<b>Invasiveness</b>	3.374 (1.893)	3.438 (1.792)	.007*	.034
<b>Pleasant†</b>	3.408 (2.294)	3.377 (2.3)	.920	-.013
<b>Sadness</b>	2.007 (1.711)	2.066 (1.726)	< .001**	.035
<b>Stress</b>	1.988 (1.605)	2.211 (1.594)	< .001**	10.140
	<b>Mean late EMA (SD)</b>	<b>Mean EDD (SD)</b>	<b>p</b>	<b>Cohen's D</b>
<b>Anger</b>	1.386 (1.5)	1.503 (1.564)	< .001**	.076
<b>Annoyance</b>	3.511 (1.857)	3.534 (1.906)	1	.013
<b>Anxiety</b>	1.248 (1.285)	1.353 (1.361)	< .001**	.079
<b>Avoidance</b>	1.142 (1.317)	1.053 (1.28)	.012*	-.068
<b>Distracting</b>	3.365 (1.826)	3.276 (1.847)	.080	.076
<b>Fear</b>	1.101 (1.259)	1.119 (1.287)	1	.014
<b>Happy†</b>	2.456 (1.164)	2.448 (1.134)	1	-.006
<b>Invasiveness</b>	3.709 (1.881)	3.762 (1.82)	.296	.029
<b>Pleasant†</b>	4.339 (2.054)	4.19 (2.11)	< .001**	-.072
<b>Sadness</b>	1.475 (1.517)	1.639 (1.572)	< .001**	.106
<b>Stress</b>	1.379 (1.397)	1.789 (1.523)	< .001**	10.279

	Mean early EMA (SD)	Mean late EMA (SD)	p	Cohen's D
<b>Anger</b>	1.87 (1.748)	1.756 (1.723)	< .001**	Paired Cohen's D could not be computed due to unequal number of measures
<b>Annoyance</b>	3.414 (1.932)	3.382 (1.931)	.003*	
<b>Anxiety</b>	1.648 (1.44)	1.575 (1.42)	< .001**	
<b>Avoidance</b>	1.422 (1.451)	1.454 (1.477)	< .001**	
<b>Distracting</b>	3.202 (1.855)	3.242 (1.891)	.004*	
<b>Fear</b>	1.425 (1.431)	1.403 (1.399)	< .001**	
<b>Happy†</b>	2.507 (1.247)	2.503 (1.308)	.163	
<b>Invasiveness</b>	3.522 (1.805)	3.461 (1.895)	< .001**	
<b>Pleasant†</b>	3.587 (2.28)	3.649 (2.271)	< .001**	
<b>Sadness</b>	1.956 (1.697)	1.869 (1.679)	< .001**	
<b>Stress</b>	2.102 (1.587)	1.831 (1.576)	< .001**	

\*p < .05; \*\*p < .001; †inverted item

Comparison of early and late EMA indicated significant differences in all but one (i.e. happiness) variables. Items on tinnitus anger, annoyance, fear, invasiveness, pleasantness, and sadness as well as levels of anxiety and stress improved after 6-pm, while tinnitus avoidance and distraction worsened.

**Figure 3:** Ecological Momentary Assessment (EMA) and End-of-Day Diary (EDD) distribution and box plot per variable.



## 4. Discussion

This study sets out to compare two similar but different daily measurement methods, namely EMA and EDD in chronic tinnitus patients during a 12-week treatment. Generally, both methods provide quite similar results. All but one item (stress,  $r = .69$ ) showed strong correlations between EMA and its EDD counterparts ( $r > .77$ ). Nevertheless, EDD stress reports are significantly higher than early-in-the-day EMA measures (where mean stress levels were at their highest). EDD painted a worse picture for another five variables when compared to EMA (i.e. tinnitus anger, anxiety, tinnitus invasiveness, tinnitus pleasantness, tinnitus sadness). EDD reports favoured negative experiences rather than recent experiences (i.e. EMA after 6-pm). Broderick et al. (2009) found similar results when comparing EMA and EDD of pain and fatigue experiences. Such occurrence is akin to the “experience memory gap” (Miron-Shatz et al., 2009), where recalled symptoms are reported as worse when compared to real-time in the moment assessments (i.e. EMA). Such memory biases were studied in a recent review (Van Den Bergh & Walentynowicz, 2016), indicating that pain and fatigue experiences are overreported when assessments rely on longer recall periods. While these findings are significant in the field of self-report assessments, no study with tinnitus complaints was included in the review and parallels must be drawn with caution.

The large sample of data provided by novel methodological approaches (e.g. EMA, EDD, SCED) present both statistical opportunities and issues not commonly encountered. Despite correcting for multiple comparisons (i.e. Holm, 1979), which decreased the threshold for significant results, the findings are still affected by the large number of data and traditional p-value selected (i.e. 0.05). As such, more conservative approaches that are beyond multiple comparison corrections may provide a more accurate picture of the results (i.e. lower p-value thresholds). In the current study, the largest significant mean difference found in tinnitus related variables was tinnitus anger (1.63%), with stress levels (3.87%) holding the largest, although small, difference in wellbeing variables. Whether these statistical differences are clinically relevant are therefore questionable. Furthermore, while EDD results may have differed from early or late EMA, the daily EMA mean accurately reflected the remaining variables (i.e. tinnitus annoyance, tinnitus avoidance, tinnitus distraction, tinnitus fear, and happiness).

An exception was found in happiness levels, which did not significantly differ between EMA and EDD measures at any point. EMA and EDD measures of happiness strongly correlated ( $r = .80$ ) even though both measures correlated weakly ( $r < .24$ ) with other

variables. Despite this seemingly independent level of happiness from other experiences (e.g. tinnitus anger, tinnitus annoyance, anxiety, stress), accurate assessment of happiness remains a challenge, with the very definition of happiness still debated (Ludwigs et al., 2019). As such, interpretations of happiness stability and independence are limited.

The choice of variables to be measured, while theoretically driven and based on specialist consensus, lacked the insight from other key stakeholders and may further benefit from initiatives acknowledging patient preferred outcomes (i.e. Hall et al., 2018). An added benefit of EMA and EDD measures is that it may conform with the push for individualized medicine (Schork, 2015; Senn, 2018) due to its flexibility in incorporating different items. Therefore, while the outcomes used in the current research are relevant within its theoretical framework, they are limited by the pool of specialist used to create the items. Further research utilizing a broader consensus of outcome variables, as suggested by Hall et al. (2018) may increase the relevance and use of EMA and EDD. Moreover, the choice of a 7-point Likert scale, while not directly inspired by standardized tinnitus self-report assessments, was made due to technical limitations of the TinNotes app. Further research incorporating other scales, specifically Visual Analog Scales (VAS), are recommended.

Additional limitations include the high proportion of men 40 years or older (88.89%) in the sample, limiting the generalizability of the findings. The homogeneous sample follows epidemiological trends in tinnitus, with 80% of tinnitus diagnosed after the age of 40 (Stohler et al., 2019) and higher incidence detected in men (Fujii et al., 2011; McCormack et al., 2014, 2016). Despite the limitations, the current results add important knowledge on long-term EMA versus EDD comparisons and provide insights into using these methods in tinnitus patients (in addition to chronic pain and fatigue)

## **5. Conclusion**

Generally, EDD and EMA provide similar data. EDD measures significantly differed from EMA daily averages for six out of eleven variables: tinnitus anger, anxiety, tinnitus invasiveness, tinnitus pleasantness, tinnitus sadness, and stress. The differences support previous literature which found that longer recall periods associate with worse symptom/experience recollection. Despite their statistical significance, the effects were small and may be attributed to the large number of data entries. As such, the minor differences may not justify EMA as the measurement of choice as the added burden to participants may be of ethical or theoretical concern. When these arise, EDD provides a viable alternative



since it accurately and closely reflects daily life experiences as measured by EMA daily mean. Nonetheless, when the use of EMA is necessary the minor differences found in the current study do not justify a correction of the data collected. EMA may better suit the need of closely investigating cyclical tinnitus patterns (e.g. sleep/awake) or possible daily correlates (e.g. work environment, presence of triggers). The knowledge of specific correlates allows for the recognition of maladaptive patterns and emotional reactions which may be addressed during treatment. Moreover, the use of repeated assessments (i.e. EMA and EDD) is vital in the application of SCEDs which are tailored to the push for individualized research and treatment (Schork, 2015).

The continuous development and understanding of tinnitus assessment must be prioritized as the lack of an objective measure of tinnitus entails an over-reliance on patient self-reports for research and treatment. Future research on accurate measurements of the underlying mechanisms of the tinnitus experience may pave the way for a broader understanding about the onset, maintenance and recovery of tinnitus disability.

## References

- Broderick, J. E., Schwartz, J. E., Schneider, S., & Stone, A. A. (2009). Can End-of-Day Reports Replace Momentary Assessment of Pain and Fatigue? *Journal of Pain*, *10*(3), 274–281. <https://doi.org/10.1016/j.jpain.2008.09.003>
- Cima, R. F. F., Maes, I. H., Joore, M. A., Scheyen, D. J. W. M., El Refaie, A., Baguley, D. M., Anteunis, L. J. C., van Breukelen, G. J. P., & Vlaeyen, J. W. S. (2012). Specialised treatment based on cognitive behaviour therapy versus usual care for tinnitus: a randomised controlled trial. *The Lancet*, *379*(9830), 1951–1959. [https://doi.org/10.1016/S0140-6736\(12\)60469-3](https://doi.org/10.1016/S0140-6736(12)60469-3)
- Cohen, J. (1988). *Statistical Power Analysis for the Behavioral Sciences*. Routledge Academic.
- de Beurs, E., Van Dyck, R., Marquenie, L. A., Lange, A., & Blonk, R. W. B. (2001). De DASS: Een vragenlijst voor het meten van depressie, angst en stress [The DASS: A questionnaire for the measurement of depression, anxiety, and stress]. *Gedragstherapie*, *34*(1), 35–53.
- Debeuf, T., Verbeken, S., Van Beveren, M.-L., Michels, N., & Braet, C. (2018). Stress and Eating Behavior: A Daily Diary Study in Youngsters. *Frontiers in Psychology*, *9*. <https://doi.org/10.3389/fpsyg.2018.02657>
- Elhai, J. D., Dvorak, R. D., Levine, J. C., & Hall, B. J. (2017). Problematic smartphone use: A conceptual overview and systematic review of relations with anxiety and depression psychopathology. *Journal of Affective Disorders*, *207*, 251–259. <https://doi.org/10.1016/j.jad.2016.08.030>
- Fujii, K., Nagata, C., Nakamura, K., Kawachi, T., Takatsuka, N., Oba, S., & Shimizu, H. (2011). Prevalence of Tinnitus in Community-Dwelling Japanese Adults. *Journal of Epidemiology*, *21*(4), 299–304. <https://doi.org/10.2188/jea.JE20100124>
- Gerull, K. M., Kallogjeri, D., Piccirillo, M. L., Rodebaugh, T. L., Lenze, E. J., & Piccirillo, J. F. (2019). Feasibility of Intensive Ecological Sampling of Tinnitus in Intervention Research. *Otolaryngology–Head and Neck Surgery*, 019459981984496. <https://doi.org/10.1177/0194599819844968>
- Goldberg, R. L., Piccirillo, M. L., Nicklaus, J., Skillington, A., Lenze, E., Rodebaugh, T. L., Kallogjeri, D., & Piccirillo, J. F. (2017). Evaluation of ecological momentary assessment for tinnitus severity. *JAMA Otolaryngology - Head and Neck Surgery*, *143*(7), 700–706. <https://doi.org/10.1001/jamaoto.2017.0020>
- Grolemund, G., & Wickham, H. (2011). Dates and Times Made Easy with lubridate. *Journal of Statistical Software*, *40*(3), 1–25. <http://www.jstatsoft.org/v40/i03/>
- Hall, D. A., Smith, H., Hibbert, A., Colley, V., Haider, H. F., Horobin, A., Londero, A., Mazurek, B., Thacker, B., & Fackrell, K. (2018). The COMiT-ID Study: Developing Core Outcome Domains Sets for Clinical Trials of Sound-, Psychology-, and Pharmacology-Based Interventions for Chronic Subjective Tinnitus in Adults. *Trends in Hearing*, *22*, 233121651881438. <https://doi.org/10.1177/2331216518814384>
- Hallam, R. S., Jakes, S. C., & Hinchcliffe, R. (1988). Cognitive variables in tinnitus annoyance. *The British Journal of Clinical Psychology*, *27* ( Pt 3)(pt 3), 213–222. <http://www.ncbi.nlm.nih.gov/pubmed/3191301>
- Holm, S. (1979). Board of the Foundation of the Scandinavian Journal of Statistics A Simple Sequentially Rejective Multiple Test Procedure A Simple Sequentially Rejective Multiple Test Procedure. *Source: Scandinavian Journal of Statistics Scand J Statist*, *6*(6), 65–70. <http://www.jstor.org/stable/4615733><http://www.jstor.org/page/info/about/policies/terms.jsp><http://www.jstor.org>

- Horwood, S., & Anglim, J. (2019). Problematic smartphone usage and subjective and psychological well-being. *Computers in Human Behavior, 97*, 44–50. <https://doi.org/10.1016/j.chb.2019.02.028>
- Jones, A., Remmerswaal, D., Verveer, I., Robinson, E., Franken, I. H. A., Wen, C. K. F., & Field, M. (2019). Compliance with ecological momentary assessment protocols in substance users: a meta-analysis. *Addiction, 114*(4), 609–619. <https://doi.org/10.1111/add.14503>
- Kazdin, A. E. (2018). Single-case experimental designs. Evaluating interventions in research and clinical practice. *Behaviour Research and Therapy, November*, 0–1. <https://doi.org/10.1016/j.brat.2018.11.015>
- Kleiman, E. M., Turner, B. J., Fedor, S., Beale, E. E., Huffman, J. C., & Nock, M. K. (2017). Examination of real-time fluctuations in suicidal ideation and its risk factors: Results from two ecological momentary assessment studies. *Journal of Abnormal Psychology, 126*(6), 726–738. <https://doi.org/10.1037/abn0000273>
- Lourenco, M., Cima, R., & Vlaeyen, J. (2019). *Tinnotes: An usability study*. OSF. [osf.io/avhb2](https://osf.io/avhb2)
- Lovibond, S. H., & Lovibond, P. F. (1995). *Manual for the Depression Anxiety & Stress Scales* (2nd Ed.). Psychology Foundation.
- Ludwigs, K., Henning, L., & Arends, L. R. (2019). *Measuring Happiness—A Practical Review* (pp. 1–34). [https://doi.org/10.1007/978-3-030-15115-7\\_1](https://doi.org/10.1007/978-3-030-15115-7_1)
- May, M., Junghaenel, D. U., Ono, M., Stone, A. A., & Schneider, S. (2018). Ecological Momentary Assessment Methodology in Chronic Pain Research: A Systematic Review. *The Journal of Pain, 19*(7), 699–716. <https://doi.org/10.1016/j.jpain.2018.01.006>
- McCormack, A., Edmondson-Jones, M., Fortnum, H., Dawes, P., Middleton, H., Munro, K. J., & Moore, D. R. (2014). The prevalence of tinnitus and the relationship with neuroticism in a middle-aged UK population. *Journal of Psychosomatic Research, 76*(1), 56–60. <https://doi.org/10.1016/j.jpsychores.2013.08.018>
- McCormack, A., Edmondson-Jones, M., Somerset, S., & Hall, D. A. (2016). A systematic review of the reporting of tinnitus prevalence and severity. *Hearing Research, 337*, 70–79. <https://doi.org/10.1016/j.heares.2016.05.009>
- Meeus, O., Blavie, C., & Van de Heyning, P. (2007). Validation of the Dutch and the French version of the Tinnitus Questionnaire. *B-ENT, 3 Suppl 7*, 11–17. <http://www.ncbi.nlm.nih.gov/pubmed/18228680>
- Miron-Shatz, T., Stone, A., & Kahneman, D. (2009). Memories of Yesterday's Emotions: Does the Valence of Experience Affect the Memory-Experience Gap? *Emotion, 9*(6), 885–891. <https://doi.org/10.1037/a0017823>
- Morley, S. (2018). *Single-Case Methods in Clinical Psychology: A Practical Guide* (M. Ciara & C. J. Main (eds.); Kindle Ed). Taylor & Francis.
- Moroń, M., & Biolik-Moroń, M. (2021). Trait emotional intelligence and emotional experiences during the COVID-19 pandemic outbreak in Poland: A daily diary study. *Personality and Individual Differences, 168*, 110348. <https://doi.org/10.1016/j.paid.2020.110348>
- Pryss, R., Reichert, M., Schlee, W., Spiliopoulou, M., Langguth, B., & Probst, T. (2018). Differences between Android and iOS Users of the TrackYourTinnitus Mobile Crowdsensing mHealth Platform. *2018 IEEE 31st International Symposium on Computer-Based Medical Systems (CBMS)*, 411–416. <https://doi.org/10.1109/CBMS.2018.00078>

- R Core Team. (2020). *R: A language and environment for statistical computing* (4.0.1). R Foundation for Statistical Computing. <https://www.r-project.org/>
- Rost, S., Van Rycckeghem, D. M. L., Koval, P., Sütterlin, S., Vögele, C., & Crombez, G. (2016). Affective instability in patients with chronic pain. *PAIN*, *157*(8), 1783–1790. <https://doi.org/10.1097/j.pain.0000000000000582>
- Schlee, W., Pryss, R. C., Probst, T., Schobel, J., Bachmeier, A., Reichert, M., & Langguth, B. (2016). Measuring the moment-to-moment variability of Tinnitus: The TrackYourTinnitus smart phone app. *Frontiers in Aging Neuroscience*, *8*(DEC), 1–8. <https://doi.org/10.3389/fnagi.2016.00294>
- Schneider, S., & Stone, A. A. (2016). Ambulatory and diary methods can facilitate the measurement of patient-reported outcomes. *Quality of Life Research*, *25*(3), 497–506. <https://doi.org/10.1007/s11136-015-1054-z>
- Schork, N. J. (2015). Personalized medicine: Time for one-person trials. *Nature*, *520*(7549), 609–611. <https://doi.org/10.1038/520609a>
- Senn, S. (2018). Statistical pitfalls of personalized medicine. *Nature*, *563*(7733), 619–621. <https://doi.org/10.1038/d41586-018-07535-2>
- Shields, A. L., Shiffman, S., & Stone, A. (2016). Recall bias: Understanding and reducing bias in PRO data collection. In *EPro: Electronic solutions for patient-reported data* (pp. 5–21).
- Shiffman, S., Stone, A. A., & Hufford, M. R. (2008). Ecological Momentary Assessment. *Annual Review of Clinical Psychology*, *4*(1), 1–32. <https://doi.org/10.1146/annurev.clinpsy.3.022806.091415>
- Stohler, N. A., Reinau, D., Jick, S. S., Bodmer, D., & Meier, C. R. (2019). A study on the epidemiology of tinnitus in the United Kingdom. *Clinical Epidemiology, Volume 11*, 855–871. <https://doi.org/10.2147/CLEPS213136>
- Stone, A. A., & Shiffman, S. (1994). Ecological Momentary Assessment (Ema) in Behavioral Medicine. *Annals of Behavioral Medicine*, *16*(3), 199–202. <https://doi.org/10.1093/abm/16.3.199>
- Tiedemann, F. (2020). *gghalves: Compose Half-Half Plots Using Your Favourite Geoms* (R package version 0.1.0.). <https://cran.r-project.org/package=gghalves>
- Tierney, N. (2017). visdat: Visualising Whole Data Frames. *The Journal of Open Source Software*, *2*(16), 355. <https://doi.org/10.21105/joss.00355>
- Vahedi, Z., & Saiphoo, A. (2018). The association between smartphone use, stress, and anxiety: A meta-analytic review. *Stress and Health*, *34*(3), 347–358. <https://doi.org/10.1002/smi.2805>
- Van Den Bergh, O., & Walentynowicz, M. (2016). Accuracy and bias in retrospective symptom reporting. *Current Opinion in Psychiatry*, *29*(5), 302–308. <https://doi.org/10.1097/YCO.0000000000000267>
- Verbrugge, L. M. (1980). Health Diaries. *Medical Care*, *18*(1), 73–95. <https://doi.org/10.1097/00005650-198001000-00006>
- Wei, T., & Simko, V. (2017). *R package “corrplot”: Visualization of a Correlation Matrix* (Version 0.84). <https://github.com/taiyun/corrplot>
- Wickham, H. (2019). *stringr: Simple, Consistent Wrappers for Common String Operations* (R package version 1.4.0.).

- Wickham, H., Averick, M., Bryan, J., Chang, W., McGowan, L., François, R., Grolemund, G., Hayes, A., Henry, L., Hester, J., Kuhn, M., Pedersen, T., Miller, E., Bache, S., Müller, K., Ooms, J., Robinson, D., Seidel, D., Spinu, V., ... Yutani, H. (2019). Welcome to the Tidyverse. *Journal of Open Source Software*, 4(43), 1686. <https://doi.org/10.21105/joss.01686>
- Wickham, H., Hester, J., & François, R. (2018). *readr: Read Rectangular Text Data* (R package version 1.3.1.). <https://cran.r-project.org/package=readr>



## CHAPTER 5

---

# Better together. Group vs individual Cognitive Behavioural Therapy (CBT) for tinnitus: A multiple-baseline Single-Case Experimental Design

Under revision:

Matheus Lourenco, Thomas Fuller, Saskia Ranson, Johan Vlaeyen & Rilana Cima.

Better together. Group vs individual Cognitive Behavioural Therapy (CBT) for tinnitus:  
A multiple-baseline Single-Case Experimental Design. *Ear and Hearing*

## **Abstract**


Chronic tinnitus is best treated through Cognitive Behaviour Therapy (CBT). Both group and individual CBT for tinnitus are effective, but no study has directly compared the two. The current study explores group vs individual CBT for tinnitus. A multiple-baseline single-case experimental design was employed in order to observe changes within/between individual and group treatments. Six participants started a 10-week CBT protocol and were equally divided into individual or group treatment. Participants were exchanged between treatments at random time-points. Diary data included 14 variables on tinnitus experience (e.g. annoyance, distraction) and wellbeing (e.g. happiness, stress). Five male participants (59-67 year old) completed treatment. Randomization tests comparing means between individual and group treatments did not reveal significant differences. Analysis of data overlap and trend (Tau-U) revealed minor significant improvements for 7 variables (50%) in group treatment as compared to individual treatment. Diminished happiness and activity levels were observed in participants who went from group to individual treatment. Low effect sizes and homogeneity of sample restrict the generalizability of data. Group CBT indicated potential benefits when compared to individual CBT. Social learning may be an underlying process in group delivery boosting tinnitus recovery. Findings are limited to male patients with chronic disabling tinnitus.



## 1. Introduction

Increasing incidence (e.g. Moore et al., 2019; Stohler et al., 2019) and high economic burden of chronic tinnitus (Maes et al., 2013) demand effective interventions to be applied. Chronic tinnitus, the persistent perception of sound (e.g. ringing, humming or hissing) in the ear(s) or head without corresponding external acoustic source, has no cure. However, Cognitive Behavioural Therapy (CBT) is strongly recommended (Cima et al., 2019) with a recent Cochrane review supporting its positive effect on participant's quality of life (Fuller et al., 2020). Based on the Fear Avoidance model (Figure 1) of chronic pain (FA; Lethem et al., 1983; Vlaeyen & Linton, 2000, 2012) adapted to tinnitus (Cima et al., 2011; Kleinstäuber et al., 2013), CBT aims to alter fear cognitions and responses by targeting different steps of the pathological cycle (e.g. catastrophizing, fear, avoidance) through different techniques (e.g. psycho-education, exposure, relaxation). CBT for tinnitus may take many forms, as different combinations of techniques may be applied.

**Figure 1:** Fear Avoidance model (reproduced from Cima et al., 2018).



Group treatment is an effective form of CBT delivery (e.g. Cima et al., 2012). Despite support for group-based CBT, it is not yet known if the group setting is vital for the effectiveness of CBT for tinnitus. Differences in method of treatment-delivery (i.e. group compared to individual therapy) have previously been explored by Fuller et al. (2020) in the context of a meta-analysis. Their analyses revealed that both individual and group-

based CBT were more effective than wait list control or “active comparison” conditions, but that there was no difference between individual or group-based delivery. In the meta-analysis, results from studies, using either individual or group-based delivery of different CBT protocols, were not compared directly in one single study. Furthermore, the studies reported mean group-level results, which limits insight in changes during treatment at the individual level. Tinnitus sufferers report high intra- and inter-individual differences (Henry et al., 2012; Schlee et al., 2016) which may unduly influence results, and which might not be accurately captured through group-level statistical analysis.

To our knowledge there is no study directly comparing method of delivery of CBT for tinnitus. The current study aims to examine the differences between individual and group treatment. A Single-Case Experimental Design (SCED) is employed to disentangle the treatment delivery methods by exploring the individual path of participants during CBT for tinnitus while alternating between group and individual treatment phases.


## **2.Method**

### ***2.1. Participants***

Patients on the waiting list for CBT treatment from the Adelante Department of Audiology and Communication (Hoensbroek, The Netherlands) were contacted and invited sequentially to participate in the study. First, patients who had previously volunteered for research on CBT for tinnitus, but could not participate due to scheduling conflicts, were contacted. All other patients on the waiting list for CBT for tinnitus were contacted next (i.e. those in the waiting list longest were invited first).

Potential participants were excluded from participation if they reportedly: (1) were currently undergoing other tinnitus or psychological treatment elsewhere; (2) commenced the use of hearing aids within the previous three months; (3) commenced or ceased the use of antidepressants, antipsychotics, anxiolytics, Ritalin, hormone replacement therapy, or medication to lower high blood pressure within the previous three months; (4) were unable to read and write in Dutch; or (5) disclosed current suicidal intent. In addition, they were excluded if (6) they showed uncorrected hearing loss of 40dB or more in one or both ears assessed by calculating the Pure Tone Average (PTA) on 500 Hz, 1000 Hz, and 2000 Hz. A total of six participants were included in the study (Figure 2). Eligible participants received study information by mail and were contacted by phone to evaluate inclusion/exclusion criteria and an appointment was set-up for signing informed consent.

**Figure 2:** Flow of recruitment for current study.



## 2.2. CBT for Tinnitus

The center provides established CBT for tinnitus (for detailed protocol see Cima et al., 2012). Relying on a stepped care approach, patients first undergo audiological assessment, psycho-educational session, and a psychological intake, as a first step in triage.


A multidisciplinary team meeting of the involved professionals determines each patient's treatment path. Patients are allocated to the more intensive second step of treatment with 2-hour sessions for approximately 10-weeks, if diagnostics showed more severe patient need. The current study examined the second step of the stepped care approach with bi-weekly treatment delivery of 20 sessions in total. Therapists experienced with the protocol utilized at the rehabilitation centre followed the pre-established descriptions of each treatment session.

### *2.3. Design*

An adapted multiple-baseline Single-Case Experimental Design (SCED) was utilized in order to examine differences in effects due to method of delivery by comparing changes in individual and group setting. Compared to traditional Randomized Control Trials (RCTs), the SCED is more appropriate for dissecting treatment components (e.g. method of delivery) as all participants undergo the isolated components (i.e. in this case, individual and group setting) during different phases of the treatment (Krasny-Pacini & Evans, 2018). More specifically, after a baseline (phase A), each participant underwent the same treatment protocol in two different sequential orders (BC vs CB): individual (phase B) or group (phase C). The usual multiple-baseline design relies on a bi-phasic A-B design and randomizes the baseline length (A) to increase power for comparisons with a subsequent treatment phase (B). The design utilized in this study randomizes the length of each treatment phase instead of baseline. The adaptation takes into consideration the research question of comparing treatment settings to each other rather than comparing treatment effectiveness to a no-treatment baseline. Three random paired schedules with a minimum of 14 observations per treatment phase (Figure 3) were randomly drawn using a dedicated Single Case Data Analysis app (SCDA; Bulté & Onghena, 2013; De et al., 2020), from a total of 477 042 permutations (minimal possible p-value =  $2.1 \times 10^{-6}$ ). For each participant commencing the treatment in the group setting, a paired participant was allocated to the individual setting. At randomly designated start-dates of the relevant treatment-setting, paired participants switched settings. This specific design guaranteed that a minimum of three participants were always allocated to group treatment, thereby maintaining the number of participants required to ensure true in-group experience. The first treatment session took place on October 9<sup>th</sup>, 2019 for all participants. The last treatment session was held on December 13<sup>th</sup>, 2019. Three-month follow-up data collection, planned for March 2020, was disrupted by the COVID-19 pandemic and therefore was slightly delayed and took place from May to July 2020. No blinding was utilized.

The research was approved by the Nationally appointed Medical Ethical Committee at Maxima Medical Centre, in Veldhoven, the Netherlands, as part of a larger project (METC; NL63262.016.18) and registered at the Dutch Trial Register (NL8056). Reporting follows guidelines established at SCRIBE (Tate et al., 2016).

**Figure 3:** Randomized paired schedules for each of the six participants.



## 2.4. Measurements

### 2.4.1. Diary outcomes

End-of-day diary (EDD) data were collected from baseline to the end of the treatment (79 days). The diary was delivered through a purpose-built app installed on participants’ smartphones (mEMA; ilumivu, Inc., Cambridge, MA, USA; www.ilumivu.com). EDD assessments took place at 8pm every night (with a 4-hour time limit for completion), and consisted of 15 questions regarding different aspects of the tinnitus experience (i.e. annoyance, interference, distraction, anger, invasiveness, sadness, activity levels, fear, avoidance, masking, pleasantness) and overall well-being (i.e. happiness, anxiety, sleep quality, social interaction and stress). All questions were rated on a 7-point Likert scale (0-6), except for social interaction (“Who did you spend time with today?”), which presented the participant with seven mutually non-exclusive options (“nobody”, “partner”, “family”, “friends”, “colleagues”, “acquaintances”, and “strangers”).

### 2.4.2. Standardized outcomes and self-reported goals

Standardized outcomes were collected prior to treatment onset (T0), after the end of the full treatment protocol (T1) and follow-up (T2).

Tinnitus Functional Index (TFI; Meikle et al., 2012) is an assessment of tinnitus impairment in daily functioning. The TFI was specifically designed as an outcome measure

for clinical trials. Consisting of 25-items, this self-report tool relies on 10-point Likert scales to classify tinnitus-related impairment experienced over the previous week (e.g. “*What percentage of your time awake were you annoyed by your tinnitus?*”). Higher scores reflect greater impairment and a 13-point change is considered clinically meaningful. The tool is divided into eight subscales: intrusiveness; sense of control; cognitive interference; sleep disturbance; auditory difficulties due to tinnitus; interference in relaxation; quality of life; and emotional distress. The Dutch version of TFI was used (Rabau et al., 2014).

The Tinnitus Questionnaire (TQ; Hallam et al., 1988) is a widely used questionnaire for the assessment of tinnitus-severity (Hall et al., 2016). Fifty-two items are rated on a three-point scale for an overall score indicating tinnitus distress (e.g. “*The noises have affected my concentration*”). Meeus, Blaivie and Van de Heyning (2007) validated the TQ into Dutch. Despite a minimal change of 5 points being commonly considered clinically relevant (Kleinjung et al., 2007), a more conservative 12 point change approach is used at present (Hall et al., 2018).

The Fear of Tinnitus Questionnaire (FTQ; Cima et al., 2011) is a 17-item list of fear-related statements regarding tinnitus experience (e.g. “*I fear that my tinnitus is the result of a tumor*”). The tool is reliable, valid and responsive to clinical change (Fuller et al., 2019), with higher scores, obtained by the number of responder-selected items (judged to be true for them) in the list, being correlated with higher interference in daily life.

The Tinnitus Catastrophizing Scale (TCS; Cima et al., 2011) is an adaptation of the Pain Catastrophizing Scale (Sullivan et al., 1995). The TCS consists of 13-items on a 5-point Likert scale, the PCS aims at assessing catastrophizing cognitions and misinterpretations related to pain (e.g. “*I feel I can't go on*”). Items have been adapted by replacing pain related items to tinnitus equivalents. The TCS has been previously used in a large RCT on CBT for tinnitus patients (Cima et al., 2012) and it was shown that increased scores on the TCS are associated with decreased quality of life, increased tinnitus severity, increased fear for tinnitus and increased negative general affect .

The Chronic Tinnitus Acceptance Questionnaire (CTAQ; Moreland, 2007) is a 20-item self-report assessment on a 7-point Likert scale adapted from the Chronic Pain Acceptance Questionnaire (CPAQ; McCracken et al., 2004, 2005). This scale assesses participation in everyday activities despite tinnitus experience as well as the acceptance of the tinnitus experience without avoiding it (e.g. “*I am getting on with the business of living no matter what my level of tinnitus is*”). Lower scores reflect poorer levels of chronic tinnitus acceptance.

The Depression Anxiety and Stress Scale-21 (DASS-21; S. H. Lovibond & Lovibond, 1995) is a widely used measure of emotional wellbeing. With excellent psychometric properties (Antony et al., 1998), the tool consists of 21-items on a 4-point Likert scale to measure symptoms of depression, anxiety and stress with higher scores indicating worsening of depression (e.g. “*I felt I wasn't worth much as a person*”), anxiety (e.g. “*I felt I was close to panic*”) and stress (e.g. “*I found it difficult to relax*”). A revised Dutch version (de Beurs et al., 2001) was utilized in the current study.

Interpersonal Needs Questionnaire (INQ-15; Van Orden et al., 2012) is a 15-item, self-report questionnaire, which uses a 7-point Likert-type scale. The questionnaire measures an individual's social disability/inclusion through two subscales: perceived interpersonal measures of relationship distress (perceived alienation; e.g. “*These days, I feel like I belong*”) and self-perceived burden to others (e.g. “*These days the people in my life would be happier without me*”). Higher scores are associated with higher levels of perceived alienation and burden.

Personal goals of treatment outcomes were set through a semi-structured interview performed at T0. Participants were asked to describe their tinnitus disability (e.g. how they were negatively impacted by their tinnitus) and then set personal therapeutic goals by answering the question: “*What do you like to be improved at the end of the therapy?*”

Goal attainment and maintenance were confirmed at T1 and T2 by asking participants to describe the personal effect of treatment, whether tinnitus disability had diminished, and where was improvement noticed. Personal goals set at T0 were then discussed and progress evaluated.

## ***2.5. Visual Inspection***

Data collected from EDD was plotted through time with level (i.e. central tendency), variability and trend visually inspected. Broadened medians (Rosenberger & Gasko, 1983), an outlier resistant alternative to means and more robust against outliers, were used to assist visual inspection of levels. Variability was inspected with the aid of range lines drawn from the highest and lowest scores per phase. Least Squares regression was used to visually inspect trend per phase. Influence of outliers was reduced conservatively by removing a singularly occurring highest and/or lowest score on a particular item per phase per participant for the visual inspection of variability (trimmed range; Morley & Adams, 1991). This approach was also applied for the inspection of trend.

*Social interaction* was separately analysed with each answer (“partner”, “family”, “friends”, “colleagues”, “acquaintances”, and “strangers”) recorded into dichotomous variables (0 = no contact; 1 = contact). Each new variable was plotted through time and visually inspected for differences in level (broadened median) between phases B and C.

## ***2.6. Statistical Analysis***

### ***2.6.1. Effect size calculations***

Non-overlap of all pairs (NAP; Parker & Vannest, 2009) was utilized to calculate effect sizes between all phases. Cut-offs for NAP effect sizes can be interpreted as small (<.66), medium (>.65 and <.93) or large (NAP > .92). At a second step, trend was taken into account when calculating NAP between individual and group phases by utilizing Tau-U (Parker et al., 2011), a combination of Kendall’s rank correlation test (Tau) and Mann Whitney statistics (U).

### ***2.6.2. Randomization Tests (RT)***

Differences between individual (B) and group (C) treatment were calculated through Randomization Tests (RT) with Monte Carlo sampling (1000) and test statistics defined as  $|\bar{B}-\bar{C}|$  (the absolute difference of the means of B phase observations and the means of C phase observations) since no specific direction of change was hypothesized (Onghena & Edgington, 2005).

## ***2.7. Standardized outcomes and self-reported goals***

Effects of the treatments were tested by examining changes in scores on the TQ and TFI. Differences between T0 and T1, as well as T1 and T2 were calculated for each participant. Personalized goals of treatment outcomes at T0 were assessed at T1 and T2.

All other standardized outcomes (i.e. FTQ, TCS, CTAQ, DASS-21, INQ) from baseline are reported.

## ***2.8. Software and output***

All visual plots and analysis, as well as RTs, were calculated through the online SCDA app (Bulté & Onghena, 2013; De et al., 2020). NAP and Tau-U analyses utilized ‘Single Case Research: web based calculators for SCR analysis (Version 2.0)’ (Vannest et al., 2016).



Visual inspection plots of the end-of-day diary scores for all items are presented in the supplementary material.

### **3.Results**

Demographic data of the 6 participants at onset of the study are presented in Table 1. One participant (P5) commenced the use of hearing aid on October 2<sup>nd</sup> and was allowed to participate. Follow-up interview of this participant revealed that the hearing aid was seldomly used, specifically when in silence. Data for this participant was analysed as planned as changes were not attributable to the hearing aid by the participant. One female participant (50 years old) dropped out during the second week of treatment due to personal reasons unrelated to tinnitus. In order to maintain a minimal of 3 participants during group treatment, one patient (male, 49-years old) who was indicated for CBT for tinnitus and next on the waiting list, was added to and participated in group treatment, while exempted from data collection. Participant P2 missed the highest number of diary entries (26.3%) due to planned holidays. Nevertheless, EDD compliance rates ranged from 73.7-98.7% among participants. No adverse events were recorded during treatment.

**Table 1:** Demographic and baseline characteristics for Tinnitus Catastrophizing Scale (TCS), Fear of Tinnitus Questionnaire (FTQ), Chronic Tinnitus Acceptance Questionnaire (CTAQ), the sub-scales of the Depression Anxiety and Stress Scale-21 (DASS), and the Burden and Belong sub-scales of the Interpersonal Needs Questionnaire (INQ).

	Age Hearing Tinnitus aid			Onset	Location	Previous treatments	Type	TQ	TFI	TCS	FTQ	CTAQ	DASS			INQ	
	(time)	Duration	aid										Stress	Anx.	Dep.		Burden
<b>P1</b>	67	10 years	2 years	No coinciding factor	Both ears	Hearing aid (no effect)	High pitch	81	66.8	20	10	59	2	0	4	6	16
<b>P2</b>	62	14 months	12 years	Airplane landing	Both sides of head	Tinnitus Masking (no effect)	High pitch	48	68	25	6	74	6	4	8	6	12
<b>P3</b>	59	No hearing aid	9 months	Concussion	Both years	CBT focused on increased physical capacity	High pitch	61	74	30	9	43	6	2	0	6	23
<b>P4</b>	65	No hearing aid	15 years	No coinciding factor	In the head	Physiotherapy Chiropractic Osteopathy	Middle pitch	68	70	33	10	71	0	0	0	7	15
<b>P5</b>	66	1 week	12 years	Traffic (pedestrian) accident	Right ear	No treatment focused on tinnitus	High pitch	57	64	21	6	75	16	12	6	6	6

Note: Tinnitus Questionnaire (TQ); Tinnitus Functional Index (TFI); Tinnitus Catastrophizing Scale (TCS); Fear of Tinnitus Questionnaire (FTQ); Chronic Tinnitus Acceptance Questionnaire (CTAQ); Depression Anxiety and Stress Scale-21 (DASS); Interpersonal Needs Questionnaire (INQ).

### ***3.1. Diary scores***


#### ***3.1.1 Visual inspection***

Visual inspection of level of all variables (broadened median) revealed that all but one participant allocated to the ABC treatment setting order demonstrated no differences between phases or only a small improved level on all variables at the group (C) phase. However, participant P1, in the ABC order, revealed slight worsening during group treatment (phase C) for fear of tinnitus (Figure 4), while all other variables remained constant or improved as well. Participants in the alternative ACB order also revealed little to no change variables between phase levels. Most variables' levels either remained constant or slightly improved at the individual phase (B). On perceived happiness and activity level, participants' levels decreased in the individual phase (B). No other consistent pattern of change could be identified. All participants' level of overall perceived social contact remained constant throughout the treatment.


No discernible patterns in variability between phases can be seen. In one participant (P5) variability decreased in all but one item (activity level), while all other participants had no consistent pattern of change in variability across items. Tinnitus avoidance (Figure 5), interference, sadness and sleep quality scores were either stable or decreased in variability for all participants. Tinnitus pleasantness decreased in variability across all participants.

Similarly, clear trend patterns between ABC and ACB orders were lacking in the visual inspection. Tinnitus avoidance trend did not change between phases in all participants while all other items presented shifts for at least one participant (e.g. tinnitus annoyance; Figure 6). Participant (P5) trend improved on five items (activity level, anxiety, tinnitus pleasantness, sleep quality and stress), while participant (P2) trend improved on 4 items (tinnitus annoyance, anxiety, tinnitus distraction and tinnitus invasiveness) after changing treatment phase. Both participants' improvements were recorded in the second treatment phase – group and individual, respectively. All other participants displayed changes in trends between phases without any discernable pattern.


**Figure 4:** Fear of tinnitus broadened medians per participant per individual (B) and group (C) treatment phase.



**Figure 5:** Tinnitus avoidance variability (ranged lines) per participant per individual (B) and group (C) treatment phase.



**Figure 6:** Tinnitus annoyance trend (Least Squares regression) per participant per individual (B) and group (C) treatment phase.



### 3.1.2. Effect sizes

Low effect sizes (NAP) between baseline (A) and individual (B) treatment were similar to those between baseline (A) and group (C) treatment (Table 2). Without controlling for trend, data overlapped between individual and group treatment. By controlling for trend, Tau-U analysis revealed different significant changes between treatment orders (Table 3). Tinnitus anger, annoyance, fear, interference, invasiveness and sadness, as well as anxiety significantly improved at the group (C) phase for the ABC treatment order. Participants in the ACB order significantly improved tinnitus sadness and worsened in happiness and activity levels when in individual (B) treatment.

**Table 2:** Non-overlap of all pairs (NAP) effect sizes, and statistical significance based on Tau-U analyses without controlling for trend.

Item	A-B	A-C	B-C
Activity level†	.462**	.423**	.461**
Anger	.410**	.403**	.445**
Annoyance	.440**	.417**	.423**
Anxiety	.394**	.340**	.443**
Avoidance	.224*	.188	.436**
Distraction	.579**	.541**	.428**
Fear	.303**	.216*	.379**
Happiness†	.524**	.393**	.300**
Invasiveness	.664**	.570**	.368**
Interference	.376**	.314*	.438**
Pleasantness†	.532**	.506**	.446**
Sadness	.379**	.367**	.464**
Sleep quality†	.456**	.410**	.477**
Stress	.360**	.350**	.500**

Note: A – Baseline, B – Individual treatment, C – Group treatment

†inverted items

\*p-value < .05

\*\*p-value < .001

**Table 3:** Weighted Tau-U effect sizes, controlling for trend.

Item	ABC (n = 3)	ACB (n = 2)
Activity Level†	-.071	-.400**
Anger	-.200*	-.222
Annoyance	-.227*	-.152
Anxiety	-.205*	-.066
Avoidance	-.088	-.083
Distraction	-.154	-.031
Fear	-.304**	-.046
Happiness†	-.080	-.667**
Interference	-.282*	-.154
Invasiveness	-.219*	.103
Pleasantness†	-.067	-.129
Sadness	-.206*	-.286*
Sleep quality†	.064	.101
Stress	.009	.040

Note: A – Baseline, B – Individual treatment, C – Group treatment

†inverted items

\*p-value < .05

\*\*p-value < .001

### 3.1.3. Randomization Tests

No significant differences between group and individual treatments for each variable were found in RTs (minimal possible p-value =  $2.1 \times 10^{-6}$ ; all  $p > .05$ ).

## 3.2. Standardized outcomes and self-reported goals

Scores of the TQ and TFI were calculated for each participant (Table 4). A consistent pattern emerged in the results from pre- to post-treatment phase, which included both individual and group settings in different orders. Four out of five participants reported improvements from baseline to after treatment end (T1). The level of improvement exceeded minimally clinically important differences for four participants on the TQ (12-point change), and three out of five participants on the TFI (13-point change), with only one participant (P1) not recording improvements in the TQ or TFI from T0 to T1. Sequential order of treatment setting did not have an effect on overall treatment outcome. Furthermore, most objectives set by participants were successfully achieved by the end of treatment (Table 5).



Follow-up assessments supported the initial patterns found with the exception of one participant (P3), where a clinically meaningful worsening was found in TQ (38 points) and TFI (19.2 points). Another participant (P4) who showed clinically meaningful improvements after treatment further improved at follow-up, reporting clinically meaningful improvements in both TQ (22 points) and TFI (19.6 points) scores.

**Table 4:** Change in scores of the Tinnitus Questionnaire (TQ) and Tinnitus Functional Index (TFI)

	TQ		TFI	
	T1 score (change)	T2 score (change)	T1 score (change)	T2 score (change)
<b>P1</b>	87 (6)	77 (-10)	78.8 (12)	70 (-8.8)
<b>P2</b>	21 (-27*)	32 (11)	36 (-32*)	37.2 (1.2)
<b>P3</b>	47 (-14*)	85 (38*)	68.4 (-5.6)	87.6 (19.2*)
<b>P4</b>	49 (-19*)	27 (-22*)	43.2 (-26.8*)	23.6 (-19.6*)
<b>P5</b>	8 (-49*)	11 (3)	7.2 (-56.8*)	8 (0.8)

*Note:* T0 = baseline; T1 = after full treatment; T2 = follow-up

\*Clinically meaningful change [TQ (>12); TFI (>13)]

Table 5: Participants' self-reported goals and outcomes.

	<b>Goals set before treatment (T0)</b>	<b>Assessment of goals after treatment (T1)</b>	<b>Follow-up (T2)</b>
<b>P1</b>	<p>Improve sleep</p> <p>Reduce fatigue</p> <p>Improve concentration</p> <p>Resume running</p> <p>Reduce neck pain complaints</p>	<p>Tinnitus disturbance remained unchanged.</p> <p>Concentration did not improve. Running resumed.</p> <p>Participant continued to apply tinnitus avoidance strategies.</p>	<p>Overall feeling that tinnitus complaints did not improve or worsened. Sleep improved with more sleeping hours. Other goals did not improve. Gave up on running and switched to walking. Did not practice techniques from treatment (i.e. exposure or relaxation)</p>
<b>P2</b>	<p>Improve concentration.</p> <p>Improve sleep.</p> <p>Improve tinnitus coping.</p>	<p>Participant is generally less concerned with tinnitus and reports more acceptance towards it (in the words of the participant: "I don't fight it anymore"). Participant experiences tinnitus more clearly in silence, but this subsides and is no longer disturbing. Sleep and concentration improved.</p>	<p>All positive changes remained. Practices exposure daily. Light sleep problems not associated with tinnitus.</p>
<b>P3</b>	<p>Less erratic tinnitus and heightened control</p> <p>Diminish influence on function</p> <p>Reduce "fight against tinnitus"</p>	<p>Feels more "present in the moment" and allows himself experience range of emotions including aversive emotions (e.g. less experiential avoidance). Participant reports using techniques learned at treatment sessions. "Fighting" tinnitus is slightly reduced, although tinnitus is still erratic.</p>	<p>Worsening of tinnitus complaints since T1. Increased perception of tinnitus triggers irritability, fatigue and concentration problems. Increased tinnitus complaints associated with increased workload. Increased work stress due to COVID-19 crisis and rest complaints due to a concussion potentially playing a role. Fear reactions to tinnitus perception and "fighting" tinnitus decreased. Frequently practices exposure.</p>
<b>P4</b>	<p>Improve sleep</p> <p>Improve concentration</p> <p>Reduce burnout feelings</p>	<p>Participant reports less experiential avoidance and open to aversive emotions. Participant allows himself to experience tinnitus through exposure exercises. Sleep and concentration improved. Increase in assertiveness associated with reduced burnout</p>	<p>Sleep deteriorated due to rumination concerning COVID-19 crisis. Tinnitus did not play a role in sleep problems. All other improvements remained. Frequently practices exposure.</p>
<b>P5</b>	<p>Restore balance in mood</p> <p>Regain sense of humor</p> <p>Enjoy silence again</p> <p>Balance vigor</p>	<p>All objectives were achieved. Tinnitus characteristics remained equal while emotional response changed. Mood is positive again. Regained sense of humor and balanced vigor. Exercises learned during therapy are applied daily.</p>	<p>All positive changes remained. Often practices techniques learned during treatment.</p>

## 4. Discussion

The current study aims to examine the differences in patient improvements when comparing individual versus group treatment delivery of CBT for chronic disabling tinnitus. The SCED employed is specifically suited to enable direct comparisons between individual and group treatment settings within subjects. Group based RCT's fail to access the high inter- and intra-individual variability found in tinnitus experience (Henry et al., 2012; Schlee et al., 2016). The SCED allowed us to control for individual conditions which accounted for participants idiosyncrasies. Moreover, SCED enabled us to follow participants undergoing both treatment conditions of interest (group and individual CBT), allowing to test differences as a result of treatment type to be carried out in detail (e.g. assessment of differences in trend, variability and level).

First, beneficial effects of CBT were confirmed on the standardized outcome measures collected before, and after treatment, and during follow-up. Moreover, clinically meaningful improvements on standardized tinnitus assessments (i.e. TQ and TFI) were found in four out of five participants. The follow-up assessments were delayed with 3 months due to the COVID-19 pandemic. Nevertheless, three of the participants improvements were not lost, with one participant improving further after end of treatment. Semi-structured clinical interviews confirmed these changes, with all but one participant reaching the personalized goals set at the beginning of treatment. These findings confirm the effectiveness of CBT for tinnitus, established by Cima et al. (2012), and are in line with current literature on CBT (Fuller et al., 2020) and guidelines (Cima et al., 2019) for tinnitus treatment.

Visual inspection of the data, as well as randomization tests between phases indicated no significant differences between group and individual treatment. Analyses of data overlap, while accounting for trend, revealed low effect sizes but significant differences between treatments. Participants who started treatment individually and subsequently joined group treatment, showed improvements in the latter when compared with the former. Half of the fourteen variables measured held significant differences between phases (tinnitus anger, annoyance, fear, interference, invasiveness and sadness, as well as anxiety), favouring group treatment. Participants who underwent the opposite treatment order, group treatment followed by individual setting later, only improved further in scores on tinnitus sadness during individual treatment compared to group treatment. Thus, while group setting increased benefits from treatment on half the variables, individual setting only provided increased improvements on tinnitus sadness. More importantly,

participants worsening happiness and activity levels was observed during subsequent individual treatment, when compared to group treatment. Despite the worsening captured by the EDD, these participants still recorded overall improvements in the standardized outcome measures and clinical interview (from pre- to post-treatment).

The small differences found between treatment settings is indicative of the potential benefits of group CBT. Unlike individual treatment, group-based treatment may enable social learning, such as observing another patient expose themselves through the feared stimulus (e.g. loud noise, tinnitus) and challenging the expected outcome (e.g. anxiety, loss of control). Social learning allows for the development of knowledge without the need for first-hand experience, avoiding possible risks of these experiences. Some studies show that seeing someone else expose themselves to a feared stimulus (i.e. vicarious extinction of fear) can lead to superior fear reduction when compared to standard extinction (Golkar et al., 2013, 2016).

While most participants improved, one participant (P1) did not benefit from the treatment in either individual or group setting. Moreover, P1 held the highest fear of tinnitus, measured through the standardized questionnaire (i.e. FTQ) and EDD, where high variability was also observed. Fear of tinnitus is known to mediate recovery (Cima et al., 2017) and partly explains treatment efficacy. Tinnitus avoidance, as reported by P1, did not abate during treatment as measured by the EDD and interview, which might explain why fear did not decrease, since the use of safety behaviours to avoid exposure to the feared stimuli may hinder fear extinction (Lovibond et al., 2009). Safety behaviours during tinnitus exposure are difficult to assess, since these might be largely cognitive or interoceptive and therefore cannot be easily observed and controlled for (e.g. patients may use imagery to distract themselves from the tinnitus experience during an exposure exercise). As such, exposure to tinnitus may better resemble Interoceptive Exposure (IE), where the more covert cognitive strategies to avoid the full experience of fear (e.g. relaxation, distraction) may disrupt fear extinction (as reported by P1).

As parallels from chronic pain research may be drawn and inspire the tinnitus field (e.g. the application of the FA model), fundamental research on the underlying mechanisms of chronic tinnitus are in their infancy. Associative fear learning paradigms have been developed and employed in the chronic pain field for decades, enabling the underlying mechanisms of change to be studied (for a review on the development and employment of fear conditioning in chronic pain see Meulders, 2020; Vlaeyen & Crombez, 2020).

While we are able to observe change in tinnitus experience through CBT, we are only able to speculate on the underlying mechanisms and associative learning models.

Beyond the inherent difficulty in controlling for covert safety behaviours in efficacious exposure exercises, a limitation may be found in the uniqueness of the sample: i.e. five male participants aged between 59 and 66 years old. This homogeneous sample serves as a double-edged sword. Naturally, our findings may not be generalizable to other tinnitus populations at the moment. On the other hand, our outcomes represent robust findings in a specific tinnitus population. The effectiveness of CBT for this specific population is supported, with group setting presenting a more beneficial treatment type, regardless of hearing aid use or tinnitus duration and location. Treatment response differences between gender have been previously observed, with men more responsive to the combined treatment of Tinnitus Retraining Therapy (TRT) and CBT (Van der Wal et al., 2020). As such, replication of the protocol with women is a natural step towards tailoring CBT for tinnitus.

## **5. Conclusion**

CBT for tinnitus is an effective treatment that may be offered in a variety of different methods. Inspired by recent interest in more personalized research and treatment development (Schork, 2015; Senn, 2018), a SCED was used to explore the differences between group and individual treatment delivery settings of an established CBT for tinnitus protocol (Cima et al., 2012). To our knowledge the current study is the first to directly compare group versus individual CBT in tinnitus patients, revealing that group treatment is potentially more beneficial to participants. While the benefits were observable though small, underlying mechanism of change (e.g. associative learning models) are yet to be fully explored through fundamental research in the tinnitus field.

## References

- Antony, M. M., Bieling, P. J., Cox, B. J., Enns, M. W., & Swinson, R. P. (1998). Psychometric properties of the 42-item and 21-item versions of the Depression Anxiety Stress Scales in clinical groups and a community sample. *Psychological Assessment, 10*(2), 176–181. <https://doi.org/10.1037/1040-3590.10.2.176>
- Bulté, I., & Onghena, P. (2013). The Single-Case Data Analysis Package: Analysing Single-Case Experiments with R Software. *Journal of Modern Applied Statistical Methods, 12*(2), 450–478. <https://doi.org/10.22237/jmasm/1383280020>
- Cima, R. F. F., Crombez, G., & Vlaeyen, J. W. S. (2011). Catastrophizing and Fear of Tinnitus Predict Quality of Life in Patients With Chronic Tinnitus. *Ear and Hearing, 32*(5), 634–641. <https://doi.org/10.1097/AUD.0b013e31821106dd>
- Cima, R. F. F., Maes, I. H., Joore, M. A., Scheyen, D. J. W. M., El Refaie, A., Baguley, D. M., Anteunis, L. J. C., van Breukelen, G. J. P., & Vlaeyen, J. W. S. (2012). Specialised treatment based on cognitive behaviour therapy versus usual care for tinnitus: a randomised controlled trial. *The Lancet, 379*(9830), 1951–1959. [https://doi.org/10.1016/S0140-6736\(12\)60469-3](https://doi.org/10.1016/S0140-6736(12)60469-3)
- Cima, R. F. F., Mazurek, B., Haider, H., Kikidis, D., Lapira, A., Noreña, A., & Hoare, D. J. (2019). A multidisciplinary European guideline for tinnitus: diagnostics, assessment, and treatment. *HNO, February 2018*. <https://doi.org/10.1007/s00106-019-0633-7>
- Cima, R. F. F., van Breukelen, G., & Vlaeyen, J. W. S. (2017). Tinnitus-related fear: Mediating the effects of a cognitive behavioural specialised tinnitus treatment. *Hearing Research, 1*–12. <https://doi.org/10.1016/j.heares.2017.10.003>
- de Beurs, E., Van Dyck, R., Marquenie, L. A., Lange, A., & Blonk, R. W. B. (2001). De DASS: Een vragenlijst voor het meten van depressie, angst en stress [The DASS: A questionnaire for the measurement of depression, anxiety, and stress]. *Gedragstherapie, 34*(1), 35–53.
- De, T. K., Michiels, B., Vlaeyen, J. W. S., & Onghena, P. (2020). *Shiny SCDA*.
- Fuller, T. E., Cima, R. F. F., Van den Bussche, E., & Vlaeyen, J. W. S. (2019). The Fear of Tinnitus Questionnaire. *Ear and Hearing, 40*(6), 1467–1477. <https://doi.org/10.1097/AUD.0000000000000728>
- Fuller, T. E., Cima, R., Langguth, B., Mazurek, B., Vlaeyen, J. W. S., & Hoare, D. J. (2020). Cognitive behavioural therapy for tinnitus. *Cochrane Database of Systematic Reviews, 4*. <https://doi.org/10.1002/14651858.CD012614.pub2>
- Golkar, A., Haaker, J., Selbing, I., & Olsson, A. (2016). Neural signals of vicarious extinction learning. *Social Cognitive and Affective Neuroscience, 11*(10), 1541–1549. <https://doi.org/10.1093/scan/nsw068>
- Golkar, A., Selbing, I., Flygare, O., Öhman, A., & Olsson, A. (2013). Other People as Means to a Safe End: Vicarious Extinction Blocks the Return of Learned Fear. *Psychological Science, 24*(11), 2182–2190. <https://doi.org/10.1177/0956797613489890>
- Hall, D. A., Haider, H., Szczepek, A. J., Lau, P., Rabau, S., Jones-Diette, J., Londero, A., Edvall, N. K., Cederroth, C. R., Mielczarek, M., Fuller, T., Batuecas-Caletrio, A., Brueggemen, P., Thompson, D. M., Norena, A., Cima, R. F. F., Mehta, R. L., & Mazurek, B. (2016). Systematic review of outcome domains and instruments used in clinical trials of tinnitus treatments in adults. *Trials, 17*(1), 270. <https://doi.org/10.1186/s13063-016-1399-9>

- Hall, D. A., Mehta, R. L., & Argstatter, H. (2018). Interpreting the Tinnitus Questionnaire (German version): what individual differences are clinically important? *International Journal of Audiology*, *57*(7), 553–557. <https://doi.org/10.1080/14992027.2018.1442591>
- Hallam, R. S., Jakes, S. C., & Hinchcliffe, R. (1988). Cognitive variables in tinnitus annoyance. *The British Journal of Clinical Psychology*, *27* ( Pt 3)(pt 3), 213–222. <http://www.ncbi.nlm.nih.gov/pubmed/3191301>
- Henry, J. A., Galvez, G., Turbin, M. B., Thielman, E. J., Mcmillan, G. P., & Istvan, J. A. (2012). *Pilot Study to Evaluate Ecological Momentary Assessment of Tinnitus (Henry et al, 2012)*. 279–290.
- Kleinjung, T., Steffens, T., Sand, P., Murthum, T., Hajak, G., Strutz, J., Langguth, B., & Eichhammer, P. (2007). Which tinnitus patients benefit from transcranial magnetic stimulation? *Otolaryngology–Head and Neck Surgery*, *137*(4), 589–595. <https://doi.org/10.1016/j.otohns.2006.12.007>
- Kleinstäuber, M., Jasper, K., Schweda, I., Hiller, W., Andersson, G., & Weise, C. (2013). The Role of Fear-Avoidance Cognitions and Behaviors in Patients with Chronic Tinnitus. *Cognitive Behaviour Therapy*, *42*(2), 84–99. <https://doi.org/10.1080/16506073.2012.717301>
- Krasny-Pacini, A., & Evans, J. (2018). Single-case experimental designs to assess intervention effectiveness in rehabilitation: A practical guide. *Annals of Physical and Rehabilitation Medicine*, *61*(3), 164–179. <https://doi.org/10.1016/j.rehab.2017.12.002>
- Lethem, J., Slade, P. D., Troup, J. D., & Bentley, G. (1983). Outline of a Fear-Avoidance Model of exaggerated pain perception--I. *Behaviour Research and Therapy*, *21*(4), 401–408. <http://www.ncbi.nlm.nih.gov/pubmed/6626110>
- Lovibond, P. F., Mitchell, C. J., Minard, E., Brady, A., & Menzies, R. G. (2009). Safety behaviours preserve threat beliefs: Protection from extinction of human fear conditioning by an avoidance response. *Behaviour Research and Therapy*, *47*(8), 716–720. <https://doi.org/10.1016/j.brat.2009.04.013>
- Lovibond, S. H., & Lovibond, P. F. (1995). *Manual for the Depression Anxiety & Stress Scales* (2nd Ed.). Psychology Foundation.
- Maes, I. H. L., Cima, R. F. F., Vlaeyen, J. W., Anteunis, L. J. C., & Joore, M. A. (2013). Tinnitus: A cost study. *Ear and Hearing*, *34*(4), 508–514. <https://doi.org/10.1097/AUD.0b013e31827d113a>
- McCracken, L. M., Vowles, K. E., & Eccleston, C. (2004). Acceptance of chronic pain: component analysis and a revised assessment method. *Pain*, *107*(1), 159–166. <https://doi.org/10.1016/j.pain.2003.10.012>
- McCracken, L. M., Vowles, K. E., & Eccleston, C. (2005). Acceptance-based treatment for persons with complex, long standing chronic pain: a preliminary analysis of treatment outcome in comparison to a waiting phase. *Behaviour Research and Therapy*, *43*(10), 1335–1346. <https://doi.org/10.1016/j.brat.2004.10.003>
- Meeus, O., Blavie, C., & Van de Heyning, P. (2007). Validation of the Dutch and the French version of the Tinnitus Questionnaire. *B-ENT*, *3 Suppl 7*, 11–17. <http://www.ncbi.nlm.nih.gov/pubmed/18228680>

- Meikle, M. B., Henry, J. A., Griest, S. E., Stewart, B. J., Abrams, H. B., McArdle, R., Myers, P. J., Newman, C. W., Sandridge, S., Turk, D. C., Folmer, R. L., Frederick, E. J., House, J. W., Jacobson, G. P., Kinney, S. E., Martin, W. H., Nagler, S. M., Reich, G. E., Searchfield, G., ... Vernon, J. A. (2012). The Tinnitus Functional Index. *Ear and Hearing, 33*(2), 153–176. <https://doi.org/10.1097/AUD.0b013e31822f67c0>
- Meulders, A. (2020). Fear in the context of pain: Lessons learned from 100 years of fear conditioning research. *Behaviour Research and Therapy, 131*(October 2019), 103635. <https://doi.org/10.1016/j.brat.2020.103635>
- Moore, B. A., Moring, J. C., Hale, W. J., & Peterson, A. L. (2019). Incidence Rates of Tinnitus in Active Duty Military Service Members Between 2001 and 2015. *American Journal of Audiology, 28*(4), 866–876. [https://doi.org/10.1044/2019\\_AJA-19-0029](https://doi.org/10.1044/2019_AJA-19-0029)
- Moreland, J. (2007). *Illness representations, acceptance, coping and psychological distress in chronic tinnitus*. University of Leicester.
- Morley, S., & Adams, M. (1991). Graphical analysis of single-case time series data. *British Journal of Clinical Psychology, 30*(2), 97–115. <https://doi.org/10.1111/j.2044-8260.1991.tb00926.x>
- Ongghena, P., & Edgington, E. S. (2005). Customization of Pain Treatments. *The Clinical Journal of Pain, 21*(1), 56–68. <https://doi.org/10.1097/00002508-200501000-00007>
- Parker, R. I., & Vannest, K. (2009). An Improved Effect Size for Single-Case Research: Nonoverlap of All Pairs. *Behavior Therapy, 40*(4), 357–367. <https://doi.org/10.1016/j.beth.2008.10.006>
- Parker, R. I., Vannest, K. J., Davis, J. L., & Sauber, S. B. (2011). Combining Nonoverlap and Trend for Single-Case Research: Tau-U. *Behavior Therapy, 42*(2), 284–299. <https://doi.org/10.1016/j.beth.2010.08.006>
- Rabau, S., Wouters, K., & Van de Heyning, P. (2014). Validation and translation of the Dutch tinnitus functional index. *B-ENT, 10*(4), 251–258. <http://www.ncbi.nlm.nih.gov/pubmed/25654947>
- Rosenberger, J. L., & Gasko, M. (1983). Comparing location estimators: Trimmed means, medians and trimean. In D. Hoaglin, F. Mosteller, & J. Tukey (Eds.), *Understanding robust and exploratory data analysis* (pp. 297–336).
- Schlee, W., Pryss, R. C., Probst, T., Schobel, J., Bachmeier, A., Reichert, M., & Langguth, B. (2016). Measuring the moment-to-moment variability of Tinnitus: The TrackYourTinnitus smart phone app. *Frontiers in Aging Neuroscience, 8*(DEC), 1–8. <https://doi.org/10.3389/fnagi.2016.00294>
- Schork, N. J. (2015). Personalized medicine: Time for one-person trials. *Nature, 520*(7549), 609–611. <https://doi.org/10.1038/520609a>
- Senn, S. (2018). Statistical pitfalls of personalized medicine. *Nature, 563*(7733), 619–621. <https://doi.org/10.1038/d41586-018-07535-2>
- Stohler, N. A., Reinau, D., Jick, S. S., Bodmer, D., & Meier, C. R. (2019). A study on the epidemiology of tinnitus in the United Kingdom. *Clinical Epidemiology, Volume 11*, 855–871. <https://doi.org/10.2147/CLEPS213136>
- Sullivan, M. J. L., Bishop, S. R., & Pivik, J. (1995). The Pain Catastrophizing Scale: Development and validation. *Psychological Assessment, 7*(4), 524–532. <https://doi.org/10.1037/1040-3590.7.4.524>



- Tate, R. L., Perdices, M., Rosenkoetter, U., McDonald, S., Togher, L., Shadish, W., Horner, R., Kratochwill, T., Barlow, D. H., Kazdin, A., Sampson, M., Shamseer, L., & Vohra, S. (2016). The Single-Case Reporting Guideline In BEhavioural Interventions (SCRIBE) 2016 Statement. *Archives of Scientific Psychology, 4*(1), 10–31. <https://doi.org/10.1037/arc0000027>
- Van der Wal, A., Luyten, T., Cardon, E., Jacquemin, L., Vanderveken, O. M., Topsakal, V., Van de Heyning, P., De Hertogh, W., Van Looveren, N., Van Rompaey, V., Michiels, S., & Gilles, A. (2020). Sex Differences in the Response to Different Tinnitus Treatment. *Frontiers in Neuroscience, 14*. <https://doi.org/10.3389/fnins.2020.00422>
- Van Orden, K. A., Cukrowicz, K. C., Witte, T. K., & Joiner, T. E. (2012). Thwarted belongingness and perceived burdensomeness: Construct validity and psychometric properties of the Interpersonal Needs Questionnaire. *Psychological Assessment, 24*(1), 197–215. <https://doi.org/10.1037/a0025358>
- Vannest, K. J., Parker, R. I., Gonen, O., & Adiguzel, T. (2016). *Single Case Research: web based calculators for SCR analysis (2.0)*. Texas A&M University. <http://www.singlecaseresearch.org/calculators/tau-u>
- Vlaeyen, J. W. S., & Crombez, G. (2020). Behavioral Conceptualization and Treatment of Chronic Pain. *Annual Review of Clinical Psychology, 16*, 187–212. <https://doi.org/10.1146/annurev-clinpsy-050718-095744>
- Vlaeyen, J. W. S., & Linton, S. J. (2000). Fear-avoidance and its consequences in chronic musculoskeletal pain: a state of the art. *Pain, 85*(3), 317–332. <http://www.ncbi.nlm.nih.gov/pubmed/10781906>
- Vlaeyen, J. W. S., & Linton, S. J. (2012). Fear-avoidance model of chronic musculoskeletal pain: 12 years on. *Pain, 153*(6), 1144–1147. <https://doi.org/10.1016/j.pain.2011.12.009>



# CHAPTER 6

---

## General Discussion

## **To see the forest from the trees**

Three main areas of interest for closer investigation were identified from the current state of the tinnitus field in Chapter 1: (1) the testing of a tinnitus fear conditioning paradigm in humans; (2) improving tinnitus assessment; and (3) the investigation of CBT components in order to better understand what works best for whom. From these, 4 specific research questions were drawn, and each question was addressed in a dedicated chapter. They were: (1) the adaptation/creation of an associative learning model for the field of tinnitus in order to establish a replicable fear learning paradigm with healthy human participants; (2) the investigation of whether Ecological Momentary Assessment (EMA) negatively affects tinnitus experience; (3) the comparison between EMA and End-of-Day Diary (EDD) in measuring tinnitus experience; and, relating to tinnitus treatment, (4) contrasting and comparing individual to group-based Cognitive Behavioural Therapy (CBT) for tinnitus. Next, we provide an overview of the research findings followed by a critical discussion and future directions.

### *Tinnitus fear conditioning*

The tinnitus Fear Avoidance (FA) model explains the role of fear in the development and maintenance of Chronic Disabling Tinnitus. Clinical studies have shown the model to be valid, nonetheless, replicable experimental paradigms to investigate underlying associative learning processes in with human participants were lacking. A novel differential fear conditioning paradigm was introduced (Chapter 2). We repeatedly presented a neutral sound (CS+), conceptualized to be tinnitus-like, followed by a short delay and the same sound at higher intensity (US) to healthy human participants. The CS+ produced fear responses (seen in self-reported expectancy ratings) in participants in absence of the US. A different, but perceptually equivalent auditory stimulus that was not paired with the US (CS-) did not elicit these fear responses. In other words, during the first phase (i.e. acquisition phase), fear of a neutral sound was acquired, after pairing it with the same louder sound. The finding provides support that classical conditioning, increased fear responding towards a neutral sound can be acquired, which may provide support for the key role of fear in triggering the pathological cycle of the FA model. Classical conditioning has been previously proposed as the underlying mechanism of the Neurophysiological model, although definition of the learning components (e.g. CS definition) have been vague and unspecific (Baguley et al., 2013). Moreover, conditioning research focused solely on animal paradigms to study tinnitus generation (Jastreboff, 1990; Jastreboff et al., 1988, 1994).

At a second phase (i.e. extinction phase) of the paradigm, the acquired fear responses decreased with repeated presentation of CS+ without US. Surprisingly, CS- evoked fear responses at the beginning of this phase also decreased. This lack of differentiation between CS+ and CS- was also observed in the third phase of the experiment (i.e. reinstatement phase), where reinstatement was tested through a single non-contingent presentation of the US. The unexpected lack of differentiation was postulated to be a consequence of conducting Acoustic Stapedial Reflex Threshold (ASRT) and Loudness Discomfort Level (LDL) tests between experiment phases for assessment purposes. These tests include presentation of acoustic probes that were similar to the CS+/- and US, potentially interfering with the CS-US associations and thereby diminishing the fear responses to the US. Moreover, the change in context, necessary to conduct the tests between phases, potentially decreased the strength of the association formed at acquisition. Beyond reconsidering the use of ASRT and LDL tests between the experimental phases, improvement of the paradigm in future studies can be achieved through limiting change of context (i.e. moving between rooms) and augmenting the number of trials during the acquisition phase, thereby increasing opportunities for a more robust and enduring fear-learning.

Findings, while not robustly establishing a fear learning paradigm, are promising and do not rule out the role of classical conditioning as a possible origin for Chronic Disabling Tinnitus. Despite the several limitations of the study, fear learning was observed in the fear expectancy ratings. On the other hand, physiological responses were inconsistent, indicating a quick habituation and indiscriminate fear learning to CS+ and CS-. Again, the limitations presented by the ASRT and LDL measurements as well as the change in context may have unduly influenced responses, specifically those after the acquisition phase (where change of context happened more often). Nevertheless, physiological measurements, including the ones used in the current study (Skin Conductance Response and Heart Rate acceleration) may be limited in explaining tinnitus disability. Given a threatening US, self-report measures of fear have stronger diagnostic, predictive and construct validity when compared to physiological measurements (Boddez et al., 2013). Therefore, an updated paradigm might take a step back in order to create a simpler approach with a lower number of outcome measures (i.e. threat expectancy only) and increased power through increased assessment points, number of trials and participants.

### *(Re)assessing tinnitus*

Self-report tools that are widely used to assess tinnitus experience may be susceptible to bias (i.e. recall bias, reading difficulty, and current psychological state). Ecological Momentary Assessments (EMA) increase ecological validity and reduce the influence of bias. However, the unwarranted attention towards the tinnitus experience has been thought to be detrimental to the patients. Increased awareness of the tinnitus percept has been shown to be associated with increased Chronic Disabling Tinnitus (e.g. lower quality of life; Cima et al., 2011). The use of EMA may exacerbate awareness of tinnitus in severe tinnitus sufferers by increasing attention to tinnitus, since it requires responding and reflecting on disability intensively on a daily basis. Chapter 3 investigated these potential negative effects through a Single-Case Experimental Design (SCED).

Our findings supported and expanded on previous findings regarding EMA reactivity in tinnitus sufferers (Henry et al., 2012; Schlee et al., 2016) as participants did not show meaningful negative reactions to EMA. Interestingly, slight improvements in tinnitus experience (i.e. decreased tinnitus avoidance, annoyance, interference, fear, sadness, distraction, masking and anger), and overall well-being (i.e. happiness, stress, sleep quality, activity level, anxiety) were observed after EMA introduction. Important to note is that EMA reactivity was only tested in participants with severe tinnitus distress. It remains to be explored if such findings are replicable in individuals with varying levels of tinnitus distress. Furthermore, attentional processes which could have further helped explain the findings, were not measured. More importantly, the surprising potential benefits of EMA remain to be explored. As research into internet and app delivered interventions is growing exponentially (e.g. Beukes et al., 2017, 2018; Hesser et al., 2012), and given the impact of the COVID-19 crisis for tinnitus patients (Beukes et al., 2020), the potential of EMA to become part of a treatment is worth exploring.

Replication studies including participants with different levels of tinnitus-severity should include whether there is an added burden on participants while using EMA, posing ethical and theoretical concerns (e.g. increased screen time). End-of-Day Diaries (EDD) could be a promising alternative to EMA by providing equivalent assessments at a lesser cost, specifically in cases where EMA data is aggregated in the form of daily scores (e.g. means).

As direct comparisons between EMA and EDD assessments in tinnitus patients had not been carried out, Chapter 4 compared EDD mean values to EMA means. Furthermore,

EMA gathered close in time to EDD was compared with earlier-in-the-day EMA and EDD. The study focused on testing if EDD accurately reflected the overall daily picture, as illustrated by EMA. Our results indicated that EDD and EMA provide similar data, though some significant differences were found. EDD measures significantly differed from EMA daily averages for six out of eleven variables: tinnitus anger, anxiety, tinnitus invasiveness, tinnitus pleasantness, tinnitus sadness, and stress. The differences indicated worse scores (e.g. lower tinnitus pleasantness, higher anxiety) on EDD. This is in line with previous literature revealing that longer recall periods are associated with worse symptom/experience recollection (Miron-Shatz et al., 2009). Despite statistical significances, the effects were small and may be attributed to the large number of data entries (4,732). Such minor differences may not justify EMA as the measurement of choice given the added burden to participants.

These findings are limited by the homogeneity of the population and external validity would be strengthened by successful replications in different age groups and in participants with varying levels of tinnitus distress. Furthermore, a more extensive set of tinnitus experiences, beyond the ones utilized in the chapter (e.g. perceived tinnitus control) can potentially produce different results due to the heterogeneity of experiences. While a preference for EDD is stated, EMA use is not to be dismissed. EMA remains a powerful tool that may be of specific benefit in understanding daily cyclical patterns in tinnitus experience (e.g. fluctuations from morning to night). Beyond these patterns, EMA may further elucidate the dynamic relationship between emotions (e.g. fear), individuals, and their environment (Shiffman et al., 2008). Reactions to specific situations and granular level insight into temporal chain reactions have the power to pinpoint clinically relevant triggers. Future research must focus on the relevant use of EMA or EDD pending research objectives and associated costs.

Beyond replicating the study with a different and wider population, other limitations of the original research may be further addressed. The use of questions developed with the patient and potentially tailored to a patient's individual need and acknowledging patient preferred outcomes is possible through the flexibility of EMA and EDD methodologies (i.e. Hall et al., 2018). Such tailoring may, for example, provide answers on how to accurately measure overt avoidance behaviour. Such as GPS location (e.g. avoidance of known restaurants, bars, concert halls), as well as the use of smartphone microphones to measure noise exposure. The use of an adapted EMA delivery, that is prompted by the participant when he/she recognizes a trigger or catastrophic thought, has potential

when integrating EMA in ongoing interventions. Naturally, while promising and exciting methodological possibilities are theoretically possible, technological and budgetary constraints may limit the adequate deployment of such tools. The use of smartphone microphones, for example, is limited to the quality of each individual smartphone used and dependent on the quality of the hardware (e.g. sensitivity of the microphone, battery life) and software (e.g. operating system) employed by the manufacturer, making it difficult to standardize and compare.

Chapters 3 and 4 highlight the potential of alternative assessment methods to standardized questionnaires, namely individualized EMA and EDD. Findings highlight that EMA can be used with severe tinnitus sufferers without negative consequences and, when EMA concerns are present, EDD is an adequate alternative for tinnitus assessment. Both represent more ecologically valid methods of assessment. Beyond a reduction in bias, these methods facilitate the use of SCEDs. As observed in this dissertation (Chapters 3 and 5), SCED relies on the frequent and repeated assessment of participants to establish a robust baseline condition. These designs are particularly useful in the field of tinnitus as they account for the high heterogeneity of tinnitus experiences. Coupling EMA or EDD with designs that focus on within-subject change and fluctuation, specifically SCED, provide promising avenues in tailored treatment research.

### *CBT for tinnitus*

Tinnitus can be treated successfully with CBT. While positive outcomes have been reported in both group and individual settings, no direct comparison with the same treatment protocol has been carried out (Fuller et al., 2020). Group treatment may provide added benefits, such as decreased treatment delivery costs and higher speed of recovery due to increased social learning opportunities. Chapter 5 aimed at comparing a successful CBT for tinnitus treatment protocol (Cima et al., 2012) between group and individual delivery setting. This chapter revealed that participants experienced increased benefits when treated in group, compared to when treated on an individual basis, as measured on over half of the variables (tinnitus anger, annoyance, fear, interference, invasiveness and sadness, as well as happiness and anxiety). Moreover, participants who switched from group to individual treatment were less happy and had lower activity levels when compared to those who went the opposite direction (from individual to group treatment).



Overall, CBT proved an effective treatment for tinnitus up to nearly six-month follow up, supporting previous findings (Cima et al., 2012, 2019; Fuller et al., 2020). Nonetheless, one single participant did not record clinically significant improvements – on neither the Tinnitus Questionnaire (TQ; Hallam et al., 1988), the Tinnitus Functional Index (TFI; Meikle et al., 2012) or self-reported goals. The participant uniquely registered higher fear levels and variability as well as self-reported nonadherence to the treatment component, namely exposure (i.e. interoceptive avoidance of exposure sessions). These observations are consistent with the FA model and appear to underscore the beneficial role of decreased fear by exposure on treatment outcomes. Conversely, the lack of treatment response also highlights the importance of individualized medicine. Studying specific effects of each individual treatment component (e.g. exposure, relaxation, psychoeducation) as well as possible combinations, contributes to achieving the most efficacious treatment geared to specific groups and tailored to the individual needs of each patient.

Cognitive Behavioural Treatments for tinnitus should continue to be explored, as we strive to understand what component works best for whom. Chapter 5 focused on only one treatment variant (i.e. delivery setting) and results were limited to men in the sixth decade of their lives. Replicability of findings in samples including women and other age groups must be carried out.

## Planting seeds

Despite growing evidence of the prominent role of fear in the acquisition, development and maintenance of Chronic Disabling Tinnitus, research within this perspective is scarce. A quick search of the literature (in March 1<sup>st</sup>, 2021) reveals that 4 articles have been published with the terms “fear” and “tinnitus” in the title, compared to over 950 in a similar search with the terms “fear” and “pain” (dating back to 1975). The earliest of the articles within the tinnitus field was published in a peer reviewed journal 10 years ago (Cima et al., 2011). During the same period 412 (of the 950) articles were published in the field of pain. Resistance to adopting a psychological tinnitus model may be encountered as patients and healthcare providers call for cures and hope for relief through sophisticated technology (McFerran et al., 2019). Yet, the majority of individuals who perceive tinnitus do not suffer from it (Davis & Refaie, 2000; Gallus et al., 2015; Kim et al., 2015). Therefore, the elimination of the tinnitus percept solely, may not directly translate to diminished suffering. Interestingly, reduction of tinnitus-related fear may be a path to diminished tinnitus perception. In the field of chronic pain, reduction in pain perception has been observed after diminished pain-related fear (de Jong et al., 2012). Growing literature on CBT for tinnitus is robust and consistent in diminishing tinnitus suffering (Fuller et al., 2020). This dissertation has highlighted the role of cognitions, behaviour and emotional reactions in Chronic Disabling Tinnitus. Current findings point towards a better understanding through the role of fear in its development (Chapter 2) and recovery (Chapter 5).

### *Hair of the dog*

It has been long believed that “like cures like” (Latin: *similia similibus curantur*). Development of vaccines, for example, lie within a concept not too far from the one in the times of Hippocrates. Exposure techniques may be used in CBT protocols, and resonate strongly with this line of thought by using the feared stimuli as treatment to the fear responses. Following the FA model, the alternative to the pathological cycle (i.e. perceiving tinnitus as a threat to health and functioning) is the recovery path (i.e. perceiving tinnitus as a harmless experience). On the path to recovery, the continuation of valued activities follows, and with it, confronting stimuli that would otherwise be avoided due to the fear of increasing in tinnitus (e.g. silent or loud environments, or avoiding it by distraction). In other words, fear can be reduced by confronting the patient with the fearful experience (e.g. silent environment evoking tinnitus experience without the

option of distraction) without the expected negative outcome (e.g. increased tinnitus). However, since exposure takes the form of confrontation with the fear-provoking stimuli, and *fear* is the catalyst of the pathology, confronting patients with the fearful stimuli can be a challenge (as highlighted in Chapter 5). The violation of the expected threatening prediction, which allows for the creation of a new and more adaptive association with tinnitus. This process however can unintentionally be interrupted and avoided (for a detailed review on underlying mechanism of inhibitory learning and exposure see Craske et al., 2014, 2018). Adding to this, despite support for exposure techniques and the strong recommendation for CBT (Cima et al., 2019), competing techniques and therapies are often recommended instead (e.g. masking, sound enrichment therapy, TRT). These require less therapeutic effort, less initial patient discomfort by confrontation with fear and less agency from the patient, with a higher perceived sense of comfort. Similar to some CBT protocols, TRT utilizes psycho-education (e.g. deconstructing tinnitus threat misconceptions), yet, contrary to CBT-Exposure, avoidance of the tinnitus experience is favoured through masking and sound enrichment therapy (Jastreboff & Hazell, 2004).

While exposure techniques have gathered support beyond the field of tinnitus, and especially in the field of chronic pain (e.g. Craske et al., 2018), findings on the detrimental effect of the utilization of safety-seeking behaviours during exposure (e.g. avoidance) are inconclusive (Meulders et al., 2016). Fundamental research in the field of tinnitus, which would provide a way into examining such effects, are lacking in part due to the absence of replicable human paradigms. Recent review findings, slightly, favour CBT over TRT (i.e. Fuller et al., 2020), however, the inconsistency, variability and availability of protocols utilized limit the interpretability of findings.

### *From whole to unit*

*One for all, all for one – Dumas (1844)*

Studies with broad inclusion criteria fail to acknowledge the idiosyncratic characteristics of tinnitus. The high inter- and intra-individual variability may not be properly acknowledged when interpreting results from group-based analysis. Strong individual effects are lost in the search for a one-size-fits-all solution. Given such limitations, there is a push for individualized medicine (Schork, 2015), warranting tailored designs for treatment and research. Following the call, half of the four studies here presented employed SCED to achieve meaningful results that would otherwise be impossible (Chapters

3 and 5). Unlike observational/non-experimental designs (e.g. case studies), SCED investigates causal relationships between variables through repeated measurements of a dependent variable over at least two different conditions of a manipulated variable (e.g. treatment onset; Morley, 2018). Practically speaking, SCED usually works by repeatedly assessing individuals before (baseline phase: A) and after the random introduction of an intervention or manipulation (phase: B). Effects are mainly examined through differences between phases in level (e.g. mean, median, broadened median), trend (e.g. Least Square Regression) and variability (e.g. range lines, trended range; Kratochwill et al., 2010). Robust statistical methods have been developed to further support the power of analyses on results from SCED (e.g. Randomization Test, NAP, Tau-U). Participants (or units) may be added to further increase power through multiple (i.e. simultaneous start) or sequential (i.e. different start) baselines. The growing use of SCED is reflected in the emergence of published guidelines, such as the Risk of Bias in N-of-1 Trials (RoBiNT; Tate et al., 2013) and The Single-Case Reporting Guideline In BEhavioural Interventions (SCRIBE; Tate et al., 2016).

As we have seen in Chapter 5, SCED may prove an adequate and powerful solution to further study the individual components of CBT for tinnitus. Beyond the capacity to isolate components (e.g. exposure, relaxation, psychoeducation), SCED allows for an intimate exploration of potential pivoting moments in treatment phases within the participant (e.g. identifying triggers of cascading events, emotions, and reactions), as well as a better understanding of current treatment options (e.g. pinpointing when change happens and potential sequence of events/treatment components) while potentially creating new research questions. Perhaps, more importantly, is the capacity for SCED to robustly investigate treatment effectiveness in small samples and single individuals. Due to the wide variety of tinnitus aetiology, longevity and experiences, SCED rises as a promising method due to the capacity to isolate and investigate identified cohorts of tinnitus patients, however small, and the possibility to perform meta-analyses over aggregated data of multiple studies with SCED. These analyses may strengthen external validity of treatment components, without losing individual case results. Creative and interesting investigations using SCEDs are continuously emerging, from analysing patterns of change within a chronic lower back pain intervention (Caneiro et al., 2019) to the efficacy of supervision in increasing CBT competencies (Alfonsson et al., 2020). Future research within tinnitus may integrate the promising research from similar fields (e.g. Caneiro et al., 2019), as well as deepen already existing findings within the field,

such as the benefits of Tinnitus Retraining Therapy (TRT; Jastreboff & Hazell, 2004) or transcranial direct current stimulation (tDCS; Jacquemin et al., 2018). Thus, SCED brings a new perspective within the field that may lead to new knowledge and the re-assessment of existing theories, while also allowing for flexible study designs without compromising the quality of findings.

Naturally, some limitations are noticeable. SCED is not adequate when time periods for phases (i.e. baseline, treatment, follow-up) are not appropriate (Vlaeyen et al., 2020), such as acute care (e.g. emergency care) or longer follow-up periods (e.g. 1 year). More importantly, the generalizability of findings provided through SCED is often questioned. While group-designs seek to increase the generalizability of findings through a larger number of participants, statistical analysis of group averages (means) may not directly translate to treatment effects at the individual level (Kazdin, 2018) and may not even translate to the individuals within that group. Such may also be due to the less idiosyncratic measures utilized in group studies when compared to SCED. By placing the patient in the center of research and treatment, as it is done in SCED, generalizability is created through the replication of treatment protocols with idiosyncratic outcome measures. In other words, it is not the use of standardized questionnaire scores which defines the outcome of an intervention, but the robust individual differences between phases of each unit studied. Replicability, a long standing issue in social sciences (e.g. “Estimating the Reproducibility of Psychological Science,” 2015; Lilienfeld, 2017), is as always, the key to the future. SCED allows for health care providers, and not only researchers, to produce robust findings which in itself may be replicable. In SCED, the feasibility of small-scale research that is rich in information may contribute to a larger set of aggregate data which can provide important insights needed into future research and treatment.

## From tinnitus to trees

“The objects of sense exist only when they are perceived; the trees therefore are in the garden... no longer than while there is somebody by to perceive them.” – Berkeley (1710)

The original thought of Berkeley has inspired the question posed by The Chautauquan which opens this PhD dissertation. The corollary: “*If a human being were to hear a tree fall, would there be any tree?*” was presented and the original thought by Berkeley may provide insight into the answer. The philosopher insisted that existence requires perception, creating the branch of idealism where *to be is to be perceived* (Latin: *esse is percipi*). While avoiding the philosophical debate that endured centuries and which is beyond the scope of this thesis, it is still possible to enrich ourselves from such a point of view. Thus, the existence of tinnitus, as the trees for Berkeley, can only exist as long as there is one to perceive it. Applying this knowledge to the postulated corollary, one might only ascertain that if a human being were to hear a tree fall, the *sound would exist*, whilst the tree remains to be perceived.

## References

- Alfonsson, S., Lundgren, T., & Andersson, G. (2020). Clinical supervision in cognitive behavior therapy improves therapists' competence: a single-case experimental pilot study. *Cognitive Behaviour Therapy*, *49*(5), 425–438. <https://doi.org/10.1080/16506073.2020.1737571>
- Baguley, D., Andersson, G., McFerran, D., & McKenna, L. (2013). Psychological Models of Tinnitus. In *Tinnitus: A Multidisciplinary Approach* (pp. 102–109). John Wiley & Sons, Ltd.
- Berkeley, G. (1710). *A Treatise Concerning the Principles of Human Knowledge*.
- Beukes, E. W., Baguley, D. M., Allen, P. M., Manchaiah, V., & Andersson, G. (2017). Audiologist-Guided Internet-Based Cognitive Behavior Therapy for Adults With Tinnitus in the United Kingdom. *Ear and Hearing*, *1*. <https://doi.org/10.1097/AUD.0000000000000505>
- Beukes, E. W., Baguley, D. M., Jacquemin, L., Lourenco, M. P. C. G., Allen, P. M., Onozuka, J., Stockdale, D., Kaldo, V., Andersson, G., & Manchaiah, V. (2020). Changes in Tinnitus Experiences During the COVID-19 Pandemic. *Frontiers in Public Health*, *8*. <https://doi.org/10.3389/fpubh.2020.592878>
- Beukes, E. W., Manchaiah, V., Davies, A. S. A., Allen, P. M., Baguley, D. M., & Andersson, G. (2018). Participants' experiences of an Internet-based cognitive behavioural therapy intervention for tinnitus. *International Journal of Audiology*, *0*(0), 1–8. <https://doi.org/10.1080/14992027.2018.1514538>
- Boddez, Y., Baeyens, F., Luyten, L., Vansteenwegen, D., Hermans, D., & Beckers, T. (2013). Rating data are underrated: Validity of US expectancy in human fear conditioning. *Journal of Behavior Therapy and Experimental Psychiatry*, *44*(2), 201–206. <https://doi.org/10.1016/j.jbtep.2012.08.003>
- Caneiro, J. P., Smith, A., Linton, S. J., Moseley, G. L., & O'Sullivan, P. (2019). 'How does change unfold?' an evaluation of the process of change in four people with chronic low back pain and high pain-related fear managed with Cognitive Functional Therapy: A replicated single-case experimental design study. *Behaviour Research and Therapy*, *xxxx*. <https://doi.org/10.1016/j.brat.2019.02.007>
- Cima, R. F. F., Crombez, G., & Vlaeyen, J. W. S. (2011). Catastrophizing and Fear of Tinnitus Predict Quality of Life in Patients With Chronic Tinnitus. *Ear and Hearing*, *32*(5), 634–641. <https://doi.org/10.1097/AUD.0b013e31821106dd>
- Cima, R. F. F., Maes, I. H., Joore, M. A., Scheyen, D. J. W. M., El Refaie, A., Baguley, D. M., Anteunis, L. J. C., van Breukelen, G. J. P., & Vlaeyen, J. W. S. (2012). Specialised treatment based on cognitive behaviour therapy versus usual care for tinnitus: a randomised controlled trial. *The Lancet*, *379*(9830), 1951–1959. [https://doi.org/10.1016/S0140-6736\(12\)60469-3](https://doi.org/10.1016/S0140-6736(12)60469-3)
- Cima, R. F. F., Mazurek, B., Haider, H., Kikidis, D., Lapira, A., Noreña, A., & Hoare, D. J. (2019). A multidisciplinary European guideline for tinnitus: diagnostics, assessment, and treatment. *HNO*, *February 2018*. <https://doi.org/10.1007/s00106-019-0633-7>
- Craske, M. G., Hermans, D., & Vervliet, B. (2018). State-of-the-art and future directions for extinction as a translational model for fear and anxiety. *Philosophical Transactions of the Royal Society B: Biological Sciences*, *373*(1742), 20170025. <https://doi.org/10.1098/rstb.2017.0025>
- Craske, M. G., Treanor, M., Conway, C. C., Zbozinek, T., & Vervliet, B. (2014). Maximizing exposure therapy: An inhibitory learning approach. *Behaviour Research and Therapy*, *58*(310), 10–23. <https://doi.org/10.1016/j.brat.2014.04.006>
- Davis, A., & Refaie, A. E. (2000). Epidemiology of Tinnitus. In R. S. Tyler (Ed.), *Handbook of Tinnitus* (pp. 1–24). Singular Thompson Learning.

- de Jong, J. R., Vlaeyen, J. W. S., van Eijsden, M., Loo, C., & Onghena, P. (2012). Reduction of pain-related fear and increased function and participation in work-related upper extremity pain (WRUEP): Effects of exposure in vivo. *Pain, 153*(10), 2109–2118. <https://doi.org/10.1016/j.pain.2012.07.001>
- Dumas, A. (1844). *The three Musketeers*.
- Estimating the reproducibility of psychological science. (2015). *Science, 349*(6251), aac4716–aac4716. <https://doi.org/10.1126/science.aac4716>
- Fuller, T. E., Cima, R., Langguth, B., Mazurek, B., Vlaeyen, J. W. S., & Hoare, D. J. (2020). Cognitive behavioural therapy for tinnitus. *Cochrane Database of Systematic Reviews, 4*. <https://doi.org/10.1002/14651858.CD012614.pub2>
- Gallus, S., Lugo, A., Garavello, W., Bosetti, C., Santoro, E., Colombo, P., Perin, P., La Vecchia, C., & Langguth, B. (2015). Prevalence and Determinants of Tinnitus in the Italian Adult Population. *Neuroepidemiology, 45*(1), 12–19. <https://doi.org/10.1159/000431376>
- Hall, D. A., Smith, H., Hibbert, A., Colley, V., Haider, H. F., Horobin, A., Londero, A., Mazurek, B., Thacker, B., & Fackrell, K. (2018). The COMiT’ID Study: Developing Core Outcome Domains Sets for Clinical Trials of Sound-, Psychology-, and Pharmacology-Based Interventions for Chronic Subjective Tinnitus in Adults. *Trends in Hearing, 22*, 233121651881438. <https://doi.org/10.1177/2331216518814384>
- Hallam, R. S., Jakes, S. C., & Hinchcliffe, R. (1988). Cognitive variables in tinnitus annoyance. *The British Journal of Clinical Psychology, 27 ( Pt 3)*(pt 3), 213–222. <http://www.ncbi.nlm.nih.gov/pubmed/3191301>
- Henry, J. A., Galvez, G., Turbin, M. B., Thielman, E. J., Mcmillan, G. P., & Istvan, J. A. (2012). *Pilot Study to Evaluate Ecological Momentary Assessment of Tinnitus*. 279–290.
- Hesser, H., Gustafsson, T., Lundén, C., Henrikson, O., Fattahi, K., Johnsson, E., Westin, V. Z., Carlbring, P., Mäki-Torkko, E., Kaldo, V., & Andersson, G. (2012). A randomized controlled trial of internet-delivered cognitive behavior therapy and acceptance and commitment therapy in the treatment of tinnitus. *Journal of Consulting and Clinical Psychology, 80*(4), 649–661. <https://doi.org/10.1037/a0027021>
- Jacquemin, L., Shekhawat, G. S., Van de Heyning, P., Mertens, G., Fransen, E., Van Rompaey, V., Topsakal, V., Moyaert, J., Beyers, J., & Gilles, A. (2018). Effects of Electrical Stimulation in Tinnitus Patients: Conventional Versus High-Definition tDCS. *Neurorehabilitation and Neural Repair, 32*(8), 714–723. <https://doi.org/10.1177/1545968318787916>
- Jastreboff, P. J. (1990). Phantom auditory perception (tinnitus): mechanisms of generation and perception. *Neuroscience Research, 8*(4), 221–254. [https://doi.org/10.1016/0168-0102\(90\)90031-9](https://doi.org/10.1016/0168-0102(90)90031-9)
- Jastreboff, P. J., Brennan, J. F., Coleman, J. K., & Sasaki, C. T. (1988). Phantom auditory sensation in rats: An animal model for tinnitus. *Behavioral Neuroscience, 102*(6), 811–822. <https://doi.org/10.1037/0735-7044.102.6.811>
- Jastreboff, P. J., & Hazell, J. W. P. (2004). *Tinnitus Retraining Therapy*. Cambridge University Press. <https://doi.org/10.1017/CBO9780511544989>
- Jastreboff, P. J., Hazell, J. W. P., & Graham, R. L. (1994). Neurophysiological model of tinnitus: Dependence of the minimal masking level on treatment outcome. *Hearing Research, 80*(2), 216–232. [https://doi.org/10.1016/0378-5955\(94\)90113-9](https://doi.org/10.1016/0378-5955(94)90113-9)
- Kazdin, A. E. (2018). Single-case experimental designs. Evaluating interventions in research and clinical practice. *Behaviour Research and Therapy, November*, 0–1. <https://doi.org/10.1016/j.brat.2018.11.015>



- Kim, H.-J., Lee, H.-J., An, S.-Y., Sim, S., Park, B., Kim, S. W., Lee, J. S., Hong, S. K., & Choi, H. G. (2015). Analysis of the Prevalence and Associated Risk Factors of Tinnitus in Adults. *Plos One*, *10*(5), e0127578. <https://doi.org/10.1371/journal.pone.0127578>
- Kratochwill, T. R., Hitchcock, J., Horner, R. H., Levin, J. R., Odom, S. L., Rindskopf, D. M., & Shadish, W. R. (2010). *Single-case designs technical documentation*. Retrieved from What Works Clearinghouse website: [http://ies.ed.gov/ncee/wwc/pdf/wwc\\_scd.pdf](http://ies.ed.gov/ncee/wwc/pdf/wwc_scd.pdf).
- Lilienfeld, S. O. (2017). Psychology's Replication Crisis and the Grant Culture: Righting the Ship. *Perspectives on Psychological Science*, *12*(4), 660–664. <https://doi.org/10.1177/1745691616687745>
- McFerran, D. J., Stockdale, D., Holme, R., Large, C. H., & Baguley, D. M. (2019). Why Is There No Cure for Tinnitus? *Frontiers in Neuroscience*, *13*. <https://doi.org/10.3389/fnins.2019.00802>
- Meikle, M. B., Henry, J. A., Griest, S. E., Stewart, B. J., Abrams, H. B., McArdle, R., Myers, P. J., Newman, C. W., Sandridge, S., Turk, D. C., Folmer, R. L., Frederick, E. J., House, J. W., Jacobson, G. P., Kinney, S. E., Martin, W. H., Nagler, S. M., Reich, G. E., Searchfield, G., ... Vernon, J. A. (2012). The Tinnitus Functional Index. *Ear and Hearing*, *33*(2), 153–176. <https://doi.org/10.1097/AUD.0b013e31822f67c0>
- Meulders, A., Van Daele, T., Volders, S., & Vlaeyen, J. W. S. (2016). The use of safety-seeking behavior in exposure-based treatments for fear and anxiety: Benefit or burden? A meta-analytic review. *Clinical Psychology Review*, *45*, 144–156. <https://doi.org/10.1016/j.cpr.2016.02.002>
- Miron-Shatz, T., Stone, A., & Kahneman, D. (2009). Memories of Yesterday's Emotions: Does the Valence of Experience Affect the Memory-Experience Gap? *Emotion*, *9*(6), 885–891. <https://doi.org/10.1037/a0017823>
- Morley, S. (2018). *Single-Case Methods in Clinical Psychology: A Practical Guide* (M. Ciara & C. J. Main (Eds.); Kindle Edi). Taylor & Francis.
- Schlee, W., Pryss, R. C., Probst, T., Schobel, J., Bachmeier, A., Reichert, M., & Langguth, B. (2016). Measuring the moment-to-moment variability of Tinnitus: The TrackYourTinnitus smart phone app. *Frontiers in Aging Neuroscience*, *8*(DEC), 1–8. <https://doi.org/10.3389/fnagi.2016.00294>
- Schork, N. J. (2015). Personalized medicine: Time for one-person trials. *Nature*, *520*(7549), 609–611. <https://doi.org/10.1038/520609a>
- Shiffman, S., Stone, A. A., & Hufford, M. R. (2008). Ecological Momentary Assessment. *Annual Review of Clinical Psychology*, *4*(1), 1–32. <https://doi.org/10.1146/annurev.clinpsy.3.022806.091415>
- Tate, R. L., Perdices, M., Rosenkoetter, U., McDonald, S., Togher, L., Shadish, W., Horner, R., Kratochwill, T., Barlow, D. H., Kazdin, A., Sampson, M., Shamseer, L., & Vohra, S. (2016). The Single-Case Reporting Guideline In BEhavioural Interventions (SCRIBE) 2016 Statement. *Archives of Scientific Psychology*, *4*(1), 10–31. <https://doi.org/10.1037/arc0000027>
- Tate, R. L., Perdices, M., Rosenkoetter, U., Wakim, D., Godbee, K., Togher, L., & McDonald, S. (2013). Revision of a method quality rating scale for single-case experimental designs and n-of-1 trials: The 15-item Risk of Bias in N-of-1 Trials (RoBiNT) Scale. *Neuropsychological Rehabilitation*, *23*(5), 619–638. <https://doi.org/10.1080/09602011.2013.824383>
- Vlaeyen, J. W. S., Wicksell, R. K., Simons, L. E., Gentili, C., De, T. K., Tate, R. L., Vohra, S., Punja, S., Linton, S. J., Snichotta, F. F., & Onghena, P. (2020). *From Boulder to Stockholm in 70 Years : Single Case Experimental Designs in Clinical Research*.



# CHAPTER 7

---

Impact statement

## Impact statement

The current body of work sought to develop three different, but related, areas in the tinnitus field. First, we focused on new tools for assessing the tinnitus more accurately. Second, we investigated tinnitus treatment, more specifically, Cognitive Behavioural Therapy (CBT) for tinnitus and whether a group delivery would be better than individually delivered CBT. Lastly, we tackled the development of research into the role of fear in tinnitus.

The investigation of new tools for assessing tinnitus was deemed necessary due to the limitations of assessing tinnitus. Due to the lack of a “definitive” measure of tinnitus, such as the results of blood sample analysis for certain conditions (e.g. hepatitis, HIV, etc), tinnitus relies solely on self-report measures. Traditional self-report measures (i.e. questionnaires) have limitations that can strongly influence results. One such limitation is that these questionnaires are subjected to biases. In other words, participants are influenced by their memory (e.g. remembering only negative experiences), mood (happy vs foul), and location (hospital vs home) when filling out these questionnaires. Newer methods that are delivered through smartphone apps allow for assessing tinnitus in daily life, where it matters most, through small questionnaires and during longer periods of time (e.g. months). Although the possible negative side-effects of such strategies (e.g. extended screen time, excessive smartphone use) were yet to be investigated within the tinnitus field.

Tinnitus treatment delivery (group vs individual) has not been directly tested within the tinnitus field before. Previous literature has demonstrated the power of group treatment, specifically the use of learning through others (i.e. observational learning). Whether if group treatment has added benefits to the participants provides guidance for future treatment design and a better understanding of the tinnitus recovery process.

It is postulated that the role of fear in the development and maintenance of tinnitus is of vital importance. Inspired in research of other fields (i.e. chronic pain), fear of tinnitus is believed to trigger a detrimental cycle where misinterpretations of the tinnitus experience (e.g. fear that tinnitus is a symptom of a tumour; i.e. catastrophic thoughts of the meaning of tinnitus), excessive and constant tinnitus monitoring, and maladaptive behavioural responses (e.g. avoiding silence or social situations; i.e. avoidance of non-harmful activities) lead to a significant decrease in quality of life. Some research on fear and tinnitus has been conducted, though research is still in its infancy.

Our findings were promising. New tools to measure the tinnitus experience were proven safe for use with those suffering from severe tinnitus. Also, given the personal or financial burden of some of those tools for the researcher (i.e. Ecological Momentary Assessment, EMA), a different method (i.e. end-of-day diary, EDD) has proven an adequate alternative. Regarding tinnitus treatment, group CBT was observed to be more beneficial when compared to individual treatment. Finally, the challenge to investigate the role of fear in tinnitus was met with a novel experimental paradigm, which provides insights into learning fear of tinnitus, and by extension the development of chronic tinnitus.

Scientific advances from these findings are considerable. We have paved the way for the safe use of new tools and methodologies within the tinnitus field. The use of EMA and EDD provides a different perspective to the understanding of the tinnitus experience that is closer to the *lived experience* of those who suffer from it. These methods also make it possible for researcher and healthcare providers to pinpoint pivoting moments in the patient's illness trajectory (e.g. trigger of emotional and behavioural cascades). This approach is congruent with personalized medicine, where treatments are tailored to the individual in question. Benefiting from the individualized treatment was the finding that group CBT provides better care when compared to individual CBT. This finding is particularly relevant for at least two reasons: (1) *tailored* treatment does not necessarily mean *individual* treatment, and as such, group treatment may be part of a tailored treatment path; and (2) healthcare resources may be optimized in order to provide the best care under the least cost.

Within the field of tinnitus, fundamental research is mainly conducted with animal samples. Animal models provide insight into the possible workings within a physical model, it cannot account for the disability and distress that tinnitus has on human beings. Perhaps more difficult to directly translate into direct patient outcome is the development of experimental research into the role of fear in tinnitus. While investigation in treatment outcomes are of paramount importance, the understanding of the underlying cognitive and behavioural mechanisms of change provide the pillars upon which such treatments stand. The understanding of the development of chronic from acute tinnitus, as well as tinnitus distress and disability all hinge in the advance of theoretical models. The contribution on this front may be the most substantial yet. While not without its limitations, our research may improve tinnitus research in human participants.

Those who suffer from tinnitus directly benefit from the research presented. All studies conducted may directly influence the treatment approaches to tinnitus patients. Beyond tinnitus, the findings of this body of work may be adapted and inspire similar fields. The growing incidence of chronic conditions may yet benefit from models adapted from parallel fields (as tinnitus has from chronic pain research). A culmination of a body of work from different fields may create a broader framework that could potentially help better understand similar disorders under a similar context. Currently and perhaps more importantly, the financial burden of such conditions (as tinnitus) to the healthcare system and the personal burden on the patients and their families may be reduced through the development of research and more efficacious treatments.

With that in mind, all the findings are (or will be) available to the public. Initiatives by the funding agencies and the author are taken to spread the knowledge produced. It is our hope to inspire future tinnitus research and push for newer ways of research which puts the individual in the center of care.







## CHAPTER 8

---

### Acknowledgments

*No man is an Iland, intire of itselſe; every man is a peece of the Continent, a part of the maine; if a Clod bee washed away by the Sea, Europe is the leſſe, as well as if a Promontorie were, as well as if a Manor of thy friends or of thine owne were; any mans death diminishes me, becauſe I am involved in Mankinde; And therefore never ſend to know for whom the bell tolls; It tolls for thee.*

—John Donne, Meditation XVII

This thesis could not have been achieved if it were not by the combined efforts of many individuals who directly and indirectly contributed to the product here presented.

First, I want to mention those who made this PhD journey possible. To Prof. Johan Vlaeyen, and Dr. Rilana Cima, thank you for the trust deposited in me to develop and deliver this thesis. In particular, thank you Johan for the crucial interventions, guidance and feedback during the process. To Rilana, thank you for the support, guidance and freedom to explore, pursue and test my own curiosity.

Moreover, this project would not exist if it were not for Prof. Winfried Schlee. Winny's enthusiasm for mentoring has touched all students of the ESIT project. For this, and for accepting the invitation to be part of the jury, I thank you.

To prof. Derek Hoare and prof. Ilse Van Diest, members of the jury, thank you for making the time to read this thesis, as well as being present here today for the defence. Prof. Derek Hoare, I also had the pleasure to travel with you and an extra thank you for providing me the necessary guidance in moments of need. Prof. Ilse van Diest, your knowledge and expertise are inspiring, and it is an honour to have you as part of such crucial moment in my career.

For the guidance, intervention and being the chair in the thesis' defense, I thank Prof. Madelon Peters. Your career, and vineyard, serve as an example to follow and hope to one day achieve.

I cannot continue my acknowledgements without mentioning where it all began: my family. My parents certainly did not imagine that one day I would defend a doctoral thesis in the Netherlands. They could not have suspected that their curiosity, integrity, and dedication would have been marked so deeply in their children. Nor could have they imagined how far it would have taken us. This thesis is a celebration of all the sacrifices

we went through as a family searching for a better life. Sacrifices that included living in completely different continents at times. But no matter how wide the oceans that would separate us, the distance could not hold a candle in light of the love my parents have for us. Portanto, aos meus pais, a minha irmã e sua nova família (Francesco, Teresa e Anna Rosa), e a Santa Teresinha, agradeço hoje, e sempre, pelo amor incondicional.

The importance of my family in my accomplishments cannot be understated and words will never be enough. I also want to acknowledge all of those who, in these past years, were my surrogate family. All of those, who in my times of need, without the comfort of my own family, made me feel loved. My older colleagues in Portugal, my in-laws in Zemst and my friends from ESIT and Maastricht, the next paragraphs are for you.

Aos meus amigos João Matos, Alexandre Teles, Alexandre Sousa, e Ricardo Simas, me faltam palavras para vos agradecer durante este processo longo e árduo. Apesar da distância, senti sempre o vosso apoio e orgulho. As palavras amigas que, em momentos difíceis, me acalmavam e me motivavam. Agradeço o humor, o apoio e a dedicação para me verem em minhas poucas visitas à Portugal. O futuro promete, e com ele novas aventuras, de Carvoeiro a Porto Covo e além, juntos a lembrar que “*vai dar tudo certo*”.

Merecedor de um parágrafo único, e sem dúvidas mais do que aqui posso escrever, é o chefe Gonçalo Santos. Mais que um amigo, é um irmão que me apoiou e continua apoiando em cada momento. A cada passo que dou ou “*por cada imponderável que surja*” posso contar com sua perspectiva realista e argumentos irrefutáveis, seja às 3:00 ou às 15:00. Não há momento que eu me sinta sozinho, porque sei que basta uma palavra e lá estarás. Obrigado chefe, em nome da minha família, por seu carinho e dedicação.

Aan de Jacquemins, ik wil jullie bedanken om mij te verwelkomen in jullie familie, de smakelijke etentjes op zondag, de strandvakanties, de hulp bij het verhuizen en zo veel meer. Er was geen enkel moment dat ik me niet thuis voelde op de “Jackie-berg”. Dus duizendmaal bedankt van mij en mijn familie.

To my ESIT colleagues, thank you for travelling the world with me. We, through the ESIT project, managed to learn, work, and even publish together. A special thanks to Axel and Constanze, thank you for the many talks and the wild nights.

To my friends who made a town in the south of the Netherlands feel like the most unusual home. A home that, because of you, was so much harder to leave when the time came. Juliane and Eveliina, thank you for walking this path with me, going over every struggle

and victory together. Thomas and Kai, thank you for forging the path, the moments of guidance and support in the past four years. For the laughs, coffees, and hangouts, I thank Krissy and Jojo. To Iris, thank you for being my therapist and running a marathon with me, in other words, I am happy that you could practice your therapeutic skills and that we could suffer together for 5 hours under the rain in Brussels. To Sieske who can always lift the mood (though in a whirlwind), Ola for proving that it could be worse, Mathilde for the ethical discussions, Astrid for the drinks, and to Andreas B. for letting me know I am not crazy, thank you. But more importantly, thank you all for the long lunches, inspiring (and not so inspiring) talks, quickly translating questionnaires and emails, travels, conferences, dinners, drinks, shopping and making Maastricht a home.

To Laure, from seeing you the first time in Regensburg, to finally meeting you in Nottingham. From reading your articles to becoming your friend. From lighting a candle in the Duomo to buying our first place together. From watching you defend your thesis to seeing you watching me do the same. I could not have imagined the role that you would come to play in achieving this goal and, more importantly, the fundamental change that you brought into my life. For all your help, from structuring my task list with post-its on the walls, to opening your family to me, thank you. There is nothing I can write here that I have not said to you before, and there is much that I will reserve for my wedding vows and the years to come. Ik hou van jou.

It is because of all of you that I can dedicate this thesis to my great-aunt, Geni. A strong woman with an unforgettable personality that despite leaving us over a decade ago, lives in my heart with memories that I will never forget. For helping me reach this conclusion and allowing me to dedicate this work to her, I will always be grateful.

