

Assessment of extracranial and intracranial atherosclerosis

Citation for published version (APA):

Jansen, J. F. A. (2018). Assessment of extracranial and intracranial atherosclerosis: Don't dismiss old school autopsy. *Atherosclerosis*, 270, 189-190. <https://doi.org/10.1016/j.atherosclerosis.2018.01.002>

Document status and date:

Published: 01/03/2018

DOI:

[10.1016/j.atherosclerosis.2018.01.002](https://doi.org/10.1016/j.atherosclerosis.2018.01.002)

Document Version:

Publisher's PDF, also known as Version of record

Document license:

Taverne

Please check the document version of this publication:

- A submitted manuscript is the version of the article upon submission and before peer-review. There can be important differences between the submitted version and the official published version of record. People interested in the research are advised to contact the author for the final version of the publication, or visit the DOI to the publisher's website.
- The final author version and the galley proof are versions of the publication after peer review.
- The final published version features the final layout of the paper including the volume, issue and page numbers.

[Link to publication](#)

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal.

If the publication is distributed under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license above, please follow below link for the End User Agreement:

www.umlib.nl/taverne-license

Take down policy

If you believe that this document breaches copyright please contact us at:

repository@maastrichtuniversity.nl

providing details and we will investigate your claim.



Editorial

Assessment of extracranial and intracranial atherosclerosis: Don't dismiss old school autopsy



Keywords:

Atherosclerosis
Carotid arteries
Cerebral arteries
Risk factors
Imaging

Atherosclerotic disease represents one of the leading causes of ischemic stroke worldwide, posing a major global public health concern [1]. The prevalence of patients with intracranial atherosclerotic disease (IAD) exceeds those of patients with extracranial atherosclerotic disease (EAD) [2]. Risk factors for IAD and EAD have been shown to differ, the former has for example been related to obesity and hyperlipidemia whereas the latter has been associated with hypertension, though not all studies concur [3]. Furthermore, ethnic differences play a major role in the location of atherosclerosis [3].

In the era of noninvasive population imaging studies of IAD such as the Atherosclerosis Risk in Communities (ARIC) initiative [4,5], there is still a necessity for old-fashioned gold standard morphometric autopsy studies. Although recent advantages in noninvasive imaging have resulted in the possibility of IAD assessment using magnetic resonance imaging and ultrasound techniques [6], autopsy studies are still preferred for accurate assessment and visualization of atherosclerotic plaques. For example, some anatomical locations, including the distal arterial branches, have proven to be a major challenge for noninvasive imaging methods, whereas they pose no challenge for autopsy morphometric evaluation. In this issue of *Atherosclerosis*, Suemoto et al. describe a large population (661 community-dwelling elderly) based autopsy study in which they examine atherosclerotic disease in the form of stenosis in the common (CCA) and internal carotid arteries (ICA), and relate their findings to several risk factors in a population with a diverse ethnic origin [7]. Interestingly, they also investigated the association of IAD with EAD.

Suemoto et al. confirmed that in their Brazilian population, IAD was indeed slightly more common than EAD (59% and 51%,

respectively). Due to the unique design of their autopsy study, subjects were included when they died from natural or unknown causes, hence the sampling was not biased towards cardiovascular disease (including stroke). Therefore, the reported prevalence and associations with risk factors are better generalizable than other imaging and autopsy studies that tend to focus on cardiovascular disease [8,9]. Importantly, it was shown that EAD was associated with more than twice the odds of having IAD. Furthermore, risk factors associated with EAD included Caucasian race, hypertension, and smoking, whereas risk factors for IAD included older age, less years of education, hypertension, diabetes, and a previous history of stroke. Due to the considerable sample size, these differences in atherosclerotic disease risk factors are not likely the result of differences in statistical power, but they rather constitute different underlying causal mechanisms. Moreover, these differences may provide valuable insights into preventive risk-factor management strategies. Future population studies might include both *in vivo* noninvasive imaging and *postmortem* autopsy morphometric assessments, although the logistics of these types of studies might provide some interesting challenges.

Conflict of interest

The author declared he does not have anything to disclose regarding conflict of interest with respect to this manuscript.

References

- [1] L.K. Wong, Global burden of intracranial atherosclerosis, *Int. J. Stroke* 1 (2006) 158–159.
- [2] C. Lei, B. Wu, M. Liu, et al., Risk factors and clinical outcomes associated with intracranial and extracranial atherosclerotic stenosis acute ischemic stroke, *J. Stroke Cerebrovasc. Dis.* 23 (2014) 1112–1117.
- [3] X. Ding, C. Li, K. Yu, et al., Different risk factors between intracranial and extracranial atherosclerotic stenosis in Asian population: a systematic review and meta-analysis, *Int. J. Neurosci.* 124 (2014) 834–840.
- [4] Y. Qiao, E. Guallar, F.K. Suri, et al., MR imaging measures of intracranial atherosclerosis in a population-based study, *Radiology* 280 (3) (2016) 860–868.
- [5] Y. Qiao, F.K. Suri, Y. Zhang, et al., Racial differences in prevalence and risk for intracranial atherosclerosis in a US community-based population, *JAMA Cardiol.* 2 (12) (2017) 1341–1348.
- [6] J.F. Arenillas, Intracranial atherosclerosis: current concepts, *Stroke* 42 (2011) S20–S23.
- [7] C.K. Suemoto, L.P. Grinberg, R.E. Leite, et al., Morphometric measurements of extracranial and intracranial atherosclerotic disease: a population-based

- autopsy study, *Atherosclerosis* 270 (2018) 218–223.
- [8] M. Mazighi, J. Labreuche, F. Gongora-Rivera, et al., Autopsy prevalence of intracranial atherosclerosis in patients with fatal stroke, *Stroke* 39 (2008) 1142–1147.
- [9] O. Alkan, O. Kizilkilic, T. Yildirim, et al., Intracranial cerebral artery stenosis with associated coronary artery and extracranial carotid artery stenosis in Turkish patients, *Eur. J. Radiol.* 71 (2009) 450–455.

Jacobus F.A. Jansen
*Department of Radiology, Maastricht University Medical Centre,
Maastricht, The Netherlands*
E-mail address: jacobus.jansen@mumc.nl.