

Classification algorithm for the International Classification of Diseases-11 chronic pain classification

Citation for published version (APA):

Korwisi, B., Hay, G., Attal, N., Aziz, Q., Bennett, M. I., Benoliel, R., Cohen, M., Evers, S., Giamberardino, M. A., Kaasa, S., Kosek, E., Lavand'homme, P., Nicholas, M., Perrot, S., Schug, S., Smith, B. H., Svensson, P., Vlaeyen, J. W. S., Wang, S. J., ... Barke, A. (2021). Classification algorithm for the International Classification of Diseases-11 chronic pain classification: development and results from a preliminary pilot evaluation. *Pain*, 162(7), 2087-2096. <https://doi.org/10.1097/j.pain.0000000000002208>

Document status and date:

Published: 01/07/2021

DOI:

[10.1097/j.pain.0000000000002208](https://doi.org/10.1097/j.pain.0000000000002208)

Document Version:

Publisher's PDF, also known as Version of record

Document license:

Taverne

Please check the document version of this publication:

- A submitted manuscript is the version of the article upon submission and before peer-review. There can be important differences between the submitted version and the official published version of record. People interested in the research are advised to contact the author for the final version of the publication, or visit the DOI to the publisher's website.
- The final author version and the galley proof are versions of the publication after peer review.
- The final published version features the final layout of the paper including the volume, issue and page numbers.

[Link to publication](#)

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal.

If the publication is distributed under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license above, please follow below link for the End User Agreement:

www.umlib.nl/taverne-license

Take down policy

If you believe that this document breaches copyright please contact us at:

repository@maastrichtuniversity.nl

providing details and we will investigate your claim.

Classification algorithm for the *International Classification of Diseases-11* chronic pain classification: development and results from a preliminary pilot evaluation

Beatrice Korwisi^{a,*}, Ginea Hay^a, Nadine Attal^b, Qasim Aziz^c, Michael I. Bennett^d, Rafael Benoliel^e, Milton Cohen^f, Stefan Evers^{g,h}, Maria Adele Giamberardino^{i,bb}, Stein Kaasa^{j,k,l}, Eva Kosek^{m,n}, Patricia Lavand'homme^o, Michael Nicholas^p, Serge Perrot^q, Stephan Schug^r, Blair H. Smith^s, Peter Svensson^t, Johan W.S. Vlaeyen^{u,v,w}, Shuu-Jiun Wang^{x,y}, Rolf-Detlef Treede^z, Winfried Rief^a, Antonia Barke^{aa}

Abstract

The *International Classification of Diseases-11 (ICD-11)* chronic pain classification includes about 100 chronic pain diagnoses on different diagnostic levels. Each of these diagnoses requires specific operationalized diagnostic criteria to be present. The classification comprises more than 200 diagnostic criteria. The aim of the Classification Algorithm for Chronic Pain in *ICD-11* (CAL-CP) is to facilitate the use of the classification by guiding users through these diagnostic criteria. The diagnostic criteria were ordered hierarchically and visualized in accordance with the standards defined by the Society for Medical Decision Making Committee on Standardization of Clinical Algorithms. The resulting linear decision tree underwent several rounds of iterative checks and feedback by its developers, as well as other pain experts. A preliminary pilot evaluation was conducted in the context of an ecological implementation field study of the classification itself. The resulting algorithm consists of a linear decision tree, an introduction form, and an appendix. The initial decision trunk can be used as a standalone algorithm in primary care. Each diagnostic criterion is represented in a decision box. The user needs to decide for each criterion whether it is present or not, and then follow the respective yes or no arrows to arrive at the corresponding *ICD-11* diagnosis. The results of the pilot evaluation showed good clinical utility of the algorithm. The CAL-CP can contribute to reliable diagnoses by structuring a way through the classification and by increasing adherence to the criteria. Future studies need to evaluate its utility further and analyze its impact on the accuracy of the assigned diagnoses.

Keywords: *ICD-11*, Chronic pain, Pain classification, Classification algorithm, Decision trees

1. Introduction

The latest revision of the *International Classification of Diseases (ICD-11)*, published by the World Health Organization (WHO), will come into effect in January 2022.⁴¹ The *ICD-11* includes a comprehensive new classification of chronic pain, developed by a taskforce of the International Association for the Study of Pain (IASP).³⁷ This classification defines chronic pain as pain that persists or recurs for more than 3 months.^{34,35} It is divided into 7 main categories, or level 1 diagnoses, all of which have been described in detail elsewhere: MG30.0 Chronic primary pain,²¹ MG30.1 Chronic cancer-related pain,² MG30.2 Chronic post-surgical or post traumatic pain,³⁰ MG30.3 Chronic secondary musculoskeletal pain,²⁴ MG30.4 Chronic secondary visceral pain,¹ MG30.5 Chronic neuropathic pain,²⁹ and MG30.6 Chronic secondary headache or orofacial pain.³

Each of these main categories has several subcategories, or child categories, on 3 or 4 diagnostic levels to provide more detailed diagnoses, and to represent the majority of chronic pain diagnoses with specific diagnostic codes. See Box 1 for details on the different diagnostic levels ("parent" and "child" diagnoses in WHO terminology). Further details can be found in the *ICD-11* Reference Guide³⁹ and the *ICD-11* User Guide⁴⁰ provided by the WHO.

In total, the *ICD-11* includes about 100 different chronic pain entities on levels 1 to 4.³⁸ For each of these diagnoses, specific

operationalized diagnostic criteria must be fulfilled. That is to say, a given diagnosis can only be assigned if all specified criteria are present in a given patient (eg, significant emotional distress or functional disability must be present to fulfil the criteria for MG30.0 Chronic primary pain). Existing criteria, such as the criteria of the third edition of the *International Classification of Headache Disorders (ICHD-3)*¹⁰ and the Rome IV criteria for functional gastrointestinal disorders,⁷ have been integrated in the *ICD-11* chronic pain classification. In total, the *ICD-11* chronic pain classification relies on more than 200 different diagnostic criteria. On average, each diagnosis is based on 4 to 7 diagnostic criteria. Child categories inherit all criteria from their parent category. For example, the above-mentioned criterion of emotional distress or functional disability applies to all level 1 to 4 diagnoses of chronic primary pain. Furthermore, more specific criteria are specified on the lower diagnostic levels only.

To guide users of the *ICD-11* chronic pain classification through the multitude of criteria, we developed a classification algorithm that provides a standardized way through the criteria leading to the specific diagnoses. The algorithm aims at structuring and facilitating the classification process, as the use of algorithms generally improves the reliability of the diagnoses.^{17,28} Within a large international field study to evaluate the *ICD-11* chronic pain classification,¹⁶ clinicians provided preliminary feedback on the algorithm.

2. Methods

2.1. Development of the classification algorithm for the International Classification of Diseases-11 chronic pain classification

The Classification ALgorithm for Chronic Pain in ICD-11 (CAL-CP) was developed and structured according to the guidelines formulated by the Society for Medical Decision Making Committee on Standardization of Clinical Algorithms.³¹ The guideline regulates the exact graphical elements and their functions. It recognizes arrows and different kinds of boxes: “clinical state boxes” (rounded rectangles), “decision boxes” (hexagons), and “action boxes” (rectangles). The boxes are linked by the arrows that flow from top to bottom and from left to right. Each arrow visualizes a given “yes” or “no” decision. All boxes are numbered consecutively with the consecutive numbering of all boxes following the flow of the arrows. Annotations on a given box may be added, for example, to add more details or clarifications to a decision box.³¹

In a first step of the development of the CAL-CP, all diagnostic criteria of the ICD-11 chronic pain classification, except for chronic headache or orofacial pain, were brought into a hierarchical order for each of the 6 main categories. The reason for the exclusion of chronic headache or orofacial pain in the present algorithm is that with the ICHD-3¹⁰ and the International Classification of Orofacial Pain (ICOP),¹¹ extensive classifications for primary and secondary headache and orofacial pain already exist. Although ICHD-3 and ICOP definitions and diagnostic criteria for the more general diagnostic levels have been harmonized with the ICD-11,^{3,21} the CAL-CP avoided interference with these existing classification efforts. Furthermore, algorithms for some of the ICHD-3 headaches are available.²⁷

Each criterion was represented in a corresponding decision box. In the next step, arrows were added to link the boxes to form a linear decision tree: each box is linked to the next box by a yes and a no arrow to represent the decision taken. Diagnoses were added in clinical state boxes. They represent the ICD-11 diagnoses reached and extend to different levels. Diagnoses to which the concept of double parenting applies (Box 1) were

added to the branch of the primary parent category. An action box within the branch of the second parent category links to the branch of the primary parent. For example, the level 3 diagnosis “Chronic central neuropathic pain associated with spinal cord injury” (ID 869493945) is a child category of MG30.50 Chronic central neuropathic pain as well as MG30.20 Chronic post traumatic pain. Therefore, this diagnosis is not only implemented in the branch for chronic neuropathic pain, but an action box within the branch of chronic post traumatic pain also links to this entry.

The resulting linear decision tree was subject to several rounds of consecutive checks and feedback. The members of the IASP taskforce who developed the classification, and who were involved in the development of the algorithm, reviewed whether the diagnostic boxes and the arrows connecting the boxes were correct. They also provided feedback on ambiguous diagnostic decisions. Three independent specialists verified the algorithm’s completeness, that is, that all diagnostic criteria and diagnoses were included. Furthermore, they checked whether all arrows were correct and whether diagnoses to which the new ICD-11 concept of “double parenting” applies were represented within all applicable branches.

The algorithm underwent 2 rounds of external review and feedback by 2 pain specialists who were not involved in its development, and who had only minimal prior knowledge regarding the ICD-11 chronic pain classification. The first pain specialist provided feedback on the general concept and the instructions (“Does it make sense?”). After the feedback implementation, the second pain specialist applied the algorithm to a mock patient case before providing detailed feedback on its use as well as any difficulties and problems encountered. The test was whether the pain specialist arrived at the correct diagnoses for the mock patient using the CAL-CP. An observer evaluated difficulties and problems that arose during the mock assessment.

In the course of these iterative rounds of checks and feedback, several additional elements were added to the decision tree: detailed instructions, an introduction form, and an appendix.

After the pilot use of the algorithm in the context of a large ecological implementation field study of the ICD-11 chronic pain

Sponsorships or competing interests that may be relevant to content are disclosed at the end of this article.

R.-D. Treede, W. Rief, and A. Barke contributed equally to this work.

^a Division of Clinical Psychology and Psychotherapy, Department of Psychology, Philipps-University Marburg, Marburg, Germany, ^b INSERM U-987, Centre d’Evaluation et de Traitement de la Douleur, CHU Ambroise Paré, Boulogne-Billancourt, France, ^c Centre for Neuroscience, Surgery and Trauma, Blizard Institute, Wingate Institute of Neurogastroenterology, Barts and The London School of Medicine and Dentistry, Queen Mary, University of London, London, United Kingdom, ^d Academic Unit of Palliative Care, University of Leeds, Leeds, United Kingdom, ^e Department of Diagnostic Sciences, Rutgers School of Dental Medicine, Rutgers, Newark, NJ, United States, ^f St Vincent’s Clinical School, UNSW, Sydney, New South Wales, Australia, ^g Department of Neurology, Krankenhaus Lindenbrunn, Copenbrügge, Germany, ^h Department of Medicine, University of Münster, Münster, Germany, ⁱ Department of Medicine and Science of Aging, CAST, G D’Annunzio University of Chieti, Chieti, Italy, ^j European Palliative Care Research Centre (PRC), Department of Cancer Treatment, University Hospital Oslo, Oslo, Norway, ^k Department of Oncology, Oslo University Hospital, Oslo, Norway, ^l Institute of Cancer Research and Molecular Medicine, Department of Medicine, Norwegian University of Science and Technology, Trondheim, Norway, ^m Department of Clinical Neuroscience, Karolinska Institute, and Department of Neuroradiology, Karolinska University Hospital, Stockholm, Sweden, ⁿ Department of Surgical Sciences Uppsala University, Uppsala, Sweden, ^o Department of Anaesthesiology, Acute Postoperative Pain Service, Saint Luc Hospital, Catholic University of Louvain, Brussels, Belgium, ^p Pain Management Research Institute, Faculty of Medicine and Health, University of Sydney and Royal North Shore Hospital, Sydney, Australia, ^q Pain Clinic, Cochin Hospital, Paris University, INSERM U987, Paris, France, ^r Department of Anaesthesiology and Pain Medicine, Medical School, University of Western Australia, Perth, Australia, ^s Division of Population Health and Genomics, University of Dundee, Dundee, United Kingdom, ^t Department of Dentistry and Oral Health, Section of Orofacial Pain and Jaw Function, Aarhus University, Aarhus, Denmark, ^u Research Group Health Psychology, Department of Psychology and Educational Sciences, University of Leuven, Leuven, Belgium, ^v TRACE, Center for Translational Health Research, KU Leuven, Ziekenhuis Oost-Limburg, Genk, Belgium, ^w Department of Clinical Psychological Science, Maastricht University, Maastricht, the Netherlands, ^x The Neurological Institute, Taipei Veterans General Hospital, Taipei, Taiwan, ^y Brain Research Center and School of Medicine, National Yang-Ming University, Taipei, Taiwan, ^z Department of Neurophysiology, Mannheim Center for Translational Neuroscience (MCTN), Medical Faculty Mannheim of Heidelberg University, Mannheim, Germany, ^{aa} Division of Clinical and Biological Psychology, Department of Psychology, Catholic University of Eichstätt-Ingolstadt, Eichstätt, Germany, ^{ab} Université Paris Saclay, Versailles, France

*Corresponding author. Address: Division of Clinical Psychology and Psychotherapy, Department of Psychology, Philipps-University of Marburg, Gutenbergstr. 18, 35032 Marburg, Germany. Tel.: +49-6421-2824048; fax: +49-6421-2826949. E-mail address: beatrice.korwisi@staff.uni-marburg.de (B. Korwisi).

Supplemental digital content is available for this article. Direct URL citations appear in the printed text and are provided in the HTML and PDF versions of this article on the journal’s Web site (www.painjournalonline.com).

PAIN 162 (2021) 2087–2096

© 2021 International Association for the Study of Pain

<http://dx.doi.org/10.1097/j.pain.0000000000002208>

Text box 1. Basic International Classification of Diseases-11 terminology.**Foundation layer**

The Foundation layer³⁹ of the *International Classification of Diseases-11 (ICD-11)* comprises all entities that are part of the *ICD-11*. It is an evolving structure and forms the basis for the linearizations. Each entity of the foundation has a uniform resource identifier, or Foundation ID.

Linearization

Linearizations are subsets of the Foundation layer. Different specific entities are summarized as diagnoses with a diagnostic code. Currently, the Mortality and Morbidity Statistics linearization (MMS) for preparing implementation³⁸ has been released by the World Health Organization. Countries that implement the *ICD-11* may create their own linearizations.

Diagnostic shoreline

The *ICD-11* MMS draws a "shoreline" at six-digit codes: Specific diagnoses on lower levels of the classification do not receive an individual code. Rather, they should be coded with the respective six-digit code of the level above (the "parent") and specified further by their individual Foundation ID. For example, both chronic primary low back pain and chronic primary cervical pain should both be coded as MG30.02 Chronic primary musculoskeletal pain. They can be specified further and distinguished from one another by their respective ID (1291385632 chronic primary low back pain, 2014134682 chronic primary cervical pain).

Parent and child diagnoses

Different levels and sublevels are referred to as parent and child categories. With regard to the *ICD-11* chronic pain classification, a level 1 diagnosis (eg, MG30.0 Chronic primary pain) is the parent of all level 2 diagnoses it comprises (eg, MG30.02 is a child category). In turn, the level 2 diagnoses are parents of the level 3 diagnoses that follow.

Double parenting

An entity in the *ICD-11* can have multiple parent categories. It may be linked to all parent diagnoses that apply. Despite multiple parents, the entities still have only one immutable diagnostic code with which they are linked to the additional parents. For example, MG30.62 Chronic neuropathic orofacial pain has MG30.6 Chronic secondary headache or orofacial pain as its primary parent. MG30.5 Chronic neuropathic pain is an additional parent. Therefore, MG30.62 Chronic neuropathic orofacial pain is linked there as well. This makes it easier for users to find a specific diagnosis that is relevant to different categories or chapters of the *ICD-11*.

Residual categories

The *ICD-11* distinguishes 2 different residual categories: "unspecified" and "other specified." Both categories are available at the general level of chronic pain as well as for each of the 7 main categories. The "other specified" diagnoses may be used when the user encounters a patient with a very specific disease or syndrome that cannot be allocated to any of the available categories. For example, chronic pain associated with a disease of the skin would be coded as MG30.Y Other specified chronic pain. If the information available is insufficient to decide whether the diagnostic criteria of a given level 1 or level 2 diagnosis are met, the "unspecified" code should be used. For example, a clinician might have sufficient information to determine that a patient has chronic secondary musculoskeletal pain, but not which level 2 diagnosis applies (eg, due to lack of resources). In this case, code MG30.3Z Chronic secondary musculoskeletal pain, unspecified should be assigned.

Coding tool

The *ICD-11* Coding Tool³⁷ should be used for the actual coding. Users enter the diagnostic term they would like to code (eg, "chronic back pain"). The coding tool shows the diagnoses and their respective *ICD-11* codes that match the term. Links to the MMS enable the user to access further information (eg, descriptions) for the different diagnoses found.

All explanations are based on the *ICD-11* Reference Guide³⁹ and the *ICD-11* User Guide.⁴⁰

classification itself¹⁶ (see below), final corrections were made, and final feedback from the pilot users as well as from all taskforce members was implemented. The final version of the CAL-CP was approved by all taskforce members who were involved in its development, and who are coauthoring the present publication.

2.2. Pilot evaluation of the Classification Algorithm for Chronic Pain in International Classification of Diseases-11

A preliminary pilot evaluation of the CAL-CP was integrated in the first phase of the ecological implementation field study of the classification itself (*ICD-11* Chronic Pain Codes Ecological Testing and Assessment [ICE TEA]). The study protocol for the ICE TEA study describes the methods in detail.¹⁶ The first phase of the ICE TEA study was conducted in different countries with varying income levels (Cuba, India, and New Zealand). Ethical approval was obtained before data collection, and all participating clinicians and patients provided their informed consent.

In total, 21 trained pain specialists in 4 different pain clinics in Cuba, India, and New Zealand used a preliminary version of the

CAL-CP to assign *ICD-11* chronic pain diagnoses to 350 patients with chronic pain. After the diagnostic assessment and code assignment, the participating clinicians rated the perceived ease of use, diagnostic confidence, and utility of the classification algorithm on 3 separate numerical rating scales (NRS) ranging from 0 *very difficult/not confident at all/not useful at all* to 10 *very easy/very confident/very useful*. Using SPPS 27 (IBM, Armonk, NY), the mean and SD for each rating scale was computed, and a compound utility score calculated. The pain specialists involved in the ICE TEA study also provided informal feedback on the algorithm, which was recorded at the time and addressed before the final version of the algorithm was approved by all authors.

3. Results

3.1. Structure of the Classification Algorithm for Chronic Pain in International Classification of Diseases-11

The complete CAL-CP is available as supplemental digital content 1 (available at <http://links.lww.com/PAIN/B277>). This PDF document includes the instructions, the introduction form,

the full decision tree including all branches, as well as the appendix.

As described above, the algorithm was developed and graphically implemented following the guidelines of the Society for Medical Decision Making Committee on Standardization of Clinical Algorithms.³¹ **Figure 1** shows examples of the different forms of boxes as implemented in the CAL-CP. The final algorithm consists of a linear decision tree comprising 26 branches and 354 boxes in total. To facilitate the use of the algorithm by providing a clear starting point, an initial decision “trunk” was added. This trunk guides the user to the first branch of the decision tree that is applicable to a given patient, as well as any following relevant branches when a patient has several comorbid chronic pain conditions, by leading to all level 1 diagnoses that apply. Thus, it forms the basis of the CAL-CP (hence, “trunk”). The initial decision trunk can be seen on page 8 of the supplemental digital content 1, available at <http://links.lww.com/PAIN/B277>. Importantly, this initial decision trunk can also be used as a standalone algorithm in less specialized settings, such as primary care. **Figure 2** shows the initial decision trunk/primary care algorithm.

Each branch of the CAL-CP begins with a clinical state box (rectangles with rounded corners). In this box, the starting point (eg, “chronic pain”) is given. In most cases, the starting point of a branch is a level 1 or level 2 diagnosis for which the diagnostic criteria have been met in a previous branch (see the branches on page 10 of supplemental digital content 1 for an example, available at <http://links.lww.com/PAIN/B277>). In these instances, the respective level 1 or level 2 diagnosis is stated in a clinical state box that represents the starting point for that respective branch. These clinical state boxes are thus called “diagnosis boxes” in the CAL-CP. **Figure 3** shows an example branch of the CAL-CP.

As mentioned above, all diagnostic criteria of the *ICD-11* chronic pain classification are represented in individual decision boxes (hexagons). Here, the user must make a dichotomous diagnostic decision, ie, judge whether the criterion is present or absent (eg, whether a given underlying disease is confirmed by a diagnostic test or not). If the diagnostic criteria require a judgement as to whether an underlying disease has been confirmed by a diagnostic test or by imaging, the user may refer to existing test results or images if he or she judges them to be conclusive. Depending on the diagnostic decision, the user then follows the yes or no arrow to the next box. If the user needs to continue in a different section (branch) of the algorithm, action boxes (rectangles) including a page reference have been implemented (eg, page 39 in supplemental digital content 1, available at <http://links.lww.com/PAIN/B277>).

Hyperlinks facilitate the navigation through the algorithm when the document is used in its PDF version in a program that enables hyperlinks. For use as a printed document and as a further means of orientation, all page references are included as comments.




Figure 1. Example boxes of the CAL-CP. A: decision box (p. 8, supplemental digital content 1, <http://links.lww.com/PAIN/B277>), B: action box (p. 16, supplemental digital content 1, <http://links.lww.com/PAIN/B277>), and C: diagnosis box for a level 1 diagnosis (p. 10, supplemental digital content 1, <http://links.lww.com/PAIN/B277>). Underlined text has been implemented as a hyperlink in the CAL-CP. CAL-CP, Classification Algorithm for Chronic Pain in *ICD-11*; *ICD-11*, International Classification of Diseases-11.

When the user follows the boxes and arrows strictly, he or she arrives at a diagnosis box that states the applicable *ICD-11* chronic pain diagnosis. All diagnostic codes listed in the CAL-CP are based on the 09/2020 version of the *ICD-11* (*ICD-11* Mortality and Morbidity Statistics, version for preparing implementation).³⁷ Different forms of lines of the diagnosis boxes (continuous line vs dotted line) indicate the diagnostic level of a given diagnosis.

The algorithm aims at arriving at the most detailed diagnosis (ie, level 3 or 4) whenever possible in a specialty setting (diagnosis box with a continuous line). Diagnoses can be assigned on a less detailed level in less specialized settings (eg, primary care) or in settings with fewer resources. In these settings, the initial decision trunk may be used to assign diagnoses on the first diagnostic level. Hyperlinks and comments for diagnosis boxes instruct the user where to continue to arrive at the diagnosis on the next level.

Entries below the *ICD-11* shoreline (Box 1) are coded with the *ICD-11* diagnostic code of the parent level and can be distinguished further by their Foundation ID. Each entry of the *ICD-11* Foundation layer has a unique Foundation ID or uniform resource identifier. The diagnosis boxes for level 3 diagnoses state the respective Foundation IDs instead of a formal *ICD-11* code. See Box 1 for further explanations.

Very specific chronic pain conditions that do not have an individual entry in the Foundation layer can still be coded with the applicable *ICD-11* code of the more general parent category. For example, chronic pain associated with carpal tunnel syndrome can be coded as MG30.51 Chronic peripheral neuropathic pain without being specified further by an individual ID. For these cases, level 2 diagnosis boxes are repeated at the end of each level 3 branch (see **Fig. 3** for an example). For a better overview, these boxes are labeled as “other specified,” although this is not part of the official *ICD-11* diagnosis. If a chronic pain condition cannot be allocated to any of the level 2 diagnoses, a specific decision box will guide the user to the respective residual category. See Box 1 for more details on the *ICD-11* residual categories of “other specified” and “unspecified.”

When arriving at a diagnosis box for a chronic secondary pain condition (Box 2), the *ICD-11* chronic pain code must be combined with the *ICD-11* code of the underlying disease associated with that given chronic pain condition. The *ICD-11* Coding Tool³⁶ provided by the WHO can be used for this complete coding.

As mentioned above, all boxes are numbered consecutively. Explanatory comments accompany some boxes. In these cases, the number of the box automatically serves as the footnote for the comment that goes with the respective box. The comments are added below each branch or on the following page. Some comments give further instructions (eg, page references in addition to the hyperlinks), whereas others list details or examples for a given diagnostic criterion (eg, examples of diseases that may be associated with a given chronic secondary pain diagnosis, such as rheumatoid arthritis and MG30.30 Chronic secondary musculoskeletal pain from persistent inflammation). Each diagnosis box has a feedback loop to remind the user to check the pain location chart (page 7, supplement digital content 1, available at <http://links.lww.com/PAIN/B277>) and initial decision trunk (page 8, supplement digital content 1, available at <http://links.lww.com/PAIN/B277>) to ensure all chronic pain syndromes of the patient have been accounted for. If additional chronic pain syndromes are present, the user must continue with the next applicable branch of the CAL-CP as highlighted in the initial decision trunk. This ensures that despite the linear structure of the decision tree, no chronic pain condition is missed in patients with several comorbid chronic pain conditions.




Figure 2. Primary care version of the CAL-CP (initial decision trunk). This initial decision trunk (p. 8, supplemental digital content 1, available at <http://links.lww.com/PAIN/B277>) can be used as a standalone algorithm to assign diagnoses on the first diagnostic level in settings where a more detailed diagnosis is not necessary (eg, primary care). Comments for some of the boxes can be found on p. 9, supplemental digital content 1 (available at <http://links.lww.com/PAIN/B277>). When a diagnosis is assigned on level 1, check if all chronic pain has been accounted for. If additional chronic pain syndromes are present, go through this “trunk” again to assign all diagnoses that apply. All boxes of the CAL-CP are numbered consecutively. Underlined text has been implemented as a hyperlink in the CAL-CP. CAL-CP, Classification Algorithm for Chronic Pain in ICD-11; ICD-11, *International Classification of Diseases-11*; ICHD, third edition of the International Classification of Headache Disorders; ICOP, International Classification of Orofacial Pain




Figure 3. Example branch of the CAL-CP. This figure shows the branch for chronic primary musculoskeletal pain. Underlined text has been implemented as a hyperlink in the CAL-CP. This branch has been implemented on page 13 of the CAL-CP (supplemental digital content 1, available at <http://links.lww.com/PAIN/B277>). The sign on top of boxes 60, 62, 64, and 66 is a reminder to check in the pain location chart and the initial decision trunk whether all chronic pain has been accounted for. All boxes of the CAL-CP are numbered consecutively. CAL-CP, Classification Algorithm for Chronic Pain in ICD-11; ICD-11, International Classification of Diseases-11.

3.2. How to use the Classification Algorithm for Chronic Pain in International Classification of Diseases-11

As outlined above, the CAL-CP consists of detailed instructions for its use (p. 1-5, supplement digital content 1, available at <http://links.lww.com/PAIN/B277>), an introduction form (p. 6-7, supplement digital content 1, available at <http://links.lww.com/PAIN/B277>), and an appendix (p. 41-45, supplement digital content 1, available at <http://links.lww.com/PAIN/B277>), besides the actual decision tree (p. 8-40, supplement digital content 1, available at

<http://links.lww.com/PAIN/B277>). With regard to the procedure for its use, the user begins with the introduction form. This comprises a general red flags question. In this context, red flags include the presence of an underlying disease (such as cancer or osteoarthritis) without reference to its seriousness. The user should assess whether the patient presents with any symptoms suggestive of an underlying disease that has not been diagnosed previously (eg, an undiagnosed cancer that might explain the pain). The assessment of these red flags should follow the standard guidelines and diagnostic routines in the respective field of expertise. Unnecessary diagnostic examinations should be avoided. If an underlying undiagnosed disease is suspected, the user should take the appropriate steps (eg, referral or further diagnostics) as he or she would usually do. The algorithm can be continued after these examinations have clarified the situation.

If a patient presents with chronic headache or orofacial pain, the user will need to refer directly to the ICHD-3¹⁰ or ICOP¹¹ for the respective diagnostic criteria, as well as to the ICD-11 Coding Tool³⁷ for diagnostic coding. Action boxes have been implemented in the CAL-CP to facilitate these references. Furthermore, the initial decision trunk/primary care version refers to the broader diagnoses that are available in the ICD-11 for chronic primary and secondary headache or orofacial pain.

Once the user has ensured that no red flags warrant further medical attention, he or she should continue with the assessment of the so-called chronic pain specifiers (Box 2). Importantly, these should be rated by the patient. Furthermore, as far as possible, the timing of the onset of the chronic pain should be documented. Then, the patient should highlight all body regions on the pain location chart where he or she experiences chronic pain. This chart guides the user through the algorithm. For example, if 2 separate body regions are highlighted, the user will see at one glance that he or she should account for both locations in his or her diagnosis. Then, the decision tree itself begins with an initial decision trunk as outlined above. The user marks all sections he or she will need to assess depending on the patient's medical history (ie, known presence or absence of any underlying disease that might be associated with the chronic pain) and first diagnostic criteria. In addition to the pain location chart, the marks on this trunk also facilitate the assessment of patients with several comorbid chronic pain conditions. The initial decision trunk also gives page references and hyperlinks to facilitate the beginning with the first branch of the decision tree that is relevant to a given patient. This initial decision tree leads to the level 1 diagnoses and can be used as a standalone algorithm in less specialized or primary care settings. In specialized pain treatment settings, such as multimodal pain treatment, the full algorithm should be used. After the initial decision trunk, the user assesses the relevant branches as described above. If the medical history of a patient does not suggest that an underlying disease is associated with the chronic pain, the user will directly begin with the assessment of chronic primary pain. No extensive exclusion diagnostics are required.

During the use of the CAL-CP, the user may refer to the appendix with a list of exemplary diseases that may be associated with chronic pain, if needed (p. 41-45, supplement digital content 1, available at <http://links.lww.com/PAIN/B277>). The appendix is based on the descriptions of each chronic pain entity in the ICD-11 Browser³⁸ as well as the publications on the different chronic pain categories^{1,2,21,24,29,30} and expert feedback by the task-force members. It aims at giving an overview of which chronic secondary pain category can be associated with which underlying diseases (eg, sickle cell disease may be associated with MG30.41 Chronic visceral pain from vascular mechanisms). The

Text box 2. New concepts of the International Classification of Diseases-11 chronic pain classification.

Pain severity

Pain severity is one of the newly defined chronic pain specifiers. It is a compound measure of pain intensity, pain-related distress, and pain-related Interference. The patient is requested to rate the 3 dimensions of pain severity on a 0 to 10 numerical rating scale or a 10 cm visual analogue scale. These ratings can be converted to the World Health Organization severity categories “mild,” “moderate,” and “severe.”

Chronic primary pain

In chronic primary pain, the pain is a health condition in its own right. Chronic primary pain is characterized by significant emotional distress or functional interference. It is multifactorial, with different biological, psychological, and social factors contributing to it. Furthermore, chronic primary pain is not better accounted for by another (secondary) chronic pain diagnosis.

Chronic secondary pain

Chronic secondary pain, as any pain, is also multifactorial. However, chronic secondary pain is pain that develops as a symptom in the context of an underlying disease (eg, rheumatoid arthritis and cancer). Nevertheless, it warrants tailored pain treatment. The severity of chronic secondary pain is not necessarily equal to the severity of the underlying disease process. In some cases, the pain persists after the underlying disease has been treated successfully.

The *International Classification of Diseases-11* chronic pain classification comprises 6 different categories of chronic secondary pain: chronic cancer-related pain (eg, associated with a tumor or cancer treatment), chronic postsurgical or post traumatic pain (eg, after spinal surgery or after bone fractures), chronic neuropathic pain (eg, associated with polyneuropathy or stroke), chronic secondary headache or orofacial pain (eg, associated with infection), chronic secondary visceral pain (eg, associated with Crohn disease or sickle cell disease), and chronic secondary musculoskeletal pain (eg, associated with spondylosis or gout). Further details can be found in Treede et al., 2019.³⁴

presence of any of these diseases is not automatically associated with chronic pain. Even if an underlying disease is present, the branch for the chronic pain condition that accompanies it should be assessed carefully.

3.3. Results of the pilot evaluation

The pain specialists participating in the ICE TEA study rated the ease of use, diagnostic confidence, and perceived utility of the algorithm as very high. **Table 1** gives the mean score and SD for each scale. Combining these 3 measures to one global utility score revealed a mean utility rating of 8.48 ± 1.67 (NRS 0-10).

4. Discussion

With the CAL-CP, we present a comprehensive classification algorithm for the *ICD-11* chronic pain classification for use in clinical practice and pain research. The CAL-CP provides a structured linear decision tree for the new classification of chronic pain, covering all 4 diagnostic levels of the classification. Its use will facilitate and standardize the process of finding the correct *ICD-11* chronic pain diagnosis, contributing to the reliability of the diagnoses.

The different diagnostic levels enable the use of the algorithm in specialty settings, where a very detailed diagnosis on level 3 or 4 of the classification is needed, as well as in less specialized settings (eg, primary care), where a diagnosis on levels 1 or 2 may be sufficient.³⁴ Importantly, the initial decision trunk (**Fig. 2**) serves as primary care version of the CAL-CP. It leads to all level 1 diagnoses that apply to a given patient, and thus represents a time-efficient and easy-to-use tool in settings where time is a limited resource, and where a more detailed diagnosis might not be necessary. In research settings, use of the full CAL-CP can contribute to increased standardization by providing a clearly defined way through the classification process. Importantly, the CAL-CP can also be used with patient records when these include all the clinical information needed.

An introduction form as well as an initial decision trunk provides guidance for the assessment of a given patient (ie, which sections [branches] are relevant). The introduction form provides helpful guidance for the use of the CAL-CP, but it might not be needed when the CAL-CP is used to code a patient based on medical records. The appendix, which gives a list of common diseases that may be associated with chronic pain, is a helpful guide when a patient who presents with a medical history extending beyond the user’s main field of expertise is assessed. For example, a clinician involved in multimodal pain treatment might not have specialist knowledge regarding the details of internal diseases that can be associated with chronic pain. Here, it might be difficult to judge whether chronic pain in the context of a given disease (eg, vasculitis) falls within the category of MG30.41 Chronic secondary visceral pain from vascular mechanisms or within the category of MG30.42 Chronic secondary visceral pain from persistent inflammation. In these instances, the user may refer to the appendix for further guidance.

Hyperlinks simplify the navigation through the algorithm. They are supplemented by complete page references, which are included in the comments. This enables the use of a digital PDF version as well as a printed version of the CAL-CP. Furthermore, a feedback loop in the comments section refers users to the pain location chart and the initial decision trunk once a diagnosis box is

Table 1
Results of the pilot evaluation of the classification algorithm.

Item	Mean (M)	SD
Ease of use	8.53	1.68
Diagnostic confidence	8.43	1.84
Utility	8.49	1.82
Global utility score	8.48	1.67

Ratings of 350 patients with chronic pain by 21 pain specialists. Ease of use, diagnostic confidence, and utility of the algorithm were rated on 3 separate NRS from 0 (very difficult/not confident at all/not useful at all) to 10 (very easy/very confident/very useful). The global utility score is the compound measure of these 3 ratings.

NRS, numerical rating scale.

reached. This ensures that all comorbid chronic pain conditions are considered during the assessment, and that no chronic pain condition is overlooked in a patient.

It should be noted that the CAL-CP does not substitute for a thorough clinical examination of the patient's pain problem. The CAL-CP will not find out whether a patient's chronic pain is caused by an underlying disease process. Rather, the algorithm facilitates navigating through the *ICD-11* chronic pain criteria when the user has all the necessary clinical information. Of course, it may also highlight in which area such information may still be lacking. If a user suspects that a hitherto undiagnosed condition causes the pain, clinical judgement is needed to take the appropriate diagnostic steps. This is in line with other medical algorithms where the developers also point out that an algorithm only assists in describing, summarizing, and classifying available information, and can never substitute for thorough clinical decision making.^{12,15,42} The CAL-CP, as with many classification algorithms, is a decision aid and cannot make the decision in lieu of a clinician. At the same time, it should be noted that the goal should not be to formally exclude any possible underlying disease with extensive diagnostics. Rather, red flags should be assessed as always in routine clinical practice, and unnecessary diagnostic procedures should be avoided.^{5,25}

Classification algorithms are common in a variety of medical fields and have been proposed for different classifications, including epilepsy,²⁸ gastrointestinal disorders,¹⁵ neuropathic pain,⁸ low back pain,²² osteoarthritis,¹⁸ periodontal diseases,³³ and abdominal pain.¹⁴ The CAL-CP hence represents an important addition to existing algorithm efforts.

The advantages of algorithms to aid classification processes in medicine, including pain medicine, have been described extensively in the literature: algorithms can provide valuable assistance for diagnostic and classification processes by illustrating the decisions to be made by the user (eg, in the case of the CAL-CP, whether a diagnostic criterion is present) in a comprehensible step-by-step sequence,¹² and by providing guidance through a classification with its different criteria.³³ The structured assessment of the diagnostic criteria is facilitated considerably by this logical decision tree.^{15,18,23} Furthermore, adherence to the diagnostic criteria increases when algorithms are used.⁴ By guiding the user through all criteria that have to be assessed, the CAL-CP ensures that none of the compulsory criteria are missed when assigning an *ICD-11* chronic pain diagnosis. This, in turn, contributes to efficiency,^{4,32} diagnostic accuracy,²⁰ diagnostic consistency,^{6,27} as well as increased reliability of the diagnoses.^{17,28} Furthermore, the use of algorithms during diagnostic and classification processes can reduce, but not eliminate, errors.¹²

Reliable diagnoses contribute to increased clinical utility of the *ICD-11* chronic pain classification itself, including patient management and documentation.⁹ Furthermore, they are essential for pain research as reliable diagnoses form the basis of accurate sample descriptions as well as data collection.

The data from the preliminary pilot evaluation of the algorithm indicate its high utility. The clinicians judged it easy to use and reported high diagnostic confidence. This is in line with other research on algorithms that also highlight their clinical utility as well as their practical value (eg, ease of interpretation and user-friendliness).^{18–20,26} Notably, the pilot evaluation was conducted in different countries with different income levels (Cuba, India, and New Zealand). Hence, the results provide an initial demonstration that the algorithm is applicable in a variety of settings.

A further important future application of the CAL-CP involves training and education. This is especially relevant as the *ICD-11*

chronic pain classification is new, and its worldwide implementation is imminent. An urgent task ahead of implementation is training: pain clinicians and pain researchers from all fields (medicine, psychology, physical therapy, among others) as well as clinicians with other specialties and professional coders will have to be trained and familiarize themselves with the new criteria and diagnoses. It has been shown that minimal training is sufficient for reliable diagnoses when decision trees are used as a way to navigate through new diagnostic criteria.¹⁷ Furthermore, decision trees similar to the CAL-CP have been demonstrated to be even more helpful to novices during a diagnostic process compared to experts.²⁰ Here, the diagnostic accuracy increased more in novices than in experts when an algorithm provided guidance through diagnostic criteria to assign a diagnosis. This highlights the importance and utility of algorithms for training purposes.

Although it was a reasoned decision not to include chronic headache or orofacial pain in the CAL-CP, some might consider this a limitation of the present algorithm. However, references to the ICHD-3¹⁰ and the ICOP¹¹ have been integrated into the decision tree using action boxes that refer to these classifications. Furthermore, comments of the respective action boxes list the *ICD-11* diagnoses for chronic primary headache or orofacial pain and chronic secondary headache or orofacial pain, respectively. The ICHD-3 and the ICOP should be available alongside the *ICD-11* in clinical settings. Furthermore, the CAL-CP in its current form is a long document, and its application will need prior training.

Future studies should investigate whether the preliminary results of the pilot evaluation are corroborated. A computer-based international evaluation study is currently in preparation. Detailed case vignettes will be implemented in the form of virtual patients. The use of standardized case vignettes allows control over patient variables.¹³ To gather the diagnostic information as needed, participants will be able to elicit information from virtual patients through chatbot technology. The computer-based implementation of this study also allows assessment of the time users need to use the CAL-CP. Future plans for the algorithm also include its preparation as a digital application or online format. However, a successful online evaluation is a prerequisite of such an effort. Future research should also include clinicians with different backgrounds such as primary care physicians who work with patients with chronic pain.

In conclusion, the CAL-CP provides a useful and easy-to-use decision aid that can guide pain clinicians as well as pain researchers through the new *ICD-11* classification of chronic pain.

Conflict of interest statement

B. Korwisi reports other from IASP (NGO), during the conduct of the study. G. Hay has nothing to disclose. N. Attal reports personal fees from Lilly, personal fees from Pfizer, personal fees from Sanofi, personal fees from Grunenthal, personal fees from Ipsen, personal fees from Air Liquide, personal fees from Novartis, personal fees from Aptinyx, personal fees from Novartis, outside the submitted work. Q. Aziz reports grants and personal fees from Grunenthal pharmaceutical, grants and personal fees from Allergan Pharmaceuticals, grants from Alimentary Health, outside the submitted work. M.I. Bennett has nothing to disclose. R. Benoliel reports personal fees from Quintessence Publishing, outside the submitted work. M. Cohen has nothing to disclose. S. Evers reports personal fees from Allergan, personal fees from Novartis, personal fees from TEVA, personal fees from Lilly, outside the submitted work. M.A. Giamberardino reports

personal fees from HELSINN HEALTHCARE, personal fees from EPITECH GROUP, personal fees from IBSA, outside the submitted work. S. Kaasa has nothing to disclose. E. Kosek reports personal fees from Lundbeck, personal fees from Eli Lilly, outside the submitted work. P. Lavand'homme has nothing to disclose. M. Nicholas has nothing to disclose. S. Perrot has nothing to disclose. S. Schug has nothing to disclose. B.H. Smith has nothing to disclose. P. Svensson has nothing to disclose. J.W.S. Vlaeyen reports grants from Asthenes long-term structural funding–Methusalem grant by the Flemish Government, Belgium (METH/15/011), outside the submitted work. S.J. Wang reports grants and personal fees from Eli-Lilly, personal fees from Daiichi-Sankyo, grants and personal fees from Novartis, grants from Minister of Science and Technology, Taiwan, grants from The Featured Areas Research Center Program within the framework of the Higher Education Sprout Project by the Ministry of Education (MOE), from null, outside the submitted work. R.-D. Treede reports grants from Teva, personal fees from Bayer, Grünenthal, GSK, Sanofi, outside the submitted work. W. Rief has nothing to disclose. A. Barke reports other from IASP (NGO), outside the submitted work.

The development of the classification algorithm was presented as a poster abstract at the 2018 IASP World Congress on Pain in Boston, MA (initial developmental stage) and at the 2020 German Pain Conference, online (preliminary version).

Acknowledgments

The authors gratefully acknowledge Christiane Blöcher for her assistance with the preparation and graphical implementation of the CAL-CP, Karoline Körfer and Tabea Kloos for their expert feedback on preliminary versions of the CAL-CP, as well as Dr Bárbara Beatriz Garrido Suárez, Dr Subrata Goswami, Dr Muralidhar Joshi, and Dr Maria Kleinstäuber for their valuable feedback and support with the preliminary pilot evaluation of the CAL-CP.

The IASP has financially supported the development of a chronic pain classification system for the *ICD-11*, by providing research grants and travel grants to the authors.

Appendix A. Supplemental digital content

Supplemental digital content associated with this article can be found online at <http://links.lww.com/PAIN/B277>.

Supplemental video content

A video abstract associated with this article can be found at <http://links.lww.com/PAIN/B278>.

Article history:

Received 23 October 2020

Received in revised form 7 December 2020

Accepted 11 December 2020

Available online 20 January 2021

References

- [1] Aziz Q, Giamberardino MA, Barke A, Korwisi B, Baranowski A, Wesselmann U, Rief W, Treede R-D; The IASP Taskforce for the Classification of Chronic Pain. The IASP classification of chronic pain for ICD-11: chronic secondary visceral pain. *PAIN* 2019;160:69–76.
- [2] Bennett MI, Kaasa S, Barke A, Korwisi B, Rief W, Treede R-D; The IASP Taskforce for the Classification of Chronic Pain. The IASP classification of chronic pain for ICD-11: chronic cancer-related pain. *PAIN* 2019;160:38–44.
- [3] Benoliel R, Svensson P, Evers S, Wang S-J, Barke A, Korwisi B, Rief W, Treede R-D; The IASP Taskforce for the Classification of Chronic Pain. The IASP classification of chronic pain for ICD-11: chronic secondary headache and orofacial pain. *PAIN* 2019;160:60–8.
- [4] Bollestad M, Grude N, Lindbaek M. A randomized controlled trial of a diagnostic algorithm for symptoms of uncomplicated cystitis at an out-of-hours service. *Scand J Prim Health Care* 2015;33:57–64.
- [5] Bouck Z, Pendrith C, Chen XK, Frood J, Reason B, Khan T, Costante A, Kirkham K, Born K, Levinson W, Bhatia RS. Measuring the frequency and variation of unnecessary care across Canada. *BMC Health Serv Res* 2019;19:446.
- [6] Coleman SH. Editorial commentary: the importance of developing an algorithm when diagnosing hip pain. *Arthrosc - J Arthrosc Relat Surg* 2016;32:1712–13.
- [7] Drossman DA, Hasler WL. Rome IV—functional GI disorders: disorders of gut-brain interaction. *Gastroenterology* 2016;150:1257–61.
- [8] Finnerup NB, Haroutounian S, Kamerman P, Baron R, Bennett DLH, Bouhassira D, Cruccu G, Freeman R, Hansson P, Nurmikko T, Raja SN, Rice ASC, Serra J, Smith BH, Treede R-D, Jensen TS. Neuropathic pain: an updated grading system for research and clinical practice. *PAIN* 2016;157:1599–606.
- [9] First MB. Clinical utility in the revision of the diagnostic and statistical manual of mental disorders (DSM). *Prof Psychol Res Pract* 2010;41:465–73.
- [10] Headache Classification Committee of the International Headache Society (IHS). The international classification of headache disorders, 3rd edition. *Cephalalgia* 2018;38:1–211.
- [11] International classification of orofacial pain, 1st edition (ICOP). *Cephalalgia* 2020;40:129–221.
- [12] Jablonski AM, DuPen AR, Ersek M. The use of algorithms in assessing and managing persistent pain in older adults. *Am J Nurs* 2011;111:34–45.
- [13] Keeley J, Reed G, Roberts M, Evans SC, Medina-Mora ME, Robles R, Rebello T, Sharan P, Gureje O, First MB, Andrews HF, Ayuso-Mateos JL, Gaebel W, Zielasek J, Saxena S. Developing a science of clinical utility in diagnostic classification systems field study strategies for ICD-11 mental and behavioral disorders. *Am Psychol* 2016;71:3–16.
- [14] Keller J, Layer P. Chronischer Oberbauchschmerz: diagnostischer und therapeutischer Algorithmus. *Dtsch Med Wochenschr* 2015;140:718–22.
- [15] Kellow JE, Drossman DA. Rome Foundation diagnostic algorithms for common gastrointestinal symptoms. *Am J Gastroenterol* 2010;75:736–801.
- [16] Korwisi B, Treede R, Rief W, Barke A. Evaluation of the International Classification of Diseases-11 chronic pain classification: study protocol for an ecological implementation field study in low-, middle-, and high-income countries. *PAIN Rep* 2020;5:e825.
- [17] Malt UF. Teaching DSM-III to clinicians. Some problems of the DSM-III system reducing reliability, using the diagnosis and classification of depressive disorders as an example. *Acta Psychiatr Scand* 1986;73:68–75.
- [18] Martel-Pelletier J, Maheu E, Pelletier JP, Alekseeva L, Mkinci O, Branco J, Monod P, Planta F, Reginster JY, Rannou F. A new decision tree for diagnosis of osteoarthritis in primary care: international consensus of experts. *Aging Clin Exp Res* 2019;31:19–30.
- [19] Moloney NA, Hall TM, Leaver AM, Doody CM. The clinical utility of pain classification in non-specific arm pain. *Man Ther* 2015;20:157–65.
- [20] Morgan RD, Olson KR, Krueger RM, Schellenberg RP, Jackson TT. Do the DSM decision trees improve diagnostic ability? *J Clin Psychol* 2000;56:73–88.
- [21] Nicholas M, Vlaeyen JWS, Rief W, Barke A, Aziz Q, Benoliel R, Cohen M, Evers S, Giamberardino MA, Goebel A, Korwisi B, Perrot S, Svensson P, Wang S-J, Treede R-D; The IASP Taskforce for the Classification of Chronic Pain. The IASP classification of chronic pain for ICD-11: chronic primary pain. *PAIN* 2019;160:28–37.
- [22] Nijs J, Apeldoorn A, Hallegraef H, Clark J, Smeets R, Malfliet A, Gírbés EL, De Kooning M, Ickmans K. Low back pain: guidelines for the clinical classification of predominant neuropathic, nociceptive, or central sensitization pain. *Pain Physician* 2015;18:E333–46.
- [23] Park CN, White PB, Meftah M, Ranawat AS, Ranawat CS. Diagnostic algorithm for residual pain after total knee arthroplasty. *Orthopedics* 2016;39:e246–52.
- [24] Perrot S, Cohen M, Barke A, Korwisi B, Rief W, Treede R-D. The IASP Taskforce for the Classification of Chronic Pain. The IASP classification of chronic pain for ICD-11: chronic secondary musculoskeletal pain. *PAIN* 2019;160:77–82.

- [25] Pichetshote N, Pimentel M. An approach to the patient with chronic undiagnosed abdominal pain. *Am J Gastroenterol* 2019;114:726–32.
- [26] Podgorelec V, Kokol P, Stiglic B, Rozman I. Decision trees: an overview and their use in medicine. *J Med Syst* 2002;26:445–63.
- [27] Prakash S, Rathore C. Side-locked headaches: an algorithm-based approach. *J Headache Pain* 2016;17:95.
- [28] Rinaldi G, Zarrelli MM, Beghi E, Apollo F, Germano M, Di Viesti P, Simone P. The international classification of the epilepsies and epileptic syndromes: an algorithm for its use in clinical practice. *Epilepsy Res* 2000;41:223–34.
- [29] Scholz J, Finnerup NB, Attal N, Aziz Q, Baron R, Bennett MI, Benoliel R, Cohen M, Cruccu G, Davis KD, Evers S, First MB, Giamberardino MA, Hansson P, Kaasa S, Korwisi B, Kosek E, Lavand'homme P, Nicholas M, Nurmikko T, Perrot S, Raja SN, Rice ASC, Rowbotham MC, Schug S, Simpson DM, Smith BH, Svensson P, Vlaeyen JWS, Wang S-J, Barke A, Rief W, Treede R-D; Classification Committee of the Neuropathic Pain Special Interest Group (NeuPSIG) Abstract. The IASP classification of chronic pain for ICD-11: chronic neuropathic pain. *PAIN* 2019;160:53–9.
- [30] Schug SA, Lavand'homme P, Barke A, Korwisi B, Rief W, Treede R-D; The IASP Taskforce for the Classification of Chronic Pain. The IASP classification of chronic pain for ICD-11: chronic postsurgical or posttraumatic pain. *PAIN* 2019;160:45–52.
- [31] Society for Medical Decision Making Committee on Standardization of Clinical Algorithms. Proposal for clinical algorithm standards. *Med Decis Mak* 1992;38:149–54.
- [32] Tallen G, Bielack S, Henze G. Musculoskeletal pain: a new algorithm for differential diagnosis of a cardinal symptom in pediatrics. *Klin Pediatr* 2014;226:86–98.
- [33] Tonetti MS, Sanz M. Implementation of the new classification of periodontal diseases: decision-making algorithms for clinical practice and education. *J Clin Periodontol* 2019;46:398–405.
- [34] Treede R-D, Rief W, Barke A, Aziz Q, Bennett MI, Benoliel R, Cohen M, Evers S, Finnerup NB, First MB, Giamberardino MA, Kaasa S, Korwisi B, Kosek E, Lavand'homme P, Nicholas M, Perrot S, Scholz J, Schug S, Smith BH, Svensson P, Vlaeyen JWS, Wang S-J; Chronic pain as a symptom or a disease: the IASP classification of chronic pain for the international classification of diseases ICD-11. *PAIN* 2019;160:19–27.
- [35] Treede R-D, Rief W, Barke A, Aziz Q, Bennett MI, Benoliel R, Cohen M, Evers S, Finnerup NB, First MB, Giamberardino MA, Kaasa S, Kosek E, Lavand'homme P, Nicholas M, Perrot S, Scholz J, Schug S, Smith BH, Svensson P, Vlaeyen JWS, Wang S-J. A classification of chronic pain for ICD-11. *PAIN* 2015;156:1003–7.
- [36] World Health Organization. ICD-11 coding tool mortality and morbidity statistics (MMS) 2020-09. 2020. Available at: https://icd.who.int/ct11/icd11_mms/en/release. Accessed December 1, 2020.
- [37] World Health Organization. ICD-11 for mortality and morbidity statistics (ICD-11 MMS) version 09/2020. 2020. Available at: <https://icd.who.int/browse11/l-m/en>. Accessed December 1, 2020.
- [38] World Health Organization. ICD-11 maintenance platform. 2020. Available at: <https://icd.who.int/dev11/l-m/en>. Accessed December 1, 2020.
- [39] World Health Organization. ICD-11 reference guide. 2019. Available at: <https://icd.who.int/icd11refguide/en/index.html>. Accessed December 1, 2020.
- [40] World Health Organization. ICD-11 user guide. 2020. Available at: <https://icd.who.int/dev11/Help/en>. Accessed December 1, 2020.
- [41] World Health Organization. World health assembly update, 25 May 2019. 2019. Available at: <https://www.who.int/news-room/detail/25-05-2019-world-health-assembly-update>. Accessed December 1, 2020.
- [42] Yelland M, Cayley WE, Vach W. An algorithm for the diagnosis and management of chest pain in primary care. *Med Clin North Am* 2010;94:349–74.