

# Comment on "Guidelines of care for the management of basal cell carcinoma"

Citation for published version (APA):

van Delft, L. C. J., Jansen, M. H. E., Mosterd, K., & Kelleners-Smeets, N. W. J. (2018). Comment on "Guidelines of care for the management of basal cell carcinoma". *Journal of the American Academy of Dermatology*, 79(5), E99-E100. <https://doi.org/10.1016/j.jaad.2018.05.1261>

## Document status and date:

Published: 01/11/2018

## DOI:

[10.1016/j.jaad.2018.05.1261](https://doi.org/10.1016/j.jaad.2018.05.1261)

## Document Version:

Publisher's PDF, also known as Version of record

## Document license:

Taverne

## Please check the document version of this publication:

- A submitted manuscript is the version of the article upon submission and before peer-review. There can be important differences between the submitted version and the official published version of record. People interested in the research are advised to contact the author for the final version of the publication, or visit the DOI to the publisher's website.
- The final author version and the galley proof are versions of the publication after peer review.
- The final published version features the final layout of the paper including the volume, issue and page numbers.

[Link to publication](#)

## General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal.

If the publication is distributed under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license above, please follow below link for the End User Agreement:

[www.umlib.nl/taverne-license](http://www.umlib.nl/taverne-license)

## Take down policy

If you believe that this document breaches copyright please contact us at:

[repository@maastrichtuniversity.nl](mailto:repository@maastrichtuniversity.nl)

providing details and we will investigate your claim.

**Comment on “Guidelines of care for the management of basal cell carcinoma”**



*To the Editor:* With interest we read the new guidelines for the management of basal cell carcinoma (BCC) by the American Academy of Dermatology.<sup>1</sup> The objective was to provide an updated review (based on studies published up to August 2016) of the evidence and recommendations for diagnostic, therapeutic, and preventative modalities for US dermatologists. The authors did a great job judging the extensive amount of available evidence used in creating these guidelines. We agree with encouraging a patient-specific approach. Surgical excision remains superior to other treatments in terms of efficacy, though practical considerations and cosmetic outcome could counterbalance this choice.<sup>2</sup> As an addition, we have some suggestions with regard to the treatment of low-risk BCC with imiquimod, and we wish to address the position of radiation therapy (RT) in treatment of BCC.

The US guidelines advise initiating additional trials to compare the efficacy of noninvasive therapies for BCC. Fortunately, after the search on which the updated guidelines are based was completed, 2 large, well-executed randomized controlled trials (RCTs) were published.<sup>3,4</sup> Williams et al. report the 5-year results of an RCT comparing imiquimod to surgery in superficial and nodular BCC.<sup>4</sup> They conclude that surgery remains the criterion standard on the basis of its efficacy in 97.7% (relative risk of imiquimod compared with that of surgery, 0.84 98% confidence interval [CI], 0.77-0.91). However, with an 82.5% clearance rate, imiquimod was deemed an important treatment modality for low-risk superficial and nodular BCC.

Our group recently described the 5-year results of an RCT comparing imiquimod, 5-fluorouracil, and methyl-aminolevulinic photodynamic therapy (MAL-PDT).<sup>3</sup> Imiquimod was superior to MAL-PDT and 5-fluorouracil in terms of efficacy. The rates of tumor-free survival were 62.7% (95% CI, 55.3-69.2%) for MAL-PDT, 70.0% (95% CI, 62.9-76.0%) for 5-fluorouracil, and 80.5% (95% CI, 74.0-85.6%) for imiquimod. Imiquimod is therefore considered the first-choice noninvasive treatment for superficial BCC.

In our opinion, both trials provide good-quality evidence to conclude that imiquimod is a reliable treatment for patients with low-risk superficial BCC as an alternative to surgery, with excellent cosmetic outcome and the possibility of at-home treatment.

With regard to the position of RT in treatment of BCC, both the US guidelines and the European

Dermatology Forum guideline<sup>2</sup> conclude that RT should be reserved for cases in which surgery is not feasible; however, the US guidelines align RT with noninvasive therapies for low-risk BCCs whereas the European Dermatology Forum guidelines reserve RT for high-risk BCCs.<sup>2</sup>

Until now, only 2 RCTs on RT for BCC have been published; both are dated, and RT techniques have changed extensively.<sup>5</sup> Thus, the current conclusion on the position of RT in treatment of BCC depends mostly on expert opinion.

There are many (noninvasive) alternatives for treatment of low-risk BCCs; therefore, in our opinion RT should be reserved for elderly patients with high-risk (facial) BCC, for whom surgery is unsuitable.<sup>2</sup>

Patients and physicians can make an educated treatment choice only after deliberation on all options and the advantages and disadvantages for their situation. We would like to add that there is appropriate evidence for the efficacy and safety of imiquimod for low-risk superficial BCC. Further evidence is needed to establish the position of RT in treatment of BCC.

Lieke C. J. van Delft, MD,<sup>a,b</sup> Maud H. E. Jansen, MD,<sup>a,b</sup> Klara Mosterd, MD, PhD,<sup>a,b</sup> and Nicole W. J. Kelleners-Smeets, MD, PhD<sup>a,b</sup>

*From the Department of Dermatology, Maastricht University Medical Centre+, Maastricht, The Netherlands,<sup>a</sup> and GROW-School for Oncology and Developmental Biology Maastricht University Medical Centre+, Maastricht, The Netherlands<sup>b</sup>*

*Funding sources: None.*

*Disclosure: Dr Mosterd has received sachets of imiquimod from Meda Connects (now called Mylan Connects) for an unrelated trial (NCT02242929 on <https://clinicaltrials.gov/>). The other authors declared no conflicts of interest.*

*Reprints not available from the authors.*

*Correspondence to: Lieke C. J. van Delft, Maastricht University Medical Centre+, PO Box 5800, 6229 HX Maastricht, The Netherlands*

*E-mail: [lieke.van.delft@mumc.nl](mailto:lieke.van.delft@mumc.nl)*

**REFERENCES**

1. Kim JYS, Kozlow JH, Mittal B, Moyer J, Olencki T, Rodgers P. Guidelines of care for the management of

- basal cell carcinoma. *J Am Acad Dermatol.* 2018;78(3):540-559.
2. Trakatelli M, Morton C, Nagore E, et al. Update of the European guidelines for basal cell carcinoma management. *Eur J Dermatol.* 2014;24(3):312-329.
  3. Jansen MHE, Mosterd K, Arits AHMM, et al. Five-Year results of a randomized controlled trial comparing effectiveness of photodynamic therapy, topical imiquimod, and topical 5-fluorouracil in patients with superficial basal cell carcinoma. *J Invest Dermatol.* 2018;138(3):527-533.
  4. Williams HC, Bath-Hextall F, Ozolins M, et al. Surgery versus 5% imiquimod for nodular and superficial basal cell carcinoma: 5-year results of the SINS randomized controlled trial. *J Invest Dermatol.* 2017;137(3):614-619.
  5. Cho M, Gordon L, Rembielak A, Woo TC. Utility of radiotherapy for treatment of basal cell carcinoma: a review. *Br J Dermatol.* 2014;171(5):968-973.

<https://doi.org/10.1016/j.jaad.2018.05.1261>