Taking control

Citation for published version (APA):

Houben, K., Dassen, F., & Jansen, A. (2016). Taking control: Working memory training in overweight individuals increases self-regulation of food intake. *Appetite*, 105, 567–574.
<https://doi.org/10.1016/j.appet.2016.06.029>

Document status and date:
Published: 01/10/2016

DOI:
[10.1016/j.appet.2016.06.029](https://doi.org/10.1016/j.appet.2016.06.029)

Document Version:
Accepted author manuscript (Peer reviewed / editorial board version)

Document license:
CC BY-NC-ND

Please check the document version of this publication:

- A submitted manuscript is the version of the article upon submission and before peer-review. There can be important differences between the submitted version and the official published version of record. People interested in the research are advised to contact the author for the final version of the publication, or visit the DOI to the publisher's website.
- The final author version and the galley proof are versions of the publication after peer review.
- The final published version features the final layout of the paper including the volume, issue and page numbers.

[Link to publication](#)

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal.

If the publication is distributed under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license above, please follow below link for the End User Agreement:

www.umlib.nl/taverne-license

Take down policy

If you believe that this document breaches copyright please contact us at:

repository@maastrichtuniversity.nl

providing details and we will investigate your claim.

RUNNING HEAD: WM training in overweight individuals

Taking control: Working memory training in overweight individuals increases self-regulation of food intake

Katrijn Houben, Fania Dassen, & Anita Jansen

Department of Clinical Psychological Science

Faculty of Psychology and Neuroscience; Maastricht University

Correspondence may be sent to: Katrijn Houben, Department of Clinical Psychological

Science, Faculty of Psychology and Neuroscience, Maastricht University, P.O. BOX

616, 6200 MD Maastricht, The Netherlands. Email:

K.Houben@maastrichtuniversity.nl, Phone: +31433881953

Abstract

Working memory (WM) plays a critical role in cognitive control by shielding self-regulatory goals from distraction by desire-related thoughts and emotions. This study examined whether training WM increases self-regulation in overweight participants. It was hypothesized that WM training would decrease psychopathological eating-related thoughts, (over)consumption of food in response to emotions and external cues, food intake and body weight. Overweight participants ($n = 50$) performed 20-25 sessions of WM training or control/sham training. The dependent measures were self-reported eating-related psychopathology, self-reported emotional/external eating behavior, food intake during a bogus taste test, and body weight, assessed before training, immediately following training, and at one-month follow-up. Relative to control, WM training reduced psychopathological eating-related thoughts and emotional eating (but not external eating). These effects were still present at follow-up, one month later. Food intake and body weight did not show an overall effect of training, though WM training did reduce food intake among highly restrained participants. WM training effectively reduced eating-related thoughts, overeating in response to negative emotions, and food intake among participants with strong dietary restraint goals. Hence, these findings indicate that WM training may strengthen self-regulation by shielding dieting goals from distraction by unwanted eating-related thoughts and emotions.

Keywords: Obesity; Working Memory; Training

48 In the last three decades, the prevalence of obesity has nearly doubled (Finucane et
49 al., 2011; Flegal, 2005; Wang & Beydoun, 2007), placing more and more individuals at
50 risk of developing cardiovascular diseases, diabetes, musculoskeletal disorders and
51 cancer (World Health Organization, 2009). In 2010, overweight and obesity were
52 estimated to cause 3,4 million deaths worldwide (Lim et al., 2012). Research further
53 shows that, unabated, the increase in obesity will lead to dramatic falls in future life
54 expectancy (Olshansky et al., 2005). A key contributor to the rapid weight gain that
55 occurred over the past 30 years is our obesogenic environment, which encourages
56 over-consumption of widely-available, inexpensive, energy-dense food and
57 discourages expenditure of energy through physical activity (Hill, Wyatt, Reed, &
58 Peters, 2003). The solution to the obesity problem therefore may lie in identifying
59 feasible ways to cope with the current food-rich environment and adhere to the
60 simple principles of the energy balance equation.

61 One strategy to reduce over-consumption could be to strengthen cognitive
62 control of consumption and body weight. Cognitive control (or executive functioning)
63 is an umbrella term that refers to three basic cognitive functions that allow for goal-
64 directed action amid the endless possibilities afforded to us in real-life situations
65 (Miyake et al., 2000): Maintaining and updating relevant information ('updating'),
66 inhibition of prepotent impulses ('inhibition'), and mental set shifting (task-switching).
67 According to contemporary dual-process models, overconsumption of palatable,
68 energy-dense foods is the result of unintentional, fast-acting impulses that are not or
69 insufficiently regulated via top-down cognitive control (Hofmann, Friese, & Strack,
70 2009; Strack & Deutsch, 2004): Individuals with both strong behavioral impulses to

consume palatable food and low levels of cognitive control are particularly susceptible to over-consumption (Frieze, Hofmann, & Wänke, 2008; Hofmann & Frieze, 2008; Hofmann, Frieze, & Roefs, 2009; Hofmann, Gschwendner, Frieze, Wiers, & Schmitt, 2008; Hofmann, Rauch, & Gawronski, 2007) and weight gain (Nederkoorn, Houben, Hofmann, Roefs, & Jansen, 2010) compared to individuals with effective cognitive control. Hence, individual differences in cognitive control may explain why some people do not succeed in resisting the lure of palatable, energy-dense foods and achieving a healthy weight.

It has been argued that working memory (WM) may very well lie at the heart of successful cognitive control (Engle et al., 1999; Kane et al., 2001). WM is closely connected to the construct of 'updating' and refers to the ability to maintain an active mental representation of (self-regulatory) goals, and shield those goals from distraction (Engle, Tuholski, Laughlin, & Conway, 1999; Kane, Bleckley, Conway, & Engle, 2001; Hofmann, Schmeichel, & Baddeley, 2012). Moreover, this ability to focus on goal-relevant information should also relate to people's ability to regulate their own thoughts and emotions (Hofmann et al., 2012). Indeed, increased WM capacity is related to less thought intrusions and mind-wandering (Brewin & Beaton, 2002; Brewin & Smart, 2005; Kane et al., 2007) and better emotion-regulation (Schmeichel & Demaree, 2010; Schmeichel, Volokhov, & Demaree, 2008). Importantly, overweight and obesity have been associated with reduced cognitive control, including WM (Smith, Hay, Campbell, & Trollor, 2011), which begs the question whether training cognitive control, and WM in particular, may translate into better behavioral self-regulation in overweight and obese individuals.

Previous studies showed that WM can be improved via adaptive training and that such training is effective in reducing clinical symptoms (for reviews see Klingberg 2010; Morrison & Chein, 2011). It is important to note, however, that there has also been criticism on the effectiveness of WM training claiming that there is yet insufficient evidence of its efficacy. Specifically, it has been argued that more scientific evidence is needed to support both near transfer of WM training to untrained WM tasks, and far transfer to WM-related abilities and behavior (e.g., Shipstead, Redick, & Engle, 2012). In the present study, it was examined whether WM training decreases over-consumption and body weight in a sample of overweight participants. Participants either performed WM training or control tasks (sham training) for 25 days. It was expected that WM training would increase self-regulation and cognitive control as evidenced by (1) reduced pathological eating-related thinking, (2) decreased (over-)consumption of food in response to emotions and external cues, (3) reduced consumption of palatable, energy-dense foods, and (4) a decrease in body weight.

Materials and Methods

Participants

Participants were recruited via advertisements in local newspapers about the possibility to participate in research exploring WM training as an intervention for overweight. The advertisements specifically asked for individuals who had overweight and who were motivated to lose weight. Eligibility required that participants were aged 18 to 65, and had a Body Mass Index (BMI) higher than 25 (i.e., A BMI above 25 indicates overweight). Of the 67 participants who responded to the advertisements

and met the eligibility criteria, 62 participants completed the pretest. Twelve participants dropped out after missing too many training sessions¹. The remaining 50 (37 female) participants completed at least 20 training sessions, the pretest, posttest and follow-up. Of the final sample, 6% received primary education, 66% received secondary education, and 28% received higher education. See also Table 1 for participant characteristics. The study was approved by the Ethical Review Committee Psychology and Neuroscience.

Materials & Measures

Working memory training. The WM training (Houben, Wiers, & Jansen, 2011; Klingberg, Forssberg, & Westerberg, 2002) consisted of three tasks: A visuospatial WM task, a backward digit span task, and a letter span task (presented in this order). During the visuospatial WM task, a sequence of squares in a 4x4 grid changed in color on the computer screen. Participants had to reproduce this sequence by clicking the squares that had changed color in the correct order using the computer mouse. During the backward digit span, a sequence of numbers was presented on the computer screen, which participants had to reproduce in reversed order, using either the computer mouse or the keyboard. In the letter span task, a sequence of letters was presented on the computer screen in a circle. One of the positions in this circle was then indicated and participants had to reproduce the corresponding letter using the keyboard. All three tasks consisted of 30 trials (one block).

In the training condition, the difficulty level of all three WM tasks was automatically adjusted on a trial-by-trial basis (cf. Houben et al., 2011; Klingberg et al., 2002): Each task initially started with a sequence of three items. When

participants correctly reproduced this sequence two times in a row, one item was added to the sequence on the next trial. When participants could not correctly reproduce the sequence on two consecutive trials, the sequence in the next trial was reduced by one item. In the control condition, the difficulty level of the WM tasks remained on the initial easy level (three items in a sequence; cf. Houben et al., 2011; Klingberg et al., 2002). Before and after training, WM was measured using the same three tasks, but these assessment tasks ended when participants were unable to reproduce a sequence on two consecutive trials. The outcome measure for each WM task was the amount of items in the sequence that could be correctly reproduced. These three scores were averaged to calculate a total WM score.

Eating Disorder Examination Questionnaire (EDE-Q). The EDE-Q (Fairburn & Beglin, 1994; Fairburn & Cooper, 1993) is a 36-item self-report measure of eating disorder psychopathology. The EDE-Q contains 23 items assessing eating disorder psychopathology over the previous 28 days. These items are answered on a 7-point Likert scale (0 = 'not one day'; 6 = 'every day'). The 23 items together comprise one global score (Cronbach's $\alpha = .90$) as well as four subscales: Restrained eating (Cronbach's $\alpha = .73$), preoccupation with food (Cronbach's $\alpha = .67$), weight concern (Cronbach's $\alpha = .78$) and body shape concern (Cronbach's $\alpha = .88$). Higher scores indicate stronger eating disorder psychopathology.

Dutch Eating Behaviour Questionnaire (DEBQ). The DEBQ (Van Strien, 2005; Van Strien, Frijters, Bergers, & Defares, 1986) is a 33-item self-report measure of emotional eating (Cronbach's $\alpha = .96$), external eating (Cronbach's $\alpha = .77$) and

restrained eating (Cronbach's $\alpha = .90$). All items are scored on a 5-point Likert scale (1 = 'Never'; 5 = 'Very often'). Item examples: 'Do you have a desire to eat when you are irritated?' (emotional eating), 'If foods smells and looks good, do you eat more than usual?' (external eating) and 'Do you try to eat less at mealtimes than you would like to eat?' (dietary restraint). Means are calculated for the three subscales with higher scores indicating increased emotional, external or dietary restraint.

Bogus taste test. Food consumption was measured using a bogus taste test. Participants were presented with four bowls containing different palatable energy-dense foods: salted potato chips (541 kcal/100gr), chocolate cookies (465 kcal/100 gr), milk chocolate (530 kcal/100gr), coated peanuts (535 kcal/100gr). Participants were told that we were interested in their taste perception of a number of food products. Participants were instructed that they were allowed to consume as much or as little of the food as they wished while completing food ratings: Participants first indicated how much they experienced hunger and desire to eat the food on a 100mm Visual Analogue Scale (0 = 'no desire/not hungry'; 100 = 'strong desire/very hungry'). Next, they compared and rated the different food products on a number of taste dimensions. After 15 minutes, the experimenter removed the bowls of food and the amount of food consumed was measured outside the test room. Total amount of calories was calculated as an index of food intake.

Dietary restraint. Dietary restraint was measured using the revised Restraint Scale (RS; Herman & Polivy, 1980). The RS is a self-report questionnaire consisting of 10 items that measure dieting concern/intentions and weight fluctuations

(Cronbach's $\alpha = .70$). Higher scores indicate an increased intention to restrict food intake.

Body Mass Index. Participants' weight and height were assessed in order to calculate participants' Body Mass Index (kg/m^2 ; BMI).

Procedure

After giving consent, participants performed the bogus taste test and the assessment WM tasks. Next, they filled out the Restraint Scale, EDE-Q and DEBQ, and their weight and height were measured. Participants were then randomly assigned to the training or control condition and were informed that they would perform a WM training consisting of 25 sessions via the Internet (so they did not have to come to the lab for the training sessions; participants were sent invitations for each training session via email together with a personalized link to start the session). Participants were given two days to complete a training session. Each session lasted about 30 minutes in total. If participants did not complete a session in time, that session was marked as missed, and participants moved on to the next session. In total, participants could miss up to 5 training sessions. Hence, the total number of training sessions varied between 20 and 25 ($M = 23.02$, $SD = 1.80$; the training sessions were on average completed within 33.92 days, $SD = 8.23$, range: $25 - 66$)². Upon completing the training, the posttest session was scheduled in the lab (on average 9.52 days, $SD = 6.19$, after the last training session). At posttest, participants again performed the bogus taste test and the WM assessment, they filled out the EDE-Q and DEBQ, and their weight was measured. The follow-up session was scheduled one month after the

posttest and included the same measures. Upon completing the experiment, participants received a gift certificate of 50€ as remuneration for their participation.

Design & statistical analyses

Participants were randomly allocated to one of two groups: Active working memory training (n = 24) or control training (n = 26). Randomization checks showed no significant differences between conditions for any potential confounding factors (Table 1). Data were analyzed using mixed-effects ANOVA with condition as between-subjects factor (training versus control) and time as within-subjects factor (pretest, posttest and follow-up)³.

Results

Manipulation Check

In the control condition, the difficulty of the training tasks always remained on the easiest level with only three to-be-remembered items in each task. Consequently, the performance of participants in the control condition remained at the same level over the course of the training period (see Figure 1). In the training condition, in contrast, the training was adjusted adaptively to participants' performance. As can be seen in Figure 1, participants in the training condition showed a steady increase in working memory performance during the training period.

Working memory

A mixed ANOVA on WM task performance showed significant main effects of time, $F(2,96) = 67.31, p < .001, \eta^2_p = .58$, and condition, $F(1,48) = 7.19, p = .01, \eta^2_p = .13$, as

well as a significant time*condition interaction, $F(2,96) = 17.29, p < .001, \eta^2_p = .27$ (see Figure 2). Follow-up analyses per condition showed a significant increase over time in WM performance in both the control condition, $F(2,50) = 17.86, p < .001, \eta^2_p = .42$, and the training condition, $F(2,46) = 57.14, p < .001, \eta^2_p = .71$. Contrasts indicated that, in both the training and control condition, the increase in WM performance from pretest to posttest was significant, $F(1,23) = 98.68, p < .001, \eta^2_p = .81$, and $F(1,25) = 28.20, p < .001, \eta^2_p = .53$, respectively, with a larger increase in the training condition than in the control condition, $F(1,48) = 6.13, p = .02, \eta^2_p = .11$. In addition, both conditions also improved in WM performance from pretest to follow-up, $F(1,23) = 66.63, p < .001, \eta^2_p = .74$, and $F(1,25) = 6.36, p = .02, \eta^2_p = .20$, respectively. At follow-up, WM performance of the training condition was still significantly higher compared to control, $F(1,48) = 23.38, p < .001, \eta^2_p = .33$.

Eating Psychopathology

For the EDE-Q, separate analyses were performed for the global EDE-Q score, and for the four separate subscales: Eating concern, shape concern, weight concern, and dietary restraint. For eating concern, results showed a significant main effect of time, $F(2,96) = 9.54, p < .001, \eta^2_p = .17$ (but no effect of condition, $F(1,48) = .14, p = .71, \eta^2_p < .01$), and a significant time*condition interaction, $F(2,96) = 3.08, p = .05, \eta^2_p = .06$. Follow-up analyses indicated a significant decrease in eating concern over time in the training condition, $F(2,46) = 7.47, p < .01, \eta^2_p = .25$, but not in the control condition, $F(2,50) = 2.47, p = .10, \eta^2_p = .10$ (see Table 2). Contrasts confirmed a significant decrease in the training condition from pretest to posttest, $F(1,23) = 16.26, p < .01$,

261 $\eta^2_p = .41$, that was still significant at one month follow-up, $F(1,23) = 4.27$, $p = .05$, $\eta^2_p =$
262 $.16$.

263 Similarly, analyses for the shape concern subscale also showed a significant
264 time effect, $F(1.77, 84.88) = 10.59$, $p < .001$, $\eta^2_p = .18$ (but not of condition, $F(1,48) =$
265 $.52$, $p = .48$, $\eta^2_p = .01$) that was qualified by a significant time*condition interaction,
266 $F(1.77, 84.88) = 5.17$, $p = .01$, $\eta^2_p = .10$ (see Table 2)⁴. Follow-up analyses indicated a
267 significant decrease in shape concern following training, $F(2,46) = 12.68$, $p < .001$, η^2_p
268 $= .36$, indicating a significant decrease in shape concern from pretest to posttest,
269 $F(1,23) = 19.81$, $p < .001$, $\eta^2_p = .46$, and to follow-up, $F(1,23) = 12.47$, $p < .01$, $\eta^2_p = .35$.
270 In the control condition, there was no significant change in shape concern over time,
271 $F(2,50) = .65$, $p = .53$, $\eta^2_p = .03$.

272 A similar pattern emerged for the weight concern subscale of the EDE-Q (see
273 Table 2), but here the interaction between time and condition did not reach
274 significance, $F(2,96) = 2.02$, $p = .14$, $\eta^2_p = .04$. There was a significant effect of time,
275 $F(2,96) = 5.56$, $p < .01$, $\eta^2_p = .10$, indicating an overall decrease in weight concern. The
276 main effect of condition was not significant, $F(1,48) = .14$, $p = .71$, $\eta^2_p < .01$. For the
277 dietary restraint subscale of the EDE-Q, no effects reached significance (time: $F(1.65,$
278 $79.26) = 2.10$, $p = .13$, $\eta^2_p = .04$; condition: $F(1,48) = .40$, $p = .53$, $\eta^2_p = .01$;
279 time*condition: $F(1.65, 79.26) = .73$, $p = .48$, $\eta^2_p = .02$)⁴.

280 This pattern of results was also illustrated in the analysis of the total EDE-Q
281 score: In addition to a significant main effect of time, $F(1.71, 82.25) = 14.34$, $p < .001$,
282 $\eta^2_p = .23$ (no significant effect of condition, $F(1,48) = .26$, $p = .61$, $\eta^2_p = .01$), the
283 time*condition interaction was significant, $F(1.71, 82.25) = 3.99$, $p = .02$, $\eta^2_p = .08$
284 (see Table 2)⁴, indicating an overall decrease in EDE-Q scores over time in the training

condition, $F(1.50, 34.42) = 16.71, p < .001, \eta^2_p = .42$, that was significant at posttest, $F(1,23) = 23.88, p < .001, \eta^2_p = .51$, as well as at follow-up one month later, $F(1,23) = 15.53, p < .01, \eta^2_p = .40$. No effect of time emerged in the control condition, $F(2,50) = 2.13, p = .13, \eta^2_p = .08$.

Emotional and external eating

Separate analyses were performed for the three subscales of the DEBQ: emotional eating, external eating and dietary restraint. For emotional eating⁴, the significant main effect of time, $F(1.73, 83.19) = 3.60, p = .04, \eta^2_p = .07$, was qualified by a significant time*condition interaction, $F(1.73, 83.19) = 3.63, p = .04, \eta^2_p = .07$ (see Table 2). The main effect of condition was not significant, $F(1,48) < .01, p = .97, \eta^2_p < .01$. Follow-up analyses per condition showed a significant decrease in emotional eating over time in the training condition, $F(2,46) = 5.87, p < .01, \eta^2_p = .20$, but no effect of time in the control condition, $F(2,50) = .81, p = .45, \eta^2_p = .03$. Contrasts confirmed that in the training condition, emotional eating was significantly reduced at posttest, $F(1,23) = 9.35, p < .01, \eta^2_p = .29$, and at follow-up, $F(1,23) = 7.37, p = .01, \eta^2_p = .24$, compared to pretest. For external eating only the main effect of time reached significance, $F(2,96) = 7.86, p < .01, \eta^2_p = .14$, indicating a decrease in external eating across both conditions (condition: $F(1,48) = .01, p = .93, \eta^2_p < .01$; time*condition: $F(2,96) = .27, p = .77, \eta^2_p = .01$). For dietary restraint there were no significant effects (time, $F(1.70, 81.43) = 2.63, p = .09, \eta^2_p = .05$; condition, $F(1,48) = .02, p = .89, \eta^2_p < .01$; time*condition, $F(1.70, 81.43) = .66, p = .49, \eta^2_p = .01$)⁴.

309 Food intake

310 Results showed no differences between the training and control condition with
311 respect to hunger (condition: $F(1, 48) = .04, p = .84, \eta^2_p = .00$; time: $F(2, 96) = 1.42, p =$
312 $.25, \eta^2_p = .03$; condition*time: $F(2, 96) = .74, p = .48, \eta^2_p = .02$) or craving (condition:
313 $F(1, 48) = .01, p = .91, \eta^2_p = .00$; time: $F(2, 96) = 61, p = .54, \eta^2_p = .01$; condition*time:
314 $F(2, 96) = 1.06, p = .35, \eta^2_p = .02$). Food intake during the bogus taste test showed no
315 significant time*condition interaction, $F(1.62, 77.52) = .38, p = .64, \eta^2_p = .01$, nor
316 significant main effects (time, $F(1.62, 77.52) = 2.04, p = .15, \eta^2_p = .04$; condition,
317 $F(1, 48) = .13, p = .72, \eta^2_p < .01$)⁴.

318 It is important to note here that research in other areas of cognitive control
319 training (i.e., inhibitory control training) indicates a moderating role of dietary
320 restraint on training effects with larger effects on food intake in participants with high
321 levels of dietary restraint (e.g., Houben & Jansen, 2011; see also Jones et al., 2016).
322 Theoretically, stronger cognitive control should result in greater concordance
323 between one's current goals and behavior (Hofmann et al., 2009). Hence, it is unlikely
324 that increasing cognitive control will lead to behavior change when there is no
325 concurrent dieting goal present. Similarly, WM is critical for sustained attention to
326 one's goals (Engle et al., 1999; Kane et al., 2001; Hofmann et al., 2012). Training WM
327 should therefore translate into lower food intake especially for participants with
328 strong dieting goals but not for participants without active dieting goals.

329 We therefore also a post-hoc analysis to examine the effect of baseline dietary
330 restraint (measured with the RS), condition and their interaction on food intake at
331 pretest, posttest and at follow-up. Restraint scores were z-standardized and entered
332 as a continuous predictor in the ANOVA (Aiken & West, 1991). One influential outlier

(Cook's > 1.5) was excluded from the analyses. For food intake at pretest, none of the effects reached significance (restraint: $F(1,45) = .48, p = .49, \eta^2_p = .01$; condition: $F(1,45) = 2.58, p = .12, \eta^2_p = .05$; condition*restraint: $F(1,45) = .23, p = .63, \eta^2_p = .01$). At posttest, the expected interaction between dietary restraint and condition was significant, $F(1,45) = 3.91, p = .05, \eta^2_p = .08$. The main effects of restraint and condition did not reach statistical significance (restraint: $F(1,45) = .04, p = .83, \eta^2_p < .01$; condition: $F(1,45) = .28, p = .60, \eta^2_p = .01$). We analyzed the effect of training separately for participants scoring high and low on the RS (respectively 1 SD above and 1 SD below the mean score; Cohen, Cohen, West, & Aiken, 2003). For participants with lower dietary restraint (-1 SD), there was no significant difference between training and control with respect to food intake, $F(1,45) = 1.26, p = .27, \eta^2_p = .03$, but among high restrained participants (+1 SD), food intake was reduced in the training condition compared to control, $F(1,45) = 3.49, p = .07, \eta^2_p = .07$ (see Figure 3). Consumption at follow-up showed a similar pattern of results, though the interaction between dietary restraint and condition was not significant, $F(1,45) = 2.11, p = .15, \eta^2_p = .05$ (restraint: $F(1,45) = .43, p = .52, \eta^2_p = .01$; condition: $F(1,45) = .61, p = .44, \eta^2_p = .01$).

BMI

None of the effects of time or condition were significant for BMI (time: $F(1.50, 71.87) = 1.00, p = .35, \eta^2_p = .02$; condition: $F(1,48) = .11, p = .74, \eta^2_p < .01$; time*condition: $F(1.50, 71.87) = .09, p = .86, \eta^2_p < .01$)⁴, indicating no significant change in body weight over time.

Discussion

Given the prominent role of cognitive control in self-regulatory behavior like food intake, the aim of the present study was to examine whether boosting WM via training would lead to better self-regulation in overweight individuals. As expected, participants who received WM training, relative to participants in the control condition, reported overall less eating-related concerns (especially less concern about eating and shape), and less emotional eating (but not external eating), immediately following training and at one-month follow-up. Food intake and body weight did not show an overall effect of training. Results, however, did show the expected effect of WM training on food intake in highly restrained participants, indicating that WM training increased correspondence between dietary goals and food intake.

WM supports self-regulation by enabling individuals to resist the attentional capture of tempting stimuli at early stages of information processing, thereby shielding self-regulatory goals from competing goals and distraction (Kane et al., 2001; Hofmann et al., 2012). As such, WM relates to the ability to regulate one's own thoughts and emotions, by focusing attention on goal-relevant information and ignoring irrelevant, distracting information (Hofmann et al., 2012). In line with this idea, WM training reduced pathological ruminative thoughts about food, weight, and body shape. This finding fits with previous research showing an association between preoccupying cognitions and WM impairment in dieters: Preoccupying thoughts about food, weight and body shape seem to consume WM resources with detrimental effects on WM performance (Green, Elliman, & Rogers, 1997; Kemps & Tiggemann, 2005; Kemps, Tiggemann, & Marshall, 2005; Vreugdenburg, Bryan, & Kemps, 2003). The present results add to these findings by demonstrating that WM

training alleviates distraction by preoccupying cognitions related to dieting, weight, food, and body shape.

Further, WM training decreased self-reported emotional eating indicating that participants who received WM training were better able to regulate their emotions in other ways than by (over)eating compared to participants in the control condition. This finding is consistent with previous studies showing that individuals with higher WM capacity, as opposed to individuals with lower levels of WM, are better able to regulate emotions and appraise emotional stimuli in an unemotional manner (Schmeichel & Demaree, 2010; Schmeichel, Volokhov, & Demaree, 2008). It was also expected that WM training would increase the resilience to temptation by food cues. Unexpectedly, self-reported external eating was reduced over time in both conditions. Previous research, however, has indicated that the external eating subscale of the DEB-Q (but not the emotional eating subscale or the dietary restraint subscale) is affected by visceral states and may thus be a state rather than a trait measure (Evers et al., 2011). It is possible that such fluctuations in visceral states over time have caused this slight, albeit significant, decrease in external eating across conditions.

These findings thus indicate that WM training might help overweight and obese individuals to create a more healthy style of thinking about their body and eating behavior. Nevertheless, the present findings did not show the expected effects of WM training on body weight, and effects on food intake were only found among highly restrained eaters. Specifically, highly restrained eaters who received WM training showed a reduction in food intake relative to participants in the control condition. Perhaps this finding is not surprising given that high WM capacity increases

the correspondence between dieting goals and eating behavior (Hofmann et al., 2007). Without (dieting) motivation, it is unlikely that cognitive control training will lead to behavioral change. Thus, the strongest effects on food intake are to be expected for participants who hold strong dietary restraint standards. It is interesting to note that training studies which have targeted a different cognitive control ability, namely response inhibition, have also shown stronger effects of inhibition training on food intake among highly restrained eaters (Houben & Jansen, 2011; Veling, Aarts, & Papies, 2011; see also Jones et al., 2016), indicating that cognitive control training may indeed be more effective for participants with high levels of dietary restraint.

While WM training did not influence body weight in the present study, it should be noted that inhibition training has been shown to reduce both food intake (e.g., Houben & Jansen, 2011; 2015; Veling et al., 2011; Lawrence, Verbruggen, Morrison, Adams, & Chambers, 2015; Veling, Aarts, & Stroebe, 2013) and body weight (e.g., Lawrence et al., 2015; Veling, van Koningsbruggen, Aarts, & Stroebe, 2014). As such, inhibition training effects appear to be stronger and more robust compared to effects of WM training. Perhaps this is due to differences in terms of the behavioral-specificity of the training. Inhibition training has been shown to be effective only when the training is focused on strengthening inhibition over food-related responses, but not when general response inhibition is targeted during training (Allom, Mullan, & Hagger, 2016). It might therefore be interesting for future research to contrast the present findings for general WM training with more applied, diet-relevant WM training.

A limitation to the present findings is that we did not measure dieting motivations, and as explained above, it is unlikely that WM training will translate into

weight loss when participants are unmotivated to lose weight. Future research should therefore screen participants for dieting motivations and test whether WM training might be more effective among overweight participants who are committed to losing weight. A second limitation is that we did not measure beliefs regarding the training in the two conditions. While both conditions received the same instructions, we cannot rule out that participants in the control condition may have become suspicious and did not believe that they were receiving WM training. It is therefore important to include measures of expectancies and beliefs regarding the training and the purpose of the study in future studies to rule out demand artefacts. Another limitation to this study concerns the fact that we did not measure transfer effects of the WM training to other non-trained tasks of executive functions (WM, task-switching, inhibition). Earlier research has shown transfer effects to non-trained tasks (see Klingberg, 2010 for a review), though the generalization to non-trained tasks has also raised considerable debate (e.g., Shipstead et al., 2012). Further, it is possible that WM training in isolation is not effective as a weight loss intervention and will only be effective in reducing weight in combination with additional (lifestyle) interventions. In this way, overweight individuals who are highly motivated to diet and who are provided with dieting strategies might profit the most from WM training that boosts self-regulatory abilities. Future research should thus further examine the effectiveness of WM training on weight in combination with other weight loss interventions.

In conclusion, WM training successfully reduced emotional eating and psychopathological eating-related concerns in a sample of overweight participants. Moreover, WM training also reduced food intake, but only among highly restrained

453 eaters, underscoring the need to further examine the effectiveness of WM training in
454 target groups of overweight individuals who are highly motivated to lose weight.
455

Footnotes

1. Of the 12 participants who dropped out, 9 participants were in the training condition and 3 participants were in the control condition. The participants who dropped out did not differ from the participants who finished the study in terms of age, or scores on WM, DEBQ, EDE-Q and RS (all $F < 1$). Participants who dropped out, however, did have a lower BMI ($M = 28.84$, $SD = 2.58$, $F(1, 60) = 5.72$, $p = .02$, compared to the rest of the sample ($M = 31.56$, $SD = 3.72$).
2. Note that the range normally should have been 25 - 50 days because participants were only allowed to do one session per day and had to complete a session every two days. However, two participants were given some extension to these rules due to personal issues.
3. We also performed an Intention To Treat (ITT) analysis on all dependent variables using the "last observation carried forward method" method. In the ITT analyses, all participants were included, rather than including only the participants who completed the study as in the Per Protocol analyses. Including all participants in the ITT analyses did not change any of the effects compared to the Per Protocol analyses.
4. Due to violation of the sphericity assumption, degrees of freedom were Greenhouse-Geisser adjusted.

476

Acknowledgements

477 This study was funded by Maastricht University. The authors declare that they have
478 no conflicts of interest.

479

References

- Allom, V., Mullan, B., & Hagger, M. (2016). Does inhibitory control training improve health behaviour? A meta-analysis. *Health Psychology Review*, 10, 168-186.
- Aiken, L. S., & West, S. G. (1991). Multiple regression: Testing and interpreting interactions. Newbury Park: Sage.
- Borella, E., Carretti, B., Riboldi, F., & De Beni, R. (2010). Working memory training in older adults: evidence of transfer and maintenance effects. *Psychology and Aging* 25, 767-778.
- Brewin, C. R., & Beaton, A. (2002). Thought suppression, intelligence, and working memory capacity. *Behaviour Research and Therapy*, 40, 923-930.
- Brewin, C. R., & Smart, L. (2005). Working memory capacity and suppression of intrusive thoughts. *Journal of Behavior Therapy and Experimental Psychiatry*, 36, 61-68.
- Cohen, J., Cohen, P., West, S. G., & Aiken, L. S. (2003). Applied multiple regression/correlation analysis for the behavioral sciences. New Jersey: Lawrence Erlbaum Associates.
- Engle, R. W., Tuholski, S. W., Laughlin, J.E., & Conway, A. R. A. (1999). Working memory capacity, short-term memory, and general fluid intelligence: A latent-variable approach. *Journal of Experimental Psychology: General*, 128, 309-331.
- Evers, C., Stok, F. M., Danner, U. N., Salmon, S. J., de Ridder, D. T. D., & Adriaanse, M. A. (2011). The shaping role of hunger on self-reported external eating status. *Appetite*, 57, 318-320.
- Fairburn, C. G., & Beglin, S. J. (1994). The assessment of eating disorders: interview or self-report questionnaire? *International Journal of Eating Disorders*, 16, 363-370.

504 Fairburn, C. G., & Cooper, Z. (1993). The Eating Disorder Examination (twelfth
505 edition). In C. G. Fairburn, & G. T. Wilson (Eds), Binge Eating: Nature, Assessment
506 and Treatment (pp. 317- 360). New York: Guilford Press.

507 Finucane, M. M., Stevens, G. A., Cowan, M. J., Danaei, G., Lin, J. K., Paciorek, C. J., ...
508 Ezzati, M. (2011). National, regional, and global trends in body-mass index since
509 1980: systematic analysis of health examination surveys and epidemiological
510 studies with 960 country-years and 9· 1 million participants. *Lancet*, 377, 557-
511 567.

512 Flegal, K. M. (2005). Epidemiologic aspects of overweight and obesity in the United
513 States. *Physiology & Behavior*, 86, 599-602.

514 Friese, M., Hofmann, W., & Wänke, M. (2008). When impulses take over. Moderated
515 predictive validity of explicit and implicit attitude measures in predicting food
516 choice and consumption behaviour. *British Journal of Social Psychology*, 47, 397-
517 419.

518 Green, M. W., Elliman, N. A., & Rogers, P. J. (1997). Impaired cognitive processing in
519 dieters: Failure of attention focus or resource capacity limitation ? *British Journal*
520 *of Health Psychology*, 2, 259-267.

521 Herman, C. P., & Polivy, J. P. (1980). Restrained eating. In A. J. Stunkard (Ed.), *Obesity*
522 (pp. 208-225). Philadelphia: Saunders.

523 Hill, J. O., Wyatt, H. R., Reed, G. W., & Peters, J. C. (2003). Obesity and the
524 environment: where do we go from here? *Science*, 299, 853-855.

525 Hofmann, W., Friese, M., & Roefs, A. (2009). Three ways to resist temptation. The
526 independent contributions of executive attention, inhibitory control and affect

527 regulation to the impulse control of eating behavior. *Journal of Experimental*
528 *Social Psychology*, 45, 431-435.

529 Hofmann, W., Friese, M., & Strack, F. (2009). Impulse and self-control from a dual-
530 systems perspective. *Perspectives on Psychological Science*, 4, 162-176.

531 Hofmann, W., & Friese, M. (2008). Impulses got the better of me. Alcohol moderates
532 the influence of implicit attitudes toward food cues on eating behavior. *Journal of*
533 *Abnormal Psychology*, 117, 420-427.

534 Hofmann, W., Gschwendner, T., Friese, M., Wiers, R. W., & Schmitt, M. (2008). Working
535 memory capacity and self-regulatory behavior: toward an individual differences
536 perspective on behavior determination by automatic versus controlled processes.
537 *Journal of Personality and Social Psychology*, 95, 962-977.

538 Hofmann, W., Rauch, W., & Gawronski, B. (2007). And deplete us not into temptation.
539 Automatic attitudes, dietary restraint, and self-regulatory resources as
540 determinants of eating behavior. *Journal of Experimental Social Psychology*, 43,
541 497-504.

542 Hofmann, W., Schmeichel, B. J., & Baddeley, A.D. (2012). Executive functions and self-
543 regulation. *Trends in Cognitive Sciences*, 16, 174-180.

544 Houben, K., & Jansen, A. (2015). Chocolate equals stop: Chocolate-specific inhibition
545 training reduces chocolate intake and go associations with chocolate. *Appetite*,
546 87, 318-323.

547 Houben, K., & Jansen, A. (2011). Training inhibitory control: A recipe for resisting
548 sweet temptations. *Appetite*, 56, 345-349.

549 Houben, K., Wiers, R. W., & Jansen, A. (2011). Getting a grip on drinking behavior:
 550 Training working memory to reduce alcohol abuse. *Psychological Science*, 22,
 551 968- 975.

552 Jones, A., Di Lemma, L. C. G., Robinson, E., Christiansen, P., Nolan, S., Tudur-Smith, C.,
 553 & Field, M. (2016). Inhibitory control training for appetitive behaviour change: A
 554 meta-analytic investigation of mechanisms of action and moderators of
 555 effectiveness. *Appetite*, 97, 16-28.

556 Kane, M. J., Bleckley, M. K., Conway, A. R. A., & Engle, R. W. (2001). A controlled-
 557 attention view of working-memory capacity. *Journal of Experimental Psychology:*
 558 *General*, 130, 169-183.

559 Kane, M. J., Brown, L. H., McVay, J. C., Silvia, P. J., Myin-Germeys, J., & Kwapil, T. R.
 560 (2007). For whom the mind wanders, and when: An experience-sampling study of
 561 working memory and executive control in daily life. *Psychological Science*, 18,
 562 614-621.

563 Kemps, E., Tiggemann, M., & Marshall, K. (2005). Relationship between dieting to lose
 564 weight and the functioning of the central executive. *Appetite*, 45, 287-294.

565 Kemps, E., & Tiggemann, M. (2005). Working memory performance and
 566 preoccupying thoughts in female dieters: Evidence for a central executive
 567 impairment. *British Journal of Clinical Psychology*, 44, 357-366.

568 Klingberg, T., Forssberg, H., & Westerberg, H. (2002). Training of working memory in
 569 children with ADHD. *Journal of Clinical Experimental Neuropsychology*, 24, 781-
 570 791.

571 Klingberg, T. (2010). Training and plasticity of working memory. *Trends in Cognitive*
 572 *Sciences*, 14, 317-324.

573 Lawrence, N. S., O'Sullivan, J., Parslowa, D., Javaid, M., Adams, R. C., Chambers, C. D.,
 574 Kos, K. & Verbruggen, F. (2015). Training response inhibition to food is associated
 575 with weight loss and reduced energy intake. *Appetite*, 95, 17-28.
 576 Lawrence, N. S., Verbruggen, F., Morrison, S., Adams, R. C., & Chambers, C. D. (2015).
 577 Stopping to food can reduce intake. Effects of stimulus-specificity and individual
 578 differences in dietary restraint. *Appetite*, 85, 91-103.
 579 Lim, S. S., Vos, T., Flaxman, A. D., Danaei, G., Shibuya, K., Adair-Rohani, H., ... Ezzati,
 580 M. (2012). A comparative risk assessment of burden of disease and injury
 581 attributable to 67 risk factors and risk factor clusters in 21 regions, 1990–2010: a
 582 systematic analysis for the Global Burden of Disease Study 2010. *Lancet*, 380,
 583 2224-2260.
 584 Miyake, A., Friedman, N. P., Emerson, M. J., Witzki, A. H., Howerter, A., & Wager T.
 585 (2000). The unity and diversity of executive functions and their contributions to
 586 complex “frontal lobe” tasks: A latent variable analysis. *Cognitive Psychology*, 41,
 587 49-100.
 588 Morrison, A. B., & Chein, J. M. (2011). Does working memory training work? The
 589 promise and challenges of enhancing cognition by training working memory.
 590 *Psychonomic Bulletin & Review*, 18, 46-60.
 591 Nederkoorn, C., Houben, K., Hofmann, W., Roefs, A., & Jansen, A. (2010). Control
 592 yourself or just eat what you like?. Weight gain over a year is predicted by an
 593 interactive effect of response inhibition and a preference for high fat foods.
 594 *Health Psychology*, 29, 389-393.

595 Olshansky, S. J., Passaro, D. J., Hershow, R. C., Layden, J., Carnes, B. A., Brody, J., ...
 596 Ludwig, D. S. (2005). A potential decline in life expectancy in the United States in
 597 the 21st century. *The New England Journal of Medicine*, 352, 1138-1145.
 598 Schmeichel, B. J., & Demaree, H. A. (2010). Working Memory Capacity and
 599 Spontaneous Emotion Regulation: High Capacity Predicts Self-Enhancement in
 600 Response to Negative Feedback. *Emotion*, 10, 739-744.
 601 Schmeichel, B. J., Volokhov, R. N., & Demaree, H. A. (2008). Working memory capacity
 602 and the self-regulation of emotional expression and experience. *Journal of*
 603 *Personality and Social Psychology*, 95, 1526-1540.
 604 Shipstead, Z., Redick, T. S., & Engle, R. W. (2012). Is working memory training
 605 effective? *Psychological Bulletin*, 138, 628-654.
 606 Smith, E., Hay, P., Campbell, L., & Trollor, J. N. (2011). A review of the association
 607 between obesity and cognitive function across the lifespan: implications for novel
 608 approaches to prevention and treatment. *Obesity Reviews*, 12, 740-755.
 609 Strack, F., & Deutsch, R. (2008). Reflective and impulsive determinants of social
 610 behavior. *Personality and Social Psychology Review*, 8, 220-247.
 611 Van Strien, T. (2005). *Nederlandse vragenlijst voor eetgedrag [Dutch Eating Behaviour*
 612 *Questionnaire, manual]*. Amsterdam: Boom Test Publishers
 613 Van Strien, T., Frijters, J. E., Bergers, G. P. A., & Defares, P. B. (1986). The Dutch Eating
 614 Behaviour Questionnaire (DEBQ) for assessment of restrained, emotional and
 615 external eating behaviour. *International Journal of Eating Disorders*, 5, 295-315.
 616 Veling, H., Aarts, H., & Papies, E. K. (2011). Using stop signals to inhibit chronic
 617 dieters' responses toward palatable foods. *Behaviour Research and Therapy*, 49,
 618 771-780.

619 Veling, H., Aarts, H., & Stroebe, W. (2013). Using stop signals to reduce impulsive
620 choices for palatable unhealthy foods. *British Journal of Health Psychology*, 18,
621 354-368.

622 Veling, H., van Koningsbruggen, G. M., Aarts, H., & Stroebe, W. (2014). Targeting
623 impulsive processes of eating behavior via the internet. Effects on body weight.
624 *Appetite*, 78, 102-109.

625 Vreugdenburg, L., Bryan, J., & Kemps, E. (2003). The effect of self-initiated weight-loss
626 dieting on working memory: The role of preoccupying cognitions. *Appetite*, 41,
627 291-300.

628 Wang, Y., & Beydoun, M. A. (2007). The obesity epidemic in the United States—
629 gender, age, socioeconomic, racial/ethnic, and geographic characteristics: a
630 systematic review and meta-regression analysis. *Epidemiologic Reviews*, 29, 6-28.

631 World Health Organization (WHO) (2009). *Global health risks: mortality and burden of*
632 *disease attributable to selected major risks*. Geneva, Switzerland: World Health
633 Organization.

634

Table 1 Participant characteristics per training condition. Means and standard deviations for baseline characteristics are provided per condition

	Training (N = 24)	Control (N = 26)	Range	F/ χ^2	p
Age	36.08 (11.28)	37.62 (10.65)	18 - 62	.24	.62
Baseline BMI (kg/m ²)	31.76 (3.79)	31.38 (3.72)	26.50 - 41.58	.13	.72
Gender (% female)	79.2%	69.2%	-	.64	.42
Dietary restraint (RS)	17.50 (3.71)	16.69 (5.19)	3 - 27	.40	.53
Working Memory (WM) - baseline	6.15 (1.54)	6.06 (1.32)	3.00 - 9.67	.05	.83

Table 2 Means and standard deviations for the dependent measures at pretest, posttest and one-month follow-up, per condition

	Training (N = 24)			Control (N = 26)		
	Pretest	Posttest	Follow-up	Pretest	Posttest	Follow-up
DEBQ emotional eating	2.90 (.86)	2.66 (.86)*	2.63 (.98)*	2.73 (1.04)	2.80 (1.01)	2.70 (1.18)
DEBQ external eating	3.07 (.54)	2.97 (.51)	2.87 (.59)	3.10 (.63)	2.94 (.51)	2.91 (.64)
DEBQ dietary restraint	3.08 (.82)	3.07 (.66)	2.93 (.63)	3.08 (.78)	2.96 (.75)	2.95 (.89)
EDE-Q eating concern	1.10 (.89)	.57 (.56)*	.74 (.75)*	.96 (.82)	.84 (.78)	.79 (.65)
EDE-Q shape concern	3.04 (1.52)	2.14 (1.28)*	2.23 (1.39)*	2.31 (1.33)	2.18 (1.30)	2.14 (1.44)
EDE-Q weight concern	2.65 (1.37)	2.10 (1.26)	2.02 (1.38)	2.22 (1.42)	2.07 (1.31)	2.08 (1.42)
EDE-Q dietary restraint	2.00 (1.41)	1.70 (.98)	1.80 (1.17)	1.84 (1.05)	1.71 (1.03)	1.45 (1.09)
EDE-Q global	2.19 (1.01)	1.63 (.78)*	1.70 (.89)*	1.83 (.98)	1.70 (.88)	1.61 (.97)
Food intake (kcal)	243.72 (203.60)	309.67 (289.46)	296.83 (233.39)	266.83 (138.19)	299.37 (174.61)	345.40 (348.40)
BMI (kg/m ²)	31.76 (3.79)	31.62 (3.76)	31.63 (3.84)	31.38 (3.72)	31.31 (3.94)	31.25 (3.96)

Note. * = Significantly different from pretest at $p < .05$.

Figure 1 The number of items that could be correctly recalled in a WM sequence at the end of each training/control session, averaged across the three training tasks (working memory span), separately for the training condition and the control condition.

Figure 2 Means and standard errors for WM performance at pretest, posttest and one-month follow-up, per condition.

Note: * indicates significant differences at $p < .05$

Figure 3 Estimated marginal means (with standard errors) for caloric intake at pretest, posttest and one-month follow-up, per condition. Means are shown separately for low restrained versus highly restrained eaters (respectively 1 SD below or above the mean restraint score).

Note: * indicates significant differences at $p < .05$