One-dimensional bargaining with a general voting rule

Citation for published version (APA):

Predtetchinski, A. (2007). *One-dimensional bargaining with a general voting rule*. METEOR, Maastricht University School of Business and Economics. METEOR Research Memorandum No. 045
<https://doi.org/10.26481/umamet.2007045>

Document status and date:

Published: 01/01/2007

DOI:

[10.26481/umamet.2007045](https://doi.org/10.26481/umamet.2007045)

Document Version:

Publisher's PDF, also known as Version of record

Please check the document version of this publication:

- A submitted manuscript is the version of the article upon submission and before peer-review. There can be important differences between the submitted version and the official published version of record. People interested in the research are advised to contact the author for the final version of the publication, or visit the DOI to the publisher's website.
- The final author version and the galley proof are versions of the publication after peer review.
- The final published version features the final layout of the paper including the volume, issue and page numbers.

[Link to publication](#)

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal.

If the publication is distributed under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license above, please follow below link for the End User Agreement:

www.umlib.nl/taverne-license

Take down policy

If you believe that this document breaches copyright please contact us at:

repository@maastrichtuniversity.nl

providing details and we will investigate your claim.

Arkadi Predtetchinski

One-dimensional bargaining with a general
voting rule

RM/07/045

JEL code: C78

Maastricht research school of **E**conomics
of **T**echnology and **O**rganizations

Universiteit Maastricht
Faculty of Economics and Business Administration
P.O. Box 616
NL - 6200 MD Maastricht

phone : ++31 43 388 3830
fax : ++31 43 388 4873

One-dimensional bargaining with a general voting rule^{*}

Arkadi Predtetchinski[†]

October 22, 2007

Abstract

We study a model of multilateral bargaining over social outcomes represented by points in the unit interval. An acceptance or rejection of a proposal is determined by a voting rule as represented by a collection of decisive coalitions. The focus of the paper is on the asymptotic behavior of subgame perfect equilibria in stationary strategies as the discount factor goes to one. We show that, along any sequence of stationary subgame perfect equilibria, as the discount factor goes to one, the social acceptance set collapses to a point. This point, called the bargaining outcome, is independent of the sequence of equilibria and is uniquely determined by the set of players, the utility functions, the recognition probabilities, and the voting rule. The central result of the paper is a characterization of the bargaining outcome as a unique zero of the characteristic equation.

JEL classification code: C78

Keywords: Bargaining, subgame perfect equilibrium, voting rules.

^{*}The author is most thankful to Jean-Jacques Herings and the participants of the Mathematical Economics Seminar in the Sobolev Institute of Mathematics (Novosibirsk, Russia).

[†]Department of Economics, Maastricht University, P.O. Box 616, 6200 MD Maastricht, The Netherlands. E-mail: A.Predtetchinski@algec.unimaas.nl

1 Introduction

This paper analyzes a model of multilateral bargaining where players must choose one alternative from a set of alternatives represented by points in the unit interval. An alternative might be a level of taxation, a location of a facility, or an index of an ideological content of a policy (left vs. right).

Bargaining proceeds as follows. At the beginning of each period, nature randomly selects one of the players as a proposer. The probability for a player to become a proposer, the so-called recognition probability, is assumed to be the same in each period. The player chosen by nature puts forward a proposal that specifies one alternative. All players then react to the proposal. Each player can either reject or accept the proposal. The votes are cast sequentially, the sequence of responses being fixed throughout the game. Whether the proposal passes or fails is then determined by a voting rule, as represented by a collection of decisive coalitions. The passing of a proposal requires an approval of it by all the players in some decisive coalition. Examples of voting rules include the unanimity acceptance rule when a passing of a proposal requires an approval of it by all the players, the quota rule, when a fixed number of votes is needed for a passing of a proposal, or the simple majority rule. If the proposal passes, it is implemented and the game ends. In this case each player receives a discounted utility of the alternative. Otherwise, a new period begins.

We consider subgame perfect equilibria in stationary strategies. Stationarity means that a proposal of any player does not depend on the history of play and a player's reaction to a proposal depends only on the proposal itself. The focus of the paper is on the asymptotic behavior of stationary subgame perfect equilibria as the discount factor approaches one.

The results are as follows. We prove that, along any sequence of subgame perfect equilibria in stationary strategies the social acceptance set collapses to a point. This point, called the bargaining outcome, is independent of the sequence of equilibria and is uniquely determined by the set of players, the utility functions, the recognition probabilities, and the voting rule. The central result of the paper is a characterization of the bargaining outcome as a unique zero of the characteristic equation. This paper is the first to provide the characterization of the limit of stationary equilibrium in a one-dimensional model of bargaining.

The results are obtained under rather minimal assumptions. Thus the instantaneous utility functions are only assumed to be single-peaked and concave. Furthermore, we require that the intersection of any two decisive coalitions contain a player with a positive recognition probability. This requirement puts but a very mild restriction on the voting rule and the recognition probabilities.

Games of bargaining with a one-dimensional space of alternatives have been previously considered in Banks and Duggan [1], Cho and Duggan [4], Kalandrakis [7], and Cardona and Ponsati [3]. Banks and Duggan [1] and Kalandrakis [7] consider bargaining in a situation where the alternatives are represented by points in a general compact convex set. For the special case where the set of alternatives is one-dimensional Banks and Duggan [1] establish the existence of stationary subgame perfect equilibria in pure strategies, while Kalandrakis [7] shows that pure strategy stationary subgame perfect equilibria are locally

unique and finite in number for almost all discount factors. Cho and Duggan [4] prove uniqueness of stationary subgame perfect equilibria in pure strategies in a model of one-dimensional bargaining under the assumption that the utility functions are quadratic. They also establish that any pair of stationary subgame perfect equilibria are nested in the sense that the social acceptance set in one equilibrium forms the subset of the social acceptance set of the other equilibrium.

The game tree used in this paper is the same as in Banks and Duggan [1], Cho and Duggan [4] and Kalandrakis [7] except that the players vote sequentially, not simultaneously. This difference turns out to be inessential, and the results of Banks and Duggan [1] and Cho and Duggan [4] are true in the setting of the present paper. The results of Kalandrakis [7] on local uniqueness and finiteness equilibria, on the other hand, are generic in the discount factors that may be different across the players. We restrict our attention to the case of a common discount factor.

Cardona and Ponsati [3] also study bargaining over a one-dimensional set of alternatives but their model differs from ours in two respects. Firstly, in Cardona and Ponsati [3] the identity of the proposer rotates deterministically in the player set, whereas we use time-independent recognition probabilities. And, secondly, Cardona and Ponsati [3] restrict attention to a quota voting rule, whereas we use a more general family of voting rules. Cardona and Ponsati [3] prove, in their setting, the asymptotic uniqueness of subgame perfect equilibria in stationary strategies, but do not provide the characterization.

Our paper complements these insights by providing the asymptotic uniqueness result, and more importantly, the explicit computation of the limit of subgame perfect equilibria in stationary strategies.

There are numerous other contributions that study stationary subgame perfect equilibria in various games of multilateral bargaining under the assumption that the the boundary of the set of feasible payoffs is of dimension $n - 1$, where n is the number of players. This assumption is satisfied in a situation when n players have to divide an amount of money amongst themselves, in which case the set of alternatives consists of all feasible redistributions of an amount among n players. Thus Merlo and Wilson [9] and Merlo and Wilson [10] give sufficient conditions for the uniqueness of stationary subgame perfect equilibria (in pure strategies) in a model where the identity of the proposal and the amount of money to be divided follow a Markov process and the unanimous approval is needed for a proposal to pass. Eraslan [5] establishes uniqueness of subgame perfect equilibria (in mixed strategies) in a game with a tree similar to the one described above and a quota voting rule. Eraslan and Merlo [6] characterize stationary subgame perfect equilibria (in mixed strategies) in a model where the amount of money to be divided is stochastic and a general agreement rule is used. It is clear that the game where the payoffs are generated by a one-dimensional space of alternatives is not covered by this analysis.

The idea of studying the asymptotic behavior of subgame perfect equilibria in the bargaining context goes back to Binmore et al [2] who prove that in a two-player game, the limit of subgame perfect equilibrium is given by the Nash bargaining solution. Miyakawa [11] generalizes the result of Binmore et al [2]. He obtains a characterization of the limit of stationary subgame perfect equilibria in the n -player game with the unanimity acceptance

rule as the probability of the breakdown of the negotiations goes to zero, under the assumption that the boundary of the set of feasible utility vectors is an $(n - 1)$ -dimensional surface. Miyakawa [11] shows that the limit is the asymmetric Nash bargaining solution, with the weights corresponding to the recognition probabilities. A related result is given in Kultti and Vartiainen [8], who also assume that the boundary of the feasible payoffs is $(n - 1)$ -dimensional surface. They obtain the (asymmetric) Nash bargaining solution as a limit of the Von Neumann-Morgenstern stable set as the discount factor vanishes. The stable set is defined with respect to a dominance relation where an alternative x dominates an alternative y if some player prefers x even with a one-period delay. Our paper provides the asymptotic results for the stationary subgame perfect equilibrium in a setting that has not been covered by the analysis in Binmore et al [2], Miyakawa [11] or Kultti and Vartiainen [8].

The outline of the paper is as follows. Section 2 introduces the game of bargaining and the concept of δ -equilibrium. Essentially, δ -equilibrium is a subgame perfect equilibrium in stationary strategies where each player is assumed to vote sincerely. Section 3 examines a relationship of the δ -equilibrium and the subgame perfect equilibrium. It is shown that each δ -equilibrium is a subgame perfect equilibrium in stationary strategies. Conversely, given a subgame perfect equilibrium in stationary strategies there exists a δ -equilibrium having the same equilibrium proposals and inducing the same equilibrium payoffs. In this sense, the set of δ -equilibria is effectively the same as the set of subgame perfect equilibria in stationary strategies.

In Section 4 we establish that, along any sequence of δ -equilibria, as the discount factor converges to one, the social acceptance set collapses to a point. Any such point is called a bargaining outcome. Section 5 introduces the characteristic function and the main result of the paper: there is a unique bargaining outcome being the (unique) generalized zero of the characteristic function. Section 6 provides the proof of the main result.

2 The model

We study a world ω described by the following variables: X , N , μ , $\mathcal{C}$ and $u_\bullet$. The symbol X denotes the unit interval $[0, 1]$. This is a space of alternatives or social states the players must choose from. The set N is a finite set of players and μ is a probability distribution on N . The probability $\mu(i)$ is a recognition probability of player i , that is the probability for player i to be chosen a proposer. The symbol $u_\bullet$ denotes a collection of utility functions, one for each player. The utility function of player $t \in N$ is $u_t : X \rightarrow [0, 1]$. We shall assume that the utility functions satisfy the following assumption.

- (A1) For each $t \in N$ the utility function $u_t : X \rightarrow [0, 1]$ is concave, continuous, and it attains its unique maximum at point $\bar{x}_t$.

The symbol $\mathcal{C}$ is a collection of subsets of N representing a voting rule. The sets in $\mathcal{C}$ are decisive coalitions in the sense that an approval of a proposal by any of these coalitions is sufficient for a passing of a proposal. For instance, if the passing of a proposal requires

the approval of it by each player, then the collection $\mathcal{C}$ consists of the set N alone. If the acceptance of a proposal requires the approval of it by at least q voters, then the collection $\mathcal{C}$ consists of all sets $C \subset N$ such that $|C| = q$. In particular, the simple majority rule is a quota rule where the quota q equals $n + 1$ in a game with $2n + 1$ players. An interesting voting rule is an adaptation of the rule used by the United Nations Security Council. The Council consists of five permanent and ten temporary members, and the passing of its resolutions requires the yes-votes of at least nine members. Furthermore, each permanent member has a veto right, which means that the no-vote by any of the permanent members overrules any majority. Letting P denote the set of permanent members, the collection $\mathcal{C}$ consists of all sets $C \subset N$ such that $P \subset C$ and $|C| = 9$. We shall assume the following.

(A2) For each C_1 and C_2 in $\mathcal{C}$ the inequality $\mu(C_1 \cap C_2) > 0$ holds.

A restriction on the collection $\mathcal{C}$ of decisive coalitions commonly employed in the literature on voting is that for each C_1 and C_2 in $\mathcal{C}$ the set $C_1 \cap C_2$ be non-empty. Assumption **(A2)** not only requires that the set $C_1 \cap C_2$ be non-empty, but also that it contain a player having a positive probability of recognition. Assumption **(A2)** is trivially satisfied in the case of the unanimity acceptance rule, i.e. when $\mathcal{C} = \{N\}$. If $|N|$ is an odd number and the collection $\mathcal{C}$ represents the simple majority rule, then assumption **(A2)** implies that all players's recognition probabilities are positive, because for each $i \in N$ there exist two decisive coalitions C_1 and C_2 such that $C_1 \cap C_2 = \{i\}$. If $\mathcal{C}$ represents the voting rule of the Security Council as described above, for the assumption **(A2)** to be satisfied it is sufficient that at least one permanent member have a positive recognition probability.

Given a discount factor $\delta \in [0, 1]$ we define a game of bargaining $\Gamma(\delta)$ as follows. The game starts in period zero. Each period τ begins with nature randomly choosing a player from the set N to make a proposal. Player i is chosen by nature with probability $\mu(i)$. The chosen player proposes an alternative x from X . All players (including the proposer) respond sequentially, according to a total order $>$ on the player set N . The order $>$ is fixed throughout the game. Each responder can either accept or reject the current proposal. If all players in some decisive coalition $C \in \mathcal{C}$ accept the proposal, the game terminates and the proposal is implemented. Otherwise, period $\tau + 1$ begins.

If alternative x is agreed upon in period τ , player i receives a payoff of $\delta^\tau u_i(x)$. The payoff of perpetual disagreement to any player is zero.

The game tree of $\Gamma(\delta)$ is similar to that in Banks and Duggan [1] with the difference that in the game $\Gamma(\delta)$ the players vote sequentially, while in [1] the votes are cast simultaneously. We insist that the players vote sequentially to rule out coordination problems in the response stage of the game, for example a situation when any point in the set X is unanimously accepted independently of preferences.

We shall restrict our attention to stationary strategies by which we mean the strategies such that (a) a proposal of any player is independent of the history of play and (b) the reaction of a player to a proposal only depends on the proposal itself. Thus a stationary strategy of player t consists of a proposal x_t and an acceptance set A_t . Player t proposes the point x_t whenever player t is chosen to be a proposer and he accepts a proposal x if

and only if x is an element of the set A_t . A joint stationary strategy $\sigma = (x_\bullet, A_\bullet)$ induces the acceptance set $A_C = \cap_{t \in C} A_t$ of coalition C and the social acceptance set $A = \cup_{C \in \mathcal{C}} A_C$. A proposal x is implemented if and only if it is an element of the social acceptance set A .

Consider a case where the proposal x_i of each player i lies in the social acceptance set A . In this case each proposal is accepted without delay. The expected utility y_t of player t can then be computed as a weighted sum of $u_t(x_i)$, where the weights are equal to the respective recognition probabilities. The utility y_t is also player t 's continuation payoff when a proposal is turned down. Therefore, player t accepts a proposal x if and only if $u_t(x) \geq \delta y_t$. Since all proposals in the set A pass, player t proposes a point x_t that maximizes player t 's utility on A . These considerations motivate the following definition.

Definition 1 A joint stationary strategy $\sigma = (x_\bullet, A_\bullet)$ is a δ -equilibrium of the world ω if there exist the numbers $(y_t)_{t \in N}$, the sets $(A_C)_{C \in \mathcal{C}}$ and the set A such that the following conditions are satisfied:

$$\begin{aligned} x_t &= \arg \max_{x \in A} u_t(x) \text{ for each } t \in N, \\ y_t &= \sum_{i \in N} \mu(i) u_t(x_i) \text{ for each } t \in N, \\ A_t &= \{x \in X | u_t(x) \geq \delta y_t\} \text{ for each } t \in N, \\ A_C &= \cap_{t \in C} A_t \text{ for each } C \in \mathcal{C}, \\ A &= \cup_{C \in \mathcal{C}} A_C. \end{aligned}$$

A δ -equilibrium can be seen to be equivalent to the definition of stationary equilibrium (in pure strategies) in Banks and Duggan [1] and Cho and Duggan [4]. In particular, the existence of δ -equilibrium is established in Theorem 2 in Banks and Duggan [1]. Cho and Duggan [4] show that δ -equilibrium is unique for each $\delta < 1$ under the assumption that the utility functions u_i are quadratic and the voting rule $\mathcal{C}$ is proper and strong. In general, there may be multiple δ -equilibria, an example being given in Cho and Duggan [?].

Since we assume the utility functions to be concave, each individual acceptance set is a closed interval. We shall use the notation $[x_t^-, x_t^+]$ to denote the individual acceptance set A_t of player t . The acceptance set of coalition C is also a closed interval, denoted by $[x_C^-, x_C^+]$. Furthermore,

$$x_C^- = \max_{t \in C} \{x_t^-\} \text{ and } x_C^+ = \min_{t \in C} \{x_t^+\}.$$

We write $E(x_\bullet)$ to denote the expected proposal, that is

$$E(x_\bullet) = \sum_{i \in N} \mu(i) x_i.$$

Proposition 1 Let $(x_\bullet, A_\bullet)$ be a δ -equilibrium inducing the social acceptance set A . Then A is a closed interval containing the point $E(x_\bullet)$.

Figure 1: Equilibrium proposals.

Proof. Each individual acceptance set A_t contains the point $E(x_\bullet)$, because

$$u_t(E(x_\bullet)) \geq E(u_t(x_\bullet)) = y_t \geq \delta y_t$$

by the concavity of the function u_t . It follows that the acceptance set A_C of each coalition C contains the point $E(x_\bullet)$. A finite union of closed intervals having a point in common is obviously closed interval. $\square$

We let the social acceptance set be $[x_-, x_+]$, where the endpoints can be computed as

$$x_- = \min_{C \in \mathcal{C}} \{x_C^-\} \text{ and } x_+ = \max_{C \in \mathcal{C}} \{x_C^+\}.$$

The equilibrium proposal x_t of player t is a point of $[x_-, x_+]$ closest to $\bar{x}_t$, the ideal point of individual t . Figure 1 illustrates. Thus

$$x_t = \begin{cases} x_- & \text{if } \bar{x}_t \leq x_- \\ \bar{x}_t & \text{if } x_- \leq \bar{x}_t \leq x_+ \\ x_+ & \text{if } x_+ \leq \bar{x}_t. \end{cases}$$

Definition 2 For each natural n let A^n be a social acceptance set in a δ^n -equilibrium of the world ω and let x^n be a point of A^n . Suppose that the sequence δ^n converges to one and x^n converges to a point x . Then the alternative x is called a bargaining outcome of ω . A collection of bargaining outcomes is called a bargaining solution.

The main result of the paper states that the bargaining outcome is unique. In particular, along any sequence of δ -equilibria, as δ converges to one, the social acceptance set collapses to a point, and in the limit all players make the same proposal.

3 Subgame perfect equilibria in stationary strategies

In this section we show that each δ -equilibrium is a subgame perfect equilibrium of the game $\Gamma(\delta)$. Conversely, given a subgame perfect equilibrium in stationary strategies there exists a δ -equilibrium having the same equilibrium proposals and inducing the same expected payoffs. In particular, each subgame perfect equilibrium in stationary strategies is a no-delay equilibrium. In this sense, the set of δ -equilibria effectively represents all subgame perfect equilibria in stationary strategies. One can say that δ -equilibrium is a subgame perfect equilibrium in stationary strategies where each player votes sincerely.

Theorem 1 *Each δ -equilibrium is a subgame perfect equilibrium of the game $\Gamma(\delta)$.*

The proof of Theorem 1 consists of two steps. The first step is to show that the strategy σ is robust to all one-shot deviations. Recall that a strategy $\bar{\sigma}_t$ for player t is said to be a one-shot deviation from σ at node h if it coincides with the strategy σ_t on all nodes but h . The second step establishes the one-shot deviation property for the game $\Gamma(\delta)$: if there is a profitable deviation from the joint strategy σ , then there is a profitable one-shot deviation from σ .

Proposition 2 *Let $\sigma = (x_\bullet, A_\bullet)$ be a δ -equilibrium. Then no player has a one-shot profitable deviation from σ at any node of the game Γ .*

Proof. Let $\bar{\sigma}_t$ be a one-shot deviation from σ_t at node h . Let $\Gamma(h)$ be the subgame that starts at node h .

Suppose player t has to make a proposal at node h . Under the strategy σ_t player t proposes the alternative x_t , which is accepted, leading to a payoff of $u_t(x_t)$ for player t . Suppose under the strategy $\bar{\sigma}_t$ player t makes a proposal x . If x is not an element of the social acceptance set $A = \cup_{C \in \mathcal{C}} \cap_{i \in C} A_i$, the proposal x will be rejected. As $\bar{\sigma}_t$ coincides with σ_t on all nodes following h , player t will receive a payoff of δy_t . By the definition of δ -equilibrium, $\delta y_t \leq u_t(x_t)$. If x is an element of A , then the proposal x is accepted and player t receives a payoff of $u_t(x)$. However, $u_t(x) \leq u_t(x_t)$, because by the definition of δ -equilibrium, x_t maximizes the function u_t on the set A . We conclude that the strategy $\bar{\sigma}_t$ does not give a higher payoff than the strategy σ_t in the subgame $\Gamma(h)$.

Suppose player t has to react to a proposal x at node h . Suppose that $u_t(x) \geq \delta y_t$. Then $x \in A_t$, and the strategy σ_t accepts the proposal x . The strategy $\bar{\sigma}_t$ rejects the proposal x , because it is assumed to be a deviation from σ_t at the node h . If in the subgame $\Gamma(h)$ the joint strategy (σ_t, σ_{-t}) leads to a social rejection of the proposal x , so does the joint strategy $(\bar{\sigma}_t, \sigma_{-t})$, since the latter gives one less yes-vote to x than the former. In that case the payoff to player t in the subgame $\Gamma(h)$ is δy_t , whether player t uses the strategy σ_t or $\bar{\sigma}_t$. Suppose that in the subgame $\Gamma(h)$ the joint strategy (σ_t, σ_{-t}) leads to a social acceptance of the proposal x , so the payoff to player t on the strategy σ_t is $u_t(x)$. The payoff on the strategy $\bar{\sigma}_t$ in the subgame $\Gamma(h)$ is either $u_t(x)$ or δy_t , depending on whether the joint strategy $(\bar{\sigma}_t, \sigma_{-t})$ leads to a social acceptance or rejection of the proposal x , but in either case is not higher than the payoff on the strategy σ_t .

Suppose that $u_t(x) < \delta y_t$. Then x is not in A_t , so the strategy σ_t rejects the proposal x , and the strategy $\bar{\sigma}_t$ accepts the proposal x at the node h . If in the subgame $\Gamma(h)$ the joint strategy (σ_t, σ_{-t}) leads to a social acceptance of the proposal x , so does the joint strategy $(\bar{\sigma}_t, \sigma_{-t})$, since under the joint strategy $(\bar{\sigma}_t, \sigma_{-t})$ the proposal x receives one more yes-vote than under the strategy (σ_t, σ_{-t}) . In that case the payoff to player t in the subgame $\Gamma(h)$ is $u_t(x)$ on both strategies σ_t and $\bar{\sigma}_t$. Suppose that in the subgame $\Gamma(h)$ the joint strategy (σ_t, σ_{-t}) leads to a social rejection of the proposal x , so the payoff to player t on the strategy σ_t is δy_t . The payoff on the strategy $\bar{\sigma}_t$ in the subgame $\Gamma(h)$ is either $u_t(x)$ or δy_t , depending on whether the joint strategy $(\bar{\sigma}_t, \sigma_{-t})$ leads to a social acceptance or rejection of the proposal x . Therefore $\bar{\sigma}_t$ does not give a higher payoff than the strategy σ_t in the subgame $\Gamma(h)$. $\square$

For the proof of Proposition 3 the reader can refer to Predtetchinski [12, Proposition 2].

Proposition 3 *Let σ be a profile of strategies. If player t has a profitable deviation from σ , then player t has a profitable one-shot deviation from σ .*

We now compute the payoffs induced by a joint stationary strategy $\sigma = (x_\bullet, A_\bullet)$. Let y_t denote the expected payoff to player t at the beginning of the game induced by σ . Let $N_a = \{t \in N | x_t \in A\}$ be the set of players whose proposal is accepted under σ and $N_r = \{t \in N | x_t \notin A\}$ denote the set of players whose proposal is rejected. If nature chooses a proposer i from N_a , player t receives a payoff of $u_t(x_i)$, while if nature chooses a proposer from N_r , then player t 's payoff is δy_t . Thus y_t satisfies the following relation:

$$y_t = \sum_{i \in N_a} \mu(i) u_t(x_i) + \delta \mu(N_r) y_t.$$

Solving the equation, we obtain the following expression for y_t :

$$y_t = \begin{cases} \frac{1}{1 - \delta \mu(N_r)} \sum_{i \in N_a} \mu(i) u_t(x_i) & \text{if } \mu(N_a) > 0, \\ 0 & \text{otherwise.} \end{cases}$$

We now turn to the converse of Theorem 1. It is clear that there always exist subgame perfect equilibria in stationary strategies that are not δ -equilibria. To illustrate, take a δ -equilibrium σ and suppose that the alternative x is not in the social acceptance set induced by σ . Let T be the set of players who reject the alternative x . Let i be a player who reacts to a proposal before any player in the set T . Then, at any node of the game Γ where player i has to respond to a proposal x , player i is indifferent between accepting the proposal x and rejecting it, since it will later be rejected by the players in the set T . Consequently, we can remove the alternative x from the individual acceptance set of player i . The resulting profile of strategies will still be a subgame perfect equilibrium, but not a δ -equilibrium.

Figure 2: The response stage of the game Γ .

Thus we will not be able to show that each subgame perfect equilibrium in stationary strategies is a δ -equilibrium. However, we shall prove that, given a subgame perfect equilibrium in stationary strategies, there exists a δ -equilibrium inducing the same expected payoffs, having the same equilibrium proposals, and (provided that there are players whose ideal points are zero and one) the same social acceptance set.

Important for the proof of Theorem 2 is the concept of pivotal vote. Suppose one is given a joint stationary strategy σ . Consider a node h of the game Γ where player t has to react to some proposal, say x . Player t 's vote is said to be pivotal at node h if it determines the social acceptance or rejection of the proposal x .

We illustrate the idea of pivotal vote by means of an example. Suppose there are three players, and the yes-votes of any two players are sufficient for a passing of a proposal. The order of response is $1 < 2 < 3$. Suppose that under the given joint stationary strategy the proposal x is rejected by all three players. Figure 2 depicts the response stage of the game Γ . The capital letters in Figure 2 indicate whether the proposal in question is socially accepted or not, and the arrows indicate the strategies.

The player 2's vote at the node h_2 is pivotal. Indeed, if player 2 accepts at the node h_2 , then the proposal x will be socially accepted, and if player 2 rejects, the proposal will be turned down. Notice that player 2's vote at node h_1 is not pivotal, because the proposal will be socially rejected, whether player 2 votes for or against it. In the same way, player 1 is not pivotal at his decision node.

Definition 3 Let σ be a joint stationary strategy. Consider a node h of the game Γ where player t has to react to some proposal, say x . Let the σ'_t and σ''_t be strategies for player t that coincide with the strategy σ_t at any node of the game Γ but h , where σ'_t accepts x and σ''_t rejects x . Player t is said to be *pivotal* at node h if the proposal x is socially accepted on the path of play induced by the joint strategy (σ'_t, σ_{-t}) in the subgame $\Gamma(h)$ and is socially rejected on the path of play induced by the joint strategy $(\sigma''_t, \sigma_{-t})$ in the subgame $\Gamma(h)$.

If all players were voting sincerely, we would expect player t to accept a proposal x if the utility of x to player t is higher than t 's continuation utility and to reject it if the utility of x is lower than the continuation utility. In fact, the players need not vote sincerely, even in a subgame perfect equilibrium. Consider again Figure 2. Since player 1's vote is irrelevant, both rejecting and accepting the proposal are player 1's best responses to player 2's and 3's strategies. Thus player 1 may well reject a proposal x even though the utility of x is higher than player 1's continuation utility. On the other hand, player 2's vote is decisive at h_2 , so player 2 receives a utility of δy_2 if he chooses to reject x and a utility of $u_2(x)$ if he chooses to accept it. Since in a subgame perfect equilibrium player 2 should maximize his utility at any node of the game, including the node h_2 , player 2 should accept the proposal x if $u_2(x) > \delta y_2$ and reject it if $u_2(x) < \delta y_2$.

It is therefore important to know whether a player is ever pivotal or not. This is answered by Proposition 4 below.

Proposition 4 *Let $\sigma = (x_\bullet, A_\bullet)$ be a joint stationary strategy in the game Γ inducing the acceptance set A_C for coalition C and the social acceptance set A . Let $x \in X$.*

- (a) *Suppose x is not an element of A . Then for each $C \in \mathcal{C}$ there is a player $t \in C$ such that x is not an element of the set A_t and player t 's vote is pivotal at some node h of the game Γ .*
- (b) *Suppose x is an element of A . Then there exists a $C \in \mathcal{C}$ such that x is an element of A_C and for each $t \in C$ player t 's vote is pivotal at some node h of the game Γ .*

The proof of Proposition 4 is very involved and is relegated to the appendix. We are now in a position to prove Theorem 2.

Theorem 2 *Suppose $0 \leq \delta < 1$. Consider a joint strategy $\sigma = (x_\bullet, A_\bullet)$ inducing the expected payoffs $y_\bullet$ and the social acceptance set A . Suppose σ is a subgame perfect equilibrium of the game $\Gamma(\delta)$. Then there exists a δ -equilibrium $(x_\bullet, A'_\bullet)$ inducing the expected payoffs $y_\bullet$ and the social acceptance set A' such that $\text{Int} A' \subset A \subset A'$. Furthermore, if there are players i^0 and i^1 such that $\bar{x}_{i^0} = 0$ and $\bar{x}_{i^1} = 1$, then $A = A'$.*

Proof. Define the following sets:

$$\begin{aligned} A'_i &= \{x \in X | u_i(x) \geq \delta y_i\} & A''_i &= \{x \in X | u_i(x) > \delta y_i\} \\ A'_C &= \cap_{i \in C} A'_i & A''_C &= \cap_{i \in C} A''_i \\ A' &= \cup_{C \in \mathcal{C}} A'_C & A'' &= \cup_{C \in \mathcal{C}} A''_C. \end{aligned}$$

Step 1. We prove the inclusion $A'' \subset A$.

Suppose x is not an element of A . Take a $C \in \mathcal{C}$. By part (a) of Proposition 4, there is a player $t \in C$ such that x is not in A_t and player t 's vote is pivotal at a node h of the game Γ . We now prove that x is not an element of the set A''_t . It then follows that it is not an element of the set A''_C , and, since C is an arbitrary coalition in $\mathcal{C}$, that it is not an element of the set A'' .

Let σ'_t and σ''_t be as in Definition 3. Thus σ'_t and σ''_t coincide with the strategy σ_t at any node of the game Γ but h , where σ'_t accepts x and σ''_t rejects x . Then the proposal x is socially accepted on the path of play of the joint strategy (σ'_t, σ_{-t}) in the subgame $\Gamma(h)$ and is rejected on the path of play of the joint strategy $(\sigma''_t, \sigma_{-t})$. It follows that the payoff to player t on his strategy σ'_t in the subgame $\Gamma(h)$ is $u_t(x)$ and the payoff on the strategy σ''_t is δy_t . Now since x is not an element of A_t , player t rejects the proposal x under the strategy σ_t , so in fact $\sigma_t = \sigma''_t$. Since σ induces a Nash equilibrium in the subgame $\Gamma(h)$, we must have that $u_t(x) \leq \delta y_t$. Thus x is not an element of the set A''_t , as desired.

Step 2. We prove the inclusion $A \subset A'$.

Let x be an element of A . By part (b) of Proposition 4, there exists a $C \in \mathcal{C}$ such that x is an element of A_C and for each $t \in C$ player t 's vote is pivotal at some node h of the game Γ . We now prove that x is an element of A'_C .

As before, let σ'_t and σ''_t be as in Definition 3. Then the payoff to player t on his strategy σ'_t in the subgame $\Gamma(h)$ is $u_t(x)$ and the payoff on the strategy σ''_t is δy_t . Since x is an element of A_t , player t accepts the proposal x under the strategy σ_t , so in fact $\sigma_t = \sigma'_t$. Since σ induces a Nash equilibrium in the subgame $\Gamma(h)$, we must have that $\delta y_t \leq u_t(x)$. It follows that $x \in A'_t$. Since t is an arbitrary player of the coalition C , we conclude that $x \in A'_C$, as desired.

Step 3. Let $N_a = \{t \in N | x_t \in A\}$. We show that $\mu(N_a) > 0$.

If $\mu(N_a) = 0$, then $y_t = 0$ for all $t \in N$. Because all functions u_t are positive on the interior of X , the inclusion $\text{Int}X \subset A''_t \subset A_t$ holds for each $t \in N$. Thus each alternative in the interior of X is unanimously accepted under the joint strategy σ . It follows that any player $t \in N \setminus N_a$ has a one-shot profitable deviation at any node where t has to make a proposal, namely to propose a point $x \in \text{Int}X$ rather than x_t . Indeed, the proposal x_t of player t is rejected, resulting in the payoff of 0, while the proposal $x \in \text{Int}X$ is socially accepted and yields a positive payoff.

From $\mu(N_a) > 0$ it follows that the set A is non-empty.

Step 4. We now prove that the set $\cap A''_t$ is non-empty, where the intersection extends over all players t in N . It then follows that both sets A'' and A' are intervals.

Let x be the average accepted proposal:

$$x = \sum_{i \in N_a} \mu(i) x_i / \mu(N_a).$$

By the concavity of the function u_t

$$u_t(x) \geq \sum_{i \in N_a} \mu(i) u_t(x_i) / \mu(N_a) \geq \sum_{i \in N_a} \mu(i) u_t(x_i) / [1 - \delta \mu(N_r)] = y_t \geq \delta y_t,$$

which shows that x is an element of the set A'_t for each t . If $y_t > 0$, then the extreme right inequality is strict, so the point x is an element of the set A''_t . And if $y_t = 0$, the the set A''_t contains the open interval $\text{Int}X$. Thus the set

$$\bigcap_{\{t \in N | y_t > 0\}} A''_t$$

is a non-empty open subset of X . Any non-empty open subset of X has a non-empty intersection with the interior of X , and therefore with the set

$$\bigcap_{\{t \in N | y_t = 0\}} A_t''.$$

We conclude that $\cap A_t''$ is non-empty.

Step 5. We show that $(x_\bullet, A'_\bullet)$ is a δ -equilibrium. First we prove that $N_a = N$.

Suppose not. Then any player $t \in N \setminus N_a$ has a one-shot profitable deviation at any node where player t is a proposer, namely to propose any point $x \in \cap A_t''$ rather than the point x_t . Indeed, the proposal x_t is rejected, resulting in the payoff of δy_t for player t . The proposal x , on the other hand, will be socially accepted, because $x \in \cap A_t'' \subset A'' \subset A$, resulting in the payoff of $u_t(x)$ for player t at node h . And, since $x \in A_t''$, we have the inequality $u_t(x) > \delta y_t$. This contradicts the assumption that σ is a subgame perfect equilibrium.

Since $N_a = N$, the expected utility of player t can be written as

$$y_t = \sum_{i \in N} \mu(i) u_t(x_i).$$

Finally, we have to show that the proposal x_t of player t maximizes the function u_t on A' . Suppose not. Then there is an $x \in \text{Int} A'$ such that $u_t(x) > u_t(x_t)$. But then player t has a one-shot profitable deviation at any node where player t has to make a proposal, namely to propose the point x rather than the point x_t . Indeed, the proposal x is socially accepted under the strategy profile σ , because $x \in \text{Int} A' \subset A'' \subset A$, and it gives a higher payoff to player t than x_t . This contradicts the hypothesis that σ is a subgame perfect equilibrium.

Step 6. Suppose there are players i^0 and i^1 such that $\bar{x}_{i^0} = 0$ and $\bar{x}_{i^1} = 1$. We show that $A = A'$. As $\text{Int} A' \subset A \subset A'$, it suffices to prove that A contains both endpoints of the interval A' . We already know that $(x_\bullet, A'_\bullet)$ is a δ -equilibrium. As the ideal point of player i^0 is zero, the proposal x_{i^0} of player s equals the left endpoint of the interval A' . The ideal point of player i^1 is one, so x_{i^1} equals the right endpoint of the interval A' . As we have already proved in Step 5, $N_a = N$, so both x_{i^0} and x_{i^1} are elements of the set A , as desired. $\square$

4 The social acceptance set

In this section we establish that along any sequence of δ -equilibria, as δ converges to one, the social acceptance set collapses to a point. The idea of the proof is very simple: we first consider the image $u_t(A)$ of the social acceptance set in the utility space. We show that for at least one player t the length of the interval $u_t(A)$ is bounded above by $(1 - \delta)/\nu$, where ν depends only on the recognition probabilities. It follows that $u_t(A)$ converges to a singleton as δ goes to one. Strict quasi-concavity of the function u_t then implies that

the set A also converges to a singleton. To highlight the importance of assumption **(A2)** for this result we consider a world where assumption **(A2)** is violated and consequently Theorem 3 fails.

At the end of the section a special case of simple majority rule is considered. If the simple majority is required for a passing of a proposal, then the unique bargaining outcome is the ideal point of the median player, i.e. that individual who divides the set of players into two coalitions of equal size, those with lower ideal points and those with higher ideal points. This result has been earlier established in Banks and Duggan [?].

Let I^n be a sequence of compact intervals and I be a compact interval. We say that the sequence I^n converges to I if the sequence of left endpoints of I^n converges to the left endpoint of I and the sequence of right endpoints of I^n converges to the right endpoint of I . We write $\text{Len}(I)$ for the length of an interval I , the difference of the upper and lower endpoints. Let $S = \{i \in N | \mu(i) > 0\}$ and let $\nu = \min_{i \in S} \{\mu(i)\}$.

Proposition 5 *Let $a_\bullet$ be an element of $[0, 1]^N$. Let $\delta \in [0, 1]$. Suppose that $\delta E(a_\bullet) \leq a_i$ for all $i \in S$. Then $a_i \leq (1 - \delta + \nu\delta)E(a_\bullet)/\nu$ for all $i \in S$.*

Proof. For each $i \in S$ we have the following chain of inequalities:

$$\begin{aligned} E(a_\bullet) &= \sum_{t \in S} \mu(t) a_t \\ &\geq \mu_i a_i + (1 - \mu_i) \delta E(a_\bullet) \\ &= \delta E(a_\bullet) + \mu_i (a_i - \delta E(a_\bullet)) \\ &\geq \delta E(a_\bullet) + \nu (a_i - \delta E(a_\bullet)) \\ &= \nu a_i + (1 - \nu) \delta E(a_\bullet). \end{aligned}$$

Rearranging yields the desired inequality. $\square$

We now estimate the length of the image of the social acceptance set in the utility space. Notice that assumption **(A2)** plays a crucial role in the argument.

Proposition 6 *Let $(x_\bullet, A_\bullet)$ be a δ -equilibrium inducing the expected payoffs $y_\bullet$ and the social acceptance set A . Then $\text{Len}(u_t(A)) \leq (1 - \delta)/\nu$ for some $t \in S$.*

Proof. Let $A = [x_-, x_+]$. There exist coalitions C^- and C^+ in $\mathcal{C}$ such that $x_- \in A_{C^-}$ and $x_+ \in A_{C^+}$. Now, by assumption **(A2)** there exists a player $t \in C^- \cap C^+$ with $\mu(t) > 0$. Thus, both points x_- and x_+ belong to the individual acceptance set A_t of player t . Since A_t is an interval, it contains the entire social acceptance set A . Fix one such player t .

We can estimate the utility of player t on the social acceptance set A as follows. Since $A \subset A_t$, the utility of player t on the set A is bounded below by δy_t . On the other hand, the proposal x_t of player t maximizes player's t utility on the set A . Thus

$$u_t(A) \subset [\delta y_t, u_t(x_t)].$$

Now we apply Proposition 5. Let $a_i = u_t(x_i)$ for all $i \in N$ and notice that $y_t = E(a_\bullet)$. Then we have $a_i \geq \delta E(a_\bullet)$ as in Proposition 5, because each x_i is an element of A , and therefore an element of A_t . Proposition 5 implies, in particular, that

$$u_t(x_t) \leq (1 - \delta + \nu\delta)y_t/\nu.$$

We can thus estimate the length of the interval $u_t(A)$ as follows:

$$\text{Len}(u_t(A)) \leq (1 - \delta + \nu\delta)y_t/\nu - \delta y_t = (1 - \delta)y_t/\nu \leq (1 - \delta)/\nu,$$

where the last inequality follows from the fact that $y_t \leq 1$. The result follows. $\square$

We are now in a position to prove the main result of this section.

Theorem 3 *For each natural number n let A^n be a social acceptance set in a δ^n -equilibrium. If the sequence A^n converges to a compact interval A and the sequence δ^n converges to 1, then A is a singleton.*

Proof. Since the set S is finite, we can assume without loss of generality that there exists a player t in S such that

$$\text{Len}(u_t(A^n)) \leq (1 - \delta^n)/\nu$$

for each n . Thus the sequence $\text{Len}(u_t(A^n))$ converges to zero. Now, $\text{Len}(u_t(I))$ is a continuous function of the endpoints of the interval I . It follows that $\text{Len}(u_t(A)) = 0$, i.e. the function u_t is constant on the interval A . By assumption (A1) the function u_t is non-constant on any non-degenerate interval. It follows that A is a singleton. $\square$

We now show, by means of an example, that Theorem 3 need not be true if assumption (A2) is violated. Let the player set N consist of player 0 with the utility function $u_0(x) = 1 - x$, player 1 with the utility function $u_1(x) = x$ and a set of players M all having the same utility function $u_i(x) = 1 - |x - 1/2|$. The utility functions are illustrated in Figure 3. Assumption (A1) is clearly satisfied.

Think of the set M as consisting of a very large number of players. It is then reasonable to assume that there exists a subset C of M such that both $C_0 = \{0\} \cup C$ and $C_1 = \{1\} \cup C$ are decisive coalitions. Furthermore, we assume that $\mu(0) = \mu(1) = 1/2$. Thus only players 0 and 1 ever make proposals. Assumption (A2) is violated because $C_0 \cap C_1 = C$ and $\mu(C) = 0$.

Now, for each $\delta \in [0, 1]$ there exists a δ -equilibrium where X is the social acceptance set. In this equilibrium each player proposes his own ideal point: player 0 proposes 0, each player in M proposes the point $1/2$ and player 1 proposes the point 1. The expected utility is $1/2$ for all players. The individual acceptance set of player 0 is $A_0 = [0, (1 + \delta)/2]$, the individual acceptance set of player 1 is $A_1 = [(1 - \delta)/2, 1]$ and that of each player in M is X . The acceptance set of coalition C_0 equals the individual acceptance set of player 0, while the acceptance set of coalition C_1 equals the individual acceptance set of player 1.

Figure 3: The utility functions in the example of Section 4.

The union of A_0 and A_1 is clearly the entire set X . Thus the social acceptance set equals X .

We conclude this section with two additional results. First we show that each bargaining outcome of the world ω lies in the interval $[\alpha, \beta]$ as defined below. We also reestablish that, under the simple majority rule, there exists a unique bargaining outcome that only depends on the set of players and their ideal points. The latter result has been established by Banks and Duggan [1], see Theorem 5 and the discussion thereafter.

Define α and β as follows:

$$\alpha = \max_{C \in \mathcal{C}} \min_{t \in C} \{\bar{x}_t\} \text{ and } \beta = \min_{C \in \mathcal{C}} \max_{t \in C} \{\bar{x}_t\}.$$

For example, in the case of the unanimity acceptance rule ($\mathcal{C} = \{N\}$) α equals the minimum of the ideal points and β is the maximal ideal point.

Proposition 7 *The inequality $\mu(\{i \in N | \alpha \leq \bar{x}_i \leq \beta\}) > 0$ holds. In particular, $\alpha \leq \beta$.*

Proof. Let $C_\alpha \in \mathcal{C}$ and $C_\beta \in \mathcal{C}$ be such that

$$\alpha = \min_{t \in C_\alpha} \{\bar{x}_t\} \text{ and } \beta = \max_{t \in C_\beta} \{\bar{x}_t\}.$$

Thus $\alpha \leq \bar{x}_t$ for all $t \in C_\alpha$ and $\bar{x}_t \leq \beta$ for all $t \in C_\beta$. By Assumption **(A2)** there exists a player $i \in C_\alpha \cap C_\beta$ such that $\mu(\{i\}) > 0$. Then $\alpha \leq \bar{x}_i \leq \beta$. This proves the result. $\square$

Proposition 8 *Let $\delta \in [0, 1]$. Let $A = [x_-, x_+]$ be the social acceptance set in a δ -equilibrium of the world ω . Then $x_- \leq \beta$ and $\alpha \leq x_+$.*

Proof. As before, let $[x_t^-, x_t^+]$ be the individual acceptance set of player t . The point $\bar{x}_t$ is the maximum of the function u_t on the set X , and so is an element of A_t . Thus

$x_t^- \leq \bar{x}_t \leq x_t^+$ for each $t \in N$. Using the formulas for x_- and x_+ of Section 2, we obtain

$$\begin{aligned} x_- &= \min_{C \in \mathcal{C}} \max_{t \in C} \{x_t^-\} \leq \min_{C \in \mathcal{C}} \max_{t \in C} \{\bar{x}_t\} = \beta, \\ x_+ &= \max_{C \in \mathcal{C}} \min_{t \in C} \{x_t^+\} \geq \max_{C \in \mathcal{C}} \min_{t \in C} \{\bar{x}_t\} = \alpha. \end{aligned}$$

□

Proposition 9 *Each bargaining outcome of the world ω lies in the interval $[\alpha, \beta]$.*

Proof. Let x be a bargaining outcome of ω . By Definition 2, the point x is a limit of a sequence x^n of points where each x^n is an element of the social acceptance set $A^n = [x_-^n, x_+^n]$ in a δ^n -equilibrium of ω and the sequence δ^n converges to one. Then by Theorem 3 both sequences x_-^n and x_+^n converge to the point x . By Proposition 8, $x_-^n \leq \beta$ and $\alpha \leq x_+^n$ for each n . Taking the limit, we obtain the inequalities $x \leq \beta$ and $\alpha \leq x$. □

We now turn to a special case of the simple majority rule.

Proposition 10 *Consider a world ω where $|N| = 2n + 1$ and the collection $\mathcal{C}$ consists of all coalitions $C \subset N$ with $|C| = n + 1$. Then $\alpha = \beta$ is the unique bargaining outcome of the world ω . Furthermore, $\alpha = \beta$ is the unique point x satisfying the inequalities*

$$|\{i \in N | \bar{x}_i \leq x\}| \geq n + 1 \text{ and } |\{i \in N | x \leq \bar{x}_i\}| \geq n + 1.$$

Proof. The fact that $\alpha = \beta$ and the inequalities are established by a direct verification. The fact that $\alpha = \beta$ is the unique bargaining outcome follows from Proposition 9. □

For example, if there are seven players four of whom have an ideal point equal to 0 and the remaining three have an ideal point 1, then 0 is the unique bargaining outcome. If no two players have the same ideal points, then the unique bargaining outcome is an ideal point of the median voter, i.e. that player who divides the set of players into two coalitions of equal size, those with a lower ideal point and those with a higher ideal point. Notice that this point only depends on the collection of players' ideal points. It does not depend on the shape of the utility functions or recognition probabilities.

In the case of the simple majority rule the computation of the bargaining outcome is trivial, because the points α and β coincide, so they must be the unique bargaining outcome by Proposition 9. One other special case where $\alpha = \beta$ is when all players' ideal points coincide, in which case $\alpha = \beta$ is the common ideal point and therefore the unique bargaining outcome. In most cases of interest, however, $\alpha < \beta$, and Proposition 9 does not pin down a specific point as a bargaining outcome. To compute the bargaining outcome we rely on a technique based on the characteristic function.

5 The characteristic function

5.1 The characteristic function and the main result

Remark 1 Each function u_t has left and right derivatives at each point x of the interior of X denoted by $l_t(x)$ and $r_t(x)$. Both the left and the right derivatives are non-increasing functions of x and $r_t(x) \leq l_t(x)$. The interval $[r_t(x), l_t(x)]$ is the subgradient of u_t at point x . Given $x \in \text{Int}X$ and $\dot{x} \in X$, the inequality $u_t(\dot{x}) - u_t(x) \leq s(\dot{x} - x)$ holds for each $s \in [r_t(x), l_t(x)]$. Furthermore, $r_t(\bar{x}_t) \leq 0 \leq l_t(\bar{x}_t)$, because $\bar{x}_t$ is the maximum of u_t on X .

For each $x \in \text{Int}X$ we define

$$\begin{aligned}\varphi_-(x) &= \max_{C \in \mathcal{C}} \min_{t \in C} \{r_t(x)/u_t(x), 0\} \text{ and } \varphi_+(x) = \min_{C \in \mathcal{C}} \max_{t \in C} \{l_t(x)/u_t(x), 0\}, \\ \mu_-(x) &= \mu(\{i \in N | x < \bar{x}_i\}) \text{ and } \mu_+(x) = \mu(\{i \in N | \bar{x}_i < x\}), \\ \xi(x) &= \mu_-(x)\varphi_-(x) + \mu_+(x)\varphi_+(x).\end{aligned}$$

These are well defined, since each function u_t is positive on the interior of the set X . The function ξ is referred to as a *characteristic function*. It will be convenient to extend the characteristic function to X by letting $\xi(0) = +\infty$ and $\xi(1) = -\infty$.

Remark 2 There is nothing particularly important about the use of left derivatives in the function φ_+ and right derivatives in φ_- . As the reader can verify, all results remain true if one replaces a derivative $l_t(x)$ or $r_t(x)$ by an arbitrary element in the subgradient $[r_t(x), l_t(x)]$ of the function u_t .

We now compute the characteristic function in a special case where the utility of an alternative x to player t depends only on the distance between x and player t 's ideal point through a function h that is the same for all players. Let N be a subset of X . Suppose that $u_t(x) = h(|x - t|)$ where $h : [0, 1] \rightarrow [0, 1]$ is a decreasing concave continuously differentiable function. Notice that the function u_t reaches its maximum on X at the point t . Let $g : [0, 1] \rightarrow [0, 1]$ be defined by $g(x) = h'(x)/h(x)$. Then

$$r_t(x)/u_t(x) = \begin{cases} -g(|x - t|) & \text{if } x < t \\ +g(|x - t|) & \text{if } t \leq x, \end{cases} \quad l_t(x)/u_t(x) = \begin{cases} -g(|x - t|) & \text{if } x \leq t \\ +g(|x - t|) & \text{if } t < x. \end{cases}$$

Since g is a non-positive decreasing function, both functions $r_t(x)/u_t(x)$ and $l_t(x)/u_t(x)$ are decreasing in the argument x for each t and are increasing in the argument t for each x . Using the latter fact we compute:

$$\begin{aligned}\varphi_-(x) &= \max_{C \in \mathcal{C}} \min_{t \in C} \{r_t(x)/u_t(x)\} = r_\alpha(x)/u_\alpha(x), \text{ whenever } \varphi_-(x) < 0, \\ \varphi_+(x) &= \min_{C \in \mathcal{C}} \max_{t \in C} \{l_t(x)/u_t(x)\} = l_\beta(x)/u_\beta(x), \text{ whenever } \varphi_+(x) > 0,\end{aligned}$$

where α and β are as in Section 4. We can restrict our attention to the interval $[\alpha, \beta]$, since we already know that it contains all bargaining outcomes. Since $r_\alpha(x)/u_\alpha(x) = g(x - \alpha)$ is non-positive and $l_\beta(x)/u_\beta(x) = -g(\beta - x)$ is non-negative on the interval $[\alpha, \beta]$, we have the following expression for the characteristic function:

$$\xi(x) = \mu_-(x)g(x - \alpha) - \mu_+(x)g(\beta - x) \text{ for each } \alpha \leq x \leq \beta.$$

Proposition 11 *The characteristic function ξ is a decreasing function on X .*

Proof. Given a real-valued function f on X let $\text{supp}(f) = \{x \in X \mid |f(x)| > 0\}$.

Consider the functions φ_+ and φ_- . Clearly, $\varphi_+(x) \geq 0$ and $\varphi_-(x) \leq 0$ for each $x \in \text{Int}X$. Using the fact that the functions $l_t(x)$ and $r_t(x)$ are non-increasing, it is easy to show that both $l_t(x)/u_t(x)$ and $r_t(x)/u_t(x)$ are decreasing functions on $\text{Int}X$ for each t . It then follows that both functions φ_+ and φ_- are non-increasing on $\text{Int}X$. Furthermore, φ_+ is a decreasing function on $\text{supp}(\varphi_+)$ and φ_- is a decreasing function on $\text{supp}(\varphi_-)$.

We show that

$$(0, \beta) \subset \text{supp}(\varphi_+) \subset [0, \beta].$$

To prove the first inclusion, let the point $0 < x < \beta$ be given. Then for each $C \in \mathcal{C}$ there exists a player $t \in C$ such that $x < \bar{x}_t$. For this player t it holds that $r_t(x) > 0$. It follows that $\varphi_+(x) > 0$, as desired. To prove the second inclusion let $\beta < x < 1$ be given. Then there exists a coalition $C \in \mathcal{C}$ such that for each $t \in C$ we have $\bar{x}_t < x$, so $r_t(x) < 0$. It follows that $\varphi_+(x) = 0$, as desired. Similarly, one shows that

$$(\alpha, 1) \subset \text{supp}(\varphi_-) \subset [\alpha, 1].$$

Consider now the function ξ . Write the function ξ as $\xi = \xi_- + \xi_+$, where $\xi_-(x) = \mu_-(x)\varphi_-(x)$ and $\xi_+(x) = \mu_+(x)\varphi_+(x)$. Of course, $\xi_-(x) \leq 0$ and $0 \leq \xi_+(x)$ for each $x \in \text{Int}X$.

We show that both functions ξ_+ and ξ_- are non-increasing on $\text{Int}X$. Indeed, both μ_+ and φ_+ are non-increasing and non-negative on $\text{Int}X$. It follows that $\xi_+ = \mu_+\varphi_+$ is non-increasing. The function μ_- is non-decreasing, while φ_- is non-positive and non-increasing. Therefore, $\xi_- = \mu_-\varphi_-$ is a non-increasing function.

Now, $\text{supp}(\xi_+) = \text{supp}(\mu_+) \cap \text{supp}(\varphi_+)$. Since the function φ_+ is decreasing on $\text{supp}(\varphi_+)$, the function ξ_+ is decreasing on $\text{supp}(\xi_+)$. Similarly, $\text{supp}(\xi_-) = \text{supp}(\mu_-) \cap \text{supp}(\varphi_-)$. Since the function φ_- is decreasing on $\text{supp}(\varphi_-)$, the function ξ_- is decreasing on $\text{supp}(\xi_-)$.

We now show that ξ is a decreasing function on $\text{Int}X$. Suppose not. Then there exist $0 < a < b < 1$ such that $\xi(a) \geq \xi(b)$. Because ξ_- and ξ_+ are non-increasing functions, all three must be constant on the interval $[a, b]$. Because the function ξ_- is decreasing on the set $\text{supp}(\xi_-)$ and the function ξ_+ is decreasing on $\text{supp}(\xi_+)$, both function must be identically equal to zero on the interval $[a, b]$.

In particular, $\xi_-(b) = 0$ and $\xi_+(a) = 0$. Now, $\xi_-(b) = 0$ means that either $\mu_-(b) = 0$ or $\varphi_-(b) = 0$. Similarly, $\xi_+(a) = 0$ means that either $\mu_+(a) = 0$ or $\varphi_+(a) = 0$. Now, we cannot have both $\mu_-(b) = 0$ and $\mu_+(a) = 0$, because $1 = \mu(N) \leq \mu_-(b) + \mu_+(a)$. Suppose

$\mu_+(a) > 0$. Then $\varphi_+(a) = 0$. Therefore, $\beta \leq a$. But then $\alpha \leq \beta \leq a < b$, so $\varphi_-(b) > 0$. Therefore, $\mu_-(b) = 0$. But then $\mu(\{i \in N | \alpha \leq \bar{x}_i \leq \beta\}) = 0$, contradicting Proposition 7.

Finally, ξ is a decreasing function on X because $\xi(0) = +\infty$ and $\xi(1) = -\infty$. $\square$

Definition 4 *The point $x \in X$ is a generalized zero of the function $f : X \rightarrow \mathbb{R} \cup \{-\infty, +\infty\}$ if there are sequences x_-^n and x_+^n of points in X converging to x such that $\lim f(x_-^n) \leq 0 \leq \lim f(x_+^n)$.*

It is clear that any point $x \in X$ such that $f(x) = 0$ is a generalized zero of the function f (take $x_-^n = x$ and $x_+^n = x$). Conversely, if x is a generalized zero of f and f is continuous at point x , then $f(x) = 0$. A decreasing function f has at most one generalized zero. We are now in a position to state the main result of the paper.

Theorem 4 (Main result) *Each world ω has a unique bargaining outcome. The bargaining outcome of ω is the unique generalized zero of the characteristic function ξ .*

5.2 Some intuition

In this subsection we wish to argue that the bargaining outcome is in some sense robust to renegotiations, when it is taken to be a status quo. All derivations are deliberately informal, the purpose being to provide some intuition behind, rather than the proof of, the main result.

Suppose that there has been a preliminary stage of negotiations that resulted in the agreement to implement some alternative x at the moment of time $t > 0$. Imagine now that the players negotiate again over the incremental change in the outcome together with an incremental reduction in length of delay, with the status quo being the implementation of the alternative x at time t . Let dx be an incremental change in the outcome and $-dt < 0$ be an incremental reduction in length of delay. The ratio dx/dt can be thought of as the speed of the adjustment. A player i evaluates the resulting time–outcome pair using the function $U_i(x, t) = \delta^t u_i(x)$. Obviously, each player prefers a change away from x in the direction of its own ideal point, however, a player also accepts an incremental change in the opposite direction, given a sufficiently large reduction in the length of delay. The proposal $(dx, -dt)$ is accepted by player i if results in a non-negative change in the utility U_i , so the condition for the acceptance of $(dx, -dt)$ is that

$$dU_i(x, t) = u'_i(x)dx - \ln(\delta)u_i(x)dt \geq 0.$$

Player i whose ideal point is to the left of the point x accepts a pair $(dx, -dt)$ provided that $dx/dt \leq \ln(\delta)u_i(x)/u'_i(x)$. Similarly, a player i whose ideal point is to the right of x accepts any pair $(dx, -dt)$ provided that $dx/dt \geq \ln(\delta)u_i(x)/u'_i(x)$. The acceptance set of coalition C is therefore an interval with the endpoints

$$\phi_C^-(x) = \max_{\substack{i \in C \\ x < \bar{x}_i}} \left\{ \frac{\ln(\delta)u_i(x)}{u'_i(x)}, -\infty \right\} \text{ and } \phi_C^+(x) = \min_{\substack{i \in C \\ \bar{x}_i < x}} \left\{ \frac{\ln(\delta)u_i(x)}{u'_i(x)}, +\infty \right\}.$$

The social acceptance set is an interval with the endpoints

$$\phi_-(x) = \min_{C \in \mathcal{C}} \phi_C^-(x) \text{ and } \phi_+(x) = \max_{C \in \mathcal{C}} \phi_C^+(x).$$

Each player with an ideal point to the left of x proposes the left endpoint of the social acceptance set, and a player with an ideal point to the right of x proposes the right endpoint of the social acceptance set. Any player whose ideal point is x makes a proposal $dx/dt = 0$. Thus the average proposed speed is

$$\psi(x) = \mu_-(x)\phi_-(x) + \mu_+(x)\phi_+(x).$$

In fact, the function ψ is related to the function ξ through the equation $\varphi_- \varphi_+ \psi = \ln(\delta)\xi$. To see this, notice that

$$\phi_C^-(x) = \ln(\delta) / \max_{i \in C} \{u'_i(x)/u_i(x), 0\} \text{ and } \phi_C^+(x) = \ln(\delta) / \min_{i \in C} \{u'_i(x)/u_i(x), 0\}$$

$$\phi_-(x) = \ln(\delta) / \varphi_+(x) \text{ and } \phi_+(x) = \ln(\delta) / \varphi_-(x).$$

Assuming that φ_- is negative and φ_+ is positive on the relevant interval, every zero of ξ is also a zero of ψ , and vice versa.

Now, it seems very intuitive that the bargaining outcome should be a zero of the function ψ . Indeed, a zero of the function ψ can be thought of being robust to a renegotiation, when taken as a status quo, in the sense that the average proposed incremental change is zero. Alternatively, one can view a zero of ψ as a rest point of a dynamics $dx/dt = \psi(x)$ that adjusts the outcome in the direction of the average proposal.

6 The proof of the main result

The central idea of the proof is to estimate the values of the characteristic function on the endpoints of the social acceptance set. Thus the main result of this section is Theorem 5 which establishes the inequalities $\xi(x_-) \geq 0$ and $0 \geq \xi(x_+)$, where x_- and x_+ are the endpoints of the social acceptance set in a δ -equilibrium for any $\delta < 1$. The inequalities imply immediately that any bargaining outcome is a generalized zero of the function ξ , and, since ξ is a decreasing function, that the bargaining outcome is unique. As a byproduct of Theorem 5 we also obtain an interesting fact that the bargaining outcome is an element of the social acceptance set of any δ -equilibrium for any $\delta \in [0, 1)$.

The following proposition is a decisive step of the proof.

Proposition 12 *Suppose $\delta \in [0, 1)$. Let $(x_\bullet, A_\bullet)$ be a δ -equilibrium inducing the expected payoffs $y_\bullet$, the acceptance set A_C for each coalition C and the social acceptance set $A = [x_-, x_+]$. Let*

$$d_- = \frac{x_- - E(x_\bullet)}{1 - \delta} \text{ and } d_+ = \frac{x_+ - E(x_\bullet)}{1 - \delta}.$$

If $x_- \in \text{Int}X$, then the following inequalities hold:

$$\begin{aligned} -1 &\leq \delta d_+ \varphi_-(x_-), \\ -1 &\geq \delta d_- \varphi_+(x_-). \end{aligned}$$

If $x_+ \in \text{Int}X$, then the following inequalities hold:

$$\begin{aligned} -1 &\leq \delta d_- \varphi_+(x_+), \\ -1 &\geq \delta d_+ \varphi_-(x_+). \end{aligned}$$

Proof. We prove the first inequality. Assume that $x_- \in \text{Int}X$.

Let t be an arbitrary player. For each $i \in N$ we have the inequalities

$$u_t(x_+) - u_t(x_i) \leq r_t(x_i)[x_+ - x_i] \leq r_t(x_-)[x_+ - x_i],$$

where the first inequality is a defining inequality for the subgradient of the function u_t , and the second inequality follows from the fact that the left derivative is a non-increasing function, so that $r_t(x_i) \leq r_t(x_-)$ for each $i \in N$ and the fact that $0 \leq x_+ - x_i$. (The right derivative at the point x_i exists if $x_i < x_+$, because in that case x_i is in the interior of X . If $x_i = x_+$, then the extreme left and right hand sides of the above inequality are both zero.) Taking the expected value with respect to i in both sides of the above inequality yields

$$u_t(x_+) - y_t \leq r_t(x_-)[x_+ - E(x_\bullet)].$$

Let $C_+ \in \mathcal{C}$ be such that $x_+ \in A_{C_+}$. Then $\delta y_t \leq u_t(x_+)$ for each $t \in C$. Let t be a player in the coalition C . Therefore,

$$\begin{aligned} 0 &\leq u_t(x_+) - \delta y_t = \\ &= (1 - \delta)u_t(x_+) + \delta[u_t(x_+) - y_t] \leq \\ &\leq (1 - \delta)u_t(x_+) + \delta r_t(x_-)[x_+ - E(x_\bullet)]. \end{aligned}$$

Dividing by $(1 - \delta)$ yields

$$0 \leq u_t(x_+) + \delta r_t(x_-)d_+.$$

Now, either $\bar{x}_t \leq x_-$ or $x_- < \bar{x}_t$. If the former, then the function u_t is non-increasing on the interval $[x_-, x_+]$, in particular $u_t(x_+) \leq u_t(x_-)$. If the latter, then $0 \leq r_t(x_-)$. In either case we obtain the inequality

$$0 \leq u_t(x_-) + \delta r_t(x_-)d_+.$$

Rearranging yields the inequality

$$-1 \leq \delta d_+ r_t(x_-)/u_t(x_-).$$

Since t is arbitrary member of coalition C_+ , we have

$$-1 \leq \min_{t \in C_+} \{\delta d_+ r_t(x_-)/u_t(x_-), 0\}.$$

Therefore,

$$\begin{aligned}
-1 &\leq \max_{C \in \mathcal{C}} \min_{t \in C} \{\delta d_+ r_t(x_-)/u_t(x_-), 0\} = \\
&= \max_{C \in \mathcal{C}} \min_{t \in C} \{r_t(x_-)/u_t(x_-), 0\} \delta d_+ = \\
&= \varphi_-(x_-) \delta d_+.
\end{aligned}$$

The proof of the third inequality is a jiggled-jagged version of the proof of the first one. Assume that $x_+ \in \text{Int}X$. Let $C_- \in \mathcal{C}$ be such that $x_- \in A_{C_-}$. Replacing every instance where x_- occurs by x_+ and vice versa, one shows that

$$-1 \leq \delta d_- l_t(x_+)/u_t(x_+).$$

for each $t \in C_-$. Therefore,

$$-1 \leq \min_{t \in C_-} \{\delta d_- l_t(x_+)/u_t(x_+), 0\}.$$

Taking into account that $d_- \leq 0$, we then obtain

$$\begin{aligned}
-1 &\leq \max_{C \in \mathcal{C}} \min_{t \in C} \{\delta d_- l_t(x_+)/u_t(x_+), 0\} = \\
&= \min_{C \in \mathcal{C}} \max_{t \in C} \{l_t(x_+)/u_t(x_+), 0\} \delta d_- = \\
&= \varphi_+(x_+) \delta d_-.
\end{aligned}$$

We prove the fourth inequality. Assume that $x_+ \in \text{Int}X$. Let C be an arbitrary coalition from $\mathcal{C}$. First we argue that there exists a player $t \in C$ such that $u_t(x_+) \leq \delta y_t$. Suppose not. Then $\delta y_t < u_t(x_+)$ for each $t \in C$. Then the acceptance set A_C of coalition C contains an open neighborhood of the point x_+ , and so does the social acceptance set A . But x_+ is an interior point of X , so any open neighborhood of x_+ contains a point $x > x_+$. This contradicts the fact that x_+ is the right endpoint of the interval A .

Fix any player $t \in C$ such that $u_t(x_+) \leq \delta y_t$. For each $i \in N$ we have the inequality

$$u_t(x_i) - u_t(x_+) \leq r_t(x_+)[x_i - x_+],$$

Taking the expected values on both sides with respect to i yields

$$u_t(x_+) - y_t \geq r_t(x_+)[x_+ - E(x_\bullet)].$$

Therefore,

$$\begin{aligned}
0 &\geq u_t(x_+) - \delta y_t = \\
&= (1 - \delta)u_t(x_+) + \delta[u_t(x_+) - y_t] \geq \\
&\geq (1 - \delta)u_t(x_+) + \delta r_t(x_+)[x_+ - E(x_\bullet)].
\end{aligned}$$

Dividing by $1 - \delta$ gives

$$0 \geq u_t(x_+) + \delta r_t(x_+)d_+.$$

Rearranging yields the inequality

$$-1 \geq \delta d_+ r_t(x_+)/u_t(x_+).$$

Thus,

$$-1 \geq \min_{i \in C} \{\delta d_+ r_i(x_+)/u_i(x_+), 0\}.$$

Since C is an arbitrary element of the collection $\mathcal{C}$, we obtain

$$\begin{aligned} -1 &\geq \max_{C \in \mathcal{C}} \min_{i \in C} \{\delta d_+ r_i(x_+)/u_i(x_+), 0\} = \\ &= \max_{C \in \mathcal{C}} \min_{i \in C} \{r_i(x_+)/u_i(x_+), 0\} \delta d_+ = \\ &= \varphi_-(x_+) \delta d_+. \end{aligned}$$

The proof of the second inequality is similar to the proof of the fourth inequality. Assume that $x_- \in \text{Int}X$. Replacing every instance where x_- occurs by x_+ and vice versa, one shows that for each coalition $C \in \mathcal{C}$ there exists a player $t \in C$ such that

$$-1 \geq \delta d_- l_t(x_-)/u_t(x_-).$$

Thus,

$$-1 \geq \min_{i \in C} \{\delta d_- l_i(x_-)/u_i(x_-), 0\}.$$

Since C is an arbitrary element of the collection $\mathcal{C}$, and taking into account that $d_- \leq 0$, we then obtain

$$\begin{aligned} -1 &\geq \max_{C \in \mathcal{C}} \min_{i \in C} \{\delta d_- l_i(x_-)/u_i(x_-), 0\} = \\ &= \min_{C \in \mathcal{C}} \max_{i \in C} \{l_i(x_-)/u_i(x_-), 0\} \delta d_- = \\ &= \varphi_+(x_-) \delta d_-. \end{aligned}$$

This completes the proof. $\square$

Proposition 13 *Suppose $\delta \in [0, 1)$. Let $(x_\bullet, A_\bullet)$ be a δ -equilibrium inducing the social acceptance set $A = [x_-, x_+]$. Let*

$$d = \frac{x_+ - x_-}{1 - \delta}.$$

If $x_- \in \text{Int}X$, then, the following inequalities hold:

$$\begin{aligned} -1 &\leq \delta d \mu_-(x_-) \varphi_-(x_-), \\ 1 &\leq \delta d \mu_+(x_-) \varphi_+(x_-). \end{aligned}$$

If $x_+ \in \text{Int}X$, then the following inequalities hold:

$$\begin{aligned} 1 &\geq \delta d \mu_+(x_+) \varphi_+(x_+), \\ -1 &\geq \delta d \mu_-(x_+) \varphi_-(x_+). \end{aligned}$$

Proof. Let d_- and d_+ be as in Proposition 12. We use the following estimates of $E(x_\bullet)$:

$$\begin{aligned} E(x_\bullet) &\leq \mu(\bar{x}_i < x_-)x_- + \mu(x_- \leq \bar{x}_i)x_+, \\ E(x_\bullet) &\leq \mu(\bar{x}_i \leq x_-)x_- + \mu(x_- < \bar{x}_i)x_+, \\ E(x_\bullet) &\geq \mu(\bar{x}_i \leq x_+)x_- + \mu(x_+ < \bar{x}_i)x_+, \\ E(x_\bullet) &\geq \mu(\bar{x}_i < x_+)x_- + \mu(x_+ \leq \bar{x}_i)x_+, \end{aligned}$$

where $\mu(\bar{x}_i < x_-)$ stands for $\mu(\{i \in N | \bar{x}_i < x_-\})$, e.t.c. To derive the first of the estimates, let z_i be equal to x_- whenever $\bar{x}_i < x_-$ and x_+ otherwise. Then $x_i \leq z_i$ for each $i \in N$. Therefore, $E(x_\bullet) \leq E(z_\bullet)$. Other estimates are derived similarly.

Use the estimates for $E(x_\bullet)$ to obtain, respectively, the following inequalities:

$$\begin{aligned} x_+ - E(x_\bullet) &\geq x_+ - [\mu(\bar{x}_i < x_-)x_- + \mu(x_- \leq \bar{x}_i)x_+] = \mu(\bar{x}_i < x_-)[x_+ - x_-], \\ x_- - E(x_\bullet) &\geq x_- - [\mu(\bar{x}_i \leq x_-)x_- + \mu(x_- < \bar{x}_i)x_+] = -\mu(x_- < \bar{x}_i)[x_+ - x_-], \\ x_- - E(x_\bullet) &\leq x_- - [\mu(\bar{x}_i \leq x_+)x_- + \mu(x_+ < \bar{x}_i)x_+] = -\mu(x_+ < \bar{x}_i)[x_+ - x_-], \\ x_+ - E(x_\bullet) &\leq x_+ - [\mu(\bar{x}_i < x_+)x_- + \mu(x_+ \leq \bar{x}_i)x_+] = \mu(\bar{x}_i < x_+)[x_+ - x_-]. \end{aligned}$$

Dividing by $1 - \delta$ gives

$$\begin{aligned} d_+ &\geq \mu(\bar{x}_i < x_-)d, \\ d_- &\geq -\mu(x_- < \bar{x}_i)d, \\ d_- &\leq -\mu(x_+ < \bar{x}_i)d, \\ d_+ &\leq \mu(\bar{x}_i < x_+)d. \end{aligned}$$

Now each inequality of Proposition 13 is obtained by combining the respective inequality of Proposition 12 with the respective inequality above.

$$\begin{aligned} -1 &\leq \delta d_+ \varphi_-(x_-) \leq \delta \mu(\bar{x}_i < x_-) d \varphi_-(x_-) = \delta d \mu_-(x_-) \varphi_-(x_-), \\ -1 &\geq \delta d_- \varphi_+(x_-) \geq -\delta \mu(x_- < \bar{x}_i) d \varphi_+(x_-) = -\delta d \mu_+(x_-) \varphi_+(x_-), \\ -1 &\leq \delta d_- \varphi_+(x_+) \leq -\delta \mu(x_+ < \bar{x}_i) d \varphi_+(x_+) = -\delta d \mu_+(x_+) \varphi_+(x_+), \\ -1 &\geq \delta d_+ \varphi_-(x_+) \geq \delta \mu(\bar{x}_i < x_+) d \varphi_-(x_+) = \delta d \mu_-(x_+) \varphi_-(x_+). \end{aligned}$$

Of course, the first two inequalities are obtained under the assumption that $x_- \in \text{Int}X$ and the last two under the assumption that $x_+ \in \text{Int}X$. $\square$

Proposition 14 below strengthens Proposition 1 in Section 2. While Proposition 1 establishes that the social acceptance set is an interval containing the average proposal, the Proposition 14 states that it contains the average proposal together an open neighborhood.

Proposition 14 *Suppose $\delta \in [0, 1)$ and let $(x_\bullet, A_\bullet)$ be a δ -equilibrium inducing the expected payoffs $y_\bullet$ and the social acceptance set A . Then A contains an open neighborhood of the point $E(x_\bullet)$.*

Proof. Each individual acceptance set A_t contains the point $E(x_\bullet)$ by the concavity of the function u_t . Indeed,

$$u_t(E(x_\bullet)) \geq E(u_t(x_\bullet)) = y_t \geq \delta y_t.$$

Furthermore, if $y_t > 0$, the last inequality is strict, so the point $E(x_\bullet)$ is contained in the interior of A_t . If $y_t = 0$, then the set A_t contains the interval $(0, 1)$. Thus, in either case, the set A_t contains the point $E(x_\bullet)$ together with an open neighborhood. It follows that for each coalition C the acceptance set A_C contains an open neighborhood of the point $E(x_\bullet)$. The result follows. $\square$

Theorem 5 *Suppose $\delta \in [0, 1)$ and let $A = [x_-, x_+]$ be the social acceptance set in a δ -equilibrium of ω . Then $\xi(x_-) \geq 0 \geq \xi(x_+)$.*

Proof. By Proposition 14 the set A is non-degenerate. In particular, $x_- < 1$. If $0 < x_- < 1$, then $\xi(x_-) \geq 0$ as can be seen by adding up the first two inequalities of Proposition 13 and dividing by δd , d being non-zero again by Proposition 14. If $x_- = 0$, then $\xi(x_-) = +\infty$ by the definition of ξ . Similarly, $x_+ > 0$ by Proposition 14. If $0 < x_+ < 1$, then $\xi(x_+) \leq 0$ is obtained by adding up the last two inequalities of Proposition 13 and dividing by δd . And if $x_+ = 1$, then $\xi(x_+) = -\infty$ by the definition of ξ . $\square$

Proposition 15 *Each bargaining outcome of the world ω is a generalized zero point of the characteristic function ξ .*

Proof. Let x be a bargaining outcome of ω . By Definition 2, there exist sequences A^n , δ^n , and x^n such that A^n is a social acceptance set in a δ^n -equilibrium of ω , x^n is a point in A^n , the sequence δ^n converges to 1 and x^n converges to x . Let $A^n = [x_-^n, x_+^n]$. Replacing, if necessary, sequences by subsequences, assume that both x_-^n and x_+^n converge. Then Theorem 3 implies that both sequences converge to x . Theorem 5 implies that $\lim \xi(x_-^n) \geq 0 \geq \lim \xi(x_+^n)$, as desired. $\square$

The following theorem follows immediately from Theorem 5 and Proposition 15.

Theorem 6 *Let x be a bargaining outcome of the world ω . Suppose $\delta \in [0, 1)$ and let A be the social acceptance set in a δ -equilibrium of ω . Then x is an element of A .*

Appendix

We now provide a proof of Proposition 4. To prove the proposition we introduce a lexicographic order on the collection $\mathcal{C}$ of decisive coalitions induced by the order $>$ on the set N of players. Recall that $>$ denotes the response sequence in the game Γ .

Definition 5 Given a natural number k and a coalition $C \in \mathcal{C}$ let $\ell_k(C)$ be the k -th lowest element of the coalition C , in the order $>$ (we let $\ell_k(C) = \ell_{|C|}(C)$ for each $k > |C|$). Define an *lexicographic order* $>$ on the set $\mathcal{C}$ by letting $C_1 > C_2$ whenever there exists a natural number K such that $\ell_k(C_1) = \ell_k(C_2)$ for each $k = 1, \dots, K$ and $\ell_K(C_1) > \ell_K(C_2)$.

To compare the coalition C_1 to the coalition C_2 we first compare their minimal elements. If the lowest element of C_1 is bigger than the lowest element of C_2 , then $C_1 > C_2$. And if the lowest element of C_1 equals the lowest element of C_2 , we go on to compare the second lowest elements. If the second lowest element of C_1 is bigger than that of C_2 , then $C_1 > C_2$. If the second lowest elements of the two coalitions are also the same, we go on to compare the third lowest elements, and so on. For example, take $C_1 = \{1, 2, 4, 5\}$ and $C_2 = \{1, 2, 3, 6, 7\}$ and let the order on the set N be the natural order ($1 < 2 < \dots < 7$). Then $C_1 > C_2$.

Clearly, the lexicographic order $>$ is a strict total order on $\mathcal{C}$. We shall be interested in the maximal elements of $>$ on a subset $\mathcal{C}'$ of $\mathcal{C}$. For each $\mathcal{C}' \subset \mathcal{C}$ there is a unique maximal element of $>$ on $\mathcal{C}'$. For the rest of this section we assume that $\mathcal{C}$ only includes the minimal decisive coalitions, i.e. there are no C_1 and C_2 in $\mathcal{C}$ such that $C_1 \subset C_2$.

Proposition 16 *Let $\mathcal{C}' \subset \mathcal{C}$ and let $C \in \mathcal{C}'$ be the unique maximal element of $>$ on $\mathcal{C}'$. Then for each $t \in C$ and each $C' \in \mathcal{C}'$ either $t \in C'$ or there is a player $i \in C' \setminus C$ such that $t > i$.*

Proof. Let $t \in C$ and $C' \in \mathcal{C}'$. Since $C > C'$, there exists a K such that $\ell_k(C) = \ell_k(C')$ for each $k = 1, \dots, K$ and $\ell_K(C) > \ell_K(C')$. If $t \leq \ell_K(C)$, then $t = \ell_k(C)$ for some $k = 1, \dots, K$, in which case $t = \ell_k(C')$ and so is an element of C' .

Suppose now that $t > \ell_K(C)$ and let $i = \ell_K(C')$. We know that $t > i$, and that the player i is an element of C' . It remains to show that i is not an element of C . Suppose it is. Then $\ell_K(C) > i$, so there is some $J < K$ such that $\ell_J(C) = i$. But we know that $\ell_J(C) = \ell_J(C')$. Thus $\ell_K(C') = i = \ell_J(C')$. This is possible only if $|C'| \leq J$. But then each element of C' can be written as $\ell_j(C')$ for some $j = 1, \dots, J$, which equals the element $\ell_j(C)$ of the set C . Thus $C' \subset C$, a contradiction. $\square$

Proposition 4 *Let $\sigma = (x_\bullet, A_\bullet)$ be a joint stationary strategy in the game Γ inducing the acceptance set A_C for coalition C and the social acceptance set A . Let $x \in X$.*

- (a) *Suppose x is not an element of A . Then for each $C \in \mathcal{C}$ there is a player $t \in C$ such that x is not an element of the set A_t and player t 's vote is pivotal at some node h of the game Γ .*
- (b) *Suppose x is an element of A . Then there exists a $C \in \mathcal{C}$ such that x is an element of A_C and for each $t \in C$ player t 's vote is pivotal at some node h of the game Γ .*

Proof. We prove part (a). Fix a coalition $C'' \in \mathcal{C}$. We show C'' contains a player t whose vote is pivotal at some node h of the game Γ .

Given a coalition $C' \in \mathcal{C}$, let $C'_r = \{i \in C' | x \notin A_i\}$. The set C'_r is non-empty for each $C' \in \mathcal{C}$. Choose a coalition $C \in \mathcal{C}$ having the following properties: (a) $C_r \subset C''$ and (b) there is no coalition $C' \in \mathcal{C}$ such that C'_r is a strict subset of C_r . Clearly, such a coalition C exists: the set C_r is a minimal element of the collection $\{C'_r | C' \in \mathcal{C} : C'_r \subset C''\}$ of sets ordered by inclusion. Let t be the last player of the set C_r in the response order $>$. Notice that t is an element of C'' and that x is not an element of A_t .

Let h be any node of the game Γ where (a) player t reacts to the proposal x , (b) each player $i \in C$ preceding player t in the response order $>$ accepts the proposal x , (c) each player $i \in N \setminus C$ preceding player t in the response order $>$ rejects the proposal x . As follows from the considerations below, the vote of player t at the node h is decisive: the proposal x is socially accepted if and only if it is accepted by player t .

Notice that under the strategy σ_t player t rejects the proposal x . We now show that the proposal x is turned down on the path of play induced by the joint strategy σ in the subgame $\Gamma(h)$. Indeed, coalition C rejects the proposal x , because player $t \in C$ rejects x at the node h . Take any $C' \in \mathcal{C}$ other than C . By the choice of the coalition C , there is a player $i \in C'_r \setminus C_r$. If $i = t$ or $i > t$, player i rejects the proposal x along the path of play induced by the joint strategy σ , because $x \notin A_i$. And if $i < t$, then player i rejects the proposal x at the node h by condition (c). Notice that player i is not an element of the coalition C , for otherwise the inclusions $i \in C$ and $i \in C'_r$ would imply the inclusion $i \in C_r$, leading to a contradiction.

Consider now a strategy σ'_t for player t that coincides with σ_t on all nodes but h , where it accepts the proposal x . We show that the proposal x is accepted by the coalition C on the path of play induced by the joint strategy (σ'_t, σ_{-t}) in the subgame $\Gamma(h)$. Since t is chosen to be the last player of C_r in the response sequence, each member i of the coalition C following player t in the response order accepts the proposal x on the path of play induced by $(\bar{\sigma}_t, \sigma_{-t})$. And each member of coalition C preceding player t in the response order accepts x at the node h by condition (b).

We now prove part (b) of the proposition. As x is assumed an element of the set A , the collection $\mathcal{C}' = \{C \in \mathcal{C} | x \in A_C\}$ is non-empty. Now choose a coalition $C \in \mathcal{C}'$ having the following property: For each $t \in C$ and each $C' \in \mathcal{C}'$ either $t \in C'$ or there is a player $i \in C' \setminus C$ such that $t > i$. The existence of coalition C is guaranteed by Proposition 16. One must take C to be the maximal element of the set $\mathcal{C}'$ in the lexicographic order induced by the response sequence $>$, as in Definition 5. We show that each player t of C has a node h where player t 's vote is pivotal.

Take a player $t \in C$. Let h be any node where (a) player t has to react to the proposal x , (b) each player $i \in C$ preceding player t in the response order $>$ accepts x , and (c) each player $i \in N \setminus C$ preceding player t in the response order $>$ rejects x .

Notice that player t accepts the proposal x under the strategy σ_t . We now show that the proposal x is accepted by the coalition C on the path of play induced by the joint strategy σ in the subgame $\Gamma(h)$. Indeed, each player $i \in C$ preceding player t in the response order $>$ accepts x by the definition of the node h . And all players $i \in C$ following player t in the response order accept the proposal x along the path of play induced by the joint strategy σ , because $x \in A_i$.

Consider the strategy σ'_t that coincides with the strategy σ_t on all nodes but h where it rejects the proposal x . We show that the proposal x is socially rejected on the path of play induced by the joint strategy (σ'_t, σ_{-t}) in the subgame $\Gamma(h)$. To see this, let $C' \in \mathcal{C}$. Suppose first that $C' \in \mathcal{C}'$. Player t , if a member of coalition C' , rejects x at the node h by the definition of the strategy σ'_t . And if t is not a member of C' , then there is a player $i \in C' \setminus C$ such that $t > i$. Any such player i rejects the proposal x at the node h by condition (c). In either case the proposal x is rejected by at least one of the players of the coalition C' . Suppose now C' is not in $\mathcal{C}'$, so there exists a player $i \in C'$ such that x is not an element of A_i . If i follows t in the response order, then i rejects the proposal x along the path of play induced by the strategy σ . Suppose player i precedes player t in the response order. In this case, note that player i is not an element of the coalition C , because $x \in A_C$, so player i rejects the proposal x by the choice of the node h . $\square$

References

- [1] JEFFREY S. BANKS AND JOHN DUGGAN: A Bargaining Model of Collective Choice. *American Political Science Review*, **94**: 73–88, 2000.
- [2] KEN BINMORE, ARIEL RUBINSTEIN, AND ASHER WOLINSKY: The Nash Bargaining Solution in Economic Modelling. *The RAND Journal of Economics*, **17**: 176–188, 1986.
- [3] DANIEL CARDONA AND CLARA PONSATÍ: Bargaining One-Dimensional Social Choices. *Journal of Economic Theory*, doi: 10.1016/j.jet.2006.12.001, 2007.
- [4] SEOK-JU CHO AND JOHN DUGGAN: Uniqueness of Stationary Equilibria in a One-Dimensional Model of Bargaining. *Journal of Economic Theory*, **113**: 118–130, 2003.
- [5] HÜLYA ERASLAN: Uniqueness of Stationary Equilibrium Payoffs in the Baron–Ferejohn Model. *Journal of Economic Theory*, **103**: 11–30, 2002.
- [6] HÜLYA ERASLAN AND ANTONIO MERLO: Majority Rule in a Stochastic Model of Bargaining. *Journal of Economic Theory*, **103**: 31–48, 2002.
- [7] TASOS KALANDRAKIS: Regularity of pure strategy equilibrium points in a class of bargaining games. *Economic Theory*, **28**: 309–329, 2006.
- [8] KLAUS KULTTI AND HANNU VARTIAINEN: Von Neumann-Morgenstern stable sets, discounting, and Nash bargaining. *Journal of Economic Theory*, doi: 10.1016/j.jet.2006.11.006, 2007.
- [9] ANTONIO MERLO AND CHARLES WILSON: A Stochastic Model of Sequential Bargaining with Complete Information. *Econometrica*, **63**: 371–399, 1995.

- [10] ANTONIO MERLO AND CHARLES WILSON: Efficient delays in a stochastic model of bargaining. *Economic Theory*, **11**: 39–55, 1998.
- [11] TOSHIJI MIYAKAWA: Non-cooperative Foundation of n -Person Asymmetric Nash Bargaining Solution. Osaka University of Economics, working paper No. 2006–2.
- [12] ARKADI PREDTETCHINSKI: One-dimensional bargaining with unanimity rule. Maastricht University, Research Memorandum 07/011.