Venous oxygen saturation and oxygen transport in the newborn period : an experimental and clinical study

Citation for published version (APA):

van der Hoeven, M. A. H. B. M. (1999). *Venous oxygen saturation and oxygen transport in the newborn period : an experimental and clinical study*. [Doctoral Thesis, Maastricht University]. Universiteit Maastricht. <https://doi.org/10.26481/dis.19990623mh>

Document status and date:

Published: 01/01/1999

DOI:

[10.26481/dis.19990623mh](https://doi.org/10.26481/dis.19990623mh)

Document Version:

Publisher's PDF, also known as Version of record

Please check the document version of this publication:

- A submitted manuscript is the version of the article upon submission and before peer-review. There can be important differences between the submitted version and the official published version of record. People interested in the research are advised to contact the author for the final version of the publication, or visit the DOI to the publisher's website.
- The final author version and the galley proof are versions of the publication after peer review.
- The final published version features the final layout of the paper including the volume, issue and page numbers.

[Link to publication](#)

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal.

If the publication is distributed under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license above, please follow below link for the End User Agreement:

www.umlib.nl/taverne-license

Take down policy

If you believe that this document breaches copyright please contact us at:

repository@maastrichtuniversity.nl

providing details and we will investigate your claim.

Venous oxygen saturation and oxygen transport in the newborn period

An experimental and clinical study

Copyright © 1999 M.A.H.B.M. van der Hoeven

All rights reserved. No part of this publication may be reproduced, stored in a retrieval system, or transmitted, in any form or by any means, electronic, mechanical, photocopying, recording or otherwise without the prior written permission of the author.

For information: M.A.H.B.M. van der Hoeven, Hoge Kanaaldijk 13, 6212 XR Maastricht.

Venous oxygen saturation and oxygen transport in the newborn period. An experimental and clinical study.

Thesis University of Maastricht - with ref.- with summary in Dutch

ISBN 90-9012779-8

Subject headings: venous oxygen saturation; oxygen transport; newborn infants

Production: Datawyse | Universitaire Pers Maastricht

Venous oxygen saturation and oxygen transport in the newborn period

An experimental and clinical study

Proefschrift

ter verkrijging van de graad van doctor aan de Universiteit Maastricht,
op gezag van de Rector Magnificus, Prof.dr. A.C. Nieuwenhuijzen Kruseman,
volgens het besluit van het College van Decanen,
in het openbaar te verdedigen op woensdag 23 juni 1999 om 16.00 uur,

door

Markus Antonius Henricus Benedictus Maria van der Hoeven

Promotor

Prof.dr. C.E. Blanco

Co-promotor

Dr. W.J. Maertzdorf

Beoordelingscommissie

Prof.dr. R.A.M.G. Donckerwolcke (voorzitter)

Prof.dr. S. Bambang Oetomo (Rijksuniversiteit Groningen)

Prof.dr. H. Kuipers

Prof.dr. S. de Lange

Prof.dr. D. Tibboel (Erasmus Universiteit, Rotterdam)

10. 10. 1911

11. 10. 1911

12. 10. 1911

13. 10. 1911

14. 10. 1911

Abbreviations

Chapter 1	Introduction and aims of the study	11
Chapter 2	Oxygen transport: Review of the literature	17
Chapter 3	Mixed venous oxygen saturation (SvO ₂) and biochemical parameters of hypoxia during progressive hypoxemia in 10-14 day old piglets. <i>Pediatric Research</i> 1997;42:878-884	89
Chapter 4	Relationship between mixed venous oxygen saturation (SvO ₂) and markers of tissue oxygenation in progressive hypoxic hypoxia and in isovolemic anemic hypoxia in 8-12 days old piglets. <i>Critical Care Medicine: In Press</i>	105
Chapter 5	Feasibility and accuracy of a fiberoptic catheter for the measurement of venous oxygen saturation (SvO ₂) in newborn infants. <i>Acta Paediatrica</i> 1995;84:122-127	123
Chapter 6	Continuous central venous oxygen saturation (ScvO ₂) measurement by a fiberoptic catheter in newborn infants. <i>Archives Diseases in Childhood; Fetal Neonatal Ed.</i> 1996;74:F177-F188	135
Chapter 7	Central venous oxygen saturation (ScvO ₂) and derived variables at different ranges of arterial oxygenation in newborn infants with respiratory insufficiency. <i>Submitted</i>	147
Chapter 8	Summary and conclusions.	163
Chapter 9	Samenvatting	173
	Dankwoord	183
	Curriculum Vitae	184

Abbreviations	1
Chapter 1 Introduction and aims of the study	1
Chapter 2 Oxygen transport: Review of the literature	2
Chapter 3 Model studies: Oxygen saturation levels and biochemical parameters of hypoxia during postoperative hypoxemia in 10 14-day-old piglets	3
Chapter 4 Relationship between arterial PaO_2 and oxygen saturation (SpO ₂) and number of tissue oxygenation microcirculatory perfusion studies in newborn piglets	4
Chapter 5 PaO_2 and SpO ₂ and tissue oxygenation in a piglet model for the assessment of tissue oxygenation during hypoxemia	5
Chapter 6 Continuous arterial oxygen saturation (SpO ₂) measurement by a fiberoptic catheter: A validation study	6
Chapter 7 Effects of different ranges of arterial oxygenation on newborn infants with respiratory distress syndrome	7
Chapter 8 Arterial and venous oxygen saturation	8
Chapter 9 Summary	9
Index	10
References	11

Abbreviations

AH	anemic hypoxia
ARDS	acquired respiratory distress syndrome
ATP	adenosine triphosphate
caO ₂	total arterial oxygen content (ml/dl)
cvO ₂	mixed venous oxygen content (ml/dl)
ccvO ₂	central venous oxygen content (ml/dl)
c(a-v)O ₂	difference between arterial and venous oxygen content (ml/dl)
COHb	carboxyhemoglobin
DO ₂	oxygen delivery (ml.min ⁻¹ .kg ⁻¹)
DO _{2 crit}	critical value of oxygen delivery (ml.min ⁻¹ .kg ⁻¹)
2,3-DPG	2,3-diphosphoglycerate (mmol/l)
ECMO	extracorporeal membrane oxygenation
FiO ₂	fractional inspiratory oxygen concentration
F.O.E.	fractional oxygen extraction
FOE _{crit}	critical value of fractional oxygen extraction
Hb	hemoglobin (mmol/l)
HbF	fetal hemoglobin (mmol/l)
HH	hypoxic hypoxia
HHb	deoxyhemoglobin
IRDS	idiopathic respiratory distress syndrome
MetHb	methemoglobin
O ₂ Hb	oxyhemoglobin
P ₅₀	partial pressure of oxygen at 50 % oxygen saturation (kPa, torr)
P ₉₀	partial pressure of oxygen at 90 % oxygen saturation (kPa, torr)
P _{ao}	mean aortal blood pressure (mmHg)
P _{pulm}	mean pulmonary blood pressure (mmHg)
paCO ₂	arterial partial pressure of carbon dioxide (kPa, torr)
p _{Alv} O ₂	alveolar partial pressure of oxygen (kPa, torr)
paO ₂	arterial partial pressure of oxygen (kPa, torr)
pvO ₂	mixed venous partial pressure of oxygen (kPa, torr)
Qt	cardiac output (ml.min ⁻¹ .kg ⁻¹)
ROP	retinopathy of prematurity
SaO ₂	arterial oxygen saturation (%)
ScvO ₂	central venous oxygen saturation (%)
SvO ₂	mixed venous oxygen saturation (%)
SvO _{2 crit}	critical value of mixed venous oxygen saturation (%)
VO ₂	oxygen consumption (ml.min ⁻¹ .kg ⁻¹)

Chapter 1

General introduction and aims of the study

General introduction and aims of the study

The transport of oxygen and carbon dioxide to and from the cells is essential for cell integrity. Oxygen is needed as the final electron acceptor to obtain energy coupled to the formation of ATP in the mitochondria. Carbon dioxide is the end product of the aerobic degradation of carbohydrate and fat molecules. In a multicellular organism gas transport constitutes of three main components including 1) gas exchange with the environment (lung), 2) the actual transport from the gas exchange unit to and from the cells (blood and heart) and 3) the gas uptake by the cells from the blood and vice versa. The aim of medical intensive care is to preserve the human organism by maintaining cellular and organ integrity. For this, 1) gas exchange with the environment is supported by oxygen supply and/or mechanical ventilation. 2) Gas transport to the cells is sustained by improving heart function and increasing hemoglobin, the main transport medium for oxygen. 3) Oxygen uptake and cell integrity is monitored by biochemical parameters of hypoxia as lactate and pyruvate.

Venous oxygen saturation

The measurement of venous oxygen saturation in the pulmonary artery - mixed venous oxygen saturation (SvO_2) - is the average of all the oxygen saturations of the venous blood from all the different organs. Within this measurement all aspects of the oxygen transport (delivery: hemoglobin, SaO_2 and cardiac output) and tissue oxygen uptake are integrated. The fact that SvO_2 is an average value and that it is an integrated measurement of different aspects of oxygen transport is at the same time the strength and the weakness of this measurement.

The strength of the integrated measurement is the fact that one is informed about the tissue oxygen uptake together with the oxygen delivery to the tissues, in contrast to other routine measurements in intensive care as hemoglobin, paO_2 , SaO_2 and cardiac output, which refer only to oxygen delivery to the tissues.

The weakness of the SvO_2 measurement as an average value is that a normal value cannot exclude an oxygen restricted metabolism in one or more organs. Another difficulty referring to the integrated aspect of the SvO_2 value is how to interpret an abnormal value, since it is dependent on so many different variables. At last, in the case of sepsis or ARDS and "pathologic oxygen supply dependency" a disorder in tissue oxygen uptake has been assumed and questionable normal or high SvO_2 values were encountered. In spite of these drawbacks many adult intensive care units have adopted the continuous SvO_2 measurement as a standard routine care, since it provides at least a trend of the balance between oxygen delivery and tissue oxygen uptake.

Neonatal intensive care

In neonatal intensive care the mentioned methods to support the integrity of the organism are as just as important as in adult intensive care.

However, in neonatal medicine several additional problems are encountered: 1) the transition from intra-uterine to extra-uterine environment associated with the switch of the gas exchange unit and with a concomitant increase of paO_2 . 2) Gas transport is affected by an increased concentration of fetal hemoglobin with a low amount of 2,3-DPG and

decreased p_{50} resulting in a left-warded shift of the oxygen dissociation curve. 3) Gas transport is affected by an increased cardiac output. 4) Oxygen uptake by the cells is increased in the neonatal period compared with adults.

Another aspect of oxygen transport in the neonatal period is the toxic effects of oxygen. Several long term consequences of prematurity as retinopathy of prematurity, bronchopulmonary dysplasia and necrotizing enterocolitis are associated with oxygen toxicity. Considering this, the monitoring of oxygen transport to the tissues has attracted a lot of attention. Oxygen transport is measured intermittently by samples taken of arterial blood to determine paO_2 and SaO_2 . The current continuously measured oxygen parameter is SaO_2 by pulse oximetry. However this parameter only assesses the delivery of oxygen to the tissues and not the tissue oxygen uptake or the oxygen extraction.

For this, it was suggested to measure venous oxygen saturation (SvO_2) to monitor the residual oxygen after tissue oxygen uptake^[1]. In neonatal medicine the knowledge of the real tissue oxygen uptake is extremely important, because both, hypoxia as well as hyperoxia have their damaging effects. An umbilical venous catheter positioned into the right atrium with a fiberoptic device should make it possible to measure central venous oxygen saturation ($ScvO_2$) continuously.

Five studies were performed to examine the possibilities and limitations of SvO_2 measurement in newborn infants and are reported in this thesis.

Outline of the thesis

Review of the literature. In this thesis four main aspects of oxygen transport will be surveyed. The review will focus on **I) physiology of oxygen transport**, subdivided into paragraphs describing definitions and terminology, oxygen cascade and venous oxygen saturation.

The consequences and compensatory mechanisms of **II) hypoxia** at cellular level and at systemic level are reviewed as far as this is related to methods used in our studies. For example, lactate and pyruvate were used in two studies to monitor tissue oxygenation. Consequently, the metabolism of lactate and pyruvate will be discussed in this review, in contrast to, for example, the ventilatory response during hypoxia, which was not related to our studies.

The assessment of the oxygen saturation in the blood as provided by **III) oximetry**, the cornerstone of the fiberoptic technique to measure SvO_2 , is reviewed featuring general principles, reliability and accuracy of fiberoptic and pulse oximetry.

IV) The management of improving oxygen transport will be described in two paragraphs containing on one hand the historical and current use of oxygen in neonatal medicine and on the other hand the debate on the DO_2 / VO_2 relationship and the strategy to optimize DO_2 and SvO_2 in adult intensive care.

Aims of the study: Since our interest in SvO_2 measurement primarily was motivated by the aspect of the tissue oxygen uptake of the integrated SvO_2 measurement we addressed the following questions:

- 1) What is the relation between tissue oxygenation and the SvO_2 value?
- 2) Is this relation between tissue oxygenation and SvO_2 different when oxygen delivery to the tissues is reduced by different methods?

To examine the clinical possibilities of the SvO_2 measurement in the newborn period, the following questions were addressed:

- 3) Is it possible to measure SvO_2 continuously in a newborn infant? Where? Are the SvO_2 values reliable and accurate?
- 4) What is the normal range of SvO_2 values in stable newborn infants breathing room air? What is the relation between SaO_2 and SvO_2 values?
- 5) What is the range of SvO_2 values during the acute phase of IRDS during mechanical ventilation and oxygen administration? What is the relation between SaO_2 and SvO_2 values and derived calculations such as venous admixture and fractional oxygen extraction? What is the value of ScvO_2 in clinical neonatal medicine.

1. Dudell G, Cornisch JD, Bartlett RH. What constitutes adequate oxygenation? *Pediatrics* 1990;85:39-41

Chapter 2

Page

Oxygen Transport: Review of the Literature

I.	Physiology of Oxygen Transport	17
II.	Hypoxia	33
III.	Monitoring Oxygen: Oximetry	49
IV.	Therapeutic Management to Improve Oxygen Transport	61
	References	71

Chapter 2

page

Oxygen Transport: Review of the Literature

I	Physiology of Oxygen Transport	19
II	Hypoxia	39
III	Monitoring Oxygen; Oximetry	49
IV	Therapeutic Management to improve Oxygen Transport	61
	References	75

Chapter 2 Oxygen Transport: Review of the Literature

I	Physiology of Oxygen Transport	19
II	Hypoxia	39
III	Monitoring Oxygen: Oximetry	49
IV	Therapeutic Management for Improved Oxygen Transport	61
	References	71

I Physiology of Oxygen Transport

1 Definitions and terminology

2 Atmosphere, oxygen and evolution

3 Oxygen Cascade

3.1 *Inspired pO_2*

3.2 *Alveolar pO_2*

3.3 *The alveolar/arterial pO_2 difference*

3.3.1 Venous admixture or shunt

3.3.2 Ventilation / perfusion (V/Q) scatter

3.4 *Oxygen transport by hemoglobin*

3.4.1 Adult Hemoglobin (HbA)

3.4.1.1 Structure and stereochemistry

3.4.1.2 Oxygen affinity

3.4.2 Hemoglobin switching

3.4.3 Fetal hemoglobin (HbF)

3.4.3.1 Structure

3.4.3.2 Oxygen affinity

3.4.4 Modulation of oxygen affinity

3.4.4.1 Clinical studies on modulating oxygen affinity

3.5 *Oxygen uptake by the tissues*

3.5.1 Tissue vascular geometry

3.5.2 Kinetics of the oxyhemoglobin dissociation

3.5.3 Intracapillary resistance to oxygen

3.5.4 Myoglobin

3.5.5 Constant and uniform oxygen consumption

3.6 *Oxygen consumption within the cells*

3.6.1 Glycolysis

3.6.2 Citric acid cycle

3.6.3 Oxidative phosphorylation

3.6.4 Fat and protein

3.7 *The mitochondrion: the end-organelle for oxygen consumption*

3.7.1 Anatomy and function

3.7.2 Uncoupling the electron transfer from the ATP synthesis.

3.7.3 The genome of mitochondria

3.7.4 The mitochondrion during perinatal development

4 Venous oxygen saturation

1 Definitions and terminology of quantities related to oxygen transport.

A set of definitions of the different quantities pertaining on the oxygen transport by the blood was described by several authors recently ^[1-5]. Some confusion has arisen about the definition of fundamental quantities as oxygen saturation and fractional oxyhemoglobin. In this chapter a set of definitions is given with the current symbols in relation to their measurement.

1.1 Oxygen tension (pO_2) is expressed in kPa or mmHg (Torr). The oxygen tension of the blood is defined as being equal to the partial pressure of oxygen in the gas mixture with which the blood is in equilibrium. Partial pressure is defined as the substance fraction of oxygen times the total pressure of the gas mixture ^[1]. The end-capillary pO_2 determines the oxygen flow to the tissues.

1.2 Oxygen content or concentration of total oxygen (ctO_2) is determined by the sum of the concentration of oxygen associated with hemoglobin ($cO_2(Hb)$) and the concentration of oxygen dissolved in the blood not associated with a protein ($cO_2(free)$).

$$ctO_2 = cO_2(Hb) + cO_2(free) \quad (1)$$

The freely dissolved oxygen is determined by the solubility constant for oxygen (α_{O_2}) and oxygen tension.

$$cO_2(free) = \alpha_{O_2} \cdot pO_2 \quad (2)$$

The solubility constant for oxygen (α_{O_2}) = 0.01 mmol/L.kPa or 0.0031 ml/dl.mmHg or 0.0225 ml/dl.kPa

1.3 Concentration of oxygen associated with hemoglobin ($cO_2(Hb)$) is determined by the concentration hemoglobin capable of binding oxygen and by oxygen saturation (SO_2).

1.4 Concentration of hemoglobin capable of binding oxygen involves only hemoglobin bound to oxygen (oxyhemoglobin, cO_2Hb) or that which can bind oxygen (deoxyhemoglobin, $cHHb$).

1.5 Oxygen saturation (SO_2) is simply the fraction or percentage of hemoglobin bound to oxygen compared with the hemoglobin capable of binding. This quantity can be measured by a spectrophotometric measurement of oxyhemoglobin (cO_2Hb) and deoxyhemoglobin ($cHHb$) ^[2].

$$SO_2 = cO_2Hb / (cO_2Hb + cHHb) \quad (3)$$

1.6 Concentration of total hemoglobin ($ctHb$) is the sum of all hemoglobin derivatives as oxyhemoglobin, cO_2Hb , deoxyhemoglobin, $cHHb$, and those derivatives, not capable of binding oxygen, the dyshemoglobins ($cdysHb$) including carboxyhemoglobin ($cCOHb$), methemoglobin ($cMetHb$), and other unnamed non-oxygen binding hemoglobin derivatives.

$$ctHb = cO_2Hb + cHHb + cCOHb + cMetHb + cdysHb \quad (4)$$

Since oxygen saturation (SO_2) involves only oxyhemoglobin and deoxyhemoglobin it can be measured spectrophotometrically without measuring carboxyhemoglobin or methemoglobin by simple spectrophotometers or by pulse oximeters and fiberoptic reflectance spectroscopy. In multiwavelengths spectrophotometers several dyshemoglobins can be determined. Oxygen saturation (SO_2) can then be determined by application of the subsequent equation ^[1,2,3]:

$$SO_2 = cO_2Hb / (ctHb - cCOHb - cMetHb - cdysHb) \quad (5)$$

1.7 **Fractional Oxyhemoglobin (FO_2Hb)** is the fraction of oxyhemoglobin compared with the concentration of total hemoglobin. This quantity is essentially different from saturation as can be seen when the two equations (5) and (6) are compared.

$$FO_2Hb = cO_2Hb / ctHb \quad (6)$$

These two quantities have been frequently confused, since in many subjects there is no or minimal dyshemoglobin present. However, when substantial amounts of dyshemoglobin are present, the results of oximeters, measuring only cO_2Hb and $cHHb$ as pulse oximeters and fiberoptic reflectance oximeters, are different to the results of multiwavelengths spectrophotometers. This discrepancy has led to the introduction of new terms as functional and fractional oxygen saturation ^[4,7]. Since the terminology and symbols used meet the international standards and recommendations these new terms were called unnecessary and self-contradictory ^[1,2,4]. The use of a special symbol for SO_2 as measured by the pulse oximeter (SpO_2) and the proposal to introduce a symbol for the oxygen saturation value given by a Hewlett Packard, for example, as $S_{hp}O_2$ or that of Oximetrix as S_xO_2 was rejected. The symbol for a measured value must be independent of the (limitations of the) method and of the manufacturer ^[5].

The fraction of each hemoglobin derivate can be defined in the same way as for fractional oxyhemoglobin.

$$\text{Fractional deoxyhemoglobin } FHHb = cHHb / ctHb \quad (7)$$

$$\text{Fractional carboxyhemoglobin } FCOHb = cCOHb / ctHb \quad (8)$$

$$\text{Fractional methemoglobin } FCOHb = cMetHb / ctHb \quad (9)$$

To illustrate the different use of the terms of saturation and fractions two definitions of the total concentrations of oxygen are given:

Firstly as the total concentration of oxygen as defined in equation (1)

$$ctO_2 = cO_2(Hb) + cO_2(\text{free}) \quad (1)$$

which may be rewritten in terms of saturation:

$$ctO_2 = ((cO_2Hb + cHHb) \cdot 1.39 \cdot SO_2) + (pO_2 \cdot \alpha O_2) \quad (10)$$

where 1.39 is an oxygen-hemoglobin binding constant based on molecular level studies of hemoglobin expressed as the maximal amount of oxygen (ml) can be bound to hemoglobin (gram). And secondly, as equation (7), which may also be expressed in terms of fractional

concentrations ^[2,3]:

$$ctO_2 = (ctHb \cdot 1.39 \cdot FO_2Hb) + (pO_2 \cdot \alpha O_2) \quad (11)$$

2 Atmosphere, Oxygen and Evolution

A discussion about oxygen transport should start with some remarks on the origin of oxygen in the atmosphere and evolution to aerobic metabolism.

The earth was formed 4.6 billion years ago by accretion of solid material condensed from the solar nebula ^[8,9]. Two crucial factors have contributed to the origin of an oxidizing atmosphere around the earth: The intermediate gravitational field, resulting both in a retention of heavier gases as oxygen, carbon dioxide and nitrogen and in an escape of lighter gases as hydrogen and helium. Smaller planets as Mercury have a weak gravitational field with no atmosphere at all, whereas large planets as Jupiter and Saturn with a strong gravitational field retain all gases including hydrogen and helium ensuring a reducing atmosphere. The other factor is the temperature that enables water to be in the liquid form.

A primitive atmosphere was formed by the decomposition of inner earth material and brought into the atmosphere by active volcanos. The water vapor in the primitive atmosphere rained out to form the oceans, when the surface of the earth had cooled sufficiently. It has been shown, that under the influence of electric discharges and ultraviolet radiation in the then existing environment a number of organic molecules could originate. Encapsulation by lipid membranes of these organic compounds has been achieved in the laboratory situation.

Although the next step in the evolution could not be reproduced, it seems likely that generated polymers of nucleotides (RNA) eventually were able to catalyze their own replication and to synthesize polypeptides ^[10]. These processes led 3.5 billion years ago to the formation of living cells. Early cells produced adenosine tri-phosphate (ATP) by the conversion of reduced compounds into organic acids as lactic, formic and succinic acid. (ATP has the unique property to store energy in high "energy bonds", which can be used everywhere in the cell.) To protect the cells from the acidic environment cells tended to develop membrane-bound H^+ pumps driven by electron transfer or / and ATP hydrolysis ^[11]. Reversal of the ATP driven H^+ pump made them function as an ATP-synthase. By combining parts of the electron transfer chains photosynthetic bacteria could employ the radiant energy of the sun to use H_2O as an electron donor to convert N_2 and CO_2 to amino acids and carbohydrates and oxygen was released to the atmosphere as a waste product. The increase of oxygen in the atmosphere was very slowly, and has been dated 2 billion years ago. By geological evidence it is assumed that the current level of atmospheric oxygen was attained between 1.5 to 0.5 billion years ago. Because of the availability of oxygen some bacteria developed the ability to use oxygen as the final electron acceptor and lost the capability to photosynthesis. Mitochondria first arose when such aerobic bacteria were endocytosed by primitive eucaryotic cells. The more favourable energy conversion supported eucaryotic cells with mitochondria to express their genetic information in different ways and to evolve to multi-cellular organisms.

3 The Oxygen Cascade ^[12]

Oxygen is vital for the effective generation of ATP. Generally it is assumed that the critical oxygen level in the mitochondrion is 0.1-0.3 kPa ^[12,13]. Consequently, an oxygen gradient exists between the air and the mitochondrion: the oxygen cascade. The different steps of this cascade will be discussed, subsequently.

3.1 Inspired pO₂ (p_iO₂)

The normal value of the concentration of atmospheric oxygen is 20.94 % or 0.2094 fractional concentration. When gas is humidified with the inhalation through the respiratory tract, the oxygen will be diluted and the pO₂ will be reduced. Therefore, the inspiratory pO₂ (P_iO₂) at a body temperature of 37° C and a barometric pressure of 760 mmHg is:

$$p_iO_2 = 0.2094 \cdot (760 - 47) = 149 \text{ mm Hg}$$

$$\text{or } p_iO_2 = 0.2094 \cdot (101.3 - 6.3) = 19.9 \text{ kPa}$$

3.2 Alveolar pO₂ (p_{Alv}O₂) ^[12,14]

Alveolar oxygen tension is the result of the oxygen delivered by the alveolar ventilation and the oxygen uptake by pulmonary blood and oxygen consumption. The relation between p_{Alv}O₂ and alveolar ventilation is described by a hyperbola with the horizontal asymptote equivalent to the p_iO₂. By increasing the inspiratory oxygen concentration (FiO₂) the relation is changed in so far that the horizontal asymptote is increased. An increase in VO₂ will also modify the relation between p_{Alv}O₂ and alveolar ventilation: as VO₂ increases, p_{Alv}O₂ is reduced even when alveolar ventilation increases.

The simplified form of the alveolar oxygen tension equation is:

$$p_{Alv}O_2 = p_iO_2 - p_{Alv}CO_2 / RQ + F$$

RQ is the respiratory exchange ratio, that corrects for the fact that usually less carbon-dioxide is produced than oxygen is consumed. F is a correction factor for the differences in volume between inspired and expired volume of gas. The p_{Alv}O₂ in a patient, whose RQ is 0.8, is 108 mmHg.

An inverse relationship exists between p_{Alv}CO₂ and alveolar ventilation. When alveolar ventilation decreases, p_{Alv}CO₂ increases proportionally. It is generally accepted that p_{Alv}CO₂ differs very little from the partial pressure of CO₂ in the arterial blood (p_aCO₂). This is why p_aCO₂ is accepted as an indicator for hypo- or hyperventilation ^[14].

3.3 The alveolar/arterial pO₂ difference

The next step in the oxygen cascade is of great importance and plays a major role in the pathophysiology of many respiratory disorders. Normally in a healthy adult the alveolar/arterial pO₂ difference will not exceed 2 kPa (15 mmHg) ^[12]. Two main causes of an increased alveolar/arterial pO₂ difference will be discussed.

3.3.1 Venous admixture or shunt ^[15]

Venous admixture is defined as the degree of admixture of mixed venous blood with pulmonary end-capillary blood, which would be required to produce the observed difference between the arterial and the pulmonary end-capillary pO₂. (usually taken to

equal ideal alveolar $p_{A_{lv}O_2}$). Venous admixture is a calculated amount of blood to produce the observed arterial pO_2 and not the actual amount of venous blood, which mingles with arterial blood. Venous admixture is thus a convenient index, but defines neither the precise volume nor the anatomical pathway of the shunt. There are different pathways of the shunt venous admixture:

- a) Venae cordis minimae (thebesian veins) are small veins of the left heart which drain directly into the left chamber. The venous blood has a low oxygen content and produces a substantial decrease of the arterial pO_2 .
- b) bronchial veins: venous drainage of the lung pass directly to the pulmonary veins; The flow is 1% or less from the cardiac output. However, it can increase in bronchiectasis, emphysema and coarctation of the aortae to values of 10 %.
- c) congenital heart disease
- d) pulmonary infection
- e) pulmonary oedema

In neonatal medicine venous admixture did not get as much attention as in adult intensive care medicine, but recently the values of venous admixture were studied in a group of preterm mechanically ventilated newborn infants ^[16]. A range between 10 % and 40 % was defined in this group of patients.

3.3.2 Ventilation / perfusion (V/Q) scatter

There are two important reasons why a scatter of V/Q values results in an increased alveolar / arterial pO_2 difference.

- a) Increased flow of blood through lung regions with a low V/Q ratio and with a low p_aO_2 cannot be balanced by the decreased amount of blood with a high p_aO_2 perfusing lung regions with a high V/Q ratio.
- b) The p_aO_2 of the blood that has passed through the low V/Q regions is disproportional low by the position of the oxygen dissociation curve in comparison to the increase of the p_aO_2 of the blood passed through the regions of the lung with a high V/Q ratio.

In neonatal medicine the scatter of V/Q ratios has been discussed by Krauss ^[17] and was determined to play a role in the first hours after birth and in respiratory disorders as meconium aspiration and transient tachypnoea of the newborn.

3.4 Oxygen transport by hemoglobin

Oxygen transport to and from the tissues is dependent on the oxygen affinity of the hemoglobin molecule. Hemoglobin is made up of four polypeptide globin chains each holding one heme. Heme consists of a porphyrin ring with an atom of a two valency ferrous iron at its centre, that is capable to form a reversible bond with oxygen without being irreversibly oxidized to the three valency ferric iron ^[18].

3.4.1 Adult Hemoglobin (HbA)

3.4.1.1 Structure and stereochemistry

Hemoglobin is a tetramer containing four polypeptide globin chains, two α and two β chains, joined by variable contacts. The contacts between α_1 and β_1 and between α_2 and β_2 are the most extensive and remain the same in oxy- and deoxyhemoglobin. However, the contacts between α_1 and β_2 and between α_2 and β_1 are changed during oxygenation. These changes are associated with a shift in the quaternary structure of hemoglobin from the tense (T) state (low-oxygen affinity, "deoxyhemoglobin") to the

relaxed (R) state (high-oxygen affinity, "oxyhemoglobin"). The T state is stabilized by electrostatic bounds between the subunits of the molecule^[19]. When oxygen is bound to the ferrous iron of the heme, the iron is pulled into the plane of the porphyrin ring. This movement is transmitted to the globin subunit and weakens these electrostatic bounds that stabilized the T state. This initiates the quaternary structure change from T state to R state.

Due to these conformational changes needed in the T state and due to the orientation of the heme in the globin the binding of oxygen in the T state is energetically unfavourable in contrast to oxygen binding in the R state. The R state has consequently an affinity for oxygen 100 fold higher than that of the T state. The switch from T state to R state normally takes place when the second or the third molecule of oxygen is bound.

As oxygen binds to hemoglobin, the equilibrium between the T and R states of hemoglobin shifts to a higher concentration of the R state and this accounts for the sigmoid shape of the oxyhemoglobin dissociation curve^[18,19].

3.4.1.2 Oxygen affinity

The affinity of oxygen for hemoglobin is illustrated by the position of the oxyhemoglobin dissociation curve: Oxygen saturation plotted against oxygen tension yields the characteristic sigmoid shape of the oxyhemoglobin dissociation curve. The affinity of oxygen for hemoglobin is reflected by the p_{50} , which expresses the oxygen tension at which 50 % of the hemoglobin is bound to oxygen. The p_{50} can be estimated with an acceptable accuracy from pH, pO_2 and oxygen saturation paired values^[20-22]. A significant error could be induced in the estimation of the p_{50} , particularly if p_{50} is calculated from an oxygen saturation value at the extremes of the oxyhemoglobin dissociation curve^[22]. The position of the oxyhemoglobin dissociation curve is affected by several factors. For example, a shift in the oxyhemoglobin dissociation curve to the right (increasing p_{50} and lowering oxygen affinity) may be caused by a decrease in pH, an increase in pCO_2 , an increase in 2,3-diphosphoglycerate (2,3-DPG) and an increase in temperature^[23].

a) pH (Bohr effect) and pCO_2

Hydrogen ions lower the oxygen affinity of the T structure and slow down the transition from the T to R structure of the hemoglobin molecule. This effect is achieved by fortifying salt bridges of the T structure. The Bohr effect once more is affected by other factors as pCO_2 , 2,3-DPG and temperature in an interrelated way. CO_2 reacts with uncharged amino-groups of both α and β globins to form carbamate. Haldane effect accounts for the increased binding of CO_2 to deoxyhemoglobin.

b) 2,3-diphosphoglycerate (2,3 DPG)

2,3-DPG is a metabolic intermediate of the anaerobic glycolysis. 2,3-DPG decreases oxygen affinity and increases p_{50} . This effect of 2,3-DPG is achieved by two different mechanisms: 2,3-DPG forms salt bridges between the two β globins of deoxyhemoglobin and consequently stabilizes the T state and 2,3-DPG is a highly charged anion and due to the impermeability of the erythrocyte cell membrane induces a decrease of the intracellular pH and consequently lower oxygen affinity by the Bohr effect.

c) Temperature

Perutz^[18] described the effect of temperature on the binding of hemoglobin and oxygen. The reaction of the heme-iron complex with oxygen is exothermic (gives off heat). Consequently, a rise in temperature hinders the reaction between heme and oxygen and shifts the oxyhemoglobin dissociation curve to the right. High temperatures decrease the affinity of hemoglobin and oxygen, whereas cold environmental temperatures increase oxygen hemoglobin affinity. This effect can also be demonstrated in hemoglobin monomers and is in fact impaired in the tetrameres of hemoglobin by increased heat absorption by heme-heme interaction. Adaptations of the hemoglobin molecule to different environmental

temperatures have been described in different animals and species^[24]. For example, in arctic ruminants, such as reindeer a normal oxygen supply can be delivered to the peripheral tissues even at very low temperatures due to a low-temperature sensitivity of the hemoglobin molecule^[25]. It was demonstrated by Condo et al.^[26] that the low temperature sensitivity of porcine blood occurred only in the presence of 2,3-DPG probably due to the interaction between 2,3-DPG, Cl⁻ and the hemoglobin molecule.

d) Other factors

Other factors as carboxyhemoglobin level, inorganic phosphate, chloride and bicarbonate ions, metabolic and endogenous states as activity, dormancy and developmental stage affect the function of hemoglobin and have been described in numerous different animals and species^[24,27,28].

3.4.2 Hemoglobin switching

Hemoglobin production in humans is characterized by two major "switches" during development in the hemoglobin composition of red cells. During the first 3 months of gestation, human red cells contain embryonic hemoglobins, whereas during the last 6 months of gestation red cells contain predominantly fetal hemoglobin. The transition from fetal to adult hemoglobin occurs during the perinatal period and first year of life^[29].

The functional genes for ϵ , γ and β globins of the embryonic, fetal and adult hemoglobin, respectively, are situated on the human β gene cluster on chromosome 11p. The globin genes are arranged according to the order of their expression during ontogeny. It is not clear how their differential expression during development is regulated. The cellular factors involved in the process of gene switching are transcription factors, globin locus control region, cis-acting sequences, trans acting factors and chromosomal gene order^[29,30]. The role of this last factor -the chromosomal gene order- was studied by Peterson et al.^[31]. The authors concluded that gene order played only a minor role in the globin gene switching.

It seems that the developmental maturity of the fetus or a developmental clock intrinsic to the cells determines the progression of hemoglobin switching^[29]. Several observations support this concept. Hybrids of fetal erythroblasts or transplanted fetal stem cells switch according to the age of the fetus from which the erythroblasts were obtained. In addition, the administration to very low birth weight infants of recombinant erythropoietin did not delay nor increase the developmentally programmed decline of hemoglobin F synthesis^[32].

However, environmental factors can modulate the developmental clock of switching. Fetal hypoxemia^[33] during late gestation increased the HbF synthesis greater than can be expected for gestational age. In infants with congenital cyanotic heart disease^[34], bronchopulmonary dysplasia^[35] or severe anemia of prematurity^[36] higher levels of hemoglobin, total HbF and HbF synthesis were demonstrated. Furthermore, Bard et al.^[37] showed that the level of HbF seemed to be dependent on glucose level. Hyperglycemia accomplishes this effect by increased fetal oxidative metabolism and increased oxygen consumption resulting in fetal hypoxemia. It was demonstrated that the hyperglycemic fetus synthesized more fetal hemoglobin than expected for the period of fetal development. In adult life fetal hemoglobin continues to be produced in several inherited and acquired conditions^[29].

3.4.3 Fetal hemoglobin (HbF)

3.4.3.1 Structure

The structure of fetal hemoglobin (HbF) is very similar to that of HbA, being a tetramer with two α and two non α globins. The non α chains do not differ in the number but in the sequence of amino acids from β chains and are designated γ chains. In the γ chain a difference has been found, that either alanine (γ^{ala}) or glycine (γ^{gly}) may be present at position 136, without any functional significance. Bard et al. [38] investigated the change in γ^{gly} to γ^{ala} chain ratio in the switch over of HbF to adult HbA. The authors concluded that during initial decrease of HbF synthesis in preterm infants the ratio of γ^{gly} and γ^{ala} ratio did not change. But after a postconceptual age of 44 wk when the HbF synthesis decreased below $42 \pm 15\%$ the ratio became more variable and decreased. The relative amount of γ^{gly} globins in fetal life is 70 %, whereas in adult life the small amount HbF synthesized contains around 40 % γ^{gly} globins.

A separate component of HbF, which is designated HbF₁, was found in 20 % of the total HbF. In HbF₁ the glycine residues in either half or all of the γ chains are bound with an acetyl group. 2,3 DPG does not interact at all with HbF₁ and therefore, HbF₁ has a very high affinity for oxygen.

3.4.3.2 Oxygen affinity

Fetal blood has a higher oxygen affinity than adult blood. However, this difference is disappeared, when HbF and HbA are isolated and stripped of phosphate. Therefore, the different interaction with 2,3-DPG is considered to be responsible for the difference in oxygen affinity between fetal and adult blood. The replacement in HbF of histidine by an uncharged serine reduces the binding sites with 2,3-DPG. In addition, a further reduction of 2,3-DPG binding sites in HbF₁ explains the lack of any effect of 2,3-DPG on oxygen affinity in HbF₁. The position of the oxyhemoglobin dissociation curve of fetal hemoglobin is affected by several factors as well.

a) pH (Bohr effect) and pCO₂ effect

The Bohr effect in fetal blood is not significantly different from that in adult blood. No separate pCO₂ effect can be discerned in fetal blood with normal 2,3-DPG levels. This means that acidosis immediately effectuates a reduction of oxygen affinity by the Bohr effect. But since acidosis induces a reduction of 2,3-DPG, changes in oxygen affinity will be reversed. The distinct effects of pH on the level of 2,3-DPG and of 2,3-DPG on oxygen affinity, together with the Bohr effect will balance one another both in fetal blood as in adult blood [23].

b) 2,3-diphosphoglycerate (2,3 DPG)

In the study of Wimberley [23] and confirmed by others the effect of 2,3-DPG on oxygen affinity (p_{50}) of HbF was 43 % compared with the effect of 2,3-DPG on the p_{50} of HbA. Half of this effect is due to the intracellular pH change within the erythrocyte, whereas the other half is due to a direct molecular interaction between hemoglobin and 2,3-DPG. 2,3-DPG levels at birth in preterm and full term infants are within the normal adult range. In a study on fresh cord blood of non-stressed, non-asphyxiated normal newborns from uncomplicated pregnancies from 24-42 weeks of gestation, it was concluded that the level of 2,3-DPG is rather constant, and that the decrease in fetal oxygen affinity as gestation progresses is related mainly to the reduction in amount of HbF [39,40]. In healthy preterm and full term infants 2,3-DPG increases postnatally in contrast with preterm infants with respiratory problems [23,39].

The level of 2,3-DPG is directly related to plasma pH. This means that acidosis is associated with low levels of 2,3-DPG and alkalosis with high levels of 2,3-DPG. 2,3-DPG

levels do not correlate with p_{aO_2} or ctO_2 . Furthermore, the level of 2,3-DPG in newborn lambs seemed to be dependent on glucose levels [40]. It was shown that during glucose infusion p_{50} and 2,3-DPG / hemoglobin ratio were significantly decreased in comparison with physiological saline infusion. One of the mechanisms to explain this effect is the inhibition or deactivation of the glucose-6- phosphatase with a concomitant suppression of 2,3-DPG synthesis. Delivoria-Papadopoulos et al. [41] described the postnatal changes in oxygen affinity in piglets. Since piglets only have one type of hemoglobin, the increasing values of p_{50} postnatally have been attributed to increased levels 2,3-DPG.

c) Temperature

The role of temperature in human fetal hemoglobin and of other modulators of oxygen affinity was investigated by Giardina et al. [42]. The overall heat production of oxygen binding was not different between HbA and HbF in the absence of 2,3-DPG. However, in the presence of 2,3-DPG and under physiological circumstances the heat production of oxygen binding of HbF was 60 % of HbA. This means that by virtue of the interplay between 2,3-DPG and temperature HbF has an increased oxygen affinity in comparison with HbA. In bovine fetal hemoglobin Clementi et al. [43] could demonstrate the lower overall heat of oxygenation in the presence 2,3-DPG. The authors speculated over the function of the decreased heat production on oxygen binding of HbF in the presence of 2,3-DPG. This difference between HbA and HbF could favour the heat exchange between the fetus and mother. After birth the decreased temperature sensitivity of HbF could facilitate a normal oxygen supply to the peripheral tissues.

3.4.4 Modulation of oxygen affinity

Modulating oxygen affinity to enable oxygen supply to the tissues have raised much interest especially in neonatal and intensive care medicine. The effects of modulating oxygen affinity on oxygenation have been studied in animal experiments during normoxic, anemic, circulatory hypoxic and hypoxic hypoxic conditions:

a) Normoxemic conditions

During normoxemic conditions the effects of low oxygen affinity (p_{50} : 29 mmHg) and high oxygen affinity (p_{50} : 18 mmHg) were examined in newborn lambs [44]. Oxygen consumption, oxygen extraction, arterial and venous oxygen saturation and cardiac output were not different between the low and high oxygen affinity group. Venous pO_2 was significantly higher in the low oxygen affinity group. The authors speculate that, since venous pO_2 is a good approximation of the mean tissue oxygen tension, lowering oxygen affinity could increase the mean tissue oxygen tension.

b) Anemic hypoxic conditions

The effect of low (p_{50} : 32 mmHg) and high (p_{50} : 19 mmHg) oxygen affinity at different degrees of anemia was compared in newborn lambs [45]. Venous pO_2 differed between the high and low oxygen affinity group, whereas arterial and venous oxygen content were not different between the high and low affinity group. Cardiac output was more increased in the low affinity group during severe anemia. In the low affinity group oxygen consumption remained stable whereas in the high affinity group oxygen consumption decreased during severe anemia. The data indicate that during severe anemia blood with a high p_{50} (low oxygen affinity) is more capable of adequately oxygenating tissues than that with a low p_{50} (high oxygen affinity), especially by increasing cardiac output. Again, mixed venous PO_2 was lower in the high affinity group during severe anemia [45].

c) Circulatory hypoxic conditions

Schumaker et al. [46] reported a study in mongrel dogs in which oxygen delivery was reduced by removing blood volume in a group with a normal p_{50} (28.4 torr) and a

reduced p_{50} (20.3 torr). The hypothesis was that a higher critical oxygen delivery was found in the group with a reduced p_{50} because of an increased oxygen affinity. However, critical DO_2 and critical oxygen extraction ratio were similar in both groups. As oxygen delivery was reduced below the critical level the extraction ratio continued to increase, which was significantly higher among normal p_{50} than among low p_{50} . Several mechanisms were mentioned to explain this finding: Redistribution of blood flow in the group with low p_{50} or the convergence of the different oxyhemoglobin dissociation curves at critically low capillary $p\text{O}_2$ could explain the similar critical oxygen delivery in both groups to some extent. Furthermore, the authors question whether the increased oxygen affinity in the group with a low p_{50} is an important determinant of oxygen unloading and whether other mechanisms as rheological properties of blood at very low cardiac output and vasoconstrictor tone may begin to limit oxygen consumption before blood-tissue oxygen diffusion becomes significant. Since the extraction ratio continued to increase in the group with a normal p_{50} , diffusion limitation seemed to play a role however, only at deliveries at which VO_2 is already depressed.

d) hypoxic hypoxic conditions

Lister^[47] examined the effects of high (p_{50} : 27 mmHg) and low (p_{50} : 35 mmHg) oxygen affinity in newborn lambs during hypoxic hypoxia. During hypoxic hypoxia ($\text{FiO}_2 = 0.1$) oxygen delivery and consumption decreased. After exchange transfusion during hypoxia oxygen delivery and oxygen consumption increased to near normoxic levels. These changes were concomitant with an increase of cardiac output and stroke volume.

3.4.4.1 Clinical studies on modulation oxygen affinity

In an observational study in 20 critically ill patients Myburg et al.^[48] found a reduced p_{50} . The value of p_{50} was strongly correlated with pH and base excess. In the discussion the authors speculate that a decreased level of 2,3-diphosphoglycerate due to a chronic acidosis or hypophosphatemia played an important role in the reduction of p_{50} in critically ill patients.

Besides this observational study two clinical trials have been performed in premature infants to determine whether an exchange transfusion with fresh adult blood could provide clinical benefits. Oski^[49] reported and reviewed these clinical trials employing exchange transfusions as an adjunct of conventional therapy. A decreased mortality was reported in those infants in whom an exchange transfusion was employed. The combined results of these trials reported an increased survival rate in the exchange group (58.2 %) vs. (25.3 %). The exchange transfusion resulted in an increase in 2,3-DPG, a decrease of fetal hemoglobin and an increase in p_{50} . However, these changes could not be correlated with the eventual outcome of the infants who received an exchange transfusion. Additionally, within 7 hours after exchange transfusion an unanticipated rise of arterial oxygen tension was found. This response could not be attributed to an increased p_{50} , but more to an improved pulmonary ventilation or pulmonary circulation. In both trials it was reported that central venous $p\text{O}_2$ was increased after exchange transfusion concomitant with the findings in animal trials. It seemed that an increased venous $p\text{O}_2$ as a consequence of a high p_{50} induced by an exchange transfusion could be one of the factors that enhance pulmonary perfusion and possibly ameliorate cardiac output.

Theoretically, the advantage of shifting the oxyhemoglobin curve is dependent on arterial $p\text{O}_2$ and venous $p\text{O}_2$ ^[50]. For example, a right shifted curve improves the maximal possible tissue oxygen uptake at any arterial $p\text{O}_2$, when venous $p\text{O}_2$ is 40 mmHg. However, when venous $p\text{O}_2$ is 10 mmHg the maximal tissue oxygen uptake is improved in the presence of hypoxemia by a left ward shifted curve, since oxygen loading in the lung and systemic oxygen transport is enhanced. Because uptake and $p_v\text{O}_2$ differ considerably from

organ to organ and within organs the benefits or drawbacks of a shift in the oxyhemoglobin curve cannot be predicted a priori.

With the theoretical possibility that shifting the oxyhemoglobin curve of the newborn to the right oxygen extraction may decrease in certain conditions, together with the risks of the exchange procedure itself and with the data from clinical trials where many other factors may have caused the increased survival, it is concluded that exchange transfusion at present cannot be recommended in the regular care of newborn infants.

3.5 Oxygen uptake by the tissues

Whereas the oxygen transport from the lungs to the tissues is accomplished by convective or bulk flow, oxygen uptake by the tissues is accomplished by diffusion of oxygen to the tissues. The process of diffusion can be described as the movement of oxygen across the cytosol and membrane of the erythrocyte, through the plasma into the tissues according to a concentration gradient^[51]. Several models have been proposed to characterize this process of oxygen uptake by the tissues.

The Krogh model is the eldest and most simplified model of oxygen diffusion to the tissues^[52,53]. This model describes oxygen transport from the centrally located capillary to the periphery of the tissues surrounding this capillary as a cylinder. Kety extended the model accounting for the pO_2 changes along the longitudinal axis of the capillary. In the skeletal muscle with the well-ordered longitudinally located capillaries the Krogh-Kety model seemed to be a good approximation. However, in this simplified model more complex microvasculature was ignored and some additional assumptions were made:

- 1) simplified tissue geometry
- 2) the instantaneous release of oxygen by the erythrocytes
- 3) neglected changes in oxygen transport within the capillary
- 4) neglected effect of myoglobin
- 5) a constant and uniform oxygen consumption

3.5.1 Tissue vascular geometry

In tissues with a more complex vascular geometry the Krogh-Kety model is not sufficient to predict tissue oxygen values. Consequently, other more complex models (multi-capillary) were developed to reproduce a more reliable tissue oxygen profile. In studies on the in the cheek retractor muscle in the hamster the influence of the microvascular geometry on oxygen exchange was studied^[54,55]. It was demonstrated that the release of oxygen from a capillary seems to be strongly influenced by its proximity to arterioles and venules, that either supply oxygen to the capillaries or augment its release. These diffusive interactions might change under different hemodynamic conditions.

3.5.2 Kinetics of the oxyhemoglobin dissociation

Gutierrez addressed the question of the time required by the erythrocytes to release oxygen to the tissues^[51]. He developed a model that took into account the time component or the kinetics of the dissociation of oxyhemoglobin into oxygen and deoxyhemoglobin. The model formulated equations for different compartments along the capillary. The model predicts that plasma oxygen is not replenished completely due to the limit of the rate of the oxyhemoglobin dissociation. The plasma pO_2 lags behind the equilibrium PO_2 ; when the blood reaches the venous pool pO_2 will equilibrate again resulting in an increase of the venous PO_2 , compared with the capillary pO_2 . According to Gutierrez this discrepancy between plasma pO_2 and venous pO_2 will increase in the case of anemic and hypoxic hypoxia due to the slower rate of dissociation, when $cO_2(Hb)$ is

reduced.

3.5.3 Intracapillary resistance to oxygen

During oxygen transport from the erythrocyte to the cell membrane several theoretical resistive forces to oxygen diffusion are encountered. The cytosol and the cell membrane of the erythrocyte have been mathematically and experimentally determined to be of a negligible resistance for oxygen diffusion^[52,56]. However, the plasma phase, with its very low oxygen solubility, limits the ability to transfer oxygen from the red cells to the tissues and affects the uniformity of oxygen flux at the capillary wall and is an important resistive component in the oxygen uptake^[52,57,58]. The effect of the low plasma oxygen solubility was demonstrated in hemodilution studies inducing an expansion of the plasma phase and an increased red cell spacing, consequently, affecting the oxygen flux to the tissues^[59]. In conclusion, a major fraction of the resistance of oxygen transport between the red blood cell and mitochondrion is intra capillary and is related to the pathway of oxygen diffusion through the plasma gaps^[60].

3.5.4 Myoglobin

The Krogh-Kety model neglects the effect of the presence of myoglobin in skeletal muscles. The myoglobin molecule is a monomer variety of the hemoglobin molecule present in the vertebrate red muscle and heart. Myoglobin has a role in the storage of oxygen and has a buffer capacity, when oxygen delivery is reduced. Another role of myoglobin, namely the facilitated diffusion of oxygen, has now been established^[52]. To illustrate the magnitude of the facilitated diffusion by myoglobin, the ratio of the facilitated oxygen diffusion to the free oxygen diffusion is reported to be 3:1 in red muscle at the p_{50} of myoglobin^[61]. Additionally, it was demonstrated that myoglobin rich skeletal muscle can sustain a high $\dot{V}O_2$ with a relatively low pO_2 due to the facilitated diffusion of oxygen.

3.5.5 Constant and uniform oxygen consumption

Inside the cell oxygen consumption is almost exclusively situated in the mitochondrion. It was demonstrated that pO_2 distribution was homogeneous in the cytosol, provided that mitochondria are uniformly distributed within the cell. However, mitochondria can be distributed heterogeneously in the tissues^[52]. Mainwood et al. showed that mitochondria were more prominently positioned around capillaries and that mitochondrial density decreased with increasing distance from the capillary^[56]. However, Hoppeler et al. concluded that these findings are largely species and muscle dependent and that it is too early to draw definitive conclusion upon the relationship between mitochondria and microvascular architecture^[62].

3.6 Oxygen consumption within the cells

Oxygen enables the energy contained in food to be converted into a form of controlled energy released by the formation of Adenosine Tri-Phosphate (ATP). ATP has the unique property to store energy in high "energy bonds" which on hydrolysis release 11-13 kcal / mole ATP. Three chemical processes can be discerned within the cell related to the conversion of energy^[12,63].

3.6.1 Glycolysis

Glucose enters the cell by a process of facilitated or carrier mediated diffusion, which is enhanced by insulin. Then a conversion to glucose-6-phosphate (enzyme: glucose-kinase, ATP consuming process) takes place. This molecule can be used for anabolic

metabolism (glycogenesis) or for the release of energy. (glycolysis). Glycolysis involves a process of ten reactions in the cytosol of the cell with the net gain of two molecules of ATP and ends with the formation of two molecules of pyruvic acid. There is also a release of protons and electrons, which combine with nicotamide adenine dinucleotide (NAD^+) to form NADH and free hydrogen ions. The whole process of glycolysis comprises the anaerobic metabolism of glucose. Thus, ATP can be produced even in the absence of oxygen by the process of glycolysis.

3.6.2 Citric acid cycle

The next step in the generation of energy is the transport of pyruvate into the mitochondrion. There acetyl-CoA is formed from pyruvate by the pyruvate dehydrogenase complex. Then, in successive steps in the citric acid cycle acetyl-CoA is transformed and 2 molecules of CO_2 and one molecule of ATP are produced. Although the citric acid cycle constitutes part of aerobic metabolism, none of the reactions makes direct use of molecular oxygen. The most important contribution of the citric acid cycle is the production of high energy electrons in the electron carriers NADH and FADH_2 .

3.6.3 Oxidative phosphorylation

The complicated transfer of these electrons along the inner membrane of the mitochondrion to oxygen is called the oxidative phosphorylation, since this process yields 34 molecules of ATP, and is dependent on oxygen.

3.6.4 Fat and protein

Other substrates as fat and protein are oxidized by transformation and deamination, respectively in order to enter the citric acid cycle. Fat is first oxidized into glycerol and fatty acids. Glycerol is transformed into glyceraldehyde-3-phosphate, which is an intermediate in the glycolytic pathway. Fatty acids are transported into the mitochondria and degraded to acetyl-CoA with the generation of NADH and FADH_2 . Amino acids are deaminated in the liver and the carbon skeleton is converted to the various intermediates of the glycolytic pathway and citric acid cycle.

3.7 The mitochondrion: the end-organelle for oxygen consumption and the final step of the oxygen cascade.

3.7.1 Anatomy and function ^[11]

Mitochondria are present in all eucaryotic cells, and are subcellular organelles, that convert energy to forms that can be used for numerous cell functions. Mitochondria are mobile and constantly changing their shape. They are bounded by two different membranes, creating two compartments: the innermembrane space or matrix and the much smaller intermembrane space between the outer and inner membrane. Because the outer membrane is permeable for all molecules of 5000 daltons or less, the inter membrane space is similar to the cytosol. The innermembrane space or matrix contains a different set of proteins because of the impermeability of the inner membrane. The phospholipid cardiolipin contributes to this impermeability, especially for ions. The inner membrane is highly convoluted, forming the cristae in the matrix and increase the surface of the inner membrane. Specific translocase proteins are needed to transfer pyruvate, fatty acids (carnithine), amino acids and intermediates for oxidative metabolism from the cytosol into the matrix space. The citric acid cycle with the generation of the electron carriers NADH⁺ and FADH_2 is located in the matrix of the mitochondrion.

The inner mitochondrial membrane has a crucial role in the oxidative phosphorylation. The process of electron transfer starts when the electrons are removed from NADH^+ and accepted by the NADH dehydrogenase complex. Then the electrons pass through 40 proteins which are grouped in three large enzyme complexes and two electron carriers. These enzyme complexes exist as independent entities in the plane of the inner membrane of the mitochondrion. Each complex has a greater electron affinity as his predecessor and electrons pass sequentially from one complex to another until they are finally transferred to oxygen, which has the greatest electron affinity. It is the last of the enzyme complexes, the cytochrome oxidase complex or cytochrome aa3, that finally catalyzes an efficient reduction of oxygen, by anchoring the oxygen molecule between a heme linked iron atom and a copper atom until it has picked up a total of four electrons [64]. The cytochrome oxidase reaction accounts for 90% of the total oxygen uptake in the cells. Cyanide is toxic to cells because it binds tightly to this complex and thereby blocks all electron transport.

This energetically favourable electron transport is coupled to a H^+ transfer across the inner membrane from the matrix to the intermembrane space. The H^+ transfer generates a gradient across the inner membrane which is composed of a membrane potential (140 mV) and of a concentration (pH) gradient (60 mV) and is called the electrochemical proton gradient (200 mV). Down the electrochemical gradient a proton flow backwards through the enzyme ATP-synthase is used to drive the synthesis of ATP and to produce large amounts of ATP. The enzyme ATP-synthase is an inner membrane bound protein complex that synthesizes ATP when protons flow through down the electrochemical proton gradient.

3.7.2 Uncoupling the electron transfer from the ATP synthesis.

Several substances have been known to uncouple the electron transport from the synthesis of ATP. These agents act as to provide additional channels for protons to flow through the inner membrane and consequently to diminish the electrochemical gradient. Remarkably, when an uncoupling agent has been added to the cells the electron transfer and oxygen consumption is increased in a large extent. It seems that the magnitude of the electrochemical gradient is a feed back for both the electron transfer as the ATP synthesis. In some specialized cells as brown fat cells the electron transfer and ATP synthesis are normally uncoupled; consequently, the energy of the electron transfer is converted into heat.

3.7.3 The genome of mitochondria [11]

In contrast to other subcellular organelles mitochondria contain a functional DNA. However, most of the proteins in mitochondria are encoded in the nuclear DNA and imported from the cytosol, but some are encoded by the organelle DNA. Mitochondrial DNA does not encode for proteins used outside the mitochondrion. Mitochondrial DNA is circular and distributed in the matrix and in contrast to nuclear DNA all DNA appears to be part of the encoding system leaving little room for regulatory DNA sequences. In addition, there are differences in the genetic code with a reduced fidelity of the DNA replication. Both DNA strands are completely transcribed, but only the transcripts made on one (heavy) strand are extensively processed.

The inheritance of characteristics encoded by mitochondrial DNA is in several ways very different from the inheritance of characteristics encoded by nuclear DNA [65,66]. The inheritance of the mitochondrial DNA in humans is uniparental and more precisely maternal, because the sperm cell does not contain mitochondria. Secondly, mutations accumulate through maternal lineage, since mitochondrial DNA does not recombine.

Mutations and normal mitochondrial DNA can coexist within the same cell, a phenomenon called heteroplasmy. By divisions and replications the proportion of mutant mitochondrial DNA can augment or diminish according to the principles of non-mendelian inheritance. Mitochondrial DNA is 10 times more prone to mutations due to several reasons: Since mitochondrial DNA has no introns, mutations will always damage an encoding segment; Furthermore, an ineffective repair system together with a continuous exposition to oxygen free radicals generated by the oxidative phosphorylation are other factors that contribute to the increased vulnerability of mitochondrial DNA.

All tissues in the body depend on integral mitochondrial function and an increasing number of diseases has been contributed to mitochondrial disorders. Aging itself was attributed to age related dysfunctions of the mitochondrial DNA mediated by accumulating oxidative damage.

3.7.4 The mitochondrion during perinatal development

Data of the fetal and postnatal mitochondrial development still remain substantially limited to animal data. In early fetal period in the rat liver mitochondria have the same dimensions and shape as in adulthood, but the number of mitochondria is much lower than in adulthood ^[67]. Later in fetal life mitochondria increase in size but not in number. Although the dimensions of mitochondria increase, this growth keeps in pace with the increasing cytoplasm and mitochondrial cell fraction remains constant. Biochemical changes during this period as a decreasing cytochrome-c oxidase and cardiolipin activity and increasing glutamate dehydrogenase reflect the increasing matrix volume relative to the inner membrane surface area.

In the postnatal period within one day after birth mitochondria decrease in size and in volume and start to proliferate rapidly. This process results in an increase in fraction and in number of mitochondria per unit of cytoplasm. The proliferation phase is characterized by an increasing mitochondrial membrane area relative to matrix space. The data on postnatal enzyme development show an increase within the first week after birth especially for the inner membrane components as cytochrome-c oxidase. The mitochondrial proliferation seems to be initiated by the synthesis of key proteins and by an accumulation of adenine nucleotide in the matrix ^[68,69]. Mitochondrial development is affected by hypothyroidism by reducing mitochondrial proliferation ^[70] and by adrenal steroid hormones by up regulating cytochrome-c oxidase expression in a tissue specific manner ^[71].

The lower activity of oxidative enzymes during the fetal period relative to postnatal and adult period was confirmed in other tissues as rat brain ^[72], rat heart tissue ^[73], rat kidney ^[74] and human skin fibroblasts ^[75].

Hence, it seems, that the perinatal period is characterized by a switch from low oxidative metabolism to efficient oxidative phosphorylation by a rapid proliferation and maturation of cellular mitochondria.

4 Venous oxygen saturation ^[76]

Venous oxygen saturation is defined as the oxygen saturation in the venous blood after it has traversed the capillary bed of a certain tissue. The value of venous oxygen saturation in the venous blood in the right heart or in the pulmonary artery (SvO_2) is the average of the venous blood from all the tissues. Kandel and Aberman ^[77] have expressed three criteria for the validity of the measurement of SvO_2 in the right heart or pulmonary artery:

- 1) SvO_2 should include all the blood that has traversed all the capillary beds

capable of extracting oxygen. Apart from the blood of the thebesian veins and bronchial arteries, the SvO_2 measurement in the right atrium, right ventricle or pulmonary artery meet this criterium. SvO_2 measurement in the superior or inferior caval vein does not fulfil this criterium and the SvO_2 value is in fact different between these two measurement sites

2) SvO_2 should exclude any blood that has not traversed capillary beds. The value of SvO_2 is unreliable when a considerable amount of blood is shunting from the left side to the right side of the heart as in the presence of an atrium - or ventricle septum defect.

3) SvO_2 should be measured in blood mixed so completely, that a single uniform oxygen saturation value can be encountered. Kandel and Aberman [77] state that only measurements in the pulmonary artery fulfils this criterium. However, some studies show evidence that the right atrium is an as proper measurement location as the pulmonary artery [78,79]. SvO_2 is determined by the oxygen transport to the tissues, or the oxygen delivery (DO_2) and by the oxygen extraction in the capillary bed of the tissues. DO_2 is determined by cardiac output (Qt) and the total oxygen content of the arterial blood (caO_2) [63,80,81].

$$DO_2 \text{ ml.kg}^{-1}.\text{min}^{-1} = caO_2 \text{ ml.dl}^{-1} \cdot 10 \cdot Qt \text{ ml.kg}^{-1}.\text{min}^{-1} \quad (1)$$

caO_2 is determined by hemoglobin concentration (Hb) and oxygen saturation (SaO_2) and the oxygen in solution expressed by the partial pressure of oxygen (paO_2).

$$caO_2 \text{ ml.dl}^{-1} = (1.39 \text{ ml.g}^{-1} \cdot SaO_2 \cdot Hb \text{ g.dl}^{-1}) + (0.0031 \text{ ml.dl}^{-1}.\text{mmHg}^{-1} \cdot paO_2 \text{ mmHg}) \quad (2)$$

Oxygen extraction or oxygen consumption (VO_2) is according to the Fick equation determined by the difference between arterial (caO_2) and venous oxygen content (cvO_2) and cardiac output (Qt).

$$VO_2 \text{ ml.kg}^{-1}.\text{min}^{-1} = (caO_2 - cvO_2) \text{ ml.dl}^{-1} \cdot 10 \cdot Qt \text{ l.kg}^{-1}.\text{min}^{-1} \quad (3)$$

When we combine and rearrange equation (2) and (3) and neglect the small volume of oxygen dissolved in the blood,

$$VO_2 \text{ ml.kg}^{-1}.\text{min}^{-1} = (1.39 \text{ ml.g}^{-1} \cdot Hb \text{ g.dl}^{-1}) \cdot (SaO_2 - SvO_2) \cdot 10 \cdot Qt \text{ l.kg}^{-1}.\text{min}^{-1} \quad (4)$$

$$SvO_2 = SaO_2 - VO_2 \text{ ml.kg}^{-1}.\text{min}^{-1} \cdot (1.39 \text{ ml.g}^{-1} \cdot Hb \text{ g.dl}^{-1} \cdot 10 \cdot Qt \text{ l.kg}^{-1}.\text{min}^{-1})^{-1} \quad (5)$$

we can see that changes of the venous oxygen saturation (SvO_2) can be due to changes in VO_2 , and in the factors that constitute DO_2 , such as arterial oxygen saturation (SaO_2), hemoglobin (Hb) and cardiac output (Qt). SvO_2 reflects the combined sufficiency of all these variables.

The effect of changes of the individual variables for the value of SvO_2 have been described by different authors [77,82,83]. SvO_2 is affected considerably by changes in oxygen consumption (VO_2), such as reductions in VO_2 during anesthesia and hypothermia, and elevations of VO_2 during convulsions or postoperative shivering [84]. Changes in the variables, that constitute the oxygen delivery (DO_2) as SaO_2 , Hb and cardiac output (Qt) have their effect on SvO_2 , as well [81]. One must be very cautious to use SvO_2 measurement to monitor exclusively one of these variables, particularly when other variables are unstable. There is a poor correlation between SvO_2 and individual determinants of the oxygen transport [76].

SvO_2 is a monitor of the balance between DO_2 and VO_2 [76,77]. When DO_2 is

decreased, several compensatory mechanisms may be called into play: increased cardiac output or increased oxygen extraction (ie, decreasing SvO_2) are the most common compensatory mechanisms [77]. For example, in the case of a severe anemia cardiac output can increase triple fold above the normal value and SvO_2 can decrease from 75 % to values below 40 % [85]. Consequently a seven to nine-fold increase in DO_2 can be accomplished, provided that cardiac performance and tissue oxygen uptake are not compromised. In the case of cardiac failure, only one of the compensatory mechanisms - increased oxygen extraction - can be called into play. Therefore, patients seem to be less tolerant to a reduction of cardiac output than they are to a reduction of SaO_2 or hemoglobin concentration.

Another compensatory mechanism in the case of hypoxia or reduced cardiac output is a redistribution of blood flow. The quantity of blood flow from the different organs determines the value of SvO_2 in addition to DO_2 and VO_2 . For example, venous blood from a high flow / low oxygen extraction organ as the kidney determines the value of SvO_2 to a greater extent than venous blood from a low flow / high extraction organ as the heart. During shock or cardiac failure, a reduction of renal blood flow together with an increase of the blood flow to coronary vascular bed of the heart will decrease the SvO_2 value substantially. SvO_2 is consequently a flow-weighted average of venous oxygen saturation values from all the tissues. In fact, redistribution of blood flow and an increased oxygen extraction or a decreased SvO_2 are partly analogous mechanisms, since by a redistribution of blood to low flow / high extraction organs oxygen extraction will increase and SvO_2 will decrease.

Monitoring SvO_2 enables the estimation of the shunt index, an estimate of the venous admixture and fractional oxygen extraction.

$$\text{Shunt Index} = (100 - SaO_2) / (100 - SvO_2) \quad (6)$$

The shunt index is an estimate of the venous admixture and is a calculated percentage of blood, that refers to the degree of admixture of mixed venous blood with pulmonary end-capillary blood, which would be required to produce the observed difference between the arterial and the pulmonary end-capillary pO_2 [15]. The shunt index as an estimate of the venous admixture obtained by continuous pulse and venous oximetry (dual oximetry) has been suggested and studied in adult intensive care medicine [86,87]. Bongard et al. [88] described a computer modelling of the shunt index and found limitations in the accuracy of the estimate of venous admixture, when errors in the SaO_2 value were present.

Fractional oxygen extraction is an assessment of the peripheral tissue oxygen utilization and an instantaneous feedback on circulatory and respiratory therapy [89].

$$\text{Fractional oxygen extraction} = (SaO_2 - SvO_2) / SaO_2 \quad (7)$$

Mathematically it can be expected that a good correlation exists between fractional oxygen extraction and SvO_2 . However, fractional oxygen extraction based on SvO_2 reflects global oxygen extraction and therefore tissue hypoxia in individual organs cannot be excluded [90].

4.1 Summary

SvO_2 is flow-weighted average of venous oxygen saturation values from all the tissues. SvO_2 is the combined sufficiency of all the variables that constitute DO_2 and VO_2 . Consequently, SvO_2 is a monitor of the balance between DO_2 and VO_2 . When the relation of DO_2 and VO_2 is out of balance, several compensatory mechanisms could come into play to restore the balance. Increased oxygen extraction or a decreased SvO_2 are one of the most common compensatory mechanism. A decreased SvO_2 is a sign, that one of the variables, that constitute DO_2 and VO_2 is impaired. SvO_2 enables the estimation of the shunt index and fractional oxygen extraction.

II *Defining* Hypoxia

1 Definitions

2 Cellular hypoxia

2.1 *Definition of cellular hypoxia*

2.2 *Alteration in substrate utilization during hypoxia*

2.3 *Anaerobic sources of energy (1): Glycolysis*

2.3.1 Control of glycolysis

2.3.2 Lactate

2.3.2.1 Lactate production

2.3.2.2 Lactate utilization

2.3.2.3 Lactate as a monitor of tissue hypoxia in adult intensive care medicine

2.3.2.4 Lactate in neonatal medicine

2.4 *Anaerobic sources of energy (2): Phosphocreatine reaction*

2.5 *Mechanisms of cell damage during hypoxia*

3 Systemic hypoxia

3.1 *Hypoxic hypoxia*

3.1.1 Acute hypoxic hypoxia and oxygen transport

3.1.2 Cardiovascular responses during development

3.1.3 Chronic hypoxic hypoxia

3.2 *Anemic hypoxia*

3.2.1 Oxygen transport and compensatory mechanisms

3.2.2 Comparing hypoxic and anemic hypoxia

3.2.3 Chronic anemic hypoxia

3.3 *Circulatory hypoxia*

3.3.1 Critical cardiac output

3.3.2 Chronic circulatory hypoxia

1 Definitions

In the past hypoxia was defined as a condition with a low paO_2 or low SaO_2 . In more recent years the definition of hypoxia has been focused to the target of oxygen transport: the peripheral tissues. In the report of a consensus conference^[91] hypoxia was defined as a condition in which the cells of a tissue have abnormal oxygen utilization, such that the tissue is experiencing anaerobic metabolism.

Terms as hypoxic, anemic, circulatory and histotoxic hypoxia are more related to the disorders that can lead to hypoxia^[92]. Connet et al.^[93] defined hypoxia as "oxygen-limited cellular energy failure". The central point is the balance between ATP demand and ATP generated by aerobic metabolism. This balance depends on the interaction between several subsystems as oxidative phosphorylation, Krebs cycle, glycolysis, substrate supply, membrane transport and cell energetics. In this review the consequences of hypoxia will be discussed both at cellular level and at systemic, multiorgan level.

2 Cellular hypoxia

2.1 Definition of cellular hypoxia

A common definition for cellular hypoxia is a "pathologic condition in which cellular energy requirements are greater than aerobic energy production"^[97,98]. Oxygen is the ultimate electron acceptor in the respiratory chain. The last enzyme complex of the respiratory chain, cytochrome oxidase, catalyses the reduction of oxygen and this reaction accounts for 90 % of the VO_2 . The *critical intracellular pO_2* has been defined as the pO_2 below which electron transport and oxidative phosphorylation are minimized. In isolated mitochondria the critical pO_2 has been estimated as low as 0.1 - 1 mmHg. However, when critical pO_2 was defined as pO_2 associated with a certain redox state of the cytochrome oxidase complex, an as 10 times higher critical pO_2 value was demonstrated in intact hepatocytes^[99]. It was hypothesized, that clustering and uneven distribution of mitochondria are the major reasons for this difference.

The *redox state of the mitochondria* is reflected by the ratio of $\text{NADH} / \text{NAD}^+$. An increased level of NADH and consequently an increased redox state is induced by a decreased oxygen availability, although other factors as substrate availability and citric acid cycle activity may also affect the NADH level. The ratio of $\text{NADH} / \text{NAD}^+$ in the cytosol is reflected by the ratio of lactate / pyruvate^[100,101].

It is known that some cells have a greater *tolerance for hypoxia* than other cells, dependent on cell activity and tissue origin. For example, skeletal muscle recover after 30 minutes complete ischaemia, whereas brain cells are irreversibly damaged after 5 minutes of hypoxia.

2.2 Alteration in substrate utilization during hypoxia or intensive exercise^[99]

The oxygen requirements for ATP formation varies between different substrates. The oxidation of free fatty acids requires 10 % more oxygen than the oxidation of carbohydrates. During aerobic metabolism the normal substrate for cellular energy production are in the myocytes, for example, lipids, lactate and glucose, in that order. However, during hypoxic conditions the oxidation of glycogen is the preferred pathway. This switch from fat to glucose oxidation during hypoxia was demonstrated in several different cell lines^[99].

2.3 Anaerobic sources of energy (1): Glycolysis

Glycolysis is a number of ten subsequent chemical reactions decomposing glucose to two molecules of pyruvate. Glycolysis occurs in the cytoplasm entirely without oxygen and with the net gain of two molecules of ATP. Although glycolysis is a very inefficient energetic process, since only 3 per cent of the total energy of the glucose molecule is used, it is a very important process to produce energy, when oxygen is not available ^[102].

2.3.1 Control of glycolysis

Glycogen is the main source for the glycolysis. The availability of glycogen and the breakdown of glycogen, (glycogenolysis), are regulating steps for the substrate availability for the glycolysis. The initial step in glycogenolysis is catalyzed by phosphorylase. The phosphorylase can be activated by hormones as epinephrine and glucagon. Factors as Ca^{2+} , AMP and inorganic phosphate also activate glycogenolysis. However, H^+ ions and acidosis inhibit glycogenolysis and muscle contraction ^[99].

Glycolysis itself is controlled by the cellular need of ATP. ATP controls one of the reactions in the glycolysis by the inhibition of the enzyme phosphofructokinase. H^+ ions and citrate potentiate this inhibition. On the other hand ADP and inorganic Phosphate increase the activity of phosphofructokinase, thereby stimulating glycolysis.

In summary, it seems that both glycogenolysis and glycolysis are increased by Ca^{2+} , ADP, AMP and inorganic phosphate and inhibited by ATP and H^+ ions.

2.3.2 Lactate

The end products of glycolysis -pyruvate, NADH and H^+ - may react with each other and lactate may be formed ^[102]. The conversion from pyruvate to lactate is catalyzed by the enzyme lactic dehydrogenase (LDH). Since lactate diffuses out of the cells to the interstitial fluid, glycolysis can be continued for around 30 to 40 seconds.

2.3.2.1 Lactate production

Lactate is produced by all tissues, but brain cells, erythrocytes and renal cells are considered as the most prominent lactate synthesizing cells ^[103]. One of the main factors determining lactate production is the balance between pyruvate production and pyruvate utilization. As pyruvate production is determined by glycolysis, pyruvate utilization is determined by the metabolism of pyruvate in the citric acid cycle or by gluconeogenesis. These two different pathways are catalyzed by pyruvate dehydrogenase and pyruvate carboxylase, respectively. These enzymes can be inhibited by insufficient quantities of cofactors, as NADH or ATP and by conditions as starvation and diabetes. All these interacting factors and conditions may affect the balance between pyruvate production and utilization and may lead to lactate acidosis ^[103].

Another factor, that has a great impact on the lactate production is the redox state of the cytosol ^[104]. As was mentioned above, the redox state of the cytosol is dependent upon the oxygen availability in the mitochondria. An increased redox state reflecting an increased ratio between the reduced and oxidized molecules in the cytosol is associated with lactate production. Decreased oxygen is the most common condition increasing the redox state, but toxins and certain drugs can also affect the redox state and stimulate lactate production ^[105].

2.3.2.2 Lactate utilization

The utilization of lactate is characterized by a reversion to pyruvate after which pyruvate is either oxidized in the citric acid cycle or reconverted in the gluconeogenesis back to glucose ^[104]. The gluconeogenesis occurs prominently in the liver

and kidney. On the other hand in the brain, heart and skeletal muscle, lactate can directly be used for energy.

2.3.2.3 Lactate as a monitor of tissue hypoxia in adult intensive care medicine^[105]

Lactate was proposed as a metabolic monitor of shock and tissue hypoxia. Wasserman et al.^[106] confirmed in a human study the concept that lactate did not increase gradually and continuously, but only when a threshold oxygen consumption is exceeded. The course of the lactate values during progressive graded hypoxia is best described by the biphasic dual-line in contrast to a single-line or polynomial line technique^[107].

The amount of lactate produced is correlated with the oxygen debt and the severity of the shock. In agreement with this lactate appeared to have a prognostic value. Several authors have demonstrated an increased mortality in shock, when lactate levels were higher than 2.4 mmol/L. Serial lactate determinations in critically ill patients can even improve the prognostic value. On the other hand, other factors as hepatic reserve and nutrition status may affect the plasma lactate concentration^[105].

2.3.2.4 Lactate in neonatal medicine

In neonatal medicine lactate was used as a monitor of tissue hypoperfusion and tissue hypoxia. In a newborn animal experiment induced group B streptococcal sepsis induced an increased arterial lactate concentration. Lactate production was most prominent in those tissues, in which oxygen uptake was reduced, as skeletal muscle and cerebral tissue^[108]. In preterm infants the effect of sepsis on carbohydrate metabolism was observed. No significant changes were seen in the glucose and insulin concentration. However, plasma lactate concentration was increased. The authors concluded that plasma lactate may be an early clinical marker of neonatal sepsis^[109]. A common problem in neonatology is when to give an erythrocyte transfusion in healthy preterm infants with anemia of prematurity. In an observational study it was concluded, that lactate could be a predictor for erythrocyte transfusion need in this group of patients^[110]. Plasma lactate concentration as a predictor of outcome was investigated in a retrospective study in neonates requiring extracorporeal membrane oxygenation (ECMO). In non-survivors lactate levels before and after ECMO were significantly higher than in survivors^[111]. Several metabolic parameters among which lactate have been studied in perinatal asphyxia to determine severity and to predict outcome. Increased plasma lactate concentration seemed to have a low sensitivity and a low positive predictive value for adverse outcome^[112].

In summary, plasma lactate concentration has been used in neonatal medicine to monitor tissue hypoxia and to estimate tissue damage after asphyxia. It seemed that elevated levels of lactate were associated with tissue hypoxia in sepsis, anemia of prematurity and in infants who need ECMO. The association between lactate and outcome after asphyxia is less obvious.

2.4 Anaerobic sources of energy (2): Phosphocreatine reaction^[99,102]

Phosphocreatine is a protein that consists of creatine and an inorganic high energy phosphate bond. By breaking down this bond enough energy is released to regenerate ATP from ADP. The break down of phosphocreatine is catalyzed by the enzyme creatine kinase and is promoted by the end products of the hydrolysis of ATP: ADP and inorganic phosphate. Therefore, the break down of phosphocreatine is connected with the hydrolysis of aerobic ATP and reflects more an increased energy demand than an anaerobic condition. Both the ATP hydrolysis and phosphocreatine breakdown result in an increased intracellular inorganic phosphate. The increased intracellular inorganic phosphate stimulate glycogenolysis, glycolysis and oxidative phosphorylation.

In summary the phosphocreatine system has an important function in the first 5 to 10 seconds of intense exercise. In addition glycolysis can be continued for a period of 30 to 40 seconds. Longer periods of exercise need to be answered by the oxidative metabolism. Phosphocreatine reaction and glycolysis produces ATP 4 and 2.5 times as rapidly as normal oxidative metabolism in the mitochondria, respectively ^[113].

2.5 Mechanisms of cell damage during hypoxia ^[97,98]

Cellular acidosis The major source of H^+ ions during hypoxia is the immediate hydrolysis of ATP formed during glycolysis. The simultaneously formed lactate will consume H^+ ions in the process of gluconeogenesis and oxidation in the Krebs cycle. When lactate utilization in these pathways is impaired (tissue hypoxia, toxicity) lactic acidosis will ensue ^[104]. Intracellular acidosis may damage the cell by the following mechanisms: loss of adenine nucleotides, as AMP precursors, ADP, ATP and NADH, from the mitochondrion and inhibition of the Na^+ / Ca^{2+} exchanger resulting in an increased intracellular calcium. **Increase in intracellular calcium concentration:** The intracellular Ca^{2+} concentration is closely regulated by membrane bound ATP controlled Ca^{2+} pumps, Na^+ / Ca^{2+} exchangers and voltage dependent Ca^{2+} pumps. In the case of hypoxia intracellular Ca^{2+} will increase by inhibition of the before mentioned Ca^{2+} pumps. Increased intracellular Ca^{2+} induces phospholipase, which damage the cell membrane, induces ATPases in the plasmamembrane, induces breaks in DNA and finally induces the expression of genes important in initiating cell death and apoptosis ^[98]. **Formation of oxygen free radicals:** Free oxygen radicals are responsible for the reperfusion damage after the hypoxic incident. One of the main mechanisms generating free oxygen radicals include the action of the O form of xanthine oxidase upon reoxygenation.

3 Systemic hypoxia

Hypoxia in terms of oxygen delivery (DO_2) and consumption (VO_2) at systemic level was investigated by Cain ^[114,115]. He found a *biphasic relationship* between DO_2 and VO_2 . VO_2 was found to be independent of the systemic DO_2 , when DO_2 decreased over a wide range due to different compensatory mechanisms. Hypoxia or anaerobic metabolism will only ensue, when DO_2 is reduced below a critical level and VO_2 will decline. In anesthetized human patients the biphasic relationship between DO_2 and VO_2 was established by Shibutani et al ^[116]. The idea of the biphasic relationship between DO_2 and VO_2 was questioned by Schlittig and Pinsky ^[117]. They argued that in most physiological situations VO_2 varies several fold during the course of a normal day and that the increased VO_2 is not supported by increased oxygen extraction but by an increased DO_2 (oxygen demand dependency). It seems that one can speak only of a biphasical relationship between DO_2 and VO_2 , when oxygen demand is kept constant.

In this review on systemic hypoxia the different disorders that can lead to tissue hypoxia as hypoxic, anemic and circulatory hypoxia will be discussed, since different compensatory mechanisms have been described.

3.1 Hypoxic hypoxia

3.1.1 Acute hypoxic hypoxia and oxygen transport in the neonatal animal.

The biphasic relationship between DO_2 and VO_2 was confirmed in different neonatal animals during acute hypoxic hypoxia ^[118-120]. Torrance et al. attempted to define the arterial pO_2 threshold for oxygen-restricted metabolism, as confirmed by metabolic

acidosis^[118] and increased lactate concentration^[119], during graded hypoxia in neonatal piglets. Normal acid-base balance was maintained with a wide range of paO_2 values to 5.3 kPa (40 mmHg). Below this paO_2 value a progressive metabolic acidosis ensued. This threshold value was confirmed in the fact that also lactate increased below a paO_2 level of 40 mmHg.

Moss et al.^[120] studied metabolic responses during graded alveolar hypoxia in conscious lambs to test whether there is a difference between the metabolic response at different age. The biphasic response, i.e. VO_2 and lactate concentration were unchanged except at the lowest inspired oxygen concentration, was confirmed in all three age groups. Since baseline DO_2 and VO_2 were higher in the youngest group of animals (< 1 week), the critical DO_2 level was higher in the youngest animals compared with the two other groups. Lactate increased when VO_2 decreased more than 15 percent below baseline. It seems that there is a bufferzone in the reduction of VO_2 , before lactate increased. For this, mechanisms of an increased lactate clearance or of an initial reduction of non-essential metabolism (for example: growth) was suggested.

3.1.2 Acute hypoxic hypoxia and cardiovascular compensatory mechanisms during development

The cardiovascular response to acute hypoxia in the *late fetus* consists of a decrease in heart rate, a variable change (often increase) of the mean arterial pressure and an increase in peripheral vascular resistance^[121,122]. The cardiac output is redistributed in favour of the cerebral and myocardial circulation at expense of the gastro-intestinal, pulmonary and renal circulation. These responses are mediated by chemoreceptors and local tissue factors. In the study of Bartelds et al.^[123] it was shown, that carotid and not aortic chemoreceptors mediate the cardiovascular response in the fetus. Sidi et al.^[124] studied the cardiovascular responses to acute hypoxemia in lambs *after birth*. The authors evaluated the responses and blood flow distribution together with the influence of age during moderate ($\text{FiO}_2 = 0.16$) and severe ($\text{FiO}_2 = 0.09$) hypoxia. Cardiopulmonary response included an early rise in heart rate, cardiac output, and arterial blood pressure during moderate hypoxemia. With severe hypoxemia cardiac output was unchanged and arterial blood pressure decreased along with a metabolic acidosis. The regional redistribution of blood flow during hypoxia changed similarly in all age groups: heart and brain flow increased; no change in flow to muscle and bone; renal, skin and GI tract flow decreased. This redistribution resulted in a increasing DO_2 to the heart, an unchanged DO_2 to the brain and a decreased DO_2 to all other organs. Another finding of this study was, that in younger lambs with a given change of oxygen consumption, metabolic acidosis and lactate was less pronounced. Younger lambs seemed to tolerate a decreased VO_2 better than older lambs. After birth, the response to acute hypoxic hypoxemia is markedly different from the response in utero: Heart rate and ventilation increase. Several mechanisms have been mentioned to explain this different response: pulmonary stretch receptors stimulated by an increased ventilation overruling chemical receptors, different responses between aortic and carotid bodies or developmental differences in chemoreceptor sensitivity^[124]. In the study of Koehler et al.^[125] regional blood flow changes during acute isocapnic hypoxia (40-50 % $\text{caO}_2(\text{Hb})$ reduction) induced by a reduced inspiratory oxygen administration were compared in newborn and *adult sheep*. The responses were qualitatively similar in cerebral, myocardial and skeletal muscle, where flow increased. In lambs renal and skin flow decreased, whereas in adult sheep it increased. In lambs small intestine flow was unchanged, but was increased in adults. It seemed, that a trend could be observed towards lesser vasoconstriction in the renal and gastro-intestinal vascular beds in the adult sheep compared with the neonatal counterpart. The authors speculate that mechanisms, as a direct effect of hypoxia on the central nervous system or an

increased sensitivity of the aortic chemoreflex possibly may play a role in the cardiovascular response in neonatal lambs ^[125].

3.1.3 Chronic hypoxic hypoxia and cardiovascular compensatory mechanisms during development

In a chronic lamb model Teitel et al. ^[126] evaluated the effects of hypoxemia produced by a right to left shunt (atrium septostomy and pulmonary artery stenosis by an inflatable balloon) with a normal alveolar pO_2 . Arterial oxygen saturations were reduced to 60 to 75 %. Systemic cardiac output decreased initially but quickly returned to normal values. DO_2 decreased concomitantly, but increased during the study period, because of the increase in hemoglobin concentration and cardiac output. Oxygen consumption indexed to weight remained within the normal range during the study period. However growth rate decreased dramatically. Again, it was speculated that oxygen consumption is redistributed from facultative or non-essential metabolism to increased cardiorespiratory work.

In a subsequent study in this chronic hypoxic lamb model Bernstein et al. ^[127] evaluated the question whether a redistribution of blood flow could account for the dramatic decrease in growth rate. Chronic hypoxemia did not alter total cardiac output. However, in contrast to the acute hypoxemia ^[124], chronic hypoxemia is not associated with increases in myocardial and cerebral blood flow. Cardiac output and cerebral blood flow initially increased but returned to control levels. Blood flow to the kidneys, spleen, gastrointestinal tract, carcass and skin remained decreased during the study period. Hemoglobin concentration gradually increased and systemic DO_2 returned to control values. However, DO_2 to kidneys, spleen, GI-tract, carcass and skin remained reduced. This decreased oxygen and substrate delivery to the gastro-intestinal tract, liver and carcass may account for the effect on growth in chronic hypoxemia.

3.2 Anemic hypoxia

3.2.1 Oxygen transport and compensatory mechanisms

Anemic hypoxia is characterized by a reduction of the arterial oxygen content ($caO_2(Hb)$) with a normal or even increased paO_2 . Compensatory mechanisms to preserve oxygen consumption include a) increased cardiac output, b) decreased systemic vascular resistance, c) an increased oxygen extraction, d) a shift of the hemoglobin oxygen dissociation curve and e) a redistribution of regional blood flow ^[128-130].

a) *Cardiac output* increases significantly in proportion to the decrease of hematocrit. The extent of the rise was reported between 40 to 125 %. The mechanisms for the increased cardiac output have been attributed in conscious animals to an increased heart rate, and in anesthetized animals to an increased stroke volume. An increased stroke volume was suggested to be generated by an increased venous return to the heart, a reduction of the left ventricular afterload and an increased myocardial contractility. As a consequence of the increased myocardial contractility it was seen that oxygen consumption may increase at the initially steps of hemodilution. The increased venous return to the heart, and the reduction of the left ventricular afterload are a direct consequence of the

b) *reduced blood viscosity* and consequently *decreased systemic vascular resistance*. Although peripheral vasodilation may play a role it seems that principally the decreased blood viscosity is responsible for an increased blood flow in the microcirculation.

c) Since further compensatory increase in flow is limited when hematocrit is reduced below 30 % to 25 %, *peripheral oxygen extraction* starts to play a more important

role below this hematocrit value. Maximal oxygen extraction values to 79 % have been encountered associated with very low SvO_2 values.

d) Chronic anemia induces a rise in the concentration of 2,3- diphosphoglycerate and therefore induces a *shift of the oxygen dissociation curve* to the right, facilitating oxygen release to the tissues.

e) A *redistribution of regional blood flow* was reported during anemic hypoxia. Together with the increased cardiac output a disproportionately increased coronary blood flow was found. An increase proportionate to the increased cardiac output was found in the vertebral and the superior mesenteric arteries and in the lower aorta. In the renal, hepatic and carotid arteries the blood flow increased proportionally less than the increase in cardiac output.

3.2.2 Comparing hypoxic and anemic hypoxia.

Cain ^[114] compared hypoxic and anemic hypoxia in dogs. He found, that oxygen uptake started to decrease and lactate started to increase at similar levels of systemic DO_2 . However, venous pO_2 at this critical level of DO_2 was higher at anemic hypoxia than in hypoxic hypoxia. Venous pO_2 is, theoretically, the lowest driving pressure of oxygen diffusion to the tissues. It seems curious, that at critical DO_2 , below which oxygen restricted metabolism will ensue, the lowest driving pressure for oxygen diffusion is different between two types of hypoxia. Explanation for this phenomenon includes the fact that the full range of the oxyhemoglobin dissociation curve is used and even at the lowest DO_2 level, high arterial SAO_2 and paO_2 , and consequently also higher values of PvO_2 and SvO_2 , are observed. In addition, differences in the redistribution of blood flow and microcirculation during anemic hypoxia compared with hypoxic hypoxia may play a role ^[131].

3.2.3 Chronic anemic hypoxia

Cardiovascular adaptations to chronic anemia were studied in the newborn lamb ^[133]. After two weeks of anemia (hemoglobin level 60 % of normal) measurements were performed. The major compensatory responses were an increase of heart rate, cardiac output and oxygen extraction. Blood flow to the heart and brain increased and flow to the viscera and carcass did not change. In chronic hypoxic hypoxia ^[127] it was found, that hemoglobin increased and systemic DO_2 normalized again within two weeks. In contrast to chronic anemic hypoxia heart rate and cardiac output did not change and blood flow to the visceral organs decreased. Normal paO_2 during anemic hypoxia may normalize blood flow to the visceral organs via peripheral chemoreceptors, whereas low paO_2 in hypoxic hypoxia may decrease bloodflow to the visceral organs.

No difference of the mixed venous oxygen tension or saturations, between anemic and hypoxemic hypoxia were found in this chronic hypoxia model.

3.3 Circulatory hypoxia

3.3.1 Critical cardiac output during development

Circulatory hypoxia in the neonatal animal model has obtained attention in studies designed to determine the critical value of cardiac output below which oxygen consumption decreased and anaerobic metabolism ensues ^[134]. Cardiac output was lowered progressively in unsedated, conscious 2 weeks, 4 weeks and 8 weeks old lambs. Baseline cardiac output showed the expected decline in the older age groups. In the 2 and 4 weeks old lambs the absolute critical level of the cardiac output was identical but in the oldest

lambs the critical levels were significantly lower. However, the critical level calculated as a fraction from baseline was significantly higher in the 4 weeks old lambs. Probably, this phenomenon is due to the nadir of the physiological anemia at this age. Subsequently, to determine the effect of hemoglobin on critical cardiac output Heusser et al. [135] reduced cardiac output at different hemoglobin concentrations. Despite different hemoglobin concentrations baseline cardiac output was not different. Since fractional oxygen extraction was increased in the lower Hgb concentration groups baseline oxygen consumption was similar in all the different Hgb groups. Critical cardiac output calculated as a fraction from baseline was significantly higher in the lower Hgb concentration groups. Critical total systemic DO_2 and fractional oxygen extraction were not different between the Hgb groups. The authors conclude that VO_2 decreases when DO_2 is reduced below a certain level, but this critical level is not dependent on the manner in which DO_2 is reduced.

Fahey et al. [136] addressed the issue of non-essential oxygen consumption and investigated the repayment of the oxygen deficit in 2 week and 8 week old lambs. They found that 2 week old lambs after a critical reduction of cardiac output maintained oxygen consumption during recovery slightly below baseline, whereas in 8-week old lambs oxygen consumption in the recovery period remained above baseline. Moreover, lactate increased compared with the oxygen deficit at a significantly lower rate in the 2-week old lambs. Consequently, 2-week old lambs showed a lower oxygen debt and a lower lactate concentration at severe reduced cardiac output. In other words it seemed, that 2-week old lambs decreased their metabolic needs by suppressing their facultative or non-essential metabolism in contrast to older lambs in comparable conditions.

3.3.2 Chronic circulatory hypoxia

Chronic circulatory (reduced cardiac output) and hypoxic hypoxia were compared, while lactate and hypoxanthine levels were determined during both types of hypoxia [137]. Lactate was linearly related to the cumulative oxygen debt in both types of hypoxia, but the slope of the regression was greater for circulatory than for hypoxic hypoxia. Hypoxanthine was only linearly related to oxygen debt in circulatory hypoxia. The authors speculated that the more pronounced redistribution of the bloodflow away from the liver and the kidneys during low cardiac output resulted in a decreased elimination of lactate and hypoxanthine, particularly.

III Monitoring Oxygen; Oximetry

- 1 General principles of oximetry
 - 1.1 *Fiberoptic reflection oximetry*
 - 1.1.1 Fiberoptic oximetry by the Oximetrix Opticath
 - 1.2 *Pulse oximetry*
- 2 Accuracy and reliability of oximetry
 - 2.1 *Fiberoptic catheter oximetry*
 - 2.1.1 Accuracy of fiberoptic oximetry in umbilical arterial catheters
 - 2.1.2 Accuracy of fiberoptic oximetry in pulmonary artery catheters for SvO₂
 - 2.1.2.1 Accuracy of fiberoptic oximetry in pulmonary artery catheter during different conditions
 - 2.1.2.2 Summary
 - 2.1.3 The position of the fiberoptic catheter for the measurement of SvO₂
 - 2.2 *Pulse oximetry*
 - 2.2.1 Accuracy studies
 - 2.2.2 Different definitions of oxygen saturation
 - 2.2.3 Dyshemoglobins, anemia and pulse oximetry
 - 2.2.4 Accuracy in neonatal medicine
 - 2.2.5 Usefulness of pulse oximetry
 - 2.2.6 Summary

1 General principles of oximetry

The term oximetry has been used for the measurement of oxygen saturation^[138]. The measurement of oxygen saturation currently includes:

- a) non-invasive monitoring of the arterial oxygen saturation of hemoglobin, now primarily by pulse oximetry;
- b) invasively monitoring both arterial, mixed venous and jugular bulb venous or intracardiac oxygen saturation by a fiberoptic catheter system;
- c) determining the oxygen saturation of drawn blood using multiple wavelengths oximeters^[139].

The basic principle underlying all oximetry is the difference in light absorption between oxyhemoglobin (O_2Hb) and deoxyhemoglobin (HHb). In the spectral range from 600–700 nm. O_2Hb has a lower absorbance than HHb , while at higher wavelengths (above 800 nm.) a reciprocal relationship is seen with a higher absorbance for O_2Hb compared with HHb ^[138,140]. If both wavelengths of light are used, a ratio (R) of absorbance at 660 nm. to absorbance > 800 nm. can be determined.

The second principle in oximetry, the Lambert-Beer's law, states that the absorbance of incident light in a solution with a solute is proportional to the concentration of that solute^[141] and that in a solution with several solutes the total absorbance is equal to the sum of the independent solutes. In the case of hemoglobin, the absorbance of light and the ratio (R) of absorbance at 660 nm. to absorbance > 800 nm. are proportional to the concentrations of O_2Hb and HHb . With this in mind a so-called calibration curve can be generated with oxygen saturation as a function of the ratio (R). Because the optics of the devices and probes do not completely fulfil the requirements of the Lambert-Beer's law and the relation between oxygen saturation and R is non-linear, the calibration curve in the currently available oximeters is derived from the relation between the value of R and invasive arterial oxygen saturation measurements^[138,142]. Since oximetry is based on the light absorbance from hemoglobin, it is therefore, unlike the oxyhemoglobin dissociation curve, unaffected by pH and pCO_2 .

1.1 Fiberoptic reflection oximetry.

The measurement of oxygen saturation of non-hemolyzed blood by reflected light was first described in 1949^[138,139]. The principle of reflection oximetry was used for fiberoptic catheters. A fiberoptic reflection oximeter measures oxygen at the tip of the catheter. The catheter contains two bundles of optical fibers, one guides light into the blood; the other one guides the reflected (back-scattered) light to one or more photocells^[143]. Usually two light emitting diodes (LED's) emit light at two different wavelengths at a rate of 200–300 Hz. The reflected light from the blood is focused on a photocell, amplified and computed to the ratio of the light reflections at the two different wavelengths. The relation ship between the ratio and oxygen saturation is represented by a slightly curved line. The light reflection is affected by hemoglobin concentration, variation in blood flow and erythrocyte shape^[143,144], dependent on wavelength and oxygen saturation.

1.1.1 Fiberoptic oximetry by the Oximetrix Opticath

Brown et al.^[22] described the fiberoptic reflection oximetry catheter, Opticath, used in our studies and manufactured by Oximetrix. (Mountain View, Ca, USA.) The light source of the Oximetrix oximeter consists of, in contrast to other fiberoptic catheter systems, three diodes that emit light (LED) at three different wavelengths $\lambda_1=670$ nm., $\lambda_2=700$ nm. and $\lambda_3=800$ nm. These wavelengths were chosen to minimize the effects

introduced by changes in hematocrit and bloodflow. They were selected after analysis of data obtained from *in vitro* experiments and *in vivo* studies conducted in anesthetized animals and human volunteers. A pulsegenerator sequentially activates each diode for 1 msec followed by an equal interval when no diodes are energized. This four period cycle occurs at a rate of 244 Hz. The light from each diode is transmitted to the blood through one of the two catheter fiberoptics. The light reflected by the O₂Hb and HHb is transmitted by the other receiving fiberoptic bundle to a photodiode. The electrical signal from the photodiode represents the reflected light intensity (I) measurement made at the wavelength of each emitting diode. The signals are amplified and sent separately to the oximeter in the same sequence in which the LED's are energized. The oximeter provides two ratios of light intensity I_1^{670}/I_2^{700} and I_3^{800}/I_2^{700} , that are used to determine oxygen saturation. One ratio was found to be independent of the hematocrit in the upper range of oxygen saturation values and the other ratio was independent of the hematocrit in the lower range of oxygen saturation values. One of the ratio's is selected when appropriate by electronic switching techniques and the use of weighing factors. By using these three wavelengths computing two ratios the computed oxygen saturation was aimed to be independent of changes in hematocrit and of other wavelength-dependent characteristics.

1.2 Pulse oximetry.

Pulse oximetry was developed based on the ingenious idea to analyze the change in light absorption produced by arterial pulsations^[140,141]. Pulse oximetry analyzes the transmitted light, that consists of two components. The first component varies with the pulsation of the blood. The second component is a constant light output level that passes through the tissues without being absorbed or scattered. These two components are referred as AC components and DC components, respectively. Dividing the AC level by the DC level gives a corrected AC level reflecting the combined light transmission of O₂Hb and HHb. The ratio R indicates the corrected AC level at 660 nm divided by the corrected AC level at 940 nm. R is then converted to a oxygen saturation value using the calibration curve. Since specifically pulsatile, arterial blood is observed, absorbances and reflections from other tissues, pigmentation and venous and capillary blood are ignored^[138,142].

2 Accuracy and reliability of oximetry

2.1 Fiberoptic catheter oximetry

2.1.1 Accuracy of fiberoptic oximetry used as an umbilical arterial catheter in neonatal medicine.

The accuracy of the fiberoptic catheter system has been tested in very different and several ways. Initially fiberoptic catheters were used as umbilical arterial catheters to measure continuously arterial oxygen saturation in newborn infants. Wilkinson et al.^[145] and Krouskop et al.^[146] assessed the use and the accuracy of a 4 Fr. fiberoptic catheter (Oximatrix, Mountain View, Ca) as an arterial umbilical catheter in sick newborn infants. In 1039 comparisons with bench measured blood samples correlation analysis revealed a correlation coefficient of $r = 0.97$. A mean error of 0.74 ± 2.17 percent saturation was found. Accuracy was not affected by bilirubin level up to 18 mg/dl, variations in hematocrit (range from 19 % - 66 %), exchange transfusion, changes in hemoglobin oxygen affinity and fetal hemoglobin. Haessler et al.^[147] reported the use of an intra-arterial fiberoptic catheter (U 425 oximetry catheter (Abbott, Chicago, Ill.) compared with pulse

oximetry and Co-oximetry during cardiopulmonary bypass. Saturation readings were obtained in 99 % of the fiberoptic catheter readings and in 60 % to 84 % of the readings by pulse oximetry. Pulse oximetry failed during cardiopulmonary bypass with non-pulsatile flow and hypothermia. The mean difference and SD were for pulse oximetry -0.8 ± 1.0 % and for catheter oximetry 0.4 ± 1.0 %. The authors concluded that the performance of the fiberoptic catheter was excellent and superior to pulse oximetry without the necessity to perform an in vitro analysis.

2.1.2 Accuracy of fiberoptic oximetry used in pulmonary artery catheters for the measurement of mixed venous oxygen saturation (SvO₂).

With the coupling of the fiberoptic device to the pulmonary artery catheters of 7.5 Fr. fiberoptic reflectance oximetry for the continuous measurement of mixed venous oxygen saturation came into the adult intensive care arena^[148,149]. Since more devices using two and three wavelengths light sources came into the market, studies comparing these devices appeared. Gettinger et al.^[150] compared the accuracy and stability of two fiberoptic (Oximetrix and Edwards) catheters with a reference Co-oximeter in an animal study. Regression line and correlation coefficient differed significantly between the two catheters. However, Reinhart et al.^[151] found little difference between these same devices in a human clinical study. Deviations of more than 5% occurred in 13 % (Edwards) and 20 % (Oximetrix) of the measurements.

Leighton et al.^[152] examined two oximetry catheters (Oximetrix and Edwards-SAT2) in a swine model over a wide range of physiological states. They concluded that both systems performed equally well over a range of oxygen saturation from 18 % to 90 %. In contrast, Rouby et al.^[153] showed that the Oximetrix system was more accurate in patients with circulatory shock and respiratory failure than the other catheter systems.

Scuderi et al.^[154,155] evaluated three mixed venous oxygen saturation catheters in thirty critically ill patients. Improvements in the catheter design, computer algorithms, and optical modules warranted a new comparison study between the most recent versions of the different fiberoptic catheter systems. The three catheter systems involved were Oximetrix, Edwards SAT 2 and Viggo Spectramed. Readings were compared with a Co-oximeter IL 282. using a four wave length spectrophotometric technique. The mean difference and SD between the three fiberoptic catheters and the Co-oximeter were -1.98 ± 3.07 , 1.80 ± 3.49 and -2.28 ± 5.24 , respectively. The authors concluded, that Oximetrix and Edwards SAT-2 demonstrated a closer agreement with Co-oximetry than Viggo Spectramed. As a continuous monitor to detect changes or trends these devices are acceptable. However, none of these catheters provided evidence that they would perform within ± 2 per cent of Co-oximetry, as was specified by the manufacturer. In a subsequent study Scuderi et al.^[156] compared the three same fiberoptic catheters in a controlled laboratory environment using a blood flow loop with whole fresh human blood. The purpose of this study was to examine whether the limitations of these devices were inherent to the oximeter technology or to the confounding effects of a given patients anatomy and physiology. They found that the mean difference $\pm$ SD were 3.20 ± 2.47 , -1.25 ± 3.36 and -9.97 ± 7.05 for the Oximetrix, Edwards SAT 2 and Viggo Spectramed, respectively. Since in the controlled laboratory environment results were similar to the results in clinical studies, the authors conclude in the discussion that the lack of agreement between the catheter readings and the Co-oximetry are inherent to the fiberoptic reflection oximetry.

2.1.2.1 Accuracy of fiberoptic oximetry used in pulmonary artery catheter during different conditions.

Mixed venous oxygen, measured by the fiberoptic reflectance oximetry has been

reported to be affected by various conditions including hypertriglyceridemia, severe anaemia, free hemoglobin solutions, methemoglobinemia, carboxyhemoglobinemia, and hyperbilirubinemia.

Howdieshell et al. [157] described the reliability of *in vivo* mixed venous oxygen saturation measured by the Oximetrix system during the infusion of progressive concentrations of Intralipid. The authors conclude that *in vivo* mixed venous oxygen saturation remained constant and reliable during hypertriglyceridemia while *in vitro* measurement (CO-Oximeter; Instrumentation Laboratories, Lexington, Ma) were artifactually reduced by the falsely high values of methemoglobin. This is explained by the light scattering properties of triglycerides and its interference of the absorbance of light at the wavelength of methemoglobin. It is concluded that continuous *in vivo* oximetry remains reliable during lipid infusion.

Theoretically, severe anemia is expected to affect the accuracy of oximeter monitors. Fiberoptic reflectance oximetry estimate relative concentrations of oxyhemoglobin by measuring the ratios of reflected light intensity. In the absence of hemoglobin or with a small concentration of hemoglobin there should be no change in signal with changes in oxygenation or only a small change relative to the background noise. Woerkens et al. [158] evaluated the accuracy of a mixed venous oxygen saturation catheter (Spectracath) during acutely induced changes in hematocrit in humans. The accuracy of the studied fiberoptic catheter was not affected by changes in hematocrit or cardiac output ($r^2 = 0.87$). However, changes in hematocrit in this study were limited to hematocrit values between 18 % and 40 %. Lee et al. [159] described the accuracy of continuous oximetry (Oximetrix) during progressive isovolemic anemia. Data were collected while fractional oxygen inspired concentration (FiO_2) was varied from 1.00 to 0.05 in seven steps. The dogs underwent isovolemic hemodilution and the FiO_2 was again varied. the sequence continued until no data could be obtained. The accuracy of the catheter system (Oximetrix) was indicated by a mean difference and SD of -0.7 ± 8.6 %. The accuracy as assessed was similar for hematocrits between 15 % to 40 %. Below a hematocrit of 10 % the accuracy of fiberoptic oximetry became less acceptable: (mean difference $\pm$ SD: -11.5 ± 11.8).

Kong et al. [160] tested two pulmonary catheters (SAT 2 and Oximetrix, Opticath) on their ability to measure oxyhemoglobin saturations in haemoglobin solutions. The performances of both catheters were excellent in the measurement of oxygen saturation in whole blood and hemodiluted blood, however, when hemoglobin solutions are used, the performance is poor and oxygen saturation catheter readings are inaccurate. Increasing methemoglobin was found in the haemoglobin solutions and could partially explain the inaccurate readings of the fiberoptic oximetry.

The Oximetrix opticath was tested as a pulmonary artery catheter in five anesthetized dogs in which increasing levels of methemoglobin were induced [6]. In the presence of increasing levels of methemoglobin SvO_2 readings overestimated progressively the value of the oxygen saturation defined as the ratio between oxyhemoglobin and the hemoglobin available for oxygen transport, also called functional oxygen saturation. For example with a methemoglobin concentration > 40 % the mean difference $\pm$ SD between the catheter reading and the functional oxygen saturation determined by Co-oximetry was 24.1 ± 10.4 %.

The same group [7] examined the accuracy of the Oximetrix catheter in dogs ventilated with increasing inspired concentrations of carbon monoxide. Carboxyhemoglobin levels ranged from 0 % to 70 %. Mixed venous oxygen saturation readings measured in the pulmonary artery overestimated the value of oxygen saturation defined as the ratio between oxyhemoglobin and total hemoglobin also including dyshemoglobins as carboxyhemoglobin. Mixed venous oxygen saturation readings approximately equal

functional oxygen saturation defined as the ratio between oxyhemoglobin and hemoglobin available for oxygen transport.

In the first reports about accuracy^[145,146] it was already described that catheter readings were not dependent on the values of bilirubin up to 18 mg/dl.

Mault et al.^[161] described the use of the 4 Fr. Oximeter fiberoptic catheter as a monitor of the venous oxygen value of the ECMO circuit. The mean difference between the catheter readings and the Co-Oximetry were: $-0.2\% \pm 4.5\%$.

2.1.2.2 Summary

Pulmonary artery oximetry relies on complex optical phenomena involving absorption, reflection and refraction of light within a suspension of erythrocytes in plasma. The accuracy of the measurements also is affected by physiological factors such as hematocrit, erythrocyte velocity and turbulence, pulsatile bloodflow and artifact induced by catheter tip interaction with the blood vessel. The pulmonary catheters are said to yield test results similar to spectro-photometers as Radiometer OSM or Co-oximeter. Most data comparing pulmonary catheter readings and hemoximeters support that PA catheters are capable of providing clinically useful trending data. The most used catheters (Oximetrix and Edwards SAT2) are comparable in accuracy and reliability. However, these catheter devices cannot provide the accuracy of 2% as specified by the manufacturers. Fiberoptic catheter oximetry can be used to monitor the mixed venous oxygen saturation during various conditions and physiological disturbances. The limitations of the fiberoptic oximetry must be taken into account especially the relative broad dispersion of the values compared with Co-oximetry. Therefore, one must be cautious to interpret absolute values.

2.1.3 The position of the fiberoptic catheter for a reliable measurement of mixed venous oxygen saturation (SvO₂).

In small newborn infants venous oxygen saturation can be monitored easily and with much less patient risk by a catheter inserted through the umbilical vein into the right atrium than by a pulmonary artery catheter. However, catheter motion, chamber wall reflection, filtering technique to calculate oxygen saturation, intra-atrial left-to-right shunting and the possible insufficient mixing of the blood may affect the feasibility of the right atrium as a measurement site for mixed venous oxygen saturation.

In several studies central venous oxygen saturation (ScvO₂) measured in the right atrium or in one of the venae cavae has been examined as a substitute for SvO₂ measured in the pulmonary artery.

Barratt-Boyes and Wood^[162] described the values of oxygen saturation of blood in the venae cavae, right heart chambers and pulmonary vessels in 26 healthy subjects. The venous blood withdrawn from the inferior vena cava, the superior vena cava, right atrium, right ventricle and pulmonary artery equalled 83 (76 to 88), 76.8 (66 to 84), 79.5 (72 to 86), 78.5 (64 to 84) and 78.4 (73 to 85) per cent oxygen saturation, respectively. There is a slight but significant reduction of the oxygen saturation of the venous blood during its passage through the ventricle probably caused by the streamline flow of the coronary sinus draining into the right ventricle. However, the mean difference between oxygen saturation in the right atrium and in the pulmonary artery in paired samples was only 0.3 ± 3.5 per cent oxygen saturation. In another study Freed et al.^[163] examined the oxygen saturation values at different locations in the central venous circulation of the heart. They found as well that right atrial oxygen saturation was only 0.7 % higher than the pulmonary arterial saturation. As in the data from Barratt-Boyes and Wood^[162] the data in this study showed less variability in the distal samples, such as in the right ventricular outflow tract, suggesting better mixing in the distal part of the central venous circulation of the heart.

In animal studies oxygen saturation values obtained in the right atrium^[78] or in the superior vena cava^[79] were compared with oxygen saturation in the pulmonary artery. The studies were performed over a broad range of cardiorespiratory conditions including hypoxia, hemorrhage and resuscitation. In 464 paired data^[78] obtained by fiberoptic oximetry mean oxygen saturation in the right atrium ($54.1 \pm 10.6\%$) was very close to the value in the pulmonary artery ($53.2 \pm 11.1\%$). A close tracking of changes across a wide range of hemodynamic conditions between the two measurement sites were found. In 179 samples^[79] the mean difference between SvO_2 and $ScvO_2$ obtained in the superior vena cava was 3.7 ± 2.9 percent saturation. Additionally, more than 20 % of the $ScvO_2$ readings differed more than 5 % from the SvO_2 value.

In human clinical studies results are more contradictory. Berridge^[164] examined the influence of cardiac output on the correlation between mixed venous and central venous oxygen saturation. In 76 samples obtained in 51 patients he found a close correlation between $ScvO_2$ and SvO_2 irrespective of the cardiac output using a standard thermodilution technique. Tahvanainen et al.^[165] estimated the value of $ScvO_2$ compared with SvO_2 to calculate pulmonary shunt and arteriovenous oxygen difference. Central venous blood was sampled through the proximal port of a pulmonary artery catheter in critically ill patients. A significant positive correlation of the variables and of the subsequent changes were found. However, the broad confidence intervals allowed remarkable errors in the estimation of pulmonary shunts and other derived variables based on central venous oxygen saturations.

Martin et al.^[166] described in a clinical study in 7 critically ill patients the reliability of the $ScvO_2$ readings obtained in the lower part of the vena cava superior. The mean difference and SD was $0.3 \pm 10\%$ in 580 comparisons between SvO_2 and $ScvO_2$. Differences of $> 5\%$ were obtained in 50 % of the compared values. The authors concluded that the measurement of $ScvO_2$ can not substitute SvO_2 . Faber^[167] examined the relation between central venous and mixed venous oxygen content in 24 critically ill patients. A relatively poor association between central venous and mixed venous oxygen saturation was found: ($r = 0.75$). A proposed algorithm to estimate the value of mixed venous oxygen content from central venous oxygen content seemed of limited value. Brandt et al.^[168] disputed the use of central venous catheters as a substitute for mixed venous oxygen saturation measurement and considered the measurement in the right atrium as unreliable and inaccurate because of catheter motion and incomplete mixing of the venous blood.

We may conclude, that only slight differences exist between the right atrium oxygen saturation compared with the pulmonary artery oxygen saturation as determined during cardiac catheterization. In animal studies and smaller clinical studies these slight differences were confirmed. In clinical studies using the superior vena cava as measurement site differences were larger and it was concluded that $ScvO_2$ could not substitute SvO_2 measurement.

The drawbacks of the oxygen measurement in the right atrium are probably also valid in neonatal medicine. However, the measurement of venous oxygen saturation in newborn infants is limited to the right atrium, since the use of a pulmonary artery catheter is difficult and hazardous. Umbilical venous catheters have been used frequently with a relatively low incidence of complications and a fiberoptic catheter could be placed in the right atrium as a estimate of mixed venous oxygen saturation^[169].

The drawbacks of right atrial oxygen saturation measurement must be taken into account by careful consideration of the position of the catheter, frequent in vivo calibration and caution in the calculation of oxygen transport variables based on central venous oxygen saturations.

2.2 Pulse oximetry

2.2.1 Accuracy studies

Yelderman and New^[170] were among the first to conclude, that the pulse oximeter in 79 pooled data from five young adults accurately measures SaO_2 in the range of 70 % - 100 %. In pre(term) infants pulse oximetry (Nellcor N200) was assessed as a monitor of oxygenation and SaO_2 readings compared with the SaO_2 measurement by the hemoximeter showed a significant correlation ($r = 0.90$)^[171]. Hay compared the SaO_2 readings obtained by pulse oximetry (Ohmeda) with the arterial samples obtained by the hemoximeter. Regression analysis yielded a good correlation ($r = 0.99$, $p < 0.001$)^[172].

In these and in other studies accuracy was studied by regression analysis. However, regression analysis was found to be inadequate because a good correlation does not imply an accurate measurement^[173]. The mean difference and standard deviation between the pulse oximeter readings and the laboratory hemoximeter values are currently the preferred indices to assess accuracy^[174].

In several reports^[174-177] pulse oximeters of different brands were described and compared. In most studies the mean difference between the pulse oximeter readings and the laboratory hemoximeter values were below ± 3 % at SaO_2 values above 70 %. Accuracy at lower levels of saturation (SaO_2 values < 80 %) have been studied by Schmitt et al.^[178] in children with congenital cyanotic heart disease. Pulse oximetry overestimated (mean difference and SD: 5.8 % $\pm$ 4.8) the true value at lower levels of saturation. In a review^[174] the overestimation of pulse oximetry values at lower level of saturation was confirmed for pulse oximeters of different brands.

2.2.2 Different definitions of oxygen saturation

Another aspect of accuracy was discussed by Thilo et al.^[179] in a study in which she compared two different pulse oximeters (Nellcor and Ohmeda). Both oximeters were used simultaneously in the same infant with an arterial umbilical catheter. The Nellcor correlated best with the oxyhemoglobin defined as $(\text{O}_2\text{Hb})/(\text{O}_2\text{Hb} + \text{HHb})$, while Ohmeda correlated best with oxyhemoglobin defined as $(\text{O}_2\text{Hb})/(\text{O}_2\text{Hb} + \text{HHb} + \text{COHb} + \text{metHb})$. Some manufacturers apparently considered in the data read out the presence of a small percentage of dyshemoglobins, whereas others considered oxygen saturation the percentage of hemoglobin bound to oxygen compared with the hemoglobin capable of binding. These two "different" oxygen saturations were called fractional and functional oxygen saturations, respectively. However, these new terms were called unnecessary and self-contradictory^[1,2,4]. In the study of Thilo et al.^[179] this difference influenced the relation between paO_2 and SaO_2 value as registered by the pulse oximeter. A paO_2 between 50 and 100 mmHg was correlated with a Nellcor SaO_2 between 92 % and 98 % and with an Ohmeda SaO_2 between 90 % and 96 %. The authors concluded that different brands of pulse oximeters are not interchangeable.

2.2.3 Dyshemoglobins, anemia and pulse oximetry

The effect of dyshemoglobins as carboxyhemoglobin (COHb) and methemoglobin (MetHb) on pulse oximetry was studied by Barker and Tremper^[6,180]. With respect to COHb the pulse oximeter continued to read an oxygen saturation of 90 %, while the actual O_2Hb fell below 30 %. Pulse oximeter readings approximate the sum of COHb and O_2Hb and consequently overestimate the O_2Hb ^[180]. With respect to metHb^[6] pulse oximeter readings overestimated progressively the value of the oxygen saturation. In the presence of significant levels of dyshemoglobins, laboratory Co-oximetry should supplement pulse oximetry.

Severe anemia is, theoretically, expected to affect the accuracy of pulse oximetry. Lee et al.^[159] described the accuracy of pulse oximetry during progressive isovolemic anemia in dogs. Data were collected while FiO_2 was varied from 1.00 to 0.05 and isovolemic hemodilution was induced. Below a hematocrit of 10 % mean difference and SD worsened from $0.2 \% \pm 7.6$ to $-5.4 \% \pm 18.8$. Decreased accuracy with increasing level of anemia was confirmed by Severinghaus et al.^[181]

2.2.4 Accuracy in neonatal medicine

In contrast to previous suggestions fetal hemoglobin (HbF) does not affect the accuracy of pulse oximetry measurements, since absorption characteristics at the measured wavelengths are the same in fetal as in adult hemoglobin^[182,183].

High bilirubin plasma levels did not affect the SaO_2 readings of the pulse oximeters as evaluated by Veyckemans^[184] and Beall^[185]. Since pulse oximeters use 660 and 940 nm wavelength and bilirubin has a broad single peak at 460 nm, interference is from a theoretical point of view not likely. However, during the catabolism of hemoglobin bilirubin and carbon monoxide are formed. In this study^[184] icteric patients had a small higher level of carboxyhemoglobin than non-icteric patients ($1.8 \% \pm 0.7$ vs. $1.3 \% \pm 0.8$). Interference by phototherapy was studied by Zubrow^[186], but he could not find any such effect in a neonatal intensive care unit.

Iyer et al.^[187] evaluated the influence of hypothermia in neonates and infants undergoing cardiac surgery. The mean difference was within an acceptable range ($< \pm 3$ %) at foot probe site temperatures above 29°C . When foot probe site temperatures decreased below 27°C , 46 % of the pulse oximetry readings differed more than 3 % from hemoximeter oxygen saturations.

Skin pigmentation seemed to cause an overestimation of the oxygen saturation of the blood^[188]. However, as the skin of a newborn baby is less pigmented, limitations of pulse oximetry in neonatal care have not been reported.

A reduced accuracy in pulse oximetry has been reported due to motion artifact^[189]. In order to reduce motion artifact the ECG-signal was coupled to the oximeter to synchronize plethysmographic signal to heart rate. Barington et al.^[190] reported a 50 % reduction of false alarms due to motion artifact when ECG coupling was added in a neonatal intensive care unit. As pulse oximeter readings that are generated by motion artefact tend to be 85 % in accordance with a ratio (R) of 1.0, the plethysmographic signal amplitude is displayed to facilitate the interpretation of the saturation data of the pulse oximeter.

2.2.5 Usefulness of pulse oximetry

The usefulness of pulse oximetry has been established in several studies performed in the operating room, intensive care unit and emergency department. Coté et al.^[191] performed a study in pediatric patients divided in two groups, in which the oximeter data were either available or unavailable to the anesthesia team. The "unavailable group" experienced a greater number of major hypoxic events (24 vs. 11) and an increased total time of hypoxia (42 min vs. 27 min). The authors concluded, that pulse oximetry is efficient to provide an early warning of desaturation, prior to the development of signs and symptoms of hypoxemia. In another study^[192] the same group compared pulse oximetry and capnography. Blinding the oximeter data increased the number of desaturations (31 vs. 12) in contrast to blinding the capnograph data. In addition, Poets et al.^[193] determined the value of pulse oximetry in an observational study in the neonatal intensive care unit. Apparently well preterm (range 28 to 36 weeks) infants showed episodes of prolonged hypoxemia unaccompanied by apneic pauses or bradycardia and therefore undetectable for

cardiorespiratory monitors. Mower et al.^[194] gave evidence that the use of pulse oximetry in children presenting to emergency department triage may result in significant changes in medical treatment in a small proportion of children.

2.2.6 Summary

In summary, pulse oximetry is a useful and accurate monitor of arterial oxygen saturation. The accuracy is limited in the presence of dyshemoglobins, severe anemia and moderate to severe hypoxemia. In these conditions laboratory oximetry should supplement pulse oximetry.

IV Therapeutic Management to improve Oxygen Transport

1 Neonatal Oxygen Therapy

1.1 Historical notes

1.2 Normal values and limits to define hyperoxia and hypoxia.

1.2.1 Arterial oxygen tension: normal values in healthy (pre)term newborn infants

1.2.2 Arterial oxygen tension: normal values defined in relation to other variables.

1.2.2.1 Hyperoxic threshold

1.2.2.2 Hypoxic threshold

1.2.3 Arterial oxygen saturation: normal values in healthy (pre)term infants

1.2.4 Arterial oxygen saturation: normal values defined in relation to other variables

1.2.5 Conclusions

2 SvO₂ and DO₂ /VO₂ Relationship in Critical Care Medicine

2.1 Observational reports on SvO₂ measurement

2.2 SvO₂ as a monitor of cardiac output

2.3 Oxygen delivery (DO₂) / Oxygen consumption (VO₂) relationship

2.3.1 Biphasic relationship

2.3.2 Pathologic oxygen supply dependency

2.3.2.1 Etiologic mechanisms studied in the animal model

2.3.2.2 Criticism on the concept of pathologic oxygen supply dependency.

2.3.2.3 New studies examining the DO₂ /VO₂ relationship

2.3.2.4 The DO₂ /VO₂ relationship in pediatric and neonatal patients

2.4 SvO₂ measurement and the DO₂ / VO₂ relationship

2.5 Supranormal DO₂ and SvO₂ as goal-oriented therapy in critically ill patients

2.5.1 Studies to validate maximal DO₂ as goal-oriented therapy

2.5.2 SvO₂ as goal-oriented therapy

2.5.3 Criticism on supranormal DO₂ as a goal-oriented therapy

2.5.4 Summary

In this chapter among the many options of improving oxygen transport two important management strategies are discussed. Neonatal oxygen therapy, so directly associated with the history of neonatal medicine, will be addressed. In particular, the estimation of the normal paO_2 and SaO_2 range. Secondly, the debate about the DO_2 and VO_2 relationship and the strategy of optimizing DO_2 and SvO_2 in adult intensive care medicine will be reviewed.

1 Neonatal oxygen therapy

1.1 Historical notes

Oxygen therapy is one of the earliest applied therapies in neonatal medicine, but also one associated with numerous dilemmas^[14] and misadventures^[195]. The observation, that periodic breathing in preterm infants could be converted to regular breathing led in the decade 1942 to 1952 to the general policy to place all small infants in high oxygen environment. When the suspicion was raised, that this routine high oxygen treatment was related to retrolental fibroplasia with the blinding of numerous children, it was recommended that oxygen in concentrations greater than 40 % only should be used in exceptional circumstances. However, during the subsequent era of restricted oxygen use mortality and morbidity increased and it was suggested that each sighted baby gained may have cost 16 deaths^[196,197].

Oxygen administration increased again in the 60's. New reported side-effects as bronchopulmonary dysplasia and pulmonary hemorrhage stressed the need for monitoring oxygenation. Intermittent measurement of arterial oxygen tension (paO_2) in samples drawn from an umbilical arterial catheter became clinical practice in the 60's and 70's. The continuous measurement of arterial oxygen saturation (SaO_2) in newborn infants by a fiberoptic catheter used as an umbilical arterial catheter was reported by Wilkinson^[145] and Krouskop^[146]. Non-invasively and continuously transcutaneously measured paO_2 revealed the variability of paO_2 , particularly during the respiratory distress syndrome. In addition, the performance of this monitoring device seemed to be dependent upon perfusion of the skin, temperature of the electrode, patient's age and probe placement^[198]. Pulse oximetry, introduced in the 80's has now become common clinical practice in neonatal intensive care and has converted the clinician's attention from oxygen tension to oxygen saturation and oxygen content again.

1.2 Normal values and limits of the normal range as to define hyperoxia and hypoxia of arterial oxygen tension and arterial oxygen saturation.

When a new monitoring device is introduced in direct patient care it is necessary to describe normal values and to define the limits of the normal range. In general, studies trying to do this use two approaches:

- 1) Studies describe and examine the 'normal range' of data in a comparable but healthy population. However, it must be stressed, that from data in normal healthy infants it cannot be inferred that values below the normal range are pathologic and subsequently, require treatment.

- 2) Studies compare the values with a related variable, of which already normal values and limits have been described, or relate the value with the occurrence of unwanted effects.

1.2.1 Arterial oxygen tension (paO_2): normal values in healthy (pre)term newborn infants

With the advent of the continuous monitoring of transcutaneous paO_2 studies were performed to describe the paO_2 in term and preterm infants, the longitudinal development of paO_2 and the modulating effect of sleep and awake states. Mok^[199] examined transcutaneously measured paO_2 in 55 term infants from birth to 6 months to obtain normal data related to age and behavioral state. The mean values in the first week for the different states (awake, feeding, quiet sleep, active sleep) were: 83.5, 73.4, 78.5 and 73.4 mm Hg, respectively. The paO_2 increased until the age of 6 weeks. In preterm infants both Mok^[200] and Hoppebrouwers^[201] found that the mean paO_2 was lower in the preterm stage. The results from preterm infants at corrected postconceptional age could not be differentiated from those of control term infants.

1.2.2 Arterial oxygen tension (paO_2): normal values defined in relation to other variables.

1.2.2.1 Hyperoxic threshold

With regard to defining a hyperoxic threshold all attention has been focused to retrolental fibroplasia or retinopathy of prematurity (ROP). Retrolental fibroplasia was first described by Terry in 1942^[202]. The etiological role of oxygen in the pathogenesis of retrolental fibroplasia was recognized in the early 50's. The threshold of paO_2 at which ROP occurred was studied by Aranda et al^[203]. They correlated retinal vessel changes with the maximal value of paO_2 . The data in this study suggested that a paO_2 value above 100 mm Hg may result in retinal vascular changes. It was suggested, that the introduction of continuous non-invasive transcutaneous measurement of paO_2 might reduce ROP or might identify a threshold paO_2 value above which ROP might occur. However, Bancalari et al.^[204] were not able to show a clear evidence that continuous monitoring of paO_2 and adjusting the FiO_2 , aiming at a paO_2 range between 50 and 70 mm Hg, had an impact on the incidence of ROP in infants < 1000 grams. Flynn et al.^[205] could present evidence supporting an association between prolonged exposure to transcutaneously monitored paO_2 of 80 mm Hg or higher and the incidence and severity of ROP. Cunningham et al.^[206] found that an increased variability of transcutaneous paO_2 was a significant predictor of severe ROP.

In conclusion, since a transcutaneous measured paO_2 value of 80 mm Hg could be identified as one of the factors in the pathogenesis of ROP and since hemoglobin is maximally bound with oxygen at a paO_2 above 70 and 80 mm Hg^[14], there is no clinical or physiological basis to increase the paO_2 above this level.

1.2.2.2 Hypoxic threshold

With regard to defining a hypoxic threshold only few studies have been performed to assess the lower limit of arterial oxygen tension. It was noted that during the era of restricted oxygen in response to the first reports about retrolental fibroplasia mortality and neurological morbidity increased^[196,197]. However, these reports were non-randomized observations without the assessment of a critical paO_2 value. In a study of Tuschmidt et al.^[207] high altitude hypoxia was induced in young adults and paO_2 was compared with lactate, hypoxanthine and ST-T changes of the ECG. In contrast to hypoxanthine, which remained unchanged, lactate started to increase significantly in the paO_2 range of 40 to 50 mm Hg. The ST-T segment of the ECG was depressed during severe hypoxia ($\text{paO}_2 = 30$ mm Hg). Hrbek et al.^[208] recorded visual evoked responses in premature infants with idiopathic respiratory distress syndrome. He found, that the responses were more affected,

when paO_2 decreased below 55 mm Hg. Other reports have documented other adverse effects of low paO_2 in newborn infants such as the effect of a low paO_2 on pulmonary vasculature and ductus arteriosus. Moss et al. [209] performed one of the few human studies in the postnatal period in newborn infants breathing 13 % oxygen resulting in oxygen saturations, that varied between 46 % and 82 % corresponding to a paO_2 of 17 to 30 mmHg [210]. It was shown that these levels of paO_2 caused a rise in pulmonary arterial pressure and resistance.

Bard et al. [210] attempted to determine whether decreasing arterial blood saturation from 95 % to 90 % could cause pulmonary vasoconstriction and dilation of the ductus arteriosus in preterm newborn infants assessed by Doppler echocardiography. Saturations of 95 % and 90 % corresponded to a mean paO_2 of 87 ± 5 mm Hg and 55 ± 3 mm Hg, respectively. The decrease did not have any effect on pulmonary hemodynamics and ductus arteriosus. In addition, no differences were found between the two extremes in these groups of patients with either a paO_2 greater than 100 mm Hg or less than 50 mm Hg.

Émond et al. [211] examined in vitro the relation between SaO_2 and paO_2 in cord blood of ten newborn infants of less than 30 weeks and less than 1000 grams by gas mixing tonometry. The mean p_{50} and p_{90} were 18.3 ± 1.9 mm Hg and 40.8 ± 3.6 mm Hg. This study demonstrated the fetal nature of the oxygen dissociation curve of the very low birth weight infant and what paO_2 is needed for a 90 % oxygen saturation. The authors concluded that, since at this saturation the end of the steep rise of the hemoglobin dissociation curve is reached and the hemoglobin is almost completely saturated the guidelines of the AAP (50 to 80 mm Hg) [212] should probably be lowered to 45 to 60 mm Hg. The transfusion of adult packed cells could increase the p_{90} in preterm infants. However, since term infants with a fetal hemoglobin of approximately 70 % had a p_{90} of 47.6 ± 2.5 mm Hg, the limits of values of paO_2 inferred in this study are also valid for infants who have received several transfusions.

At last, also Schultze et al. [16] performed a non-randomized cross-over study by targeting the higher (97 ± 1.7 %) versus the lower (92 ± 3.8 %) end of the recommended range of pulse oximetry. These ranges of oxygen saturations corresponded to a mean paO_2 of 78 ± 14 mm Hg and 52 ± 6 mm Hg, respectively. VO_2 and cardiac output remained unchanged during the lower oxygen condition, whereas SvO_2 decreased and oxygen extraction increased. Targeting the lower range of SaO_2 resulted in an increased oxygen extraction whereas other potential compensatory mechanisms may not need to be called into play.

In conclusion, the lower limit of paO_2 during oxygen therapy in neonatal care could probably be defined at 45 mm Hg (6 kPa) since at this level approximately 90 % of the hemoglobin is bound to oxygen and since no adverse effects have been found such as pulmonary vasoconstriction or decreased VO_2 .

1.2.3 Arterial oxygen saturation (SaO_2): normal values in healthy (pre)term newborn infants.

Poets et al. [213] examined baseline SaO_2 monitored by pulse oximetry in combination with the analysis of breathing pattern in healthy full term infants in the second month of life. The pulse oximeter used was the Nellcor N 100 (beat to beat mode). The baseline SaO_2 during regular breathing was 99.8 % (97 -100) (median and range) with a low variability. The desaturation episodes ($\text{SaO}_2 < 80$ % during > 4 seconds) occurred particularly during non-regular breathing or apnoeic pause. The different reference range 92 to 93 % (CI 88 - 97) described by Thilo [214] in healthy term infants measured in the first 48 hrs could be explained by the use of a pulse oximeter of a different brand (Ohmeda) and the fact that the measurements were performed at an altitude of 1610 m.

In 66 preterm infants baseline SaO_2 values were examined at term age (37 weeks) by Poets et al. ^[215]. The median baseline SaO_2 was 99.4 % (88.9 - 100) with a higher variability than in full term infants. 12 % Of the preterm infants had a median SaO_2 value below 97 %. The desaturation episodes were more frequent in infants with a shorter gestation. The authors concluded, that during stable regular breathing preterm infants had baseline SaO_2 in the same range as the full term infants.

This last study was extended by a population of preterm infants and follow up recordings six weeks after discharge ^[216]. Baseline SaO_2 at a gestational age of 37 weeks in 160 infants was 99.6 % (88.7 - 100). In the follow up recordings SaO_2 increased to values of 100% (95.3 - 100) with a lower variability. When the follow up recordings were compared with the recordings made in full term infants at the same postconceptional age ^[213] a higher baseline SaO_2 in the preterm infants was found without a difference in the number of desaturation episodes. Maturation of respiratory control and improvement in the matching of ventilation and perfusion are thought to explain the increase of baseline SaO_2 and the decrease in desaturations.

All these studies were performed in infants at the time of discharge. Richard ^[217] obtained baseline SaO_2 measured by pulse oximetry in 55 preterm infants at 24 hours of age, breathing room air. The median gestational age was 35 weeks. Median baseline SaO_2 , measured only during regular breathing was 99.4 % (90.7 - 100). These results are similar to the results in older preterm infants. Less desaturation episodes were found compared with the study of preterm infants at discharge ^[216]. This can be explained by the fact that apnoeas with episodes of desaturation rarely are found before the third day of life.

The authors question in the discussion whether recommendations to keep SaO_2 at for example 87 to 92 % ^[218,219] can be considered appropriate, if 95 % of healthy preterm infants maintain a baseline SaO_2 at a level greater than 95%. They concluded that the results suggest that baseline SaO_2 in preterm infants is maintained at lower values than normal. On the other hand, the authors recognized that for infants receiving ventilatory support other recommendations have to be made than based upon the examination of healthy preterm infants. For example, in order to avoid hyperoxemia with oxygen tensions above 80 mm Hg the authors cannot recommend SaO_2 values above 95 %, although obviously these values are normal for healthy preterm infants during the first days of life.

To estimate normal SaO_2 values one has to take into account environmental factors as residency at high altitude and genetic adaptations of the population to a reduced oxygen availability. In a study ^[220] in Tibet (3658 m above sea level) SaO_2 was compared in 15 native Tibetan infants, whose ancestors inhabited the Himalayan plateau for 25.000 years and in 15 Han infants whose ancestors moved to Tibet after 1951. SaO_2 values were lower in Han infants than in Tibetan infants at all times, in all conditions and during all activities. At four month of age SaO_2 values of Han infants were $85 \% \pm 4$ while awake and $76 \% \pm 5$ during quiet sleep. In the Tibetan infants SaO_2 values were $88 \% \pm 4$ while awake and $86 \% \pm 5$ during quiet sleep.

1.2.4 Arterial oxygen saturation (SaO_2): normal values and alarm limits defined in relation to paO_2 .

Hay et al. ^[172,221] reviewed oxygen therapy in relation with the arrival of pulse oximetry in the neonatal intensive care unit. In this report SaO_2 values of $92 \pm 3 \%$ obtained by pulse oximetry (Ohmeda) excluded all paO_2 values below 45 mm Hg and above 100 mm Hg. Southall ^[222] compared pulse oximeter SaO_2 values, measured by the Nellcor N100, with direct arterial line oxygen saturation and with direct in line PO_2 . The patient population included 24 neonatal and 53 pediatric intensive care patients. Hypoxia defined as paO_2 values < 50 mm was detected by SaO_2 value below 90 % with a sensitivity

of 88 %, and hyperoxia defined as paO_2 values > 100 mm Hg was detected by SaO_2 above 98 % with a sensitivity of 90 %.

The problem how to detect the **hyperoxic threshold** by pulse oximetry in relation to arterial oxygen tension was considered in several reports^[223-226]. Walsh et al.^[223] evaluated the accuracy of pulse oximetry in acute and chronic preterm newborn infants by assessing the relation between SaO_2 and paO_2 . The authors concluded that in chronic infants pulse oximetry saturations and their derived paO_2 values were well correlated with measured paO_2 , whereas in acute infants pulse oximetry reflects paO_2 , unreliably. These infants had higher SaO_2 values, positioned at the flat upper part of the oxygen dissociation curve, where small changes in SaO_2 reflect large changes in the paO_2 . Bucher^[224] stated that the reliability of pulse oximetry to detect hyperoxemia is controversial, because small changes in oxygen saturation above 90 % are associated with relatively large changes in paO_2 . He examined the sensitivity and specificity of two different types of pulse oximeters (Nellcor and Ohmeda) in detecting hyperoxemia defined as $\text{paO}_2 > 90$ mm Hg. Sensitivity was defined as the proportion of hyperoxemia correctly classified by pulse oximetry and specificity the proportion of nonhyperoxemia correctly classified by pulse oximetry. In his study setting the alarm limit at 95 % the Nellcor N100 had 100 % sensitivity and 38 % specificity, whereas Ohmeda had a lower sensitivity (37 %) with a specificity of 83 %. To detect hyperoxemia the optimal alarm limit defined as a sensitivity of 95 % and a maximal specificity was for Nellcor 96 % and for Ohmeda 90 %.

Poets^[225] evaluated the detection of hyperoxia defined as a $\text{paO}_2 > 80$ mm Hg in 50 patients of the pediatric intensive care. From 202 similar measurements it was concluded that the upper alarm limit set at 95 %, 95 % of hyperoxemia, defined as a $\text{PO}_2 > 80$ mm Hg could be identified. These conclusions were drawn for the Nellcor N200 and in a population with a relatively right warded shift of the oxygen dissociation curve in comparison with a preterm newborn population. Therefore, one must be cautious to extrapolate these results to pulse oximeters of other manufacturers and to a neonatal intensive care setting.

Wasunna et al.^[218] evaluated pulse oximetry in preterm infants (range 26 - 32 weeks) at a postnatal age of 8 to 48 hrs. The pulse oximeter evaluated was the Ohmeda 3700. SaO_2 of 86 to 92 % corresponded to a paO_2 of 5-13 kPa. paO_2 was measured by an umbilical PO_2 electrode. SaO_2 values above 92 % were sometimes associated with paO_2 values to 14 kPa.

Paky and Koeck^[226] reached the same conclusions in a study performed to detect hyperoxemia and hypoxemia and to define alarm limits. A total of 792 SaO_2 readings and corresponding paO_2 from 146 infants were compared. Predictive value analysis of pulse oximetry readings related to arterial oxygen tension confirmed the ability to identify both hyperoxemia ($\text{paO}_2 > 90$ mm Hg) and hypoxemia ($\text{paO}_2 < 40$ mm Hg). However, alarm limits could not be established at a sensitivity level of 0.90. At a sensitivity level of 0.85 an alarm range was 92.5 - 95 % could identified. The authors concluded, that pulse oximetry alarm limits should be used with caution and one must be aware of the limitation to detect with a high level of sensitivity both hyperoxemia and hypoxemia.

1.2.5 Conclusions

At present pulse oximetry is the most used continuous oxygen monitor in neonatal intensive care medicine. In the observational studies, according to the studies on transcutaneously measured pO_2 , SaO_2 values were observed to increase with age. In these studies in healthy (pre)term infants rather high SaO_2 values are encountered. The current guidelines as published by manuals^[219], working groups^[227] and national pediatric associations^[212,228] provide limits for paO_2 values ranging between 40 (5.5 kPa) and 80 (11

kPa). From the studies mentioned it must be concluded that alarm limits with a sufficient sensitivity and specificity could hardly be deduced. In neonatal clinical medicine SaO_2 limits between 85-90 % and 92-96 % have been used. The discrepancy between the high SaO_2 values in healthy infants and the lower values in therapeutical guidelines led to some debate^[229,230]. In addition one must remember that different brands have used different calibration curves resulting in a different correlation between SaO_2 and paO_2 . Also blood transfusions and changes in pH can change this relationship. With these thoughts in mind the mentioned limits of SaO_2 can only be used supplemented by regular monitoring of blood gases with the measurement of paO_2 .

2 SvO_2 and DO_2 / VO_2 relationship in critical care medicine

2.1 Observational reports on SvO_2 measurement

Continuous SvO_2 measurement by a fiberoptic catheter was initially reported as a valuable warning system for deterioration in cardiopulmonary function and as an indicator of the effects of various therapeutic maneuvers^[148,77]. Divertie et al.^[149] reported that the ability to measure SvO_2 continuously has reduced the need for repeated measurements of cardiac output and blood gases.

2.2 SvO_2 as a monitor of cardiac output

In three small studies in patients with respiratory failure^[231] and with chronic congestive failure^[232,233] the relation between SvO_2 and cardiac output was evaluated and individual analysis yielded a good correlation in the majority of the patients. However, a number of other studies concluded that continuous monitoring of SvO_2 was not a sensitive measure of cardiac output. Kyff et al.^[234] described that decreases in SvO_2 by 5 % and 10 % corresponded with an decrease in cardiac index in only 45.5 and 61 % of measurements, respectively. Changes in cardiac index corresponded only in 30 % with changes in similar direction in patients with myocardial infarction. In another study during aortic surgery no significant correlation between SvO_2 and cardiac index could be found during the most critical periods of surgery^[235]. In two other studies^[236,237] in critically ill patients it was concluded that SvO_2 measurement could not substitute intermittent cardiac output and VO_2 determinations. It was recognized that changes in SvO_2 values may be assumed to be due to changes in cardiac output but also to changes in arterial oxygen saturation, O_2 uptake, hemoglobin, shift of the oxyhemoglobin dissociation curve or a combination of these variables. As mentioned by Nelson^[82], SvO_2 monitoring does not yield specific information regarding any individual oxygen transport variable. In addition, Jain et al.^[238] considered that the relation between SvO_2 and cardiac output is nonlinear according to the Fick equation. In his study on critically ill patients with cardiac failure a curvi-linear relationship was observed. Using linear regression the mean slope and correlation coefficient were dependent on whether the baseline SvO_2 and cardiac output were located on the steep or flat portion of the curvi-linear SvO_2 -cardiac output curve.

The curvi-linear relation between SvO_2 and cardiac index was established by Buheitel et al.^[240] in his study performed on a pediatric ICU. This group showed a uniform VO_2 and $\text{caO}_2(\text{Hb})$, which could explain the good correlation coefficient ($r = 0.90$). In a less uniform population Mahutte et al.^[241] compared the linear and curvi-linear regression analysis upon the relation between SvO_2 and cardiac output. Using the curvi-linear regression analysis the correlation coefficient improved, but remained rather poor ($r = 0.45$), due to the individual variability in SaO_2 , hemoglobin and VO_2 .

In summary, SvO_2 reflects the combined sufficiency of all the variables, that

constitute the oxygen delivery and uptake and cannot be used to monitor exclusively one of these variables, particularly when other variables are unstable. Due to the curvi-linear relationship between SvO_2 and cardiac output it is difficult to predict the change of one based on the other.

2.3 Oxygen delivery (DO_2) / Oxygen consumption (VO_2) relationship

2.3.1 Biphasic relationship

Animal experimental data in which the DO_2 has been progressively reduced either by hypoxia or by anemia, show a biphasic relationship between the DO_2 and VO_2 [114,115]. When DO_2 is reduced, the VO_2 remains constant over a wide range of DO_2 . Below a critical value of DO_2 , VO_2 starts to decrease and becomes dependent upon DO_2 . Shibutani et al. [116] established the biphasic relationship between DO_2 and VO_2 in anesthetized patients. The critical DO_2 could be identified at a DO_2 value of 300 ml/min/m². The phenomenon, that below the critical DO_2 a reduction of DO_2 leads to a decrease of VO_2 , is called "physiologic oxygen supply dependency".

2.3.2 Pathologic oxygen supply dependency

Komatsu et al. [242] compared the relationship between DO_2 and VO_2 in patients directly after cardiopulmonary bypass with and without increased lactate levels. In patients with low lactate levels VO_2 decreased proportionally, when DO_2 decreased below the critical DO_2 level of 300 ml/min/m². However, in patients with an increased lactate level no apparent critical DO_2 level could be identified. For the entire range of DO_2 values VO_2 decreased proportionally and a plateau phase could not be discerned. This phenomenon, that VO_2 is dependent upon DO_2 above the normal critical DO_2 is called "pathologic oxygen supply dependency" (see figure).

Pathologic oxygen supply dependency was observed in patients with ARDS [243,244], respiratory failure not due to ARDS [245,246], sepsis [247,248] and with advanced malignancy [249], after burn injury [250] cardiopulmonary resuscitation [242] and after acute liver failure and liver transplantation [251], and during congestive heart failure [252].

2.3.2.1 Etiologic mechanisms studied in the animal model

Several etiologic mechanisms that might account for this phenomenon of pathologic oxygen supply dependency have been suggested [253]. One of the main features is the inability to increase oxygen extraction in response to a reduction of DO_2 . In the experimental dog model induced bacteremia and endotoxemia produced an increased critical DO_2 and a decreased oxygen extraction at the whole body level and at the level of individual organs such as the gut [254] and in a lesser degree the hind limb muscle [107]. A peripheral oxygen extraction defect was explained by Bredle and Cain [255], who emphasized the importance of vasoconstrictor tone for oxygen extraction. They found an association between a lower oxygen extraction ratio and a flattening of the pressure / flow slope in the limb vasculature of endotoxin injected dogs indicating a loss of vasoconstrictor tone and vascular reactivity. Other mechanisms, such as a disorder of the redistribution of blood flow during a reduction of DO_2 and arteriovenous shunting of blood past nutritional capillaries [256] have been mentioned. Microembolization was demonstrated in experimental animals to limit the capillary surface reducing the oxygen extraction and to stimulate the release of the thromboxane A_2 [257], enhancing the effect of microembolization.

figure In "pathologic oxygen supply dependency" VO_2 is dependent upon DO_2 above the normal critical DO_2 in contrast to "physiologic oxygen supply dependency".

2.3.2.2 Criticism on the concept of pathologic oxygen supply dependency.

The question has been raised whether the disorders in the biphasic relationship between DO_2 and VO_2 as obtained in the endotoxic animal model by reducing DO_2 , are similar to the phenomenon of pathologic oxygen supply dependency in critically ill patients [258]. In animals the oxygen extraction failed to increase during a reduction in DO_2 , whereas in critically ill patients it was observed, that oxygen extraction failed to decrease when DO_2 was increased [253].

Another criticism was expressed by Russell and Phang [259] reviewing the methodology of the measurement of DO_2 and VO_2 in the clinical studies and the documented pathologic dependence of VO_2 upon DO_2 . Their criticism consisted of 1) the use of the indirect determination of VO_2 calculated as the product of cardiac output and arteriovenous oxygen content difference, 2) of the mathematical coupling of the shared variables (cardiac output, hemoglobin and arterial saturation) in the Fick equation of VO_2 and DO_2 and 3) of the use of pooled data of different patients.

2.3.2.3 New studies examining the $\text{DO}_2 / \text{VO}_2$ relationship

Therefore, clinical studies have been performed in patients with severe sepsis and/or acute respiratory failure [260-264] measuring VO_2 independently by calorimetry. In these studies the phenomenon of pathologic oxygen supply dependency could not be established. Vermeij [260] and Mira [261] comparing the indirect VO_2 with the direct VO_2 measured by calorimetry concluded that the use of indirect VO_2 led to an erroneous

conclusion concerning the relationship between DO_2 and VO_2 . Hanique et al. [263] reported that in 32 critically ill patients after volume loading the mean indirect VO_2 increased, whereas the mean direct VO_2 remained unaltered. Analysis of their indirect and direct determined VO_2 data [264] revealed that the two determinations did not provide similar results. Moreover, indirect calculations of the VO_2 were not as reproducible as the direct measurements. This was explained by the higher variability of the calculated VO_2 as a consequence of the cumulative effects of the errors of measurements of the different variables of the Fick equation.

In two more recent studies [265,266] carefully designed in relation with the direct measurement of VO_2 , pathologic oxygen supply dependency could be demonstrated in some patients. In these patients a critical DO_2 and plateau of VO_2 could not be observed. DO_2 was augmented by a dobutamine infusion [265] or by crystalloid and colloid infusions [266]. It was discussed that by increasing blood flow, organ metabolism and myocardial work may increase and consequently VO_2 .

2.3.2.4 The DO_2 / VO_2 relationship in pediatric and neonatal patients

In pediatric and neonatal intensive care studies on the relationship between DO_2 and VO_2 are limited. Lucking et al. [267] found in septic pediatric patients a linear relationship between DO_2 and VO_2 determined indirectly by calculation. Mink [268] and Dietrich [269] examined the metabolic response to increasing DO_2 by erythrocyte transfusion. VO_2 did not change, whereas oxygen extraction decreased. Sear et al. [270] studied the relationship in post-cardiac surgery infants receiving blood transfusions or adrenaline infusions. In post cardiac surgery infants VO_2 was not increased when DO_2 was increased by erythrocyte transfusion, whereas when DO_2 was increased by adrenaline infusion VO_2 increased significantly. This rise was thought to be due to increased cell metabolism analogous to exercise physiology. The authors concluded that the results did not support the concept of pathologic oxygen supply dependency.

2.4 SvO_2 measurement and the DO_2 / VO_2 relationship

These considerations have implications for the measurement of SvO_2 , since SvO_2 reflects the residual oxygen content after tissue VO_2 . When, in the plateau phase of the biphasic relationship, DO_2 is reduced, VO_2 can maintain at the same level by an increase of the oxygen extraction and consequently by a decrease of SvO_2 . Shibutani et al. [116] reported, that at DO_2 levels greater than 300 ml/min/m² a reduction of SvO_2 was proportional to a decrease of DO_2 . Below the critical DO_2 value VO_2 decreased and SvO_2 was not related to DO_2 . In several studies describing the phenomenon of pathologic oxygen supply dependency the poor correlation between DO_2 and SvO_2 was reported [242-250]. The measurement of SvO_2 , initially seen as an index for over-all tissue oxygenation was considered not accurately to reflect tissue oxygenation or changes in DO_2 or cardiac output [243]. In studies, in which VO_2 independently was measured, SvO_2 either increased proportionally [261,269] to DO_2 or was not related to DO_2 [265,266]. SvO_2 related to so different variables seemed to be of limited value.

In summary, the biphasic relationship between DO_2 and VO_2 has been confirmed in human clinical studies. The question remains whether the phenomenon of a pathologic oxygen supply dependency really exists. With the use of the direct independent measurement of the VO_2 pathologic oxygen supply dependency seemed to be of less frequent occurrence than initially was assumed. However, in individual critical ill patients VO_2 was shown to be dependent upon DO_2 with an impairment of the oxygen transport from the capillaries to the tissues and, consequently, a decreased oxygen extraction ratio. The measurement of SvO_2 will be of limited value, when the phenomenon of pathologic

oxygen supply dependency is present. According to the pulmonary artery catheter consensus statement ^[271] SvO₂ measurement may be considered in intensive care patients, when a pulmonary artery catheter is used and when therapy will be based upon oxygen transport end points, although randomized clinical outcome studies did not support the value of continuous venous oximetry.

2.5 Supranormal DO₂ and SvO₂ as goal-oriented therapy in critically ill patients

One of the therapeutic consequences of the concept of oxygen supply dependency was the idea to augment DO₂ to supranormal values, thereby increasing VO₂ and reversing tissue hypoxia and oxygen debt. This so-called goal-oriented supranormal DO₂ therapy was examined in several clinical trials ^[272-283].

Bland and co-workers ^[272-275] identified the different values of oxygen transport parameters between survivors and non-survivors in adult critical ill patients. In survivors cardiac index, DO₂ and VO₂ were greater than in non-survivors. It is noteworthy, that the mean value of cardiac index, DO₂ and VO₂ in non-survivors were within the normal range, whereas in survivors supranormal values were encountered. Therefore, it was concluded, that the goals of therapy should not be the normal values of unstressed, healthy subjects, but the supranormal values of survivors of life-threatening illness. Volume therapy and inotropic agents were used to increase cardiac index and DO₂.

2.5.1 Studies to validate maximal DO₂ as a goal-oriented therapy

Shoemaker et al. ^[276] tested the hypothesis, that the supranormal values of oxygen transport parameters may be valuable therapeutic goals for high-risk critically ill patients. He compared three groups of patients: a) with a central venous catheter and a group b) with a pulmonary artery catheter and c) with a pulmonary artery catheter and a treatment protocol to achieve supranormal oxygen transport values. Mortality and rate of complications were not different between the first two groups, but mortality was decreased in the pulmonary artery supranormal treatment group. The authors concluded, that for high-risk post-operative patients treatment aiming at supranormal oxygen transport values is more appropriate. Boyd et al. ^[277] confirmed in a prospective trial in high-risk surgical patients the reduced mortality (22 % vs. 6 %) of the supranormal DO₂ treatment strategy. The group of Shoemaker et al. ^[278,279] stressed the importance of the time frame in which the supranormal values are reached. In a prospective study ^[278] the protocol patients (supranormal DO₂ values within 24 hrs after admission) had a significant lower mortality (18 % vs. 37 %) than control patients.

Tuchsmidt et al. ^[280] and Yu et al. ^[281] did not find a significant reduction between the control and the supranormal DO₂ treatment group in a septic and a mixed group of patients, respectively. However, in a subset analysis between patients who actually achieved supranormal DO₂ values, whether self generated or as result from therapy, and patients with normal DO₂, a significant difference in mortality was found.

However, Hayes et al. ^[282] found in a prospective randomized trial even an increased mortality in the supranormal DO₂ treatment group (54 % vs. 34 %). The authors gave several reasons for these results: 1) VO₂ associated with the reversal of tissue hypoxia and oxygen debt was not increased in the treatment group; 2) in this study a patient population with a higher APACHE II score was encountered than in other studies; 3) in contrast to other studies patients in whom the therapeutic goals were only achieved by fluid administration were not assigned to the study. When all the patients, including those patients who were not assigned to the study, were considered in whom the target DO₂ levels were achieved, 90 % of the patients survived and left hospital. This supports the idea that the ability to achieve a high level of DO₂ and VO₂ indicates a larger physiological

reserve, less severe illness and a better prognosis ^[282]. In a retrospective study ^[283] of a subgroup of patients with sepsis this idea was confirmed. Survivors were characterized by the ability to increase both DO_2 and VO_2 , whereas non-survivors failed to increase VO_2 .

2.5.2 Normal SvO_2 values as goal-oriented therapy

Since SvO_2 reflects the balance between DO_2 and VO_2 consumption, studies evaluating treatment targeting supranormal levels of DO_2 and VO_2 included the measurement of SvO_2 . Normal SvO_2 values were defined as one of the treatment goals in critically ill patients. Gattinoni et al. ^[284] performed a randomized trial to determine whether targeting a normal value for SvO_2 ($> 70\%$) or supranormal value for cardiac index would improve morbidity and mortality in critically ill patients. 762 Patients were randomly assigned to the control group, to the cardiac index group or to the SvO_2 group. No differences in mortality and morbidity were found at the time of discharge or six month later. Subgroup analysis of the patients in whom hemodynamic targets were achieved, did not show any difference in mortality.

Kremzar et al. ^[285] determined in a retrospective non-randomized study in 40 patients with multiple injuries whether maintaining normal values for SvO_2 (70 -75 %) is more relevant to survival than maintaining supranormal oxygen transport parameters per se. In the SvO_2 group normal SvO_2 values were maintained by correcting hypovolemia, anaemia and hypoxemia and by the administration of dobutamine, which resulted in an average DO_2 value of 605 - 688 ml/min/m². In the supranormal DO_2 group an average DO_2 value of 835 ml/min/m² was achieved by continuous administration of fluids and increasing doses of dobutamine. An increased survival and decreased proportion of organ failure was found in the SvO_2 group (96 % *vs.* 65 %) and (4 % *vs.* 35 %), respectively. The authors concluded, that maintaining a normal SvO_2 and increasing DO_2 only when required is better than aiming at supranormal DO_2 levels.

2.5.3 Criticism on supranormal DO_2 as a goal-oriented therapy

Besides the conflicting results of the randomized trials more criticism on supranormal DO_2 treatment have been reported:

- 1) The use of catecholamines may be associated with an enhancement of VO_2 by direct stimulation of the tissue metabolism independent of its effect on DO_2 .
- 2) Arrhythmias and EKG changes have been reported in patients in whom high doses of catecholamines were used ^[282].
- 3) The frequency of myocardial infarction, a potential harm of the use catecholamines to maximize DO_2 , may increase in patients who receive catecholamines.
- 4) Aggressive fluid administration can induce fluid accumulation with impairment of the microcirculation and with lactate acidosis.
- 5) Some authors have argued that increasing DO_2 seems useless, since the uptake of oxygen at the tissue level is disturbed.

2.5.4 Summary

Debate about the efficacy of increasing DO_2 in critically ill patients still continues. The difficulty in several prospective trials to reach the supranormal target level of the oxygen transport parameters was mentioned. This difficulty seemed to reflect the different characteristics of the patients and consequently the different ability to achieve the aimed hemodynamic goal. However, adult intensive care centers have adopted this controversial treatment strategy. Several consensus statements ^[271,287] and a meta analysis the clinical trials ^[288] did not support aggressive attempts to increase DO_2 , although immediate resuscitation and achievement of normal oxygen transport variables are called essential. The SvO_2 as an

ingredient of the goal-oriented therapy has not introduced a new element in this debate. In the SvO₂ collaborative trial the oxygen transport parameters were indistinguishable in the control group and the SvO₂ group. The care in the control group resulted already in a normal SvO₂ value and was not better or worse than the care aimed at supranormal DO₂ levels. In the retrospective non-randomized trial the monitoring and targeting of normal SvO₂ values in combination with the dobutamine test seemed to be more promising. The conflicting results warrant further studies to validate normal SvO₂ values as a goal oriented therapy.

References

- 1 Zijlstra WG, Maas AHJ, Moran RF. Definition, significance and measurement of quantities pertaining to the oxygen carrying capacity of human blood. *Scand J Clin Lab Invest* 1996;56, Suppl 224:27-45
- 2 Moran RF. The case for standardized terminology: Oxygen saturation values can trick the unwary and lead to clinical misjudgment. *Crit Care Med* 1993;21:805-807
- 3 Wimberley PD, Siggaard-Andersen O, Fogh-Andersen N, Zijlstra WG, Severinghaus JW. Haemoglobin oxygen saturation and related quantities: definitions, symbols and clinical use. *Scand J Clin Lab Invest* 1990;50, Suppl 203:455-459
- 4 Zijlstra WG. Quantities and units in medical science: a plea for standardization. *Scand J Clin Lab Invest* 1993;53, Suppl 214:131-136
- 5 Zander R, Mertzluft F. Oxygen parameters of blood: definitions and symbols. *Scand J Clin Lab Invest* 1990;50, Suppl 203:177-185
- 6 Barker SJ, Tremper KK, Hyatt J. Effects of methemoglobin on pulse oximetry and mixed venous oximetry. *Anesthesiology* 1989;70:112-117
- 7 Haney M, Tait AR, Tremper KK. Effect of carboxyhemoglobin on the accuracy of mixed venous oximetry monitors in dogs. *Crit Care Med* 1994;22:1181-1185
- 8 Nunn JF. The atmosphere. In: Nunn's applied respiratory physiology. Oxford, Butterworth, Heinemann. 1993, pp 3-12
- 9 Kasting JF. Earth's early atmosphere. *Science Wash. DC.* 1993;259:920-926
- 10 Alberts B, Bray D, Lewis J, Raff M, Roberts Watson JD. The evolution of the cell. In: *Molecular biology of The Cell*, New York & London, Garland Pub Inc, 1993;pp 3-40
- 11 Alberts B, Bray D, Lewis J, Raff M, Roberts Watson JD. Energy conversion: Mitochondria and chloroplast. In: *Molecular biology of The Cell*, New York & London, Garland Pub Inc, 1993; pp 653-720
- 12 Nunn JF. Oxygen. In: Nunn's applied respiratory physiology. Oxford, Butterworth, Heinemann. 1993, pp 247-305
- 13 Ledingham IMcA, Naguib M. Overview: Evolution of the concept of Fick to the present day. In: Edwards JD, Shoemaker WC, Vincent JL. (eds.) *Oxygen transport. Principles and practice*. London, W.B. Saunders Co Ltd. 1993; pp 3-21
- 14 Duc G. Oxygen therapy. In: Vert P, Stern L. eds. *Neonatal Medicine*. Paris: Masson 1987: 505-537
- 15 Nunn JF. Distribution of pulmonary ventilation and perfusion. In: Nunn's applied respiratory physiology. Oxford, Butterworth, Heinemann. 1993, pp 178-197
- 16 Schultze A, Whyte RK, Clifton Way R, Sinclair JC. Effect of the arterial oxygenation level on cardiac output, oxygen extraction and oxygen consumption in low birth weight infants receiving mechanical ventilation. *J Pediatr* 1995;126:777-784
- 17 Krauss AH. Ventilation - perfusion relationships in neonates. In: Thibeault DW, Gregory GA. (eds.) *Neonatal pulmonary care*. Norwalk, Cn, Appleton. 1986: pp 123-148
- 18 Perutz MF. Molecular anatomy, physiology, and pathology of hemoglobin. In: Stamatoyanopoulos G, Nienhuis AW, Majerus PW, Varmus H. (eds): *The molecular basis of blood diseases*, Philadelphia, PA, Saunders, 1987, p.127-177
- 19 Wardrop CAJ, Holland BM, Jones JG. Red-cell physiology. In: Gluckman PJ, Heymann MA. (Eds.): *Perinatal physiology*. Philadelphia, PA, Saunders, 1997, p.868-876.
- 20 Severinghaus JW. Simple, accurate equations for human blood O₂ dissociation computations. *J Appl Physiol* 1979;43:599-602
- 21 Siggaard-Andersen O, Wimberley PD, Gothgen IH, Siggaard-Andersen M. A mathematical model of the hemoglobin-oxygen dissociation curve of human blood and of the oxygen partial pressure as a function of temperature. *Clin Chem* 1984;30:1646-1651

- 22 Brown EG, Krouskop RW, McDonnell FE, Monge CC, Winslow RM. A technique to continuously measure arteriovenous oxygen content difference and p_{50} in vivo. *J Appl Physiol* 1985;58:1383-1389
- 23 Wimberley PD. Fetal hemoglobin, 2,3-diphosphoglycerate and oxygen transport in the newborn premature infant. *Scand J Clin Lab Invest* 1982;42:Supp 160:13-149
- 24 Weber RE, Jensen FB. Functional adaptations in hemoglobin from ectothermic vertebrates. *J Exp Biol*
- 25 Giardina B, Condo SG, Petruzzelli R, Bardgard A, Brix O. Thermodynamics of oxygen binding to arctic hemoglobins: the case of reindeer. *Biophys Chem* 1990;37:281-286
- 26 Condo SG, Corda M, Sanna MT, Pellegrini MG, Ruiz MP, Castagnola M, Giardina B. Molecular basis of low-temperature instability in pig hemoglobins. *Eur J Biochem* 1992;209:773-776
- 27 Gustin P, Detry B, Cao ML, Chenut F, Robert A, Ansay M, Frans A, Clerbeaux T. Chloride and inorganic phosphate modulate binding of oxygen to bovine red blood cells. *J Appl Physiol* 1994;77:202-208
- 28 Zwart A, Kwant B, Oeseburg B, Zijlstra WG. Human whole-blood oxygen affinity: effect of carbon monoxide. *J Appl Physiol* 1984;57:14-20
- 29 Stamatoyannopoulos G, Nienhuis AW. Hemoglobin switching. In: Stamatoyannopoulos G, Nienhuis AW, Majerus PW, Varmus H. (eds): *The molecular basis of blood diseases*, Philadelphia, PA, Saunders, 1994, pp. 107-155
- 30 Cunningham JM, Jane SM. Hemoglobin switching and fetal hemoglobin reactivation. *Semin Hematol* 1996;33:9-23
- 31 Peterson KR, Stamatoyannopoulos G. Role of gene order in development control of human gamma- and beta-globin gene expression. *J Mol Cell Biol* 1993;13:4836-4843
- 32 Bard H, Widness JA. Effect of recombinant human erythropoietin on the switch over from fetal to adult hemoglobin synthesis in preterm infants. *J Pediatr* 1995;127:478-480
- 33 Bard H, Fouron JC, Prosmanne J, Gagnon J. Effect of hypoxemia on fetal hemoglobin synthesis during late gestation. *Pediatr Res* 1992;31:483-485
- 34 Bard H, Fouron JC, Gagnon C, Gagnon J. Hypoxemia and increased fetal hemoglobin synthesis. *J Pediatr* 1994;124:941-943
- 35 Bard H, Prosmanne J. Elevated levels of fetal hemoglobin synthesis in infants with bronchopulmonary dysplasia. *Pediatrics* 1990;86:193-196
- 36 Bard H, Lachance C, Widness JA, Gagnon C. The reactivation of fetal hemoglobin synthesis during anemia of prematurity. *Pediatr Res* 1994;36:253-256
- 37 Bard H, Fouron JC. The increase in fetal hemoglobin synthesis in the fetal lamb during hyperglycemic hypoxemia. *Am J Obstet Gynecol* 1988;159:1269-1272
- 38 Bard H, Widness JA, Ziegel EE, Gagnon C, Peri KG. The proportions of G-gamma and A-gamma globins in the fetal hemoglobin synthesized in preterm and term infants. *Pediatr Res* 1995;37:361-364
- 39 Bard H, Faesdale F. Red cell oxygen affinity, hemoglobin type, 2,3-diphosphoglycerate, and pH as a function of fetal development. *Pediatrics* 1979;64:483-487
- 40 Bard H, Fouron JC, Prosmanne J. The inhibition of the postnatal rise of 2,3-diphosphoglycerate in newborn lambs as a result of glucose perfusion. *Pediatr Res* 1988;24:470-472
- 41 Delivoria-Papadopoulos M, Martens RJ, Forster II RE, Oski FA. Postnatal changes in oxygen-hemoglobin affinity and erythrocyte 2,3-diphosphoglycerate in piglets. *Pediatr Res* 1974;8:64-66
- 42 Giardina B, Scatena R, Clementi M, Cerroni L, Nuutinen M, Brix O, Sletten SN, Castagnola M, Condo SG. Physiological relevance of the overall delta H of oxygen binding to fetal human hemoglobin. *J Mol Biol* 1993;229:512-516
- 43 Clementi ME, Scatena R, Mordente A, Condo SG, Castagnola M, Giardina B. Oxygen transport by fetal bovine hemoglobin. *J Mol Biol* 1996;255:229-234

- 44 Fouron JC, Bard H, Guennec Le JC, Van Ameringen MR. Effect of fetal or adult red cells on tissue oxygenation and myocardial function in normoxemic newborn lambs. *Pediatr Res* 1981;15:967-970
- 45 Van Ameringen MR, Fouron JC, Bard H, Le Guennec JC, Prossmanne J. Oxygenation in anemic newborn lambs with high or low oxygen affinity red cells. *Pediatr Res* 1981;15:1500-1503
- 46 Schumaker PT, Long R, Wood LDH. Tissue oxygen extraction during hypovolemia: role of hemoglobin p_{50} . *J Appl Physiol* 1987;62:1801-1807
- 47 Lister G. Oxygen transport in the intact hypoxic newborn lamb: Acute effects of increasing p_{50} . *Pediatr Res* 1984;16:172-177
- 48 Myburgh JA, Webb RK, Worthley LG. The p_{50} is reduced in critically ill patients. *Intensive Care Med* 1991;17:355-358
- 49 Oski FA. Clinical implications of the oxyhemoglobin dissociation curve in the neonatal period. *Crit Care Med* 1979;7:412-418
- 50 Woodson RD. Physiological significance of oxygen dissociation curve shifts. *Crit Care Med* 1979;7:368-373
- 51 Gutierrez G. The rate of oxygen release and the effect on capillary oxygen tension: a mathematical study. *Resp Physiol* 1986;63:79-96
- 52 Popel AS. Theory of oxygen transport to tissues. *Crit Rev Biomed Engin* 1989;17:257-321
- 53 Bos K. Mathematical modeling of oxygen transport from capillaries to tissue. Thesis. Catholic University Nijmegen, The Netherlands. 1997
- 54 Ellsworth ML, Pittman RN. Arterioles supply oxygen to capillaries by diffusion as well by convection. *Am J Physiol* 1990;258:H1240-H1243
- 55 Pittman RN. Influence of microvascular architecture on oxygen exchange in skeletal muscle. *Microcirculation* 1995;2:1-18
- 56 Hoofd L, Kreuzer F. Oxygen transfer from blood to mitochondria. In: Crystal RG, West JB, Weibel ER, Barnes PJ. (Eds.) *The lung. Scientific foundations. vol 2.* Lippincott-Raven, New York, pp 1913-1923
- 57 Fan FC, Chen RYZ, Schuessler GB. Effects of hematocrit variations on regional hemodynamics and oxygen transport in the dog. *Am J Physiol* 1980;238:H545-H552
- 58 Biro GP, Anderson PJ, Curtis SE. Stroma free hemoglobin: its presence does not improve oxygen supply to the resting hindlimb vascular bed of hemodiluted dogs. *Can J Physiol Pharmacol* 1991;69:1656-1662
- 59 Federspiel WJ, Salerius RH. An examination of the contribution of red cell spacing to the uniformity of oxygen flux at the capillary wall. *Microvasc Res* 1984;27:273-285
- 60 Popel AS. Invited editorial on "A finite model of oxygen diffusion in the pulmonary capillaries". *J Appl Physiol* 1997;82:1717-1718
- 61 Honig CR, Gayeski TEJ, Groebe K. Myoglobin and oxygen gradients. In: Crystal RG, West JB, Weibel ER, Barnes PJ. (Eds.) *The lung. Scientific foundations. vol 2.* Lippincott-Raven, New York, pp 1925-1933
- 62 Hoppeler H, Mathieu-Costello O, Kayar SR. Mitochondria and microvascular design. In: Crystal RG, West JB, Weibel ER, Barnes PJ. (Eds.) *The lung. Scientific foundations. vol 2.* Lippincott-Raven, New York, pp 1913-1923
- 63 Little RA, Edwards JD. Applied Physiology. In: Edwards JD, Shoemaker WC, Vincent JL. (Eds) *Oxygen Transport; principles and practice.* London, Saunders, 1993, pp 21-40
- 64 Babcock GT, Wilkström M. Oxygen activation and the conservation of energy in cell respiration. *Nature* 1992;356:301-308
- 65 Sokol RJ. Expanding spectrum of mitochondrial disorders. *J Pediatr* 1996;128:597-599
- 66 Johns DR. Mitochondrial DNA and disease. *N Engl J Med* 1995;333:638-644
- 67 Aprille JR. Perinatal development of mitochondria in rat liver. In *Physiology and pathology of mitochondria.* ed: Fiskum G. Van Vostrand Reinhold, New York 1986: 66-94

- 68 Valcarce C, Navarette RM, Encabo P, Loeches E, Satrustegui J, Cuezva JM. Postnatal development of rat liver mitochondrial functions. The roles of protein synthesis and of adenine nucleotides. *J Biol Chem* 1988;263:7767-7775
- 69 Joyal JL, Hagen T, Aprille JR. Intramitochondrial protein synthesis is regulated by matrix adenine content and requires calcium. *Arch Biochem Biophys* 1995;319:322-330
- 70 Izquierdo JM, Jimenez E, Cuezva JM. Hypothyroidism affects the expression of the β -F₁-ATPase gene and limits mitochondrial proliferation in rat liver at all stages of development. *Eur J Biochem* 1995;232:344-350
- 71 Prieur B, Bismuth J, Delaval E. Effects of adrenal steroid hormones on mitochondrial maturation during late fetal period. *Eur J Biochem* 1998;252:194-199
- 72 Bates TE, Almeida A, Heales SJ, Clark JB. Postnatal development of the complexes of the electron transport chain in isolated rat brain mitochondria. *Dev Neurosci* 1994;16:321-327
- 73 Schagger H, Noack H, Halngk W, Brandt U, von Jagow G. Cytochrome-C oxidase in developing rat heart. Enzymic properties and amino-terminal sequences suggest identity of the fetal heart and the adult liver isoform. *Eur J Biochem* 1995;230:235-241
- 74 Delaval E, Razanoelina M, Bastin J, Bismuth J, Geloso JP. Mitochondrial activity of rat kidney during ontogeny. *J Dev Physiol* 1990;14:1-5
- 75 Allen RG, Keogh BP, Tresini M, Gerhard GS, Volker C, Pignolo RJ, Horton J, Cristofalo VJ. Development and age associated differences in electron transport potential and consequences for oxidant generation. *J Biol Chem* 1997;272:24805-24812
- 76 Cernaianu AC, Nelson LD. The significance of mixed venous oxygen saturation and technical aspects of continuous measurement. In: Edwards JD, Shoemaker WC, Vincent JL. (Eds) *Oxygen Transport; principles and practice*. London, Saunders, 1993, pp 99-124
- 77 Kandel G, Aberman A. Mixed venous oxygen saturation. Its role in the assessment of the critical patient. *Arch Int Med* 1983;143:1400-1402
- 78 Davies GG, Mendenhall J, Symreng T. Measurement of right atrial oxygen saturation by fiberoptic oximetry accurately reflects mixed venous oxygen saturation in swine. *J Clin Monit* 1988;4:99-102
- 79 Reinhart K, Rudolph T, Bredle DL, Hanneman L, Cain SM. Comparison of central venous to mixed venous oxygen saturation during changes in oxygen supply/demand. *Chest* 1989;95:1216-1221
- 80 Schweiss JF. Mixed venous hemoglobin saturation: Theory and application. *Int Anesthesiol Clin* 1987;25:113-136
- 81 Kupeli IA, Satwicz PR. Mixed venous oximetry. *Int Anesthesiol Clin* 1989;27:176-183
- 82 Nelson LD. Continuous venous oximetry in surgical patients. *Ann Surg* 1986;329:329-333
- 83 Lehot JJ, Durand PG. Mixed venous oxygen saturation monitoring in surgery. In: Edwards JD, Shoemaker WC, Vincent JL. (Eds). *Oxygen Transport; principles and practice*. London, Saunders, 1993, pp 125-136
- 84 Zwischenberger JB, Kirsh MM, Dechet RE. Suppression of shivering decreases oxygen consumption and improves hemodynamic stability during postoperative rewarming. *Ann Thorac Surg* 1987;43:428-431
- 85 Schlittig R, Cowden LW, Chatman BR. Tolerance of unusually low mixed venous oxygen saturation: Adaptations in the chronic low cardiac output syndrome. *Am J Med* 1986;80:813-818
- 86 Räsänen J, Downs JB, Malec DJ, Seidman P. Estimation of oxygen utilization by dual oximetry. *Ann Surg* 1987; 206: 621-623
- 87 Carrol GC. A continuous monitoring technique for management of acute pulmonary failure. *Chest* 1987;92:467-469
- 88 Bongard FS, Leighton TA. Continuous dual oximetry in surgical critical care. Indications and limitations. *Ann Surg* 1992;216:60-68
- 89 Räsänen J. Role of dual oximetry in the assessment of pulmonary function. In: Reinhart K, Eylich K. (Eds) *Clinical aspects of O₂ transport and tissue oxygenation*. Springer Verlag, Berlin, 1989; 230-240

- 90 Reinhart K. Monitoring O_2 transport and tissue oxygenation in critically ill patients. In: Reinhart K, Eyrich K. Editors. Clinical aspects of O_2 transport and tissue oxygenation. Berlin: Springer Verlag, 1989; pp 193-211
- 91 Consensus conference. Tissue hypoxia. How to detect, how to correct how to prevent. Am J Respir Crit Care Med 1996;154:1573-1578
- 92 Finch CA, Lenfant C. Oxygen transport in man. New Engl J Med 1970;286:407-415
- 93 Connet RJ, Honig CR, Gayeski TEJ, Brooks GA. Defining hypoxia: a systems view of VO_2 , glycolysis, energetics, and intracellular pO_2 . J Appl Physiol 1990;68:833-842
- 94 Robin ED. Of men and mitochondria: Coping with hypoxic dysoxia. Am Rev Respir Dis 1980;122:517-531
- 95 Shapiro BA, Cane RD. Blood gas monitoring: Yesterday, today, and tomorrow. Crit Care Med 1989;17:573-581
- 96 Shapiro BA. Assessment of oxygenation: Today and tomorrow. Scan J Clin Invest 1990;50:197-202
- 97 Gutierrez G. Cellular energy metabolism during hypoxia. Crit Care Med 1991;19:619-626
- 98 Gutierrez G. Cellular effects of hypoxia and ischemia. In: Crystal RG, West JB, Weibel ER, Barnes PJ. eds. The Lung. Scientific foundations. Philadelphia, Lippincott-Raven, 1997, pp 1969-1987
- 99 Sahlin K. Aerobic and anaerobic processes of energetics. In: Crystal RG, West JB, Weibel ER, Barnes PJ. eds. The Lung. Scientific foundations. Philadelphia, Lippincott-Raven, 1997, pp 1889-1899
- 100 Cain SM, Curtis SE. Systemic and regional oxygen uptake and delivery and lactate flux in endotoxic dogs infused with dexamethasone. Crit Care Med 1991;19:1552-1560
- 101 Balaban RS. Regulation of oxidative phosphorylation in the mammalian cell. Am J Physiol 1990;258:C377-C385
- 102 Guyton AC. Metabolism of carbohydrates and formation of adenosine triphosphate. In Guyton AC. Textbook of medical physiology. 7th edition. Philadelphia, Saunders 1986, pp 808-817
- 103 Kreisberg RA. Lactate homeostasis and lactic acidosis. Ann Intern Med 1980;92(1):227-237
- 104 Mizock BA. Lactic Acidosis. Disease-a-Month 1989;4:235-300
- 105 Mizock BA, Falk JL. Lactic acidosis in critical illness. Crit Care Med 1992;20:80-93
- 106 Wasserman K, Koike A. Is the anaerobic threshold truly anaerobic. Chest 1992;101:2115-2185
- 107 Samsel RW, Schumaker PT. Determination of the critical O_2 delivery from experimental data: sensitivity to error. J Appl Physiol 1988;64:2074-2082.
- 108 Barefield ES, OH W, Stonestreet BS. Group B streptococcus-induced acidosis in newborn swine: regional oxygen transport and lactate flux. J Appl Physiol 1992;72:272-277
- 109 Fitzgerald MJ, Goto M, Myers TF, Zeller WP. Early metabolic effects of sepsis in the preterm infant: Lactic acidosis and increased glucose requirement. J Pediatr 1992;121:951-955
- 110 Izraeli S, Ben-Sira L, Harel D, Naor N, Ballin A, Davidson S. Lactic acid as a predictor for erythrocyte transfusion in healthy preterm infants with anemia of prematurity. J Pediatr 1993;122:629-631.
- 111 Cheung PY, Finer NN. Plasma lactate concentration as a predictor of death in neonates with severe hypoxemia requiring extracorporeal membrane oxygenation. J Pediatr 1994;125:753-758
- 112 Ruth V, Raivio KO. Perinatal brain damage: predictive value of metabolic acidosis and the apgar score. BMJ 1988;297:24-27
- 113 Wilmore JH, Costill DL. Energy for movement. Basic energy systems. In: Wilmore JH, Costill DL. Physiology of sport and exercise. Campaign, Human kinetics. 1994; pp 94-120.
- 114 Cain SM. Oxygen delivery and uptake in dogs during anemic and hypoxic hypoxia J Appl Physiol 1977;42:228-234
- 115 Cain SM. Effects of time and vasoconstrictor tone on oxygen extraction during hypoxic hypoxia. J Appl Physiol 1978;45:219-223
- 116 Shibutani K, Komatsu T, Kubal K, Sanchala V, Kumar V, Bizzari DV. Critical level of oxygen delivery in anesthetized man. Crit Care Med 1983;11:640-643

- 117 Schlichtig R, Pinsky MR. Defining the hypoxic threshold. *Intensive Care Med* 1991;19:147-149
- 118 Torrance SM, Wittnich C. The effect of varying arterial oxygen tension on neonatal acid-base balance. *Pediatr Res* 1992;31:112-116
- 119 Torrance SM, Wittnich C. Blood lactate and acid-base balance in graded neonatal hypoxia: evidence for oxygen-restricted metabolism. *J Appl Physiol* 1994;77:2318-2324
- 120 Moss M, Moreau G, Lister G. Oxygen transport and metabolism in the conscious lamb: the effects of hypoxemia. *Pediatr Res* 1987;22:177-183
- 121 Walker AM, Cannata JP, Dowling MH, Ritchie BC, Maloney JE. Age-dependent pattern of autonomic heart rate control during hypoxia in fetal and newborn lambs. *Biol Neonate* 1979;35:198-208
- 122 Giussani DA, Spencer JAD, Hanson MA. Fetal cardiovascular responses to hypoxemia. *Fetal and Maternal Medicine Review* 1994;6:17-37.
- 123 Bartelds B, van Bel F, Teitel DF, Rudolph AM. Carotid not aortic, chemoreceptors mediate the fetal cardiovascular response to acute hypoxemia in lambs. *Pediatr Res* 1993;34:51-55
- 124 Sidi D, Kuipers JRG, Teitel D, Heymann MA, Rudolph AM. Developmental changes in oxygenation and circulatory responses to hypoxemia in lambs. *Am J Physiol* 1983;245:H674-H682
- 125 Koehler RC, Traystman RJ, Jones Jr MD. Regional bloodflow and O₂ transport during hypoxic and CO hypoxia in neonatal and adult sheep. *Am J Physiol* 1985;245:H118-H124
- 126 Teitel D, Sidi D, Bernstein D, Heymann MA, Rudolph AM. Chronic hypoxemia in the newborn lamb: Cardiovascular, hematopoietic and growth adaptations. *Pediatr Res* 1985;19:1004-1010
- 127 Bernstein D, Teitel D, Sidi D, Heymann MA, Rudolph AM. Redistribution of regional bloodflow and oxygen delivery in experimental cyanotic heart disease in newborn lambs. *Pediatr Res* 1987;22:389-393
- 128 Trouwborst A, Tenbrink R, Fennema M, Bucx M, Broek WGM v.d, Trouwborst-Weber BK. Cardiovascular responses, hemodynamics and oxygen transport to tissue during moderate isovolemic hemodilution in pigs. In Piiper J. eds. *Oxygen transport to the tissues XII*. Plenum Press, New York, 1990:873-878
- 129 Trouwborst A, Woerkens ECSM van, Tenbrink R. Hemodilution and oxygen transport. In Erdman W, Bruley DF. eds. *Oxygen transport to the tissues XIV*. Plenum Press, New York, 1992:431-440
- 130 Robertie PG, Gravlee GP. Safe limits of isovolemic hemodilution and recommendations for erythrocyte transfusion. *Int Anesthesiol Clin* 1990;28(4):197-204
- 131 Cain SM. Peripheral oxygen uptake and delivery in health and disease. *Clinics in Chest Med* 1982;4:139-148
- 132 Kreuzer F, Cain SM. Regulation of the peripheral vasculature and tissue oxygenation in health and disease. *Critical Care Clin* 1985;1:453-470
- 133 Bernstein D, Teitel DF, Rudolph AM. Chronic anemia in the newborn lamb: Cardiovascular adaptations and comparison to chronic hypoxemia. *Pediatr Res* 1988;23:621-627
- 134 Fahey JT, Lister G. Postnatal changes in critical cardiac output and oxygen transport in conscious lambs. *Am J Physiol* 1987;253:H100-H106
- 135 Heusser F, Fahey JT, Lister G. Effect of hemoglobin concentration on critical cardiac output and oxygen transport. *Am J Physiol* 1989;253:H527-H532
- 136 Fahey JT, Lister G. Response to low cardiac output: Developmental differences in metabolism during oxygen deficit and recovery in lambs. *Pediatr Res* 1989;26:180-187
- 137 Moss M, Kurzner S, Razlog Y, Lister G. Hypoxanthine and lactate concentrations in lambs during hypoxic and stagnant hypoxia. *Am J Physiol* 1988;255:H53-H59
- 138 Zijlstra W. G. Oximetry: A historical introduction. In: *Fetal and neonatal physiological measurements*. Ed: Lafeber HN. Elsevier Science Publisher. Amsterdam 1991:97-103
- 139 Severinghaus JW, Astrup P. History of blood gas analysis. *Oximetry*. *Int Anesthesiol Clin* 1987;25(3):167-204

- 140 Thilo EH, Brockway Curlander J, Hay WW. Pulse oximetry. In: Physiological monitoring and instrumental diagnosis in perinatal and neonatal medicine. Eds: Brans YW, Hay WW. Cambridge University Press, Cambridge. 1995:147-161
- 141 Hay WW, Thilo EH, Brockway Curlander J. Pulse oximetry in neonatal medicine. *Clinics in perinatology* 1991;18:441-472
- 142 Pologe JA. Pulse oximetry: Technical aspects machine design. *Int Anesthesiol Clin* 1987;25(4):137-175
- 143 Landsman MLJ, Knop N, Kwant G, Mook GA, Zijlstra WG. A fiberoptic reflection oximeter. *Pfluegers Arch* 1978;373:273-282
- 144 Mook GA, Osypka P, Strum RE, Wood EH. Fiberoptic reflection photometry on blood. *Cardiovasc Res* 1968;2:199-209
- 145 Wilkinson AR, Phibbs RH, Gregory GA. Continuous measurement of oxygen saturation in sick newborn infants. *J Pediatr* 1978;93:1016-1090
- 146 Krouskop R, Cabatu EE, Chelliah BP, McDonnell FE, Brown EG. Accuracy and clinical utility of an oxygen saturation catheter. *Crit Care Med* 1983;11:744-749
- 147 Haessler R, Brandl F, Zeller M, Briegel J, Peter K. Continuous intra-arterial oximetry, pulse oximetry and Co-oximetry during cardiac surgery. *J Cardiathorac Vasc Anesth* 1992;6:668-673
- 148 Baele PL, McMichan JC, Marsh HM, Sill JC, Southorn PA. Continuous monitoring of mixed venous oxygen saturation in critically patients. *Anesth Analg* 1982;61:513-517
- 149 Divertie MB, McMichan JC. Continuous monitoring of mixed venous oxygen saturation. *Chest* 1984;85:423-428
- 150 Gettinger A, DeTrailia MC, Glass D. In vivo comparison of two mixed venous saturation catheters. *Anesthesiology* 1987;66:373-375
- 151 Reinhart K, Moser N, Rudolph T, Bredle D, Specht M, Gramm HJ, Goeke J, Eyrich K. Accuracy of two mixed venous saturation catheters during long-term use in critically ill patients. *Anesthesiology* 1988;69:769-773
- 152 Leighton T, Liu SY, Lee TS, Klein S, Bongard F. Simultaneous in-vivo comparison of 2-versus 3-wavelength mixed venous oximetry catheters. *Anesthesiology* 1991;75:A408
- 153 Rouby JJ, Poete P, Bodin L, Bourgeois JL, Arthaud M, Viars P. Three mixed venous oxygen catheters in patients with circulatory shock and respiratory failure. *Chest* 1990;98:954-958
- 154 Scuderi PE, Bowton DL, Meredith JW, Harris LC, Brockschmidt J. Clinical evaluation of three mixed venous oxygen saturation catheters. *Anesthesiology* 1990;V73:A484
- 155 Scuderi PE, Bowton DL, Meredith JW, Harris LC, Brockschmidt Evans J, Anderson RL. A comparison of three pulmonary artery oximetry catheters in intensive care unit patients. *Chest* 1992;102:896-905
- 156 Scuderi PE, MacGregor DA, Bowton DL, James RL. A laboratory comparison of three pulmonary artery oximetry catheters. *Anesthesiology* 1994;81:245-253
- 157 Howdieshell TR, Sussman A, Dipiro J, McCarten M, Mansbergen A. Reliability of in vivo mixed venous oximetry during experimental hypertriglyceridemia. *Crit Care Med* 1992;20:999-1004
- 158 Woerkens ECSM, Trouwborst A, Tenbrink R. Accuracy of a mixed venous oxygen saturation catheter during acutely induced changes in hematocrit in humans. *Crit Care Med* 1991;19:1025-1029
- 159 Lee S, Tremper KK, Barker SJ. Effects of anemia on pulse oximetry and continuous mixed venous hemoglobin saturation monitoring in dogs. *Anesthesiology* 1991;75:118-122
- 160 Kong CS, Ryder IG, Kahn R, Gregory L, Mackenzie CF. In vitro oxyhemoglobin saturation measurements in haemoglobin solutions using fiberoptic catheters. *Br J Anaesth* 1995;74:201-208
- 161 Mault J.R, Santoro-Nease A, Leonard R.A, Ungerleider R.M. Continuous fiberoptic venous oximetry during neonatal ECMO: Analysis of accuracy and longevity. *Crit Care Med* 1992;20:S11
- 162 Barratt-Boyes BG, Wood EH. The oxygen saturation of blood in the venae cavae, right-heart chambers and pulmonary vessels of healthy subjects. *J Lab Clin Med* 1957;50:93-106
- 163 Freed M, Miettinen OS, Nadas A. Oximetric detection of intracardiac left to right shunts. *Br Heart J* 1979;42:690-694

- 164 Berridge JC. Influence of cardiac output on the correlation between mixed venous and central venous oxygen saturation. *Br J Anaesthesia* 1992;69:409-410.
- 165 Tahvanainen J, Meretoja O, Nikki P. Can central venous blood replace mixed venous blood samples. *Crit Care Med* 1982;10:758-761.
- 166 Martin C, Auffray JP, Badetti G, Papazian L, Gouin F. Monitoring of central venous oxygen saturation versus mixed venous oxygen saturation in critically ill patients. *Intensive Care Med* 1992;18:101-104.
- 167 Faber T. Central venous versus mixed venous oxygen content. *Acta Anaesthesiol Scand* 1995;39:S107:33-36.
- 168 Brandt L, Mertzlufft F. Zur aussagekraft "zentral venoser" blutproben. "Zentralvenoser" vs. gemischtvenoser O₂ status. *Anaesthesist* 1991;40:131-144.
- 169 Dudell G, Cornisch JD, Bartlett RH. What constitutes adequate oxygenation? *Pediatrics* 1990;85:39-41.
- 170 Yelderman M, New W. Evaluation of pulse oximetry. *Anesthesiology* 1983;349-352.
- 171 Jennis MS, Peabody JL. Pulseoximetry: an alternative method for the assessment of oxygenation in newborn infants. *Pediatrics* 1987;79:524-528.
- 172 Hay WW, Brockway Curlander J, Eyzaguirre M. Neonatal pulse oximetry: Accuracy and reliability. *Pediatrics* 1989;83:717-722.
- 173 Bland JM, Altman DG. Statistical methods for assessing agreement between two methods of clinical measurement. *Lancet* 1986;i:307-317.
- 174 Severinghaus JW, Kelleher JF. Recent developments in pulse oximetry. *Anesthesiology* 1992;76:1018-1038.
- 175 Choe H, Tashiro C, Fukumitsu K, Masahru Y, Yoshiya I. Comparison of recorded values from six pulse oximeters. *Crit Care Med* 1989;17:678-681.
- 176 Nickerson BG, Sarkisan C, Tremper K. Bias and precision of pulseoximeters and arterial oximeters. *Chest* 1988;93:515-517.
- 177 Bowes WA, Corke BC, Hulka J. Pulse oximetry: A review of the theory, accuracy, and clinical applications. *Obstet Gynec* 1989;74:541-546.
- 178 Schmitt HJ, Schüetz WH, Proeschel PA, Jaklin C. Accuracy of pulse oximetry in children with cyanotic congenital heart disease. *J Cardiothoracic Vasc Anesth* 1993;7:61-65.
- 179 Thilo EH, Anderson D, Wasserstein ML, Schmidt J, Luckey D. Saturation by pulse oximetry: Comparison of the results obtained by instruments of different brands. *J Pediatr* 1993;122:620-626.
- 180 Barker SJ, Tremper KK. The effect of carbon monoxide inhalation on pulseoximetry and transcutaneous pO₂. *Anesthesiology* 1987;66:677-679.
- 181 Severinghaus JW, Koch SO. Effect of anemia on pulse oximeter accuracy at low saturation. *J Clin Monit* 1990;6:85-88.
- 182 Pologe J, Raley D. Effects of fetal hemoglobin on pulse oximetry. *J Perinatol* 1987;7:324-326.
- 183 Rajadurai VS, Walker AM, Yu VY, Oates A. Effect of fetal hemoglobin on the accuracy of pulse oximetry in preterm infants. *J Paediatr Child Health* 1992;28:43-46.
- 184 Veyckemans F., Baele P, Guillaume JE, Willems E, Robert A, Clerboux T. Hyperbilirubinemia does not interfere with hemoglobin saturation measured by pulse oximetry. *Anesthesiology* 1989;70:118-122.
- 185 Beall SN, Moorthy SS. Jaundice, oximetry, and spurious hemoglobin desaturation. *Anesth Analg* 1989;68:806-807.
- 186 Zubrow AB, Henderson GW, Imaizumi SO, Pleasure JR. Pulse oximetry in sick premature infants and the effect of phototherapy and radiant warmers in the oxygen saturation read out. *Am J Perinatol* 1990;7:75-78.
- 187 Iyer P, McDougall P, Loughan P, Mee RB, Al-Tawill K, Carlin J. Accuracy of pulse oximetry in hypothermic neonates and infants cardiac surgery. *Crit Care Med* 1996;24:507-511.
- 188 Volgyesi GA, Spahr-Schopfer I. Does skin pigmentation affect the accuracy of pulse oximetry? An in vitro study. *Anesthesiology* 1991;75:A407.

- 189 Barker SJ, Shah NK. Effects of motion on the performance of pulse oximeters in volunteers. *Anesthesiology* 1996;85:774-781
- 190 Barington KJ, Finer NN, Ryan CA. Evaluation of pulse oximetry as a continuous monitor technique in the neonatal intensive care unit. *Crit Care Med* 1988;16:1147-1153
- 191 Coté CJ, Goldstein EA, Coté MA, Hoaglin DC, Ryan JF. A single blind study of pulse oximetry in children. *Anesthesiology* 1988;68:184-188
- 192 Coté CJ, Rolf N, Guodsouzian NG, Ryan JF, Zavlavnsky A, Gore R, Todres TD, Vasallo S, Polaner D. A single blind study of combined pulse oximetry and capnography in children. *Anesthesiology* 1991;74:980-987
- 193 Poets CF, Stebbens VA, Richard D, Southall DP. Prolonged episodes of hypoxemia in preterm infants undetectable by cardiorespiratory monitors. *Pediatrics* 1995;95:860-863
- 194 Mower WR, Sachs C, Nicklin EL, Baraff LJ. Pulse oximetry as a fifth pediatric vital sign. *Pediatrics* 1997;99:681-686
- 195 Silverman WA. Setting a limit in the treatment of neonates: all or none? In: Duc G, Huch A, Huch R. eds. The very low birth weight infant; A challenge to neonatology and obstetrics. New York: Georg Thieme Verlag, 1990:111-128
- 196 Cross KW. Costs of preventing retrolental fibroplasia. 1973;15:954-956
- 197 Bolton DPG, Cross KW. Further observations on costs of preventing retrolental fibroplasia. *Lancet* 1974;445-448
- 198 Poets CF, Southall DP. Non-invasive monitoring of oxygenation in infants and children: practical considerations and areas of concern. *Pediatrics* 1994;93:737-745
- 199 Mok JYQ, McLaughlin FJ, Pintar M, Hak H, Amaro-Galvez R, Levison H. Transcutaneous monitoring of oxygenation: what is normal? *J Pediatr* 1986;108:365-371
- 200 Mok JYQ, Hak H, McLaughlin FJ, Pintar M, Canny GJ, Levison H. Effect of age and state of wakefulness on transcutaneous oxygen values in preterm infants; A longitudinal study. *J Pediatr* 1988;113:706-709
- 201 Hoppenbrouwers T, Hodgeman JE, Arakawa K, Durand M, Cabal LA. Transcutaneous oxygen and carbon dioxide during the first half year of life in premature and normal term infants. *Pediatr Res* 1992;31:73-79
- 202 Terry TL. Extreme prematurity and fibroblastic overgrowth of persistent vascular sheath behind each crystalline lens. I. Preliminary report. *Amer J Ophthal* 1942;25:203-204
- 203 Aranda JV, Saheb N, Stern L, Avery ME. Arterial oxygen tension and retinal vasoconstriction in newborn infants. *AJDC* 1971;122:189-194
- 204 Bancalari E, Flynn J, Goldberg RN, Bawol R, Cassady J, Schiffman J, Feuer W, Roberts J, Gillings D, Sim E. Influence of transcutaneous oxygen monitoring on the incidence of retinopathy of prematurity. *Pediatrics* 1987;79:663-669
- 205 Flynn J, Bancalari E, Sim Snyder E, Goldberg RN, Bawol R, Feuer W, Cassady J, Schiffman J, Feldman HI, Bachinsky B, Buckley E, Roberts J, Gillings D. A cohort study of transcutaneous oxygen tension and the incidence and severity of retinopathy of prematurity. *N Engl J Med* 1992;326:1050-1054
- 206 Cunningham S, Fleck BW, Elton RA, McIntosh N. Transcutaneous oxygen levels in retinopathy of prematurity. *Lancet* 1995;346:1464-1465
- 207 Tuchschnid PE, Boutellier U, Koller EA, Duc GV. Comparison of hypoxanthine, lactate, and ECG signs as indicators of hypoxia. *Pediatr Res* 1981;15:28-31
- 208 Hrbek A, Karlberg P, Kjellmer I, Olsson T, Riha M. Clinical application of evoked EEG responses in newborn infants II: Idiopathic respiratory distress syndrome. *Develop Med Child Neurol* 1978;20:619-626
- 209 Moss AF, Emmannouillides GC, Adams FH. Response of ductus arteriosus and pulmonary and systemic arterial pressures to changes in oxygen environment in the newborn infant. *Pediatrics* 1964;33:937-944
- 210 Bard H, Belanger S, Fouron JC. Comparison of effects of 95 % and 90 % oxygen saturations in respiratory distress syndrome. *Arch Dis Child* 1996;75:F94-F96

- 211 Émond D, Lachance C, Gagnon J, Bard H. Arterial partial pressure of oxygen required to achieve 90 % saturation of hemoglobin in very low birth weight newborns. *Pediatr* 1993;91:602-604
- 212 American Academy of Pediatrics and American College of Obstetricians and Gynecologists. Guidelines for perinatal care. 3rd ed. Washington, DC. 1992:197-203
- 213 Poets CF, Stebbens VA, Alexander JR, Arrowsmith WA, Salfield SAW, Southall DP. Oxygen saturation and breathing patterns in infancy. 1: Full term infants in the second month of life. *Arch Dis Child* 1991;66:569-573
- 214 Thilo EH, Park-Moore B, Berman ER, Carson BS. Oxygen saturation by pulse oximetry in healthy infants at an altitude of 1610 m (5280 ft). *AJDC* 1991;145:1137-1140
- 215 Poets CF, Stebbens VA, Alexander JR, Arrowsmith WA, Salfield SAW, Southall DP. Oxygen saturation and breathing patterns in infancy. 2: Preterm infants at discharge from special care. *Arch Dis Child* 1991;66:574-578
- 216 Poets CF, Stebbens VA, Alexander JR, Arrowsmith WA, Salfield SAW, Southall DP. Arterial oxygen saturation in preterm neonates at discharge from the hospital and six weeks later. *Arch Dis Child* 1992;62:882-888
- 217 Richard D, Poets CF, Neale S, Stebbens VA, Alexander JR, Southall DP. Arterial oxygen saturation in preterm neonates without respiratory failure. *J Pediatr* 1993;123:963-968
- 218 Wasunna A, Whitelaw AG. Pulse oximetry in preterm infants. *Arch Dis Child* 1987;62:882-888
- 219 Stark AR, Richardson D. Blood gas monitoring. In Cloherty JP, Stark AR. *Manual of neonatal care*. Boston: Little, Brown, 1991:209-214
- 220 Niemeyer S, Yang P, Shanmina; Drolkar; Zhuang J, Moore LG. Arterial oxygen saturation in Tibetan and Han infants born Lhasa, Tibet. *N Engl J Med* 1995;333:1248-1252
- 221 Hay WW, Thilo E, Brockway Curlander J. Pulseoximetry in neonatal medicine. *Clin.in Perinatol*. 1991;18:441-72
- 222 Southall DP, Bignall S, Stebbens VA, Alexander JR, Rivers RPA, Lissauer T. Pulse oximeter and transcutaneous arteria; oxygen measurements in neonatal and paediatric intensive care. *Arch Dis Child* 1987;62:882-888
- 223 Walsh MC, Noble LM, Carlo WA, Martin RJ. Relationship of pulse oximetry to arterial oxygen tension in infants. *Crit Care Med* 1987;15:1102-1105
- 224 Bucher HU, Fanconi S, Baekert P, Duc G. Hyperoxemia in newborn infants: Detection by pulseoximetry. *Pediatrics* 1989;84:226-230
- 225 Poets CF, Wilken M, Seidenberg J, Southall DP, Hardt H von der. Reliability of a pulseoximeter in the detection of hyperoxemia. *J Pediatr* 1993;122:87-90
- 226 Paky F, Koeck CM. Pulse oximetry in ventilated preterm newborns: reliability of detection of hyperoxaemia and hypoxaemia, and feasibility of alarm settings. *Acta Paediatr* 1995;84:613-616
- 227 Joint working group of the British Association of Perinatal Medicine and the Research Unit of the Royal College of Physicians. Development of audit measures and guidelines for good practice in the management of neonatal respiratory distress syndrome. *Arch Dis Child* 1992;67: 1221-1227
- 228 Swyer PR, Boston RW, Murdock A, Pare C, Rees EP, Segal S, Sinclair JC. Oxygen therapy in the newborn infant. Statement by the fetus and newborn committee of the canadian pediatric society. *Can Med Assoc J* 1975;113:750-763
- 229 Southall DP, Samuels MP, Poets CF. Neonatal respiratory distress syndrome [letter]. *Arch Dis Child* 1993;69:330
- 230 Cochran DP, Shaw NJ. Neonatal respiratory distress syndrome [letter]. *Arch Dis Child* 1994;70:F230
- 231 Fahey PJ, Harris K, Vanderwarf C. Clinical experience with continuous monitoring of mixed venous oxygen saturation in respiratory failure. *Chest* 1984;85:748-752
- 232 Richard C, Thuilez C, Pezzano M, Bottineau G, Giudicelli JF, Auzepy P. Relationship between mixed venous oxygen saturation and cardiac index in patients with chronic congestive heart failure. *Chest* 1989;95:1289-1294

- 233 Hassan E, Green JA, Nara AR, Jarvis RC, Kasmer RJ, Pospisil R. Continuous monitoring of mixed venous oxygen saturation as an indicator of pharmacological intervention. *Chest* 1989;95:406-409
- 234 Kyff JV, Vaughn S, Yang SC, Raheja R, Puri VK. Continuous monitoring of mixed venous oxygen saturation in patients with acute myocardial infarction. *Chest* 1989;95:607-611
- 235 Shenaq SA, Casar G, Chelly JE, Ott H, Crawford ES. Continuous monitoring of mixed venous oxygen saturation during aortic surgery. *Chest* 1987;92:796-799
- 236 Vaughn S, Puri VK. Cardiac output changes and continuous mixed venous oxygen saturation in the critically ill. *Crit Care Med* 1988;16:495-498
- 237 Schmidt CR, Frank LR, Forsythe SB, Estafanous FG. Continuous SvO₂ measurement and oxygen transport patterns in cardiac surgery patients. *Crit Care Med* 1984;12:523-527
- 238 Jain A, Shroff SG, Janicki JS, Reddy HK, Weber KT. Relation between mixed venous oxygen saturation and cardiac index. Nonlinearity and normalization for oxygen uptake and hemoglobin. *Chest* 1991;99:1403-1409
- 239 Silance PG, Simon C, Vincent JL. The relation between cardiac index and oxygen extraction in acutely ill patients. *Chest* 1994;105:1190-1197
- 240 Buheitel G, Scharf J, Hofbeck M, Singer H. Estimation of cardiac index by means of the arterial and the mixed venous oxygen content and pulmonary oxygen uptake determination in the early post-operative period following surgery of congenital heart disease. *Intensive Care Med* 1994;20:500-503
- 241 Mahutte CK, Jaffe MB, Sasse SA, Chen PA, Berry RB, Sassoon CSH. Relationship of thermomodulation cardiac output to metabolic measurements and mixed venous oxygen saturation. *Chest* 1993;104:1236-1242
- 242 Komatsu T, Shibutani K, Okamoto K, Kumar V, Kubal K, Sanchala V, Lees DE. Critical level of oxygen delivery after cardiopulmonary bypass. *Crit Car Med* 1987;15:194-197
- 243 Danek SJ, Lynch JP, Weg JG, Danzker DR. The dependence of oxygen uptake on oxygen delivery in the adult respiratory distress syndrome. *Am Rev Resp Dis* 1980;122:387-395
- 244 Clarke C, Edwards JD, Nightingale P, Mortimer AJ, Morris J. Persistence of supply dependency of oxygen uptake at high levels of delivery in adult respiratory distress syndrome. *Crit Care Med* 1991;19:497-502.
- 245 Dorinsky PM, Costello JL, Gadek JE. Relationships of oxygen uptake and oxygen delivery in respiratory failure not due to the adult respiratory distress syndrome. *Chest* 1988;93:1013-1019
- 246 Ronco JJ, Montaner JSG, Fenwick JC, Ruedy J, Russel JA. Pathologic dependence of oxygen consumption on oxygen delivery in acute respiratory failure secondary to AIDS-related pneumocystis carinii pneumonia. *Chest* 1990;98:1463-1466
- 247 Astiz ME, Rackow EC, Falk JL, Kaufman BS, Weil MH. Oxygen delivery and consumption in patients with hyperdynamic septic shock. *Crit Care Med* 1987;15:26-28
- 248 Guttierrez G, Pohil RJ. Oxygen consumption is linearly related to O₂ supply in critically ill patients. *J Crit Care* 1986;1:45-53
- 249 Silverman HJ, Abrams J, Rubin LJ. Effects of interleukin-2 on oxygen delivery and consumption in patients with advanced malignancy. *Chest* 1988;94:816-821
- 250 Demling RH, Lalonde C, Fogt F, Zhu D, Liu Y. Effect of increasing oxygen delivery postburn on oxygen consumption and oxidant-induced lipid peroxidation in the adult sheep. *Crit Care Med* 1989;17:1025-1030
- 251 Steib A, Freys G, Gohard R, Curzola U, Ravanello J, Lutun P, Boudjema K, Ottenni JC. Tissue oxygenation during liver transplantation. *Crit Care Med* 1992;20:977-983
- 252 Mohsenifar Z, Amin D, Jasper AC. Dependence of oxygen consumption on oxygen delivery in patients with chronic congestive heart failure. *Chest* 1987;92:447-452
- 253 Cain SM, Curtis SE. Experimental models of pathologic oxygen supply dependency. *Crit Care Med* 1991;19:603-643
- 254 Nelson DP, Samsel RW, Wood LDH, Schumacker PT. Pathological supply dependence of systemic and intestinal oxygen uptake during endotoxemia. *J Appl Physiol* 1988;64:2410-2419

- 255 Bredle DL, Cain SM. Systemic and muscle oxygen uptake/delivery after dopexamine infusion in endotoxic dogs. *Crit Care Med* 1991;19:198-204
- 256 Schumacker PT, Cain SM. The concept of a critical oxygen delivery. *Intensive Care Med* 1987;13:223-229
- 257 Ellsworth ML, Goldfarb RD, Alexander RS. Microembolization induced oxygen utilization impairment in the canine gracilis muscle. *Adv Shock Res* 1981;193:296
- 258 Dantzer DR, Foresman B, Gutierrez G. Oxygen supply and utilization relationships. A reevaluation. *Am Rev Respir Dis* 1991;143:675-679
- 259 Russell JA, Phang PT. The oxygen delivery / consumption controversy. Approaches to management of the critically ill. *Am J Respir Crit Care Med* 1994;149:533-537
- 260 Vermeij CG, Feenstra BWA, Bruining HA. Oxygen delivery and oxygen uptake in postoperative and septic patients. *Chest* 1990;98:415-420
- 261 Mira JP, Fabre JE, Baigorri F, Coste J, Annat G, Artigas A, Nitenberg G, Dhainaut JFA. Lack of oxygen supply dependency in patients with severe sepsis. *Chest* 1994;106:1524-1531
- 262 Ronco JJ, Fenwick JC, Tweedale MG, Wiggs BR, Phang T, Cooper DJ, Cunningham KF, Russell JA, Walley KR. Identification of the critical oxygen delivery for anaerobic metabolism in critically ill septic and non-septic humans. *JAMA* 1993;270:1724-1730
- 263 Hanique G, Dugernier T, Laterre PF, Dougnac A, Roeseler J, Reynaert MS. Significance of pathologic oxygen supply dependency in critically ill patients: Comparison between measured and calculated methods. *Intensive Care Med* 1994;20:12-18
- 264 Hanique G, Dugernier T, Laterre PF, Roeseler J, Dougnac A, Reynaert MS. Evaluation of oxygen uptake and delivery in critically ill patients: a statistical reappraisal. *Intensive Care Med* 1994;20:19-26
- 265 Backer De D, Moraine JJ, Berre J, Kahn RJ, Vincent JL. Effects of dobutamine on oxygen consumption in septic patients. direct versus indirect determinations. *Am J Respir Crit Care Med* 1994;150:95-100
- 266 Yu M, Burchell S, Takiguchi SA, McNamara JJ. The relationship of oxygen consumption measured by indirect calorimetry to oxygen delivery in critically ill patients. *J Trauma* 1996;41:41-47
- 267 Lucking SE, Williams TM, Chaten FC, Metz RI, Mickell JJ. Dependence of oxygen consumption in pediatric septic shock and low oxygen extraction. *Crit Care Med* 1990;18:1316-1319
- 268 Mink RB, Pollack MM. Effect of blood transfusion on oxygen consumption in pediatric septic shock. *Crit Care Med* 1990;18:1087-1091
- 269 Dietrich KA, Conrad SA, Herbert CA, Levy GF, Romero MD. Cardiovascular and metabolic response to red blood cell transfusion in critically ill volume resuscitated non-surgical patients. *Crit Care Med* 1990;18:940-944
- 270 Seear M, Wensley D, MacNab A. Oxygen consumption-oxygen delivery relationship in children. *J Pediatr* 1993;123:208-214
- 271 Pulmonary artery catheter consensus conference participants. Pulmonary artery catheter consensus conference: consensus statement. *Crit Care Med* 1997;6:910-925
- 272 Bland RD, Shoemaker WC. Probability of survival as a prognostic and severity of illness score in critically ill patients. *Crit Care Med* 1985;13:91-96
- 273 Bland RD, Shoemaker WC, Abraham E, Cobo JC. Hemodynamic and oxygen transport patterns in surviving and non-surviving patients. *Crit Care Med* 1985;13:85-92
- 274 Shoemaker WC. Hemodynamic and oxygen transport patterns in shock: Pathophysiology, monitoring, outcome prediction and therapy. In: Edwards JD, Shoemaker WC, Vincent JL. (Eds) *Oxygen Transport: principles and practice*. London, Saunders, 1993, pp 70-98
- 275 Prielipp RC, MacGregor DA, Butterworth JF, Meredith JW, Levy JH, Wood KE, Coursin DB. Pharmacodynamics and pharmacokinetics of Milrinone administration to increase oxygen delivery in critically ill patients. *Chest* 1996;109:1291-1301
- 276 Shoemaker WC, Appel PL, Kram HB, Waxman K, Lee TS. Prospective trial of supranormal values of survivors as therapeutic goals in high-risk surgical patients. *Chest* 1988;94:1176-1186

- 277 Boyd O, Grounds M, Bennet D. A randomized clinical trial of the effect of deliberate perioperative increase of oxygen delivery on mortality in high-risk surgical patients. *JAMA* 1993;270:2699-2707
- 278 Bishop MH, Schoemaker WC, Appel PL, Meade P, Ordog GJ, Wasserberger J, Wo CJ, Rile DA, Kram HB, Umali R, Kennedy F, Shuleshko J, Stephen CM, Shori SK, Thadepalli HD. Prospective, randomized trial of survivor values of cardiac index, oxygen delivery, and oxygen consumption as resuscitation endpoints in severe trauma. *J Trauma* 1995;38:780-787
- 279 Schoemaker WC, Belzberg H. Maximizing oxygen delivery in high-risk surgical patients. *Crit Care Med* 1997;25:714-715
- 280 Tuschmidt J, Fried J, Astiz M, Rackow E. Elevation of cardiac output and oxygen delivery improves outcome in septic shock. *Chest* 1992;102:216-220
- 281 Yu M, Levy MM, Smith P, Tagikuchi SA, Miyasaki A, Hasaniya N, Myers SA. Effect of maximizing oxygen delivery on morbidity and mortality rates in critically ill patients: A prospective randomized trial. *Crit Care Med* 1993;21:830-838
- 282 Hayes MA, Timmins AC, Yau EHS, Palazzo M, Hinds CJ, Watson D. Elevation of systemic oxygen delivery in the treatment of critically ill patients. *N Engl J Med* 1994;330:1717-1722
- 283 Hayes MA, Timmins AC, Yau EHS, Palazzo M, Watson D, Hinds CJ. Oxygen transport patterns in patients with sepsis syndrome or septic shock: influence of treatment and relationship to outcome. *Crit Care Med* 1997;25:926-936
- 284 Gattinoni L, Brazzi L, Pelosi P, Latini R, Tognoni G, Pesenti A, Fumagalli R, for the SvO₂ collaborative group. A trial of goal-oriented hemodynamic therapy in critically ill patients. *N Engl J Med* 1995;335:1025-1032
- 285 Kremzar B, Spec-Marn A, Kompan L, Cerovic O. Normal values of SvO₂ as therapeutic goal in patients with multiple injuries. *Intensive Care Med* 1997;23:65-70
- 286 Yu M, Takanishi D, Myers SA, Tagikuchi SA, Severino R, Hasaniya N, Levy MM, McNamara J. Frequency of mortality and myocardial infarction during maximizing oxygen delivery: A prospective randomized trial. *Crit Care Med* 1995;23:1025-1032
- 287 Third european consensus conference in intensive care medicine. Tissue hypoxia: How to detect, how to correct, how to prevent. *Am J Respir Crit Care Med* 1996;154:1573-1578
- 288 Heyland DK, Cook DJ, King D. Maximizing oxygen delivery in critically ill patients: A methodologic appraisal of excellence. *Crit Care Med* 1996;24:317-324

Chapter 3

Mixed Venous Oxygen Saturation (SvO₂) and biochemical parameters of hypoxia during progressive hypoxemia in 10-14 day old piglets.

Mark A.H.B.M. van der Hoeven, Wiel J. Maertzdorf, Carlos E. Blanco.

Pediatric Research 1997;42:878-884

Abstract

In this study we wanted to assess the relationship between mixed venous oxygen saturation (SvO_2) and tissue oxygenation. For that, we compared the values of SvO_2 with oxygen delivery (DO_2), oxygen consumption (VO_2) and markers of tissue hypoxia as lactate and pyruvate during progressive hypoxemia. Eight 10 - 14 days old piglets were anesthetized, tracheotomized, intubated and ventilated. A fiberoptic catheter was placed in the carotid artery to monitor arterial oxygen saturation (SaO_2). A thoracotomy was performed and a fiberoptic catheter was placed in the pulmonary artery to monitor SvO_2 . A transit time ultrasound flow probe was positioned around the ascending aorta to measure aorta flow. Progressive graded hypoxemia was induced by decreasing fractional inspiratory oxygen concentration (FiO_2) from 1.0 to 0.30, 0.21, 0.15 and 0.10. After each FiO_2 interval blood samples were taken for blood gases, lactate and pyruvate. DO_2 and VO_2 were calculated. SvO_2 decreased similarly to SaO_2 . A value of SvO_2 of more than 40 % excluded oxygen restricted metabolism. When DO_2 decreased below a critical range (8,4 - 12,8 ml/kg.min), SvO_2 decreased below 15 %, and lactate and lactate/pyruvate ratio increased. We conclude, a) that baseline SvO_2 values excluded oxygen restricted metabolism, b) that SvO_2 values between 15 % and 40 % were not a marker for oxygen restricted metabolism and c) that SvO_2 values below 15 % were associated with oxygen restricted metabolism. Reduced SvO_2 values must be interpreted as a change of the factors, that determine the balance between DO_2 and VO_2 and as a warning that, with further reduction of SvO_2 , oxygen restricted metabolism can develop.

Introduction

Mixed venous oxygen saturation (SvO₂) reflects the residual oxygen after tissue oxygen extraction and represents the combined sufficiency of arterial oxygen content (caO₂), cardiac output (Qt) and tissue oxygen consumption (VO₂)^[1]. In adult intensive care SvO₂ has been used to monitor cardiac output. In neonatal medicine SvO₂ monitoring is used during extracorporeal support to monitor the sufficiency of oxygen delivery (DO₂)^[2]. Continuous SvO₂ measurement was proposed in neonatal intensive care to monitor the oxygen sufficiency at the tissue level, but it is only used occasionally^[1,3].

We have determined the feasibility and the accuracy of a fiberoptic catheter system in the newborn, providing a continuous central SvO₂ measurement in the right atrium^[4]. We have also described SvO₂ values in 1-7 days old preterm infants breathing room air^[5]. However, critical values of SvO₂ below which oxygen sufficiency at the tissue level is threatened, have not been properly evaluated in the newborn period.

It is known, that, when DO₂ at the tissue level is insufficient, oxygen restricted metabolism will develop and the concentration of biochemical markers of tissue hypoxia such as lactate and pyruvate will appear in increased concentrations in the blood^[6,7].

In this study we attempted to assess the relation between SvO₂ and tissue oxygen sufficiency. Hence, we examined the values of SvO₂ in comparison with DO₂, VO₂ and biochemical markers of tissue hypoxia in order to determine a normal and safe range of SvO₂ and to determine a critical value of SvO₂ below which oxygen restricted metabolism will develop.

Materials and methods:

This study was approved by the animal care committee of the University of Maastricht.

Animal preparation

Eight piglets between 10 and 14 days of age were anesthetized using halothane for induction, tracheotomized, intubated and ventilated with room air. Anesthesia was maintained by continuous intravenous infusion of pentobarbital sodium and pancuronium in glucose 10 % as required. A 4 Fr. fiberoptic catheter (Abbott Lab, Chicago Ill. USA) was placed in the right carotid artery for continuous monitoring of arterial oxygen saturation (SaO₂), for blood sampling and for monitoring arterial blood pressure. A 5 Fr. catheter was placed in the jugular or femoral vein for fluid administration. A thoracotomy was performed and the ductus arteriosus was ligated. A transit time ultrasound blood volume flow probe (Transonic System Inc, Ithaca, NY, USA) was positioned around the ascending aorta to monitor continuously aorta blood flow. The pulmonary artery was catheterized with a 4 Fr. fiberoptic catheter (Abbott Lab, Chicago Ill. USA) for monitoring SvO₂ continuously and for blood sampling. A rectal probe was placed to monitor core body temperature.

Protocol

The piglets were ventilated to maintain P_aCO₂ at 5.0 ± 1.0 kPa, with a pressure limited, time cycled ventilator (Bourns BP-200, Medical Systems, Inc. Ca. USA). Inspiratory gas was warmed and humidified. We induced progressive graded hypoxemia by changing the proportion of oxygen/nitrogen in the inspiratory gas resulting in a fractional inspiratory oxygen concentration (FiO₂) of 1.0, 0.30, 0.21, 0.15, and 0.10. Every fixed FiO₂ was kept for an interval of 15 minutes. When blood pressure dropped or bradycardia occurred at the lowest FiO₂, the interval was shortened or discontinued. At the end of each interval blood samples were taken simultaneously from the carotid artery for blood gases,

hemoglobin, lactate, pyruvate and from the pulmonary artery for blood gases and hemoglobin. The study included two series of the different FiO_2 intervals with a one hour period in between for recovery. We have repeated the series of graded hypoxia, because all animals except one regained a stable condition after the first series of the FiO_2 intervals and because more data could be obtained per animal. A blood transfusion was given for accidental blood loss during the operative procedure or when hemoglobin level decreased due to blood sampling during the experiment to maintain a baseline hemoglobin level of 5.0 ± 1.0 mmol/L.

Measurements

The fiberoptic catheter system (Oximetrix System, Abbott Laboratories, North Chicago Ill. U.S.A.) includes a fiberoptic catheter and a microprocessor based oximeter. The light back scattered from the oxygenated and deoxygenated blood is transmitted by the receiving fiberoptic bundle to a photodetector. The oximeter computes percentage of oxygen saturation values based on the electrical signals from the photodetector.

SaO_2 , SvO_2 , arterial blood pressure, blood flow in the aorta, heart rate and core body temperature were sampled every 10 seconds, displayed and stored by a software system (Hemodynamic Data Acquiring System (HDAS) developed at the Service Center, University of Maastricht). Blood samples for blood gases were analyzed on an ABL 3 acid base analyzer (Radiometer, Copenhagen, Denmark) corrected to the animal core body temperature. SaO_2 , arterial hemoglobin (Hb) (mmol/l) and SvO_2 were analyzed on the OSM2 (Radiometer, Copenhagen, Denmark). Lactate and pyruvate were analyzed in arterial blood samples taken simultaneously with the samples for acid base analysis. 1 ml Arterial blood was pipetted into a tube containing 1 ml 1.2 M perchloric acid to obtain proteinfree solution. The enzyme lactate dehydrogenase (LDH) added to this solution, catalyses the conversion of pyruvic acid in the presence of NADH to lactic acid and catalyses the conversion from lactic acid to pyruvic acid in the presence of NAD. The reduced or increased absorbance at 340 nm becomes a measure of the amount of pyruvic acid and of lactic acid, respectively ^[8].

Calculations

- Arterial oxygen content (caO_2) and venous oxygen content (cvO_2) were calculated using the equations:

caO_2 (ml/dl) = $(1.36 \times \text{SaO}_2 \times \text{Hb (mmol/l)} \times 1.611) + (0.0031 \times \text{paO}_2 \text{ (kPa)} \times 7.5)$ and
 cvO_2 (ml/dl) = $(1.36 \times \text{SvO}_2 \times \text{Hb (mmol/l)} \times 1.611) + (0.0031 \times \text{pvO}_2 \text{ (kPa)} \times 7.5)$, respectively.

- Cardiac output (Qt) was determined by the mean value of the bloodflow in the aorta during the last five minutes of each FiO_2 interval measured by the flowsensor.

- DO_2 was calculated from its definition:

DO_2 (ml/kg.min) = caO_2 (ml/dl) $\times$ Qt (ml/kg.min) / 100 .

- VO_2 was calculated using the Fick equation:

VO_2 (ml/kg.min) = $\text{Qt}/100 \times \text{caO}_2 - \text{cvO}_2$.

- Fractional oxygen extraction (F.O.E.) was calculated as the VO_2 divided by DO_2 .

- P_{50} was calculated from venous blood gas samples using SvO_2 and pvO_2 as has been described in the literature ^[9]. O_2 debt was calculated by multiplying the difference between the mean baseline VO_2 at $\text{FiO}_2 = 1.0$ and the estimated VO_2 in the FiO_2 interval by the duration of that same FiO_2 interval.

- Cumulative O_2 debt was calculated by summing up the O_2 debt of each interval.

- The change of pHa, P_aCO_2 , paO_2 , pvO_2 , p_{50} , Hemoglobin (Hb), temperature, heart rate, Qt, lactate, pyruvate, lactate/pyruvate ratio, VO_2 c(a-v) O_2 , and F.O.E. was calculated in comparison with the mean baseline value of these items at $\text{FiO}_2 = 1.0$ in each series of FiO_2 intervals in each individual animal. Absolute values were used for SaO_2 , SvO_2

and DO₂.

Data analysis

Mean, standard deviation, median value and percentile were calculated by descriptive statistical methods. Data are presented as mean $\pm$ SD. Data at the different FiO₂ intervals are presented as absolute values and as a mean change in comparison with the mean baseline values at FiO₂ = 1.0 in each individual animal. To test whether a difference existed between the two series of induced hypoxemia, a Mann-Whitney U test was performed for each FiO₂ interval. Baseline data were compared with the data at the different low FiO₂ intervals using oneway analysis of variance and contrasted according to Tukey B method.

To determine the critical value of DO₂, that is the DO₂ below which the change in lactate concentration increased rapidly, we adapted a technique from Mellits^[10,11] to find the section point of two lines of regression. This included constructing the best pair of regression lines that included all points only once. The best pair of regression lines was defined as to minimize the total of the residual sum of squares of each line in the pair. Their intersection point was determined by solving the equations of the pair of regression lines. The x-coordinate of the intersection point is the critical DO₂^[11]. This technique was also applied to change in lactate/pyruvate ratio and change in VO₂ versus DO₂.

To test whether a value of SvO₂ was predictive for a change in lactate, a change in lactate/pyruvate ratio and a change in VO₂, a 2 x 2 contingency table was derived for each item and sensitivity, specificity and positive predictive accuracy were calculated. Tests were considered to be significant when $p \leq 0.05$.

Results:

Eight piglets were studied with a mean weight of 4.08 ± 0.8 kg and at a mean age of 11 ± 2 days. In seven piglets a complete protocol of 2 series of experiments could be performed. No significant differences were found between the results of corresponding FiO₂ intervals in the two consecutive series of induced hypoxemia. Measured and calculated data during the five FiO₂ intervals are shown in table 1. Baseline values regarding cardiac output (Qt), calculated VO₂, hemoglobin, caO₂, and p₅₀ were consistent with values in 10-14 day old piglets, described by others^[12,13].

Hemodynamic responses during the different intervals consisted of an increased heart rate in the last two FiO₂ intervals and a decreased bloodpressure at the last interval, whereas cardiac output remained unaltered (table 1).

Progressive graded hypoxemia induced a decrease in SaO₂ and paO₂ from 98.0 % to 24.7 % and from 51.4 kPa to 2.7 kPa, respectively. SvO₂ decreased similarly to the decreasing SaO₂ (figure 1). Therefore in all but the lowest FiO₂ interval the difference between arterial and venous oxygen content ($c(a-v)O_2$) remained unaltered and a constant VO₂ was preserved (table 1). In the lowest FiO₂ interval DO₂ decreased further, and VO₂ decreased significantly (table 1, figure 1).

Lactate and pyruvate during baseline showed a broad dispersion of values reflecting different postoperative status: (accidental blood loss, intubation difficulties etc.). The change in lactate and lactate/pyruvate ratio was significantly in the last FiO₂ interval. Pyruvate changed not significant from baseline value (table 1).

The relationships between DO₂ and change lactate, change lactate/pyruvate ratio and change VO₂ were analyzed in figure 2 according to the technique from Mellits^[10]. The critical DO₂ value was 12.86 ml/kg.min for lactate, 8.43 ml/kg.min for lactate/pyruvate ratio and 11.82 ml/kg.min for VO₂.

table 1. Measured and calculated data during baseline and change from baseline during the different FiO₂ intervals.

	FiO ₂ = 1,0 baseline	FiO ₂ = 0,3 change	FiO ₂ = 0,21 change	FiO ₂ = 0,15 change	FiO ₂ = 0,10 change
pHa	7,43 ± 0,12	0,03 ± 0,06	0,05 ± 0,07	0,01 ± 0,09	-0,03 ± 0,11
paCO ₂ (kPa)	4,5 ± 1,71	-0,45 ± 0,87	-0,49 ± 0,92	-0,49 ± 0,58	-0,56 ± 1,24
paO ₂ (kPa)	51,5 ± 17,34	-35,3 ± 12,3 *	-43,8 ± 10,9 *	-48,7 ± 10,1 *	-47,6 ± 10,6 *
SaO ₂ (%)	98,0 ± 2,1	-3,6 ± 5,3	-15,6 ± 12,8 *	-42,2 ± 12,7 *	-73,3 ± 10,5 *
pvO ₂ (kPa)	5,58 ± 0,97	-1,13 ± 0,65 *	-2,16 ± 0,85 *	-3,23 ± 0,83 *	-4,59 ± 0,93 *
SvO ₂ (%)	59,5 ± 9,6	-10,5 ± 7,9 *	-22,9 ± 13,5 *	-40,9 ± 12,6 *	-59,4 ± 8,2 *
p ₅₀ (kPa)	4,68 ± 0,67	-0,24 ± 0,65	-0,49 ± 0,83	-0,37 ± 1,05	0,29 ± 2,68
Hb. (mmol/L)	5,6 ± 1,1	0,19 ± 0,38	0,15 ± 0,45	0,54 ± 0,49 *	1,10 ± 0,73 *
Qt (ml/kg.min)	165 ± 49	-4 ± 17	-2 ± 21	-0,7 ± 32	-18 ± 36
Mean P _{ao} (mm Hg)	94,2 ± 14,4	-0,7 ± 8,8	-4,4 ± 9,5	-6,5 ± 10,3	-21,4 ± 15,2 *
Lactate (mmol/l)	2,75 ± 1,97	0,27 ± 0,48	0,58 ± 1,11	1,17 ± 1,78	4,18 ± 2,61 *
Pyruvate (μmol/l)	189,6 ± 99,2	19,0 ± 65,3	45,8 ± 95,3	29,7 ± 76,6	66,6 ± 104,5
lact/pyr. ratio	13,5 ± 3,4	1,1 ± 2,4	1,2 ± 3,0	4,9 ± 6,3	19,2 ± 11,9 *
DO ₂ (ml/kg.min)	22,3 ± 6,4	-2,2 ± 2,1	-5,8 ± 3,1 *	-9,8 ± 3,5 *	-18,1 ± 5,2 *
VO ₂ (ml/kg.min)	9,3 ± 2,0	0,3 ± 1,0	0,08 ± 1,9	-0,91 ± 1,4	-3,9 ± 2,6 *
c(a-v)O ₂ (ml/dl)	6,14 ± 2,02	0,29 ± 0,85	-0,17 ± 1,66	-0,70 ± 1,46	-2,34 ± 1,67 *
F.O.E.	0,44 ± 0,09	0,05 ± 0,07	0,10 ± 0,14 *	0,26 ± 0,13 *	0,47 ± 0,11 *

* = significant different from baseline. Note that biochemical markers, VO₂ and c(a-v)O₂(Hb) only during the last FiO₂ interval increase significantly from baseline.

figure 1. SaO_2 , SvO_2 , DO_2 and change VO_2 in relation to the different FiO_2 intervals. Note that, SvO_2 decreased similarly to SaO_2 ; Change VO_2 was constant in all but the last interval. In the lowest FiO_2 interval DO_2 further decreased, and change VO_2 was significantly reduced (*). Data are presented as mean $\pm$ SD.

figure 2. Change lactate, change lactate/pyruvate ratio and change VO_2 in relation with DO_2 . A critical DO_2 (vertical broken line) could be observed at 12.86, 8.46 and 11.81 $\text{ml.kg}^{-1}.\text{min}^{-1}$, respectively.

figure 3. SvO₂ in relation with DO₂. Regression line with confidence intervals is: $2.51 \times \text{DO}_2 - 4.01$ ($r = 0.87$; $p < 0.0001$). Vertical lines represent critical DO₂ calculated for change lactate/pyruvate ratio and change lactate, respectively. Below this critical range of DO₂ (8.43 - 12.86 ml/kg.min) SvO₂ values of less than 15 % were observed.

Above the critical DO₂, the change of lactate, lactate/pyruvate ratio and VO₂ were only slight dependent of DO₂. Below the critical DO₂ a significant regression coefficient (slope) was obtained.

Linear regression analysis showed a good correlation between SvO₂ in relation with DO₂ ($r = 0.87$; $p < 0.0001$). When DO₂ decreased below the critical range (8.43 and 12.86 ml/kg.min), SvO₂ values less than 15 % were observed. (figure 3).

In figure 4 the relationship between SvO₂ and the change in lactate, the change in lactate/pyruvate ratio and the change in VO₂ was examined. We considered a change in lactate of more than 1,03 mmol/l, a change in lactate/pyruvate ratio of more than 2,45 and a change in VO₂ of less than -2,25 ml/kg.min as significant, because these values represented the mean change $\pm$ 2SD at FiO₂ = 1.00. SvO₂ values of more than 40 % were considered as baseline, as it represented the mean value minus 2 SD in the highest FiO₂ interval. SvO₂ values of less than 15 % were considered as critical value, as it represented the values encountered below the critical range of DO₂. Lactate, lactate/pyruvate ratio and VO₂ remained mainly unaltered until SvO₂ was less than 15 %.

To test whether values of SvO₂ above or below 40 % were predictive of a significant change in lactate, in lactate/pyruvate ratio and in VO₂, a 2 X 2 contingency table was derived for each item; a change in lactate of ≥ 1.03 mmol/l and of < 1.03 mmol/l, and a change in lactate/pyruvate ratio of ≥ 2.45 and of < 2.45 and a change in VO₂ of > -2.25 ml/kg.min and ≤ -2.25 ml/kg.min constituted the columns and SvO₂ values above 40

figure 4. Change lactate, change lactate/pyruvate ratio, and change VO_2 in relation with SvO_2 . Vertical broken lines indicate a SvO_2 value of 15 % (mean - 2SD at $FiO_2 = 0.10$) and a SvO_2 value of 40 % (mean - 2SD at $FiO_2 = 1.00$). Horizontal broken line indicates change lactate of 1.03 mmol/l (mean change + 2SD at $FiO_2 = 1.0$), change lactate/pyruvate ratio of 2.45 (mean change + 2SD at $FiO_2 = 1.0$) and change VO_2 of -2.25 ml/kg.min (mean change - 2SD at $FiO_2 = 1.0$). Note, that lactate, lactate/pyruvate ratio and VO_2 significantly change, when SvO_2 values below 15 % are encountered.

% and below 40 % were used for the rows. A value of SvO₂ above 40 % was 68 % sensitive and 92 % specific for a change within baseline $\pm$ 2SD in lactate concentration and 79 % sensitive and 96 % specific for a change in lactate/pyruvate ratio and 63 % sensitive and 94 % specific for a change VO₂ within baseline $\pm$ 2SD. The positive predictive accuracy of SvO₂ above 40 % was 94 %, 97 % and 98 % for a change in lactate, lactate/pyruvate ratio and VO₂ within baseline $\pm$ 2SD. However, the positive predictive accuracy of a SvO₂ below 40 % for a significant change above baseline $\pm$ 2SD was only 57 %, 76 % and 37 %.

In the same way we tested whether SvO₂ values below or above 15 % were predictive of a significant change in lactate, lactate/pyruvate ratio and VO₂. A value of SvO₂ below 15 % was 84 % sensitive and 49 % specific for a significant increase in lactate, 77 % sensitive and 100 % specific for a significant increase in lactate/pyruvate ratio and 84 % sensitive and 96 % specific for a change in VO₂. The positive predictive accuracy of SvO₂ below 15 % was 87 %, 100 % and 84 % for a significant change in lactate, lactate/pyruvate ratio and VO₂. The positive predictive accuracy of a SvO₂ above 15 % for a change within baseline $\pm$ 2SD was 92 %, 86 % and 96 %. The positive predictive accuracy of a SvO₂ between 15 % and 40 % for a significant change from baseline $\pm$ 2SD was 13 %, 60 % and 12 %.

In other words SvO₂ values above 40 % were very frequently associated with baseline lactate, lactate/pyruvate ratio and baseline VO₂. SvO₂ values below 15 % were very frequently associated with increased lactate, lactate/pyruvate ratio and decreased VO₂. SvO₂ values between 15 % and 40 % were very occasionally associated with an increased lactate, lactate/pyruvate ratio and decreased VO₂.

Discussion

We assessed the relationship between SvO₂ and tissue oxygenation in piglets by inducing graded progressive hypoxemia. The main results of this study were: a) SvO₂ values of more than 40 % excluded oxygen restricted metabolism, b) SvO₂ values of less than 15 % were associated with oxygen restricted metabolism and c) SvO₂ value between 15 % and 40 % were very frequently associated with normal tissue oxygen metabolism. The aim of this study was neither to determine the value of SvO₂ measurement in neonatal intensive care medicine, nor to describe the limits of the range of SvO₂ values for direct use in neonatal intensive care. The critical values of 15 % and 40 %, found in this study can not be extrapolated to patient care in a neonatal intensive care unit.

This hypoxia model consists of progressive graded hypoxemia by reducing FiO₂ in order to reduce DO₂. In our study the circulatory adjustments as increased cardiac output to compensate for a decreased caO₂ were variable, as it is known from the literature [14]. Microcirculatory adjustments were not measured in this study, but may affect the actual oxygen flow from the capillary bed to the cells and may consequently affect the SvO₂ value. As other types of hypoxia (anemic hypoxia, circulatory failure hypoxia), could have other patterns of circulatory adjustments and compensations, one must be cautious to extrapolate the findings from this hypoxia model to other types of hypoxia.

The series of graded hypoxemia were started with hyperoxia in the FiO₂ = 1.0 interval with a wide variation in baseline conditions, probably due to animal preparation. This is also illustrated by the fact, that in the third FiO₂ = 0.21 interval a mean SaO₂ value of 82 % was present. However, these variations in baseline conditions did not affect the changes induced by the different FiO₂ intervals: SaO₂, the main factor of caO₂, decreased significantly in the FiO₂ intervals, resulting in a significant reduction of DO₂. SvO₂ decreased similarly with SaO₂, which maintained a stable difference between arterial and

venous oxygen content ($c(a-v)O_2$) and preserved a constant VO_2 in all but the last FiO_2 interval (Figure 1). When FiO_2 further decreased, $c(a-v)O_2$ decreased and VO_2 was reduced with the development of oxygen restricted metabolism and lactic acidosis. The initial reduction of SvO_2 was not accompanied by a reduction of VO_2 or an increased concentration of lactate. Only with a severe reduction of SvO_2 below 15 %, regardless of the basal condition of the animal, lactate increased and VO_2 decreased.

In our study SvO_2 values above 15 % were found with only slight changes in lactate concentration, in lactate/pyruvate ratio and in VO_2 . In adult intensive care medicine tolerance of unusually low SvO_2 values have been described^[15]. Therefore, the reduction of SvO_2 per se is not an indication of oxygen restricted metabolism and lactic acidosis. However, in some instances in our study SvO_2 values above 15 % were associated with oxygen restricted metabolism and lactic acidosis. Since SvO_2 reflects only the global tissue oxygen extraction, moderately reduced SvO_2 values with a normal $c(a-v)O_2$ cannot exclude a reduced VO_2 in individual organs, primarily in organs with a high oxygen extraction, like the myocardium and the brain^[16,17]. In accordance with this we found that the critical DO_2 calculated for change in lactate concentration was higher than the critical DO_2 calculated for change in VO_2 . This supports the hypothesis, that oxygen restricted metabolism in individual organs could initiate before a low SvO_2 value is observed and global VO_2 is reduced.

In adult intensive care medicine the relation between SvO_2 and lactic acidosis was studied in relation to prognosis and mortality. Astiz et al. failed to identify a threshold value for SvO_2 below which lactate increased. Lactic acidosis developed over a wide range of both DO_2 and SvO_2 ^[18]. In a retrospective analysis of critical ill patients Rashkin et al. showed that mixed venous oxygen tension was not a reliable indicator of blood lactate^[19]. Thus, a low SvO_2 value can be associated with both a normal tissue oxygen metabolism and an oxygen restricted tissue metabolism. In our study only SvO_2 values below 15 % were very frequently associated with oxygen restricted metabolism. However, these findings do not imply, that SvO_2 is not useful in the estimation of tissue oxygenation. We found that SvO_2 values of more than 40 % (mean - 2SD at $FiO_2 = 1.0$) were not associated with oxygen restricted metabolism. Therefore, this range of SvO_2 values can be interpreted as an indication of normal tissue oxygenation i.e. a normal balance between tissue DO_2 and VO_2 . However, even normal and high SvO_2 values have been associated with lactic acidosis in diseases with disturbances of the microcirculation with precapillary shunting and inability of the tissues to extract enough oxygen, like sepsis and ARDS^[15,20]. This was not the case in our study.

Not only the reduction of DO_2 and VO_2 , but also the duration and cumulation of the effects of the different FiO_2 intervals must be considered. The cumulation of the oxygen debt in the last three FiO_2 intervals can be a reason for the increased lactate concentration in the last FiO_2 interval. Cumulative O_2 debt was calculated and the correlation between cumulative O_2 debt and lactate and lactate / pyruvate ratio analyzed to control for the cumulative effect of different FiO_2 intervals. Cumulative O_2 debt was linearly related to the change in lactate and in lactate/pyruvate ratio, but the correlation coefficient ($r = 0.53$; $r = 0.70$) was not different from the correlation coefficient between the change in VO_2 and the change in lactate and in lactate/pyruvate ratio ($r = 0.53$; $r = 0.53$). The linear correlation analysis between the change in VO_2 and cumulative O_2 debt yielded a good correlation ($r = 0.69$, $p < 0.001$). Cumulative O_2 debt was in this hypoxic piglet model not a better parameter of oxygen restricted metabolism than change in VO_2 . Additionally no differences were found in changes in lactate concentration and in lactate/pyruvate ratio between the two series of FiO_2 intervals.

In this study VO₂ was calculated according to Fick's principle and it was dependent on the precision of caO₂, cvO₂ and of cardiac output measurements. The SvO₂ values are reliable, because after closure of the ductus arteriosus during the procedure, blood-samples and fiberoptic catheter readings reflect truly oxygen saturation values of the mixed venous blood. Whereas oxygen saturation measurements and oxygen content are reliable, the different methods of measurement of cardiac output have many error sources [21]. The value of calculated VO₂ has been questioned [22]. In particular when calculated DO₂ and calculated VO₂ are compared as independent and dependent variables and a pathologic dependence of VO₂ on DO₂ was demonstrated, erroneous mathematical coupling of shared variables was suggested [23]. In our study cardiac output was measured directly by a 7-mm transit-time ultrasonic flow transducer positioned around the aorta. The transducer was precalibrated in vitro. This transit-time ultrasonic technique is described as more precise and as easier to apply than formerly used techniques [24]. The transit-time flow measurements were used as a "gold standard" and confirmed to be accurate compared with a radionuclide-labeled microsphere technique [25]. Therefore, we considered the calculated DO₂ and VO₂, derived from the mentioned data, both reliable and accurate.

The biochemical markers as lactate and pyruvate were used as the indicator for oxygen restricted metabolism. The use of lactate as an index for tissue oxygen sufficiency has several limitations. Lactic acidosis originates only, when lactate production exceeds hepatic, renal and muscle lactate uptake [7]. Thus, lactate value is among others dependent on the capacity of liver and kidney to extract and utilize lactate [6]. Lactate / pyruvate ratio was measured to estimate the ratio between the reduced and oxidized molecules in the cytosol: this intracellular redox state is dependent upon oxidative function in the mitochondrion [26] and consequently upon an adequate O₂ supply to the mitochondrion [27]. In contrast, an increase of lactate and pyruvate without a change in lactate / pyruvate ratio could be a result of inactivation or downregulation of the pyruvate dehydrogenase complex, which controls the entry of two carbon fragments from pyruvate into the citric acid cycle [28]. It was suggested that the measurement of lactate / pyruvate ratio would discriminate between hypoxic and non-hypoxic induced changes of lactate [28].

We conclude that in this hypoxia model, a) SvO₂ values above 40 % excluded oxygen restricted metabolism, b) that SvO₂ values between 15 % and 40 % in general were not associated with oxygen restricted metabolism or lactic acidosis and c) that values of SvO₂ below 15 % were associated with oxygen restricted metabolism. When SvO₂ is used in patient care, reduction of SvO₂ must be interpreted as a reduction of the residual oxygen after tissue oxygen extraction without necessarily affecting VO₂ and without a definite indication of lactic acidosis. Reduced SvO₂ values must be interpreted as a change of the factors, that determine the balance between DO₂ and VO₂ and as a warning that, with further reduction of SvO₂, oxygen restricted metabolism can develop.

References

- 1 O'Connor TA, Hall RT. Mixed venous oxygenation in critically ill neonates. *Crit Care Med* 1994; 22:343-346
- 2 Bartlett RH, Cilley RE. Physiology of extracorporeal life support. In: Arensman RM, Cornish JD (eds). *Extracorporeal life support*. Blackwell, Boston, 1993 pp 89-104
- 3 Dudell G, Cornish JD, Bartlett RH. What constitutes adequate oxygenation? *Pediatrics* 1990;86:39-41

- 4 van der Hoeven MAHBM, Maertzdorf WJ, Blanco CE. Feasibility and accuracy of a fiberoptic catheter for the measurement of venous oxygen saturation in newborn infants. *Acta Paediatr* 1995;84:122-127
- 5 van der Hoeven MAHBM, Maertzdorf WJ, Blanco CE. Continuous central venous oxygen saturation (ScvO₂) measurement using a fibreoptic catheter in newborn infants. *Arch Dis Child* 1996;74:F177-F181
- 6 Kreisberg RA. Lactate homeostasis and lactic acidosis. *Ann Intern Med* 1980;92(1):227-237
- 7 Mizock BA, Falk JL. Lactic acidosis in critical illness. *Crit Care Med* 1992;20:80-93
- 8 Sigma chemical Company. The quantitative determination of pyruvic acid and lactic acid in whole blood at 340 nm. Sigma technical bulletin. No. 726-UV, No. 826-UV, St Louis, 1976 pp 1-21
- 9 Brown EG, Krouskop RW, McDonnell FE, Monge CC, Winslow RE. A technique to continuously measure arteriovenous oxygen content difference and p_{50} in vivo. *J Appl Physiol* 1985;58(4):1383-1389
- 10 Mellits ED. Statistical methods. In: Cheek DB (ed) *Human Growth*. Lea & Febiger, Philadelphia, 1968 pp 19-38
- 11 Fahey JT, Lister G. Postnatal changes in critical cardiac output and oxygen transport in conscious lambs. *Am J Physiol* 1987;253:H100-H106
- 12 Delivoria-Papadopoulos M, Martens RJ, Forster II RE, Oski FA. Postnatal changes in oxygen-hemoglobin affinity and erythrocyte 2,3-diphosphoglycerate in piglets. *Pediatr Res* 1974;8:64-66.
- 13 Mount LE, Ingram DL. The pig as a laboratory animal. Academic Press, London, 1971 pp 65-78
- 14 Moss M, Moreau G, Lister G. Oxygen transport and metabolism in the conscious lamb: the effects of hypoxemia. *Pediatr Res* 1987;22:177-183
- 15 Schlichtig R, Cowden WL, Chaitman BR. Tolerance of unusually low mixed venous oxygen saturation. Adaptations in the chronic low cardiac output syndrome. *Am J Med* 1986;80:813-818
- 16 Reinhart K. Monitoring O₂ transport and tissue oxygenation in critically ill patients. In: Reinhart K, Eyrich K. (Eds) *Clinical aspects of O₂ transport and tissue oxygenation*. Springer Verlag, Berlin, 1989 pp 193-211
- 17 Ceraianu AC, Nelson LD. The significance of mixed venous oxygen saturation and technical aspects of continuous measurement. In: Edwards JD, Shoemaker WC, Vincent JL (Eds) *Oxygen transport; Principles and practice*. Saunders, London, 1993 pp 99-124
- 18 Astiz ME, Rackow EC, Kaufman B, Fay JL, Weil MH. Relationship of oxygen delivery and mixed venous oxygen saturation in patients with sepsis and acute myocardial infarction. *Crit Care Med* 1988;16:655-658
- 19 Rashkin MC, Bosken C, Baughman RF. Oxygen delivery in critically ill patients. Relationship to blood lactate and survival. *Chest* 1985;87:580-584
- 20 Danek SJ, Lynch JP, Weg JG, Dantzker DR. The dependence of oxygen uptake on oxygen delivery in the adult respiratory distress syndrome. *Am Rev Respir Dis* 1980;122:387-395
- 21 Welch E, Duara S, Suguihara C, Bandstra E, Bancalari E. Validation of cardiac output measurement with non-invasive Doppler echocardiography by thermodilution and Fick methods in newborn piglets. *Biol Neonate* 1994;66:137-145
- 22 Bizouarn P, Soulard D, Blanloeil Y, Guillet A, Goarin Y. Oxygen consumption after cardiac surgery - a comparison between calculation by Fick's principle and measurement by indirect calorimetry. *Intens Care Med* 1992;18:206-209
- 23 Russell JA, Phang PT. The oxygen delivery / consumption controversy. Approaches to management of the critically ill. *Am J Respir Crit Care Med* 1994;149:533-537
- 24 Rubertson S, Grenvick Å, Wicklund L. Blood flow and perfusion pressure during open-chest versus closed-chest cardiopulmonary resuscitation in pigs. *Crit Care Med* 1995;23:715-725
- 25 Shiraishi H, Silverman NH, Rudolph AM. Accuracy of right ventricular output estimated by Doppler echocardiography in the sheep fetus. *Am J Obstet Gynecol* 1993;168:947-953
- 26 Mizock B. Lactic Acidosis. *Disease-a-Month* 1989;4:235-300
- 27 Wasserman K, Beaver WL, Whipp BJ. Gas exchange theory and the acidosis (anaerobic) threshold.

- 27 Wasserman K, Beaver WL, Whipp BJ. Gas exchange theory and the acidosis (anaerobic) threshold. *Circulation* 1990;81(suppl II):14-30
- 28 Cain SM, Curtis SE. Systemic and regional oxygen uptake and delivery and lactate flux in endotoxic dogs infused with dopexamine. *Crit Care Med* 1991;19:1552-1560

Acknowledgement

This study was supported by Abbott Inc. The Netherlands.

Relative oxygen saturation (SvO₂) and markers of tissue oxygenation in progressive hypoxic hypoxia and in isovolemic anemic hypoxia in 8-12 days old piglets.

Mark A F.R.M. van der Hoeve, Willem A. Meuwinkel, Carlos E. Blanco

Critical Care Medicine, in press

Chapter 4

Relationship between mixed venous oxygen saturation (SvO₂) and markers of tissue oxygenation in progressive hypoxic hypoxia and in isovolemic anemic hypoxia in 8-12 days old piglets.

Mark A.H.B.M. van der Hoeven, Wiel J. Maertzdorf, Carlos E. Blanco

Critical Care Medicine: in press

Abstract

Objective: To examine the hypothesis that SvO_2 values, which reflect the residual oxygen after tissue oxygen extraction, would be similar during hypoxic and anemic hypoxia.

Design: SvO_2 values, oxygen delivery (DO_2), arterial oxygen content (CaO_2), fractional oxygen extraction (FOE) were compared and critical values were determined based on lactate, lactate/pyruvate ratio and oxygen consumption (VO_2) during hypoxic and anemic hypoxia.

Setting: Laboratory of Physiology at a University Hospital.

Subjects: Two groups of eight piglets, 8 - 12 days old.

Interventions: Piglets were anesthetized, tracheotomized, intubated and ventilated. A thoracotomy was performed and a fiberoptic catheter was placed in the pulmonary artery to monitor SvO_2 . A transit time ultrasound flow probe was positioned around the ascending aorta to measure aorta flow. Progressive hypoxic hypoxia was induced by decreasing fractional inspiratory oxygen concentration (FiO_2) from baseline (0.30 - 0.75) to 0.21, 0.15 and 0.10. Progressive anemic hypoxia was induced by a repeated isovolemic exchange transfusion with 50 ml pasteurized plasma.

Measurements and Main Results: Fifteen or 30 minutes after each intervention, samples were taken from the carotid artery for blood gases, hemoglobin, lactate and pyruvate, and from the pulmonary artery for blood gases and hemoglobin. Hemodynamic, SaO_2 and SvO_2 measurements were made. The calculated DO_2 and VO_2 decreased in both hypoxic and anemic hypoxia. At the lowest DO_2 level of anemic hypoxia, the decrease in SvO_2 was less than that in hypoxic hypoxia (-26 % vs. -55 %). The range of critical values for SvO_2 calculated for each individual piglet below which lactate, lactate/pyruvate ratio and VO_2 rapidly changed from baseline value, was significantly lower in hypoxic hypoxia (11 to 24 %) than in anemic hypoxia (26 to 48 %). F.O.E. increased significantly, but not with a change as high as in hypoxic hypoxia 0.31 (0.20, 0.41) vs. 0.49 (0.41 to 0.54).

Conclusions: In comparison with hypoxic hypoxia, critical values of SvO_2 are higher in anemic hypoxia, indicating that oxygen unloading from blood to tissues is impaired in anemic hypoxia. These characteristics in oxygen transport and capillary hemodynamics should be taken into consideration when SvO_2 is used in clinical critical care.

Introduction

Mixed venous oxygen saturation (SvO₂) reflects the residual oxygen after tissue oxygen extraction and represents the combined sufficiency of arterial oxygen content (CaO₂), cardiac output (Qt) and tissue oxygen consumption (VO₂)^[1]. In the adult intensive care, SvO₂ has been used to monitor cardiac output. In neonatal medicine, SvO₂ monitoring is used during extracorporeal support to monitor the sufficiency of oxygen delivery (DO₂)^[2]. Central SvO₂ measurement in the right atrium has been proposed in neonatal intensive care to monitor the oxygen sufficiency at the tissue level^[3]. We have determined the feasibility and the accuracy of a fiberoptic catheter system in the newborn, providing a continuous central SvO₂ measurement^[4] and described the "normal" range of SvO₂ values in 1 to 7 days old preterm infants breathing room air^[5]. The critical range of SvO₂ below which oxygen restricted metabolism will develop and the concentration of biochemical markers of tissue hypoxia such as lactate and pyruvate will increase^[6,7] was described during hypoxic hypoxia in newborn piglets^[8].

As different types of hypoxia (hypoxic hypoxia, anemic hypoxia or circulatory failure hypoxia) could have other patterns of circulatory adjustments and compensations, the actual oxygen flow from the capillary bed to the cells may be affected. Consequently, different SvO₂ values may be observed during different types of hypoxia. However, SvO₂ and PvO₂ reflect the residual oxygen after tissue oxygen extraction and form the lowest driving pressure for oxygen diffusion to the tissues. Therefore, we hypothesized that critical SvO₂ values, below which oxygen diffusion to the tissues starts to decrease, would not be different during different types of hypoxia. The aim of this study was to compare the critical range of SvO₂ values in progressive hypoxic hypoxia with those in anemic hypoxia. SvO₂ values were examined in comparison with DO₂, VO₂ and biochemical markers of tissue hypoxia during hypoxic hypoxia and anemic hypoxia. Critical values of DO₂, FOE and SvO₂ were determined and compared in both types of hypoxia.

Materials and methods

This study was approved by the animal care committee of the University of Maastricht.

Animal preparation

Two groups of eight piglets between 8 and 12 days of age were anesthetized using halothane for induction, tracheotomized, intubated and ventilated with room air. Anesthesia was maintained by continuous intravenous infusion of pentobarbital sodium and pancuronium in glucose 10 % as required. A 4 Fr. fiberoptic catheter (Abbott Laboratories, North Chicago IL) was placed in the right carotid artery for continuous monitoring of arterial oxygen saturation (SaO₂), blood sampling and monitoring arterial blood pressure. A 5 Fr. catheter was placed in the jugular or femoral vein for fluid administration. A thoracotomy was performed and the ductus arteriosus was ligated. A transit time ultrasound blood volume flow probe (Transonic Systems, Ithaca, NY) was positioned around the ascending aorta to monitor continuously aorta blood flow. The pulmonary artery was catheterized with a 4 Fr. fiberoptic catheter so as to continuously monitor SvO₂ for blood sampling and pulmonary arterial blood pressure. A rectal probe was placed to monitor core body temperature.

Protocol

The piglets were ventilated with a pressure limited, time cycled ventilator (Bourns BP-200, Medical Systems, Riverside, CA). Inspiratory gas was warmed and humidified. PaCO_2 was maintained in a range between 27 and 45 torr.

In the hypoxic hypoxia group, progressive graded hypoxia was induced in eight piglets by changing the proportion of oxygen/nitrogen in the inspiratory gas. A variable fractional inspiratory oxygen concentration (FiO_2) of 0.30 to 0.75 was used during baseline. Progressive graded hypoxia was induced by a FiO_2 of 0.21, 0.15, and 0.10, respectively. Every fixed FiO_2 was kept for an interval of 15 minutes. At the end of each interval, blood samples were simultaneously taken from the carotid artery for blood gases, hemoglobin, lactate and pyruvate and from the pulmonary artery for blood gases and hemoglobin. The study included two series of different FiO_2 intervals with a one hour period in between for recovery. The series of graded hypoxia, could be repeated because all animals except one regained a stable condition after the first series of the FiO_2 intervals and consequently more data could be obtained per animal. A blood transfusion was given for accidental blood loss during the operative procedure or when the hemoglobin level decreased because of blood sampling during the experiment to maintain a baseline hemoglobin level between 7.0 and 12 gr/dl.

In the anemic hypoxia group, progressive anemia was induced in eight piglets by a repeated isovolemic exchange transfusion with 50 ml pasteurized plasma. Thirty minutes after each exchange transfusion, blood samples were taken for the same determinations as in the hypoxic hypoxia group. During this procedure, piglets were ventilated with a FiO_2 of 1.0.

Measurements

The fiberoptic catheter system (Oximatrix System, Abbott Laboratories) includes a fiberoptic catheter and a microprocessor-based oximeter. The back scattered light from the oxygenated and deoxygenated blood is transmitted by the receiving fiberoptic bundle to a photodetector. The oximeter computes percentage of oxygen saturation values based on the electrical signals from the photodetector.

SaO_2 , SvO_2 , arterial and pulmonary artery blood pressures, blood flow in the aorta, heart rate and core body temperature were sampled every 10 secs, displayed and stored by a software system (Hemodynamic Data Acquiring System (HDAS), University of Maastricht, the Netherlands). Blood samples for blood gases were analyzed on an ABL 3 acid-base analyzer (Radiometer, Copenhagen, Denmark) corrected for the animal core body temperature. SaO_2 , SvO_2 and hemoglobin (gr/dl) were analyzed on the OSM2 oximeter (Radiometer, Copenhagen, Denmark). Lactate and pyruvate were analyzed in arterial blood samples by an enzymatic spectrophotometric method (Sigma Chemical St Louis, MO).

Calculations

- Arterial oxygen content (CaO_2) and venous oxygen content (CvO_2) were calculated as the product of hemoglobin concentration, hemoglobin oxygen saturation and a hemoglobin oxygen binding capacity of 1.36 ml/dl.

- Cardiac output (Qt) was determined by the mean value of the blood flow in the aorta during the last five minutes of each FiO_2 interval or hemoglobin interval measured by the flowsensor.

- DO_2 and VO_2 were calculated as the product of the cardiac output and arterial oxygen content and the difference between arterial and venous oxygen content, respectively.

- Fractional oxygen extraction (F.O.E.) was calculated as the VO_2 divided by DO_2 .
- P_{50} was calculated from venous blood gas samples using SvO_2 and PvO_2 as has been described in the literature.^[9]

- The change of the parameters was calculated in comparison with the baseline value of these items in each individual animal.
- Oxygen debt was calculated by multiplying the change in VO₂ in a certain interval by the duration of that interval.

Data analysis

Data were retrospectively standardized to four DO₂ levels: DO₂ values above 18 ml.min⁻¹.kg⁻¹, DO₂ values between 12 and 18 ml.min⁻¹.kg⁻¹, DO₂ values between 6 and 12 ml.min⁻¹.kg⁻¹, and DO₂ values below 6 ml.min⁻¹.kg⁻¹.

Data at baseline are presented as the median (25th - 75th percentile) of the absolute values. Data standardized to the four DO₂ levels are presented as the median value (25th - 75th percentile) of the changes in comparison with the baseline value in each individual animal.

To test whether a difference existed in the hypoxic hypoxia between the two consecutive series of induced hypoxemia the results in the corresponding FiO₂ intervals were compared using a Mann-Whitney U test.

Data of the different DO₂ levels within each type of hypoxia group were compared using the Kruskal-Wallis one way of variance. When the null hypothesis that a parameter had an equal distribution in all DO₂ levels, could be rejected, a Mann-Whitney U test between the different DO₂ levels and the highest DO₂ was performed adjusting the significance level for the number of comparisons by performing the Bonferroni test.

Between the types of hypoxia data of the same DO₂ level were compared using a Mann-Whitney U test.

The critical value of DO₂ was calculated for each piglet, that is the DO₂ below which the change in lactate concentration increased rapidly. To do this, we adapted a technique from Mellits^[10-12] to find the section point of two lines of regression. This included constructing the best pair of regression lines that included all points only once. The best pair of regression lines was defined as to minimize the total of the residual sum of squares of each line in the pair. Their intersection point was determined by solving the equations of the pair of regression lines. The x-coordinate of the intersection point is the critical DO₂^[11]. The critical DO₂ for each piglet was also determined for change in lactate/pyruvate ratio and change in VO₂ applying this same technique. Likewise, critical values for each piglet of CaO₂, SvO₂ and of FOE were determined. To compare these critical values between hypoxic hypoxia and anemic hypoxia a Mann-Whitney U test was performed.

Linear regression analysis was performed to relate DO₂ and SvO₂ (least squares method) in the hypoxic hypoxia and the anemic hypoxia group. To test whether linear regression was appropriate, an analysis of the residuals was performed. The linear regression equations in hypoxic hypoxia and anemic hypoxia were contrasted using covariant analysis.

Results

Two groups of eight piglets were studied; they had a median weight of 3.2 (2.8 , 4.1) Kg and a median age of 9 (8 , 12) days.

No significant differences were found between the results of corresponding FiO₂ intervals in the two consecutive series of induced hypoxemia.

Table 1 shows the measured and calculated data during hypoxic hypoxia standardized for DO₂. SaO₂ and SvO₂ decreased in the lowest DO₂ level 77 % and 55 % from baseline, respectively.

table 1: Measured and calculated data during baseline and the change from baseline standardized to four levels of DO₂ during hypoxic hypoxia. Values are median (25th, 75th percentile)

DO ₂ levels (ml.min ⁻¹ .kg ⁻¹)		DO ₂ levels (ml.min ⁻¹ .kg ⁻¹)			
baseline		> 18 change	12 - 18 change	6 - 12 change	< 6 change
pH _a	7.45 (7.40, 7.47)	0.01 (-0.05, 0.05)	0.07 (0.03, 0.10)	0.03 (-0.14, 0.08)	-0.02 (-0.09, 0.00)
PaCO ₂ (torr)	33 (28, 41)	0 (-4, 4)	3 (-7, -1)	-6 (-12, 3)	-3 (-12, 2)
PaO ₂ (torr)	315 (97, 465)	-13 (-127, 12)	-201 (-375, -15) *	-275 (-360, -43) *	-292 (-435, -82) *
SaO ₂ (%)	97 (89, 98)	0 (-2, 2)	3 (-17, 0) *	-48 (-53, -11) *	-77 (-78, -69) *
CaO ₂ (ml/dl)	12.4 (11.0, 14.5)	0 (-0.5, 0.9)	-1.6 (-3.6, -0.8) *	-5.6 (-7.6, -1.7) *	-10.4 (-11.0, -8.6) *
PvO ₂ (torr)	37 (35, 43)	-3 (-7, 3)	-12 (-18, -6) *	-19 (-26, -13) *	-30 (-39, -25) *
SvO ₂ (%)	54 (47, 61)	-2 (-6, 3)	-15 (-26, -7) *	-40 (-52, -25) *	-55 (-63, -50) *
CvO ₂ (ml/dl)	7.2 (6.1, 8.2)	-0.2 (-0.6, 0.6)	-2.3 (-3.9, -1.0) *	-4.3 (-5.8, -2.3) *	-7.2 (-8.3, 5.8) *
Ca-vO ₂ (ml/dl)	5.4 (4.5, 7.7)	0 (-0.3, 0.3)	0.0 (0.5, 0.5)	-0.7 (-1.7, 0.7)	-2.9 (-4.2, -2.3) *
P ₅₀ (torr)	34 (32, 37)	0 (0, 0)	2 (-7, 0)	-1 (-5, 6)	-1 (-10, 6)
Hb (g/dl)	9.3 (8.1, 9.7)	0 (-1.2, 0.12)	-0.0 (0.3, 0.6)	0.8 (0.1, 1.6)	2.2 (0.6, 3.0) *
Qt (ml.min ⁻¹ .kg ⁻¹)	165 (118, 218)	0 (-10, 7)	2 (-9, 8)	2 (-9, 18)	-22 (-54, -2)
mean P _{ao} (mmHg)	90 (80, 105)	0 (-3, 5)	2 (-7, 1)	-7 (-15, 1) *	-25 (-37, -16) *
mean P _{pulm} (mmHg)	17 (14, 26)	0 (-1, 4)	2 (-1, 4)	5 (0, 11)	5 (2, 10) *
Lactate (mmol/l)	2.05 (1.2, 3.1)	0 (0, 0)	0.40 (0.00, 0.68) *	1.50 (0.53, 3.31) *	3.85 (2.02, 7.12) *
Pyruvate (μmol/l)	169 (105, 205)	0 (-1, 1)	13 (-14, 63)	36 (5, 66)	33 (-15, 132)
lact/pyr.ratio	11.0 (9.5, 13.5)	0 (0, 0.8)	1.1 (0.5, 3.3) *	7.1 (3.4, 13.9) *	24.4 (16.1, 32.9) *
DO ₂ (ml.min ⁻¹ .kg ⁻¹)	19.6 (16.4, 21.3)	0 (-0.9, 1.0)	-2.9 (-3.9, -1.2) *	-6.2 (-11.7, -3.8) *	-16.4 (-18, -11) *
VO ₂ (ml.min ⁻¹ .kg ⁻¹)	8.7 (7.5, 10.0)	0.0 (-0.2, 0.6)	-0.4 (-0.7, 0.1)	-1.3 (-1.8, -0.4) *	-4.4 (-5.1, -3.7) *
FOE	0.47 (0.39, 0.52)	0 (-0.04, 0.04)	0.11 (0.0, 0.19) *	0.26 (0.15, 0.35) *	0.49 (0.41, 0.54) *
O ₂ debt (ml.kg ⁻¹)		-1.5 (-16, 3.6)	-1.0 (-3.9, 10.1)	22.6 (4.9, 40.1) *	51.3 (35.2, 89.4) *

table 2: Measured and calculated data during baseline and the change from baseline standardized to four levels of DO₂ during anemic hypoxia. Values are median (25th, 75th percentile)

DO ₂ levels (ml.min ⁻¹ .kg ⁻¹)	baseline	> 18 change	12 - 18 change	6 - 12 change	< 6 change
pHa	7.44 (7.42, 7.49)	0.00 (-0.01, 0.00)	-0.02 (-0.06, 0.0) §	-0.09 (-0.18, -0.05) *§	0.22 (-0.45, 0.06) *§
PaCO ₂ (torr)	32 (28, 36)	0 (0, 1)	0 (0, 3) §	1 (0, 5) §	-3 (-6, 2)
PaO ₂ (torr)	490 (450, 510) §	0 (-9, 16) §	1 (-3, 3) §	9 (-32, 27)	13 (-32, 57)
SaO ₂ (%)	99 (98, 99) §	0 (0, 0)	-1 (-2, 0) §	-2 (-3, 0) §	-6 (-10, 0) §
CaO ₂ (ml/dl)	10.3 (9.8, 11.7)	0 (-1.6, 0.1)	-3.2 (-4.5, 0.1) *	-6.3 (-7.1, -5.6) *	-8.4 (-9.8, -7.1) *
PvO ₂ (torr)	43 (37, 54)	0 (-1, 2)	-4 (-6, 1) §	-7 (-13, -2) *§	3 (-12, 7) §
SvO ₂ (%)	54 (45, 71)	0 (-1, 2)	-6 (-10, -1) *§	-23 (-30, -16) *§	-26 (-36, -6) *§
CvO ₂ (ml/dl)	5.1 (4.8, 6.4) §	0 (-1.1, 0.1)	-2.6 (-3.5, -0.5) *	-4.7 (-5.7, 3.7) *	-5.2 (-7.3, -4.1) *§
Ca-vO ₂ (ml/dl)	5.5 (3.7, 6.4)	0 (-0.3, 0.3)	-0.4 (-0.9, 0.0)	-1.3 (-2.4, 0.9)	-3.2 (4.1, -1.4) *
P ₅₀ (torr)	39 (36, 42) §	0 (-1, 1)	2 (0, 6) §	7 (3, 13) *§	19 (7, 40) *§
Hb (gr/dl)	6.6 (6.3, 7.6) §	0 (-1.2, 0.1)	-2.3 (-3.4, -0.1) *§	-4.6 (-5.3, 3.9) *§	-6.2 (-7.2, -5.2) *§
Qt (ml.min ⁻¹ .kg ⁻¹)	180 (127, 215)	3 (-4, 17)	19 (0, 40) §	45 (1, 64) §	-39 (-78, 24)
mean P _{ao} (mm Hg)	81 (77, 89)	0 (-3, 2)	-5 (-15, 1)	-28 (-35, -17) *§	-35 (-60, -13) *
mean P _{pulm} (mm Hg)	20 (18, 24)	0 (-3, 6)	0 (-3, 1)	0 (-1, 1)	1 (-5, 22)
Lactate (mmol/l)	1.03 (0.65, 1.25)	-0.03 (-0.1, 0.05)	-0.12 (-0.22, 0.01) §	0.47 (-0.03, 2.40) *	8.52 (5.07, 11.2) *§
Pyruvate (μmol/l)	107 (86, 120)	-2 (-11, 1)	-9.5 (-26, 1)	10 (-18, 49)	106 (-6.5, 197) *§
Lact/pyr.ratio	9.1 (7.7, 10.2) §	0.0 (-0.6, -1)	0.0 (-0.6, 0.6) §	3.5 (1.4, 12.3) *	24.0 (20.1, 29.1) *
DO ₂ (ml.min ⁻¹ .kg ⁻¹)	18.5 (14.5, 22.1)	0 (-1, 0.5)	-4.7 (-6.6, -0.6) *	-9.5 (-12.8, -5.5) *	-14.4 (-17.6, -12.5) *
VO ₂ (ml.min ⁻¹ .kg ⁻¹)	8.3 (7.1, 10.4)	0 (-0.3, 0.2)	-0.4 (-0.7, 0.0)	-1.0 (-2.0, -0.5) *	-5.4 (-6.8, -3.5) *
FOE	0.52 (0.37, 0.59)	0 (0, 0.02)	0.09 (0.03, 0.15) *	0.29 (0.20, 0.32) *	0.31 (0.20, 0.41) *§
O ₂ debt (ml.kg ⁻¹)		0 (-0.7, 7)	5.3 (-0.8, 15.1)	31.4 (14.3, 62.1) *§	135.9 (100.4, 214) *§

To convert torr to kPa multiply the value by 0.1333

DO₂, oxygen delivery; PaCO₂, arterial partial pressure of carbon dioxide; PaO₂, arterial partial pressure of oxygen; SaO₂, arterial oxygen saturation; CaO₂, arterial oxygen content; PvO₂, venous partial pressure of oxygen; SvO₂, mixed venous oxygen saturation; CvO₂, venous oxygen content; Ca-vO₂, difference between arterial and venous oxygen content; P₅₀, partial pressure of oxygen at 50 % oxygen saturation; Hgb, hemoglobin; Qt, cardiac output; mean P_{ao}, mean blood pressure in the aorta; mean P_{pulm}, mean bloodpressure in the pulmonary artery; lact/pyr.ratio, ratio between lactate and pyruvate; VO₂, oxygen consumption; F.O.E, fractional oxygen extraction;

* = significant different from the highest DO₂ level. § = significant different between alveolar and anemic hypoxia in the similar DO₂ level.

figure 1. Relationship of oxygen delivery (DO_2) and mixed venous oxygen saturation (SvO_2) during hypoxic (A) and anemic hypoxia (B). The linear regression equation for hypoxic hypoxia ($2.93 \cdot \text{DO}_2 - 9.93$) was significantly different from the regression line during anemic hypoxia ($1.91 \cdot \text{DO}_2 + 21.70$). ($p < 0.001$, covariant analysis).

VO_2 decreased significantly 1.3 (0.4, 1.8) $\text{ml.kg}^{-1}.\text{min}^{-1}$ and 4.4 (3.7, 5.1) $\text{ml.kg}^{-1}.\text{min}^{-1}$, respectively in the two lowest levels of DO_2 . Consequently, in these DO_2 levels, lactate and the lactate/pyruvate ratio increased significantly, whereas pyruvate did not change from baseline. Hemodynamic responses during the different DO_2 levels consisted of an unaltered cardiac output and systemic mean blood pressure except at the lowest DO_2 level, at which systemic blood pressure decreased. In contrast to systemic blood pressure, pulmonary mean blood pressure increased with increasing hypoxic hypoxia.

figure 2: Oxygen delivery (DO_2) and change lactate during hypoxic and anemic hypoxia. The median of the critical values of DO_2 determined in each individual piglet with regard to change lactate were not different between hypoxic and anemic hypoxia: $11.5 \text{ ml.kg}^{-1}.\text{min}^{-1}$ (10.4 , 12.8) vs. $11.1 \text{ ml.kg}^{-1}.\text{min}^{-1}$ (10.2 , 12.8), respectively.

Table 2 shows measured and calculated data for the anemic hypoxia group standardized for DO_2 . Baseline values for PaO_2 (490 torr (450 , 510)) were significantly higher in comparison with baseline values in the hypoxic hypoxia group (315 torr (97 , 465)). Baseline hemoglobin and CvO_2 were significantly higher in the hypoxic hypoxia group in comparison with the anemic hypoxia group. The baseline value of p_{50} was higher in the anemic hypoxia group. SvO_2 and PvO_2 decreased during the progressive decrease in DO_2 , whereas SaO_2 and PaO_2 remained largely unaltered. At the lowest DO_2 level of anemic hypoxia, the change in SvO_2 was less than that in hypoxic hypoxia (-26 % (-36 , -6) vs. -55 % (-63 , -50)). In fact, in anemic hypoxia, the median change of PvO_2 at the lowest DO_2 level showed an increase (3 torr (-12 , 7)) compared with the baseline value. O_2 debt increased and was significantly different from baseline in the two lowest DO_2 levels during both types of hypoxia. The increase of O_2 debt was more pronounced during anemic hypoxia than during hypoxic hypoxia.

In figure 1, we plotted the relationship between DO_2 and SvO_2 and a linear regression analysis was performed. Residual analysis showed an equal variance and a normal distribution. When the residuals were plotted against the values predicted by the regression line, a random distribution was observed. This analysis showed that necessary assumptions for linear regression were not violated.

figure 3. Arterial oxygen content (CaO_2) and change lactate during hypoxic and anemic hypoxia. The median of the critical values of CaO_2 determined in each individual piglet below which lactate rapidly increased was significantly different between hypoxic and anemic hypoxia: 8.6 ml/dl (6.4 , 10.9) vs. 4.1 ml/dl (4.0 , 6.3), respectively; $p = 0.002$

The regression equation for hypoxic hypoxia ($2.93 \cdot \text{DO}_2 - 9.93$; $r^2 : 0.72$) was significantly different from the regression equation for anemic hypoxia ($1.91 \cdot \text{DO}_2 + 21.70$; $r^2 : 0.41$). ($p < 0.001$). This means that the decrease in SvO_2 for an identical decrease in DO_2 is significantly less during anemic hypoxia than during hypoxic hypoxia.

VO_2 was decreased in both types of hypoxia. Consequently, lactate and the lactate/pyruvate ratio were significantly increased at the two lowest DO_2 -levels. However, in anemic hypoxia, lactate and pyruvate increased more at the lowest DO_2 level. In the anemic hypoxia group, F.O.E. did not change in equal amount as in hypoxic hypoxia: 0.31 (0.20 , 0.41) vs. 0.49 (0.41 , 0.54). Hemodynamic responses on decreasing CaO_2 consisted of an increase in cardiac output and a decrease in systemic blood pressure. At the lowest DO_2 level, cardiac output and systemic blood pressure decreased below baseline values. Pulmonary artery pressure remained stable at all DO_2 levels in the anemic hypoxia group in contrast to the increased pulmonary pressure in the hypoxic hypoxia group.

Figure 2 and 3 show the relationships between DO_2 and CaO_2 and lactate, respectively. The critical values of DO_2 and CaO_2 for each individual piglet concerning the change in lactate, in the lactate/pyruvate ratio and in VO_2 were calculated according to the technique of Mellits^[10]. There were no differences in the critical range of DO_2 in hypoxic hypoxia and anemic hypoxia, whereas the critical range of CaO_2 was lower in the anemic

table 3. Critical values of arterial oxygen content (caO₂), of oxygen delivery (DO₂), of mixed venous oxygen saturation (SvO₂) and of fractional oxygen extraction (F.O.E.), below which lactate, lactate / pyruvate ratio increased and oxygen consumption decreased. Values are median (25th , 75th percentile).

Oxygen transport variables		Hypoxic hypoxia		Anemic hypoxia	
caO ₂ crit (ml/dl) *	lactate	8.6	(6.4 , 10.9)	4.1	(4.0 , 6.3)
	lactate / pyruvate ratio	7.5	(6.3 , 11.8)	4.5	(3.5 , 6.9)
	VO ₂	7.2	(6.6 , 9.6)	5.4	(4.0 , 6.0)
DO ₂ crit (ml.min ⁻¹ .kg ⁻¹)	lactate	11.5	(10.4 , 12.8)	11.1	(10.2 , 12.8)
	lactate / pyruvate ratio	12.4	(12.0 , 13.0)	12.0	(10.6 , 13.9)
	VO ₂	13.1	(11.4 , 13.8)	12.9	(10.0 , 13.8)
SvO ₂ crit (%) *	lactate	14	(11 , 24)	38	(30 , 48)
	lactate / pyruvate ratio	16	(14 , 21)	41	(26 , 48)
	VO ₂	15	(14 , 21)	37	(27 , 47)
F.O.E.crit	lactate	0.73	(0.66 , 0.75)	0.69	(0.67 , 0.78)
	lactate /pyruvate ratio	0.72	(0.67 , 0.73)	0.68	(0.68 , 0.77)
	VO ₂	0.72	(0.71 , 0.74)	0.71	(0.64 , 0.76)

** = significant different between alveolar and anemic hypoxia. DO₂crit, critical value of oxygen delivery; SvO₂crit, critical value of mixed venous oxygen saturation; caO₂crit, critical value of arterial oxygen content; F.O.E.crit, critical value of fractional oxygen extraction; VO₂, oxygen consumption.

hypoxia group (3.5 to 6.9 ml/dl) than in the hypoxic hypoxia group (6.3 to 11.8 ml/dl) (table 3). The critical range for SvO₂ below which lactate, the lactate/pyruvate ratio and VO₂ rapidly changed from baseline value was significantly lower in hypoxic hypoxia than in anemic hypoxia (11 to 24 %) *vs* (26 to 48 %), respectively (table 3). Lactate and the lactate/pyruvate ratio increased and VO₂ decreased at a low and small range of SvO₂ values during hypoxic hypoxia, whereas during anemic hypoxia lactate and the lactate/pyruvate ratio increased and VO₂ decreased at a higher and larger range of SvO₂ values. (figure 4). No differences were found in the critical value of F.O.E.

Discussion

This study showed that in the case of anemic hypoxia, measuring SvO₂ reflects better a reduced DO₂ than does measuring SaO₂. It also demonstrated that the critical range of SvO₂ - the main parameter of residual oxygen in venous blood after tissue oxygen extraction - was 26 % to 48 % in anemic hypoxia, whereas the critical range in hypoxic hypoxia was 11 % to 24 %. However, the critical values of DO₂ and FOE showed no difference in hypoxic and anemic hypoxia. The decrease in SvO₂ for an identical reduction of DO₂ was significantly less in anemic hypoxia than in hypoxic hypoxia. These findings suggest that in anemic hypoxia, oxygen unloading from the blood to tissue is impaired. In agreement with this, our study found that fractional oxygen extraction (FOE) above the critical value at the lowest DO₂ level was decreased in anemic hypoxia in comparison with hypoxic hypoxia.

figure 4. Change lactate, lactate/pyruvate ratio and change oxygen consumption (VO_2) as a function of mixed venous oxygen saturation (SvO_2) during hypoxic and anemic hypoxia. Note, that during anemic hypoxia higher SvO_2 values are associated with increased lactate, lactate/pyruvate ratio and decreased VO_2 .

Before addressing the possible reasons for these findings, the model we used will be discussed. To compare the critical values of SvO₂ in hypoxic and anemic hypoxia, a progressive graded hypoxia was induced either by administering different nitrogen/oxygen mixtures or performing repeated exchange transfusions. To compare these two different types of hypoxia, a FiO₂ of 1.0 was maintained during anemic hypoxia to eliminate any element of hypoxemia.

Differences between the two groups with respect to hemoglobin level and P₅₀ during baseline condition, could affect the response on graded progressive hypoxia. However, these differences did not have any influence on the baseline values of lactate, pyruvate, VO₂, F.O.E. or of SvO₂. The low hemoglobin level was compensated by an increased cardiac output. The shift of the oxygen dissociation curve affected possibly the value of PvO₂ but not of SvO₂.

Because of the prolonged procedure of exchange transfusion, the duration of the intervals was longer in the anemic hypoxia group (30 minutes) than in the hypoxic hypoxia group (15 minutes). The concept of O₂ debt was introduced to account for this time aspect of the protocol. We analyzed the relation between O₂ debt and lactate to control for the different durations of the intervals during the two types of hypoxia. O₂ debt was linearly related to the change in lactate, but the coefficient of determination during hypoxic hypoxia and anemic hypoxia ($r^2 = 0.31$; $r^2 = 0.73$, respectively), was not different from the coefficient of determination between the change in VO₂ and the change in lactate ($r^2 = 0.37$; $r^2 = 0.67$). O₂ debt introduced to consider the different intervals in the model was not a better parameter of oxygen restricted metabolism than change in VO₂. In both types of hypoxia, lactate increased significantly in the two lowest DO₂ levels from baseline (table 1,2). The values of lactate and the lactate/pyruvate ratio were only used to determine the critical values of different parameters such as DO₂, CaO₂ and SvO₂, below which lactate and the lactate/pyruvate increased. The increase itself was less important than the moment when the lactate and lactate pyruvate ratio started to increase. In agreement with this, critical values of the different parameters were not different when calculated either for a significant change in lactate or for a change in VO₂, (table 3). Hence, we think that the findings in this study are a consequence of the differences in oxygen transport and not of the model we used.

There are several possible reasons to explain the higher values of SvO₂ and PvO₂ in anemic hypoxia in comparison with hypoxic hypoxia:

a) The reliability and accuracy of the fiberoptic catheter readings can be questioned at these low hemoglobin values. However, in several reports ^[13,14] different fiberoptic catheter systems, including the one we used, reflected SvO₂ values accurately during abrupt changes in cardiac output and hematocrit.

b) In anemic hypoxia, the full range of the oxyhemoglobin dissociation curve is used because even at the lowest DO₂ level, high arterial SaO₂ and PaO₂, and consequently also higher values of PvO₂ and SvO₂, are found ^[15,16].

c) Another possible explanation is the rightward shift in the oxyhemoglobin dissociation curve in anemic hypoxia. In the present study, P₅₀ was increased during anemic hypoxia in comparison with hypoxic hypoxia, especially in the lower DO₂ levels. However, one must be cautious to interpret the calculated P₅₀ values at oxygen saturation values below 20 %, because the changing value of n in the Hill equation (because then only one heme group is involved in the oxygen binding) could cause a significant error in the calculated value of P₅₀ ^[9,17].

The position of the oxyhemoglobin dissociation curve can be modulated by Cl⁻ and inorganic phosphate ions, by the level of 2,3-diphosphoglycerate, by the value of pH and by the hemoglobin level. In a bovine animal model it was demonstrated that the

oxygen dissociation curve can be shifted to the right by the infusion of Cl and inorganic phosphate ions containing fluids [18]. In our study exchange transfusions were performed with pasteurized plasma containing normal electrolyte concentrations, but changes in Cl and inorganic phosphate ions after the exchange transfusions cannot be excluded. In adult patients anemia was associated with an increased level of 2,3-diphosphoglycerate, although in neonatal blood 2,3-diphosphoglycerate was not correlated with the hemoglobin concentration [17,19,20]. In the present study pH decreased significantly in the last two DO₂ levels during anemic hypoxia in comparison with hypoxic hypoxia. This could be another reason for a rightward shift of the oxyhemoglobin curve during anemic hypoxia. However, studies examining the effect of a rightward shift in the oxyhemoglobin dissociation curve during normoxic and anemic hypoxic conditions demonstrated that differences in oxygen affinity have consequences for PvO₂, but not or in a much lesser degree for SvO₂ values or CvO₂ [21,22,23].

d) Higher SvO₂ values in anemic hypoxia can be explained through a redistribution of blood flow to different organs. SvO₂ does not reflect the oxygen transport balance of any one individual organ, but is an average of saturation data from all perfused tissues, strongly affected by the blood flow through these tissues [24]. When the perfusion through high-blood flow and low-oxygen extraction organs increases, the SvO₂ value also increases.

During hypoxic hypoxia, a variable and transient response of cardiac output was reported consisting of a redistribution of the regional blood flow in favor of heart and brain and a decrease to the skin, gastrointestinal tract, kidneys and spleen, whereas blood flow to the carcass remained unchanged [25,27]. The response in lambs and adult sheep was qualitatively similar for heart and brain with an increase in blood flow, whereas the response differed for the skin, kidney and gastrointestinal tract, in which flow decreased in neonatal sheep and was unchanged or increased in adult sheep [28].

During anemic hypoxia on anesthetized dogs [29], a marked increase in cardiac output was found, with a redistribution in favor of the myocardium and brain. Kidney, liver and gastro-intestinal tract blood flow remained unchanged, which resulted in a linear decrease in oxygen transport to these organs. In anesthetized pigs [30], isovolemic anemic hypoxia resulted in a two-fold increase in cardiac output, with a redistribution in favor of the heart and brain. However, flow increased in all organs including the liver, gastrointestinal tract and kidney.

Interpreting these different studies performed in different species at different ages is difficult. It is clear that in all species, in both hypoxic hypoxia and anemic hypoxia, a regional redistribution of blood flow occurs in favor of the heart and brain. During hypoxic hypoxia, the response in other organs seems to be dependent on age of the animal, with a trend toward more vasoconstriction in younger animals [28]. During anemic hypoxia, regional redistribution of the blood flow to other organs than the heart and brain was studied in adult animals and is reported to be unchanged or increased [29,30]. The difference in regional redistribution and the unchanged or increased flow to low-oxygen extraction organs in anemic hypoxia can explain to a certain extent the higher SvO₂ value observed in anemic hypoxia.

e) Another reason for higher values of PvO₂ and SvO₂ during anemic hypoxia is an impairment of oxygen uptake at the tissue level. At the lowest DO₂ level, we found a lower FOE during anemic hypoxia in comparison with the lowest DO₂ level during hypoxic hypoxia (table 2). Tissue oxygen uptake in anemic hypoxia has been related to increased red cell spacing and capillary architecture [31]. Hemodilution induces an increased red cell spacing in the capillary bed. The induced expansion of the plasma phase, with its very low oxygen solubility, limits the ability to transfer oxygen from the red cells to the tissues and affects the uniformity of oxygen flux at the capillary wall [32,33].

The influence of microvascular architecture of the tissues on oxygen exchange was studied in the cheek retractor muscle of the hamster^[34,35]. In these studies, a more complex tissue oxygenation was demonstrated than Krogh's oxygen transport model, which postulated that oxygen is transferred to the capillaries by convection and released from these capillaries by diffusion. The release of oxygen from a capillary seems to be strongly influenced by its proximity to arterioles and venules, which either supply oxygen to the capillaries or augment its release. These diffusive interactions might change under different hemodynamic conditions. The effect of limited isovolemic hemodilution on microcirculatory hemodynamics and the rate of oxygen transport was studied^[36]. Because the red blood cell velocity is increased and the red blood cell residence time is reduced during isovolemic hemodilution, a decrease in precapillary oxygen release and in diffusional oxygen transfer to other microvessels was found.

These proposed mechanisms could explain to some extent the higher PvO₂ and SvO₂ values found in our study during anemic hypoxia in comparison with hypoxic hypoxia. Therefore, our hypothesis that similar critical SvO₂ values would be observed in both hypoxic hypoxia and anemic hypoxia was not confirmed.

Previous studies described and compared oxygen transport parameters during different types of hypoxia. The effect of hemoglobin concentration on critical oxygen transport was examined during progressive reductions of cardiac output in one-month old lambs^[15]. The authors reported a range of critical PvO₂ values (31 - 36 mmHg), irrespective of the variation of hemoglobin concentrations. The difference between these results and ours may be explained in several ways. In this study hemoglobin was only moderately reduced, two factors of oxygen delivery -hemoglobin and cardiac output- were simultaneously modulated and SaO₂ was equivalent under all hypoxic conditions. Our results were consistent with the findings of Cain^[16,37], who compared hypoxic hypoxia and anemic hypoxia in adult anesthetized dogs and reported different critical values of PvO₂ of 17 mmHg and 45 mmHg, respectively. A critical value of SvO₂ of 44 %, similar with our findings, was reported by Trouwborst et al.^[38] during anemic hypoxia in adult pigs, but was not compared with critical values during hypoxic hypoxia. Developmental aspects could account for the different critical F.O.E. value of 0.57 in this adult pig model.

We conclude that the critical value of SvO₂ is dependent on the type of hypoxia and is higher in anemic hypoxia. Although developmental and interspecies aspects should be not be disregarded, these characteristics in oxygen transport and capillary hemodynamics should be taken into consideration, when SvO₂ is used in clinical critical care.

References

- 1 O'Connor TA, Hall RT: Mixed venous oxygenation in critically ill neonates. *Crit Care Med* 1994;22:343-346
- 2 Bartlett RH, Cilley RE: Physiology of extracorporeal life support. In: Arensman RM, Cornish JD (eds). *Extracorporeal life support*. Boston, Blackwell, 1993: pp 89-104
- 3 Dudell G, Cornish JD, Bartlett RH: What constitutes adequate oxygenation? *Pediatrics* 1990;85:39-41
- 4 Hoeven MAHBM van der, Maertzdorf WJ, Blanco CE: Feasibility and accuracy of a fiberoptic catheter for the measurement of venous oxygen saturation in newborn infants. *Acta Paediatr* 1995;84:122-127
- 5 Hoeven MAHBM van der, Maertzdorf WJ, Blanco CE: Continuous central venous oxygen saturation (ScvO₂) measurement using a fibreoptic catheter in newborn infants. *Arch Dis Child* 1996;74:f177-f181

- 6 Kreisberg RA: Lactate homeostasis and lactic acidosis. *Ann Intern Med* 1980;92(1):227-237
- 7 Mizock BA, Falk JL: Lactic acidosis in critical illness. *Crit Care Med* 1992;20:80-93
- 8 Hoeven MAHBM van der, Maertzdorf WJ, Blanco CE: Mixed venous oxygen saturation (SvO₂) and biochemical parameters of hypoxia during progressive hypoxemia in 10-14 day old piglets. *Pediatr Res* 1997;42:878-884
- 9 Brown EG, Krouskop RW, McDonnel FE et al: A technique to continuously measure arteriovenous oxygen content difference and p₅₀ in vivo. *J Appl Physiol* 1985;58:1383-1389.
- 10 Mellits ED: Statistical methods. In: Cheek DB (ed) *Human Growth*. Philadelphia, Lea & Febiger, 1968: 19-38
- 11 Fahey JT, Lister G: Postnatal changes in critical cardiac output and oxygen transport in conscious lambs. *Am J Physiol* 1987;253:H100-H106
- 12 Samsel RW, Schumaker PT: Determination of the critical O₂ delivery from experimental data: sensitivity to error. *J Appl Physiol* 1988;64:2074-2082
- 13 Woerkens ESM van, Trouwborst A, Tenbrinck R: Accuracy of a mixed venous saturation catheter during acutely induced changes in hematocrit in humans. *Crit Care Med* 1991;19:1025-1029
- 14 Woerkens ECSM van, Trouwborst A, Tenbrinck R: Comparative study of the accuracy of two fiberoptic mixed venous saturation catheters (Spectracath vs Opticath) during acute in hematocrit and cardiac output in humans. In: Erdmann W, Bruley DF. eds. *Oxygen transport to tissue XIV*. New York, Plenum Press, 1992: 509-514
- 15 Heusser F, Fahey JT, Lister G: Effect of hemoglobin concentration on critical cardiac output and oxygen transport. *Am J Physiol* 1989;253:H527-H532
- 16 Cain SM: Peripheral oxygen uptake and delivery in health and disease. *Clinics in Chest Med* 1982;4:139-148
- 17 Wimberley PD: Fetal hemoglobin, 2,3-diphosphoglycerate and oxygen transport in the newborn premature infant. *Scand J Clin Lab Invest* 1982;42:Supp 160:1-149
- 18 Gustin P, Detry B, Cao ML, et al: Chloride and inorganic phosphate modulate binding of oxygen to bovine red blood cells. *J Appl Physiol* 1994;77:202-208
- 19 Nunn JF: The carriage of oxygen in the blood. in Nunn's applied respiratory physiology 4th ed. Oxford, Butterworth-Heinemann, 1993: 269-282
- 20 Robertie PG, Gravlee GP: Safe limits of isovolemic hemodilution and recommendations for erythrocyte transfusion. *Int Anesth Clin* 1990;28:197-204
- 21 Van Ameringen MR, Fouron JC, Bard H, et al: Oxygenation in anemic newborn lambs with high or low oxygen affinity red cells. *Pediatr Res* 1981;15:1500-1503
- 22 Fouron JC, Bard H, Guennec Le JC, et al: Effect of fetal or adult red cells on tissue oxygenation and myocardial function in normoxemic newborn lambs. *Pediatr Res* 1981;15:967-970.
- 23 Lister G: Oxygen transport in the intact hypoxic newborn lamb: Acute effects of increasing p₅₀. *Pediatr Res* 1984;16:172-177
- 24 Ceraianu AC, Nelson LD: The significance of mixed venous oxygen saturation and technical aspects of continuous measurement. In: Edwards JD, Shoemaker WC, Vincent JL (Eds) *Oxygen transport: Principles and practice*. London, Saunders, 1993: 99-124
- 25 Moss M, Moreau G, Lister G: Oxygen transport and metabolism in the conscious lamb: the effects of hypoxemia. *Pediatr Res* 1987;22:177-183
- 26 Sidi D, Kuipers JRG, Teitel D et al: Developmental changes in oxygenation and circulatory responses to hypoxemia in lambs. *Am J Physiol* 1983;245:H674-H682
- 27 Kuwahira I, Gonzales NC, Heisler N et al: Changes in regional blood flow distribution and oxygen supply during hypoxia in conscious rats. *J Appl Physiol* 1993;74:211-214
- 28 Koehler RC, Traystman RJ, Jones Jr MD: Regional blood flow and O₂ transport during hypoxic and CO hypoxia in neonatal and adult sheep. *Am J Physiol* 1985;245:H118-H124.
- 29 Fan FC, Chen RYZ, Schuessler GB et al: Effects of hematocrit variations on regional hemodynamics and oxygen transport in the dog. *Am J Physiol* 1980;238:H545-H552

- 30 Woerkens ECSM van, Trouwborst A, Duncker DJGM, et al: Catecholamines and regional hemodynamics during isovolemic hemodilution in anesthetized pigs. *J Appl Physiol* 1992;72:760-769
- 31 Biro GP, Anderson PJ, Curtis SE et al: Stroma free hemoglobin: its presence does not improve oxygen supply to the resting hindlimb vascular bed of hemodiluted dogs. *Can J Physiol Pharmacol* 1991;69:1656-1662
- 32 Federspiel WJ, Salerius RH: An examination of the contribution of red cell spacing to the uniformity of oxygen flux at the capillary wall. *Microvasc Res* 1984;27:273-285.
- 33 Popel AS: Theory of oxygen transport to tissues. *Crit Rev Biomed Engin* 1989;17:257-321.
- 34 Ellsworth ML, Pittman RN: Arterioles supply oxygen to capillaries by diffusion as well by convection. *Am J Physiol* 1990;258:H1240-H1243.
- 35 Pittman RN: Influence of microvascular architecture on oxygen exchange in skeletal muscle. *Microcirculation* 1995;2:1-18.
- 36 Kuo L, Pittman RN: Effect of hemodilution on oxygen transport in arteriolar networks of hamster striated muscle. *Am J Physiol* 1988;254:H331-H339.
- 37 Cain SM: Oxygen delivery and uptake in dogs during anemic and hypoxic hypoxia. *J Appl Physiol* 1977;42:228-234.
- 38 Trouwborst A, Tenbrinck R, Woerkens ECSM van: Blood gas analysis of mixed venous blood during normoxic acute isovolemic hemodilution in pigs. *Anaesth Analg* 1990;70:523-529

Acknowledgement: Supported by Abbott Inc, The Netherlands

20. Westlake KCM and Townsend A. (1968) *Quinone*. In: *Comprehensive Chemical Kinetics*, Vol. 1, Ed. Bamford, C., pp. 1-100. London: Butterworths.
21. Westlake KCM and Townsend A. (1968) *Quinone*. In: *Comprehensive Chemical Kinetics*, Vol. 1, Ed. Bamford, C., pp. 1-100. London: Butterworths.
22. Westlake KCM and Townsend A. (1968) *Quinone*. In: *Comprehensive Chemical Kinetics*, Vol. 1, Ed. Bamford, C., pp. 1-100. London: Butterworths.
23. Westlake KCM and Townsend A. (1968) *Quinone*. In: *Comprehensive Chemical Kinetics*, Vol. 1, Ed. Bamford, C., pp. 1-100. London: Butterworths.
24. Westlake KCM and Townsend A. (1968) *Quinone*. In: *Comprehensive Chemical Kinetics*, Vol. 1, Ed. Bamford, C., pp. 1-100. London: Butterworths.
25. Westlake KCM and Townsend A. (1968) *Quinone*. In: *Comprehensive Chemical Kinetics*, Vol. 1, Ed. Bamford, C., pp. 1-100. London: Butterworths.
26. Westlake KCM and Townsend A. (1968) *Quinone*. In: *Comprehensive Chemical Kinetics*, Vol. 1, Ed. Bamford, C., pp. 1-100. London: Butterworths.
27. Westlake KCM and Townsend A. (1968) *Quinone*. In: *Comprehensive Chemical Kinetics*, Vol. 1, Ed. Bamford, C., pp. 1-100. London: Butterworths.
28. Westlake KCM and Townsend A. (1968) *Quinone*. In: *Comprehensive Chemical Kinetics*, Vol. 1, Ed. Bamford, C., pp. 1-100. London: Butterworths.
29. Westlake KCM and Townsend A. (1968) *Quinone*. In: *Comprehensive Chemical Kinetics*, Vol. 1, Ed. Bamford, C., pp. 1-100. London: Butterworths.
30. Westlake KCM and Townsend A. (1968) *Quinone*. In: *Comprehensive Chemical Kinetics*, Vol. 1, Ed. Bamford, C., pp. 1-100. London: Butterworths.

Received for publication, September 15, 1968; accepted, November 15, 1968.

Chapter 5

Feasibility and accuracy of a fiberoptic catheter for the measurement of venous oxygen saturation in newborn infants.

Mark A.H.B.M. van der Hoeven, Wiel J. Maertzdorf, Carlos E. Blanco

Acta Paediatrica 1995;84:122-127

Abstract

The current parameters of oxygenation, arterial oxygen saturation and arterial oxygen pressure represent the availability of oxygen to the tissues. Venous oxygen saturation reflects oxygen extraction and could be a more useful indicator of tissue oxygenation. Therefore, we tested the feasibility and accuracy of a fiberoptic catheter, inserted into the right atrium via the umbilical vein, to measure venous oxygen saturation, continuously. In 24 of 36 (67%) infants the catheter could be placed into the right atrium. Blood samples were withdrawn through the catheter, analyzed with a hemoximeter and compared with simultaneously recorded oximeter values. The fiberoptic catheter readings correlated significantly with hemoximeter values and the mean difference of the 116 paired samples was -0.37%.

We conclude that the success rate and complications of the fiberoptic catheter were similar to previously published results. The oxygen saturation readings accurately represented oxygen saturation in the right atrium in newborn infants.

Introduction

For more than a decade continuous venous oxygen saturation measurements through a fiberoptic catheter in the pulmonary artery have been used in critically ill adult patients^[1,2,3] and postoperative cardiac surgical patients^[4,5]. In neonatal medicine the current clinical parameters of oxygenation, arterial oxygen pressure (paO_2) and arterial oxygen saturation (SaO_2) are poor indicators of tissue oxygenation, because they represent only the availability of oxygen to the tissues. The actual amount of oxygen in the blood, the arterial O_2 content (caO_2), is determined by the paO_2 , the hemoglobin concentration (Hb), O_2 carrying capacity and the SaO_2 .

$$caO_2 = paO_2 \times 0.0031 + 1.36 \times Hb \times SaO_2$$

Oxygen delivery (DO_2) to the tissues is determined by arterial oxygen content (caO_2) and the cardiac output (Qt).

$$DO_2 = caO_2 \times Qt$$

Overall tissue oxygen extraction (VO_2) is the product of cardiac output and the difference between arterial and mixed venous oxygen content (cvO_2).

$$VO_2 = Qt \times (caO_2 - cvO_2)$$

As the contribution of paO_2 to oxygen content can be neglected, the equation for VO_2 can be expressed as:

$$VO_2 = Qt \times 1.36 \times Hb \times 10 \times (SaO_2 - SvO_2)$$

$$SvO_2 = SaO_2 - VO_2 / Qt \times 1.36 \times Hb \times 10$$

Therefore, venous saturation (SvO_2) is directly related to arterial oxygen saturation, cardiac output and hemoglobin. However, SvO_2 also reflects tissue oxygen extraction and could be a more useful indicator of tissue oxygenation^[6].

However, measurement of mixed venous oxygen saturation in small newborn infants through a catheter in the pulmonary artery is hazardous and difficult. On the other hand, catheter placement into the right atrium through the umbilical vein via the ductus venosus is common practice in neonatal care and has a low incidence of complications^[7,8]. Therefore, we tested the feasibility of a fiberoptic catheter, placed in the right atrium via the umbilical vein, to measure central venous oxygen saturation ($ScvO_2$). Furthermore we determined the accuracy of the continuous oxygen saturation readings.

Patients and methods

This study was approved by the Ethics Committee of the Academic Hospital, Maastricht. The patient population of the feasibility study included 36 infants receiving care on the neonatal intensive care unit. Their clinical condition (Table 1) warranted insertion of an umbilical vein, according to our practice.

The fiberoptic catheter system (Oximetrix System, Abbott Laboratories, North Chicago, IL, USA) includes an optical module, a fiberoptic catheter and a microprocessor-

based oximeter. The light source in the optical module of the Oximetrix system consists of three diodes emitting light at three different wavelengths $\lambda_1=670$ nm, $\lambda_2=700$ nm and $\lambda_3=800$ nm.

A pulse generator sequentially activates each diode for 1 ms followed by an equal interval when no diodes are energized. This four period cycle occurs at a rate of 244 Hz [9].

The four french fiberoptic catheter consists of two lumens: one lumen includes the two fiberoptics to transmit the light to and from the blood; the other internal lumen leads from a distal end hole to a standard Luer-Lock hub. This lumen is used for administration of medication and fluids, blood sampling and blood pressure measurement. Performance is similar to that of a 3.5 French single lumen catheter.

Light reflected from the oxygenated and unoxygenated blood is transmitted by the receiving fiberoptic bundle to a photodetector in the optical module. The oximeter computes percentage oxygen saturation values based on the electrical signals from the optical module. The mean value for oxygen saturation over a 5 s period is displayed digitally, and this value is updated every 1 s.

The Oximetrix computer monitors continuously the optical intensity of the reflected light to warn for abnormal signal conditions. A change of $\pm 30\%$ in the mean received intensity signal gives a light intensity alarm message as it may indicate a significant change in light reflectance conditions at the catheter tip.

Before insertion the system was calibrated according to the manufacturer's instructions and methodology. This calibration procedure was performed to a disposable standard optical reference, packaged with each catheter. The proper location of the catheter was facilitated by the continuous display of the oxygen saturation value at the tip of the catheter during the insertion procedure. The catheter position was defined as correct when the catheter was positioned in the mid-atrial level or at the transition of the inferior vena cava and the right atrium; the position was evaluated by X-ray and ultrasound techniques. An in vivo calibration was performed after stabilization and establishment of the proper position of the catheter. This calibration procedure was performed at least every 12 hour to check for incorrect oxygen saturation readings. A correction was only performed only when the hemoximeter values of the oxygen saturation in correlated samples showed a discrepancy of $\geq 5\%$. A constant low rate (0.1 cc/h), heparinized saline (0.5 U/cc) infusion was administered continuously through the catheter to avoid fibrin formation in the fiberoptic catheter. After removal of the catheter the tip was inspected for fibrin clot formation.

The objective variables to assess the feasibility of the fiberoptic catheter system included: (a) catheter placement in the proper position; (b) total time the catheters were in place; (c) fibrin clot formation; and (d) incidence of complications.

To determine the accuracy of the oxygen saturation values given by the fiberoptic catheter system, blood samples (0.4 ml) were withdrawn through the catheter and immediately analyzed with a hemoximeter (Radiometer OSM3), which was calibrated routinely every hour.

Data are presented as mean $\pm$ SD. The unpaired Student's *t*-test or Yates' corrected chi-square test was used, where appropriate, to evaluate statistical difference between the groups of patients in which the catheter was properly placed. The in-vivo fiberoptic catheter values and the in-vitro hemoximeter values were analyzed by the least squares method, which yielded a regression line and correlation coefficient. Furthermore, data were analyzed comparing the difference against the mean of the two values [10]. The mean difference and SD were calculated and the change in the mean difference with time was estimated. The effect of recalibration was calculated and analyzed using the Student's *t*-test. Tests were considered to be significant for $p \leq 0.05$.

Results

In 24 patients (67%) the fiberoptic catheter was placed in the correct position, i.e. at the mid-atrial level or at the transition of the vena cava inferior and the right atrium. The characteristics and diagnostic categories of the 36 patients are summarized in table 1. There were no significant differences between the groups in which the catheter was placed successfully and those in which it was not placed.

Table 1: Birth weight (BW), gestational age (GA), gender, diagnosis and outcome in patients with successful compared with unsuccessful placement of the fiberoptic catheter system. Data are presented as mean $\pm$ SD or number. There were no significant differences between the two groups.

	Success (n = 24)	Failure (n = 12)
BW (g)	1484 $\pm$ 883	1747 $\pm$ 883
GA (weeks)	30.3 $\pm$ 4	31.6 $\pm$ 5
Sex (M/F)	14/10	6/6
Mortality	7	3
IRDS	16	7
Asphyxia	5	2
Lung hypoplasia	2	1
Streptococcus group B	1	2

The catheters were left in place for an average of 107 h (range 1 h 20 min to 172 h). In six infants, at the end of the indwelling period, the light intensity alarm of the fiberoptic catheter system indicated that the signal was affected by unusually small light reflectance at the cathetertip; at that time saturation readings were not considered to be reliable. On two occasions fibrin clot formation was seen by ultrasound in the right atrium around the tip of the catheter. The clots disappeared gradually without therapeutic intervention after removal of the catheter. Small fibrin clots at the tip of the catheter were seen in five infants (18%) at inspection after removal.

Complications such as pericardial effusions, cardiac tamponade, cardiac arrhythmias or perforations were not seen.

In order to establish the accuracy of the fiberoptic catheter system, 116 samples were withdrawn through the catheter. These measurements were performed 1-161 h after birth. The median time interval between the samples was 6 h (10th and a 90th percentiles of 2 and 19 h, respectively).

Correlation analysis of the 116 paired measurements showed a significant correlation ($r = 0.93$, $p < 0.001$), while regression analysis gave the following equation: $y = 0.96x + 2.96$ (Fig. 1). The statistical method for assessing agreement between two methods of measurement (10) showed a mean difference of -0.36 percentage points with 95% confidence intervals of +0.01 to -0.71. The limits of agreement (mean $\pm$ 2 SD) were +7.28 and -8.00 (Fig 2).

The first blood sample after pre-insertion calibration showed a comparable mean difference of 0.57%, with limits of agreement of +7.09 and -5.95 between the hemoximeter and fiberoptic catheter recorded readings.

Figure 3 shows the mean difference and SD between the fiberoptic catheter and the hemoximeter measurements with time. There was no significant effect with time during the study.

figure 1: Comparison of oxygen saturation values for the fibreroptic catheter with those obtained with the hemoximeter.
(linear regression)

In 30 of the 116 samples a discrepancy of $\geq 5\%$ or more was noted and only then was a correction performed. The actual degree to which the catheter values varied from the hemoximeter readings before they were recalibrated was mean 6.3% (SD 1.2%). The effect of recalibration was tested by ignoring the sample values taken after recalibration. The mean difference and SD (-0.54 ± 3.70) of the uncorrected remaining samples were compared with the mean difference of the values taken after recalibration (0.45 ± 4.42) and with the previously described mean difference and SD (-0.36 ± 3.82) of all 116 sample values. Using the unpaired Student's *t*-test a significant difference could not be demonstrated ($p = 0.29$ and 0.74 , respectively).

Figure 2: Difference between the fiberoptic catheter (SvO_2) and the hemoximeter (HbO_2) values against their means. The mean difference ($\pm 2SD$) was -0.36 ± 7.64 .

Discussion

This study reports the feasibility and accuracy of a fiberoptic catheter inserted via the umbilical vein into the right atrium for the measurement of venous oxygen saturation in newborn infants. This study, however, was not designed to describe critical values of $ScvO_2$ in newborn infants or its clinical applications in neonatal intensive care. Although umbilical venous catheterization is a long established and widely used technique in critically ill infants, two drawbacks of this procedure have been reported. Ideally, the tip of the umbilical venous catheter should be placed in the inferior vena cava near the right atrium. The success rate of passing the catheter through the ductus venosus is about 60%^[7,8,11,12]. In this study the fiberoptic catheter was placed successfully into the right atrium in 67% of infants. This of course could be improved by placing the catheter through another vein by a more invasive technique; however, since its clinical application has not been proved, we decided to place the catheter using the well established umbilical venous catheterization. The other problem is the incidence of fibrin clot formation, varying from small thrombi in the catheter to severe portal vein thrombosis^[13]. The reported incidence varies from 3 to 44%^[14]. In our study we found fibrin clot formation in seven (29%) infants.

Despite these drawbacks, umbilical venous catheterization is a common procedure because it is a rapid and simple technique^[15].

figure 3: Mean difference and SD between the fiberoptic catheter (SvO_2) and the hemoximeter (HbO_2) values with time (note that there was no drift with time).

The fiberoptic catheter did not differ in its usefulness from the standard umbilical catheters regarding administration of fluid, medication and monitoring central venous blood pressure. The catheters were left in place as long as was necessary for clinical management.

This study is the first report on the accuracy of venous oxygen saturation readings in the right atrium in a population of neonatal intensive care patients. The accuracy of the fiberoptic catheter has been demonstrated previously in the umbilical and pulmonary artery, for which the fiberoptic catheter originally was designed.

Previous studies on accuracy in adult intensive care medicine reported that SvO_2 , analyzed using the least squares method, was significantly correlated with hemoximeter values [16,17]. In our study the same correlation coefficient and regression equation were found as in those previous reports [16-18]. The correlation coefficient and regression analysis must be used with caution when it is used to compare two methods measuring the same parameter [10,19].

Another possibility and perhaps more appropriate is the use of a plot of the difference between the two values against their mean, with calculation of the mean difference and SD (see Fig. 2). The limits of agreement (mean $\pm$ 2SD) (+7.28 and -8.00) were small enough to consider the readings of venous oxygen saturation reliable. These limits of agreement were comparable with previous reports using this statistical method [20,21].

Because of the similar mean difference in the first blood sample, the oxygen saturation readings after preinsertion calibration can be considered to have been accurate and reliable and facilitated the proper location of the catheter during the insertion procedure. Spectrophotometric measurement of oxygen saturation may be biased by

changes in pH of blood^[22]. The direct effect of the pH on the absorption spectrum of the oxyhemoglobin has been described at pH 6.8 and 7.6 with dissimilar absorption peaks at 535 nm, 577 nm and 600 nm. No effect was observed on the absorption spectrum of deoxyhemoglobin. However, in the pH range (median 7.34: 5th and 95th percentile 7.19 and 7.47, respectively) we have encountered, measurement of oxygen saturation in the hemoximeter may reflect an inaccuracy of up to 1%, which is acceptable^[22]. Moreover, the fiberoptic catheter measures absorbance at wavelengths that are not influenced by these extreme pH values.

The multiwave length spectrophotometer Radiometer OSM (3) showed very high accuracy and precision in the measurement of oxygen saturation. A potential small inaccuracy due to the high sensitivity of the presence of fetal hemoglobin may occur^[23]. Although we did not measure fetal hemoglobin, we assumed that the fraction of fetal hemoglobin in the samples was variable due to the intensive care practice of giving blood transfusions after sampling a definite amount of blood.

The saturation values > 88% shown in Figs 1 and 2, were from samples taken before the catheter was withdrawn from the left atrium. In this study we described the correlation between values obtained from the catheter system and samples analyzed by the hemoximeter; therefore we decided to include these values in the study.

Other reports^[16,17] have shown a tendency for SvO₂ measurement to drift with time and become increasingly inaccurate. In our study no significant change in the difference and SD with time was found (Fig. 3).

Measurements in our study were performed in the right atrium in contrast with previous reports in adults where venous oxygen saturation was measured in the pulmonary artery. No evidence was found for a systematic difference between right atrial and pulmonary arterial oxygen saturations over a wide range of hemodynamic conditions both in an animal model and in adults^[24,25,26]. Other reports have described a poor correlation between right atrial and pulmonary arterial oxygen saturations when absolute numerical values were compared^[27]. However, a better correlation was observed when subsequent changes of the right atrial and pulmonary arterial saturations were compared^[27]. Therefore venous oxygen saturation monitoring in the right atrium is useful for following trends and changes in oxygenation status^[28]. Intracardiac left-to-right shunting through the foramen ovale is common in premature infants, but it is not responsible for differences in right atrial and pulmonary arterial saturations. However, variable shunting across the foramen ovale and/or incomplete mixing of blood in the right atrium can be a problem and is another argument for cautions with the use of exact numerical values of venous saturation.

This study demonstrated the feasibility of the use of a fiberoptic catheter for measurement of venous oxygen saturation in the right atrium with a low incidence of complications. The oxygen saturation readings using this system represented accurately and reliably blood oxygen saturation over a wide range.

In neonatal medicine, knowledge of real tissue oxygen needs is very important because hypoxia, as well as hyperoxia, have their deleterious effects. The current clinical parameters (paO₂ and SaO₂) represent only partially the availability of oxygen to the tissues, and disregard other parameters as blood flow, hemoglobin and tissue oxygen consumption. ScvO₂, however, reflects the residual oxygen after tissue oxygen extraction and represents the adequacy of all the above-mentioned parameters of the oxygen transport system.

ScvO₂ is measured after mixing of venous blood from all organs in the right heart and reflects global oxygen extraction. Therefore, normal ScvO₂ cannot exclude regional tissue ischaemia or hypoxia. Despite these known limitations ScvO₂ measurement in the right atrium could be useful as a complementary monitor of tissue oxygen needs in the

newborn infant.

The results of this study encouraged us to consider this fiberoptic catheter for measuring continuously the venous oxygen saturation in critically ill newborn infants. We are now testing for clinical relevance and describing the "safe ranges" for ScvO₂ in neonates requiring intensive care.

References

- 1 Kandel G, Aberman A. Mixed Venous Oxygen Saturation; its role in the assessment of the critically ill patient. *Arch Intern Med* 1983;143:1400-1402
- 2 Baele PL, McMichan JC, Marsh HM, Sill JC, Southorn PA. Continuous monitoring of mixed venous oxygen saturation in critically ill patients. *Anesth Analg* 1982;61:513-517
- 3 Divertie MB, McMichan JC. Continuous monitoring of mixed venous oxygen saturation. *Chest* 1984;85:423-428
- 4 Shenaq SA, Casar G, Chelly JE, Ott H, Crawford ES. Continuous monitoring of mixed venous oxygen saturation during aortic surgery. *Chest* 1987;92:796-799
- 5 Magilligan DJ, Taesdall R, Eisingminger R, Peterson E. Mixed venous oxygen saturation as a predictor of cardiac output in the postoperative cardiac surgical patient. *Ann Thorac Surg* 1987;44:260-262
- 6 Dudell G, Cornisch JD, Bartlett RH. What constitutes adequate oxygenation? *Pediatrics* 1990;85:39-41
- 7 Kitterman JA, Phibbs RH, Tooley WH. Catheterization of umbilical vessels in newborn infants. *Ped Clin of North America* 1970;17:895-912
- 8 Ewing CK, Durand DJ. Use of umbilical catheters: A study of 100 patients. *Pediatr Res* 1989;25:214A
- 9 Brown EG, Krouskop RW, McDonnel FE, Monge CC, Winslow RE. A technique to continuously measure arteriovenous oxygen content difference and p₅₀ in vivo. *J Appl Physiol* 1985;58(4):1383-1389
- 10 Bland JM, Altman DG. Statistical methods for assessing agreement between two methods of clinical measurement. *Lancet* 1986;i:307-317
- 11 Baker DH, Berdon MD, James LS. Proper localization of umbilical arterial and venous catheters by lateral roentgenogram. *Pediatrics* 1969;43:34-39
- 12 Rosen MS, Reich SB. Umbilical venous catheterization in the newborn: Identification of correct positioning. *Radiology* 1970;95:335-340
- 13 Oski FA, Allen DM, Diamond LK. Portal hypertension; a complication of umbilical vein catheterization. *Pediatrics* 1963;39:297-302
- 14 Wigger HJ, Bransilver BR, Blanc WA. Thromboses due to catheterization in infants and children. *J Pediatr* 1970;76:1-12
- 15 Dorand RD, Cook LN, Andrews BF. Umbilical vessel catheterization; The low incidence of complications in a series of 200 newborn infants. *Clin Pediatr* 1977;16:569-572
- 16 Reinhart K, Moser N, Rudolph T, Bredle D, Specht M, Gramm H-J, et al. Accuracy of two mixed venous saturation catheters during long term use in critically ill patients. *Anesthesiology* 1988;69:769-773
- 17 Gettinger A, De Traglia MC, Glass D. In vivo comparison of two mixed venous saturation catheters. *Anesthesiology* 1987;66:373-375
- 18 Vaughn S, Puri VK. Cardiac output changes and continuous mixed venous oxygen saturation measurement in the critically ill. *Crit Care Med* 1988;16:495-498
- 19 Healy MJR. Statistics from the inside; Regression and correlation. *Arch Dis Child* 1992;67:1306-1309

- 20 Scuderi PE, Bowton DL, Meredith JW, Harris LC, Brockschmidt J. Clinical evaluation of three mixed-venous oxygen saturation catheters. *Anesthesiology* 1990;73:A484
- 21 Boeden G, Spies C, Zaune U, Gabriel H, Martin E. Longterm-accuracy of continuous fiberoptic SvO₂ monitoring. *Anesthesiology* 1990;73:A495
- 22 Wimberley PD, Fogh-Andersen N, Siggaard-Andersen O, Lundsgaard FC, Zijlstra WG. Effect of pH on the absorption spectrum of human oxyhemoglobin: a potential source of error on the measuring the oxygen saturation of hemoglobin. *Clin Chem* 1988;34:750-754
- 23 Wimberley PD, Helledie NR, Friis-Hansen B, Siggaard-Andersen O, Fogh-Andersen N. Some problems involved in using hemoglobin oxygen saturation in arterial blood to detect hypoxemia and hyperoxemia in newborn infants. *Scand J Clin Lab Invest* 1988; 48, Suppl 189: 45-48
- 24 Barratt-Boyes BG, Wood EH. The oxygen saturation of blood in the venae cavae, right-heart chambers and pulmonary vessels of healthy subjects. *J Lab Clin Med* 1957;50:93-106
- 25 Reinhart K, Rudolph T, Bredle DL, Hannemann L, Cain SM. Comparison of central-venous to mixed-venous oxygen saturation during changes in oxygen supply/demand. *Chest* 1989;95:1216-1221
- 26 Davies GG, Mendenhall J, Symreng T. Measurement of right atrial oxygen saturation by fiberoptic oximetry accurately reflects mixed venous oxygen saturation in swine. *J Clin Monit* 1988;4:99-102
- 27 Auffray JP, Martin C, Prima F, et al. Venous oxygen saturation: Central venous (CV) or Pulmonary artery (PA) catheter? *Anesthesiology* 1991;75:A455
- 28 Tahvanainen J, Meretoja O, Nikki P. Can central venous blood replace mixed venous blood samples? *Crit Care Med* 1982;10:758-761

Acknowledgments

This study was supported by Abbott Inc., the Netherlands.

Chapter 6

Continuous Central Venous Oxygen Saturation (ScvO₂) measurement by a fiberoptic catheter in newborn infants.

Mark A.H.B.M. van der Hoeven, Wiel J. Maertzdorf, Carlos E. Blanco.

Archives of Diseases in Childhood; Fetal and Neonatal Edition 1996;74:F177-F188

Abstract

Background: Central venous oxygen saturation ($ScvO_2$) can be a valuable monitor, reflecting residual oxygen after tissue oxygen extraction and indicating the combined sufficiency of tissue oxygen supply and demand.

Objective: a) To describe the range of $ScvO_2$ values in stable newborn infants breathing room air. b) To examine the relationship between $ScvO_2$ and arterial oxygen saturation (SaO_2) c) to describe fractional oxygen extraction (F.O.E.) and d) the shunt index, an estimate of the venous admixture.

Study Design: a prospective clinical observational study

Patients: 10 preterm infants breathing room air after the acute phase of respiratory distress syndrome and with an umbilical venous catheter in situ. A fiberoptic catheter remained in the right atrium for continuous measurement of oxygen saturation.

Measurements and Main Results: $ScvO_2$, SaO_2 , bloodpressure and heart rate were registered every 15 minutes. F.O.E. and shunt index were calculated. SaO_2 and $ScvO_2$ were $93.4\% \pm 3.7$ and $73.56\% \pm 5.25$ (mean $\pm$ SD.), respectively. In seven patients $ScvO_2$ values correlated significantly with SaO_2 . F.O.E. was 0.21 ± 0.04 (mean $\pm$ SD.) and was significantly correlated with $ScvO_2$. The shunt index was $24\% \pm 12$ (mean $\pm$ SD) and was significantly correlated with SaO_2 .

Conclusions: Stable preterm infants, breathing room air had a $ScvO_2$ range from 65% to 82% (5th and 95th percentile), which corresponded to $SaO_2 \geq 86\%$. $ScvO_2$ values were significantly correlated with SaO_2 in most patients.

Introduction

The optimal level of oxygenation in sick newborn infants is not well established. Hyperoxia has been associated with oxygen toxicity related cell injury [1], retinopathy of prematurity [2] and bronchopulmonary dysplasia [3], while hypoxia has been associated with vasoconstriction of the pulmonary vascular bed [4,5], redistribution of the cardiac output [6] and oxygen insufficiency at tissue level with anaerobic metabolism leading to cell damage.

Arterial partial pressure of oxygen (paO₂) and/or arterial oxygen saturation (SaO₂) are widely used in judging oxygen needs. These are limited because they do not measure other factors that determine tissue oxygenation such as hemoglobin concentration, cardiac output, local tissue blood flow and oxygen consumption.

Central venous oxygen saturation (ScvO₂) reflects residual oxygen after tissue oxygen extraction and accounts for all the mentioned variables [7]. Measuring the ScvO₂ in newborn babies may help minimize unnecessary oxygen administration, and thus decreasing long term morbidity associated with oxygen injury [8]. ScvO₂ measurement could define insufficient oxygen delivery to tissues when arterial oxygen content and cardiac output are decreased or when oxygen consumption is increased.

In this study we attempted to define the normal range of ScvO₂ values in stable newborn infants, breathing room air after a period of respiratory insufficiency.

The measurement of both SaO₂ and ScvO₂ (dual oximetry) made it possible to calculate fractional oxygen extraction (F.O.E.) and the shunt index [9]. F.O.E. is an estimation of the peripheral tissue oxygen extraction coefficient. As oxygen delivery falls, oxygen consumption can be maintained by an increase in the extraction of oxygen delivered to the tissues [10]. Shunt index represents an estimate of the venous admixture. Pulmonary venous admixture reflects the degree of mixture of arterial blood and mixed venous blood. We examined the range of these two variables and their relation with SaO₂ and ScvO₂.

Materials and Methods

This study was approved by the ethics committee of the Academic Hospital of Maastricht.

We used a 4 Fr. fiberoptic catheter (Oximetrix System, Abbott Lab. Chicago Ill.), which allowed both a continuous measurement of oxygen saturation at the tip of the catheter and infusion of fluids and medication. The continuous measurement of oxygen saturation is based on reflection spectrometry. The light source, consisting of three diodes, emits light at three different wavelengths through a fiberoptic bundle. The back scattered light from the oxygenated and unoxygenated blood is transmitted by the receiving fiberoptic bundle to a photodetector in the optical module. The oximeter computes percent of oxygen saturation values based on the electrical signals from the optical module. The average value for oxygen saturation over a five second period is displayed digitally, and the value updated each second.

The fiberoptic catheter was placed in the right atrium via the umbilical vein in infants who needed a central venous line for clinical management. The catheter was positioned at the midatrial level or at the transition of the inferior vena cava to the right atrium. The position was confirmed by X-ray and ultrasound. Heparinized saline (0.5 U./cc.) was infused continuously (0.1 cc/hr.) to avoid fibrin formation over the catheter. Bloodsamples were withdrawn through the catheter at least every twelve hours, immediately analyzed with a hemoximeter (Radiometer OSM3, Copenhagen) and compared

with simultaneously recorded fiberoptic catheter values. A correction was made when oxygen saturation in correlated samples showed a discrepancy of $\geq 5\%$. We have previously reported the feasibility and accuracy of the fiberoptic device in newborn infants [11].

The values of ScvO_2 were obtained from preterm patients breathing room air after the acute phase of RDS. Infants were allowed to breathe room air when SaO_2 monitored by pulse oximetry was $\geq 86\%$ [12,13]. ScvO_2 values were accepted when hemoglobin (Hb) was ≥ 7.0 mmol/L, with normal arterial pH (7.30-7.45) and pCO_2 (4.5-8 kPa), and arterial bloodpressure [14].

SaO_2 was continuously measured by pulse oximetry (Hewlett Packett). Each value is the average of samples taken every 12 seconds over one minute. Arterial blood pressure was monitored through an umbilical or radial artery catheter; these catheters were also used for sampling arterial pH, PaCO_2 , paO_2 , and base excess for clinical purposes.

ScvO_2 , SaO_2 values, bloodpressure and heart rate were registered every 15 minutes. Arterial oxygen content (caO_2) and central venous oxygen content (ccvO_2) were calculated using the equations:

$$- \text{caO}_2 \text{ (ml/dl)} = \text{SaO}_2 \times \text{Hb (mmol/L)} \times 1.61 \times 1.36$$

and

$$- \text{ccvO}_2 \text{ (ml/dl)} = \text{ScvO}_2 \times \text{Hb (mmol/L)} \times 1.61 \times 1.36, \text{ respectively.}$$

Fractional oxygen extraction (F.O.E.) was calculated using the equation

$$- \text{F.O.E.} = (\text{SaO}_2 - \text{ScvO}_2) / \text{SaO}_2. \text{ The shunt index was calculated using the equation:}$$

$$- \text{shunt index} = 100 \times \{(100 - \text{SaO}_2) / (100 - \text{ScvO}_2)\}.$$

Oxygen saturation and pcvO_2 were determined from samples withdrawn from the central venous fiberoptic catheter and the p_{50} was calculated as described previously [15].

Statistics

The paired Student's t test was used to compare the hemoximeter values with the simultaneously recorded fiberoptic catheter readings. For each patient we determined mean value and standard deviation (mean $\pm$ SD) for SaO_2 and ScvO_2 . The range was defined by the 5th and 95th percentile. To compare SaO_2 with ScvO_2 , a separate linear regression analysis within each individual was performed to avoid erroneous combining of observations from different individuals [16]. A slope (the regression coefficient) for each individual was calculated to obtain an average slope for the group [16]. To compare F.O.E and shunt index with SaO_2 and ScvO_2 the same statistical analysis was performed. Tests were considered to be significant when $p < 0.05$.

Results

In 10 preterm infants breathing room air, ScvO_2 could be monitored continuously. They are described in table 1. The monitoring time ranged from 10-82 hrs. The mean age of the infants when ScvO_2 was monitored was 52 hrs. (range 18-168 hrs.). A total of 1061 ScvO_2 and SaO_2 values were obtained simultaneously with bloodpressure and heart rate. The validity of the ScvO_2 values was confirmed, since hemoximeter values and simultaneously recorded fiberoptic catheter readings were not significantly different: (74.7 % $\pm$ 6.8 vs. 73.8 % $\pm$ 6.3). SaO_2 and ScvO_2 were 93.4 % $\pm$ 3.7 and 73.56 % $\pm$ 5.25 (mean $\pm$ SD.), respectively. The SaO_2 and ScvO_2 values ranged respectively from 87 to 99 % and 65 % to 82 % (5th and 95th percentile). When analyzed for each patient separately, ScvO_2 (lowest 5th and highest 95th percentile) ranged from 60.1 % to 83.5 % (table 2).

table 1. Patient characteristics, physiological parameters, arterial pH, paCO₂, paO₂ and base excess and oxygen saturation, calculated arterial and venous oxygen content.

	Mean	(SD)	Range (5th..95th perc)	
birthweight (grams)	938	(246)		
gestational age (weeks)	27.0	(1.3)		
hemoglobin (mmol/L)	8.4	(1.8)	7.0	10.2
MAP (mm Hg.)	41.5	(7.4)	30	54.7
heart rate (bpm)	165	(15)	144	192
p ₅₀ (kPa)	3.1	(0.4)	2.7	3.9
pH	7.36	(0.04)	7.30	7.41
pCO ₂	5.5	(1.0)	4.4	7.8
pO ₂	6.6	(0.1)	3.5	10.2
Base Excess (mmol/L)	-2.8	(2.3)	-7.3	0.1
caO ₂ (ml/dl)	16.9	(2.2)	12.6	20.0
cvO ₂ (ml/dl)	13.2	(2.0)	9.6	16.1

table 2. Mean $\pm$ SD, range (5th and 95th percentile), correlation coefficient and slope of SaO₂ and ScvO₂.
*: $p < 0.05$

Case No	Samples	SaO ₂		ScvO ₂			Correlation	
		Mean (SD)	Percentile 5th 95th	Mean (SD)	Percentile 5th 95th		Correlation coefficient	Slope
1	59	91.1 (2.1)	87 99	70.5 (4.4)	61 78		0.34*	0.71*
2	67	93.1 (1.8)	90 96	71.3 (3.4)	65 77		0.05	0.10
3	57	90.2 (2.3)	87 94	70.6 (4.4)	63 77		0.65*	1.2 *
4	56	92.1 (3.5)	86 98	70.5 (4.3)	62 77		0.25	0.30
5	40	93.2 (3.0)	87 97	69.9 (4.6)	60 76		0.60*	0.92*
6	45	90.5 (3.2)	86 98	70.6 (5.0)	62 78		0.18	0.28
7	81	91.4 (2.2)	88 95	72.6 (4.7)	64 80		0.73*	1.6 *
8	184	93.3 (3.3)	87 99	72.5 (4.9)	65 81		0.35*	0.78*
9	144	89.7 (1.9)	86 93	72.2 (2.2)	69 75		0.69*	0.69*
10	328	97.2 (2.0)	93 100	77.9 (4.5)	68 83		0.57*	1.33*

Linear regression analysis comparing SaO₂ and ScvO₂ for each patient yielded a slope (regression coefficient) and a correlation coefficient r . (table 2, figure 1). In seven patients a significant correlation was found (see figure 1). F.O.E was 0.21 ± 0.04 (mean $\pm$ SD.). The 5th and 95th percentile were 0.15 and 0.29, respectively. When analyzed for each patient separately, a range was found (lowest 5th and highest 95th percentile) from 0.13 to 0.33. Figure 2 shows the relation between F.O.E. and ScvO₂. F.O.E. and ScvO₂ were significantly correlated in all patients. SaO₂ and F.O.E. were significantly correlated in only 6 patients with a lower range of correlation coefficients (range r : -0.28..-0.48). Analysis of the relation between F.O.E and Hb yielded a correlation coefficient $r = -0.14$. ($p < 0.01$).

The shunt index was $24\% \pm 12$ (mean $\pm$ SD). A range was found (lowest 5th and highest 95th percentile) from 7% to 53% (table 2). When analyzed for each patient separately, we found a significant negative correlation with the SaO₂ value in all patients (range r : -0.72..-0.95; see figure 3). Analysis of the relation with ScvO₂ showed a significant correlation in five patients (range r : -0.37..0.36).

figure 1. Central venous oxygen saturation ($ScvO_2$) values corresponding to simultaneously obtained arterial oxygen saturation (SaO_2) values. Linear regression lines are drawn for each patient. In seven patients a significant correlation was found. An average regression equation was obtained: $ScvO_2 = 0.46 + 0.77 \times SaO_2$.

Discussion

Continuous monitoring of mixed venous oxygen saturation ($ScvO_2$) has been used in critically ill adult patients as a valuable alarm signal, reflecting the residual oxygen after tissue oxygen extraction and indicating the combined sufficiency of the tissue oxygen supply and demand [17,18,19]. Venous oxygen saturation is used very frequently during extracorporeal life support to determine pump blood flow [20] and has been advocated in the care for sick preterm infants [8]. In neonatal medicine the use of $ScvO_2$ could allow to minimize unnecessary oxygen administration and hence to decrease long term morbidity associated with oxygen toxicity related injury [8]. However, normal and safe values have not been defined for newborn infants.

In our study we observed a range from 60.1 % to 83.5 % (5th and 95th percentile) in 10 stable preterm infants breathing room air, while SaO_2 was $\geq 86\%$, hemoglobin was ≥ 7.0 mmol/L. Arterial pH and arterial blood pressure were in normal range. This $ScvO_2$ range is similar to the normal range defined for adults; however, oxygen consumption in newborn infants is 5-8 ml/kg/min, whereas in resting adults oxygen consumption is 3-5 ml/kg/min [20]. Increased oxygen consumption is not compensated for by increased extraction within tissues, since $ScvO_2$ does not apparently differ from adult values.

figure 2. Fractional Oxygen Extraction (F.O.E.) corresponding to central venous oxygen saturation (ScvO₂). Linear regression lines are drawn for each patient. F.O.E. was significantly correlated with the ScvO₂ in each patient. An average regression equation was obtained: $F.O.E. = 0.83 - 0.0085 \times ScvO_2$

The oxygen delivery is determined by cardiac output and caO₂, which in turn is determined by hemoglobin concentration (Hb), O₂ carrying capacity, oxygen saturation (SaO₂) and paO₂. Cardiac output and hemoglobin concentration in the newborn compensates for higher oxygen consumption. Indeed, hemoglobin concentration and caO₂ are higher in newborn babies than in adults [21]. Cardiac output in newborn infants determined by various methods is 230 ± 70 ml/kg/min [22], whereas cardiac output in adults is 70-90 ml/kg/min [23]. Therefore, the increased oxygen consumption in the newborn infant is counterbalanced by an increased oxygen delivery.

In seven patients a significant correlation was found between SaO₂ and ScvO₂. This indicates that, when SaO₂ is reduced, oxygen consumption is preserved by a fixed oxygen extraction. In three patients SaO₂ and ScvO₂ were not significantly correlated. We might speculate, that in these babies oxygen consumption was preserved by increased cardiac output. It is unlikely that oxygen consumption was decreased by SaO₂ reduction, because in stable patients with a normal blood pressure, normal hemoglobin level and normal oxygen saturation oxygen consumption is independent of oxygen delivery [24]. Oxygen delivery is normally four to five times oxygen consumption as shown the normal fractional oxygen extraction (F.O.E.) of 0.20 - 0.25. In adults, a normal range of 0.22 to 0.30 of the F.O.E. has been described [25]. An increased F.O.E. can partly compensate for imbalance of oxygen consumption and delivery. Increased oxygen consumption, low hemoglobin and a low cardiac output could explain the high F.O.E. found in our study.

Figure 3. The shunt index, an estimate for the venous admixture corresponding to the arterial oxygen saturation (SaO_2). Linear regression lines are drawn for each patient. Shunt index and SaO_2 were correlated in each patient. An average regression equation was obtained: shunt index = $2.85 - 0.0278 \times \text{SaO}_2$

Indeed, we found a significant negative correlation between Hb and F.O.E. The highly significant correlation between F.O.E. and ScvO_2 in all patients confirms that ScvO_2 is an excellent monitor of the balance between oxygen delivery and tissue oxygen consumption.

The shunt index is an estimate of venous admixture and can be calculated when both SaO_2 and ScvO_2 (dual oximetry) are available. It is a derived formula, $(100 \times (100 - \text{SaO}_2) / (100 - \text{ScvO}_2))$. Dissolved oxygen is discounted, and it is assumed that pulmonary end capillary blood is fully saturated.

Bongard et al. ^[26] demonstrated a linear relation between this index and venous admixture; when paO_2 was low, it was largely accurate but with some overestimation of the value. Concomitantly with F.O.E. it has been used to titrate continuous positive airway pressure in adult intensive care ^[27]. In our patients the shunt index was significantly correlated with SaO_2 with values up to 50% in the lower SaO_2 range (fig. 3). Recently, Schultze et al. ^[28] described approximately similar values of venous admixture of $31\% \pm 9$ at a SaO_2 range between 89% and 92% in premature infants. Obviously an increased venous admixture is present when lower SaO_2 values are encountered. As can be expected and was demonstrated in a computermodel ^[26], changes in SaO_2 alter the value of the shunt index to a greater extent than changes in ScvO_2 .

The measurement site of venous oxygen saturation is still in debate^[29]. In adult intensive care medicine venous oxygen saturation has been monitored continuously in the pulmonary artery. Measurement of the ScvO₂ in the right atrium instead of in the pulmonary artery has been reported^[30,31]. No evidence was found for a systematic difference between right atrial and pulmonary arterial oxygen saturations over a wide range of hemodynamic conditions both in the animal model and in the adult human^[30,31,32]. Other reports describe a poor correlation between right atrial and pulmonary arterial oxygen saturation when comparing absolute numerical values^[33]. However, a better correlation was observed comparing subsequent changes of right atrial and pulmonary arterial saturations^[34].

In neonatal medicine the use of a pulmonary artery catheter is difficult and hazardous, but umbilical venous catheters have been used frequently with a low incidence of complications. Hence, measurements of venous oxygen saturation in newborn infants are limited to the right atrium. Intracardiac left-to-right shunting through the foramen ovale is common in preterm infants, and can be a reason for the higher ScvO₂ values found in preterm infants compared with adults. However, intracardiac shunting at the right atrium does not produce a difference in right atrial and pulmonary arterial oxygen saturations, and, therefore, is not an argument for questioning the validity of the right atrium as measurement site.

ScvO₂ measurement has some drawbacks: since it reflects global oxygen extraction, normal values cannot exclude tissue hypoxia in individual organs^[35]. In sepsis and multi-organ failure normal ScvO₂ values may occur despite global hypoxia due to precapillary shunting and the incapacity of tissues to extract enough oxygen^[36]. These drawbacks do not imply, that measuring ScvO₂ is not useful, but that normal and abnormal values must be interpreted together with other physiologic variables in conditions such as perfusion disturbances and sepsis^[8].

In summary, we have defined a range of ScvO₂ values in preterm newborn infants when breathing room air with a SaO₂ ≥ 86%. Whether these values can be used in sick newborn infants to evaluate tissue oxygenation and whether this range can be used to regulate oxygen administration remains to be established. Further study is needed to determine the clinical utility of ScvO₂ measurements in neonatal intensive care.

References

- 1 Frank L, Bucher JR, Roberts RJ. Oxygen toxicity in neonatal and adult animals of different species. *J Appl Physiol* 1978;45:699-704
- 2 Flynn JT, Bancalari E, Snyder EM, Goldberg RN, Feuer W, Cassady J. A cohort study of transcutaneous oxygen tension and the incidence and severity of retinopathy of prematurity. *N Engl J Med* 1992;326:1050-1054
- 3 Goetzman BW. Understanding bronchopulmonary dysplasia. *AJDC* 1986;140: 332-334
- 4 Fishman AP. Respiratory gases in the regulation of the pulmonary circulation. *Physiol Rev* 1961;41:214-280
- 5 Huage A. Hypoxia and pulmonary resistance. The relative effects of pulmonary arterial and alveolar pO₂. *Acta Physiol Scand* 1969; 76: 121-130
- 6 Sheldon R, Peeters LLH, Jones Jr. MD, Makowski EL, Meschia G. Redistribution of cardiac output and oxygen delivery in the hypoxemic fetal lamb. *Am J Obstet Gynecol* 1979;135:1071-1078
- 7 O'Connor TA, Hall RT. Mixed venous oxygenation in critically ill neonates. *Crit Care Med* 1994;22:343-346

- 8 Dudell G, Cornisch JD, Bartlett RH. What constitutes adequate oxygenation? *Pediatrics* 1990;85:39-41
- 9 Räsänen J. Role of dual oximetry in the assessment of pulmonary function. In: Reinhart K, Eyrich K. Eds. *Clinical aspects of O₂ transport and tissue oxygenation*. Berlin: Springer Verlag, 1989; pp 193-211
- 10 Little RA, Edwards JD. Applied physiology. In: Edwards JD, Shoemaker WC, Vincent JL. *Oxygen transport. principles and practice*. Philadelphia: Saunders, 1993; pp 21-40
- 11 van der Hoeven MA, Maertzdorf WJ, Blanco CE. Feasibility and accuracy of a fiberoptic catheter for the measurement of venous oxygen saturation in newborn infants. *Acta Paed* 1995;84:122-127
- 12 Wasunna A, Whitelaw AG. Pulse oximetry in preterm infants. *Arch Dis Child* 1987;62:882-888
- 13 Joint working group of the British Association of Perinatal Medicine and the Research Unit of the Royal College of Physicians. Development of audit measures and guidelines for good practice in the management of neonatal respiratory distress syndrome. *Arch Dis Child* 1992;67:1221-1227
- 14 Shortland DB, Evans DH, Levene MI. Blood pressure measurements in very low birth weight over the first week of live. *J Perinat Med* 1988;16:93-97
- 15 Brown EG, Krouskop RW, McDonnell FE, Monge CC, Winslow RE. A technique to continuously measure arteriovenous oxygen content difference and p_{50} in vivo. *J Appl Physiol* 1985;58(4):1383-1389
- 16 Smith EO. Analysis of repeated measures designs. *J Pediatr* 1987;111:723-725
- 17 Kandel G, Aberman A. Mixed Venous Oxygen Saturation; Its role in the assessment of the critically ill patient. *Arch Intern Med* 1983;143:1400-1402
- 18 Baele PL, McMichan JC, Marsh HM, Sill JC, Southorn PA. Continuous monitoring of mixed venous oxygen saturation in critically ill patients. *Anesth Analg* 1982;61:513-517
- 19 Divertie MB, McMichan JC. Continuous monitoring of mixed venous oxygen saturation. *Chest* 1984;85:423-428
- 20 Bartlett RH, Cilley RE. Physiology of extracorporeal life support. In: Arensman RM, Cornish JD. editors. *Extracorporeal life support*. Boston: Blackwell, 1993; pp 89-102
- 21 Delivoria-Papadopolous M, Wagerle LC. Oxygen diffusion and transport. In: Scarpelli EM. editor. *Pulmonary Physiology*. Philadelphia: Lea & Febiger, 1990; pp 281-305
- 22 Hudson I, Houston A, Aitchison T, Holland B, Turner T. Reproducibility of measurements of cardiac output in newborn infants by doppler ultrasound. *Arch Dis Child* 1990;65:15-19
- 23 Guyton AC. Cardiac output, venous return, and their regulation. In: *Textbook of medical physiology*. Philadelphia: Saunders, 1986; pp 272-284
- 24 Sear M, Wensley D, MacNab A. Oxygen consumption-oxygen delivery relationship in children. *J Pediatr* 1993;123:208-214
- 25 Shoemaker WC. Haemodynamic and oxygen transport in shock: Pathophysiology, monitoring, outcome prediction and therapy. In: Edwards JD, Shoemaker WC, Vincent JL. eds. *Oxygen transport. principles and practice*. Philadelphia, Saunders, 1993; pp 70-98
- 26 Bongard FS, Leighton TA. Continuous dual oximetry in surgical critical care. indications and limitations. *Ann Surg* 1992;216:60-68
- 27 Räsänen J, Downs JB, Malec DJ, Seidman P. Estimation of oxygen utilization by dual oximetry. *Ann Surg* 1987;206:621-623
- 28 Schultze A, Whyte RK, Clifton Way R, Sinclair JC. Effect of the arterial oxygenation level on cardiac output, oxygen extraction and oxygen consumption in low birth weight infants receiving mechanical ventilation. *J Pediatr* 1995;126:777-784
- 29 Vincent JL. Does central venous oxygen saturation accurately reflect mixed venous oxygen saturation? Nothing is simple, unfortunately. *Int Care Med* 1992;18:386-387
- 30 Reinhart K, Rudolph T, Bredle DL, Hannemann L, Cain SM. Comparison of central-venous to mixed-venous oxygen saturation during changes in oxygen supply/demand. *Chest* 1989;95:1216-1221

- 31 Davies GG, Mendenhall J, Symreng T. Measurement of right atrial oxygen saturation by fiberoptic oximetry accurately reflects mixed venous oxygen saturation in swine. *J Clin Monit* 1988;4:99-102
- 32 Barratt-Boyes BG, Wood EH. The oxygen saturation of blood in the venae cavae, right-heart chambers and pulmonary vessels of healthy subjects. *J Lab Clin Med* 1957;50:93-106
- 33 Martin C, Auffray JP, Badetti C, Perrin G, Papazian L, Gouin F. Monitoring of central venous oxygen saturation versus mixed venous oxygen saturation in critically ill patients. *Int Care Med* 1992;18:101-104
- 34 Tahvanainen J, Meretoja O, Nikki P. Can central venous blood replace mixed venous blood samples? *Crit Care Med* 1982;10:758-761
- 35 Reinhart K. Monitoring O_2 transport and tissue oxygenation in critically ill patients. In: Reinhart K, Eyrich K. Editors. *Clinical aspects of O_2 transport and tissue oxygenation*. Berlin: Springer Verlag, 1989; pp 193-211
- 36 Shah DM, Weil MH, Gazmuri RJ, Rackow EC. O_2 transport in low flow states of circulatory shock and cardiac arrest. In: Reinhart K, Eyrich K. Editors. *Clinical aspects of O_2 transport and tissue oxygenation*. Berlin: Springer Verlag, 1989; pp 445-461

Acknowledgements

This study was supported by Abbott Inc. the Netherlands.

Chapter 7

Central venous oxygen saturation (ScvO₂) and derived variables at different ranges of arterial oxygenation in newborn infants with respiratory insufficiency.

Mark A. van der Hoeven, Wiel J. Maertzdorf, Carlos E. Blanco.

Submitted for publication

Abstract

Aims: To observe ScvO₂ values at different SaO₂ values < 86 %, at the lower end (86 - 90 %) and higher (91 - 96 %) end of the recommended range of SaO₂ values measured by pulse oximetry in newborn infants with respiratory insufficiency. Other variables as arterial oxygen content (caO₂) and venous oxygen content (cvO₂), fractional oxygen extraction and the shunt index, an estimate of the venous admixture were calculated.

Design: A non-randomized clinical observational study.

Setting: Level III neonatal intensive care unit of a university hospital.

Patients: 12 (pre)term newborn infants with mechanical ventilation and oxygen need.

Birthweight: 1130 gm (830 , 1355); **Gestational age:** 28 wks (26.4 , 30).

Measurements and main results: On line continuous ScvO₂ measurement in the right atrium by an umbilical venous fiberoptic catheter. ScvO₂, SaO₂ and FiO₂ were registered every 15 minutes. F.O.E. and shunt index were calculated. ScvO₂ was significantly reduced in the SaO₂ range of 86 - 90 % in nine of the twelve patients. Linear regression analysis yielded a significant correlation between SaO₂ and ScvO₂ in all but one of the patients. The slope of the regression line between the two levels of oxygenation were not different in any of the infants. ScvO₂ values in SaO₂ values < 86 % were significantly reduced compared with the values between 86 % and 96 %. The shunt index, an estimate of the venous admixture, increased significantly in the lower level of oxygenation in all the patients. In four of the five patients with SaO₂ values < 86 % the slope of the regression line was significantly reduced compared to that obtained at SaO₂ values > 86 %.

Conclusions: On line measurement of ScvO₂ at different levels of SaO₂ values showed a significant reduction of ScvO₂. This reduction of ScvO₂ seems to be the primary compensatory mechanism to stabilize tissue oxygenation.

Introduction

The optimal level of oxygenation in sick newborn infants is not well established. The currently measured parameters of oxygenation include arterial oxygen tension (paO₂) and arterial oxygen saturation (SaO₂).

Attempts have been made to define the limits of the normal range of paO₂ and SaO₂. For this, values of paO₂ and SaO₂ were correlated to other variables.

PaO₂ levels between 2.5 to 4 kPa (SaO₂ between 46 to 82 %) in newborn infants breathing 13 % oxygen were associated with a rise in pulmonary arterial pressure and resistance [1]. Additionally, it was found that visual evoked responses in preterm infants with idiopathic respiratory distress syndrome (IRDS) were more affected, when PaO₂ decreased below 7.5 kPa [2]. On the other hand Émond et al. [3] demonstrated that approximately 90 % of the hemoglobin of preterm newborn infants is bound to oxygen (p₅₀) at a level of paO₂ of 5.5 ± 0.5 kPa. Furthermore, it was shown, that decreasing SaO₂ from 95 % to 90 %, corresponding to paO₂ values decreasing from 13 kPa to 6.6 kPa, did not have any effect on pulmonary hemodynamics, ductus arteriosus, oxygen consumption and cardiac output [4,5]. These data have led to different recommendations [6,7] and debate [8,9].

The current clinical parameters of oxygenation, PaO₂ and SaO₂, are rather poor indicators of tissue oxygenation, because they represent only the availability of oxygen to the tissues. Tissue oxygenation is also determined by other variables as hemoglobin, cardiac output and oxygen uptake of the tissues. Venous oxygen saturation reflects the residual oxygen after tissue oxygen extraction and indicates the combined sufficiency of all these variables [10,11,12]. In a previous study we have determined the feasibility and the accuracy of a fiberoptic catheter system in the newborn, providing a continuous central venous oxygen saturation (ScvO₂) measurement in the right atrium [13]. We have also described a "normal" range of ScvO₂ values between 58 % to 85 % in 1-7 days old (pre)term infants breathing room air [14].

The aim of the present work was to describe the ScvO₂ values at three levels of SaO₂ values: below 86 %, between 86 to 90 % and between 91 to 96 %; these last two values are the lower and the higher end of the recommended range of SaO₂ values in newborn infants with respiratory insufficiency, respectively. Other oxygen transport variables, as arterial (caO₂) and venous oxygen content (cvO₂), fractional oxygen extraction, an estimate of the venous admixture: the shunt index, p₅₀ and p₉₀ are described.

Patients and methods

This study was approved by the ethics committee of the Academic Hospital of Maastricht.

Patients included 12 newborn infants with respiratory insufficiency, oxygen need and endotracheal mechanical ventilation. Umbilical catheters were placed for clinical management.

measurements

ScvO₂ was continuously measured by a 4 Fr. fiberoptic catheter (Oximetrix System, Abbott Lab. Chicago Ill.), which allowed both a continuous measurement of oxygen saturation at the tip of the catheter and infusion of fluids and medication. The continuous measurement of oxygen saturation is based on reflection spectrometry. The light source, consisting of three diodes, emits light at three different wavelengths through a fiberoptic bundle. The back scattered light from the oxygenated and unoxygenated red cells is transmitted by the receiving fiberoptic bundle to a photodetector in the optical module.

The oximeter computes percent of oxygen saturation values based on the electrical signals from the optical module. The average value for oxygen saturation over a five second period is displayed digitally, and the value updated each second.

The fiberoptic catheter was placed in the right atrium via the umbilical vein. The catheter was positioned at the midatrial level or at the transition of the inferior vena cava to the right atrium. The position was confirmed by X-ray and ultrasound. Routinely, heparinized saline (0.5 U./cc.) was infused continuously (0.1 cc/hr.) to avoid fibrin formation over the catheter. Bloodsamples were withdrawn through the catheter at least every twelve hours and immediately analyzed with a hemoximeter (Radiometer OSM3, Copenhagen) and compared with simultaneously recorded fiberoptic catheter readings. When oxygen saturation in correlated samples showed a discrepancy of $\geq 5\%$, a correction was made.

SaO₂ was monitored continuously by pulse oximetry (Hewlett Packard). Each value is the average of samples taken every 12 seconds over one minute.

Arterial blood pressure was monitored through an umbilical or radial artery catheter; these catheters were also used for sampling arterial pH, PaCO₂, PaO₂, and base excess for clinical purposes.

calculations

Arterial oxygen content (caO₂) and central venous oxygen content (ccvO₂) were calculated using the equation:

$$- \text{caO}_2 \text{ (ml/dl)} = \text{SaO}_2 \times \text{Hb (mmol/L)} \times 1.61 \times 1.36$$

$$- \text{ccvO}_2 \text{ (ml/dl)} = \text{ScvO}_2 \times \text{Hb (mmol/L)} \times 1.61 \times 1.36, \text{ respectively.}$$

Fractional oxygen extraction (F.O.E.) was calculated using the equation: F.O.E. =

$$- (\text{SaO}_2 - \text{ScvO}_2) / \text{SaO}_2.$$

The venous admixture was estimated using the equation for the shunt index:

$$- 100 \times (100 - \text{SaO}_2) / (100 - \text{ScvO}_2), \text{ assuming that the pulmonary end-capillary blood was fully saturated with oxygen.}$$

The alveolar pO₂ (p_{Alv}O₂) was calculated using a simplified alveolar air equation:

$$- \text{p}_{\text{Alv}}\text{O}_2 = (\text{p}_{\text{barometric}} - \text{p}_{\text{(H}_2\text{O)}}) \cdot \text{FiO}_2 - \text{p}_{\text{aCO}_2} / 0.8$$

The pO₂ at an oxygen saturation of 50 % (p₅₀) and of 90 % (p₉₀) were estimated using the oxygen status algorithm [16].

Data analysis

SaO₂, ScvO₂, arterial blood pressure, heart rate and the related calculated variables as venous admixture and fractional oxygen extraction were registered every 15 minutes. Arterial blood gas measurements and hemoglobin values were determined when appropriate and related variables as caO₂, ccvO₂, p_{Alv}O₂, p₅₀ and p₉₀ were calculated. These different parameters were compared for the different levels of oxygenation based on whether the actual SaO₂ value did belong to the lower (86 - 90 %) or higher (91 - 96 %) end of the recommended range of pulse oximetry. In patients with sufficient SaO₂ values below 86 %, the comparison included this low level of oxygenation.

Data are reported as median values with the 10th and 90th percentile range. ScvO₂ values were compared with the described normal range in infants breathing room air. Data were compared for the different level of oxygenation using the Mann-Whitney U test when data were not normally distributed. To describe the relation between SaO₂ and ScvO₂ a linear regression analysis was performed in every patient. The slope (regression coefficient) of the regression line was compared for the two levels of oxygenation within each patient using covariant analysis. The identical analysis was performed for the relation between SaO₂ and venous admixture and ScvO₂ and fractional oxygen extraction.

Results

In 12 newborn infants the ScvO₂ could be monitored continuously by a fiberoptic catheter in the right atrium confirmed by X-ray or ultrasound and reliable oxygen saturation readings were obtained confirmed by correlated blood samples. Birthweight and gestational age are given in table 1. Diagnoses in all but one patients included idiopathic respiratory distress syndrome (IRDS). One patient (4) had oligohydramnion sequence and lung hypoplasia.

table 1. Gestational age (GA), birth weight (BW) and oxygen transport variables. median (10th , 90th percentile)

Patient No	GA (weeks)	BW (gram)	FiO ₂	p _{Alv} O ₂ (kPa)	p ₅₀ (kPa)	p ₉₀ (kPa)
1	26	920	0.30 (0.25 , 0.50)	28 (18 , 43)	3.1 (2.8 , 3.5)	6.9 (6.3 , 7.6)
2	27	720	0.39 (0.21 , 0.65)	31 (13 , 48)	3.9 (3.5 , 4.9)	8.5 (7.5 , 10.3)
3	28.6	940	0.40 (0.32 , 0.68)	32 (22 , 56)	3.0 (2.5 , 3.8)	6.8 (5.7 , 8.3)
4	39	3400	1.00 (1.00 , 1.00)	90 (85 , 92)	3.0 (1.9 , 3.5)	6.8 (4.8 , 7.7)
5	30	1000	0.40 (0.29 , 0.57)	34 (17 , 54)	4.1 (3.6 , 4.9)	8.8 (7.8 , 10.3)
6	28	1355	0.38 (0.21 , 0.60)	22 (13 , 31)	3.2 (2.7 , 4.0)	7.1 (6.2 , 8.6)
7	28	1180	0.25 (0.21 , 0.35)	14 (13 , 20)	3.2 (2.7 , 4.0)	7.1 (6.2 , 8.7)
8	29	1350	0.31 (0.24 , 0.37)	21 (16 , 27)	3.7 (2.7 , 4.2)	8.0 (6.2 , 9.1)
9	25	840	0.26 (0.21 , 0.33)	18 (14 , 29)	3.2 (2.5 , 4.2)	8.1 (6.0 , 9.0)
10	28.3	1130	0.33 (0.21 , 0.56)	30 (14 , 54)	2.9 (2.8 , 3.2)	6.6 (6.4 , 7.2)
11	26.4	840	0.26 (0.21 , 0.30)	16 (14 , 32)	3.4 (3.3 , 3.8)	7.7 (7.4 , 8.3)
12	26.4	830	0.25 (0.21 , 0.45)	19 (13 , 25)	3.3 (2.8 , 3.7)	7.4 (6.3 , 8.8)

Oxygen transport variables as p_{Alv}O₂, p₅₀ and p₉₀ are shown in table 1. In none of the infants a difference between the two levels of oxygenation with regard to the p_{Alv}O₂, p₅₀ and p₉₀ values was found. The plot of the paO₂ and HbO₂ values is shown in figure 1. The median paO₂ values for each level of oxygenation 5.6 kPa (4.6 , 7.3) and 6.8 kPa (5.8 , 8.4) vs. 8.1 kPa (6.6 , 11.0) are significantly different ($p < 0.001$). The different values of the SaO₂ at the levels of oxygenation resulted in eight patients to a significant difference of the caO₂. In all but three patients ScvO₂ values were significantly reduced in the lower compared to the higher level of oxygenation. Linear regression analyses comparing SaO₂ and ScvO₂ yielded a slope (regression coefficient) and a correlation coefficient r (table 2, figure 2). In all but one patient a significant correlation was found. Comparing the slope of the regression line between the two levels of oxygenation did not yield a significant difference in any of the infants. In five patients sufficient SaO₂ values below 86 % could be registered to include the data in table 2 and to perform a regression analysis. ScvO₂ values were significantly reduced in this low level of oxygenation compared to higher levels. In two of these five patients linear regression analysis showed a significant increase of the slope in the oxygenation level below 86 % compared with the level between 86 % and 96 % (figure 2).

The shunt index, an estimate of the venous admixture, increased significantly in the lower level of oxygenation in all the patients. In the five patients with SaO₂ values below 86 % the shunt index increased additionally.

table 2. Arterial oxygen saturation (SaO_2), oxygen content (caO_2) and central venous oxygen saturation (ScvO_2), oxygen content (ccvO_2) at different levels of oxygenation defined by SaO_2 measured by pulse oximetry. Slope and correlation coefficient (r) between SaO_2 and ScvO_2 at the oxygenation level below 86 and above 86 %. '*' : the slope increased significantly at the oxygenation level below 86 %. Median values (10th , 90th percentile).

Patients No	Oxygen transport variables		Correlation			
	Arterial		Central venous		between SaO_2 and ScvO_2	
	SaO_2 (%)	caO_2 (ml/dl)	ScvO_2 (%)	ccvO_2 (ml/dl)	coefficient (r)	slope
1	78 (67 , 84)	13.3 (9.5 , 14.3)	64 (55 , 74)	10.7 (7.8 , 12.6)	0.63	0.68 (0.38 , 0.98)
	89 (86 , 90)	14.7 (13.6 , 15.1)	72 (65 , 78)	12.0 (11.2 , 12.7)	0.72	0.97 (0.69 , 1.15)
	93 (91 , 95)	15.2 (13.4 , 16.2)	77 (69 , 82)	12.4 (10.9 , 13.9)		
2	89 (87 , 90)	17.6 (11.8 , 19.9)	75 (65 , 80)	12.4 (10.0 , 15.8)	0.17	0.33 (0.02 , 0.63)
	93 (91 , 96)	15.6 (12.4 , 20.4)	75 (67 , 85)	13.3 (9.6 , 15.9)		
3	83 (72 , 85)	17.1 (14.4 , 21.3)	70 (64 , 78)	15.6 (10.4 , 17.3)	0.75	0.98 (0.59 , 1.38)
	89 (97 , 90)	17.3 (15.7 , 23.2)	78 (72 , 84)	16.0 (13.1 , 20.9)	0.16	0.35 (0.07 , 0.63)
	92 (91 , 94)	19.4 (16.3 , 24.0)	78 (71 , 86)	20.4 (14.5 , 22.1)		
4	83 (61 , 85)	17.2 (11.2 , 19.8)	69 (19 , 78)	11.6 (8.6 , 15.6)	0.95	1.62 (1.46 , 1.78)*
	88 (86 , 90)	17.3 (16.5 , 20.9)	77 (66 , 81)	15.5 (11.7 , 16.8)	0.35	0.69 (0.49 , 0.89)
	93 (91 , 95)	18.1 (16.6 , 21.3)	80 (71 , 83)	15.8 (13.8 , 16.6)		
5	88 (86 , 90)	18.1 (16.9 , 19.5)	76 (67 , 83)	15.6 (13.8 , 16.7)	0.12	0.23 (-0.09 , 0.56)
	92 (91 , 94)	19.3 (17.6 , 20.6)	79 (63 , 84)	16.5 (13.2 , 17.5)		
6	89 (86 , 90)	15.3 (14.9 , 25.0)	67 (58 , 73)	11.8 (9.9 , 18.1)	0.79	1.40 (1.27 , 1.53)
	93 (91 , 95)	16.8 (15.7 , 23.0)	75 (69 , 80)	14.3 (12.1 , 17.9)		
7	89 (88 , 90)	14.4 (14.1 , 14.5)	66 (58 , 76)	11.7 (11.3 , 12.1)	0.59	0.96 (0.82 , 1.08)
	93 (95 , 96)	20.2 (12.2 , 20.9)	75 (66 , 80)	16.3 (8.5 , 16.7)		
8	88 (86 , 90)	18.5 (14.2 , 22.7)	69 (64 , 73)	13.9 (10.8 , 18.5)	0.76	1.19 (1.07 , 1.31)
	93 (91 , 95)	16.9 (14.8 , 23.9)	73 (69 , 78)	13.4 (11.6 , 19.4)		

9	83 (76 , 85)	14.9 (11.9 , 15.9)	64 (53 , 72)	11.6 (7.7 , 12.8)	0.53	0.75 (0.30 , 1.20)
	89 (86 , 90)	16.9 (13.6 , 18.2)	70 (65 , 73)	13.2 (10.6 , 14.1)	0.17	0.26 (0.08 , 0.42)*
	94 (91 , 96)	17.3 (10.7 , 15.8)	73 (65 , 77)	13.2 (10.7 , 15.9)	0.05	0.33 (0.08 , 0.58)
10	83 (78 , 85)	18.1 (17.4 , 18.8)	67 (57 , 73)	15.1 (13.9 , 16.4)	0.30	0.57 (-0.12 , 1.17)
	88 (87 , 90)	18.5 (16.6 , 19.9)	73 (63 , 80)	15.2 (13.3 , 17.1)	0.34	0.73 (0.47 , 0.98)
	93 (91 , 95)	18.8 (17.9 , 20.6)	75 (70 , 83)	15.0 (13.6 , 18.2)	0.34	0.73 (0.47 , 0.98)
11	89 (87 , 90)	18.9 (16.4 , 20.5)	72 (69 , 73)	15.3 (13.1 , 16.8)	0.79	0.87 (0.77 , 0.97)
	92 (91 , 93)	19.6 (17.6 , 21.2)	74 (72 , 76)	16.0 (14.0 , 17.4)	0.79	0.87 (0.77 , 0.97)
12	89 (86 , 90)	14.4 (13.6 , 15.3)	70 (67 , 76)	11.5 (11.4 , 13.0)	0.47	0.42 (0.29 , 0.56)
	93 (91 , 96)	15.8 (14.1 , 16.3)	73 (67 , 79)	12.6 (10.5 , 13.5)	0.47	0.42 (0.29 , 0.56)

(100% - 100%)

WVQ (100%)

(20%)

WVQ (100%)

WVQ (100%)

tabel 3. Arterial oxygen saturation (SaO₂), fractional oxygen extraction (FOE) and shunt index at different levels of oxygenation defined by SaO₂ measured by pulse oximetry. Correlation coefficient (*r* (ScvO₂) between ScvO₂ and FOE at the oxygenation level below 86 and above 86 %. Correlation coefficient (*r* (SaO₂) between SaO₂ and shunt index at the oxygenation level below 86 and above 86 %. ** : *r* increased significantly at the oxygenation level above 86 %. Median values (10th , 90th percentile).

patient No	Oxygenation level	Fractional oxygen extraction		Shunt index		
	SaO ₂ (%)	median (10th , 90th)	<i>r</i> (ScvO ₂)	Median (%) (10th , 90th)	<i>r</i> (SaO ₂)	slope (95 % CI)
1	78 (67 , 84)	0.17 (0.10 , 0.24)	0.63	60 (47 , 79)	0.72	-1.4 (-1.9 , -0.9)*
	89 (86 , 90)	0.19 (0.13 , 0.26)		42 (31 , 52)	0.73	-2.5 (-2.9 , -2.1)
	93 (91 , 95)	0.17 (0.12 , 0.25)		30 (10 , 40)		
2	89 (87 , 90)	0.16 (0.11 , 0.27)	0.91	45 (33 , 53)	0.77	-3.4 (-3.8 , -3.0)
	93 (91 , 96)	0.20 (0.08 , 0.27)		26 (15 , 50)		
3	83 (72 , 85)	0.12 (0.03 , 0.20)	0.82	64 (48 , 87)	0.29	-0.6 (-1.5 , 0.3)*
	89 (97 , 90)	0.11 (0.06 , 0.19)		52 (39 , 67)	0.61	-3.7 (-4.3 , -3.1)
	92 (91 , 94)	0.12 (0.06 , 0.19)		36 (23 , 56)		
4	83 (61 , 85)	0.18 (0.08 , 0.64)	0.94	60 (47 , 73)	0.03	-0.03 (-0.16 , 0.10)*
	88 (86 , 90)	0.13 (0.09 , 0.25)		47 (35 , 64)	0.74	-3.3 (-3.7 , -2.9)
	93 (91 , 95)	0.14 (0.10 , 0.24)		35 (20 , 45)		
5	88 (86 , 90)	0.14 (0.07 , 0.25)	0.89	48 (32 , 67)	0.73	-3.6 (-4.1 , -3.0)
	92 (91 , 94)	0.15 (0.09 , 0.32)		35 (19 , 50)		
6	89 (86 , 90)	0.23 (0.18 , 0.34)	0.94	35 (28 , 40)	0.78	-1.5 (-1.7 , -1.4)
	93 (91 , 95)	0.20 (0.15 , 0.26)		26 (20 , 34)		
7	89 (88 , 90)	0.26 (0.15 , 0.34)	0.89	31 (25 , 48)	0.87	-2.1 (-2.3 , -2.0)
	93 (95 , 96)	0.21 (0.16 , 0.28)		20 (15 , 27)		
8	88 (86 , 90)	0.22 (0.17 , 0.28)	0.87	37 (31 , 44)	0.92	-2.7 (-2.8 , -2.6)
	93 (91 , 95)	0.21 (0.16 , 0.25)		28 (20 , 34)		

9	83 (76 , 85)	0.23 (0.12 , 0.31)	0.75	51 (39 , 70)	0.60	-1.3 (-2.0 , -0.6)*
	89 (86 , 90)	0.20 (0.17 , 0.27)		38 (32 , 46)	0.90	-3.1 (-3.3 , -2.9)
	94 (91 , 96)	0.22 (0.16 , 0.30)		24 (14 , 34)		
10	83 (78 , 85)	0.19 (0.11 , 0.31)	0.84	50 (40 , 66)	0.59	-2.0 (-3.3 , -0.7)
	88 (87 , 90)	0.18 (0.10 , 0.28)		42 (31 , 56)	0.71	-3.1 (-3.5 , -2.7)
	93 (91 , 95)	0.19 (0.11 , 0.25)		28 (20 , 45)		
11	89 (87 , 90)	0.19 (0.17 , 0.21)	0.70	40 (36 , 44)	0.92	-2.4 (-2.5 , -2.2)
	92 (91 , 93)	0.20 (0.18 , 0.22)		32 (27 , 36)		
12	89 (86 , 90)	0.19 (0.15 , 0.23)	0.60	39 (33 , 47)	0.96	-3.0 (-3.1 , -2.8)
	93 (91 , 96)	0.22 (0.17 , 0.28)		24 (15 , 33)

figure 1. Arterial pO_2 (paO_2) corresponding to arterial oxygen saturation (HbO_2). The paO_2 values: 5.6 kPa (4.6 , 7.3), 6.8 kPa (5.8 , 8.4) vs. 8.1 kPa (6.6 , 11.0) for each level of oxygenation: 83 % (79 , 85), 88 % (86 , 90) and 93 % (91 , 95) are significantly different ($p < 0.001$).

Linear regression analyses for each patient separately showed a negative correlation with SaO_2 in all patients (table 3, figure 4).

No differences were found in the slope of the regression lines between the two recommended SaO_2 levels above and below 90 %. In four of the five patients with SaO_2 values below 86 % the slope of the regression line was significantly reduced in comparison with the slope of the regression line obtained at SaO_2 values above 86 %.

Fractional oxygen extraction (FOE) was significantly increased in five of the twelve patients in the lower level of oxygenation. Linear regression analysis yielded only a significant correlation in six patients with a low range of correlation coefficients: r : 0.25 (0.04 , 0.42). However, linear regression analysis between $ScvO_2$ and fractional oxygen extraction showed a significant correlation in all patients (table 3, figure 4). Comparing the three SaO_2 levels fractional oxygenation remained stable, when $ScvO_2$ decreased.

Discussion

In this study we report on line monitoring $ScvO_2$ by an umbilical fiberoptic catheter in 12 newborn infants with moderate to severe respiratory failure. $ScvO_2$ values and derived variables, as the shunt index and fractional oxygen extraction were compared at low values (SaO_2 below 86 %) and the lower (86 - 90 %) and the higher (91 - 96 %) end of the recommended range of SaO_2 values.

figure 2. Arterial oxygen saturation values obtained by pulse oximetry (SaO_2) corresponding to central venous oxygen saturation values simultaneously obtained by fiberoptic catheter readings ($ScvO_2$). Linear regression lines are drawn for each patient and regression analysis showed in all but one patient a significant correlation in at SaO_2 values above 86 %. **: In two of the five patients the slope of the regression line at SaO_2 values below 86 % was significantly different from the slope at SaO_2 values above 86 %. The arcuated area is the normal range of $ScvO_2$ values as defined in infants breathing room air.

$ScvO_2$ was significantly reduced in the lower level of the recommended range of oxygenation in nine of the twelve patients. Linear regression analysis yielded a significant correlation between SaO_2 and $ScvO_2$ in all but one of the patients. However, it must be recognized that changes in $ScvO_2$ values may not only be due to changes in SaO_2 but also to changes in cardiac output, O_2 uptake, hemoglobin, shift of the oxyhemoglobin dissociation curve or a combination of these variables^[15]. For example, since in our present study the different SaO_2 value in the different levels of oxygenation did not always result in a different caO_2 , due to differences in hemoglobin concentration, $ScvO_2$ values may not be affected by changes in SaO_2 .

In addition, $ScvO_2$ value may also be affected by cardiac output and tissue oxygen uptake. $ScvO_2$ monitoring does not yield specific information regarding any individual oxygen transport variable. But in prospective studies in newborn infants^[4,5] it was reported that the cardiac output and oxygen consumption did not change neither in the low nor in the high level of the recommended range of oxygenation.

figure 3. Arterial oxygen saturation values obtained by pulse oximetry (SaO_2) corresponding to the shunt index, an estimate of the venous admixture. Linear regression lines are drawn for each patient. Regression analysis showed a significant correlation in all patients. "": in four of the five patients with SaO_2 values below 86 % the slope of the regression line at SaO_2 values below 86 % was significantly different from the slope of the regression line obtained at SaO_2 values above 86 %

Consequently, the reduction of ScvO_2 reported in our study seemed at these levels of oxygenation the primary compensatory mechanism to stabilize the tissue oxygenation at lower SaO_2 values. This is in agreement with our animal study in which moderate reductions of ScvO_2 were not associated with a reduced oxygen consumption or an increased lactate concentration^[19]. The amount of ScvO_2 reduction did not change since the slope of the regression line did not differ between the two levels of oxygenation in any of the infants. In two of the five patients with SaO_2 values below 86 % the rate of ScvO_2 reduction was significantly increased compared with the change in the recommended range of SaO_2 values. Apparently, in these two patients oxygen uptake increased whereas other factors which determine oxygen delivery as cardiac output failed to counterbalance the reduced oxygen saturation. When the present ScvO_2 values were compared with the "normal range" defined in infants breathing room air^[14], most ScvO_2 values in both recommended levels were included within this range. Lower ScvO_2 values were mainly encountered with SaO_2 values below 86 %.

The shunt index is an estimate of the venous admixture that refers to the degree of admixture of mixed venous blood with pulmonary end-capillary blood. The venous admixture is a calculated percentage of blood, which would be required to produce the observed difference between the arterial and the pulmonary end-capillary pO_2 , which is

figure 4. Central venous oxygen saturation (ScvO₂) values obtained by fiberoptic catheter readings corresponding to the fractional oxygen extraction.

usually taken to equal ideal alveolar $p_{\text{alv}}\text{O}_2$ [20]. The calculated $p_{\text{alv}}\text{O}_2$ values in our study (table 1) are such that the assumptions of the shunt index, that the pulmonary end-capillary blood is fully saturated and that the dissolved oxygen can be discounted, are warranted. In our study we found an increased shunt index in the lower end of the recommended range of oxygenation, which even further increased in the patients in whom SaO₂ values below 86 % were obtained. Linear regression analysis yielded a significant correlation in all the patients.

The shunt index as an estimate of the venous admixture obtained by continuous pulse and venous oximetry (dual oximetry) has been suggested and studied in adult intensive care medicine [21,22,23]. It was used to titrate rapidly continuous positive pressure (CPAP) in adults [24]. In neonatal medicine the association between venous admixture and biochemical lung maturity was determined [25] and after administration of surfactant a rapid fall of venous admixture was described [26]. In agreement with our study Schultze et al. [5] found an increased venous admixture in the low level of oxygenation in preterm infants targeting different levels of oxygenation. Although it has been shown, that decreasing SaO₂ from 95 % to 90 % corresponding to $p\text{aO}_2$ values of 11.6 ± 0.7 kPa and 7.5 ± 0.4 kPa, respectively, did not have any effect on pulmonary circulatory hemodynamics [4], a certain grade of intrapulmonary shunting, ventilation perfusion mismatch or pulmonary vasoconstriction must be assumed to explain the increased venous admixture in the lower levels of oxygenation. At SaO₂ values below 86 % the shunt index increased at a reduced

rate since the slope of the regression lines significantly decreased in four of the five patients compared with the lines obtained at SaO_2 values above 86 %. (table 3, figure 3) As other factors which determine oxygen delivery as cardiac output failed to counterbalance the reduced oxygen saturation, ScvO_2 starts to decrease at a higher rate (figure 2). Consequently the shunt index will increase at a lower rate when arterial oxygen saturation was further reduced.

Fractional oxygen extraction was in six patients significantly correlated with SaO_2 , albeit weak. ($r = 0.18$ to 0.39). Mathematically it can be expected that a good correlation exists between fractional oxygen extraction and ScvO_2 . In figure 4 it can be seen that for the same level of ScvO_2 fractional oxygen extraction increased facing the increased levels of oxygenation. In other terms fractional oxygen extraction remained at the same level when with a reducing oxygenation level ScvO_2 decreased concomitantly. The clinical value of fractional oxygen extraction has been considered as an estimate of the peripheral oxygen utilization by the tissues and an instantaneous feedback on circulatory and respiratory therapy [27]. However, fractional oxygen extraction based on ScvO_2 reflects global oxygen extraction and therefore tissue hypoxia in individual organs cannot be excluded [28].

In summary on line measurement of ScvO_2 at different levels of SaO_2 values showed a significant reduction of ScvO_2 . Moreover, this reduction of ScvO_2 seems to be the primary compensatory mechanism to stabilize tissue oxygenation. On line ScvO_2 measurement allowed an estimate of the venous admixture and fractional oxygen extraction. At the lower level of SaO_2 values an increased venous admixture seems to indicate a decreased effectiveness in gas exchange. Fractional oxygen extraction was only slightly affected by the level of SaO_2 values.

References

- 1 Moss AF, Emmannouillides GC, Adams FH. Response of ductus arteriosus and pulmonary and systemic arterial pressures to changes in oxygen environment in the newborn infant. *Pediatrics* 1964;33:937-944
- 2 Hrbek A, Karlberg P, Kjellmer I, Olsson T, Riha M. Clinical application of evoked EEG responses in newborn infants II: Idiopathic respiratory distress syndrome. *Develop Med Child Neurol* 1978;20:619-626
- 3 Émond D, Lachance C, Gagnon J, Bard H. Arterial partial pressure of oxygen required to achieve 90 % saturation of hemoglobin in very low birth weight newborns. *Pediatrics* 1993;91:602-604
- 4 Bard H, Belanger S, Fouron JC. Comparison of effects of 95 % and 90 % oxygen saturations in respiratory distress syndrome. *Arch Dis Child* 1996;75:F94-F96
- 5 Schultze A, Whyte RK, Clifton Way R, Sinclair JC. Effect of the arterial oxygenation level on cardiac output, oxygen extraction and oxygen consumption in low birth weight infants receiving mechanical ventilation. *J Pediatr* 1995;126:777-784
- 6 Joint working group of the British Association of Perinatal Medicine and the Research Unit of the Royal College of Physicians. Development of audit measures and guidelines for good practice in the management of neonatal respiratory distress syndrome. *Arch Dis Child* 1992;67:1221-1227
- 7 American Academy of Pediatrics and American College of Obstetricians and Gynecologists. Guidelines for perinatal care. 3rd ed. Washington, DC. 1992:197-203
- 8 Southall DP, Samuels MP, Poets CF. Neonatal respiratory distress syndrome [letter]. *Arch Dis Child* 1993;69:330
- 9 Cochran DP, Shaw NJ. Neonatal respiratory distress syndrome [letter]. *Arch Dis Child* 1994;70:F230
- 10 O'Connor TA, Hall RT. Mixed venous oxygenation in critically ill neonates. *Crit Care Med* 1994;22:343-346

- 11 Dudell G, Cornisch JD, Bartlett RH. What constitutes adequate oxygenation? *Pediatrics* 1990;85:39-41
- 12 White RK. Mixed venous oxygen saturation in the newborn: can we and should we measure it? *Scand J Clin Lab Invest Suppl* 1990;203:203-211
- 13 van der Hoeven MAHBM, Maertzdorf WJ, Blanco CE. Feasibility and accuracy of a fiberoptic catheter for the measurement of venous oxygen saturation in newborn infants. *Acta Paediatr* 1995;84:122-127
- 14 van der Hoeven MAHBM, Maertzdorf WJ, Blanco CE. Continuous central venous oxygen saturation (ScvO₂) measurement using a fiberoptic catheter in newborn infants. *Arch Dis Child* 1996;74:F177-F181
- 15 Nelson LD. Continuous venous oximetry in surgical patients. *Ann Surg* 1986;203:329-333
- 16 Siggaard-Andersen O, Fogh-Andersen N, Gøthgen IH, Højtkjær Larsen V. Oxygen status of arterial and mixed venous blood. *Crit Care Med* 1995;23:1284-1293
- 17 Bartlett RH, Cilley RE. 1993 Physiology of extracorporeal life support. In: Arensman RM, Cornish JD (eds). *Extracorporeal life support*. Blackwell, Boston, pp 89-104
- 18 Wasunna A, Whitelaw AG. Pulse oximetry in preterm infants. *Arch Dis Child* 1987;62:882-888
- 19 van der Hoeven MA, Maertzdorf WJ, Blanco CE. Mixed venous oxygen saturation (SvO₂) and biochemical parameters of hypoxia during progressive hypoxemia in 10-14 day old piglets. *Pediatr Res* 1997;42:878-884
- 20 Nunn JF. Distribution of pulmonary ventilation and perfusion. In: Nunn's applied respiratory physiology. Oxford, Butterworth, Heinemann. 1993, pp 156-197
- 21 Bongard FS, Leighton TA. Continuous dual oximetry in surgical critical care. Indications and limitations. *Ann Surg* 1992;216:60-68
- 22 Räsänen J, Downs JB, Malec DJ, Seidman P. Estimation of oxygen utilization by dual oximetry. *Ann Surg* 1987;206:621-623
- 23 Carrol GC. A continuous monitoring technique for management of acute pulmonary failure. *Chest* 1987;92:467-469
- 24 Räsänen J, Downs JB, Dehaven B. Titration of continuous positive airway pressure by real time dual oximetry. *Chest* 1987;92:853-856
- 25 de Winter JP, Egberts J, Merth IT, Peerdeman A, van Bel F, Bakker GC, Quanjer PH. Biochemical lung maturity static respiratory compliance, and pulmonary gas transfer in intubated preterm infants with and without respiratory distress syndrome. *Pediatr Pulmonol* 1995;20:152-159
- 26 de Winter JP, Merth IT, van Bel F, Egberts J, Brand R, Quanjer PH. Changes of respiratory system mechanics in ventilated lungs of preterm infants with two different schedules of surfactant treatment. *Pediatr Res* 1994;35: 541-549
- 27 Räsänen J. Role of dual oximetry in the assessment of pulmonary function. In: Reinhart K, Eyrich K. (Eds) *Clinical aspects of O₂ transport and tissue oxygenation*. Springer Verlag, Berlin, 1989; 230-240
- 28 Reinhart K. Monitoring O₂ transport and tissue oxygenation in critically ill patients. In: Reinhart K, Eyrich K. Editors. *Clinical aspects of O₂ transport and tissue oxygenation*. Berlin: Springer Verlag, 1989; pp 193-211

Acknowledgements

This study was supported by Abbott Inc. the Netherlands.

the same, the rate of oxygenation was significantly lower in the patients with severe hypoxemia compared with the group obtained at SaO_2 values above 80%. In the latter group, the rate of oxygenation was stable or slightly increased after the first 10 min of reduced oxygenation. In the group with severe hypoxemia, the rate of oxygenation decreased after the first 10 min of reduced oxygenation. The rate of oxygenation was significantly lower in the patients with severe hypoxemia compared with the group obtained at SaO_2 values above 80%.

The rate of oxygenation was significantly lower in the patients with severe hypoxemia compared with the group obtained at SaO_2 values above 80%. In the latter group, the rate of oxygenation was stable or slightly increased after the first 10 min of reduced oxygenation. In the group with severe hypoxemia, the rate of oxygenation decreased after the first 10 min of reduced oxygenation. The rate of oxygenation was significantly lower in the patients with severe hypoxemia compared with the group obtained at SaO_2 values above 80%.

Fig. 1. Rate of oxygenation (ml/min) versus time (min) for two groups of patients. The group with severe hypoxemia (n=10) shows a rapid decrease in oxygenation rate, while the group with $\text{SaO}_2 > 80\%$ (n=10) shows a more stable or slightly increased rate.

Summary and Conclusions

In the introduction in this thesis it was stated that it is the objective of intensive care to preserve the human organism by supporting gasexchange with the environment, by improving oxygen transport to the tissues and by monitoring and promoting oxygen uptake by the tissues. It is not astonishing, therefore, that the measurement of SvO_2 raised so much interest among intensive care physicians, since SvO_2 is an integrated value of all these objectives together.

In neonatal medicine oxygen monitoring and oxygen supply has remained a topic during the history of this discipline. Since both hyperoxia and hypoxia have their damaging effects, the questions, how much oxygen the tissues really need, and how to monitor these oxygen needs, are still not answered completely.

We performed five studies to investigate whether the measurement of SvO_2 may give some answers on these issues. The main questions related to these studies were formulated in the introduction and are subsequently discussed in the following paragraphs:

1. What is the relation between tissue oxygenation and the SvO_2 value?
2. Is this relation between tissue oxygenation and SvO_2 different when oxygen delivery to the tissues is reduced by different manners?

To examine the clinical possibilities of the SvO_2 measurement in the newborn period, the following questions were addressed:

3. Is it possible to measure SvO_2 continuously in a newborn infant? Where? Are the SvO_2 values reliable and accurate?
4. What is the normal range of SvO_2 values in stable newborn infants breathing room air? What is the relation between SaO_2 and SvO_2 values?
5. What is the range of SvO_2 values during the acute phase of IRDS during mechanical ventilation and oxygen administration? What is the relation between SaO_2 and SvO_2 values and derived calculations as venous admixture and fractional oxygen extraction? What is the value of $ScvO_2$ in clinical neonatal medicine?

1. SvO_2 and tissue oxygenation

We have assessed the relationship between SvO_2 and tissue oxygenation in an acute neonatal piglet model during hypoxic hypoxia. During the graded steps of hypoxic hypoxia SvO_2 decreases parallel to SaO_2 , oxygen extraction increases and consequently a stable VO_2 can be preserved during a rather prolonged period of graded hypoxia. The reduction of SvO_2 and the concomitant increase of oxygen extraction seem to be one of the mechanisms to preserve VO_2 during moderate hypoxia. Other mechanisms to compensate are an increased cardiac output and a redistribution of blood flow between different (parts of) organs. A redistribution of blood flow was not measured in our experiments. In fact, the value of SvO_2 and the distribution of blood flow are related entities, since by a redistribution of blood to high extraction organs the average oxygen extraction will increase and SvO_2 will decrease.

This finding means that a reduced SvO_2 value in itself is not a sign of anaerobic metabolism but is a sign that an organism is calling on its compensatory mechanisms. When more severe hypoxic hypoxia is induced, the decrease of SaO_2 is not any more paralleled by SvO_2 , oxygen consumption is reduced and anaerobic metabolism and lactic acidosis will develop.

At what SvO_2 value exactly anaerobic metabolism will develop was determined in

this study: When the SvO_2 value decreased below 15 %, tissue oxygen consumption decreased and anaerobic metabolism developed. A value of SvO_2 above 40 % excluded anaerobic metabolism. SvO_2 values between 15 % and 40 % were rarely a marker for anaerobic metabolism.

Some remarks have to be made to delineate these findings before they can be extrapolated to clinical medicine:

The time aspect in this acute study must be taken into consideration. The above mentioned values could become different in experiments with a more prolonged duration. Secondly, the experiments were performed in anesthetized piglets with mechanical ventilation and pancuronium infusion and consequently a stable and rather low VO_2 . In clinical medicine in conditions with a more variable and higher VO_2 , critical values of SvO_2 could become different. Thirdly, one must be cautious to use SvO_2 to monitor exclusively tissue oxygenation, as this latter is only one of the many variables, that can modify the SvO_2 value. Fourth, the SvO_2 is a concept defined by the averaged value of the venous oxygen saturations and of the averaged blood flow from all the different organs. This means that normal SvO_2 values do not exclude an imbalance between DO_2 and VO_2 in one or two organs. Finally, the relation between SvO_2 values and tissue oxygenation can be even more complex in conditions as sepsis and ARDS, when tissue oxygen uptake is impaired and lactic acidosis exists together with normal or even high SvO_2 values.

Probably because of all these reasons aiming at normal SvO_2 values was shown not to affect mortality and other oxygen variables in prospective clinical trials in critical ill adult patients, as was discussed in the review on SvO_2 and DO_2 / VO_2 relationship. However, it must be recognized that in these trials patients with complex hemodynamic problems were studied. With respect to the question on the relation between tissue oxygenation and SvO_2 during hypoxic hypoxia more direct answers could be given.

Summarizing, reduced SvO_2 values during hypoxic hypoxia in itself are not an sign for a reduced VO_2 and for anaerobic metabolism. As long as a reduction of SaO_2 is paralleled by a equivalent reduction of SvO_2 , a stable VO_2 can be preserved. When severe hypoxic hypoxia is induced, the difference between SaO_2 and SvO_2 will decrease and VO_2 will reduce. Consequently, the relation between SvO_2 and tissue oxygenation during hypoxic hypoxia is characterized by a low threshold SvO_2 value, below which anaerobic metabolism will ensue. In addition, the above mentioned considerations and remarks must be taken into account, when the relation between SvO_2 and tissue oxygenation is evaluated.

2. The relation between tissue oxygenation and SvO_2 during anemic hypoxia compared to hypoxic hypoxia.

Another study, assessing the relation between SvO_2 and tissue oxygenation, was performed to compare the range of SvO_2 values during hypoxic hypoxia with the SvO_2 values during anemic hypoxia. As different types of hypoxia could have other patterns of circulatory adjustments and compensations, tissue oxygen uptake and consequently SvO_2 may be affected.

The critical range of DO_2 and oxygen extraction, however, showed no difference between the two types of hypoxia. Hemodynamic responses differed between hypoxic and anemic hypoxia, consisting of an unaltered and increased cardiac output, respectively. We found higher and more variable SvO_2 values during anemic hypoxia than during hypoxic hypoxia within and below the critical DO_2 range. In our opinion, the most important reasons for this finding include *a)* the use of the whole range of the oxygen dissociation curve in anemic hypoxia, *b)* the induced expansion of the plasma phase with its very low

oxygen solubility, limiting the ability to transfer oxygen from the red cells to the tissues and c) the increased red cell velocity with a reduced residence time during anemic hypoxia decreasing precapillary oxygen release and diffusional oxygen transfer to other microvessels.

Summarizing, the relation between SvO_2 and tissue oxygenation seems to be even more complex than initially was considered, since a critical SvO_2 range, below which anaerobic metabolism develops, is shown to be dependent upon the type of hypoxia. However, it can be concluded that during both types hypoxia VO_2 could be preserved as long as a stable difference between arterial and venous oxygen content was maintained. Consequently, a threshold $ScvO_2$ value, below which VO_2 decreased, could only be discerned at quite low $ScvO_2$ values.

3a. The continuous measurement of SvO_2 in newborn infants.

It has been suggested to measure SvO_2 in clinical neonatal medicine to monitor the oxygen needs of the tissues and, consequently, to minimize on one hand oxygen toxicity related diseases as retinopathy of prematurity, bronchopulmonary dysplasia and necrotizing enterocolitis and on the other hand hypoxia related diseases such as cerebral palsy. Since the SvO_2 measurement in the pulmonary artery is difficult and hazardous, SvO_2 measurements were performed in the right atrium (central venous oxygen saturation: $ScvO_2$) by a catheter capable of measuring oxygen saturation by fiberoptic oximetry continuously.

3b. The feasibility of the fiberoptic catheter. The reliability and accuracy of the $ScvO_2$ values. The position of the catheter.

We tested the feasibility and accuracy of the continuous $ScvO_2$ measurement with a 4 Fr. fiberoptic catheter, inserted into the right atrium via the umbilical vein to measure $ScvO_2$, continuously.

With regard to feasibility, we conclude that the success rate and complications of the fiberoptic catheter are similar to previously published results. In 67% of the patients the fiberoptic catheter could be placed in the correct position, i.e. at the mid-atrial level or at the transition of the vena cava inferior and the right atrium. In some infants at the end of the indwelling period, the signal was affected by unusually small light reflectance at the catheter tip and saturation readings were not considered to be reliable any more. Complications such as pericardial effusions, cardiac tamponade, cardiac arrhythmias or perforations were not seen.

With regard to the accuracy it was concluded that the oxygen saturation readings accurately and reliably represented oxygen saturation in the right atrium of newborn infants. The mean difference (± 2 SD) between the fiberoptic catheter $ScvO_2$ readings and the hemoximeter HbO_2 values was -0.36 ± 7.64 . This value of difference and SD is comparable with the data mentioned in the review on accuracy of the fiberoptic oximetry used in pulmonary artery catheters.

The position of the catheter was extensively discussed in the review. It was concluded that the drawbacks (incomplete mixing, susceptibility to catheter motion and atrium wall reflection) are probably as valid in neonatal medicine as in adult medicine. However, the measurement of venous oxygen saturation in newborn infants is limited to the right atrium, since the use of a pulmonary artery catheter is difficult and hazardous. The drawbacks of right atrial oxygen saturation measurement must be taken into account

by consideration of the position of the catheter, frequent in vivo calibration and caution in the calculation of oxygen transport variables based on central venous oxygen saturations.

Some remarks have to be made upon the limitations and drawbacks of the use of a single lumen catheter in practical neonatal intensive care medicine. Recently, in the care of very small newborn infants the use of double lumen or even triple lumen umbilical vein catheters have become a common procedure. The use of these catheters is warranted by a simultaneous use of continuous medications and parenteral nutrition. The use of a single lumen catheter as the 4 Fr. fiberoptic catheter seems now to be limited to more stable infants in whom the continuous ScvO₂ measurement does not seem directly necessary. Another limitation is the fact that, as was shown in our feasibility study, the positioning of the catheter through the ductus venosus in the right atrium was only successful in 60 % of the cases. The limitation of the measurement of the ScvO₂ in the right atrium and the risk of incomplete mixing have been discussed extensively. In conclusion, the limitations of a single lumen umbilical venous catheter makes one reluctant to consider ScvO₂ measurement as a standard procedure in neonatal intensive care medicine. On the other hand the fiberoptic catheter readings do accurately represent ScvO₂ values reflecting the residual oxygen after tissue oxygen extraction.

4. The normal range of ScvO₂ values in stable newborn infants breathing room air. The relation between SaO₂ and ScvO₂ values. The value of fractional oxygen extraction (F.O.E.) and the shunt index.

In a prospective clinical observational study the range of ScvO₂ was described in 10 stable newborn infants breathing room air. In this study a ScvO₂ range from 60.1 % to 83.5 % (5th and 95th percentile) was observed, when other variables affecting oxygen transport were in the normal range. i.e. SaO₂: $\geq 86\%$; hemoglobin: ≥ 7.0 mmol/L; pH and blood pressure: normal. This ScvO₂ range is similar to the normal range defined for adults, whereas VO₂ in newborn infants is increased. Increased VO₂ is apparently compensated for by an increased DO₂. And indeed both cardiac output and hemoglobin concentration are increased in the newborn period.

The relationship between ScvO₂ and SaO₂ was examined. In seven patients ScvO₂ values correlated significantly with SaO₂. This indicates that, when SaO₂ is reduced, VO₂ is preserved by a fixed oxygen extraction. In three patients SaO₂ and ScvO₂ were not correlated. We might speculate, that in these babies VO₂ was preserved by increased cardiac output. It is unlikely that VO₂ was decreased by SaO₂ reduction, because in stable patients with a normal blood pressure, normal haemoglobin level and normal oxygen saturation VO₂ is independent of DO₂ over a wide range.

DO₂ is normally four to five times VO₂ as shown by the normal fractional oxygen extraction (F.O.E.) of 0.20 - 0.25, as described in adults. When analyzed for each patient separately, a range (5th and 95th percentile) was found between 0.13 to 0.33. The highly significant correlation between F.O.E. and ScvO₂ in all patients confirms that ScvO₂ is an excellent monitor of the balance between oxygen delivery and tissue oxygen consumption.

In addition, the measurement of ScvO₂ enables the calculation of the shunt index, an estimate of the venous admixture. A range of the shunt index was found between 7 % and 53 %. We found a significant negative correlation between the shunt index and SaO₂ in all patients. Obviously, an increased shunt index is present when lower SaO₂ values are encountered.

5. The range of $ScvO_2$ values during the acute phase of IRDS during mechanical ventilation and oxygen supply at different levels of oxygenation. The relation between SaO_2 and $ScvO_2$ values and derived calculations as shunt index and F.O.E. The value $ScvO_2$ in clinical neonatal medicine.

A non-randomized clinical observational study was performed in twelve newborn infants with mechanical ventilation and oxygen administration to determine $ScvO_2$ values at different oxygenation levels namely: at SaO_2 values below 86 % and at the lower (86 - 90 %) and higher (91 - 96 %) end of the recommended range of SaO_2 values measured by pulse oximetry.

The $ScvO_2$ values in both recommended levels are not beyond the "normal range" defined in infants breathing room air. Lower $ScvO_2$ values are mainly encountered in the lowest oxygenation level with a SaO_2 below 86 %.

A significant correlation between SaO_2 and $ScvO_2$ was found in all but one of the patients. The slope of the regression lines did not show a significant difference between the two levels of oxygenation in any of the infants. In the SaO_2 level below 86 % $ScvO_2$ values were significantly reduced compared with the level between 86 % - 96 %. In two patients the rate of $ScvO_2$ reduction (the slope) was significantly increased compared with the change in the SaO_2 level between 86 % - 96 %. Apparently, these two patients tried to increase oxygen uptake whereas other factors which determine oxygen delivery as cardiac output failed to counterbalance the reduced oxygen saturation.

An increased shunt index was found in our study in the lower end of the recommended range of oxygenation, which even further increased in the patients in whom SaO_2 values below 86 % were obtained. Linear regression analysis yielded a significant correlation in all the patients. It may be assumed that a certain grade of intrapulmonary shunting or ventilation / perfusion mismatch and pulmonary vasoconstriction must exist to explain the increased venous admixture in the lower levels of oxygenation. On the other hand it can be questioned, whether a calculated item as the shunt index really reflects these clinical problems, although the assumptions were not violated.

Fractional oxygen extraction was in six patients significantly correlated with SaO_2 , albeit weak. ($r = 0.18$ to 0.39). When F.O.E. was compared for the different levels of oxygenation, F.O.E. remained at the same level when $ScvO_2$ decreased concomitantly with a reducing oxygenation level.

In conclusion, on line measurement of $ScvO_2$ at decreasing levels of SaO_2 showed a significant reduction of $ScvO_2$. The $ScvO_2$ values were not beyond the range described in infants breathing room air. This reduction of $ScvO_2$ seems to be the primary compensatory mechanism to stabilize oxygen extraction and tissue oxygenation. Referring to the debate on SaO_2 alarm limits in oxygen therapy in the newborn period the measurement of $ScvO_2$ may introduce a new element in this debate. Since $ScvO_2$ decreased parallel with the reduction of SaO_2 , we have shown that even at the lowest SaO_2 levels tissue oxygen uptake is preserved. Consequently, it can be concluded that the difference between SaO_2 and $ScvO_2$ is more important than the absolute value of $ScvO_2$.

On the question whether it is of interest to know the $ScvO_2$ value, three cases will be presented as examples that continuous $ScvO_2$ measurement could be a valuable complimentary monitor of oxygenation:

case 1. A preterm infant with a gestational age of 28 weeks and a birth weight of 890 gram. Diagnoses: IRDS and E. Coli sepsis and meningitis. The infant required positive pressure ventilation, later followed by high frequency ventilation and intratracheal surfactant. The clinical course was complicated by an intraventricular hemorrhage. The umbilical

vein was catheterized with a fiberoptic catheter for central venous access and ScvO₂ measurement. The tip of the catheter was located just above the junction of the right atrium and the inferior caval vein. The fiberoptic readings correlated with the hemoximeter values ($77.16\% \pm 3.25$ vs $76.16\% \pm 4.41$). The hemoglobin level decreased from 8.5 mmol/L to 7.6 mmol/L and further to 6.8 mmol/L due to multiple blood sampling. SaO₂ monitored by pulse oximetry did not show any alteration, however, ScvO₂ decreased significantly with an increase in fractional oxygen extraction.

case 2. A preterm infant with a gestational age of 27 weeks and a birth weight of 720 gram. The delivery was complicated by a moderate asphyxia. The umbilical vein was catheterized with a fiberoptic catheter for central venous access and ScvO₂ measurement. The course was complicated by severe hypotension and respiratory insufficiency. Intermittent positive pressure ventilation and surfactant replacement therapy were required. Blood pressure was supported by volume replacement and the use of dopamine and dobutamine. After increasing dopamine up to a level of $30\text{ }\mu\text{g/kg/min}$, the ScvO₂ level increased from $70.5\% \pm 5.0$ to $78.6\% \pm 6.7$ in contrast to the SaO₂, which decreased from $91.8\% \pm 3.4$ to $90.0\% \pm 2.6$.

case 3. A preterm infant with a gestational age of 28 weeks and a birth weight of 1350 gram. The delivery was complicated by a severe asphyxia: arterial umbilical cord pH was 6.89. Immediately after delivery ventilator support was initiated. After stabilization, a fiberoptic catheter was inserted into the right atrium for central venous access. The infant stabilized and the ventilatory support could be discontinued. However sudden epileptic spells occurred associated with an elevation of the heart frequency and the blood pressure and a small decrease in the SaO₂. The ScvO₂ decreased during these episodes from 75 % to 55%. Anticonvulsive therapy was started and these episodes did not repeat.

The three cases here reported illustrate that the continuous monitoring of the ScvO₂ allowed to detect an alteration in the balance between DO₂ and VO₂, while SaO₂ did not show changes. The first patient demonstrated a decrease in arterial oxygen content by a low hemoglobin, counterbalanced by an increasing oxygen extraction. The second patient was treated with volume replacement and inotropic support. The improvement of DO₂ by increasing heart frequency and cardiac output resulted in a higher ScvO₂ value. In the third patient the ScvO₂ decreased because of an increased oxygen consumption due to convulsions.

Although we are aware that these cases do not support in a scientific way the clinical utility of the ScvO₂ measurement, we think that these cases illustrate the effectiveness of the ScvO₂ measurement. These cases illustrate the different variables, that can affect ScvO₂. Consequently, the interpretation of abnormal values may be difficult in clinical practice. In addition, one has to consider that even normal ScvO₂ values do not exclude disturbances in oxygen transport in individual organs and impairments in tissue oxygen uptake as has been described in sepsis.

In these conclusive remarks we want to touch briefly on some ideas for further research in this field. Future research should be directed to simplifying the interpretation of ScvO₂ values, for example, by additional continuous monitoring of other variables that determine ScvO₂. It seems that continuous measurement of cardiac output has arrived in adult intensive care medicine. This monitoring tool is not ready yet for neonatal medicine but it is worth to observe closely the results of pending clinical trials. In the same direction is the measurement of the gastric intramucosal pH to evaluate tissue (an)aerobic

metabolism. Recently a device is produced suitable for neonatal care medicine.

The review on oxygen transport learns that much of the perinatal development of tissue microvasculature and mitochondria is unknown or only occasionally investigated. Future research should be directed to determine the influence of the development of microvasculature on oxygen uptake by the tissues. Since in newborns and especially in preterm infants vascular system is still in development, oxygen uptake in these infants could be affected by a still less extended tissue vasculature.

The fact that the perinatal period is characterized by a switch from low oxidative metabolism to efficient oxidative phosphorylation by a rapid proliferation and maturation of cellular mitochondria, has raised our interest. Is this switch controlled by an intrinsic cellular clock or triggered by the birth itself and what are the consequences of a preterm birth?

SvO_2 is a flow-weighted average of venous oxygen saturation values from all the tissues and reflects the combined sufficiency of all the variables that constitute DO_2 and VO_2 . In neonatal medicine $ScvO_2$ measurement in the right atrium together with SaO_2 by pulse oximetry is useful as a complementary monitor of tissue oxygen needs in newborn infants.

Hoofdstuk 9

Sameenvatting en Conclusies

metabolism. Recently a device is produced suitable for neonatal care medicine.

The review on oxygen transport learns that much of the perinatal development of tissue microvasculature and mitochondria is unknown or only occasionally investigated. Future research should be directed to determine the influence of the development of microvasculature on oxygen uptake by the tissues. Since in newborns and especially in preterm infants vascular system is still in development, oxygen uptake in these infants could be affected by a still less extended tissue vasculature.

The fact that the perinatal period is characterized by a switch from low oxidative metabolism to efficient oxidative phosphorylation by a rapid proliferation and maturation of cellular mitochondria, has raised our interest. Is this switch controlled by an intrinsic cellular clock or triggered by the birth itself and what are the consequences of a pattern birth?

$\dot{SvO}_2$ is a flow-weighted average of venous oxygen saturation values from all sites, and reflects the combined influence of all the variables that determine $\dot{Q}$ and $\dot{VO}_2$. In neonatal medicine best measurement of the right ventric together with $\dot{Q}_{\text{pul}}$ by pulse oximetry is vital in a comprehensive assessment of tissue oxygen needs in neonatal infants.

Hoofdstuk 9

Samenvatting en Conclusies

Samenvatting en Conclusies

In de inleiding van dit proefschrift is de stelling naar voren gebracht dat het uiteindelijke doel van intensieve medische zorg is om het menselijke organisme in stand te houden door de gasuitwisseling te ondersteunen, door het zuurstoftransport naar de weefsels te verbeteren en door het monitoren en bevorderen van de zuurstofopname door de perifere weefsels. Het is niet verwonderlijk, dat de gemengd veneuze zuurstofsaturatie (SvO_2) zoveel belangstelling heeft gewekt bij intensieve care artsen, omdat de SvO_2 een geïntegreerde waarde is, waarin zowel de gasuitwisseling, als het zuurstoftransport en de zuurstofopname in tegenwoordig zijn.

SvO_2 geeft namelijk het overblijvende zuurstofgehalte weer nadat de weefsels zuurstof hebben opgenomen. De waarde van de SvO_2 wordt dan ook bepaald door de arteriële zuurstofsaturatie, het hemoglobine, de cardiac output en door het zuurstofverbruik van de weefsels.

In de neonatale medische zorg zijn het monitoren van het zuurstofgehalte en het toedienen van zuurstof altijd belangrijke discussiepunten geweest. Omdat zowel een tekort als een overmaat aan zuurstof een schadelijk effect kunnen hebben, zijn vragen over de exacte zuurstofbehoefte en over het meten van deze zuurstofbehoefte nog steeds actueel en niet volledig beantwoord.

Ons onderzoek heeft zich gericht op de meting van SvO_2 en op de vraag of SvO_2 een antwoord kan geven naar de exacte zuurstofbehoefte van de weefsels. De belangrijkste vragen zijn in de inleiding van dit proefschrift geformuleerd en worden hier kort gememoreerd en de conclusies zullen vervolgens in de volgende paragrafen worden besproken.

1. Welke is de relatie tussen het zuurstofverbruik van de weefsels en de SvO_2 waarde?

2. Is deze relatie tussen het zuurstofverbruik van de weefsels en de SvO_2 waarde anders wanneer het tekort aan zuurstof op een andere manier tot stand komt?

Om de klinische mogelijkheden van de continue SvO_2 meting in de neonatale periode te onderzoeken, werden de volgende vragen behandeld.

3. Is het mogelijk om de SvO_2 continue te meten in een pasgeboren kind? Waar kan de SvO_2 gemeten worden? Zijn de SvO_2 waarden betrouwbaar en nauwkeurig?

4. Welke zijn de normale SvO_2 waarden in stabiele pasgeboren kinderen, die geen extra zuurstof toegediend krijgen. Welke is de relatie tussen de arteriële en de veneuze zuurstofsaturatie waarden?

5. Wat zijn de SvO_2 waarden tijdens de acute fase van de IRDS tijdens kunstmatige beademing en extra zuurstofvoorziening. Welke is de relatie tussen de arteriële en veneuze zuurstofsaturatie en afgeleide waarden als veneuze admixture en fractionele zuurstof extractie? Wat is het belang van SvO_2 in de zorg voor ernstig zieke pasgeboren kinderen?

1. SvO_2 en het zuurstofverbruik van de weefsels.

Wij hebben de relatie tussen de zuurstofverbruik van de weefsels en SvO_2 onderzocht in een acuut dierenmodel -het neonatale varken-, waarbij de zuurstof toediening stapsgewijze werd verminderd. Tijdens de geleidelijke vermindering van de zuurstof verminderden de arteriële en de veneuze zuurstofsaturatie in dezelfde mate, de fractionele zuurstof extractie nam toe en diensgevolge bleef het zuurstofverbruik van de weefsels gedurende lange tijd gelijk. Het verminderen van de SvO_2 en de toename in fractionele

zuurstof extractie zijn mechanismen om het zuurstofverbruik van de weefsels te stabiliseren. Andere mechanismen die een verminderd zuurstofaanbod kunnen compenseren zijn een verhoogde cardiac output en een redistributie van de bloeddorstrooming naar de verschillende (delen van) organen. Een redistributie van de bloeddorstrooming werd in onze experimenten niet gemeten. Maar de waarde van SvO_2 en de distributie van de bloeddorstrooming zijn gerelateerde waarden, omdat door een redistributie van bloed naar organen met een hoge extractie de gemiddelde zuurstof extractie zal toenemen en de SvO_2 dus zal afnemen.

Deze bevinding betekent dat een lage SvO_2 waarde op zich zelf niet een signaal van anaërobe stofwisseling is, maar een signaal dat een organisme een beroep doet op zijn compensatie mechanismen om het zuurstofverbruik gelijk te houden. Wanneer een meer verdergaande hypoxie wordt geïnduceerd, wordt de vermindering van SaO_2 niet meer in gelijke mate gevolgd door de SvO_2 , vermindert het zuurstofverbruik van de weefsels en ontstaat een anaërobe metabolisme.

Bij welke SvO_2 waarden het anaërobe metabolisme ontstaat is ook in ons onderzoek bepaald. Als de SvO_2 waarde beneden de 15 % kwam, verminderde het zuurstofverbruik van de weefsels en ontstond er een anaëroob metabolisme met lactaat acidose. Een SvO_2 waarde boven de 40 % sloot een anaëroob metabolisme uit. SvO_2 waarden tussen de 15 % en 40 % waren zelden geassocieerd met anaëroob metabolisme.

Enkele opmerkingen moeten gemaakt worden om bovenstaande bevindingen te nuanceren. Ten eerste moet het tijdsaspect van dit onderzoek in ogenschouw worden genomen. De bovenvermelde waarden zouden verschillend kunnen zijn in experimenten met een langere tijdsduur. Ten tweede werden de experimenten uitgevoerd in biggetjes onder narcose en met een kunstmatige beademing. Daardoor was er sprake van een constant en tamelijk laag zuurstofverbruik. In de klinische geneeskunde in patiënten met een meer variabel en hoger zuurstofverbruik zou de SvO_2 drempelwaarde voor anaëroob metabolisme kunnen veranderen. Ten derde moet men omzichtig zijn om de SvO_2 te gebruiken als een parameter exclusief voor het monitoren van één variabele, zoals het zuurstofverbruik van de weefsels. De waarde van SvO_2 is afhankelijk van veel verschillende variabelen. Ten vierde is de SvO_2 een begrip, gedefinieerd door de gemiddelde waarde van de veneuze saturaties van de verschillende organen en door de bloeddorstrooming door die verschillende organen. Dit betekent dat een "normale" SvO_2 waarde een afwijkende verhouding tussen het zuurstofaanbod en het zuurstofverbruik in één bepaald orgaan niet uitsluit. Ten slotte kan de relatie tussen SvO_2 en zuurstofverbruik nog complexer zijn in condities zoals sepsis en ARDS, als de opname van zuurstof door de weefsels is verminderd en lactaat acidose ontstaat samen met normale of zelfs hoge SvO_2 waarden.

Waarschijnlijk omwille van de bovenvermelde argumenten is men er nooit in geslaagd in prospectief klinisch onderzoek aan te tonen dat het klinisch handelen gericht op het verkrijgen van "normale" SvO_2 waarden de uitkomst van ernstig zieke volwassen patiënten heeft beïnvloed of verbeterd. Wel moet gezegd worden dat in deze trials patiënten met zeer complexe hemodynamische problemen bestudeerd werden. In ieder geval kunnen wat betreft de relatie tussen SvO_2 en zuurstofverbruik van de weefsels tijdens hypoxische hypoxie de volgende conclusies worden getrokken.

Tijdens hypoxische hypoxie zijn verlaagde SvO_2 waarden op zich zelf niet een signaal voor een verminderd zuurstofverbruik van de weefsels of voor een anaëroob metabolisme. Zo lang als een verlaging van de SaO_2 gepaard gaat met een evenredige verlaging van de SvO_2 kan een constant en stabiel zuurstofverbruik in stand worden gehouden. Bij een diepere hypoxische hypoxie zal het verschil tussen de SaO_2 en SvO_2 verminderen en het zuurstofverbruik van de weefsels zal lager worden. Daarom wordt de relatie tussen zuurstofverbruik en SvO_2 gekenmerkt door een lage kritische drempel waarde

van SvO_2 , beneden welke een anaëroob metabolisme ontstaat. Bovendien moet men rekening houden met bovengenoemde overwegingen en opmerkingen, als men de relatie tussen SvO_2 en zuurstofverbruik evalueert.

2. De relatie tussen zuurstofverbruik van de weefsels en SvO_2 vergeleken tijdens enerzijds hypoxische en anderzijds anemische hypoxie.

Een volgend onderzoek werd gedaan om de relatie tussen zuurstofverbruik van de weefsels en SvO_2 vast te stellen tijdens hypoxische en anemische hypoxie. Omdat verschillende soorten van hypoxie kunnen verschillen in circulatoire compensatie mechanismen, kan de opname van zuurstof door de weefsels verschillen en dus ook de SvO_2 . Ons belangrijkste resultaat was dat er hogere en meer variabele SvO_2 waarden tijdens de anemische hypoxie werden gevonden in vergelijking met de hypoxische hypoxie. De drempelwaarde van het zuurstofaanbod en van de zuurstof extractie, beneden welke een anaëroob metabolisme ontstaat, liet geen verschillen zien tussen de hypoxische en anemische hypoxie. Naar onze mening zijn de belangrijkste oorzaken voor deze bevinding *a)* het ten dienste staan van de volledige zuurstof dissociatie curve tijdens de anemische hypoxie, *b)* de relatieve toename van het plasma met zijn lage zuurstof oplosbaarheid tijdens de anemische hypoxie beperkt de mogelijkheid voor het transport van zuurstof van de erythrocyet naar de weefsels en *c)* de toegenomen snelheid van de bloeddorstroming tijdens anemische hypoxie met een kortere verblijfstijd in de weefsel capillairen vermindert de precapillaire zuurstof afgifte naar andere weefselvaten.

Samenvattend lijkt de relatie tussen SvO_2 en zuurstofverbruik van de weefsels nog complexer, omdat de SvO_2 drempelwaarde blijkbaar ook beïnvloed wordt door het soort hypoxie. Maar de conclusie kan getrokken worden, dat tijdens beide soorten hypoxie het zuurstofverbruik van de weefsels in stand gehouden kan worden, zolang als een bestendig verschil tussen de arteriële en het veneuze zuurstofgehalte kan worden gehandhaafd. Daarom werd een SvO_2 drempelwaarde, waar beneden het zuurstofverbruik van de weefsels verminderde, alleen gevonden bij relatief lage SvO_2 waarden.

3a De continue meting van SvO_2 in pasgeboren kinderen

Het voorstel om de SvO_2 te meten in de klinische zorg voor ernstig zieke pasgeboren kinderen was naar voren gebracht om de precieze zuurstofbehoefte van pasgeboren kinderen te kunnen bepalen. Men hoopte dan aandoeningen te verminderen samenhangend met de toxiciteit van zuurstof, zoals retinopathie van de prematuriteit, bronchopulmonale dysplasie en necrotiserende enterocolitis. Ook aandoeningen samenhangend met een tekort aan zuurstof, zoals cerebraal bepaalde spasticiteit en mentale schade hoopte men zo te kunnen voorkomen. Omdat de SvO_2 meting in de arteria pulmonalis in kleine pasgeboren kinderen moeilijk en gevaarlijk is, werden de SvO_2 metingen verricht in het rechteratrium (centraal veneuze zuurstofsaturatie; ScvO_2) met een fiberoptic catheter, die een continue meting van zuurstofsaturatie mogelijk maakt.

3b De toepasbaarheid van de fiberoptic catheter. De betrouwbaarheid en de nauwkeurigheid van de ScvO_2 waarden. De positie van de catheter.

Wij hebben de toepasbaarheid en de nauwkeurigheid van de continue ScvO_2 meting getest zoals die gerealiseerd werd door een 4 French fiberoptic catheter, die inge-

bracht was in het rechteratrium via de navelvene.

Ten eerste wat betreft de toepasbaarheid van deze catheter, hebben wij geconcludeerd dat het percentage met succes ingebrachte catheters en het aantal complicaties niet verschiden van de routinematig gebruikte catheters. In 67 % van de patiënten kon de fiberoptic catheter geplaatst worden in de juiste positie d.w.z. in het midden van het rechteratrium of op de overgang van de vena cava inferior en het rechteratrium. In een paar kinderen was het signaal aan het einde van de periode, dat de catheters in situ waren, verminderd door een lage reflectie aan de tip van de catheter en de weergave van de zuurstofsaturatie kon niet meer als betrouwbaar worden beschouwd. Ernstige complicaties zoals pericardiale effusies, cardiale tamponades, perforaties en aritmieën hebben zich niet voorgedaan.

Ten tweede wat betreft de nauwkeurigheid van de ScvO₂ weergave werd door ons geconcludeerd, dat de zuurstofsaturaties van het bloed in het rechteratrium van pasgeboren kinderen nauwkeurig en betrouwbaar werden weergegeven. Het gemiddelde verschil ($\pm$ SD) tussen de weergave van de fiberoptic catheter en de laboratorium hemoximeter zuurstofsaturatie waarden bedroeg -0.36 ± 7.64 . Deze waarde van het gemiddelde verschil en de standaard deviatie is vergelijkbaar met de gegevens uit de "volwassen" literatuur over de nauwkeurigheid van de fiberoptic oximetrie in de arteria pulmonalis.

De positie van de catheter is uitgebreid bediscussieerd in het literatuur overzicht inzake de oximetrie. Hierin werd geconcludeerd, dat de nadelen (niet volledige menging van het bloed, grotere gevoeligheid voor catheter bewegingen en reflectie van de atrium wand) waarschijnlijk even geldig zijn in de neonatologie als in de volwassen klinische zorg. Maar men moet zich bedenken, dat de meting van de gemengd veneuze zuurstofsaturatie (SvO₂) in de neonatologie alleen mogelijk is in het rechteratrium, omdat het gebruik van een arteria pulmonalis catheter in een (prematuur) pasgeboren kind moeilijk en gevaarlijk is. Wel dient men te anticiperen op de genoemde nadelen van de ScvO₂ meting in het rechteratrium door a) frequent een in vivo calibratie uit te voeren en b) door bedachtzaam te zijn bij het berekenen van zuurstoftransport variabelen gebaseerd op centraal veneuze zuurstofsaturatie metingen.

Enkele opmerkingen dienen nog gemaakt te worden over de beperkingen en nadelen van het gebruik van een "single" lumen navelvene catheter in de praktische neonatale intensieve zorg. Sinds kort is het in de neonatale intensieve zorg routine geworden om dubbel of zelfs triple lumen navelvene catheters te gebruiken. Het routinematig toepassen van deze catheters wordt gerechtvaardigd door het gelijktijdig en continue toedienen van medicijnen en parenterale voeding. Het gebruik maken van een "single" lumen catheter zoals de door ons gebruikte 4 Fr. catheter lijkt zich nu te beperken tot meer stabiele kinderen, in wie nu juist de meting van de veneuze saturatie niet direct noodzakelijk is. Een andere beperking van een navelvene catheter is het feit dat slechts in twee derde van de patiënten een catheter correct via de ductus venosus in het rechteratrium gepositioneerd kon worden.

Samenvattend zal men aan de ene kant met een zekere terughoudendheid de ScvO₂ meting in overweging nemen gezien de beperkingen van een "single" lumen catheter terwijl men aan de andere kant zich dient te realiseren dat de fiberoptic catheter betrouwbaar en nauwkeurig de ScvO₂ waarde van pasgeboren kinderen weergeeft.

4. De "normale" reeks van ScvO_2 waarden in stabiele pasgeboren kinderen zonder extra zuurstofbehoefte. De relatie tussen SaO_2 en ScvO_2 waarden. De waarde van de fractionele zuurstof extractie en de "shunt index".

We hebben in een prospectief klinisch observationeel onderzoek de reeks van ScvO_2 waarden beschreven van 10 stabiele pasgeboren kinderen die geen extra zuurstof (meer) nodig hadden. Hierbij werd een reeks ScvO_2 waarden gevonden van 60 % tot 83 % (5e en 95ste percentiel). Andere variabelen, die het zuurstoftransport kunnen beïnvloeden waren normaal: SaO_2 : > 86 %; hemoglobine: > 7.0 mmol/L; pH en bloeddruk: normaal. Deze reeks ScvO_2 waarden in stabiele niet zieke kinderen is gelijk aan de waarden beschreven bij volwassenen, hoewel het zuurstofverbruik bij pasgeborenen duidelijk hoger is dan bij volwassenen. Het hogere zuurstofverbruik wordt kennelijk gecompenseerd door een toegenomen zuurstofaanbod aan de weefsels. En inderdaad zowel de cardiac output als de hemoglobine concentratie zijn hoger in de neonatale periode in vergelijking met de volwassen leeftijd.

De relatie tussen ScvO_2 en SaO_2 werd eveneens onderzocht. In zeven patiënten vertoonden de ScvO_2 waarden een significante correlatie met de SaO_2 waarden. Dit toont aan dat, wanneer SaO_2 vermindert, ScvO_2 in evenredige mate vermindert, zodat een stabiel zuurstofverbruik in stand kan worden gehouden. In drie patiënten waren de SaO_2 en ScvO_2 niet met elkaar gecorreleerd. Speculerend kunnen wij aannemen dat het zuurstofverbruik bij deze kinderen in stand werd gehouden door een toename van de cardiac output, omdat in stabiele patiënten het zuurstofverbruik onafhankelijk is over een groot traject van het zuurstofaanbod.

Het zuurstofaanbod is gewoonlijk vier tot vijf keer zo groot als het zuurstofverbruik, zoals naar voren komt in de normale waarde van de fractionele zuurstof extractie van 0.20 tot 0.25. In ons onderzoek geanalyseerd voor iedere patiënt afzonderlijk werd een reeks (5e en 95ste percentiel) van waarden tussen de 0.13 en 0.33 werd gevonden. Er bestond een zeer goede correlatie tussen de fractionele zuurstof extractie en de ScvO_2 . Hierin wordt nogmaals het opmerkelijke belang van de ScvO_2 aangetoond als een monitor van de balans tussen het zuurstofaanbod en zuurstofverbruik.

Verder maakt de ScvO_2 het mogelijk om de shunt index te berekenen: de shunt index is een vereenvoudigde bepaling van de veneuze admixture. Wij vonden in deze 10 stabiele pasgeborenen een reeks waarden tussen 7 % en 53 %. Er was sprake van een duidelijke negatieve correlatie tussen de shunt index en SaO_2 in alle patiënten: in andere woorden bij lagere SaO_2 waarden was er sprake van een toegenomen shunt index.

5. De reeks van ScvO_2 waarden tijdens de acute fase van respiratoire problemen in pasgeboren kinderen met kunstmatige beademing en extra zuurstofbehoefte. De relatie tussen SaO_2 en ScvO_2 waarden. Fractionele zuurstof extractie en shunt index. Het belang van de ScvO_2 meting in de neonatale klinische zorg.

Wij hebben tot slot een observationele studie verricht in twaalf pasgeboren kinderen met kunstmatige beademing en extra zuurstof om de ScvO_2 waarden te bepalen op verschillende niveaus van oxygenatie gedefinieerd door de SaO_2 waarde: SaO_2 waarden onder 86 %, SaO_2 waarden tussen 86 en 90 % en SaO_2 waarden tussen 91 en 96 %. Deze laatste twee niveaus behoren, respectievelijk, tot het lage en hoge gedeelte van de in de literatuur aanbevolen reeks van SaO_2 waarden, zoals die gemeten wordt door de pols-oximetrie.

De ScvO_2 waarden in het lage en in het hoge gedeelte van de aanbevolen SaO_2

waarden komen overeen met de "normale" ScvO₂ waarden zoals deze door ons zijn beschreven in kinderen, die geen extra zuurstof nodig hadden. Lagere ScvO₂ waarden worden meestal gevonden in het laagste oxygenatie niveau bij SaO₂ waarden onder 86 %.

Een significante correlatie tussen SaO₂ en ScvO₂ werd in alle kinderen op één na gevonden. De helling van de regressielijnen vervaardigd voor ieder kind afzonderlijk toonde geen significant verschil tussen de twee gedeelten van de aanbevolen oxygenatie waarden. In het laagste oxygenatie niveau bij SaO₂ waarden onder de 86 % waren de ScvO₂ waarden duidelijk verminderd ten opzichte van het oxygenatie niveau met SaO₂ waarden tussen 86 % en 96 %. In twee patiënten was de mate van de ScvO₂ vermindering (de helling van de regressielijn) significant toegenomen in vergelijking met de ScvO₂ vermindering in de hogere oxygenatie niveaus. Blijkbaar was er in deze patiënten sprake van een toegenomen zuurstof extractie, terwijl andere compensatie mechanismen zoals cardiac output niet in staat waren om de verminderde zuurstofsaturatie te compenseren.

In ons onderzoek werd een toegenomen shunt index gevonden in het lagere gedeelte van de aanbevolen SaO₂ waarden. Bij SaO₂ waarden onder de 86 % was de shunt index verder toegenomen tot zelfs waarden van 60 % tot 80 %. Aangenomen moet worden dat er een zekere mate van een intrapulmonale shunting of een minder goed op elkaar afgestemd zijn van long perfusie en ventilatie moet bestaan om deze toegenomen veneuze admixture te kunnen verklaren in de lagere niveaus van oxygenatie. Aan de andere kant kan men zich afvragen of een berekend item als de shunt index werkelijk een betrouwbare afspiegeling is van klinische problemen, hoewel aan de voorwaarden en vooronderstellingen om de shunt index te gebruiken wel voldaan was.

Fractionele zuurstof extractie was, al was het in geringe mate ($r = 0.18$ tot 0.39) in zes patiënten significant gecorreleerd met SaO₂.

Samenvattend liet de "on line" monitoring van ScvO₂ zien dat ScvO₂ daalde bij afnemende niveaus van oxygenatie. De ScvO₂ waarden waren niet buiten de reeks van waarden, zoals die beschreven is bij kinderen, die geen extra zuurstof nodig hadden. Deze vermindering van ScvO₂ waarden lijkt het eerste compensatiemechanisme te zijn om het zuurstofverbruik van de weefsels in stand te houden. Betreffende de discussie over de alarmgrenzen van SaO₂ inzake de neonatale zuurstof therapie kan de meting van de ScvO₂ een nieuw element in deze discussie brengen: Een lagere grenswaarde voor SaO₂ zou eventueel geaccepteerd kunnen worden, wanneer gelijktijdige gemeten ScvO₂ waarden in evenredige mate verminderd zijn, en dus het zuurstofverbruik van de weefsels stabiel is gebleven. Men kan dan ook concluderen, dat het verschil tussen de SaO₂ en ScvO₂ waarden van groter belang lijkt te zijn dan de absolute waarde van ScvO₂ alleen.

Aangaande de vraag welke het klinisch belang is om de ScvO₂ waarde te kennen, zullen hier in het kort drie klinische casus worden gepresenteerd als voorbeelden dat de continue meting van ScvO₂ een waardevolle complementaire monitor van de oxygenatie kan zijn:

Casus 1. Prematuur kind; zwangerschapsduur: 28 weken; geboortegewicht: 890 gram. Diagnose: IRDS en E Coli sepsis, meningitis. Behandeling: positieve druk beademing; hoge frequentie beademing; intratracheale surfactant suppletie. Het klinisch beloop werd gecompliceerd door een intraventriculaire bloeding. De navelvene werd gecatheteriseerd met een fiberoptic catheter voor centraal veneuze toegang en de meting van de ScvO₂. De punt van de catheter was gelegen op de overgang van het rechteratrium naar de vena cava inferior. De zuurstofsaturatie weergave van de fiberoptic catheter kwam overeen met de waarden van de hemoximeter ($77.16 \% \pm 3.25$ vs $76.16 \% \pm 4.41$) Het hemoglobine daalde van 8.5 mmol/L tot 7.6 mmol/L en verder tot 6.8 mmol/L, waarschijnlijk vanwege veel bloedafnames. De SaO₂ waarden lieten geen enkele verandering zien, maar de ScvO₂

verminderde significant met een stijging van de fractionele zuurstof extractie.

Casus 2. Prematuur kind; zwangerschapsduur: 27 weken; geboortegewicht: 720 gram. De bevalling werd gecompliceerd door een matig ernstige asphyxie. De navelvene werd gecatheteriseerd met een fiberoptic catheter voor centraal veneuze toegang en de meting van de ScvO₂. Het klinisch beloop werd gecompliceerd door een respiratoire insufficiëntie en ernstige hypotensie. Positieve drukbeademing en intratracheale surfactant suppletie waren noodzakelijk. De bloeddruk werd ondersteund door middel van volume therapie en dopamine en dobutamine. Na de dopamine medicatie opgevoerd te hebben tot een dosering van 30 µg/kg/min nam de ScvO₂ toe van 70.5 % ± 5.0 tot 78.6 % ± 6.7 in tegenstelling tot de SaO₂ die zelfs iets verminderde van 91.8 % ± 3.4 tot 90.0 % ± 2.6.

Casus 3. Prematuur kind; zwangerschapsduur: 28 weken; geboortegewicht: 1350 gram. De bevalling werd gecompliceerd door een ernstige asphyxie. Arteriële navelstreng pH: 6.89. Onmiddellijk na de bevalling werd begonnen met kunstmatige beademing. Na stabilisatie werd de navelvene gecatheteriseerd met een fiberoptic catheter voor een centraal veneuze toegang en de meting van de ScvO₂. Het beloop liet een respiratoire verbetering zien, zodanig dat de kunstmatige beademing gestopt kon worden. Maar er ontstonden epileptische trekkingen die geassocieerd waren met een verhoging van de hartfrequentie en de bloeddruk en een kleine vermindering van de SaO₂. De ScvO₂ daalde gedurende deze episodes van 75 % tot 55 %. Antiepileptische medicatie werd begonnen en deze episodes deden zich niet meer voor.

Deze drie casus illustreren dat de continue meting van de ScvO₂ ons in staat stelt om veranderingen in de balans tussen het zuurstofaanbod en het zuurstofverbruik waar te nemen, terwijl SaO₂ geen grote veranderingen liet zien. De eerste patiënt laat een vermindering zien van het zuurstofgehalte in het arteriële bloed door een laag hemoglobine, hetgeen wordt gecompenseerd door een toegenomen zuurstof extractie. De tweede patiënt werd behandeld met een volumetherapie en inotropica ondersteuning. De verbetering van het zuurstofaanbod door een toegenomen cardiac output leidde tot een verhoogde ScvO₂ waarde. In de derde patiënt verminderde de ScvO₂ vanwege het toegenomen zuurstofverbruik door de epileptische convulsies.

Ofschoon wij ons realiseren dat deze casus niet op een wetenschappelijke manier de klinische bruikbaarheid van de ScvO₂ meting aantonen, denken we dat zij wel de effectiviteit illustreren van deze meting. Tevens laten deze casus de verschillende variabelen zien, die de waarde van ScvO₂ bepalen. De interpretatie van abnormale waarden kan daarom soms moeilijk zijn. Bovendien kunnen ook normale waarden stoornissen in het zuurstoftransport niet volledig uitsluiten bij voorbeeld in organen afzonderlijk of in het geval van sepsis.

In dit afsluitend hoofdstuk willen wij nog kort ingaan op enkele ideeën voor toekomstig onderzoek in deze richting. Toekomstig onderzoek kan gericht worden naar het vereenvoudigen van de interpretatie van de ScvO₂ waarden in de klinische praktijk. Bijvoorbeeld door het continue monitoren van ander variabelen, die de ScvO₂ meebepalen. Het lijkt erop, dat de continue meting van de cardiac output tot de mogelijkheden van de volwassen intensive care kan gaan behoren. Hoewel deze meting nog niet klaar is voor de neonatale intensieve care is het de moeite waard om de vooruitgang en de klinische trials in dezen goed in de gaten te houden. In dezelfde richting ligt de meting van de intramucosale pH meting in de maag voor het evalueren van het (an)aërobe weefsel metabolisme. Onlangs is er een maagsonde op de markt gekomen, die ook geschikt is voor kleine pasgeboren

kinderen.

Het literatuuroverzicht in dit proefschrift leerde ons dat er nog veel over de perinatale ontwikkeling van de microvasculatuur nog onbekend of onvoldoende onderzocht is. Ook de laat foetale en de perinatale ontwikkeling van het eind-orgaan van het zuurstoftransport, het mitochondrion is nog onvoldoende onderzocht. Het feit dat de perinatale ontwikkeling wordt gekarakteriseerd door een "switch" van een metabolisme met een lage zuurstofverbruik naar een efficiënte oxidatieve phosphorylatie en dat deze "switch" gepaard gaat met een snelle proliferatie en maturatie van de mitochondria heeft onze interesse gewekt. Wordt deze "switch" gecontroleerd door een intrinsieke klok in de cel of wordt hij getriggerd door de geboorte zelf. En welke zijn dan de consequenties bij een premature geboorte?

Ik dank ScvO₂ is gedefinieerd als de gemiddelde waarde van de veneuze saturaties afkomstig van alle organen, en is mede bepaald door de bloeddoorstroming door die verschillende organen. De ScvO₂ weerspiegelt alle factoren die het zuurstofaanbod en het zuurstofverbruik bepalen. In de neonatologie is de ScvO₂ meting in het rechteratrium samen met de SaO₂ meting door middel van de pols oximetrie bruikbaar als een aanvullende monitor van het zuurstoftransport bij pasgeboren kinderen.

Dankwoord

Carlos Blanco bedank ik voor zijn ideeën bij het opzetten en uitvoeren van het onderzoek. Zijn kritische kanttekeningen (letterlijk en figuurlijk) bij de manuscripten van de artikelen waren van grote waarde en bovendien erg leerzaam. Kortom, ik ben hem dankbaar voor het "leren onderzoek te doen". Ik ben er trots op door hem tot neonatoloog te zijn opgeleid en ik beschouw het als een voorrecht om met hem dit vak uit te kunnen oefenen.

Wiel Maertzdorf ken ik vanaf mijn sollicitatie in Maastricht. Hij is dan ook nadrukkelijk verbonden met mijn vorming tot kinderarts, neonatoloog en "import Maastrichtenaar". Ik ben hem dankbaar voor het meedenken, meediscussieren en meelesen met dit proefschrift.

Ik dank mijn collega's neonatologen Pieter Degraeuwe, jarenlang mijn kamergenoot, Danilo Gavilanes, Twan Mulder en Eduardo Villamor voor de goede en vriendelijke sfeer en voor de belangstelling naar de vorderingen van het boekje.

Ik dank de verpleegkundigen van de neonatologie afdeling E2 voor hun geïnteresseerde, soms wat ongelovige vragen naar het proefschrift. Ja, het is echt waar, ik ga promoveren op "die catheters".

Theo Hoorntje wil ik bedanken voor zijn hulp en betrokkenheid bij de biggetjesexperimenten.

Ik wil mijn ouders bedanken voor hun vertrouwen en ondersteuning tijdens de studie jaren. Ik ben blij, dat zij deze afsluiting van de studie kunnen meemaken.

Tot slot bedank ik Brigitte: aan haar en aan onze kinderen is dit boek opgedragen.

Curriculum Vitae

De schrijver van dit proefschrift werd geboren in 's-Gravenhage op 18 november 1954. In 1973 werd op het Aloysius college te 's-Gravenhage het Gymnasium β eindexamen behaald.

Van 1973 tot 1981 studeerde hij geneeskunde aan de Rijksuniversiteit Leiden. Het doctoraal examen werd behaald in 1980. Op 18 september 1981 behaalde hij zijn artsexamen.

Van 1982 tot 1986 volgde hij de opleiding kindergeneeskunde bij Prof.dr. LHJ. Ramaekers in het Annadal ziekenhuis in Maastricht. De datum van inschrijving in het register als kinderarts is 1 februari 1986. Hierna was hij werkzaam tot maart 1988 als chef de clinique in het De Weverziekenhuis en de St. Elisabeth kliniek, de vroedvrouwenschool, te Heerlen.

Van 1988 tot 1990 volgde hij de opleiding tot kinderarts-neonatoloog bij Prof.dr. CE. Blanco in het Academisch Ziekenhuis Maastricht.

Vanaf maart 1990 is hij werkzaam als kinderarts-neonatoloog bij de capaciteitsgroep kindergeneeskunde in het Academisch Ziekenhuis Maastricht.

Binnen de Universiteit Maastricht is hij werkzaam als blokcoördinator van het blok 2.6 (Geboren en getogen).

Hij is getrouwd met Brigitte van Wijk en zij hebben drie kinderen.