

Effects of carbohydrate (CHO) and fat supplementation on CHO metabolism during prolonged exercise.

Citation for published version (APA):

Jeukendrup, A. E., Saris, W. H. M., Brouns, F. J. P. H., Halliday, D., & Wagenmakers, A. J. M. (1996). Effects of carbohydrate (CHO) and fat supplementation on CHO metabolism during prolonged exercise. *Metabolism-Clinical and Experimental*, 45, 915-921. [https://doi.org/10.1016/S0026-0495\(96\)90169-9](https://doi.org/10.1016/S0026-0495(96)90169-9)

Document status and date:

Published: 01/01/1996

DOI:

[10.1016/S0026-0495\(96\)90169-9](https://doi.org/10.1016/S0026-0495(96)90169-9)

Document Version:

Publisher's PDF, also known as Version of record

Please check the document version of this publication:

- A submitted manuscript is the version of the article upon submission and before peer-review. There can be important differences between the submitted version and the official published version of record. People interested in the research are advised to contact the author for the final version of the publication, or visit the DOI to the publisher's website.
- The final author version and the galley proof are versions of the publication after peer review.
- The final published version features the final layout of the paper including the volume, issue and page numbers.

[Link to publication](#)

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal.

If the publication is distributed under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license above, please follow below link for the End User Agreement:

www.umlib.nl/taverne-license

Take down policy

If you believe that this document breaches copyright please contact us at:

repository@maastrichtuniversity.nl

providing details and we will investigate your claim.

Effects of Carbohydrate (CHO) and Fat Supplementation on CHO Metabolism During Prolonged Exercise

Asker E. Jeukendrup, Wim H.M. Saris, Fred Brouns, David Halliday, and Anton J.M. Wagenmakers

The aim of the study was to examine carbohydrate (CHO) utilization in subjects receiving CHO or CHO + medium-chain triglycerides (MCT) supplements during 180 minutes of exercise at 50% maximal aerobic work rate ($[W_{max}]$ 57% maximal oxygen consumption $[\dot{V}O_{2max}]$). In a double-blind crossover design, nine trained athletes cycled four times. Subjects received a bolus of $4 \text{ mL} \cdot \text{kg}^{-1}$ at the start and $2 \text{ mL} \cdot \text{kg}^{-1}$ every 20 minutes during exercise of either a $150\text{-g} \cdot \text{L}^{-1}$ CHO solution (CHO trial), an equicaloric 70 energy% (en%) CHO–30 en% MCT suspension containing 29 g MCT (CHO + MCT trial), or a $150\text{-g} \cdot \text{L}^{-1}$ CHO (high-CHO [HCHO]) solution plus 29 g MCT (HCHO + MCT trial). A fourth trial consisted of a ^{13}C -background control trial (CON). The four trials were randomized. Before and after the exercise bout, muscle biopsies were taken from the quadriceps muscle and muscle glycogen levels were determined. During exercise, breath samples were collected for estimation of exogenous and endogenous CHO oxidation. No significant differences were detected in glycogen breakdown among the trials (277 ± 14 $\text{mmol} \cdot \text{kg dry weight}^{-1}$ CHO, 249 ± 20 CHO + MCT, and 240 ± 18 HCHO + MCT) or in the respiratory exchange ratio during exercise. Mean exogenous CHO oxidation rates during the final hour of exercise were 0.79 , 0.63 , and $0.73 \text{ g} \cdot \text{min}^{-1}$, respectively. No differences were observed between the trials regarding exogenous or endogenous CHO oxidation. Plasma free fatty acid (FFA) concentrations were elevated during exercise to a level of approximately $500 \mu\text{mol} \cdot \text{L}^{-1}$ and were comparable in all trials, whereas plasma ketone concentrations significantly increased after MCT ingestion as compared with the CHO trial. It is concluded that 29 g MCT co-ingested with CHO during 180 minutes of exercise does not influence CHO utilization or glycogen breakdown.

Copyright © 1996 by W.B. Saunders Company

ORAL INGESTION of carbohydrate (CHO) during exercise has been shown to better maintain plasma glucose levels¹ and high rates of plasma glucose oxidation,² resulting in improved endurance capacity.¹ Oxidation rates of orally ingested CHO have never been observed to be higher than 1.0 to $1.1 \text{ g} \cdot \text{min}^{-1}$, which seems to be the maximum oxidation rate. Even ingestion rates of up to $2 \text{ g} \cdot \text{min}^{-1}$ during exercise at 70% maximal oxygen consumption ($\dot{V}O_{2max}$) resulted in oral glucose oxidation rates of no more than $1 \text{ g} \cdot \text{min}^{-1}$.³ To supply an additional exogenous energy source for the working muscle, substrates other than or in addition to CHO might be ingested. One potential candidate for such an energy source could be medium-chain triglycerides (MCT). MCT have been shown to be rapidly hydrolyzed, absorbed,⁴ and subsequently oxidized.^{5,6,6a} In addition, a preliminary study showed that MCT co-ingested with CHO do not inhibit gastric emptying.⁷ Moreover, it has been suggested that MCT ingestion may improve exercise performance by elevating plasma free fatty acid (FFA) levels and sparing muscle glycogen.⁸ Several studies have reported this relationship between increased availability of plasma FFA, reduced rate of muscle glycogen breakdown,⁹⁻¹³ and delayed onset of exhaustion.^{9,10,13}

However, increased supply and subsequent oxidation of FFA or ketone bodies have been shown to result in an inhibition of plasma glucose utilization,^{14,15} as well as overall CHO utilization. From these observations, one might hypothesize that increasing the FFA availability by oral fat intake may also limit the utilization of the exogenously supplied CHO.

The aim of the present study therefore was to investigate the effects of MCT co-ingested with CHO on exogenous and endogenous CHO utilization during prolonged exercise, and to determine what effect this regimen had on muscle glycogen breakdown. The CHO + MCT mixture was based on the CHO + MCT suspension that emptied

most rapidly from the stomach in a previous comparative study.⁷ The amount ingested was based on the maximal amount of MCT that could be ingested without causing gastrointestinal problems.

SUBJECTS AND METHODS

Subjects

Nine male trained triathletes or cyclists aged (mean \pm SEM) 26.0 ± 5.0 years with a weight of $75.7 \pm 4.3 \text{ kg}$, height $185 \pm 8 \text{ cm}$, maximal aerobic work rate (W_{max}) $5.48 \pm 0.24 \text{ W} \cdot \text{kg}^{-1}$, $\dot{V}O_{2max}$ $64.7 \pm 2.3 \text{ mL} \cdot \text{kg}^{-1}$, and maximal heart rate $194 \pm 5.3 \text{ bpm}$ participated in this study. The nature and the risks of the experimental procedures were explained to the subjects, and their written informed consent was obtained. The study was approved by the local medical ethics committee.

Pretrials

W_{max} was measured on an electronically braked ergometer (Lode Excalibur, Groningen, The Netherlands) during an incremental exhaustive exercise test¹⁶ 1 week before the first experimental trial. The results of this initial test were used to determine 50% W_{max} , which was later used in the experimental trials.

Experimental Trials

Nine subjects performed four trials, each separated by at least 7 days. A trial consisted of 180 minutes of cycling at 50% W_{max} ($57\% \pm 2\% \dot{V}O_{2max}$). Drinks were given in a randomized order

From the Department of Human Biology, Nutrition Research Centre, University of Limburg, Maastricht, The Netherlands.

Submitted December 12, 1995; accepted January 18, 1996.

Supported by an Isostar Research Grant from Sandoz Nutrition, Berne, Switzerland.

Address reprint requests to Asker E. Jeukendrup, Department of Human Biology, University of Limburg, PO Box 616, 6200 MD Maastricht, The Netherlands.

Copyright © 1996 by W.B. Saunders Company

0026-0495/96/4507-002/\$03.00/0

and double-blind; all drinks were vanilla-flavored (Sandoz Nutrition, Berne, Switzerland). Subjects abstained from training and were instructed to consume a similar diet for the 3 days before each trial. In addition, they were instructed not to consume any products with a high natural abundance of ^{13}C during the entire experimental period.

Protocol

Subjects reported to the laboratory at 8:00 AM after an overnight fast, and a standardized breakfast of two crackers with cheese was provided (14 g CHO, 4 g fat, and 6 g protein). A Teflon catheter (Baxter Quick Cath; Uden, The Netherlands) was inserted into an antecubital vein, and at 8:30 AM a resting blood sample was drawn. Also, a muscle biopsy was taken from the lateral part of the vastus lateralis. Resting breath gases were collected for measurement of oxygen consumption (2900 analyzer; SensorMedics, Anaheim, CA), and vacutainer tubes were filled directly from the mixing chamber in duplicate to determine the $^{13}\text{C}/^{12}\text{C}$ ratio in expired CO_2 . At 8:50 AM, subjects started cycling for 10 minutes at 100 W as a warm-up. At 9:00 AM, exercise intensity was increased to 50% \dot{V}_{max} for 180 minutes. Blood samples were drawn at 30-minute intervals until the end of exercise. Expiratory gases were collected every 15 minutes. Two subjects were tested on the same day, starting the protocol 10 minutes apart. Directly after the exercise bout, a second muscle biopsy was taken 2 cm proximal to the first biopsy.

Drinks

Subjects received a bolus of $4 \text{ mL} \cdot \text{kg}^{-1}$ at the start ($t = 0$) and $2 \text{ mL} \cdot \text{kg}^{-1}$ every 20 minutes during exercise of either a 15% CHO solution (CHO) or an equicaloric CHO + MCT suspension ([CHO + MCT] 70 energy% [en%] as CHO [149 g/180 min] and 30 en% as MCT [29 g/180 min]). To study the effect of MCT added to the CHO instead of the equicaloric CHO + MCT suspension, a third trial (high-CHO [HCHO] + MCT) was included in which a suspension was ingested containing the same amount of CHO (214 g/180 min) as in the CHO trial and the same amount of MCT (29 g/180 min) as in the CHO + MCT and MCT trials. Therefore, it could be investigated whether differences between the CHO + MCT trial and the CHO trial are due to the MCT or to the differential amount of CHO. The CHO in these trials were corn-derived long-chain glucose polymers of high natural ^{13}C abundance (-11.31δ per mil ν PDB, 0.0111101 $^{13}\text{C}/^{12}\text{C}$ ratio). To enable correction for possible shifts in background ^{13}C enrichment during exercise, a fourth trial was included in which tapioca-derived long-chain glucose polymers of low ^{13}C natural abundance (-26.12δ per mil ν PDB, 0.0109437 $^{13}\text{C}/^{12}\text{C}$ ratio; Sandoz Nutrition; Berne, Switzerland) were ingested (CON). This CON trial was used only for ^{13}C -background measurements; no other measurements are presented.

MCT contained fatty acids with a chain length of C8 (Estasan GT8-99; Unichema, Barcelona, Spain) and had a ^{13}C enrichment of -29.81δ per mil versus PDB (0.0109222 $^{13}\text{C}/^{12}\text{C}$ ratio). NaCl ($20 \text{ mmol} \cdot \text{L}^{-1}$) was added to all drinks. Meal temperature was kept constant at 20°C .

Analysis

Blood (10 mL) was collected into EDTA-containing tubes and centrifuged for 4 minutes. Aliquots of plasma were frozen immediately in liquid nitrogen and stored at -40°C until analyses of glucose (Uni Kit III, 0710970; La Roche, Basel, Switzerland), lactate,¹⁷ β -hydroxybutyrate,¹⁸ FFA (Wako FFA-C test kit; Wako Chemicals, Neuss, Germany), and glycerol (GPO-trinder 337; Sigma, St Louis, MO), which were performed with the COBAS BIO analyzer (La Roche). Muscle biopsies were freeze-dried, and glycogen content was assayed spectrophotometrically after hydrolysis with HCl.¹⁹ Glycogen concentration was expressed as millimoles of glycosyl units per kilogram dry weight of tissue. Total energy expenditure and oxidation rates of total fat, total CHO, and exogenous MCT were calculated from indirect calorimetry (respiratory quotient and $\dot{V}\text{O}_2$) and stable-isotope measurements ($^{13}\text{CO}_2/\text{CO}_2$) (GC continuous-flow IRMS; Finnigan MAT 252, Bremen, Germany). Enrichments of the substrates (in drinks) were measured with an elemental analyzer-IRMS combination (Carlo-Erba on-line [EA 1108CHN; Fisons, Milan, Italy] connected to the Finnigan MAT 252).

Calculations

CHO and fat oxidation rates were calculated from $\dot{V}\text{CO}_2$ and $\dot{V}\text{O}_2$ using the nonprotein respiratory quotient²⁰: CHO oxidation = $4.585\dot{V}\text{CO}_2 - 3.226\dot{V}\text{O}_2$, and fat oxidation = $1.695\dot{V}\text{O}_2 - 1.701\dot{V}\text{CO}_2$. Isotopic enrichment of expired air was expressed as the delta per mil difference between the $^{13}\text{C}/^{12}\text{C}$ ratio of the sample and a known laboratory reference standard according to the formula, $\delta^{13}\text{C}$ (per mil) = $([^{13}\text{C}/^{12}\text{C} \text{ sample}]/[^{13}\text{C}/^{12}\text{C} \text{ standard}] - 1) \times 10^3$. The $\delta^{13}\text{C}$ was then related to the international standard, Pee Dee Bellemnitella (PDB-I). The amount of CHO oxidized was calculated according to the formula, exogenous CHO oxidation = $\dot{V}\text{CO}_2 \cdot (\delta \text{ ref} - \delta \text{ exp}) / (\delta \text{ ref} - \delta \text{ ing}) \cdot 1/k$, in which $\delta \text{ ref}$ is the ^{13}C enrichment of expired air in the reference test CON (background), $\delta \text{ exp}$ is the ^{13}C enrichment of expired CO_2 during exercise with CHO(+MCT) ingestion at different time points, $\delta \text{ ing}$ is the ^{13}C enrichment of the CHO in the ingested CHO + MCT suspension, and k is the amount of CO_2 (in liters) produced via oxidation of 1 gram glucose ($k = 0.7466 \text{ L CO}_2/\text{g glucose}$).

In the present study and in previous studies from our laboratory,^{3,5,21,22} it was shown that instructing the subjects not to eat any products of high natural ^{13}C abundance during the experimental period was effective in reducing the background shift (change in $^{13}\text{CO}_2$) from endogenous substrate stores.²² However, although the

Table 1. $\dot{V}\text{O}_2$ and Respiratory Exchange Ratio (mean \pm SEM) During 3 Hours of Cycling Exercise at 57% $\dot{V}\text{O}_{2\text{max}}$ with ingestion of CHO, CHO + MCT, or HCHO + MCT

	Time (min)					
	30	60	90	120	150	180
$\dot{V}\text{O}_2$ ($\text{L} \cdot \text{min}^{-1}$)						
CHO	2.78 \pm 0.08	2.79 \pm 0.10	2.79 \pm 0.10	2.77 \pm 0.10	2.80 \pm 0.08	2.75 \pm 0.10
CHO + MCT	2.76 \pm 0.08	2.76 \pm 0.09	2.79 \pm 0.07	2.76 \pm 0.08	2.77 \pm 0.09	2.73 \pm 0.09
HCHO + MCT	2.77 \pm 0.07	2.80 \pm 0.07	2.80 \pm 0.08	2.84 \pm 0.09	2.73 \pm 0.08	2.80 \pm 0.09
Respiratory exchange ratio						
CHO	0.89 \pm 0.01	0.88 \pm 0.01	0.87 \pm 0.01	0.88 \pm 0.01	0.87 \pm 0.01	0.86 \pm 0.02
CHO + MCT	0.86 \pm 0.02	0.87 \pm 0.01	0.86 \pm 0.01	0.86 \pm 0.01	0.87 \pm 0.01	0.86 \pm 0.01
HCHO + MCT	0.87 \pm 0.02	0.88 \pm 0.01	0.87 \pm 0.01	0.86 \pm 0.01	0.87 \pm 0.01	0.88 \pm 0.01

NOTE. No significant differences were observed among the trials or over time.




Fig 1. Breath ¹³CO₂ enrichment during exercise in (●) CHO, (□) CHO + MCT, and (△) HCHO + MCT trials and during the (■) CON trial used for background correction (mean ± SEM, n = 6).

background shift was small in the present study, background correction was made by using the ¹³C enrichment of breath samples in the CON trial. We assumed that the low ¹³C abundance of MCT was similar enough to the ¹³C abundance of endogenous substrate stores, so that the observed change in ¹³CO₂ production can be attributed to the CHO and not to the MCT. To check whether this was a valid assumption, we performed a pilot study in which subjects underwent the same protocol described herein while ingesting CHO or CHO + MCT beverages containing tapioca-derived CHO (-26.12 δ per mil v PDB). Any difference between the two trials could be attributed to the lower ¹³C abundance of the MCT (-29.81 δ per mil v PDB). However, the ¹³C enrichment of expired CO₂ was identical in both conditions at all time points, and thus, we concluded that the low ¹³C abundance of MCT would not influence the calculated exogenous CHO oxidation rates in the present study.

Another methodological consideration when using ¹³CO₂ in expired gases to calculate exogenous substrate oxidation is the trapping of exogenous ¹³CO₂ in the bicarbonate pool, a very large




Fig 2. Exogenous MCT oxidation rates over 180 minutes of exercise (mean ± SEM, n = 6). Symbols are as in Fig 1.

Table 2. Oxidation of Endogenous and Exogenous Substrates (in grams) Over the 60- to 120-minute and 120- to 180-Minute Periods (mean ± SEM)

Interval	CHO	CHO + MCT	HCHO + MCT
60-120			
CHO	132.2 ± 11.4	122.5 ± 10.3	127.3 ± 10.3
Exogenous	44.2 ± 3.7	34.2 ± 3.7	38.6 ± 5.0
Endogenous	88.0 ± 8.3	88.3 ± 7.3	88.7 ± 9.3
Fat	34.3 ± 3.7	37.5 ± 3.7	36.7 ± 3.2
120-180			
CHO	124.1 ± 12.8	119.6 ± 9.8	130.8 ± 9.9
Exogenous	50.3 ± 3.2	41.5 ± 5.0	48.5 ± 4.8
Endogenous	73.8 ± 9.9	78.1 ± 6.5	82.3 ± 9.3
Fat	33.8 ± 4.2	38.7 ± 3.0	35.1 ± 3.1

NOTE. No significant differences in substrate utilization were observed among the trials or time periods.

and slowly exchanging pool in which some CO₂ arising from decarboxylation of energy substrates is temporarily trapped.²³ However, during exercise, CO₂ production increases severalfold so that a physiological steady-state situation will occur and ¹³CO₂ in expired air will be rapidly equilibrated with the ¹³CO₂/H¹³CO₃⁻ pool. The dilution of ¹³CO₂ becomes negligible and recovery of ¹³CO₂ approaches 100% after 60 minutes of exercise.³⁸ Therefore, in the present study, data from the initial 60 minutes were not used for calculation of exogenous MCT oxidation.

Gastrointestinal Discomfort

A questionnaire assessing gastrointestinal discomfort was provided after each exercise test. Subjects had to score the following items on a scale from 1 to 5 (1 = not at all to 5 = very severe): nausea, intestinal cramps, belching, vomiting, diarrhea, flatulence, stomach ache, abdominal pressure, and eructation.

Statistics

ANOVA for repeated measures was used to compare differences in substrate utilization and in blood-related parameters




Fig 3. (□) Preexercise and (▤) postexercise glycogen concentrations and (■) glycogen breakdown (decrease) for the CHO, CHO + MCT, and HCHO + MCT trials. No significant differences were observed among the trials (mean ± SEM, n = 6).




Fig 4. Plasma (A) FFA, (B) glycerol, and (C) β-hydroxybutyrate concentrations during exercise for the CHO, CHO + MCT, and HCHO + MCT trials. *Significant difference ($P < .05$) between CHO and CHO + MCT (mean \pm SEM, $n = 6$). Symbols are as in Fig 1.

among the CHO, CHO + MCT, and HCHO + MCT trials. A Scheffé post hoc test was used in the event of a significant ($P < .05$) F ratio. All results are expressed as the mean \pm SEM.

RESULTS

Breath Analysis

Oxygen consumption (Table 1) during exercise was relatively constant, with exercise intensity of the subjects maintained at close to 57% $\dot{V}O_2$ max and not significantly different among the trials. The respiratory exchange ratio also was not significantly different among the trials.

The mean ¹³C enrichment of the resting breath samples

was -26.36 ± 0.25 δ per mil versus PDB (0.0109410 ¹³C/¹²C ratio). Changes in the isotopic composition of expired CO₂ in response to exercise are depicted in Fig 1. During CON (with ingestion of CHO of low ¹³C natural abundance), there was a small nonsignificant increase of ¹³C in the expired air (0.2 to 0.6 δ per mil ν PDB). In the CHO(+MCT) trials, the increase in ¹³C was significant, reaching a difference of 3 to 4 δ per mil versus PDB toward the end of 180 minutes' exercise (compared with resting breath sample; Fig 1). Exogenous CHO oxidation increased during the first hour and leveled off during the final 90 minutes (Fig 2). At the end of the exercise bout, exogenous CHO oxidation

rates were 0.89, 0.73, and 0.91 g · min⁻¹ for CHO, CHO + MCT, and HCHO + MCT, respectively. Mean oxidation rates over the 60- to 180-minute period were 0.79, 0.63, and 0.73 g · min⁻¹, respectively. Exogenous CHO oxidation tended to be slightly lower in the CHO + MCT trial than in the CHO trial. However, this difference did not reach statistical significance. It was estimated that during the 60- to 180-minute period in the CHO trial, 94.5 g exogenous CHO was oxidized, versus 75.7 g in CHO + MCT and 87.1 g in HCHO + MCT. The amounts of CHO (exogenous and endogenous) and fat oxidation during the second and third hour of exercise are presented in Table 2). Energy expenditure was comparable in all trials.

Muscle Glycogen

Preexercise muscle glycogen concentrations were comparable among the trials: 516 ± 32, 518 ± 30, and 505 ± 26 mmol glucose · kg dry weight (dw)⁻¹ for CHO, CHO + MCT, and HCHO + MCT, respectively. Muscle glycogen levels decreased significantly in the three trials: 277 ± 14 mmol glucose · kg dw⁻¹ for CHO, 249 ± 20 for CHO + MCT, and 240 ± 18 for HCHO + MCT. Glycogen breakdown was not significantly different among the three trials (Fig 3).

Plasma Variables

Compared with rest, plasma FFA levels were decreased after 30 minutes of exercise and were significantly elevated during the final 30 minutes of exercise (Fig 4A). No differences in plasma FFA levels were observed among the trials. There was a significant increase in plasma glycerol levels after 30 minutes in all trials, but no differences were observed among the three trials (Fig 4B). Plasma β-hydroxybutyrate concentrations increased to approximately 500 μmol · L⁻¹ during the first 30 minutes in the CHO + MCT and HCHO + MCT trials; thereafter, concentrations remained stable (Fig 4C). In the CHO trial, no changes in β-hydroxybutyrate were observed throughout exercise. Plasma glucose levels were maintained during exercise (Fig 5A), whereas plasma lactate concentrations tended to decrease as compared with the resting value (Fig 5B). No differences were observed in plasma glucose or lactate concentrations between the trials.

Gastrointestinal Discomfort

Subjects reported some gastrointestinal discomfort in all tests. The occurrence and severity (mean score, 2.8) of the complaints were not different among the trials. Most frequently reported were intestinal cramps, nausea, and belching.

DISCUSSION

Muscle Glycogen and Plasma FFA

It has been reported in several studies that increased availability of plasma FFA resulted in muscle glycogen sparing^{9-13,24} and hence increased performance.¹⁰ In humans, plasma FFA levels have been elevated by injecting




Fig 5. Plasma (A) glucose and (B) lactate concentrations during exercise for the CHO, CHO + MCT, and HCHO + MCT trials. *Significant difference between CHO and CHO + MCT ($P < .05$). @Significant difference between CHO and HCHO + MCT ($P < .05$) (mean ± SEM, $n = 6$).

heparin, which stimulates lipoprotein lipase activity, after feeding subjects a high-fat meal (long-chain fatty acids) or infusing a triglyceride emulsion (Intralipid).^{9,13,24} In these studies, a glycogen-sparing effect was observed when FFA availability was high. Although the infusion of fats and consumption of triglycerides in combination with an injection of heparin are interesting approaches to test the

interaction between CHO and fat metabolism during exercise, this method has little practical value. It has therefore been suggested that ingestion of MCT will increase FFA availability.^{5,6,8,25-28} MCT are delivered into the blood more rapidly than ingested long-chain triglycerides,^{4,29,30} and can cross the mitochondrial membrane without carnitine.³¹ Therefore, it has been argued that MCT might be a readily available energy source for the working muscle. In addition, it has been suggested that MCT might spare muscle glycogen and improve time-trial cycling performance.⁸ In the present study and in a recent study with a similar experimental protocol,⁵ although the maximally tolerable amount of MCT was ingested, it did not affect plasma FFA levels. Hence, no effect on muscle glycogen utilization was observed. This is in agreement with the only other available study in which muscle glycogen concentrations were determined after MCT ingestion.²⁶ In it, MCT were ingested 1 hour before exercise and did not result in glycogen-sparing during 1 hour of exercise at 60% $\dot{V}O_2$ max. It may be that the amount of MCT provided was too small to influence plasma FFA concentrations, or that medium-chain fatty acids are oxidized rapidly in the liver and/or in skeletal muscle so that plasma FFA concentration remains the same.

Plasma Glycerol and Ketone Bodies

Increased plasma levels of glycerol and ketone bodies have been frequently observed after MCT feeding.^{5,6,25-28} In the present study, glycerol concentrations were not significantly elevated after MCT ingestion. Plasma β -hydroxybutyrate was elevated to moderate levels (400 to 500 mmol \cdot L⁻¹) after MCT ingestion. This may indicate that part of the MCT are metabolized in the liver, resulting in the production of ketone bodies, while the glycerol from hydrolysis of MCT is rapidly utilized for gluconeogenesis.

Substrate Utilization

We found that MCT ingestion did not affect total CHO or fat utilization. Peak oxidation rates of the oral ingested

long-chain glucose polymer were 0.80 to 0.94 g \cdot min⁻¹, which is in line with previous studies.^{3,21,32-37} Peak oxidation rates of glucose, glucose polymers, and starch have been found to be between 0.8 and 1.0 g \cdot min⁻¹ with comparable ingestion rate, feeding schedule, and exercise intensity.^{2,3,21} The oxidation rate of orally ingested CHO seemed to be slightly lower (although not significantly) in the CHO + MCT trial as compared with CHO and HCHO + MCT. The reason for this small difference may be that the amount of CHO ingested was lower (149 v 214 g). As shown previously,^{3,37} the oxidation rate of orally ingested CHO did not increase to the same magnitude as the amount of CHO ingested. The rate of exogenous CHO oxidation seems to be limited by a yet-undetermined factor. The ingested CHO contributed 15% to 25% to energy expenditure during the 120- to 180-minute period. The contribution of MCT to energy expenditure has been shown to be approximately 7% during the same period (120 to 180 minutes) with similar experimental conditions.⁵

However, according to the literature, the amount of MCT ingested in the present study (29 g) is reported to be the maximal amount of MCT that can be ingested without causing gastrointestinal problems.²⁷

Gastrointestinal Discomfort

The MCT (29 g) seemed to have no influence on palatability of the beverages. However, since in all tests some such discomfort was reported, this may be attributed to the long-chain glucose polymers that were co-ingested with the MCT. Ivy et al²⁷ reported that administration of 30 g MCT in combination with cereal caused some minor distress in 10% of the subjects.

In summary, 29 g MCT co-ingested with CHO during 180 minutes of exercise at 57% $\dot{V}O_2$ max does not influence exogenous or endogenous CHO utilization or muscle glycogen breakdown.

REFERENCES

1. Coyle EF, Coggan AR, Hemmert MK, et al: Muscle glycogen utilization during prolonged strenuous exercise when fed carbohydrate. *J Appl Physiol* 61:165-172, 1986
2. Bosch AN, Dennis SC, Noakes TD: Influence of carbohydrate ingestion on fuel substrate turnover and oxidation during prolonged exercise. *J Appl Physiol* 76:2364-2372, 1994
3. Wagenmakers AJM, Brouns F, Saris WHM, et al: Oxidation rates of orally ingested carbohydrates during prolonged exercise in man. *J Appl Physiol* 75:2774-2780, 1993
4. Bach AC, Babayan VK: Medium-chain triglycerides: An update. *Am J Clin Nutr* 36:950-962, 1982
5. Jeukendrup AE, Saris WHM, Schrauwen P, et al: Metabolic availability of medium chain triglycerides co-ingested with carbohydrates during prolonged exercise. *J Appl Physiol* 79:756-762, 1995
6. Massicotte D, Péronnet F, Brisson GR, et al: Oxidation of exogenous medium-chain free fatty acids during prolonged exercise—Comparison with glucose. *J Appl Physiol* 73:1334-1339, 1992
- 6a. Jeukendrup AE, Saris WHM, Van Diesen R, et al: Effect of endogenous carbohydrate availability on oral medium chain triglyceride oxidation during prolonged exercise. *J Appl Physiol* 80:949-954, 1996
7. Beckers EJ, Jeukendrup AE, Brouns F, et al: Gastric emptying of carbohydrate-medium chain triglyceride suspensions at rest. *Int J Sports Med* 13:581-584, 1992
8. Van Zeyl C, Lambert EV, Noakes TD, et al: Effects of medium-chain triglyceride ingestion on carbohydrate metabolism and cycling performance. *Clin Sci* 87:30, 1994 (abstr)
9. Costill DL, Coyle E, Dalsky G, et al: Effects of elevated plasma FFA and insulin on muscle glycogen usage during exercise. *J Appl Physiol* 43:695-699, 1977
10. Hickson RC, Rennie MJ, Conlee RK, et al: Effect of increased plasma fatty acids on glycogen utilization and endurance. *J Appl Physiol* 43:829-833, 1977
11. Issekutz B, Miller HI, Rodahl K: Lipid and carbohydrate metabolism during exercise. *Fed Proc* 25:1415-1420, 1966
12. Rennie MJ, Winder WW: A sparing effect of increased plasma fatty acids on muscle and liver glycogen content in the exercising rat. *Biochem J* 156:647-655, 1976

13. Vukovich MD, Costill DL, Hickey MS, et al: Effect of fat emulsion infusion and fat feeding on muscle glycogen utilization during cycle exercise. *J Appl Physiol* 75:1513-1518, 1993
14. Hargreaves M, Kiens B, Richter EA: Effect of increased plasma free fatty acid concentrations on muscle metabolism in exercising men. *J Appl Physiol* 70:194-201, 1991
15. Randle PJ, Hales CN, Garland PB, et al: The glucose-fatty acid cycle: Its role in insulin sensitivity and the metabolic disturbances of diabetes mellitus. *Lancet* 1:785-789, 1963
16. Kuipers H, Verstappen FTJ, Keizer HA, et al: Variability of aerobic performance in the laboratory and its physiologic correlates. *Int J Sports Med* 6:197-201, 1985
17. Gutmann I, Wahlefeld AW: L-(+)-Lactate determination with lactate dehydrogenase and NAD, in Bergmeyer HU: *Methods of Enzymatic Analysis* (ed 2). New York, NY, Academic, 1974, pp 1464-1468
18. Moore JJ, Marcus M, Sax SM: Kinetic assay of β -hydroxybutyrate in plasma with COBAS BIO centrifugal analyzer. *Clin Chem* 73:1334-1339, 1982
19. Passenneau JV, Lauderdalet VR: A comparison of three methods of glycogen measurement in tissue. *Anal Biochem* 60:404-412, 1974
20. Péronnet F, Massicotte D: Table of nonprotein respiratory quotient: An update. *Can J Sport Sci* 16:23-29, 1991
21. Saris WHM, Goodpaster BH, Jeukendrup AE, et al: Exogenous carbohydrate oxidation from different carbohydrate sources during exercise. *J Appl Physiol* 75:2168-2172, 1993
22. Wagenmakers AJM, Rehrer NJ, Brouns F, et al: Breath $^{13}\text{CO}_2$ background enrichment at rest and during exercise: Diet related differences between Europe and America. *J Appl Physiol* 74:2353-2357, 1993
23. Robert JJ, Koziat J, Chauvet D, et al: Use of ^{13}C -labeled glucose for estimating glucose oxidation: Some design considerations. *J Appl Physiol* 63:1725-1732, 1987
24. Dyck DJ, Putman CT, Heigenhauser GJF, et al: Regulation of fat-carbohydrate interaction in skeletal muscle during intense aerobic cycling. *Am J Physiol* 265:E852-E859, 1993
25. Auclair E, Satabin P, Servan E, et al: Metabolic effects of glucose, medium chain triglyceride and long chain triglyceride feeding before prolonged exercise in rats. *Eur J Appl Physiol* 57:126-131, 1988
26. Decombaz J, Arnaud M-J, Milon H, et al: Energy metabolism of medium chain triglycerides versus carbohydrate during exercise. *Eur J Appl Physiol* 52:9-14, 1983
27. Ivy JL, Costill DL, Fink WJ, et al: Contribution of medium and long chain triglyceride intake to energy metabolism during prolonged exercise. *Int J Sports Med* 1:15-20, 1980
28. Satabin P, Portero P, Defer G, et al: Metabolic and hormonal responses to lipid and carbohydrate diets during exercise in man. *Med Sci Sports Exerc* 19:218-223, 1987
29. Greenberger NJ, Skillman TG: Medium-chain triglycerides: Physiologic considerations and clinical implications. *N Engl J Med* 280:1045-1058, 1969
30. Metges CC, Wolfram G: Medium- and long-chain triglycerides labeled with ^{13}C : A comparison of oxidation after oral or parenteral administration in humans. *J Nutr* 121:31-36, 1991
31. Bremer J: Carnitine-metabolism and functions. *Physiol Rev* 63:1420-1479, 1983
32. Hawley JA, Dennis SC, Nowitz A, et al: Exogenous carbohydrate oxidation from maltose and glucose ingested during prolonged exercise. *Eur J Appl Physiol* 64:523-527, 1992
33. Leijssen DPC, Saris WHM, Jeukendrup AE, et al: Oxidation of orally ingested [^{13}C]-glucose and [^{13}C]-glucose during exercise. *J Appl Physiol* 79:720-725, 1995
34. Massicotte D, Péronnet F, Allah C, et al: Metabolic response to [^{13}C]-glucose and [^{13}C]-fructose ingestion during exercise. *J Appl Physiol* 61:1180-1184, 1986
35. Moodley D, Noakes TD, Bosch AN, et al: Oxidation of exogenous carbohydrate during prolonged exercise: The effects of the carbohydrate type and its concentration. *Eur J Appl Physiol* 64:328-334, 1992
36. Pallikarakis N, Jandrain B, Pirnay F, et al: Remarkable metabolic availability of oral glucose during long-duration exercise in humans. *J Appl Physiol* 60:1035-1042, 1986
37. Rehrer NJ, Wagenmakers AJM, Beckers EJ, et al: Gastric emptying, absorption and carbohydrate oxidation during prolonged exercise. *J Appl Physiol* 72:468-475, 1992
38. Pallikarakis N, Sphiris N, Lefèbvre P: Influence of the bicarbonate pool on the occurrence of $^{13}\text{CO}_2$ in exhaled air. *Eur J Appl Physiol* 63:179-183, 1991