Determination of heparin in blood

Citation for published version (APA):

Hemker, C. H., & Beguin, S. (1990). Determination of heparin in blood. *Medicinski razgledi*, 29(Suppl. 1), 95-110.

Document status and date:

Published: 01/01/1990

Document Version:

Publisher's PDF, also known as Version of record

Please check the document version of this publication:

- A submitted manuscript is the version of the article upon submission and before peer-review. There can be important differences between the submitted version and the official published version of record. People interested in the research are advised to contact the author for the final version of the publication, or visit the DOI to the publisher's website.
- The final author version and the galley proof are versions of the publication after peer review.
- The final published version features the final layout of the paper including the volume, issue and page numbers.

[Link to publication](#)

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal.

If the publication is distributed under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license above, please follow below link for the End User Agreement:

www.umlib.nl/taverne-license

Take down policy

If you believe that this document breaches copyright please contact us at:

repository@maastrichtuniversity.nl

providing details and we will investigate your claim.

DETERMINATION OF HEPARIN IN BLOOD

C. H. Hemker and S. Béguin

INTRODUCTION

Probably the most important antithrombotic drug in use at the moment is heparin (1-10). Actually the annual sale in the Netherlands is 6.10^{10} U, that is 2000 U per head of the population. Its administration by injection and the fact that the prescribing doctors are well aware of the inherent risk of bleeding (11) make that it is not likely to be used inadvertently. The consumption in the Netherlands is probably the minimum that can be used in any medical community that takes the prevention and treatment of thrombosis seriously because especially in this country the use of the only serious alternative - i.e. oral anticoagulation - is very well organized and hence very common (12,13).

In recent years we have witnessed the advent of low molecular weight heparins (LMWH), there are indications that they may be more convenient drugs than unfractionated heparin (UFH) is (14-23). This promises that heparin usage will still increase. In fact the discussion that is going on on LMWH already seems to bring about a rise in the use of UFH.

This situation makes that the question of how to determine heparin in the plasma of a patient becomes more and more urgent. Logically another question should precede the question of *how* to determine heparin, and that is *why* to determine heparin. Is it necessary to determine heparin levels (or a related parameter) or will giving standard doses do the job? That fundamental question can be answered only from controlled clinical trials and for such trials an adequate heparin test must be available. This is a circular argument. Therefore the question of what makes a good heparin test may well be the one to start with.

CURRENT TYPES OF HEPARIN TESTS

All current determinations of heparin are based on its influence on the blood coagulation process i.e. its enhancing effect on the inactivation of activated clotting factors by antithrombin III (AT III). Several other actions of heparin have been described however. It binds to the vessel wall (24-27), influences fibrinolysis (28-32), causes the release of lipoprotein lipase (33-38) etc. We keep thinking that its main therapeutic action is on blood coagulation for a simple reason: vitamin K antagonists also have a large spectrum of actions apart from that on coagulation. Yet coagulation is the only one that it has in common with heparin. It seems more likely than not that it is indeed this common denominator that has the antithrombotic effect. Until proof of the contrary we therefore

think that a test of the coagulation system is the most adequate to reflect the therapeutic action of heparin.

What are the current coagulation based tests of heparin activity? They come in several kinds:

1. **Thrombin time (TT)** (39-42). These tests are based upon the clotting of plasma by a fixed amount of added thrombin.
2. **Activated partial thromboplastin time (aPTT)** (41-50). Based upon starting coagulation in the patient's plasma via the intrinsic pathway.
3. **Heptest** (39,51,52). A clotting time started with activated factor X among other things.
4. **Anti-thrombin tests** (39, 53-59, 62). An amount of thrombin is added and the test measures the amount that remains active after a fixed incubation time.
5. **Anti-factor Xa tests** (39,55-62). Like 4, but now with factor Xa rather than with thrombin.
6. **Neutralisation tests** (63, 64).

Under 6 we group a number of more specific and more precise tests that are not in routine use, often because they are relatively time consuming or require techniques not readily available in the clinical laboratory. The neutralization tests, based upon titration of the heparin in the sample with a heparin neutraliser such as protamine are mainly used for determining precisely the high heparin concentrations met in extracorporeal circulation. These will not be discussed further because we focus on tests for heparin treatment and prevention here.

Tests 1, 2 and 3 are clotting tests; 4 and 5 are spectrophotometric tests based on the use of thrombin- or factor Xa-specific chromogenic substrates. Tests 1 and 4 measure the anti-thrombin effect, tests 3 and 5 the anti-factor Xa effect. The way in which heparin influences the aPTT has until recently remained relatively unclear. Especially the reason why the aPTT is sensitive to heparin whereas the PT (Quick-time) is not, remained obscure. We demonstrated that the aPTT is primarily dependent upon the time necessary for trace amounts of thrombin to activate so much factor VIII that a burst of thrombin formation ensues (65,66). This positive feedback effect of thrombin on its own formation is the main feature of the kinetics of the aPTT. Heparin prolongs the aPTT by enhancing the action of AT III on the trace amounts of thrombin that act as a feedback activator. Once the burst of massive thrombin generation occurs coagulation follows immediately, largely independent of the *amount* of thrombin that is formed. In the PT the feedback activation of factor VIII does not play a role, as would be expected, but, possibly because of the large excess of factor Xa and the availability of factor V (not bound to von Willebrand factor), the feedback of thrombin on factor V does not play a role either. Heparin will decrease the peak amount of thrombin formed in both aPTT and PT. That the PT is insensitive to heparin illustrates the fact that decreasing the peak *amount* does not influence the *clotting time* (see later).

The Hestest is a kind of aPTT that is started by the addition of activated factor X to the sample. If factor Xa is rapidly broken down, as in the presence of heparin (either UFH or LMWH) this test will prolong. On the other hand it also is sensitive to thrombin inactivation. The first effect is more important though and the Hestest therefore may be regarded as "the poor man's anti Xa" (64).

For several years it has been thought that the antithrombin action of heparin was responsible for bleeding whereas the anti-Xa action was thought to reflect antithrombotic power (67-69). More recent research very much doubts whether this is the case (15, 70-71). In fact it could be demonstrated that UFH in the concentration range used for prophylaxis or treatment of thrombosis has only a very small effect on the conversion velocity of prothrombin in clotting plasma (65, 66, 72, 73) while its antithrombotic action is not questioned. On the other hand a trial on a LMWH that had a high anti-Xa action had to be stopped because of the many bleeding complications in the treated group (21).

PLATELETS AND HEPARIN

Before discussing this point any further it is necessary to consider the important effect of blood platelets on heparin. Whereas the inverse, the effect of heparin on blood platelets is often the subject of discussion, without for that matter leading to a very clear picture (75-84), the effect of platelets on heparin is straightforward and clearcut (65, 85). Blood platelets, when activated, shed heparin neutralizing material such as platelet factor 4 (PF4), a protein with strong antiheparin properties (86-92). The 3.10^8 pL of normal platelet rich plasma (PRP) can neutralize about 0.3 U/ml of UFH. Normal venepuncture will always cause some platelet activation, especially when carried out under routine clinical conditions and even more so when vacuum tubes are used. This means that important amounts of PF4 may be shed so that between 0 and 0.4 U/ml of heparin will escape attention! Collecting blood on a platelet inhibitor is a partial solution to this problem. Unless this is done any exact heparin determination is illusory and 0.18 $\pm$ 0.07 U/ml will be neutralised in the sample (93). Even when collecting on platelet inhibitor the difference between heparin in a patient's sample and spiked heparin remains, be it to a lesser extent (94,95).

The very fact that platelet activation is an important event in haemostasis and thrombosis makes that the PF4 susceptibility of a heparin will be one of the factors that determine its *in vivo* action. If thrombin generation in PRP is triggered by small amounts of tissue thromboplastin, a test system is obtained that is one step nearer to the *in vivo* situation than tests in platelet poor plasma (PPP) are. Under these circumstances a burst of thrombin formation is observed about five minutes after starting the reaction. In the presence of UFH this burst is dose-dependently postponed but the amount of thrombin formed is not diminished. This is because trace amounts of thrombin that are generated during the lag-phase activate the platelets. Once the platelets are activated they provide

procoagulant phospholipids and factor V which allow a massive increase of thrombin formation. The situation is completely comparable to the feedback activation of factor VIII discussed above but even more outspoken. The amount of thrombin available during the lagphase is less in the presence of heparin and hence the lagphase will prolong. Once the platelets are activated they shed so much PF4 that the heparin is neutralized and the thrombin then formed is not or hardly inhibited at heparin concentrations up to 0.4 U/ml. None of the current tests takes this effect into account. This is the more regrettable because here an important difference between UFH and LMWH shows up. LMW Heparins are much less susceptible to PF4 inactivation than UFH and each of them to a different extent (96,97).

HOW TO CHOOSE A HEPARIN TEST?

For a good heparin test there is one minimal requirement: it should indicate the amount of active material in the circulation of the patient. Indeed there need not even be a relation between the property measured and the property that carries the pharmacological effect, a correlation will do. If heparins would strongly adsorb at a convenient wavelength or carry detectable radioactivity, then these properties could be used for measuring heparin concentrations in plasma. Even if the molecules that carry the signal would not be the same as those carrying the pharmacological effect.

On the other hand one dreams of the ideal test, which does not inform us about the amount of material in the blood but about the effect of that material on the relevant properties of the blood. In other words: the test that would not be a concentration-indicator but a true model of the therapeutic efficacy.

It looks as if, in the heparin field, we do have an acceptable test of the first type, whereas we are far from having one of the second type. This is all the more regrettable because heparins are notoriously heterogeneous. This makes that any property that we measure and that has not been proven to indicate therapeutic efficacy directly, may fool us as soon as the proportion of the indicator property to the effector property changes.

There are at this moment good reasons to believe that anti-factor Xa activity is an acceptable indicator for concentrations of circulating heparin. In many clinical studies it has been shown to correlate with the effect, at least when LMWHs are compared and within the rather broad limits that are set by clinical dose finding studies (98,104). From animal studies one has good grounds to doubt whether it also holds for a purely anti-factor Xa heparin like the Choay pentasaccharide (105).

Particularly interesting is the study of Cadroy in baboons that presents a model of the arterial platelet thrombus and that compares the dose-effect curves of UFH and of the very low molecular weight heparin CY222 in that model (106). The results of this study have been interpreted as to indicate the practical usefulness of the anti-thrombin test (107). This seems indeed a sound conclusion. Concluding however on basis of these tests that anti-thrombin measurements or even weight would be less useful seems precocious.

The matter can be conveniently discussed in terms of IC₅₀ concentrations (108). When anti-factor Xa activity is used to compare the IC₅₀'s of both heparins, then they appear to be much closer than when they are expressed in terms of either weight or anti-thrombin activity (table 1). This is the core of reasoning of ref. (107).

Table 1. Comparison of IC₅₀ values of two different heparins.

	UFH	CY222	Ratio
IC ₅₀ (μg/mL)	6.25	33	0.2
Anti Xa (U/mL)	1	1.4	0.7
Anti IIa(U/mL)	1	0.15	6.7
High affinity mat.(%)	36	10,	
IC ₅₀ H.A.mat (mg/mL)	2.25	3.3	0.7
Decay constant at IC ₅₀			
Factor Xa	37.5	26.4	1.4
Factor IIa	81.2	29.7	2.7

This certainly proves the practicability of the anti-factor Xa activity test. Does it however settle the argument in disfavour of either weight or anti-thrombin activity tests? Let us discuss weight first. What substance should be weighed? Obviously the active substance. Even though it has been described that non-AT III binding heparins contribute to the antithrombotic activity (109), one is rather tempted to think that the amount of AT III binding material is the important thing. We determined, by fluorescence-titration of pure AT III the percentage high affinity material in UFH and CY 222. If the IC₅₀s are calculated in weight of AT III binding material, it appears that the difference between UFH and CY222 is mainly due to a difference in content of high affinity material (table 1). Thus the apparently strong case against weight as a basis of comparison seems to weaken. Below we will argue that also the anti-thrombin activity may not be as easily dismissed as would be thought on basis of a quick interpretation of Cadroy's results.

Nevertheless one must admit that the presently available anti-thrombin tests are less sensitive than the anti-test, especially in the case of LMWHs. It remains an interesting question whether its sensitivity could be increased by rigorous prevention of platelet activation in the samples.

The aPTT test measures principally the effect of heparin on the feedback activation of factor VIII by thrombin and on factor IXa inactivation during the lag phase of thrombin formation (65,66,110). It is an indirect test of antithrombin action that is more sensitive than the direct test is because it takes advantage of the amplifying effect of the feedback mechanism. The main drawback is the large individual variation. It has been shown that spiking individual patient plasmas with the same amount of heparin results in very different prolongation ratios of the aPTT. This is probably due to the fact that the test is sensitive to the levels of all clotting factors, all heparin binding proteins and also to pathological inhibitors (111-116). Also different types of the test (as to contact activator used and phospholipid preparation) will give different outcomes.

We conclude that the anti-factor Xa activity test seems to fulfill the minimal requirements for a heparin determination. In comparing it to other tests, it must be said in its favour, that it seems to be more sensitive and less subject to inter patient variation than the trombin time and the aPTT, especially with LMWHs. The Hepptest seems to hold an intermediary position here.

HOW TO FIND A MODEL FOR CLINICAL EFFICACY?

After having argued that the anti-factor Xa activity test at this moment is probably the best available practical shortcut to heparin measurement, we will now show that it can by no means be accepted as a model for clinical efficacy. This may seem a superfluous remark because it never has been claimed that it would be. It is well known however that any test that is in general use gets a respectability of its own that might make it slip stealthily in the position of the accepted standard (117).

We found a method that enabled us to measure the velocity of prothrombin conversion in clotting plasma, independant of the simultaneous thrombin inactivation. Surprisingly CY222 and other substances that show a high conventional anti-factor Xa activity, do not show a parallel inhibition of prothrombin conversion in the extrinsic system. This is an unexpected result because the only physiological effect of factor Xa is prothrombin conversion. The reason is that factor Xa is produced during coagulation in such excess that it can be inhibited quite markedly before any effect on prothrombinase activity can be observed. In the intrinsic system heparin does inhibit prothrombin conversion, but this must be attributed to a secondary effect on the feedback activation of factor VIII by thrombin and to inhibition of factor IXa (64,65,110).

The main effect of heparins, be it fractionated or unfractionated, is inactivation of thrombin. The only exception are very low molecular weight products like pentasaccharide and certain subfractions of the usual LMWHs (64,96,97).

If indeed anti-thrombin activity is the main action, then why do the IC50s of that property not compare? (table 1). The reason is probably that the anti-thrombin activity and the anti-factor Xa activity as they are routinely measured bear only a distant relationship to the same properties in clotting plasma. In the study of Cadroy the anti-thrombin activity was measured by the aPTT and by the TT. These can not be accepted as exact quantitative estimations of the anti-thrombin effect (see above). Neither is the anti-factor Xa activity as measured routinely a good indicator of what happens to factor Xa in plasma. It has been shown that the decay of factor Xa in plasma varies both with the species from which the factor Xa is prepared and with the concentration of Ca^{++} ; moreover these variations are not the same with different types of heparin. It therefore will make an enormous difference whether bovine factor Xa decay is measured in the absence of Ca^{++} , as in the anti-Xa test, or human Xa in the presence of Ca^{++} , as in vivo.

We measured the specific decay constants of standard UFH and of CY222 in human plasma with human factor Xa and at physiological Ca^{++} concentration. Then we calcu-

lated what the effect of the IC₅₀s of Cadroy would be in PPP on the decay of thrombin and factor Xa. We see that the almost tenfold difference between anti-factor Xa activity and anti-thrombin activity of the original data tend to diminish to a bare twofold (table 1). We are conscious that we are applying data from human plasma to the *in vivo* situation in baboons, but it is the best approximation that we can offer for the moment.

THE SEARCH FOR IDEAL TEST

We must conclude that the ideal test for anticoagulant treatment does not yet exist. We therefore should try and make one. Such a test should meet a number of stiff demands, it should:

- a.* Be equally sensitive to comparable therapeutic levels of oral anticoagulation, and of treatment by heparin or heparinlike drugs.
- b.* Be sensitive to the inhibition of factor Xa (or any other clotting factor (notably factor Xa)) only insofar as this factor influences thrombin generation.
- c.* Be either not hampered by platelet activation (PPPtest) or take physiological platelet activation into account (PRPtest).

There is only one procedure that, as far as we can see, approaches these requirements. It has been in practical use for many decades and is often tacitly assumed to be the golden standard for effects on the coagulation system. We mean the thrombin generation test (TGT). It is indeed influenced by all anticoagulation procedures. It reflects inhibition of individual clotting factors, but not proportionally to their percentual variation, except for prothrombin (117). It can be carried out equally well in PRP as in PPP.

The thrombin generation curve of course has a mayor drawback: it takes an experienced laboratory, much time and a lot of reagent. This makes that until now it has practically been used for research purposes only.

In order to develop a routine test on this basis one should first decide what type of value one wants to measure. A classical TGT shows a number of different features. One can measure the lagtime of thrombin formation, the peak value of thrombin obtained or the surface under the curve. The lagtime is equivalent to a clotting time, because clotting is invariably seen at the beginning of the thrombin burst. In our experience the surface and the peakvalue correlate strongly. There is a theoretical argument to favour the surface under the TGT as an essential variable: thrombin is an enzyme. One nanomolar for ten minutes will in principle have the same effect as ten nanomolar for one minute. This means that the product of time and thrombin concentration is important, which is equivalent to the surface under the curve (118). As the next section will show, we developed a method that directly renders this surface from an easily automated test.

A CANDIDATE: THE THROMBIN POTENTIAL AND HOW TO MEASURE IT

As discussed above, it is a common property of all antithrombotic drugs that they diminish the amount of thrombin generated in clotting plasma, either because there are less clotting factors present (oral anticoagulation) or because the thrombin formed disappears faster (heparins etc.). Whether heparins also inhibit (the formation of) prothrombinase and inhibit thrombin formation also in that way is of no importance for the present purpose. If they do this effect is also measured by the procedure that we are going to discuss. Under antithrombotic treatment as described above the form of the thrombin generation curve changes in one or more of the following respects: it starts later, i.e. the lagphase before explosive thrombin formation occurs is longer, the peak is lower or the decay is faster so that the thrombin present is inactivated earlier. It has been shown that a lower generation (oral anticoagulation) as well as a faster decay (heparin treatment) cause efficacious antithrombotic treatment. Both result in a smaller surface under the thrombin generation curve. This surface we define as the endogenous thrombin potential (ETP) of the plasma. We here present the outlines of a method to determine the ETP and we think that the ETP thus measured can, among other things, serve to measure the effect of any known anticoagulant drug: heparins and heparin likes, hirudin and oral anticoagulation.

In the assay that we propose a test sample of blood, platelet rich plasma, platelet poor plasma or defibrinated plasma is added to a solution containing a trigger of the coagulation process, a source of Ca^{++} ions, a preparation of a natural antithrombin and a substrate of thrombin. The natural antithrombin may be antithrombin III or heparin cofactor II or both. They serve the purpose of rendering the test less sensitive to variations in the concentration of these inhibitors in the test sample and to minimise the side reaction with other antithrombins such as α_2 -Macroglobulin. By selecting either ATIII or HCII the test can be made specifically sensitive to drugs of the heparin, resp. of the dermatan sulfate type. It is not even excluded that under certain circumstances it may not be absolutely necessary to add extra inhibitors to the reaction medium. What is essential to the present test is that a substrate is used that is not exhausted during the lifetime of the thrombin that generates and disappears in the clotting plasma. The mixture with the test sample is incubated during a specified time under specified conditions. Then the amount of thrombin substrate is read directly or, in opaque solutions the reaction is stopped by adding a thrombin inhibitor and the mixture is treated, e.g. by centrifugation in order to allow determination of the amount of product formed or the amount of substrate that has disappeared. Also is it possible in optically clear solutions to monitor spectrophotometrically the amount of product formed. This is of particular interest because the first derivative of that curve will represent the course of thrombin generation in the sample.

The following reactions further illustrate the present method:

- Prothrombinase
2. Prothrombin —————> Thrombin
3. Thrombin + Antithrombins ———> Inactive complexes
- Thrombin
4. Indicator Agent —————> Signal Molecule

Reactions 2 and 3 go to completion so that thrombin is only transitorily present in the reaction mixture. During its presence it partakes in reaction 4, so that the extent of the conversion of the indicator agent reflects the time during which- and the amount with which, thrombin has catalysed that reaction. It is essential that the indicator agent is not exhausted before the thrombin has disappeared. The reaction velocity should be proportional to the concentration of thrombin at any moment, i.e. the concentration of substrate has to be several times higher than K_m .

In a test set up according to this principle a variety of different triggers may be used: thromboplastins of any origin for investigating the ETP of the extrinsic pathway, contact activators etc.

In performing the assay a great variation of protein concentrations, incubation times, coagulation triggers, reagent concentrations and temperatures may be used. These parameters may be varied according to the special purpose that the determination of the ETP may serve e.g. determination of low molecular weight heparins, monitoring of oral anticoagulation, detection of hyper-coagulable states etc.

The end level of substrate conversion is not necessarily completely fixed under all circumstances, because the complex of thrombin to α_2 -Macroglobulin has a persistent amidolytic activity. It may therefore be necessary to measure the optical density at a fixed moment after starting the reaction, or, if the reaction mixture is not optically clear, such as with the use of whole blood, to stop the reaction with a quenching solution and subject the reaction mixture to manipulations (such as centrifugation or precipitation) that allow measurement of the amount of the signal molecule present. Any chemical that will stop the action of thrombin on the thrombin substrate can in principle serve this purpose. Benzamidine, hirudin and α -NAPAP are obvious candidates.

The indicator agent can be any molecule capable of reacting with thrombin. In such reaction products should be formed which produce a measurable signal. P-nitroaniline, easily detectable photometrically at 405 nm is an obvious choice but other indicator agents may be used, such as those producing fluorescent or electrochemical signals.

REFERENCES

1. Gallus AS, Hirsh J, Tuttle RJ, Trebilcock R, O'Brien SE, Carrol JJ, Minden JH, Hudecki SM. Small Subcutaneous Doses of Heparin in Prevention of Venous Thrombosis. *N Engl J Med* 1973; 288: 545.

2. Gallus AS, Hirsh J, O'Brien SE. Prevention of Venous Thrombosis with Small Subcutaneous Doses of Heparin. *JAMA* 1975; 235: 980.
3. Gallus AS, Hirsh J. Treatment of Venous Thromboembolic Disease. *Semin Thromb Hemost* 1976; 2: 291.
4. Gordon-Smith IC, Quense-Le LP, Gundy DJ, Newcomber JF. Controlled-trial of two Regimes of Subcutaneous heparin in prevention of post-operative deep vein thrombosis. *Lancet* 1972; 1: 1133.
5. Manucci PM, Citterio LA, Panajotopolus M. Low-dose Heparin and Deep-vein Thrombosis after Total Hip Replacement. *Thromb Haemostas* 1976; 36: 157.
6. Rosenberg IL, Evans M, Pollock AV. Prophylaxis of Post-operative Leg Vein Thrombosis by Low-dose Subcutaneous Heparin or Preoperative Calf Muscle Stimulation: A Controlled Clinical Trial. *Br Med J* 1975; 1: 649.
7. Salzman EW, Hirsh J. Prevention of Venous Thromboembolism. In: Colman RW, Hirsh J, Marder VJ, Salzman EW, eds. *Hemostasis and Thrombosis: Basic Principles and Clinical Practice*. 2nd edition. Philadelphia: Lippincott, 1987: 1252-65.
8. Wessler S. Medical Management of Venous Thrombosis. *Annu Rev Med* 1987; 27: 313-9.
9. Basu D, Gallus A, Hirsh J, Cade J. A Prospective Study of the Value of Monitoring Heparin Therapy with the Activated Partial Thromboplastin Time. *N Engl J Med* 1972; 287: 324-7.
10. Adar R, Salzman EW. Treatment of Thrombosis of Veins of the Low Extremities. *N Engl J Med* 1975; 292: 348.
11. Kelton JG, Hirsh J. Bleeding Associated with Antithrombotic Therapy. *Semin Hematol* 1980; 17: 259-379.
12. Dijk-van-Wierda CA, Loeliger EA, Meilof J, Roos J. Anticoagulant Control in the Netherlands. *Lancet* 1981; 1321-2.
13. Loeliger EA, Hensen A. Grundlagen erfolgversprechender Langzeitbehandlung mit Cumarin. In: *Langzeitbehandlung mit Antikoagulantien VI. Hamburger Symposium über Blutgerinnung* mai 1963. Kelton JG, Hirsh J. Bleeding Associated with Antithrombotic Therapy. *Semin Hematol* 1980; 17: 259-379.
14. Bergquist D, Burmark US, Frisell J, Hallboeek T, Lindblad B, Risberg B, Toerngren S, Wallin G. Prospective double blind comparison between Fragmin and conventional low-dose heparin: Thromboprophylactic effect and bleeding complications. *Haemostasis* 1986; 16: Suppl 2: 11-8.
15. Carter CJ, Kelton JG, Hirsh J, Cerskus A, Santos AV, Gent M. The relationship between the hemorrhagic and antithrombotic properties of low molecular weight heparin in rabbits. *Blood* 1982; 59(6): 1239-45.
16. Cifonelli JA. The relationship between molecular weight and sulfate content and distribution to the anticoagulant activity of heparin preparations. *Carbohydr Res* 1974; 37: 145-54.
17. Dahan R, Houlbert D, Caulin C, Cuzin E, Viltart C, Woler M, Segrestaq JM. Prevention of deep vein thrombosis in elderly medical in-patients by a low molecular weight heparin: a randomized double blind trial. *Haemostasis* 1986; 16: 159-61.
18. Hirsh J, Ofosu F, Buchanan M. Rationale behind the development of low molecular weight heparin derivatives. *Semin Thromb Hemost* 1985; 11(1): 13-6.
19. Kakkar VV, Murray MJG. Efficacy and Safety of Low Molecular Weight Heparin (CY216) in Preventing Post-Operative Venous Thromboembolism: A Cooperative study. *Br J Surg* 1985; 72: 786-91.

20. Ofosu FA, Modi GJ, Blajchman MA, Buchanan MR, Johnson EA. Increased Sulfation Improves the Ability of Vessel Wall Glycosaminoglycans to Regulate Thrombin Activity and Prothrombin Activation in Plasma. *Thromb Haemostas* 1990. In Press.
21. Schmitz-Huebner U, Bunte H, Freise G, Reers B, Ruschemeyer C, Scherer R, Schulte H, Loo van de J. Clinical efficacy of low molecular weight heparin in postoperative thrombosis prophylaxis. *Klin Wochenschr* 1984; 62: 349-53.
22. Thomas DP, Merton RE, Barrowcliffe TW, Thunberg L, Lindahl U. Effects of Heparin Oligosaccharides with High Affinity for Antithrombin III in Experimental Venous Thrombosis. *Thromb Haemostas* 1982; 47: 244-8.
23. Turpie AGG, Levine MN, Hirsh J, Carter CJ, Jay RM, Powers PJ, Andrew M, Hull RD, Gent M. A randomized controlled trial of a low-molecular-weight heparin (enoxaparin) to prevent deep-vein thrombosis in patients undergoing elective hip surgery. *New Eng J Med* 1986; 315: 925-9.
24. Rijn-van JLML, Trillou M, Mardiguian J, Tobelem G, Caen J. Selective Binding of Heparins to Human Endothelial Cells. Implications for Pharmacokinetics. *Thromb Res* 1987; 45: 211-22.
25. Homandberg GA, Kramer-Bjerke J, Grant D, Christianson G, Eisenstein R. Heparin-binding fragments of fibronectin are potent inhibitors of endothelial cell growth: structure-function correlations. *Biochim Biophys Acta* 1986; 874(1): 61-71.
26. Barzu T, Molho P, Tobelem G, Petitou M, Caen J. Binding and Endocytosis of Heparin by Human Endothelial Cells in Culture. *Biochim Biophys Acta* 1985; 845: 196-203.
27. Barzu T, Rijn-van JLML, Petitou M, Molho P, Tobelem G, Caen J. Endothelial Binding Sites for Heparin: Specificity, Role in Heparin Neutralization. *Biochem J* 1986. In press.
28. Halse Th. Aktivierung der Fibrinolyse und Thrombolyse durch Polysaccharidschwefelsäureester (Heparin, Heparinoide). *Arzneim Forsch* 1962; 12: 574.
29. Klocking HP, Markwardt F. The influence of heparin and heparinoids on the fibrinolytic system. 3rd Danube symposium on thrombosis and haemostasis. Bratislava, June 1983.
30. Gaffney PJ, Marsh NA, Thomas DP. The influence of heparin and heparin-like substances on the fibrinolytic system in vivo. 7th International Congress on thrombosis. Valencia, October 1982.
31. Gaffney PJ. Fibrinolytic enhancement by heparin-related material. 8th International Congress on thrombosis. Istanbul, June 1984.
32. Vinazzer H, Stemberger A, Haas S, Blumel G. Influence of Heparin ; of Different Heparin Fractions and of a Low Molecular Weight Heparin-like Substance on the Mechanism of Fibrinolysis. *Thromb Res* 1982; 27: 341-52.
34. Barrowcliffe TW. Lipolytic activities of low molecular weight heparins (letter). *Thromb Res* 1986; 42(4): 583-4.
35. Merton RE, Curtis AD, Thomas DP. A comparison of heparin potency estimates obtained by activated partial thromboplastin time and british pharmacopoeial assays. *Thromb Haemostas* 1985; 53(1): 116-7.
36. Swart de CAM, Nijmeyer B, Andersson L, Holmer E, Verschoor L, Bouma BN, Sixma JJ. Elimination of High Affinity Heparin Fractions and their Anticoagulant and Lipase Activity. *Blood* 1984; 6(4): 836-42.

37. Casu B, Johnson EA, Mantovani M, Mulloy B, Oreste P, Pescador R, Prino G, Torri G, Zopetti G. Correlation between structure, Fat-clearing and anticoagulant properties of heparins and heparan sulphates. *Arzneim forsch* 1983; 33: 135-42.
38. Markwardt F. Antipemic action of heparin. *Thromb Diath Haemorrh* 1974; 33: 73-6.
39. Putten van JJ, Ruit van de M, Beunis M, Hemker HC. Interindividual Variation in Relationships between Plasma Heparin Concentration and the Results of Five Heparin Assays. *Clin Chim Acta* 1982; 122: 261-70.
40. Penner JA. Experience with a Thrombin Clotting Time for Measuring Heparin Activity. *Am J Clin Pathol* 1974; 61: 645-53.
41. Lang M, Fey MF, Furlan M, Beck EA. Überwachung der Heparin-therapie mit der Thrombinzeit und der aktivierten partiellen Thromboplastinzeit - ein Vergleich. *Schweiz Med Wschr* 1986; 116: 1681-4.
42. Bounaemaux H, Marbet GA, Lammle B, Eichlisberger R, Duckert F. Monitoring of heparin treatment. Comparison of thrombin time, activated partial thromboplastin time and plasma heparin concentration and analysis of the behaviour of antithrombin III. *Am J Clin Path* 1980; 74: 68-73.
43. Brandt JT, Triplett DA. Laboratory monitoring of heparin. Effects of reagents and instruments on the activated partial thromboplastin time. *Am J Clin Path* 1981; 76: Suppl: 530-7.
44. Poller L, Thomson JM, Yee KF. Heparin and Partial Thromboplastin Time: an International Survey. *Brit J Haemat* 1980; 44: 161-165.
45. Suchman AL, Griner PF. Diagnostic Uses of the Activated Partial Thromboplastin Time and Prothrombin Time. *Ann Int Med* 1986; 104: 810-6.
46. Teien AN, Abildgaard U. On the Value of the Activated Partial Thromboplastin Time (APTT) in Monitoring Heparin Therapy. *Thromb Res* 1976; 35: 592-6.
47. Chen AL, Hershgold EJ, Wilson DE. One-stage assay of heparin. *J Lab Clin Med* 1975; 85: 843-54.
48. Griffiths BL. Monitoring the intrinsic pathway of coagulation during heparin therapy-reply (letter). *Thromb Haemostas* 1985; 53(2): 287.
49. Hoffmann JJ. Monitoring the intrinsic pathway of coagulation during heparin therapy (letter). *Thromb Haemostas* 1985; 53(1): 161.
50. Besselaar AMHP, van den Meeuwisse-Braun J, Jansen-Gruter R, Bertina RM. Monitoring Heparin Therapy by the Activated Partial Thromboplastin Time- The Effect of Pre-Analytical Conditions. *Thromb Haemostas* 1987; 57: 226-31.
51. Yin ET. Appraisal of Clot Based and Amidolytic Methods for the Monitoring of Heparin and its Derivatives. *Semin Thromb Hemost* 1985; 11: 243-4.
52. Yin ET. Procedure for the determination of heparin. *Thromb Diath Haemorrh* 1975; 33: 393-5.
53. Larsen ML, Abildgaard U, Teien AN, Gjersdal K. Assay of Plasma Heparin using Thrombin and Chromogenic Substrate H-D-Phe-Pip-Arg-pNA (S-2238). *Thromb Res* 1978; 13: 285-8.
54. Handeland G. Simplified chromogenic substrate assay for low molecular weight heparin. *Thromb Res* 1986; 42(1): 105-8.
55. Hoppensteadt DA, Walenga JM, Fareed J. Validity of serine protease inhibition tests in the evaluation and monitoring of the effect of heparin and its fractions. *Semin Thromb Hemost* 1985; 11(2): 112-20.

56. Putten van J, Ruit van de M, Beunis M, Hemker HC. Automated Determination of Heparin with Chromogenic Substrates. *Haemostasis* 1984; 14: 184-94.
57. Putten van J, Ruit van de M, Beunis M, Hemker HC. Automated Spectrophotometric Heparin Assays. *Haemostasis* 1984; 14: 195-204.
58. Teien AN, Lie M, Abildgaard U. Assay of Heparin in Plasma using a Chromogenic Substrate for Activated Factor X. *Thromb Res* 1976; 8: 413-6.
59. Putten van J, Ruit van de M, Beunis M, Hemker HC. Determination of Low Molecular Weight Heparin in Clinical Laboratory. *Haemostasis* 1984; 14: 205-10.
60. Walenga JM, Bara L, Samama MM, Fareed J. Amidolytic antifactor Xa assays in the laboratory evaluation of heparin and low molecular weight fractions. *Semin Thromb Hemost* 1985; 11(2): 100-7.
61. Aiach M, Roncato M. Practical considerations on the measurement of low molecular weight heparin anti-Xa activity. *Semin Thromb Hemost* 1985; 11(2): 108-11.
62. Delahousse B, Gruel Y, Moalic P, Toulemonde F, Leroy J. Comparative Efficacy of a New Clot-Based Anti-Xa/IIa Assay and an Amidolytic Anti-Xa Method during a very Low Molecular Weight Heparin Fragment (CY 222) Treatment. *Thromb Res* 1987; 46: 405-7.
63. Nenci GG, Berretti M, Agnelli G, Parise P. Heparin Assay: A Comparison between Titrimetric and Amidolytic Methods. *Boll Soc It Biol Sper* 1980; LVI:519-25.
64. Dawes J, Pepper DS. A Sensitive Competitive Binding Assay for Exogenous and Endogenous Heparins. *Thromb Res* 1982; 27: 387-96.
65. Hemker HC. The Mode of Action of Heparin in Plasma. In: Verstraete M, Vermeylen J, Lijnen HR, Arnout J, eds. *Thrombosis and Haemostasis*. Leuven: International Society on Thrombosis and Haemostasis and Leuven University Press, 1987.
66. Béguin S, Lindhout T, Hemker HC. The Mode of Action of Heparin in Plasma. *Thromb Haemostas* 1988; 60(3): 457-62.
67. Lane DA, MacGregor IR, Ross-Van M, Cella G, Kakkar VV. Molecular Weight Dependence of the Anticoagulant Properties of Heparin: Intravenous and Subcutaneous Administration of Fractionated Heparins to Man. *Thromb Res* 1979; 16: 651-62.
68. Thomas DP, Merton RE, Lewis ME, Barrowcliffe TW. Studies in Man and Experimental Animals of a Low Molecular Weight heparin Fraction. *Thromb Res* 1981; 45: 214-8.
69. Holmer E, Mattsson C, Nilsson S. Anticoagulant and Antithrombotic Effects of Heparin and Low Molecular Weight Heparin Fragments in Rabbits. *Thromb Res* 1982; 25: 475-85.
70. Buchanan MR, Boneu B, Ofosu FA, Hirsh J. The Relative Importance of Thrombin Inhibition to the Antithrombotic Activity of an Ultra-Low Molecular Weight Heparin Fraction. *Thromb Res* 1982; 28: 401-9.
71. Ockelford PA, Carter CJ, Mitchell L, Hirsh J. Discordance between the Anti-Xa Activity and the Antithrombotic Activity of an Ultra-Low Molecular Weight Heparin Fraction. *Thromb Res* 1982; 28: 401-9.
72. Ofosu FA, Blajchman MA, Modi GJ, Smith LM, Buchanan MR, Hirsh J. The Importance of Thrombin Inhibition for the Expression of the Anticoagulant Activities of Heparin, Dermatan Sulphate, Low Molecular Weight Heparin and Pentosan Polysulphate. *Brit J Haematol* 1985; 60: 695-704.
73. Ofosu FA, Smith LM, Anvari N, Blajchman MA. An Approach to Assigning in Vitro Potency to Low Molecular Weight Heparins based on the Catalysis of Thrombin Inhibition and the Inhibition of Prothrombin Activation. *Thromb Haemostas* 1988; 60: 193-8.

74. Ofosu FA, Modi G, Blajchman MA, Hirsh J. The Activation of Factor X and Prothrombin in Antithrombin III-dependent plasma: The Effects of Heparin and Antithrombin III. *Thromb Res* 1981; 23: 331-345.
75. Dunn FW, Soria C, Thomaidis A, Soria J. Interactions of Platelets with Standard Heparin and Low Molecular Weight Fractions. *Nouv Rev Fr Hematol* 1984; 26: 249-53.
76. Heiden D, Mielke CH, Rodvien R. Impairment by Heparin of Primary Haemostasis and Platelet (14C) 5-Hydroxytryptamine Release. *Br J Haematol* 1977; 36: 427-36.
77. Ljungberg B, Johnsson H. In Vivo Effects of a Low Molecular Weight Heparin Fragment on Platelet Aggregation and Platelet Dependent Hemostasis in Dogs. *Thromb Haemostas* 1988; 60(2): 232-5.
78. Saba SR, Saba HI, Morelli GA. Effect of Collagen and Citrate on Heparin-related Antiaggregatory Activity. *Thromb Haemostas* 1987; 58: 18 (Abstract).
79. Westwick J, Scully MF, Poll C, Kakkar VV. Comparison of the effects of low molecular weight heparin and unfractionated heparin on activation of human platelets in vitro. *Thromb Res* 1986; 42: 435-47.
80. Brace LD, Fareed J. An objective assessment of the interaction of heparin and its fractions with human platelets. *Semin Thromb Hemost* 1985; 11(2): 190-8.
81. Brace LD, Fareed J. Heparin-induced platelet aggregation: dose/response relationships for a low molecular weight heparin derivative (PK10169) and its subfractions. *Thromb Res* 1986; 42(6): 769-82.
82. Sie P, Fernandez F, Caranobe C, Gabaig AM, Boneu B. Inhibition of thrombin-induced platelet aggregation and serotonin release by antithrombin III and heparin cofactor II in the presence of standard heparin, dermatan sulfate and pentosan polysulfate. *Thromb Res* 1984; 35: 231-6.
83. O'Brien JR, Path FRC. Heparin and Platelets. *Current therapeutic Research* 1975; 18(1): 79-90.
84. Fernandez F, N'guyen P, Van Ryn J, Ofosu FA, Hirsh J, Buchanan MR. Hemorrhagic doses of heparin and other glycosaminoglycans induce a platelet defect. *Thromb Res* 1986; 43(4): 491-5.
85. Béguin S, Lindhout T, Hemker HC. The Effect of Trace Amounts of Tissue Factor on Thrombin Generation in Platelet Rich Plasma, its Inhibition by Heparin. *Thromb Haemost* 1989; 61: 25-9.
86. Walz DA, Hung GL. In vivo studies on the binding of heparin and its fractions with platelet factor 4. *Semin Thromb Hemost* 1985; 11(1): 40-7.
87. Lane DA. Platelet-derived heparin neutralizing proteins. *Adv Exp Med Biol* 1985; 192: 427-38.
88. Ofosu FA, Cerskus AL, Hirsh J, Smith LM, Modi GJ, Blajchman MA. The Inhibition of the Anticoagulant Activity of Heparin by Platelets, Brain Phospholipids, and Tissue Factor. *Brit J Haematol* 1984; 57: 229-38.
89. Conley CL, Hartmann RC, Lalley JS. The Relationship of Heparin Activity to Platelet Concentration. *Proc Soc Exp Biol Med* 1948; 69: 284-7.
90. Lane DA, Denton J, Flynn AM, Thunberg L, Lindahl U. Anticoagulant activities of heparin oligosaccharides and their neutralization by platelet factor 4. *Biochem J* 1984; 218: 725-32.
91. Lane DA, Pejler G, Flynn AM, Thompson EA, Lindahl U. Neutralization of heparin-related saccharides by histidine-rich glycoprotein and platelet factor 4. *J Biol Chem.* 1986; 261(9): 3980-6.

92. Levine SP, Sorenson RR, Harris MA, Knieriem LK. The effect of platelet factor 4 (PF4) on assays of plasma heparin. *Brit J Haematol* 1984; 57: 585-96.
93. Putten van J, Ruit van de M, Beunis M, Hemker HC. Heparin Neutralization during Collection and Processing of Blood Inhibited by Pyridoxal 5'-Phosphate. *Haemostasis* 1984; 14: 253-61.
94. van de Besselaar AMHP, Meeuwisse-Braun J, Bertina RM. Monitoring heparin therapy: relationships between the activated partial thromboplastin time and heparin assays based on ex-vivo heparin samples. *Thromb Haemostas* 1990; 63: 16.
95. Bain B, Forster T, Sleight B. Heparin and the activated partial thromboplastin time - a difference between in vivo and in vitro effects and implications for the therapeutic range, 1980.
96. Béguin S, Mardiguian J, Lindhout T, Hemker HC. The Mode of Action of Low Molecular Weight Heparin Preparation (PK10169) and Two of its Major Components on Thrombin Generation in Plasma. *Thromb Haemostas* 1989; 61(1): 30-4.
97. Béguin S, Choay J, Hemker HC. The Action of a Synthetic Pentasaccharide on Thrombin Generation in Whole Plasma. *Thromb Haemostas* 1989; 61(3): 397-401.
98. Tew CJ, Lane DA, Thompson E, Ireland H, Curtis JR. Relationship between ex vivo anti-proteinase (factor Xa and thrombin) assays and in vivo anticoagulant effect of very low molecular weight heparin, CY 222. *Br J Haematol* 1988; 70: 335-40.
99. Albada J, Nieuwenhuis HK, Sixma JJ. Treatment of acute venous thromboembolism with low molecular weight heparin (Fragmin): Results of a double blind study. *Thromb Haemostas* 1989. In press.
100. Levine MN, Planes A, Hirsh J, Goodyear M, Vochelle N, Gent M. The relationship between anti-factor Xa level and clinical outcome in patients receiving enoxaparin low molecular weight heparin to prevent deep vein thrombosis after hip replacement. *Thromb Haemostas* 1989. In press.
101. Dechavanne M, Ville D, Berruyer M, Trepo F, Dalery F, Clermont N, Lerat JL, Moyen B, Fischer LP, Kher A, Barbier P. Randomized trial of a low-molecular weight heparin (Kabi 2165) versus adjusted-dose subcutaneous standard heparin in the prophylaxis of deep-vein thrombosis after elective hip surgery. *Haemostasis* 1989; : 5-12.
102. Kakkar VV, Djazaeri B, Fok J, Fletcher M, Scully MF, Westwick J. Low molecular weight heparin and prevention of postoperative deep vein thrombosis. *Br Med J* 1982; 284: 375-9.
103. Harenberg C, Giese C, Knodler A, Zimmerman R. Comparative study on a new one-stage clotting assay for heparin and its low molecular weight derivatives. *Haemostasis* 1989; 1: 13-20.
104. Ireland H, Lane DA, Flynn A, Pegrum AC, Curtis JR. Low molecular weight heparin in haemodialysis for chronic renal failure. *Thromb Haemostas* 1988; 59: 240-7.
105. Walenga JM, Petitou M, Lormeau JC, Samama MM, Fareed J, Choay J. Antithrombotic activity of a synthetic heparin pentasaccharide in a rabbit stasis thrombosis model using different thrombogenic challenges. *Thromb Res* 1987; 46: 187-98.
106. Cadroy Y, Harker LA, Hanson SR. Inhibition of platelet dependent thrombosis by low molecular weight heparin (CY222); comparison with standard heparin. *J Lab Clin Med* 1989; 114: 349-57.
107. Lane DA, Ryan K. Heparin and low molecular weight heparin: Is anti-factor Xa activity important? *J Lab Clin Med* 1989; 114: 331-3.

108. Hemker HC. Is anti-factor Xa important? *J Lab Clin Med* 1990. In press.
109. Barrowcliffe TW, Merton RE, Havercroft SJ, Thunberg L, Ludahl U, Thomas DP. Low-affinity heparin potentiates the action of high-affinity heparin oligosaccharides. *Thromb Res* 1984; 34: 125-34.
110. Béguin S, Dol F, Hemker HC. Factor IXa inhibition contributes to the heparin effect. Submitted, 1990.
111. Colman RW, Wong PY. Participation of Hageman Factor Dependent Pathways in Human Disease States. *Thromb Haemos* 1977; 38(4): 751-75.
112. Green D, Hougie C, Kazmier FJ, Lechner K, Mannucci PM, Rizza CR, Sultan Y. Report of the Working Party on Acquired Inhibitors of Coagulation: Studies of the "Lupus" Anticoagulant. *Thromb Haemostas* 1983; 49: 144-6.
113. Lämmle B, Tran TH, Ritz R, Duckert F. Plasma Prekallikrein, Factor XII, Antithrombin III, C1-Inhibitor and Alpha-2-macroglobulin in Critically Ill Patients with suspected Disseminated Intravascular Coagulation (DIC). *Am J Clin Pathol* 1984; 82: 396-404.
114. Mason JW, Kleeberg U, Dolan P, Colman RW. Plasma Kallikrein and Hageman Factor in Gram-negative Bacteremia. *Ann Int Med* 1983; 73: 545-51.
115. Ragni MV, Lewis JH, Hasiba U, Spero JA. Prekallikrein (Fletcher factor) Deficiency in Clinical Disease States. *Thromb Res* 1980; 18: 45-54.
116. Saito H, Poon MC, Vicic W, Glodsmith GH, jr, Menitove JE. Human Plasma Prekallikrein (Fletcher Factor) Clotting Activity and Antigen in Health and Disease. *J Lab Clin Med* 1978; 92: 84-95.
117. Hemker HC. A standard for low molecular weight heparin? *Haemostasis* 1989; 19: 1-4.
118. Ma Xi, Béguin S, Hemker HC. Importance of factor IX dependent protrombinase formation - The Jasso Pathway - in clotting plasma. *Haemostasis* 1989; 19: 301-8.
119. Hemker HC, Béguin S. The mode of action of heparin and related drugs. *Semin Thromb Haemostas* 1990. In press.