Inhibition of thrombin-catalyzed reactions in blood coagulation and platelet activation by heparin fractions in the absence of antithrombin III

Citation for published version (APA):

Baruch, D., Lindhout, T., Wagenvoort, R., & Hemker, H. C. (1986). Inhibition of thrombin-catalyzed reactions in blood coagulation and platelet activation by heparin fractions in the absence of antithrombin III. *Haemostasis*, 16(2), 71-81. <https://doi.org/10.1159/000215276>

Document status and date:

Published: 01/01/1986

DOI:

[10.1159/000215276](https://doi.org/10.1159/000215276)

Document Version:

Other version

Please check the document version of this publication:

- A submitted manuscript is the version of the article upon submission and before peer-review. There can be important differences between the submitted version and the official published version of record. People interested in the research are advised to contact the author for the final version of the publication, or visit the DOI to the publisher's website.
- The final author version and the galley proof are versions of the publication after peer review.
- The final published version features the final layout of the paper including the volume, issue and page numbers.

[Link to publication](#)

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal.

If the publication is distributed under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license above, please follow below link for the End User Agreement:

www.umlib.nl/taverne-license

Take down policy

If you believe that this document breaches copyright please contact us at:

repository@maastrichtuniversity.nl

providing details and we will investigate your claim.

Download date: 26 Apr. 2023

Inhibition of Thrombin-Catalyzed Reactions in Blood Coagulation and Platelet Activation by Heparin Fractions in the Absence of Antithrombin III

Dominique Baruch, Theo Lindhout, Rob Wagenvoord, H. Coenraad Hemker

Department of Biochemistry, University of Limburg, Maastricht, The Netherlands

Key Words. Thrombin · Heparin · Factor V · Factor VIII · Platelets

Abstract. The antithrombin-III-independent effect of heparin was studied in the following thrombin-catalyzed reactions: activation of purified plasma factor V and partially purified plasma factor VIII:C, generation of factor V_a from the platelets and, in the presence of collagen, of the platelet procoagulant activity. Five heparin fractions and a heparinoid were compared to crude heparin. Crude heparin was a more potent inhibitor of these reactions than the fractions or the heparinoid. The inhibitory action of heparin (fractions) appeared to be the result of the formation of a complex between heparin and thrombin that alters the specificity of thrombin towards high molecular weight substrates. The inhibition of these thrombin-dependent feedback reactions in blood coagulation might be of importance in the mechanisms for the dissociation between the antithrombotic and hemorrhagic properties of low molecular weight heparins.

Introduction

Heparin is one of the most important drugs for the treatment and prevention of venous thrombosis and thromboembolism. However, it still presents many puzzling questions on different levels to clinicians and clinical chemists, biologists, biochemists and certainly not least for the pharmaceutical industry. Clinicians and clinical chemists know that the major drawback to the therapeutical use of heparin is the absence of any reliable laboratory test to keep the patient in the narrow range that ensures an efficient anti-

thrombotic action but will not cause a hemorrhagic tendency. From the literature it is evident that the properties of heparin to catalyze factor X_a and/or thrombin inactivation by antithrombin III do not sufficiently reflect the complexity of in vivo heparin actions. Research over the last few years has led to the development of heparin fractions of known compositions which allowed considerable progress in understanding the structure-function relationships. Our contribution, as biochemists of blood coagulation, is to identify the complete spectrum of heparin-sensitive reactions, to develop a relevant in vitro de-

termination of the action of heparin and to establish their correlation with *in vivo* effects.

As thrombin formation is the central process in hemostasis and thrombosis, the inhibition by heparin of thrombin formation has been our main focus of interest. It should be appreciated that heparin can inhibit thrombin by 2 separate mechanisms, i.e. the antithrombin activity can be either antithrombin-III-dependent or -independent.

Thrombin Generation and Feedback Mechanisms

The mechanisms leading to thrombin formation are, on the one hand, a series of proenzyme activation reactions (factor VII $\rightarrow$ factor X $\rightarrow$ factor II) and on the other hand interactions at a phospholipid-solute interface [1]. Procoagulant, negatively charged phospholipids are provided by the membrane of activated platelets and wounded cells [2]. The assembly of an enzymatic complex at the phospholipid surface is completed by the presence of non-enzymatic protein co-factors (respectively tissue factor, factor VIII_a and factor V_a) [3–5]. The interaction of thrombin and platelets is a complicated process. Thrombin once formed in trace amounts governs a set of positive feedback reactions. Thrombin is a potent platelet activator and induces the release reactions by which factor V becomes available. Platelet factor V seems to contribute more to thrombin generation and hence to hemostasis and coagulation than plasma factor V does. This was demonstrated within a group of factor-V-deficient patients, where the severity of bleeding was shown to depend on the platelet rather than on the plasma factor V deficiency [6]. Thrombin also converts factor V and factor VIII into their activated forms. Together

with collagen, thrombin generates the appearance of a procoagulant surface at the outer membrane of the platelets [2]. If the thrombin formation were not controlled by natural inhibitors and regulatory processes (such as inactivation of factors V_a and VIII_a by activated protein C), these coagulation factors and platelet interactions would result in an explosive thrombin formation.

The Antithrombin-III-Dependent Action of Heparin

The main anticoagulant function of heparin is to enhance the inactivation of the serine proteases of blood coagulation (thrombin, factor X_a, IX_a, XI_a and XII_a) by antithrombin III (AT III). Three models are currently described. (a) Heparin binds to AT III, inducing a conformational change [7, 8]; (b) the reactivity of thrombin with AT III is enhanced by the binding of heparin to thrombin [9, 10]; and (c) a ternary complex is formed and stabilized by the interaction of AT III and thrombin with heparin, the so-called template model [11–13]. These models are not mutually exclusive.

Factor X_a inactivation by the heparin-AT III complex is accounted for by the interaction of factor X_a with AT III and not by the interaction of factor X_a with heparin bound to AT III. Factor XII_a is inhibited in the same way. The saccharide structure required to potentiate the inactivation of factor X_a and factor XII_a is contained within the AT-III-binding region [14]. Heparin fractions with a molecular weight higher than 3,400 are equally active [15]. On the contrary, thrombin and factors IX_a and XI_a can bind to the heparin molecule as well as to the AT III molecule. The potency of heparin in catalyzing the AT III thrombin reaction increases with the molecular weight of heparin [16]. At

high concentrations, heparin has an additional effect by catalyzing the inhibition of thrombin by heparin cofactor II [17].

However, because thrombin formation is a phospholipid-bound process, the inactivation of the serine proteases of blood coagulation has to be considered as it occurs at the phospholipid-solute interphase, where the enzyme is located when it is physiologically active, rather than on the free enzyme. It has already been known for several years [18, 19] that factor X_a bound to phospholipids is much less accessible to the inactivation by AT-III-heparin than free factor X_a is. This was recently demonstrated to be also the case when factor X_a and factor V_a bind to activated platelets, therefore requiring that these complexes are cleared from blood before the heparin therapy can be effective [20, 21].

The Antithrombin-III-Independent Effect of Heparin

Evidence for an AT-III-independent effect of heparin in reactions of blood coagulation and hemostasis is three-fold.

(1) Some heparin fractions with a low affinity for AT-III, which are virtually devoid of any antithrombin or antifactor X_a activity in vitro significantly increased hemorrhage without influencing the in vivo antithrombotic effect in animal experimental models [22]. Furthermore this low affinity heparin fraction potentiated the antithrombotic action of a high affinity heparin oligosaccharide of a molecular weight lower than 3,500 which only had an antifactor X_a activity in vitro [23].

(2) In a system consisting of purified coagulation factors and synthetic phospholipids, heparin can inhibit prothrombin and factor X activation. This effect was attributed to the displacement of the bound proteins from the phospholipid surface [24-26]. However,

these experimental results can alternatively be explained by an effect of heparin on the thrombin-catalyzed factor VIII activation. As the effect of heparin in these experiments was practically abolished if the activated factor VIII was present, the latter explanation seems more probable. Also the fact that in an AT-III-depleted plasma the effect of heparin on the inhibition of factor X_a or thrombin generation is almost negligible, supports this notion [27]. Comparison of crude heparin and low affinity heparin in prothrombin activation in AT-III-depleted plasma showed that the inhibitory effect of both heparins was similar. These experiments do not exclude an effect due to other serine protease inhibitors. No further insight was brought in the mechanism underlying the displacing effect by heparin of protein-lipid interactions [28].

(3) Conflicting data have been reported showing that heparin either induced [29] or inhibited platelet aggregation [30]. It was shown that some of the heparin effects were related to an interaction with platelet functions [31]. Bleeding caused by heparin has been attributed to a direct effect on platelets. The original observation of *Salzman et al.* [32] that high molecular weight, low AT-III affinity heparin fractions, were more reactive on platelets than low molecular weight, high AT-III affinity fractions, suggested a correlation between the hemorrhagic effect of heparin and the anticoagulant properties in vitro. However, this correlation is far from being a strong one [33].

In this study, we show that an AT-III-independent effect of heparin in blood coagulation and platelet reactions might be the result of an interaction between thrombin and heparin. To this end, we used purified proteins, synthetic phospholipids or washed platelets under well defined kinetic condi-

tions, together with different heparin fractions of known antithrombotic/hemorrhagic properties [34–37].

Materials and Methods

Experimental procedures for protein and phospholipid preparation as well as platelet isolation are as described earlier [38]. Purification of bovine factor VIII:C was performed by differential precipitation by polyethylene glycol. The determination of the platelet factor V_a generation upon the addition of thrombin, and of the platelet procoagulant activity in the presence of thrombin and collagen, has been reported previously [38].

Heparin Fractions

We were interested in measuring the effect of heparin fractions that had already been used in clinical trials and in vivo animal experimental models [34–37]. They were supplied by the following laboratories: Choay Institute (CY 216, an ethanol precipitation fragment; CY 222, a fragment obtained by nitrous acid depolymerization), Pharmuka Laboratories (FF 32038, a crude heparin; PK 10169, obtained by controlled depolymerization; EMT 966, EMT 967, obtained by gel filtration of PK 10169) and Organon Laboratories (crude heparin and Org 10172, a sulphated mucopolysaccharide). The average molecular weights, antifactor X_a and antithrombin activities are listed in table I.

Purified Plasma Factor V Activation

Bovine factor V (100 nM) was incubated for 5 min at 37 °C in the presence of 3 mM CaCl₂. The reaction was started by the addition of thrombin (usually 0.3 nM). Aliquots were taken in time, diluted 1:1,000 in buffer containing 50 mM Tris, 175 mM NaCl, 5 mM CaCl₂ and 0.5 mg/ml HSA, pH 7.9. Factor V_a activity was measured as previously described [38].

Partially Purified Factor VIII:C Activation

Partially purified factor VIII:C was activated by bovine thrombin in the following way: factor VIII:C (20 µl, 27 U/ml) was incubated in 1 ml Michaelis buffer containing 29 mM sodium barbiturate, 29 mM sodium acetate and 116 mM NaCl, pH 7.3. The reaction was started with the addition of thrombin (3 nM).

Table I. Characteristics of heparin fractions

Compound	Mean molecular weight	Anti X _a activity U/mg	Anti II activity U/mg
Crude heparin	15,000	175	175
FF 32038	15,000	165	165
Org 10172	6,400	8	0.32
CY 216	4,500	200	10–20
CY 222	2,500	250	5–15
PK 10169	4,500	120	30
EMT 966	6,500	160	70
EMT 967	3,300	100	0

Aliquots (10 µl) of a factor-VIII_a-containing sample were added to the clotting assay containing 60 µl factor-VIII-deficient plasma, 4 nM factor IX_a, 50 µM phospholipid [(25% phosphatidylserine (PS), 75% phosphatidylcholine (PC)] and 10 mM CaCl₂. The clotting time was recorded on an automatic coagulometer. A calibration curve was established with normal pool plasma activated with 9 nM thrombin in order to determine the amount of factor VIII_a present in the sample.

Inhibition Studies

The assessment of a heparin effect on thrombin-catalyzed reactions can be made provided that the following conditions are met.

(1) Thrombin formation by the prothrombinase complex can be measured as a function of varying concentrations of an accessory component (factor V_a or phospholipids) under the conditions where the accessory component under investigation is the only rate-limiting factor of the reaction. Factor VIII_a can be determined in a clotting assay in a comparable manner.

(2) An effect of heparin on a thrombin-catalyzed reaction can be assessed if the manner in which this reaction depends on the thrombin concentration is known. This is shown, in figure 1, for the thrombin-catalyzed factor VIII activation and in figure 2 for the thrombin-catalyzed factor V activation. Throughout this study, excess thrombin concentrations were avoided in order to be able to quantitate the heparin

Fig. 1. Dependency of the factor VIII activation on thrombin concentration. Normal plasma is incubated at 37 °C with varying bovine thrombin concentrations for 15 s. A sample is brought to the factor X activation mixture containing phospholipids vesicles (25% PS/75% PC, 5×10^{-5} M), CaCl_2 (10 mM) and factor IX_a (4.0×10^{-9} M). The rate of factor X activation is measured after 2 and 4 min. The results are expressed as rate of the factor X_a formation per minute per unit factor VIII_a present in 1 ml plasma.

Fig. 2. Dependency of factor V activation on thrombin concentration. Purified bovine factor V (100 nM) was incubated for 5 min at 37 °C in a buffer containing 50 mM Tris, 175 mM NaCl, 3 mM CaCl_2 , 0.5 mg human serum albumin/ml in the presence of varying thrombin concentrations. Factor V_a activity was determined as described in 'Materials and Methods'.

effects. We previously showed that a similar relationship exists in the thrombin-catalyzed platelet V_a generation and thrombin-plus collagen-induced platelet procoagulant activity [38].

(3) The interaction between heparin and thrombin results in the formation of a 1:1 complex. This complex formation leaves unaffected the catalytic site of thrombin towards the synthetic chromogenic substrate, but has an altered enzymatic activity towards macromolecular substrates. In contrast with others [39], we found that heparin and heparin fractions up to 200 µg/ml had no effect on the catalytic efficiency of thrombin on the chromogenic substrate S2338 when the substrate concentration is above or near the K_m (data not shown).

Upon the formation of a heparin-thrombin complex, the inhibition of the thrombin activity on its macromolecular substrate results in a decrease in the rate of the thrombin-catalyzed reaction. The IC_{50} is the heparin concentration required to obtain half-maximal reaction velocity. The IC_{50} values can be

accurately determined by a plot of the reciprocal total heparin concentration versus the ratio $v_0/(v_0-v)$, where v_0 and v are the velocities of the reaction in the absence and the presence of heparin, respectively.

Results and Discussion

Demonstration of the Absence of Interaction of Heparin with Other Components than Thrombin

Before we could attribute to heparin the observed effects on the thrombin-catalyzed reactions, we had to exclude any effect of heparin carried over into the assay mixtures. Therefore, as an effect of heparin on the protein-phospholipid interaction has been reported [22], we had to estimate the effect of

heparin on separate protein-lipid interactions. This was done by use of a protein-lipid equilibrium technique according to *van de Waart et al.* [40]. Factor X_a ($0.1 \mu M$), factor V_a ($0.1 \mu M$) or prothrombin ($0.05 \mu M$) were incubated with large volume phospholipid vesicles, $25 \mu M$ (20% PS – 80% PC), in 50 mM Tris, 100 mM NaCl, pH 7.5, at $20^\circ C$. After centrifugation for 30 min at 30,000 g at $20^\circ C$, the concentrations of the bound proteins were estimated by determination of the remaining activity in the supernatant (factor X_a was determined on S2337, factor V_a by the factor V_a assay and prothrombin after activation by *Echis carinatus* venom coagulant). Both heparin FF 32038 and PK 10169 at a final concentration of 0.5 mg/ml could not displace factor V_a , factor X_a or prothrombin from the phospholipid surface. The concentrations of bound protein in the presence and absence of heparin (fraction) were: $0.02 \mu M$, $0.10 \mu M$, $0.03 \mu M$ for factor X_a , factor V_a and prothrombin, respectively.

The effect of heparin on the thrombin-catalyzed factor VIII activation is measured in a clotting assay using a factor-VIII-deficient plasma; therefore heparin is carried over and indeed influences the clotting times. This problem is circumvented by making a calibration curve for the factor VIII_a formation in the presence of the same amounts of heparin as those that are carried over. In that way, heparin does not affect the relative changes occurring during the thrombin-catalyzed factor VIII activation. In all other studies, we used a completely purified system, in which no effect of heparin carried over into the assay was observed. This was demonstrated by letting the substrate of thrombin be fully activated in the absence of heparin and adding a high heparin concentration immediately before transferring a sample to the

measuring system. No effect of heparin or heparin fractions at final concentrations of up to 0.5 mg/ml was observed after a 10-min incubation, showing that the amount of thrombin-activated substrate was independent of the presence of heparin in the activation mixture. This demonstrates that heparin did not interfere with the assembly or the functioning of the prothrombinase complex.

In the reactions where platelets are involved, we had to rule out the presence of AT III in the washed platelet preparations [41]. This was done by incubation of washed platelet suspension (10^7 /ml) with 5 nM thrombin in the presence of heparin (fractions) (10 μg /ml). After 20 min, the amidolytic activity of thrombin appeared to be unaffected. In order to discriminate between an effect of heparin on the thrombin-induced factor V_a generation and an effect related to the binding of heparin to platelets, we measured the effect of heparin on the beta-thromboglobulin release, induced by either thrombin or collagen. Only the thrombin-induced release of beta-thromboglobulin was affected by heparin (90% inhibition of the release reaction at 50 μg heparin/ml), showing that the reaction was occurring between thrombin and heparin and was not due to an effect of heparin on platelets.

Effect of Heparin (Fractions) on the Thrombin-Catalyzed Plasma Factor VIII Activation

We used a semi-quantitative estimation of the effect of heparin on the thrombin-catalyzed factor VIII activation. Figure 3 shows the time course of the factor VIII activation in the absence and presence of varying amounts of unfractionated heparin (FF 32038). Heparin has an effect both on the initial rate of the factor VIII activation and on

Fig. 3. Effect of crude heparin FF 32038 on the thrombin-catalyzed factor VIII activation. Isolated factor VIII:C was incubated with thrombin (3 nM) and varying concentrations of heparin FF 32038. ● = No heparin; ○ = 0.14 $\mu\text{g/ml}$; ▲ = 0.70 $\mu\text{g/ml}$; △ = 1.43 $\mu\text{g/ml}$; ■ = 2.9 $\mu\text{g/ml}$; □ = 5.7 $\mu\text{g/ml}$. At timed intervals, factor VIII_a-containing samples were transferred to the assay. Factor VIII_a was determined by using a calibration curve established with normal plasma as described in 'Materials and Methods'.

the rate of inactivation of factor VIII_a. It was observed that the level of factor VIII_a after 4 min was in all cases lower in the absence than in the presence of heparin. This stabilizing effect of heparin could be due to an effect either on thrombin, and/or on the factor VIII_a. Further experiments are required for a better understanding. We estimated the effect on the initial rate of activation, of heparin FF 32038, PK 10169 and Org 10172 (fig. 4). The IC₅₀ values were found to be 0.5 $\mu\text{g/ml}$, 1.75 $\mu\text{g/ml}$ and 25 $\mu\text{g/ml}$, respectively.

Effect of Heparin (Fractions) on the Thrombin-Catalyzed Plasma Factor V Activation

Bovine factor V (100 nM) was incubated at 37 °C with calcium (3 mM) and varying concentrations of heparin (fractions). The reaction was initiated by the addition of thrombin (0.3 nM). After 6 min, 1:1,000 diluted samples were assayed for the factor V_a activity. The percentage of the residual factor V_a concentration was plotted as a function of the heparin concentration (fig. 5). Determination of the IC₅₀ values was performed

by use of the double reciprocal plot as described above (fig 5, inset). It is clear that unfractionated heparin is a more efficient inhibitor (IC₅₀ = 10 $\mu\text{g/ml}$) than the fractions CY 216 (IC₅₀ = 25 $\mu\text{g/ml}$), CY 222 (IC₅₀ = 25 $\mu\text{g/ml}$) or the heparinoid Org 10172 (IC₅₀ = 125 $\mu\text{g/ml}$). The IC₅₀ values of EMT 966, EMT 967 and PK 10169 were 28 $\mu\text{g/ml}$, 12.5 $\mu\text{g/ml}$ and 62.5 $\mu\text{g/ml}$, respectively.

Effect of Heparin Fractions on the Thrombin-Induced Platelet Factor V_a Generation and on the Thrombin-plus Collagen-Induced Platelet Procoagulant Activity

We have previously described a direct inhibitory effect of heparin on thrombin in platelet activation [38]. We tested the effect of heparin on the thrombin-induced platelet factor V_a generation and on the thrombin-plus collagen-induced platelet procoagulant activity. We have extended the previous study to the effect of other heparin fractions, PK 10169 and 2 fractions obtained by gel filtration of PK 10169: EMT 966 and EMT 967. The results of these studies are depicted

Fig. 4. Effect of heparin (fractions) on the initial rate of the thrombin-catalyzed factor VIII activation. Factor VIII:C was incubated with thrombin in the presence of varying amounts of heparin. The initial rate of the factor VIII activation was measured at 30 s as described in the legend to figure 3. a ● = FF 32038; ○ = PK 10169. b ● = Org 10172.

Fig. 5. Effect of heparin on the initial rate of the thrombin-catalyzed factor V activation: factor V (100 nM) was activated for 5 min by thrombin (0.3 nM) in the presence of $CaCl_2$ (3 mM) and varying concentrations of heparin. ○ = Crude heparin; ● = CY 216; ▲ = CY 222; ■ = Org 10172. Factor V_a activity was determined as explained in 'Materials and Methods'.

in table II. Whereas in the platelet factor V_a generation the IC_{50} values are very similar (± 20 µg/ml), there are striking differences for the procoagulant activity of thrombin-plus collagen-activated platelets. Heparin FF 32038 inhibited the thrombin-plus collagen-induced procoagulant activity with an IC_{50} of 2 µg/ml, whereas much higher concentrations of PK 10169 and EMT 967 would be required to obtain the same extent of inhibition.

Conclusions

From in vitro studies we have been able to demonstrate a direct inhibitory effect of heparin on several thrombin-dependent reactions in blood coagulation. An AT-III-independent effect of heparin is shown on the thrombin-catalyzed factor VIII activation.

Table II. Effect of heparin on the thrombin-induced platelet factor V_a generation and on the thrombin-plus collagen-induced platelet procoagulant activity

Compound	IC ₅₀ (μg/ml)	
	factor V_a generation	procoagulant activity
FF 32038	18	2
PK 10169	22	> 150
EMT 966	25	17
EMT 967	16	> 150

Human washed platelets (4.6×10^6 /ml) were incubated for 3 min at 37 °C with 3 mM calcium and varying concentrations of heparin (fractions). The reaction was started with the addition of the activator (thrombin 0.5 nM or thrombin 0.3 nM plus collagen 5 μg/ml) and after 5 min the respective activities were determined.

This is in agreement with the results of *Oforu et al.* [26] obtained by a completely different method.

In purified systems, the same effect was shown on the plasma factor V activation and, as demonstrated earlier [38], on the platelet factor V_a generation and platelet procoagulant activity. We have demonstrated that this effect was due to a nonspecific heparin-thrombin interaction. We have not been able to confirm the observation of *Walker and Esmon* [24] that the AT-III-independent effect of heparin is due to interference with the protein-lipid interaction, as we performed binding experiments that allow direct determination of the protein bound to phospholipids and were unable to find any effect of heparin on this binding.

Structural characteristics of the heparin molecule have been described that are required for protease-binding: the charge den-

sity of heparin is independent of the affinity for AT III, but correlates with the thrombin-binding [42], and a recent report describes that the surface charge as well as the surface charge density of heparin [43] is important in thrombin-binding.

Sulphatation seems to be an important parameter for the nonspecific binding of polysaccharides to various proteins. This not only holds for the heparin-protease interaction, but also for the inhibiting action of pentosane polysulphate [44], recently shown to be caused by inhibition of the thrombin-induced factor VIII:C activation [*Wagenvoord*, unpublished results]. Also the interaction of heparin fractions with endothelium is seen to be strongly dependent upon the degree of sulphatation [*Barzu*, personal communication]. Further studies are required to better characterize the in vivo significance of this effect, as to the dissociation between antithrombotic and hemorrhagic effect of low molecular weight heparin fractions. Inhibition by heparins of the feedback activation reactions of thrombin by a process independent of AT III anyhow seems to be one of the mechanisms by which heparins modulate the thrombin formation and hence hemostasis and thrombosis.

Acknowledgements

We are grateful to *Jo Franssen* and *Hans Hendrix* for excellent technical assistance and to Mrs. *T. Camp-huisen* for preparing the manuscript. *D. Baruch* was supported by a fellowship from the Institut National de la Santé et de la Recherche Médicale, Paris.

References

- 1 Hemker, H.C.; Lindhout, Th.: A clotting scheme for 1984. *Nouv. Revue fr. Hémat.* 26: 222-231 (1984).

- 2 Bevers, E.M.; Comfurius, P.; van Rijn, J.L.M.L.; Hemker, H.C.; Zwaal, R.F.A.: Generation of prothrombin-converting activity and the exposure of phosphatidylserine at the outer surface of platelets. *Eur. J. Biochem.* 122: 429-436 (1982).
- 3 Rosing, J.; Tans, G.; Govers-Riemslog, J.W.P.; Zwaal, R.F.A.; Hemker, H.C.: The role of phospholipids and factor V_a in the prothrombinase complex. *J. biol. Chem.* 255: 274-283 (1980).
- 4 Van Dieijen, G.; Tans, G.; Rosing, J.; Hemker, H.C.: The role of phospholipid and factor VIII_a in the activation of bovine factor X. *J. biol. Chem.* 256: 3433-3442 (1981).
- 5 Nemerson, Y.: Regulation of the initiation of coagulation by factor VII. *Hemostasis* 13: 150-155 (1983).
- 6 Miletich, J.P.; Kane, W.H.; Hofman, S.L.; Stanford, N.; Majerus, P.W.: Deficiency of factor X_a-factor V_a binding site on the platelets of a patient with a bleeding disorder. *Blood* 54: 1015-1022 (1979).
- 7 Jordan, R.; Beeler, D.; Rosenberg, R.: Fractionation of low molecular weight heparin species and their interaction with antithrombin. *J. biol. Chem.* 254: 2902-2913 (1979).
- 8 Olson, S.T.; Shore, J.D.: Binding of high affinity heparin to antithrombin III. Characterization of the protein fluorescence enhancement. *J. biol. Chem.* 256: 11065-11072 (1981).
- 9 Machovich, R.; Blasko, G.; Palos, L.A.: Action of heparin on thrombin-antithrombin reaction. *Biochim. biophys. Acta* 379: 193-200 (1975).
- 10 Griffith, M.J.: Kinetic analysis of the heparin-enhanced antithrombin III-thrombin reaction. *J. biol. Chem.* 254: 12044-12049 (1979).
- 11 Pomerantz, M.W.; Owen, W.G.: A catalytic role for heparin. Evidence for a ternary complex of heparin cofactor thrombin and heparin. *Biochim. biophys. Acta* 535: 66-77 (1978).
- 12 Nesheim, M.E.: A simple rate law that describes the kinetics of the heparin-catalyzed reaction between antithrombin III and thrombin. *J. biol. Chem.* 258: 14708-14717 (1983).
- 13 Griffith, M.J.: The heparin-enhanced antithrombin III/thrombin reaction is saturable with respect to both thrombin and antithrombin III. *J. biol. Chem.* 257: 13899-13902 (1982).
- 14 Choay, J.; Petitou, M.; Lormeau, M.C.; Sinay, P.; Casu, B.J.; Gatti, G.: Structure activity relationship in heparin: a synthetic pentasaccharide with high affinity for antithrombin III and eliciting high anti factor X_a activity. *Biochem. biophys. Res. Commun.* 116: 492-499 (1983).
- 15 Holmer, E.; Kurachi, K.; Soderstrom, G.: The molecular weight dependence of the rate-enhancing effect of heparin on the inhibition of thrombin, factor X_a, factor IX_a, factor XI_a, factor XII_a and kallikrein by antithrombin III. *Biochem. J.* 193: 395-400 (1981).
- 16 Lane, A.; Denton, J.; Flynn, A.M.; Thunberg, L.; Lindahl, U.: Anticoagulant activities of heparin oligosaccharides and their neutralization by platelet factor 4. *Biochem. J.* 218: 725-732 (1984).
- 17 Tollefsen, D.M.; Blank, M.K.: Detection of a new heparin-dependent inhibitor of thrombin in human plasma. *J. clin. Invest.* 68: 589-596 (1981).
- 18 Marciniak, E.: Factor X_a inactivation by antithrombin III. Evidence for biological stabilization of factor X_a by factor-V-phospholipid complex. *Br. J. Haemat.* 24: 391-400 (1973).
- 19 Joso, F.; Béguin, S.: Changes in the antithrombin III activity at the interface plasma-phospholipids (Abstract). *Thromb. Haemostasis* 46: 285 (1981).
- 20 Ellis, V.; Scully, M.F.; Kakkar, V.V.: Inhibition of prothrombinase complex by plasma proteinase inhibitors. *Biochemistry* 23: 5882-5887 (1984).
- 21 Teitel, J.M.; Rosenberg, R.D.: Protection of factor X_a from neutralization by the heparin-antithrombin complex. *J. clin. Invest.* 71: 1383-1391 (1983).
- 22 Ockelford, P.; Carter, C.J.; Cerskus, A.; Smith, C.A.; Hirsh, J.: Comparison of the in vivo haemorrhagic and antithrombotic effects of a low antithrombin III affinity heparin fraction. *Thromb. Res.* 27: 679-690 (1982).
- 23 Barrowcliffe, T.W.; Merton, R.E.; Havercroft, S.J.; Thunberg, U.; Thomas, D.P.: Low affinity heparin potentiates the action of high affinity heparin oligosaccharides. *Thromb. Res.* 34: 125-133 (1984).
- 24 Walker, F.J.; Esmon, C.T.: Interactions between heparin and factor X_a. Inhibition of prothrombin activation. *Biochim. biophys. Acta* 585: 405-415 (1979).
- 25 Brown, J.E.; Baugh, R.F.; Hougie, C.: The inhibition of the intrinsic generation of activated factor X by heparin and hirudin. *Thromb. Res.* 17: 267-272 (1980).
- 26 Ofosu, F.A.; Blajchman, M.A.; Hirsh, J.: The inhibition by heparin of the intrinsic pathway activation of factor X in the absence of antithrombin III. *Thromb. Res.* 20: 391-403 (1980).

- 27 Ofosu, F.A.; Blajchman, M.A.; Modi, G.; Cerskus, A.L.; Hirsh, J.: Activation of factor X and prothrombin in antithrombin-III-depleted plasma: the effects of heparin. *Thromb. Res.* 23: 331-345 (1981).
- 28 Ofosu, F.A.; Modi, G.; Cerskus, A.L.; Hirsh, J.; Blajchman, M.A.: Heparin with low affinity to antithrombin III inhibits the activation of prothrombin in normal plasma. *Thromb. Res.* 28: 487-497 (1982).
- 29 Lecrubier, C.; Conard, J.; Horellou, M.H.; Kher, A.; Samama, M.: Spontaneous platelet aggregation in heparin-treated patients. *Acta Haemat.* 71: 63-65 (1984).
- 30 Kelton, J.G.; Hirsh, J.: Bleeding associated with antithrombotic therapy. *Semin. Hematol.* 17: 375-379 (1980).
- 31 Harker, L.A.; Molpass, T.W.; Branson, H.E.; Hessel, E.A.; Slichter, S.J.: Mechanism of abnormal bleeding in patients undergoing cardiopulmonary bypass: acquired transient dysfunction associated with selective alpha-granule release. *Blood* 56: 824-834 (1980).
- 32 Salzman, E.W.; Rosenberg, R.D.; Smith, M.H.; Lindon, J.N.; Favreau, L.: Effect of heparin and heparin fractions on platelet aggregation. *J. clin. Invest.* 65: 64-73 (1980).
- 33 Hirsh, J.: Heparin-induced bleeding. *Nouv. Revue fr. Hémat.* 26: 261-266 (1984).
- 34 Kakkar, V.V.; Djazaeri, D.B.; Webb, P.; Scully, M.; Westwick, T.; MacGregor, I.: Low molecular weight heparin and prevention of postoperative deep vein thrombosis. *Br. med. J.* 284: 375-379 (1982).
- 35 Huisse, M.G.; Guillin, M.C.; Bezeaud, A.; Toulemonde, F.; Kitzis, M.; Adreassian, B.: Heparin-associated thrombocytopenia. In vitro effect of different low molecular weight heparin fractions. *Thromb. Res.* 27: 485-490 (1982).
- 36 Henny, C.P.; Ten Cate, J.W.; van Bronswijck, H.; Ten Cate, H.; Surachino, S.; Wilmink, J.M.; Ockelford, P.A.: Use of a new heparinoid as anticoagulant during acute hemodialysis of patients with bleeding complications. *Lancet* i: 890-893 (1983).
- 37 Aiach, M.; Michaud, A.; Balian, J.L.; Lefebvre, M.; Woler, M.; Fourtillan, J.: A new low molecular weight heparin derivative. In vitro and in vivo studies. *Thromb. Res.* 31: 611-621 (1983).
- 38 Baruch, D.; Franssen, J.; Hemker, H.C.; Lindhout, T.: Effect of heparin and low molecular weight heparins on thrombin-induced blood platelet activation in the absence of antithrombin III. *Thromb. Res.* 38: 447-458 (1985).
- 39 Griffith, M.J.; Kingdon, H.S.; Lundblad, R.L.: The interaction of heparin with human alpha-thrombin: effect on the hydrolysis of aniline tripeptide substrates. *Archs Biochem. Biophys.* 195: 378-384 (1979).
- 40 Van de Waart, P.; Bruls, H.; Hemker, H.C.; Lindhout, T.: Interaction of bovine blood clotting factor V_a and its subunit with phospholipid vesicles. *Biochemistry* 22: 2427-2432 (1983).
- 41 Kim, B.K.; Steiner, M.; Baldini, M.G.: Antithrombin III binds to human platelets. *Biochem. Biophys. Res. Commun.* 97: 1227-1232 (1980).
- 42 Hurst, R.E.; Poon, M.C.; Griffith, M.J.: Structure-activity relationships of heparin. Independence of heparin charge density and antithrombin-binding domains in thrombin inhibition by antithrombin and heparin cofactor II. *J. clin. Invest.* 72: 1042-1045 (1983).
- 43 Heuck, C.C.; Schiele, U.; Horn, D.; Fronda, D.; Ritz, E.: The role of surface charge in the accelerating action of heparin on the antithrombin-III-inhibited activity of thrombin. *J. biol. Chem.* 260: 4598-4603 (1985).
- 44 Fischer, A.M.; Barrowcliffe, T.W.; Thomas, D.P.: A comparison of pentosan polysulphate and heparin. I. Mechanism of action on blood coagulation. *Thromb. Haemostasis* 47: 104-108 (1982).

Dr. D. Baruch,
Department of Biochemistry,
Biomedical Center,
University of Limburg,
PO Box 616,
NL-6200 MD Maastricht (The Netherlands)