

Characterization of Two Forms of Human Factor Va with Different Cofactor Activities

Citation for published version (APA):

Rosing, J., Bakker, H. M., Christella, M., Thomassen, L. G. D., Hemker, H. C., & Tans, G. (1993). Characterization of Two Forms of Human Factor Va with Different Cofactor Activities. Journal of Biological Chemistry, 268(28), 21130-21136. https://doi.org/10.1016/S0021-9258(19)36902-9

Document status and date: Published: 05/10/1993

DOI: 10.1016/S0021-9258(19)36902-9

Document Version: Publisher's PDF, also known as Version of record

Please check the document version of this publication:

 A submitted manuscript is the version of the article upon submission and before peer-review. There can be important differences between the submitted version and the official published version of record. People interested in the research are advised to contact the author for the final version of the publication, or visit the DOI to the publisher's website.

• The final author version and the galley proof are versions of the publication after peer review.

 The final published version features the final layout of the paper including the volume, issue and page numbers.

Link to publication

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these riahts.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
 You may freely distribute the URL identifying the publication in the public portal.

If the publication is distributed under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license above, please follow below link for the End User Agreement:

www.umlib.nl/taverne-license

Take down policy

If you believe that this document breaches copyright please contact us at:

repository@maastrichtuniversity.nl

providing details and we will investigate your claim.

Characterization of Two Forms of Human Factor Va with Different Cofactor Activities*

(Received for publication, April 19, 1993, and in revised form, June 18, 1993)

Jan Rosing‡, Harry M. Bakker, M. Christella L. G. D. Thomassen, H. Coenraad Hemker, and **Guido Tans**

From the Department of Biochemistry, Cardiovascular Research Institute, University of Limburg, Maastricht, The Netherlands

ibc

lyzed prothrombin activation. Purified human factor Va appears to consist of a heavy chain $(M_{\rm r} \approx 105,000)$ and a light chain doublet with $M_{\rm r} \approx 74,000$ and \approx 71,000. We separated factor Va by chromatography on a Mono-S column into two fractions, designated factors Va₁ and Va₂. Factor Va₁ contains the light chain with $M_{\rm r} \approx 74,000$, and factor Va₂ exclusively contains the light chain with $M_{\rm r} \approx 71,000$. The two forms of factor Va express different cofactor activities when prothrombin is activated at low phospholipid concentrations or on membranes containing low amounts of phosphatidylserine in phosphatidylcholine. Compared with factor Va₂, much higher amounts of factor Va₁ are required for factor Xa. Va complex formation at the membrane surface. Once incorporated into the prothrombinase complex, factors Va1 and Va2 are equally active in prothrombin activation. This indicates that the two forms of factor Va do not differ in their ability to promote the catalytic activity of factor Xa or to interact with prothrombin. Direct binding experiments show that the different cofactor activities are explained by a greatly impaired ability of factor Va_1 to bind to negatively charged membranes. Factor V is also separated into two protein peaks after chromatography on a Mono-S column. Upon incubation with thrombin, the first peak yields factor Va₁ and the second peak factor Va₂. The same two forms of factor Va were generated when freshly prepared plasma samples or platelet suspensions were treated with thrombin. This shows that the heterogeneity of the light chain domain is an intrinsic property of both plasma and platelet factor V. It is hypothesized that the heterogeneity is caused by small differences in the carboxylterminal C2 domain of factor V that are introduced as the result of post-ribosomal processing.

Factor Va is an essential cofactor in factor Xa-cata-

Activated blood coagulation factor V (factor Va) plays an essential role as cofactor in the conversion of prothrombin into thrombin by the serine protease factor Xa. Factor Va accelerates factor Xa-catalyzed prothrombin activation four to five orders of magnitude by causing a 1000-fold increase of the catalytic capacity (k_{cat}) of factor Xa (1, 2) and by promoting the binding of both factor Xa and prothrombin to procoagulant membranes (1, 3, 4).

Factor Va is generated from the pro-cofactor, factor V, which circulates in plasma as a single chain glycoprotein with M_r of about 330,000 (5, 6). Factor V has little or no procoagulant activity (1) and is converted into factor Va after proteolysis of specific peptide bonds, which, among others, can be catalyzed by thrombin (5–7). Thrombin-activated factor V is a two-chain molecule (7), that consists of a heavy and a light chain held together by a tightly bound calcium ion (8, 9). The heavy chain is derived from the amino-terminal region of factor V and has an M_r of 105,000. The light chain originates from the carboxyl-terminal region and appears on SDS-polyacrylamide gels as a doublet with apparent M_r of 71,000 and 74,000, both in case of human (7) and bovine factor Va (10, 11).

There is as yet no good explanation for the heterogeneity of the light chain of factor Va. It is likely, however, that the structural basis for the light chain doublet has to be sought in its carboxyl-terminal region, since large parts of the aminoterminal region of the light chain (with M_r up to 62,000) appear as single bands after SDS-gel electrophoretic analysis (11, 12).

Downloaded from www.jbc.org at UKB - Universiteit Maastricht on July 27, 2007

In this paper, we show that the heterogeneity of the carboxyl-terminal domain of the light chain region of factor Va is already observed in factor V present in fresh plasma and platelet samples. We further report the separation of factor Va into two fractions that are homogeneous with respect to their light chain composition. The factor Va preparation with the M_r 71,000 light chain is characterized by a more efficient incorporation in the prothrombinase complex than the factor Va that contains the light chain with $M_r = 74,000$. This difference in cofactor function appears to be due to different binding affinities for procoagulant membranes.

EXPERIMENTAL PROCEDURES

Materials-Bovine serum albumin, chicken egg albumin (ovalbumin), soybean trypsin inhibitor (type IS), bovine brain PS,¹ egg-yolk PC, Russel's viper venom, Hepes, and Tris were purchased from Sigma. DOPC and DOPS were obtained from Avanti Polar Lipids, Pelham, AL. S2238 (D-Phe-(pipecolyl)-Arg-pNA), S2337 (Ile-Glu-(piperidyl)-Gly-Arg-pNA), and I2581 (N-dansyl-(p-guanidino)-phenylalanine-piperidide hydrochloride) were supplied by AB Kabi Diagnostica, Stockholm, Sweden, PPACK (D-Pro-Phe-Arg-CH₂Cl) was obtained from Calbiochem, and p-NPGB was from Nutritional Biochemicals. Column materials and FPLC equipment used for protein

^{*} This work was supported by Program Grant 900-526-192 from the Dutch Organization for Scientific Research (NWO). The costs of publication of this article were defrayed in part by the payment of page charges. This article must therefore be hereby marked "advertisement" in accordance with 18 U.S.C. Section 1734 solely to indicate this fact.

[‡] To whom correspondence should be addressed: Dept. of Biochemistry, University of Limburg, P. O. Box 616, 6200 MD Maastricht, The Netherlands. Tel.: 31-43-881678; Fax: 31-43-670988.

¹ The abbreviations used are: PS, phosphatidylserine; PC, phosphatidylcholine; DOPS, 1,2-Dioleoyl-sn-glycero-3-phosphoserine; DOPC, 1,2-dioleoyl-sn-glycero-3-phosphocholine; BSA, bovine serum albumin; FPLC, fast liquid protein chromatography; PAGE, polyacrylamide gel electrophoresis; p-NPGB, p-nitrophenyl-p'-guanidinobenzoate hydrochloride; RVV-X, purified factor X activator from Russel's viper venom.

purification were purchased from Pharmacia, Uppsala, Sweden.

Proteins-The human coagulation factors used in this study were purified from fresh frozen plasma. Human prothrombin and factor X were purified according to DiScipio et al. (13). Human thrombin was prepared from prothrombin activation mixtures by the method of Pletcher and Nelsestuen (14). Human factor Xa was obtained from purified factor X after activation with RVV-X and isolation from the activation mixture by affinity chromatography on soybean trypsin inhibitor-Sepharose (15). RVV-X was purified from Russel's viper venom according to Schiffman et al. (16). Human factor V was purified essentially as described by Dahlbäck (5) with minor modifications (17). Factor Va was prepared by incubating factor V (0.3 mg/ ml) for 20 min with 30 nM thrombin in a buffer containing 10 mM Hepes pH 7.5, 50 mM NaCl, and 5 mM CaCl₂. After activation, 60 nM PPACK was added to inhibit thrombin. The factor Va preparation was subsequently applied to a 1-ml prothrombin-CL4B-Sepharose column and eluted with a linear gradient of 50-500 mM NH4Cl in 10 mM Hepes pH 7.5. Factor Va eluted at 260 mM NH4Cl. The factor Va-containing fractions were pooled and diluted in a buffer containing 25 mM Hepes pH 7.5, 100 mM NH₄Cl, 5 mM CaCl₂, and 5 mg/ml BSA. Prothrombin, factor Xa, factor V, and factor Va were stored at -80 °C. Protein preparations were homogeneous and >95% pure as judged by SDS-PAGE according to Laemmli (18).

Separation of the Two Forms of Factor Va (Va₁ and Va₂)-The factor Va obtained from the prothrombin-CL4B-Sepharose column (see previous paragraph) was further subjected to fast protein liquid chromatography (FPLC) on a Mono-S column (HR 5/5) at room temperature. The flow rate during the whole purification procedure was 0.5 ml/min. After application of 0.6-mg factor Va, the column was washed with 15 ml of a buffer containing 25 mM Hepes pH 7.5, 50 mM NH4Cl, and 5 mM CaCl₂. No factor Va activity eluted during the application and wash procedure. The Mono-S column was then developed with 15 ml of a linear gradient (50-1000 mM NH₄Cl) in the same buffer. Factor Va activity eluted from the column in two well separated protein peaks at 450 and 750 mM NH₄Cl. The first factor Va peak contained the light chain with $M_r = 74,000$ (factor Va₁), and the second peak consisted of factor Va with a light chain with $M_r = 71,000$ (factor Va₂). The two factor Va-containing peaks were pooled separately and contained pure and homogeneous factor Va as judged by SDS-PAGE according to Schägger and von Jagow (19). In a large number of factor Va preparations isolated from different batches of human plasma, the amount of protein present in peak 1 was about half that present in peak 2.

Separation and identification of different factor V forms present in purified factor V preparations and in plasma and platelet samples were accomplished by the same procedure described in the legends to the figures.

Gel Electrophoretic Techniques—Factor Va preparations and the activation patterns of factor V activated with thrombin were analyzed by SDS-PAGE in the presence of Tricine according to Schägger and von Jagow (19). This electrophoretic technique has the advantage that the separation of protein bands in the gel is hardly affected by the ionic strength of the samples.

Protein Concentrations—Protein concentrations were determined according to Lowry *et al.* (20). Molar thrombin and factor Xa concentrations were determined by active site titration with *p*-NPGB (21, 22). Prothrombin concentrations were determined after complete activation of prothrombin with *Echis carinatus* venom and quantitation of thrombin with *p*-NPGB. Factor V concentrations were estimated from the absorbance at 280 nm using an A_{280} 1% of 8.9 (6). Factor Va concentrations were determined as described by Lindhout *et al.* (3).

Phospholipid Vesicle Preparations—Appropriate quantities of phospholipids dissolved in CHCl₃/CH₃OH (1/1, v/v) were mixed in a glass tube and dried under a mild flow of N₂. The phospholipids were suspended in 2 ml of buffer (25 mM Hepes pH 7.5, 175 mM NaCl) and vigorously vortexed for 1 min. The phospholipid suspension was subsequently sonicated for 10 min at 0 °C using an MSE Mark II 150 W ultrasonic disintegrator set at 8 μ m peak-to-peak amplitude. Phospholipid concentrations were determined by phosphate analysis (23).

Factor Va Assay—Factor Va was quantitated by measuring its cofactor activity in factor Xa-catalyzed prothrombin activation at saturating concentrations of factor Xa, phospholipid vesicles, and prothrombin and at a limiting amount of factor Va (17). The factor Va concentration in the assay mixture was calculated from the rate of prothrombin activation using a turnover number of 6000 mol of prothrombin activated/min/mol of factor Xa · Va complex (24).

It should be emphasized that the quantitation of the two forms of

factor Va (Va₁ and Va₂) that are studied in this paper is correct despite the fact that there are reaction conditions at which they express different cofactor activities in prothrombin activation. This is inherent to the fact that the high concentrations of factor Xa, prothrombin, and phospholipids used in the assay system nullify the differences in cofactor activities (see "Results," Table I).

Kinetic Data Analysis-Complex formation between membranebound factors Xa and Va was determined by measuring the rate of prothrombin activation in the presence of phospholipid vesicles at a fixed limiting concentration of factor Xa and varying amounts of factor Va. Factor Va, factor Xa, and phospholipid vesicles were preincubated for 5 min at 37 °C in 25 mM Hepes pH 7.5, 175 mM NaCl, 2 mM CaCl₂, and 5 mg/ml BSA. Activation was started by addition of prothrombin (preincubated at 37 °C in the same buffer). Rates of prothrombin activation were determined with the chromogenic substrate S2238 (2). The apparent K_d for dissociation of the membrane-bound factor Xa · Va complex $(K_{i_{2}Va})$ and the prothrombinconverting activity of this complex at [factor Va] $\rightarrow \infty$ ($V_{Va \rightarrow \infty}$) were obtained from a plot of the rate of prothrombin activation as a function of the factor Va concentration that was fitted to the equation for a single site binding isotherm (hyperbola) in the computer program "Enzfitter.

The kinetic parameters of factor Xa-catalyzed prothrombin activation $(K_m \text{ for prothrombin and } V_{\max} \text{ of prothrombin activation})$ were determined by measuring the rate of thrombin formation at varying prothrombin concentrations in the presence of a fixed phospholipid concentration, a limiting amount of factor Xa, and a saturating concentration factor Va under conditions described in the previous paragraph. The kinetic parameters were obtained by fitting the data to the Michaelis-Menten equation using the Enzfitter computer program.

Binding Studies—Binding parameters for factor Va-membrane association were determined on planar phospholipid bilayers. The concentrations of membrane-bound factor Va were determined by ellipsometry (25), and the free factor Va concentrations were determined with the functional assay described above. Amounts of phospholipid-bound factor Va were plotted as a function of the free factor Va concentration, and the binding parameters (K_d and number of membrane binding sites) were obtained by fitting the data to the equation for a single site binding isotherm in the computer program Enzfitter.

RESULTS

Separation of Two Forms of Factor Va by FPLC on a Mono-S Column—Purified human factor V was converted into factor Va with thrombin. Factor Va was subsequently separated from the activation fragments by chromatography on a prothrombin-Sepharose column (cf. Ref. 26). SDS-gel electrophoretic analysis showed that this procedure yielded a pure factor Va preparation that consisted of a heavy chain with M_r of 105,000 and the characteristic light chain doublet migrating in the 71,000–74,000 molecular weight region (Fig. 1, lane 2).

Although anion exchangers are usually employed in the purification of factor V and Va, it appears that the light chain of factor Va is a basic protein with a high affinity for cation exchangers (3, 11). Fig. 2 shows that factor Va also bound to a Mono-S column. Subsequent elution with a NH₄Cl gradient yielded two protein peaks with factor Va activity. SDS-PAGE analysis showed that the first peak consisted of factor Va molecules with a light chain of $M_r = 74,000$ (Fig. 1, fractions 79-81) and that the factor Va that eluted in the second peak contained the light chain with $M_r = 71,000$ (Fig. 1, fractions 87-90). In the text that follows, we will designate the factor Va present in the first peak, factor Va₁, while the factor Va present in the second peak will be called factor Va₂.

Separation of Two Forms of Factor V on a Mono-S Column—Nonactivated human factor V showed a chromatographic behavior that strongly resembled factor Va. Factor V also bound to the Mono-S column and eluted as a double peak when the column was developed with an NH₄Cl gradient (Fig. 3A). Incubation of the protein fractions with thrombin yielded two factor Va activity peaks, which, as shown by SDS-PAGE



FIG. 1. Gel electrophoretic analysis of factor V and Va preparations. Factor V, thrombin-activated factor V, and factors Va₁ and Va₂ that eluted from the Mono-S column were subjected to SDS-PAGE on slab gels containing 7.5% acrylamide (19). 20 μ l of a factor V- or Va-containing sample (0.6-5 μ g protein) was mixed with 5 μ l of electrophoresis sample buffer and applied to the gel. Lane 1, factor V; lane 2, thrombin-activated factor V (factor Va); fractions 79-81 = factor Va₁; fractions 87-90 = factor Va₂. Gels were stained with Coomassie Brilliant Blue R-250.



FIG. 2. Chromatography of factor Va on Mono-S. Factor Va obtained from prothrombin CL 4B-Sepharose was diluted to 0.6 mg/ 20 ml in a buffer containing 25 mM Hepes pH 7.5, 50 mM NH₄Cl, and 5 mM CaCl₂ and was applied to a Mono-S column (HR 5/5) that was equilibrated with the same buffer. After application of factor Va, the column was washed with 15 ml of buffer. Factor Va was subsequently eluted from the column with 15 ml of a linear gradient (50-1000 mM NH₄Cl) in 25 mM Hepes pH 7.5, 5 mM CaCl₂. Fractions (0.5 ml) were collected and assayed for factor Va activity ($\textcircled{\bullet}$) as described under "Experimental Procedures." Fractions from peak 1 (79-81) and peak 2 (87-90) were subjected to SDS-PAGE (Fig. 1).

analysis (Fig. 3B), were highly enriched in the 74,000 and 71,000 Da light chains. This experiment indicates that the light chain doublet present on SDS gels of factor Va preparations does not result from additional peptide bond cleavages that occur during the activation of factor V by thrombin but shows that the heterogeneity in the light chain domain is already present in factor V.

Structural Properties of Plasma and Platelet Factor V Present in Freshly Collected Blood Samples—The experiments presented in the previous paragraph strongly suggest that purified factor V preparations already contain two forms of factor V. However, this does not necessarily mean that there are also two forms of factor V circulating in plasma, since it is possible that the heterogeneity of purified factor V preparations results from reactions that occurred during the purification procedure. In order to exclude this possibility, we performed the following experiment. Freshly collected blood samples were subjected to two centrifugation steps, which successively removed the erythrocytes and blood platelets and yielded platelet-poor plasma. The platelet pellet was resuspended in buffer, and thrombin was added to both the platelet-poor plasma and to the platelet suspension. The generation of factor Va activity was followed, and, after complete activation of factor V, the plasma and platetet suspensions were diluted and applied to a Mono-S column. Development of this column with a linear NH₄Cl gradient yielded, both for the plasma (Fig. 4) and the platelet sample (data not shown), two factor Va activity peaks that eluted between 450–500 and 700–750 mM NH₄Cl, respectively.

From the fact that the time interval between taking the blood samples and separating the plasma and platelet factor Va samples on the Mono-S column was only 50 min and since the activity ratios of the two factor Va peaks in the plasma and platelet samples were approximately the same as those observed in purified factor V(a) preparations (the purification of which took 4 days), we conclude that the heterogeneity of factor V does not result from reactions that have occurred during the purification procedure. The presence of two forms of factor V in fresh plasma and platelet samples indicates that the heterogeneity is an intrinsic property of factor V.

Cofactor Activities of the Two Forms of Factor Va—Expression of the cofactor activity of factor Va is dependent on direct interactions with prothrombin, factor Xa, and phospholipid. In the factor Va assay (see "Experimental Procedures"), the reaction conditions are chosen such that factor Va is the limiting factor and all other components are present in excess. This means that under these conditions, differences in protein-protein or protein-lipid interactions between the two forms of factor Va may nullify and escape attention.²

To compare the cofactor activities of the two forms of factor Va, we have, therefore, performed a kinetic analysis of the stimulatory effects of the factor Va preparations on prothrombin activation under suboptimal reaction conditions, i.e. at a limiting amount of factor Xa and at a low concentration of phospholipid vesicles containing a low mole percentage of PS (2 µM DOPS/DOPC, 5/95, mol/mol). Fig. 5 shows that there was a rather large difference between the amounts of factors Va1 and Va2 required for the assembly of a functionally active membrane-bound factor Xa. Va complex. Half-maximal prothrombinase complex formation was observed at 5.93 nM factor Va1 and 0.45 nM factor Va2. A double-reciprocal plot of these data (Fig. 5B) shows that the rates of prothrombin activation attained at saturating levels of factor Va were approximately the same (Table I), which indicates that the factor $Xa \cdot Va_1$ and $Xa \cdot Va_2$ complexes, once they are formed, have equal catalytic activities.

The differences in factor Xa · Va complex formation disappeared when the same experiment was performed with phospholipid vesicles that contained a high mole percentage of PS (*i.e.* 2 μ M PS/PC, 20/80 mol/mol). On these membranes, halfmaximal prothrombinase complex formation required 0.047 nM factor Va₁ and 0.044 nM factor Va₂ (Table I). This exper-

The Journal of Biological Chemistry

² In the factor Va assay, reaction conditions were such that factor Va was present at a limiting concentration, and all other components were added in excess. The phospholipid concentration and composition (100 μ M PS/PC, 10/90, mol/mol) and the high factor Xa concentration (5 nM) ensures full incorporation of factor Va into the membrane-bound factor Xa · Va complex. Complete incorporation of both factors Va₁ and Va₂ into the prothrombinase complex is shown by the fact that rates of prothrombin activation were not further increased when the prothrombin, phospholipid, and factor Xa concentrations in the assay mixture were increased. The validity of this assay for quantitating factor Va₁ and Va₂ preparations is also shown by the fact that preparations that were set at the same protein concentration on the basis of A₂₈₀ yielded the same concentrations in the functional assay.

FIG. 3. A, chromatography of factor V on Mono-S. 5 ml of purified factor V (5 mg) in 10 mM Hepes pH 7.5, 50 mM NaCl was applied to a Mono-S column (HR 5/5). The column was washed with 15 ml of a buffer containing 25 mM Hepes pH 7.5, 50 mM NH₄Cl, and 5 mM CaCl₂ and subsequently developed with 30 ml of a linear gradient from 50 to 1000 mM NH₄Cl in 25 mM Hepes pH 7.5 and 5 mM CaCl₂. Samples from the column fractions were incubated for 1 h at 37 °C with thrombin (final concentration, 25 nm) and the factor Va generated (\bullet) was assayed as described under "Experimental Procedures." B, gel electrophoretic analysis of factor Va generated in the factor V fractions that eluted from the Mono-S column. Aliquots (50 µl) from the column fractions 64, 66, 72, 76, 78, and 82) were activated with thrombin (Fig. 3A) and subsequently mixed wih 12.5 μ l of gel electrophoresis sample buffer (19). 20 μ l of the gel samples were then applied to a 7.5% slab gel and subjected to SDS-PAGE (19). Sm is a lane with starting material (factor V) activated with thrombin; M_r is a lane with molecular weight markers. The proteins on the gel were stained with Coomassie Brilliant Blue R-250.

The Journal of Biological Chemistry

ibc



FIG. 4. Chromatography of thrombin-activated plasma on a Mono-S column. Platelet-poor plasma derived from blood collected in citric acid was brought to 5 mM free CaCl₂ and incubated for 10 min at 37 °C with thrombin (final concentration 10 nM). The fibrin clot that was formed was removed, and 1 ml of the serum was diluted five times with column buffer (25 mM Hepes pH 7.5, 5 mM CaCl₂) and applied to a Mono-S column (HR 5/5). The column was washed with 13 ml of a buffer containing 25 mM Hepes pH 7.5, 150 mM NH₄Cl, and 5 mM CaCl₂, and the proteins bound to the column were subsequently eluted with 15 ml of a salt gradient from 50 to 1000 mM NH₄Cl in 25 mM Hepes pH 7.5 and 5 mM CaCl₂. Column fractions (0.5 ml) were assayed for factor Va activity (\bullet) as described under Experimental Procedures."

iment indicates that the different efficiencies by which factors Va1 and Va2 incorporate in the prothrombinase complex are likely compensated by an increased binding affinity and an increased number of binding sites for factors Xa and Va on membranes containing 20 mole percent of PS (3, 27, 28-31). This argumentation predicts that differences between factors Va1 and Va2 can also be observed on membranes with 20 mole percent of PS, provided that the experiment is performed at a lower phospholipid concentration. Indeed titration of a limited amount of factor Xa with factors Va1 or Va2 on 0.1 µM PS/PC (20/80, mol/mol) vesicles again showed a difference between the two forms of factor Va (Table I). At this phospholipid concentration, a 9-fold higher concentration of factor Va1 was required to obtain half-maximal rates of prothrombin activation (0.389 nM factor Va1 versus 0.044 nM factor Va₂).

With respect to the different functional properties of factors Va_1 and Va_2 , the possibility has to be ruled out that factor Va_1 lacks functional activity and that the cofactor activity of factor Va_1 is due to a small (~8%) contamination with factor Va_2 . This possibility is excluded by the observations that (*a*) the amounts of factor Va_1 and factor Va_2 required for factor $Xa \cdot Va$ complex formation on 2 μ M PS/PC (20/80) were the same (Table I), and (*b*) equal protein amounts of factors Va_1

1.0 1.0 1.0 0.8 0.8 0.8 0.6 0.6 0.6 [Factor Va] (µM) A280 nm 0.4 0.4 0.4 0.2 0.2 0.2 0.0 0.0 0.0 60 100 0 20 40 80 Fraction number В

66

64

Sm

72

Fraction

76

78

82

Mr

Α

- 92,000

- 67,000

- 43,000

30,000

- 21,500 - 14,400

Table I

Kinetic analysis of the functional activities of Factors Va1 and Va2

Cofactor activities of factors Va₁ and Va₂ were determined by measuring rates of prothrombin activation at a limiting factor Xa concentration in a reaction mixture containing 25 mM Hepes pH 7.5, 175 mM NaCl, 2 mM CaCl₂, 5 mg/ml BSA, 3 pM factor Xa, 1 μ M prothrombin, varying amounts of factors Va₁ or Va₂, and phospholipid vesicles as indicated in the Table. Rates of prothrombin activation were plotted against the factor Va concentrations and K_{WVa} and $V_{Va \to \infty}$ were obtained by fitting the data to a hyperbola (see Fig. 5). Further experimental conditions are given under "Experimental Procedures."

Phoenholinid	Factor Va ₁		Factor Va ₂	
rnosphoupia	K.,val	V _{Va→∞}	K _{MVa2}	$V_{Va \rightarrow \infty}$
	nM	пм IIa/min	nM	п м IIa/min
2 µm PS/PC (5/95)	5.93	6.35	0.45	6.15
2 µM PS/PC (20/80)	0.047	18.6	0.044	17.9
0.1 µM PS/PC (20/80)	0.39	12.8	0.044	13.5



FIG. 5. Comparison of cofactor activities of factors Va₁ and Va₂. Rates of prothrombin activation at limiting factor Xa were determined in a reaction mixture containing 25 mM Hepes pH 7.5, 175 mM NaCl, 2 mM CaCl₂, 5 mg/ml BSA, 2 μ M phospholipid vesicles (DOPS/DOPC, 5/95, mol/mol), 3 pM factor Xa, 1 μ M prothrombin, and amounts of factor Va₁ (\oplus) or factor Va₂ (\blacktriangle) indicated in the figure. Further experimental conditions are described under "Experimental Procedures." A, rate of prothrombin activation as a function of the concentration factor Va₁ or Va₂. The solid lines represent hyperbola obtained after fitting the data with $K_{16}(Va_1) = 5.93$ nM and $V_{Va \rightarrow \infty} = 6.35$ nM IIa/min (\bigoplus) or $K_{16}(Va_2) = 0.45$ nM and $V_{Va \rightarrow \infty} = 6.15$ nM IIa/min (\bigstar). B, double-reciprocal plots of the same data.

or Va_2 yield the same concentrations in the quantitative functional assay.²

Table II summarizes the kinetic parameters for prothrombin activation determined at saturating concentrations of factors Va₁ or Va₂. Similar values were obtained for the K_m for prothrombin and the V_{max} of prothrombin activation for both forms of factor Va. This indicates that factors Va₁ and Va₂ neither differ in their interaction with prothrombin nor in their capacity to promote the catalytic activity of factor Xa but exhibit considerable differences in their ability to assemble into a membrane-bound factor Xa · Va complex.

Membrane Binding Properties of Factors Va_1 and Va_2 —The impaired incorporation of factor Va_1 into the membrane-

TABLE II

Kinetic parameters of prothrombin activation

Rates of prothrombin activation at a limiting factor Xa concentration were determined in a reaction mixture containing 25 mM Hepes pH 7.5, 175 mM NaCl, 2 mM CaCl₂, 5 mg/ml BSA, 10 μ M phospholipid vesicles (DOPS/DOPC, 5/95, mol/mol), 3 pM factor Xa, 25 nM factor Va₁ or factor Va₂, and varying amounts of prothrombin. Kinetic parameters (K_m and V_{max}) were obtained by fitting the data to the Michaelis-Menten equation using the Enzfitter computer program. Further experimental conditions are given under "Experimental Procedures."

	K _m	Vmax	
	μM	mol/min/mol Xa	
Factor Va ₁	0.32	5386	
Factor Va ₂	0.16	5687	



FIG. 6. Binding of factors Va₁ and Va₂ to planar phospholipid bilayers. Binding of factors Va₁ (\bullet) and Va₂ (\blacktriangle) to planar phospholipid bilayers (DOPS/DOPC, 20/80, mol/mol) was determined by ellipsometry (25) in a reaction mixture containing 25 mM Hepes pH 7.5, 150 mM NH₄Cl, 5 mM CaCl₂, and 5 mg/ml BSA. The solid lines represent hyperbolas obtained by a non-linear least-squares fit of the data. The binding parameters describing these hyperbola are $K_d(Va_1) = 24$ nM and 0.397 µg factor Va₁ bound/cm² phospholipid at saturating factor Va₁ or $K_d(Va_2) = 0.53$ nM and 0.364 µg factor Va₂ bound/cm² phospholipid at saturating factor Va₂.

bound factor Xa. Va complex most likely results from a decreased affinity for either phospholipid or factor Xa. To distinguish between these possibilities, we have determined the affinity of both forms of factor Va for PS-containing membranes (Fig. 6). Factor Va₂ bound with high affinity to membranes composed of 20 mole percent of PS in PC. A hyperbolic fit of the binding data yielded a K_d of 0.53 nM and 0.364 μ g factor Va₂ bound/cm² phospholipid at saturating factor Va. Factor Va₁ had a much lower affinity for phospholipid, and, with the amounts of factor Va1 available, we were unable to add sufficiently high concentrations to achieve saturation of the membrane surface (Fig. 6). From a hyperbolic fit of the binding curve, it can be estimated, however, that the interaction between factor Va1 and the membrane is characterized by a K_d of about 24 nM and 0.397 μ g factor Va₁ bound/cm² phospholipid at saturating factor Va₁. From the molecular weights of factor Va (~180,000 Da) and the phospholipid used (~800 Da) and considering that 1 cm² of a bilayer phospholipid surface contains 0.4 μ g phospholipid, it can be calculated that phospholipid to factor Va ratios at saturation are 227 and 247 for factors Va1 and Va2, respectively.

These observations show that the two forms of factor Va differ considerably in their affinity for phospholipids, while the number of binding sites for factors Va₁ and Va₂ on the phospholipid surface are the same. The large affinity difference very likely explains the different cofactor activities of factors Va₁ and Va₂ in prothrombin activation.

DISCUSSION

Purified human factor Va preparations have been reported to consist of a heavy chain with $M_r \approx 105,000$ and a doublet

light chain with $M_r \approx 71,000$ and $\approx 74,000$ (7). The data presented in this paper show that it is possible to bind factor Va to a Mono-S column and to separate two forms of factor Va by developing the column with an NH₄Cl gradient. The peak eluting at the low salt concentration consisted of factor Va (Va₁) with $M_r \approx 74,000$ light chain, whereas factor Va with $M_r \approx 71,000$ light chain (factor Va₂) eluted at a much higher salt concentration. It is likely that the factor Va bound to the Mono-S column via its light chain, since this subunit is considered to be a basic protein that exhibits a high affinity for cation exchangers (3, 11). The elution profile from the Mono-S column is, therefore, indicative of a charge difference between the light chains of factors Va₁ and Va₂, in the sense that the light chain of factor Va₂ is more positively charged than the light chain of factor Va₁.

The interaction of factor Va with procoagulant membranes appears to depend on both hydrophobic interactions (32–35) and on electrostatic interactions between the positively charged light chain and negatively charged phospholipids (27, 30, 36). Considering that electrostatic interactions are important for factor Va-membrane association and that there may be a considerable charge difference between the light chains of the two factor Va forms, we hypothesized that factor Va₁ may exhibit a lower affinity for negatively charged membranes than factor Va₂. Direct binding experiments show that this is indeed the case. Factor Va_2 binds with a high affinity ($K_d =$ 0.53 nm) to PS-containing membranes (PS/PC, 20/80), whereas the affinity of factor Va_1 for such membranes is much less ($K_d \approx 24$ nM). The lipid/factor Va ratios (n) at protein saturation were approximately the same (n = 227 for factor) Va_1 and n = 247 for factor Va_2), which shows that the membrane surface contains the same number of binding sites for both forms of factor Va. It is difficult to compare these binding parameters with literature values, since the interaction of human factor Va with membranes is not well documented, whereas a wide range of binding parameters ($K_d =$ 0.05-10 nm and n = 42-250) is reported to describe the interaction of bovine factor Va with PC membranes that contain 25 mole percent of PS (27, 29-31, 36).

Factor Va plays an important role in both the assembly and expression of the catalytic activity of the prothrombinase complex on procoagulant membrane surfaces (1-4). Since proper cofactor function requires the binding of factor Va to the membrane, it is to be expected that the different affinities of factors Va1 and Va2 for negatively charged membranes will result in different efficiencies to act as a cofactor in prothrombin activation. Kinetic experiments showed that the cofactor function of factor Va₁ is indeed impaired. Much higher amounts of factor Va₁ were required for full assembly of the prothrombinase (factor Xa.Va) complex at the membrane surface. Once formed, the factor Xa · Va1 and Xa · Va2 complexes exhibit the same catalytic activity in prothrombin activation. This can be inferred from the fact that the kinetic parameters $(k_{cat} \text{ and } K_m)$ for prothrombin activation are similar (Table II). Differences in efficiencies by which factors Va_1 and Va_2 assemble into a membrane-bound factor $Xa \cdot Va$ complex were observed only on membranes that contain low amounts of PS or when prothrombin is activated at low phospholipid concentrations. This is not surprising if one considers that factor Xa. Va complex formation on the phospholipid surface is a reflection of the concentrations of surface-bound factors Va and Xa (3, 37). At high mole percentages of PS and at high phospholipid concentrations, differences between factors Va1 and Va2 will be nullified when the concentration of factor Va binding sites exceeds the K_d values for the membrane-factor Va complexes, since this condition

results in virtually complete binding of both forms of factor Va.

The heterogeneity in the light chains of factor Va is not a unique property of factor Va but is already present in factor V. It appears that the purified factor V preparations and even freshly prepared plasma samples and platelet suspensions contain two forms of factor V, which, after activation with thrombin, are converted into factors Va1 and Va2. This shows that the heterogeneity is an intrinsic property of factor V and is not caused by reactions occurring during the purification procedure or to additional peptide bond cleavages that take place when factor V is activated with thrombin.

Our experiments give no direct insight to the structural basis underlying the difference between the light chain regions of the two forms of factor V. However, combination of information presented in this paper with data recently reported by Ortel et al. (12) strongly suggests that the observed heterogeneity originates from the carboxyl-terminal C2 domain of the light chain region of factor V. Ortel et al. (12) isolated recombinant human factor V and showed that activation with thrombin resulted in the formation of a light chain doublet. Activation of mutant molecules that lack the carboxyl-terminal C2 domain (i.e. amino acid residues 2037-2196) yield a factor Va molecule with a single truncated light chain with $M_r \approx 62,000$. Deletion of the C2 domain caused an essentially complete loss of the ability of the mutant factor Va molecule to bind to negatively charged membranes. This indicates that the occurrence of two forms of factor Va (Va₁ and Va₂) with different affinities for procoagulant membranes results from the presence of two light chains with structural differences in the C2 domain. The fact that heterogeneity is also observed in the factor V present in freshly prepared platelet and plasma samples and in recombinant factor V (12) strongly suggests that the heterogeneity is due to differences in post-ribosomal processing or post-ribosomal modification (e.g. glycosylation) of the two factor V molecules. Whether the presence of two forms of factor V in plasma has a physiological function remains to be clarified.

Acknowledgments-We thank Rita Janssen and Dr. George Willems for performing the binding experiments on the ellipsometer.

REFERENCES

- 1. Nesheim, M. E., Taswell, J. B., and Mann, K. G. (1979) J. Biol. Chem. 254, 10952-10962
- 254, 10952-10962
 Rosing, J., Tans, G., Govers-Riemslag, J. W. P., Zwaal, R. F. A., and Hemker, H. C. (1980) J. Biol. Chem. 255, 274-283
 Lindhout, T., Govers-Riemslag, J. W. P., van de Waart, P., Hemker, H. C., and Rosing, J. (1982) Biochemistry 21, 5494-5502
 van Rijn, J. L. M. L., Govers-Riemslag, J. W. P., Zwaal, R. F. A., and Rosing, J. (1984) Biochemistry 23, 4557-4564
 Dahlbäck, B. (1980) J. Clin. Invest. 66, 583-591
 Kane, W. H., and Majerus, P. W. (1981) J. Biol. Chem. 256, 1002-1007
 Suzuki, K., Dahlbäck, B., and Stenflo, J. (1982) J. Biol. Chem. 257, 6556-6564

- 6564
- Esmon, C. T. (1979) J. Biol. Chem. 254, 964–973 Guinto. E. R., and Esmon, C. T. (1982) J. Biol. Chem. 257, 10038–10043 Nesheim, M. E., Foster, W. B., Hewick, R., and Mann, K. G. (1984) J. Biol.
- 10. Chem. 259, 3187-3196
- Odegaard, B., and Mann, K. (1987) J. Biol. Chem. 262, 11233-11238
 Ortel, T. L., Devore-Carter, D., Quinn-Allen, M., and Kane, W. H. (1992) J. Biol. Chem. 267, 4189-4198
- DiScipio, R. G., Hermodson, M. A., Yates, S. G., and Davie, E. W. (1977) Biochemistry 16, 698-706
 Pletcher, C. H., and Nelsestuen, G. L. (1982) J. Biol. Chem. 257, 5342-
- 5345
- Bock, P. E., Craig, P. A., Olsen, S. T., and Singh, P. (1989) Arch. Biochem. Biophys. 273, 375–388 16. Schiffman, S., Theodor, J., and Rapaport, S. J. (1969) Biochemistry 8,
- 1397-1405 17. Tans, G., Rosing, J., Thomassen, M. C. L. G. D., Heeb, M. J., Zwaal, R. F.

- Ians, G., Rosing, J., Hormassen, M. C. L. G. D., Heeb, M. J., Zwaai, K. F. A., and Griffin, J. H. (1991) Blood 77, 2641–2648
 Laemmli, U. K. (1970) Nature 227, 680–685
 Schägger, H., and von Jagow, G. (1987) Anal. Biochem. 166, 368–379
 Lowry, O. H., Rosebrough, N. J., Farr, A. L., and Randall, R. J. (1951) J. Biol. Chem. 102, 265–275. Biol. Chem. 193, 265–275 hase, T., Jr., and Shaw, E. (1969) Biochemistry 8, 2212–2224
- 21. Chase,
- 22. Smith, R. L. (1973) J. Biol. Chem. 248, 2418-2423

- 23. Böttcher, C. J. F., van Gent, C. M., and Pries, C. (1961) Anal. Chim. Acta 24, 203-207

- 24, 203-207
 24. Tans, G., Janssen-Claessen, T., Hemker, H. C., Zwaal, R. F. A., and Rosing, J. (1991) J. Biol. Chem. 266, 21864-21873
 25. Cuypers, P. A., Corsel, J. W., Janssen, M. P., Kop, J. M. M., Hermens, W. T., and Hemker, H. C. (1983) J. Biol. Chem. 258, 2426-2431
 26. Guinto, E. R., and Esmon, C. T. (1984) J. Biol. Chem. 259, 13986-13992
 27. van de Waart, P., Bruls, H., Hemker, H. C., and Lindhout, T. (1983) Biochemistry 22, 2427-2432
 28. Nelsestuen, G. L., and Broderius, M. (1977) Biochemistry 16, 4172-4177
 29. Bloom, J. W., Nesheim, M. E., and Mann, K. G. (1979) Biochemistry 18, 4419-4425

- Pusey, M. L., Mayer, L. D., Jason Wei, G., Bloomfield, V. A., and Nelsestuen, G. L. (1982) Biochemistry 21, 5262-5269
 Krishnaswamy, S., and Mann, K. G. (1988) J. Biol. Chem. 258, 6503-6508
 Lecompte, M. F., Krishnaswamy, S., Mann, K. G., Nesheim, M. E., and Gitler, C. (1987) J. Biol. Chem. 262, 1935-1937
 Krieg, U. C., Isaacs, B. S., Yemul, S. S., Esmon, C. T., Bailey, H., and Johnson, A. E. (1987) Biochemistry 26, 103-109
 Kalafatis, M., Jenny, R. J., and Mann, K. G. (1984) Biochemistry 23, 6202-6210
 Fusey, M. L., and Nelsestuen, G. L. (1984) Biochemistry 23, 6202-6210
 Krishnaswamy, S., Jones, K. C., and Mann, K. G. (1988) J. Biol. Chem. 263, 3823-3834