Laboratory assessment of antithrombotic therapy

Citation for published version (APA):

Kher, A., Al Dieri, R., Hemker, H. C., & Beguin, S. (1997). Laboratory assessment of antithrombotic therapy: What tests and if so why? *Haemostasis*, 27(5), 211-218. <https://doi.org/10.1159/000217459>

Document status and date:

Published: 01/01/1997

DOI:

[10.1159/000217459](https://doi.org/10.1159/000217459)

Document Version:

Publisher's PDF, also known as Version of record

Please check the document version of this publication:

- A submitted manuscript is the version of the article upon submission and before peer-review. There can be important differences between the submitted version and the official published version of record. People interested in the research are advised to contact the author for the final version of the publication, or visit the DOI to the publisher's website.
- The final author version and the galley proof are versions of the publication after peer review.
- The final published version features the final layout of the paper including the volume, issue and page numbers.

[Link to publication](#)

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal.

If the publication is distributed under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license above, please follow below link for the End User Agreement:

www.umlib.nl/taverne-license

Take down policy

If you believe that this document breaches copyright please contact us at:

repository@maastrichtuniversity.nl

providing details and we will investigate your claim.

André Kher^a
Raed Al Dieri^b
H. Coenraad Hemker^b
Suzette Béguin^b

Laboratory Assessment of Antithrombotic Therapy: What Tests and If So Why?

^a Euthemis, Saint Mandé, France;

^b Department of Biochemistry,
Cardiovascular Research Institute
and Medical Faculty, Maastricht
University, Maastricht,
The Netherlands

Abstract

A critical review is given of the tests available for the assessment of the action of anticoagulants, such as heparins, oral anticoagulants and direct thrombin inhibitors, in patients under antithrombotic therapy. The principle of action and the performance of the thromboplastin time (PT), the activated partial thromboplastin time (aPTT), the whole blood clotting time, the thrombin time, the ecarin clotting time and the endogenous thrombin potential (ETP) is discussed, as well as the evidence behind the accepted therapeutic ranges. The two most common tests, PT and aPTT, respond in an essentially different way to clinically effective anticoagulation with heparin and with oral anticoagulants. This means that they covariate with, but do not themselves represent the essential parameter influenced by anticoagulation. The experimental basis for the widely accepted two times prolongation of the aPTT as an indicator for adequate anticoagulation is shown to be meagre in the case of unfractionated heparin and lacking for the other anticoagulants. Common sources for error in the interpretation of anti-factor Xa- and anti-thrombin activity of heparins are indicated. Extensive experience with new tests like the ecarin clotting time and the ETP is still lacking. On the basis of preliminary data and in view of the importance of the enzymatic action of thrombin in the pathogenesis of thrombosis, the ETP is considered a possible candidate for a common parameter to assess different types of anticoagulants.

KARGER

Fax +41 61 306 12 34
E-Mail karger@karger.ch
www.karger.com

© 1998 S. Karger AG, Basel
0301-0147/97/0275-0211\$15.00/0

This article is also accessible online at:
<http://BioMedNet.com/karger>

Prof. Dr. H.C. Hemker
Maastricht University, Dept. of Biochemistry
Faculty of Medicine II, Level 4, Room 4342
Universiteitssingel 50
NL-6220 ER Maastricht (The Netherlands)

Two classes of anticoagulant agents are in current use today, the heparins and the oral anticoagulants. Within the heparin group, we distinguish unfractionated heparin (UFH) that accelerates the inactivation of both thrombin and factor Xa and low-molecular-weight heparins that lack the very high molecular weight fractions (>10,000) and contain significant amounts of a component that only fosters the inactivation of factor Xa. Oral anticoagulants inhibit the synthesis of functionally active carboxylated vitamin-K-dependent coagulation factors (factors II, VII, IX, X, proteins C and S). During the last decade, a third class of substances has been developed: direct specific thrombin inhibitors, such as hirudin, hirulog, argatoban and several others [1].

One of the key points in the clinical development of a new antithrombotic agent is to determine the optimal dosage. The proof of the pudding remains in the eating: i.e. the ultimate test are clinical dose-finding studies that relate the antithrombotic effect to the plasma levels of the anticoagulant. Such studies are extremely costly in time and money and fairly inaccurate in determining the range of concentrations that enables adequate treatment. It is therefore of utmost importance to dispose of a laboratory model that can serve to relate pharmacokinetics to the anticoagulant effect.

There were two fundamentally different types of such models. The animal thrombosis model, in which thrombosis is provoked in any of a number of standardized ways, and the relationship between drug level and thrombosis prevention can be determined, if required to a high degree of accuracy. The problem with this approach is that the data on artificial thrombosis in an animal should be used to guess the dose relationship in clinical thrombosis in man. This article is not meant to discuss this problem further.

The second type of model is an ex vivo test that can be carried out on the blood of a patient and that varies with the level of antithrombotic action obtained on the pathogenetic mechanism in the patient. An ideal test would be that one that directly reflects the effect of the drug on the relevant pathogenetic mechanism. Tests that only covariate with the antithrombotic action are still useful as long as the correlation with the pathogenetic mechanism is high and the tests meet general standards of reproducibility and accuracy. This raises the question of what indeed is the mechanism of action of the anticoagulant drugs. The fact that the three groups of drugs listed above each have an entirely different mode of action, facilitates an educated guess: it will not seem far fetched to surmise that the one thing that they have in common is the essential property that determines their antithrombotic action. That evidently is their capacity to diminish the amount of free thrombin that develops in clotting blood.

The next question, and the main one that we want to discuss in this review is: what are the relative merits of the currently used ex vivo tests for monitoring anticoagulant/antithrombotic treatment [2].

Thromboplastin Time or Prothrombin Time Test (PT)

The principle of the PT is that it measures the time necessary to clot a plasma sample after recalcification in the presence of an excess of tissue factor and procoagulant phospholipid, i.e. the clotting time in the extrinsic system, in which the factors I, II, V, VII and X are involved. Since the introduction of oral anticoagulants in the early 1940s, the PT has been the primary and exclusive means of monitoring the effectiveness and safety of oral anticoagulant therapy. Its results can be ex-

pressed in seconds, in the prolongation ratio (i.e. seconds in the sample/seconds in a normal control) or in percent activity, i.e. the clotting factor concentration in diluted normal plasma that will cause an identical clotting time. The results vary with the type of thromboplastin used, no matter what way of expression is used. Also, as a rule, calibration curves of diluted normal plasma are not parallel to those of diluted samples of anticoagulated blood and are dependent upon the dilution medium used. Hence, it is not possible to determine percent activity unequivocally. Therefore, the ratio method is commonly preferred. Variabilities in the results obtained with different thromboplastin reagents may cause oral anticoagulant therapy to be managed inappropriately [3]. In response to these problems, the World Health Organization (WHO) recommends a mathematical correction of the PT results known as the International Normalized Ratio (INR). The INR relies upon the calibration of commercial thromboplastins against a standard reagent with a known sensitivity to the antithrombotic effect of oral anticoagulants. By reporting PT as INR, results from laboratory to laboratory can be compared with acceptable accuracy. Today, this reporting technique is recommended for the monitoring of oral anticoagulant therapy. Extensive studies have been carried out to express the level of safe and efficient oral anticoagulant therapy in terms of INR [4, 5]. Nevertheless, a number of potential problems with the INR were identified. On the one hand, even the most extensive studies leave large margins of uncertainty as to the relationship between INR and bleeding or rethrombosis [6]. On the other hand, a lack of reliability was noticed at the onset of warfarin therapy. Also, the ISI provided by the manufacturer may be a source of error. On the other hand, a loss of accuracy was reported when thromboplastins with high INR values and/or

automatic clot detectors were used [7]. Finally, it should be noted that the PT is not sensitive to the effect of heparin. This in itself shows that it is a correlate to antithrombotic action only if this is obtained by oral anticoagulation, but cannot be the indicator of the pathophysiological mechanism behind it, otherwise it would have to react to heparin as well.

Activated Partial Thromboplastin Time (aPTT)

Principle of the Test

The aPTT represents the lag phase before explosive thrombin generation in an intrinsically triggered plasma sample, i.e. after recalcification in the presence of sufficient procoagulant phospholipid and a trigger of contact activation. The clotting factors involved are XII, XI, X, IX, VIII, V, II and I. The rate-limiting step in this process in all probability is the feedback activation of factor VIII by thrombin. This makes it an indicator of antithrombin activity, at least in plasma in which the participating clotting factors are normally present. Factor VIII cannot participate in the clotting cascade unless it is bound to phospholipid. Plasmatic, unactivated factor VIII does not bind to phospholipid because it is bound to von Willebrand factor. Its activation therefore can only be done by thrombin in solution. The presence of heparin decreases the lifetime of thrombin in solution and hence makes it more difficult for thrombin to activate factor VIII and thus retards the thrombin burst and prolongs the lag time [8, 9]. Factor V can adsorb to phospholipid and can be activated by membrane-bound meizothrombin. These membrane-bound processes are essentially immune to heparin action. This explains why thromboplastin-induced clotting times are insensitive to heparin whereas the aPTT is.

Performance

aPTT is an aspecific test, sensitive to all disturbances of the intrinsic clotting system. As a consequence of its mechanism of action, it is particularly sensitive to variations in the concentration and/or activity of factor VIII. It shows large interindividual variation and is strongly reagent and instrument dependent [10].

Interlaboratory variability of aPTT is excessive, as revealed by national and international surveys [11]. A collaborative study was aiming to standardize aPTT monitoring of UFH by use of a reference aPTT reagent. The study concluded that each laboratory should perform its own local aPTT system calibration. The study also shows that aPTT standardization is obviously not simple, if at all possible. It is strongly felt that a reliable overall test to monitor heparin therapy is strongly needed [12], but there is considerable doubt whether the approach that was successful with the PT in oral anticoagulant will serve as useful purpose in this case.

aPTT in the Control of Heparin Therapy

Despite these drawbacks, the aPTT is universally used as an indicator of the therapeutic level of heparin-mediated anticoagulation. As a rule of thumb, doubling the aPTT is thought to represent adequate heparin administration. This concept of a defined therapeutic range is based on experimental studies in animals [13] and on a limited subgroup analysis of two prospective cohort studies in man [14, 15]. The results of these studies have led to the recommendation that the therapeutic range of heparin should be an aM of 1.5 to 2.5 times the control, which is equivalent to a heparin level (measured by protamine titration of the thrombin time) of 0.1–0.4 IU/ml or, in terms of anti-factor Xa activity, of 0.3–0.7 U/ml. For the most commercial aPTT reagents in use at present, the lower limit of

the therapeutic range is over 1.5 and close to 2.0 times prolongation.

How does the aPTT perform as a monitor of heparin administration? In a recent study, significant differences in respect of heparin responsiveness were found between six aPTT reagents [16]. In addition, the limitation of aPTT for heparin monitoring was highlighted by the poor correlation between heparin concentration by protamine titration and aPTT ratio [16]. In some cases, high aPTT values are associated with subtherapeutic heparin levels [17]. This lack of association in these circumstances could be due to underlying diseases which affect the aPTT only.

It has been observed that a similar dosage of UFH produces significantly different prolongations in aPTT in different patients. The most likely explanation is the nonspecific binding of UFH to plasma proteins that neutralizes its anticoagulant effect. A number of these heparin-binding proteins are acute-phase reactants that tend to be elevated in sick patients. Another protein that neutralizes heparin is platelet factor 4, a platelet constituent that is released during platelet activation in vivo (thrombosis!) and in vitro (blood handling and storage). In patients with heparin resistance as assessed by the aPTT, the heparin level measured by the anti-Xa assay is a more relevant marker of the effectiveness of the treatment, since this test is relatively unaffected by the high levels of procoagulants that shorten the aPTT [16].

Recently, a formal review of the literature was performed on all available studies that provided information on the relationship between the risk of recurrent venous thromboembolism and the early aPTT response (within 24–48 h of initiation of treatment) to continuous intravenous heparin. More than 700 studies were identified, of which 50 were considered sufficiently informative for the overview. No convincing evidence shows that the

risk of recurrent venous thromboembolism is critically dependent on achieving a therapeutic aPTT result within 24–48 h [17].

aPTT as a General Indicator of Anticoagulant Therapy

The criterion of doubling the aPTT is sometimes, thoughtlessly, applied to other anticoagulants like direct thrombin inhibitors. This overlooks the fact that the mechanism by which such inhibitors prolong the aPTT is significantly different from that of heparin. Unlike heparin, such inhibitors do act on meizothrombin and therefore on the prothrombinase complex. This shows, even without going into details, that the dose-effect relationship must be quite different.

It is generally recognized that low-molecular-weight heparins with a high anti-factor Xa activity, and the synthetic pentasaccharides a fortiori, exert their antithrombotic action at doses that hardly prolong the aPTT. This in the first place indicates that the aPTT does not directly represent the antithrombotic effect, so that it is not allowed to assume that two times prolongation of the aPTT can be used with other anticoagulants than with UFH.

Thrombin Clotting Time (TCT)

TCT is based on the fact that a certain amount of thrombin is required over a certain time in order to clot a fibrin-containing plasma sample. The more thrombin is present, the shorter the time. Thrombin added to plasma will disappear with a velocity dependent upon the antithrombin activity, i.e. upon the presence of heparin. Clotting will be faster when thrombin persists longer, i.e. the less heparin there is. This test is very sensitive and relatively specific for the antithrombin effect of heparin. There are differences in heparin sen-

sitivity between various forms of TCT, that can be ascribed to variations in thrombin concentrations, thrombin type (bovine or human), and addition or not of calcium [18]. Very low concentrations of UFH can be detected when a low thrombin concentration and no calcium is used. Conversely, high concentration of thrombin and addition of calcium is necessary to assess high concentrations of UFH [19].

TCT is influenced by other factors than heparin, such as fibrin degradation products. Also variations in pH or temperature influence the results [20]. The reliability of TCT in monitoring of heparin treatment has been a matter of dissension. It has been advocated by some authors, while others have underlined cautiousness and doubt [21, 22]. One point seems clear: if only one single thrombin concentration is used, only a narrow range of heparin concentrations can be accurately assessed. This range is somewhat enlarged by the addition of calcium [23].

Activated Clotting Time

The activated clotting time is a whole blood clotting time in the presence of a contact activator such as celite [24]. It is a rapid and simple test that can be carried out at the bedside and in the operating theater. Indeed its widest use, in the USA, is to monitor heparin therapy during extracorporeal circulation and coronary angioplasty [25]. The test is essentially an aPTT carried out on whole blood. Like the aPTT it covers an acceptable range of heparin levels. It carries the same drawbacks as the aPTT and is also strongly influenced by the conditions of blood sampling [26].

Ecarin Clotting Time

Ecarin is a snake venom enzyme derived from the viper *Echis carinatus*, which converts prothrombin into meizothrombin and other intermediates. The thrombin intermediates exhibit thrombin-like activity on the fibrinogen in the sample. While direct thrombin inhibitors are able to inhibit these thrombin intermediates, antithrombin has no effect and consequently heparin not either. Recently, ecarin clotting time has been proposed for the specific measurement of the effect of hirudin and related direct antithrombin agents [27]. This method is reported to show linear correlation with the anti-IIa activity of hirudin, to be highly reproducible, simple to perform and reliable for bedside monitoring of hirudin [28, 29]. Critical reviews of its performance in practice are still lacking.

Heparin Anti-IIa and Anti-Xa Activities

The specific action of heparin is that it accelerates the inactivation of thrombin and factor Xa in plasma by antithrombin. This property can be used to estimate these specific activities of heparin in a sample. These methods are based on inhibition of a known amount of factor Xa or thrombin, and are usually based on measuring residual factor Xa (thrombin) activity after a fixed incubation time, using a photometric determination with a synthetic peptide substrate to determine residual activities. The use of a photometer reduces the observer bias. These tests are as sensitive as and more specific than the aPTT [30]. Assay of anti-Xa activity is the most widely used method for assessing low-molecular-weight heparin *ex vivo*. The results are expressed in anti-Xa international units per milliliter of plasma. Here an important artifact must be noted. In plasma, in the presence

of Ca^{2+} ions, the anti-factor Xa activity of UFH, and hence of the international heparin standard, is twice higher than in an anti-Xa test situation where Ca^{2+} is absent. This is not the case, however, for low-molecular-weight heparins. Hence, the latter are measured against a 'handicapped' standard and overestimated. The effect causes 1.5–2.0 times too high anti-factor Xa values [31]. It should be noted that the use of the international standard for low-molecular-weight heparins does not abolish this error because the activity of that standard itself has been determined against the UFH standard. Amidolytic methods may differ as to the reagents used. The addition of purified antithrombin to the test plasma is recommended by some authors and rejected by others [32]. A well-designed test renders, however, always the catalytic activity of the heparin present in the sample, independent of other components that may be present.

Endogenous Thrombin Potential (ETP)

The thrombin-generating capacity of plasma is one of the main determinants of hemostasis and thrombosis [33]. Clotting times measure the onset of thrombin generation, the moment that about 1 unit of thrombin (~ 10 nM) is formed. Most thrombin, up to 100–250 nM, appears after the clot is formed. *In vivo* this thrombin will still act on its numerous substrates and exert a prothrombotic action. It has been shown that the enzymatic potency of generated thrombin is best expressed as the area under the thrombin generation curve, or ETP. The ETP can be determined from a classical thrombin generation curve, but this is unsuitable for routine purposes. A simplified and automated test has been described [34]. This parameter might possibly be representing the essential variable

decreasing in drug-induced hypocoagulability and increasing in hypercoagulability [34].

Preliminary results were obtained with this ETP assay as a routine procedure. The ETP appears to be decreased to between 20 and 40% of normal by oral anticoagulation in the therapeutic range (INR = 1.5–3.0) and by heparin administration in the therapeutic range (aPTT = 1.5–2.5 control). It also indicates the hypocoagulability obtained in mixed treatment, common in the 1st week of treatment of venous thrombosis, and in which none of the currently used tests gives reliable guidelines. The ETP is increased in untreated subjects with congenital antithrombin deficiency and in women using contraceptives. It is also increased in patients with deep vein thrombosis and in coronary infarction after the acute phase [28]. Extensive large-scale clinical trials will still be necessary to demonstrate the feasibility and the ability of the ETP as a general indicator of anticoagulation or hypercoagulable states.

Conclusion

Laboratory monitoring of anticoagulant therapy has its roots in the past, and is more often than not founded on tradition rather than on well-established data. Secondary rigorous standardization may (as in the case of PT) or may not (as in the case of aPTT) ameliorate the situation. In any case, the need of an adequate assay to monitor anticoagulant/antithrombotic therapy is obvious. This test should be easy and cheap and should represent an essential parameter of the coagulation system and therefore be applicable to all anticoagulants in the same way. No such test is available at this moment, but the ETP proposes as a candidate worthy of clinical testing.

References

- 1 Lefkovits J, Topol EJ: Direct thrombin inhibitors in cardiovascular medicine. *Circulation* 1994;90:1522–1536.
- 2 Fitzgerald GA: The human pharmacology of thrombin inhibition. *Coron Artery Dis* 1996;7:911–918.
- 3 van-Bergen PF, Deckers JW, Jonker JJ, van Domburg RT, Azar AJ, Hofman A: Efficacy of long-term anticoagulant treatment in subgroups of patients after myocardial infarction. *Br Heart J* 1995;74:117–121.
- 4 Azar AJ, Cannegieter SC, Deckers JW, Briet E, van Bergen PF, Jonker JJ, Rosendaal FR: Optimal intensity of oral anticoagulant therapy after myocardial infarction. *J Am Coll Cardiol* 1996;27:1349–1355.
- 5 Azar AJ, Deckers JW, Rosendaal FR, van Bergen PF, van der Meer FJ, Jonker JJ, Briet E: Assessment of therapeutic quality control in a long-term anticoagulant trial in post-myocardial infarction patients. *Thromb Haemost* 1994;72:347–351.
- 6 Azar AJ, Koudstaal PJ, Wintzen AR, van Bergen PF, Jonker JJ, Deckers JW: Risk of stroke during long-term anticoagulant therapy in patients after myocardial infarction. *Ann Neurol* 1996;39:301–307.
- 7 Hirsh J, Dalen JE, Deykin D, Poller L, Bussey H: Oral anticoagulants. Mechanism of action, clinical effectiveness, and optimal therapeutic range. *Chest* 1995;108(suppl):231S–246S.
- 8 Béguin S, Lindhout T, Hemker HC, et al: The mode of action of heparin in plasma. *Thromb Haemost* 1988;60:457–462.
- 9 Béguin S, Dol F, Hemker HC: Factor IXa inhibition contributes to the heparin effect. *Thromb Haemost* 1991;66:306–309.
- 10 Kitchen S, Preston FE: The therapeutic range for heparin therapy: Relationship between six activated partial thromboplastin time reagents and two heparin assays. *Thromb Haemost* 1996;75:734–739.
- 11 D'Angelo A, Seveso MP, D'Angelo SV, Gilardoni F, Dettori AG, Bonini P: Effect of clot-detection methods and reagents on activated partial thromboplastin time (APTT). *Am J Clin Pathol* 1990;94:297–306.

- 12 Van der Velde EA, Poller L: The APTT monitoring of heparin - The ISTH/ICSH Collaborative Study. *Thromb Haemost* 1995;73:73-81.
- 13 Chiu HM, Hirsh J, Yung WL, Regoeczi E, Gent M: Relationship between the anticoagulant and antithrombotic effects of heparin in experimental venous thrombosis. *Blood* 1977;49:171-184.
- 14 Basu D, Gallus A, Hirsh J, Cade JF: A prospective study of the value of monitoring heparin treatment with the activated partial thromboplastin time. *N Engl J Med* 1972;287:324-327.
- 15 Hull RD, Raskob GE, Hirsh J: Continuous intravenous heparin compared with intermittent subcutaneous heparin in the initial treatment of proximal vein thrombosis. *N Engl J Med* 1986;315:1109-1114.
- 16 Levine MN, Hirsh J, Gent M, Turpie AG, Cruickshank M, Weitz J, Anderson D, Johnson M: A randomized trial comparing activated thromboplastin time with heparin assay in patients with acute venous thromboembolism requiring large daily doses of heparin. *Arch Intern Med* 1994;154:49-56.
- 17 Anand S, Ginsberg JS, Kearon C, Gent M, Hirsh J: The relation between the activated partial thromboplastin time response and recurrence in patients with venous thrombosis treated with continuous intravenous heparin. *Arch Intern Med* 1996;156:1677-1681.
- 18 Teien AN, Lie M: Heparin assay in plasma. A comparison of five clotting methods. *Throm Res* 1975, pp 777-788.
- 19 Eika C, GoM HC, Kierulf P: Detection of small amounts of heparin by the thrombin clotting time. *Lancet* 1972;ii:376.
- 20 Godal HC: The assay of heparin in thrombin systems. *Scand J Clin Lab Invest* 1961;13:153-166.
- 21 Soloway HB, Cox SP: In vitro comparison of the thrombin time and activated partial thromboplastin time in the laboratory control of heparin therapy. *Am J Clin Pathol* 1973;60:648-650.
- 22 Ts'ao C, Raymond J, Kolb T, Lo R: Effects of source and concentration of thrombin, and divalent cations, on thrombin time of heparinized plasma. *Am J Clin Pathol* 1976;65:206-212.
- 23 Denson KW, Bonnar J: The measurement of heparin. A method based on the potentiation of anti-factor Xa. *Thromb Diath Haemorrh* 1973;30:471-479.
- 24 Hattersley PG: Progress Report: The activated coagulation time of whole blood (ACT). *Am J Clin Pathol* 1976;66:899-904.
- 25 Blakely JA: A rapid bedside method for the control of heparin therapy. *Can Med Assoc J* 1968;99:1072-1076.
- 26 Forman WB, Bayer G: A simplified method for monitoring heparin therapy at the bedside: the activated whole blood clotting time. *Am J Hematol* 1981;11:277-281.
- 27 Nowak G, Bucha E: A new method for the therapeutical monitoring of hirudin. *Thromb Haemost* 1993;69:1306-1314.
- 28 Esslinger HU, Dubbers K, Radziwon P, Breddin HK: Monitoring of the anticoagulant effects of PEG-hirudin administered as a continuous 24-hour infusion following an I.V. bolus injection in healthy volunteers using the ecarin clotting time. *Thromb Haemost* 1997; (suppl):277-278.
- 29 Bode C, Kohler B, Steg G, Parow C, Rubsamen K: The ecarin clotting time but not the aPTT is a reliable indicator for PEG-hirudin blood levels in patients with unstable angina pectoris. *J Am Coll Cardiol* 1997; 29:401A-411A.
- 30 Teien AN, Lie M, Abildgaard U: Assay of heparin in plasma using a chromogenic substrate for activated factor X. *Thromb Res* 1976;8:413-416.
- 31 Hemker HC, Béguin S: The activity of heparin in the presence and absence of Ca^{++} ions; why the anti-Xa activity of LMW heparins is about two times overestimated. *Thromb Haemost* 1993;70:717-718.
- 32 Walenga JM, Bara L, Samama M, Fareed J: Amidolytic antifactor Xa assays in the laboratory evaluation of heparin and low molecular weight fractions. *Semin Thromb Haemost* 1985;11:100-107.
- 33 Hemker HC, Béguin S: Thrombin generation in plasma: Its assessment via the endogenous thrombin potential. *Thromb Haemost* 1995;74:134-138.
- 34 Wielders S, Mukherjee M, Michiels J, Rijkers DTS, Cambus JP, Knebel RWC, Kakkar VV, Hemker HC, Béguin S: The routine determination of the endogenous thrombin potential. First results in different forms of hyper- and hypocoagulability. *Thromb Haemost* 1997;77:629-636.