Current perspective: Osimertinib-induced QT prolongation

Citation for published version (APA):

Schiefer, M., Hendriks, L. E. L., Dinh, D., Lalji, U., & Dingemans, A-M. C. (2018). Current perspective: Osimertinib-induced QT prolongation: new drugs with new side-effects need careful patient monitoring. *European Journal of Cancer*, 91, 92-98. <https://doi.org/10.1016/j.ejca.2017.12.011>

Document status and date:

Published: 01/03/2018

DOI:

[10.1016/j.ejca.2017.12.011](https://doi.org/10.1016/j.ejca.2017.12.011)

Document Version:

Publisher's PDF, also known as Version of record

Document license:

Taverne

Please check the document version of this publication:

- A submitted manuscript is the version of the article upon submission and before peer-review. There can be important differences between the submitted version and the official published version of record. People interested in the research are advised to contact the author for the final version of the publication, or visit the DOI to the publisher's website.
- The final author version and the galley proof are versions of the publication after peer review.
- The final published version features the final layout of the paper including the volume, issue and page numbers.

[Link to publication](#)

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal.

If the publication is distributed under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license above, please follow below link for the End User Agreement:

www.umlib.nl/taverne-license

Take down policy

If you believe that this document breaches copyright please contact us at:

repository@maastrichtuniversity.nl

providing details and we will investigate your claim.

Available online at www.sciencedirect.com

ScienceDirect

journal homepage: www.ejancer.com

Current Perspective

Current perspective: Osimertinib-induced QT prolongation: new drugs with new side-effects need careful patient monitoring

Mart Schiefer^a, Lizza E.L. Hendriks^{b,*}, Trang Dinh^c, Ulrich Lalji^d,
Anne-Marie C. Dingemans^b

^a Dept. of Pulmonary Diseases, Zuyderland Medical Center Location Heerlen, PO Box 5500, 6130 BM, Sittard-Geleen, The Netherlands

^b Dept. of Pulmonary Diseases, GROW – School for Oncology and Developmental Biology, Maastricht University Medical Center+, PO Box 5800, 6202 AZ, Maastricht, The Netherlands

^c Dept. of Cardiology, Maastricht University Medical Center+, PO Box 5800, 6202 AZ, Maastricht, The Netherlands

^d Dept. of Radiology, Maastricht University Medical Center+, PO Box 5800, 6202 AZ, Maastricht, The Netherlands

Received 24 November 2017; accepted 6 December 2017

KEYWORDS

Non-small cell lung cancer;
Tyrosine kinase inhibitor;
QT interval;
Disease flare;
Drug monitoring

Abstract An increasing number of tyrosine kinase inhibitors (TKIs) are available for the treatment of non-small cell lung cancer (NSCLC). QT prolongation is one of the known, but relatively rare, adverse events of several TKIs (e.g. osimertinib, crizotinib, ceritinib). Screening for QT prolongation in (high risk) patients is advised for these TKIs. When a QT prolongation develops, the physician is challenged with the question whether to (permanently) discontinue the TKI. In this perspective, we report on a patient who developed a grade III QT prolongation during osimertinib (a third-generation epidermal growth factor receptor [EGFR]-TKI) treatment. On discontinuation of osimertinib, she developed a symptomatic disease flare, not responding to subsequent systemic treatment. The main aim of this perspective is to describe the management of QT prolongation in stage IV *EGFR* driver mutation NSCLC patients. We also discuss the ethical question of how to weigh the risk of a disease flare due to therapy cessation against the risk of sudden cardiac death. A family history of sudden death and a prolonged QT interval might indicate a familiar long QT syndrome. We have summarised the current monitoring advice for TKIs used in the treatment of lung cancer and the most common drug–TKI interactions to consider and to optimise TKI treatment in lung cancer patients.

© 2017 Elsevier Ltd. All rights reserved.

* Corresponding author. Fax: +31 (0) 433875051.

E-mail addresses: m.schiefer@zuyderland.nl (M. Schiefer), lizza.hendriks@mumc.nl (L.E.L. Hendriks), trang.dinh@mumc.nl (T. Dinh), u.lalji@mumc.nl (U. Lalji), a.dingemans@mumc.nl (A.-M.C. Dingemans).

1. Introduction

More and more targetable molecular drivers of non-small cell lung cancer (NSCLC) are being discovered [1]. The targeted agents for these driver mutations are often tyrosine kinase inhibitors (TKIs). The European Medicines Agency (EMA)- and US Food and Drug Administration (FDA)-approved TKIs for activating epidermal growth factor receptor (*EGFR*) mutations include erlotinib, gefitinib, afatinib and osimertinib. Examples of TKIs approved for anaplastic lymphoma

kinase rearrangements are crizotinib, ceritinib and alectinib and for *ROS1* translocations, crizotinib is approved [1]. *EGFR* mutations are found in approximately 10–15% of Caucasian patients with stage IV non-squamous NSCLC [2], and this percentage increases to more than 60% in non-smoking Asian females [3]. An activating *EGFR* mutation is predictive for response to EGFR-TKI [4]. For example, osimertinib is a third-generation EGFR-TKI and EMA approved for *EGFR*-mutated patients with a T790M mutation. Recently, results of the first line randomised phase III

Fig. 1. Primary tumour in the left lower lobe before, during and after osimertinib therapy. Consecutive CT scans demonstrating the primary tumour in the left lower lobe, adjacent to a pleural thickening and calcification caused by previous empyema. (A) Before osimertinib therapy. (B) Day 32 of osimertinib therapy, partial response. (C) Day 183, stable situation. (D) Day 241, slow progression. (E) Day 301, slow progression. (F) Day 350, rapid progression after cessation of osimertinib. CT, computed tomography.

FLAURA trial were reported, and it was shown that osimertinib had a superior progression free survival (PFS) compared with gefitinib [5]. For osimertinib, most common adverse events (almost all grade I or II) are diarrhoea (41%), rash (34%), dry skin (23%) and paronychia (22%). Although not frequent, QT prolongation is one of the known adverse events, and this occurs in approximately 4% of patients (almost all patients grade I or II) [6]. For several other TKIs, QT prolongation has been described [7–10]. For a grade III QT prolongation, it is advised to temporarily discontinue the causative agent as there is an increased risk of sudden cardiac death due to ventricular tachycardia and torsade de pointes [11]. However, discontinuation of a TKI can result in a rapid symptomatic disease flare [12]. In the case reported below, we discuss that a disease flare due to therapy cessation should be weighed against the risk of sudden cardiac death. Furthermore, we provide recommendations for QT management in a patient treated with a TKI. In a broader perspective, we summarise the current monitoring advice for TKIs (almost) approved for the treatment of lung cancer and the most common drug–TKI interactions to consider and to optimise TKI treatment of lung cancer patients.

1.1. Case report

A 62-year-old female known with stage IV pulmonary adenocarcinoma with an *EGFR* exon 19 deletion was treated with fourth line osimertinib 80 mg once daily. This was well tolerated and resulted in a rapid clinical and radiological response. After 9 months, a very slow asymptomatic progression was observed (Fig. 1); osimertinib was continued because of persisting clinical benefit. Regular (2 weekly) electrocardiograms (ECGs)

were performed for the first 2 months, thereafter monthly. These ECGs showed no abnormalities. However, after 11 months, there was a sudden grade III QT prolongation (560 ms) (Fig. 2). Electrolytes were normal; no other QT-prolonging drugs were used. Osimertinib was discontinued, and on advice of the cardiologist, a detailed family history was obtained. A brother and uncle had sudden cardiac deaths. Because of the family history, the patient was referred for genetic counselling for a workup of familial QT prolongation. Genetic counselling showed no specific mutations related to the long QT syndrome (no pathogenic mutations in *KCNQ1*, *KCNH2*, *SCN5A*, *KCNE1*, *KCNE2*, *KCNJ8*, *CAV3*, *KCNJ2* and exon 8 of the *CACNA1C* gene) [13,14]. Within 5 days, the QT interval normalised, and osimertinib was not reintroduced because of the already occurring slow progressive disease. A new tumour biopsy was performed directly after discontinuing osimertinib. Within 2 weeks of discontinuing osimertinib, while awaiting biopsy results, a rapid symptomatic disease progression occurred, not responding to fifth line pemetrexed monotherapy. Biopsy results became available during pemetrexed treatment but revealed no druggable target and no T790M. The patient died from symptomatic brain metastasis 2 months after discontinuation of osimertinib.

1.2. Discussion: monitoring of TKI treatment

For several TKIs, grade III or higher QT prolongation has been described (e.g. osimertinib: 1%, dacomitinib: 1%, crizotinib: 3%, ceritinib: 3–6%) [7–10]. Animal models suggest that this is caused by inhibition of PI3K signalling [15]. QT prolongation has been reported in 4% of patients receiving osimertinib and was classified a

Fig. 2. QT interval during osimertinib therapy. (A) Graphic presentation of the development of corrected QT time during and after osimertinib therapy, which was discontinued at day 302. (B) Electrocardiogram (ECG) at the start of osimertinib therapy, demonstrating a QTc of 444 ms. (C) ECG at day 302 of osimertinib therapy, demonstrating a QTc of 560 ms.

Table 1
Summary of advised monitoring per TKI.

Monitoring	Renal function & electrolytes ^e	Liver function ^e	Fasting plasma glucose ^e	Creatine kinase, ^d myalgia symptoms	Pancreatic enzymes	Lipid profile ^a	Blood count and differentiation	Close INR control in coumarin therapy	Pulse and blood pressure	ECG (in patients at risk)	Echocardiography (in patients at risk)	ILD symptoms	Ocular symptoms	Bullous skin conditions	Heart failure symptoms
Gefitinib (Iressa) [20]	x	x						x				x	x		
Erlotinib (Tarceva) [21]	x	x						x				x	x	x	
Aflatinib (Giotrif) [22]	x	x										x	x	x	x ^b
Dacomitinib ^a	x	x					x			x	x	x	x		
(PF-00299804) [23]															
Osimertinib (Tagrisso) [11]	x	x								(x)	x	x	x		x
Crizotinib (Xalkori) [16]	x	x					x		x	x	x	x	x		x
Ceritinib (Zykadia) [17]	x	x	x		x ^c				x	x	x	x	x		x
Alectinib (Alecensa) [24]	x	x		x					x			x			
Brigatinib (Alunbrig) [25]	x	x	x	x	x ^c				x			x	x		
Lorlatinib ^a (PF-6463922) [26]	x	x			x ^d		x	x		x		x			

INR, international normalized ratio; ECG, electrocardiogram; ILD, interstitial lung disease; TKI, tyrosine kinase inhibitor; FDA, US Food and Drug Administration; EMA, European Medicines Agency.

^a Not yet FDA or EMA approved.

^b Preclinical toxicology studies did not show cardiotoxicity.

^c At baseline, monthly and as clinically indicated in patients reporting symptoms.

^d At baseline, every two weeks during the first month of treatment and as clinically indicated in patients reporting symptoms.

^e At baseline and as clinically indicated in patients reporting symptoms.

grade III adverse event in 1%. Although the QT prolongation is rare, for osimertinib, it is advised to regularly perform an ECG in patients at risk for QT prolongation [11]. For crizotinib and ceritinib, it is advised to perform baseline ECGs in all patients and regular ECGs at least in patients at risk for QT prolongation [16,17]. As the TKIs used in the treatment of driver mutated NSCLC differ in their adverse event profile, monitoring of patients should be tailored according to the prescribed TKI. Furthermore, interaction with other drugs can cause an increase or decrease in the plasma concentration of the TKI, resulting in a higher chance of adverse events or in possible undertreatment of the patient. Monitoring advice is summarised in Table 1, and possible drug–TKI interactions with the resulting effects on the TKI are presented in Table 2.

A prolonged QT interval can lead to ventricular tachycardia and torsade de pointes with potentially fatal outcome. It can present as a congenital 'long QT syndrome' and can be provoked by conditions such as bradycardia or hypokalemia or exposure to certain drugs. In general, a QT prolongation beyond 500 ms should prompt cessation of QT-prolonging drugs to prevent sudden cardiac death as the relative risk of a cardiac death is more than 2.5 times that of a person without QT prolongation [14,18]. Predisposing factors for QT prolongation and their risk management are summarised in Table 3.

In our case, the risk of fatal arrhythmia due to QT prolongation occurred in the context of a stage IV *EGFR*-mutated pulmonary adenocarcinoma that was treated with an *EGFR*-TKI. Disease flare after TKI discontinuation has been described and occurred after a median of 8 days [12]. This unfavourable prognostic consequence of therapy cessation should be weighed against the risk of sudden cardiac death. An ethical question is whether a risk of iatrogenic death is acceptable in palliative oncologic therapy. In case of therapy continuation, preventive measures to limit the risk of cardiac arrhythmias consist of early patient reporting of palpitations and (near) syncope, early recognition and treatment of hypokalemia, beta-blocker therapy and avoidance of concomitant QT-prolonging or interacting drugs (Table 3) [13,14]. Furthermore, attention should be paid to the family history as a prolonged QT interval might indicate a familial long QT syndrome [13,14]. When there is an indication for a familial long-QT syndrome, first degree family members with a (suspicion of) a long-QT syndrome are eligible for a periodical cardiology evaluation [19]. In general, one should be aware of the monitoring advice for TKI use and possible drug–TKI interactions to optimise the TKI treatment of lung cancer patients.

At this point, we do not know if dose reduction is associated with a decreased risk of QT prolongation or if QT prolongation would recur or not with the introduction of a lower dose. However, as our patient already

Table 2

Summary of important drug interactions per TKI.

Type of drug	CYP3A4 inhibitors ^b	CYP3A4 inducers ^c	CYP1A2 inhibitors ^d	P-gp inhibitors ^e	P-gp inducers ^f	P-gp–dependent drugs ^g	BCRP–dependent drug ^h	CYP3A–dependent drugs ⁱ	CYP2C9–dependent drugs ^j	CYP2B6–dependent drugs ^k	CYP2D6–dependent drugs ^l	QT–prolonging drugs ^m	Gastrointestinal toxic drugs	Hepatotoxic drugs	Gastric pH–modifying agents	Coumarins (INR increase)	Statins (myopathy risk)	Oral contraception unsafe
Gefitinib (Iressa) [20]	x	x	x								x				x	x		
Erlotinib (Tarceva) [21]	x	x	x	x									x			x	x	
Afatinib (Giotrif) [22]				x	x		x											
Dacomitinib ^a (PF-00299804) [23]						x					x	x		x				
Osimertinib (Tagrisso) [11]		x					x					x						x
Crizotinib (Xalkori) [16]	x	x				x		x		x		x	x	x				x
Ceritinib (Zykadia) [17]	x	x		x	x	x	x					x			x			
Alectinib (Alecensa) [24]						x	x											x
Brigatinib (Alunbrig) [25]	x	x						x										
Lorlatinib ^a (PF-6463922) [26]	x	x				x		X ⁿ	x	x						x		

FDA, US Food and Drug Administration; TKI, tyrosine kinase inhibitor; EMA, European Medicines Agency.

^a Not yet FDA or EMA approved.^b CYP3A4-inhibiting drugs increase TKI concentration, i.e. grapefruit juice, azoles, erythromycin, clarithromycin, boceprevir and HIV protease inhibitors.^c CYP3A-inducing drugs decrease TKI concentration, i.e. phenytoin, rifampicin, carbamazepine, barbiturates and St. John's wort.^d CYP1A2 inhibitors increase TKI concentration, i.e. ciprofloxacin and fluvoxamine.^e P-glycoprotein inhibitors increase TKI concentration, i.e. oritonavir, cyclosporine A, ketoconazole, itraconazole, erythromycin, verapamil, quinidine, tacrolimus, nelfinavir, saquinavir and amiodarone.^f P-glycoprotein inducers decrease TKI concentration, i.e. rifampicin, carbamazepine, phenytoin, phenobarbital and St. John's wort.^g P-glycoprotein–dependent drugs increase in plasma concentration, i.e. digoxin, colchicine, dabigatran, paclitaxel, pravastatin and vincristine.^h Breast cancer resistance protein (BCRP)–dependent drugs increase in plasma concentration, i.e. rosuvastatin, methotrexate, sulfasalazine and topotecan.ⁱ CYP3A-dependent drugs increase in plasma concentration, i.e. astemizole, cisapride, ciclosporin, ergotamine, fentanyl, pimozone, quinidine, tacrolimus, alfentanil and sirolimus.^j CYP2C9-dependent drugs increase in plasma concentration, i.e. phenytoin and warfarin.^k CYP2B6-dependent drugs increase in plasma concentration, i.e. bupropion and efavirenz.^l CYP2D6-dependent drugs increase in plasma concentration, i.e. metoprolol.^m QT-prolonging drugs, i.e. amiodarone, kinidine, disopyramide, sotalol, domperidone, methadone, tricyclic antidepressants, antipsychotics, macrolide antibiotics, fluorochinolones, antimycotics, granisetron and ondansetron.ⁿ CYP3A-dependent drugs decrease in plasma concentration.

Table 3
Risk factors and preventive measures regarding drug-induced QT prolongation.

Predisposing factors for QT prolongation	Risk management
Electrolyte disturbances (hypoK, –Mg, –Ca)	Laboratory monitoring (in vomiting, diarrhoea) Caution using provoking drugs (i.e. diuretics) Pharmacologic surveillance; drug avoidance
Provoking or interacting drugs	Regular ECGs; screening for gene polymorphisms
Heart failure, left ventricular hypertrophy, extreme bradycardia	Cardiologic screening and optimisation ECG monitoring
Strenuous physical activity, hypothermia	Avoidance, patient education
Hypothyroidism	Laboratory screening and suppletion
Female sex, increasing age	Awareness
Family history LQT syndrome, sudden death	Awareness Genetic counselling

hypoK, hypokaliemia; Mg, magnesium; Ca, calcium; ECG, electrocardiogram; LQT, long QT.

had progressive disease on full dose osimertinib, we did not think that a lower dose would give us disease control or a reassurance about the effect on the QT interval.

2. Conclusion

New drugs have new side-effects that require more general medical input, often in the course of a busy oncology clinic. We think that a prolonged QT interval due to palliative cancer therapy should not automatically result in therapy cessation, but should be weighed against the risk of disease flare. Genetic testing should be considered especially with a family history of sudden death.

We have developed some tables that should help busy clinicians to make decisions.

Conflict of interest statement

None declared.

Funding

This research did not receive any specific grant from funding agencies in the public, commercial or not-for-profit sectors.

Financial disclosures

Lizza E.L. Hendriks received personal fees from Roche, MSD, AstraZeneca and BMS, outside the submitted work. Anne-Marie C. Dingemans received personal fees from AstraZeneca, Roche and Pfizer, outside the submitted work. The remaining authors have no financial disclosures to declare.

References

- [1] Hirsch FR, Suda K, Wiens J, Bunn jr PA. New and emerging targeted treatments in advanced non-small-cell lung cancer. *Lancet* 2016;388(10048):1012–24.
- [2] Rosell R, Carcereny E, Gervais R, Vergnenegre A, Massuti B, Felip E, et al. Erlotinib versus standard chemotherapy as first-line treatment for European patients with advanced EGFR mutation-positive non-small-cell lung cancer (EORTC): a multicentre, open-label, randomised phase 3 trial. *Lancet Oncol* 2012;13(3):239–46.
- [3] Shi Y, Au JS, Thongprasert S, Srinivasan S, Tsai CM, Khoa MT, et al. A prospective, molecular epidemiology study of EGFR mutations in Asian patients with advanced non-small-cell lung cancer of adenocarcinoma histology (PIONEER). *J Thorac Oncol* 2014;9(2):154–62.
- [4] Mok TS, Wu YL, Thongprasert S, Yang CH, Chu DT, Saijo N, et al. Gefitinib or carboplatin-paclitaxel in pulmonary adenocarcinoma. *N Engl J Med* 2009;361(10):947–57.
- [5] Soria JC, Ohe Y, Vansteenkiste J, Reungwetwattana T, Chewaskulyong B, Lee KH, et al. Osimertinib in Untreated EGFR-Mutated Advanced Non-Small-Cell Lung Cancer. *N Engl J Med* 2018;378:113–25. <https://doi.org/10.1056/NEJMoa1713137> [Epub ahead of print].
- [6] Mok TS, Wu Y-L, Ahn M-J, Garassino MC, Kim HR, Ramalingam SS, et al. Osimertinib or Platinum-Pemetrexed in EGFR T790M-positive lung cancer. *N Engl J Med* 2017;376(7):629–40.
- [7] Sullivan I, Planchard D. Osimertinib in the treatment of patients with epidermal growth factor receptor T790M mutation-positive metastatic non-small cell lung cancer: clinical trial evidence and experience. *Ther Adv Respir Dis* 2016;10(6):549–65.
- [8] Abdul Razak AR, Soulieres D, Laurie SA, Hotte SJ, Singh S, Winquist E, et al. A phase II trial of dacomitinib, an oral pan-human EGF receptor (HER) inhibitor, as first-line treatment in recurrent and/or metastatic squamous-cell carcinoma of the head and neck. *Ann Oncol* 2013;24(3):761–9.
- [9] Kazandjian D, Blumenthal GM, Chen HY, He K, Patel M, Justice R, et al. FDA approval summary: crizotinib for the treatment of metastatic non-small cell lung cancer with anaplastic lymphoma kinase rearrangements. *Oncologist* 2014;19(10):e5–11.
- [10] Raedler LA. Zykadia (ceritinib) approved for patients with crizotinib-resistant ALK-positive non-small-cell lung cancer. *Am Health Drug Benefits* 2015;8(Spec Feature):163–6.
- [11] EMA. Summary of the risk management plan (RMP) for Tagrisso (osimertinib). 03-Oct-2017; Available from: http://www.ema.europa.eu/docs/en_GB/document_library/EPAR_-_Product_Information/human/004124/WC500202022.pdf.
- [12] Chaft JE, Oxnard GR, Sima CS, Kris MG, Miller VA, Riely GJ. Disease flare after tyrosine kinase inhibitor discontinuation in patients with EGFR-mutant lung cancer and acquired resistance to erlotinib or gefitinib: implications for clinical trial design. *Clin Cancer Res* 2011;17(19):6298–303.
- [13] Priori SG, Barhanin J, Hauer RN, Haverkamp W, Jongsma HJ, Kleber AG, et al. Genetic and molecular basis of cardiac

- arrhythmias: impact on clinical management parts I and II. *Circulation* 1999;99(4):518–28.
- [14] Roden DM. Drug-induced prolongation of the QT interval. *N Engl J Med* 2004;350(10):1013–22.
- [15] Lu Z, Wu CY, Jiang YP, Ballou LM, Clausen C, Cohen IS, et al. Suppression of phosphoinositide 3-kinase signaling and alteration of multiple ion currents in drug-induced long QT syndrome. *Sci Transl Med* 2012;4(131). 131ra50.
- [16] EMA. EPAR crizotinib. 10-Oct-2017; Available from: http://www.ema.europa.eu/docs/en_GB/document_library/EPAR_-_Product_Information/human/002489/WC500134759.pdf.
- [17] EMA. Summary of the risk management plan (RMP) for Zykadia (ceritinib). 06-Oct-2017; Available from: http://www.ema.europa.eu/docs/en_GB/document_library/EPAR_-_Risk-management-plan_summary/human/003819/WC500183128.pdf.
- [18] Zhang Y, Post WS, Dalal D, Blasco-Colmenares E, Tomaselli GF, Guallar E. QT-interval duration and mortality rate: results from the third National Health and Nutrition Examination Survey. *Arch Intern Med* 2011;171(19):1727–33.
- [19] Skinner, J. et al. Guidelines for the diagnosis and management of Familial Long QT Syndrome. 2011 01-nov-2017; Available from: http://www.csanz.edu.au/documents/guidelines/clinical_practice/Familial_Long_QT_Syndrome.pdf.
- [20] EMA. EPAR gefitinib. access date 01-nov-2017; Available from: http://www.ema.europa.eu/docs/en_GB/document_library/EPAR_-_Product_Information/human/001016/WC500036358.pdf.
- [21] EMA. EPAR erlotinib. access date 01-nov-2017; Available from: http://www.ema.europa.eu/docs/en_GB/document_library/EPAR_-_Product_Information/human/000618/WC500033994.pdf.
- [22] EMA. EPAR afatinib. access date 01-nov-2017; Available from: http://www.ema.europa.eu/docs/en_GB/document_library/EPAR_-_Product_Information/human/002280/WC500152392.pdf.
- [23] Wu YL, Cheng Y, Zhou X, Lee KH, Nakagawa K, Niho S, et al. Dacomitinib versus gefitinib as first-line treatment for patients with EGFR-mutation-positive non-small-cell lung cancer (ARCHER 1050): a randomised, open-label, phase 3 trial. *Lancet Oncol* 2017 Nov;18(11):1454–66.
- [24] EMA. EPAR alectinib. 01-nov-2017; Available from: http://www.ema.europa.eu/docs/en_GB/document_library/EPAR_-_Product_Information/human/004164/WC500225707.pdf.
- [25] FDA. FDA prescribing information brigatinib. 01-nov-2017; Available from: https://www.accessdata.fda.gov/drugsatfda_docs/label/2017/208772lbl.pdf.
- [26] Pfizer. PF-06463922 named-patient early access reference guide for physicians, Version 2.0, 05 August 2016. 2016.