Higher Serum Vitamin D Concentrations Are Longitudinally Associated with Better Global Quality of Life and Less Fatigue in Colorectal Cancer Survivors up to 2 Years after Treatment

Citation for published version (APA):

Koole, J. L., Bours, M. J. L., van Roekel, E. H., Breedveld-Peters, J. J. L., van Duijnhoven, F. J. B., van den Ouweland, J., Breukink, S. O., Janssen-Heijnen, M. L. G., Keulen, E. T. P., & Weijenberg, M. P. (2020). Higher Serum Vitamin D Concentrations Are Longitudinally Associated with Better Global Quality of Life and Less Fatigue in Colorectal Cancer Survivors up to 2 Years after Treatment. *Cancer Epidemiology Biomarkers & Prevention*, 29(6), 1135-1144. <https://doi.org/10.1158/1055-9965.epi-19-1522>

Document status and date:

Published: 01/06/2020

DOI:

[10.1158/1055-9965.epi-19-1522](https://doi.org/10.1158/1055-9965.epi-19-1522)

Document Version:

Publisher's PDF, also known as Version of record

Document license:

Taverne

Please check the document version of this publication:

- A submitted manuscript is the version of the article upon submission and before peer-review. There can be important differences between the submitted version and the official published version of record. People interested in the research are advised to contact the author for the final version of the publication, or visit the DOI to the publisher's website.
- The final author version and the galley proof are versions of the publication after peer review.
- The final published version features the final layout of the paper including the volume, issue and page numbers.

[Link to publication](#)

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal.

If the publication is distributed under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license above, please follow below link for the End User Agreement:

www.umlib.nl/taverne-license

Take down policy

If you believe that this document breaches copyright please contact us at:

repository@maastrichtuniversity.nl

providing details and we will investigate your claim.

Download date: 26 Apr. 2023

Higher Serum Vitamin D Concentrations Are Longitudinally Associated with Better Global Quality of Life and Less Fatigue in Colorectal Cancer Survivors up to 2 Years after Treatment

Janna L. Koole¹, Martijn J.L. Bours¹, Eline H. van Roekel¹, José J.L. Breedveld-Peters¹, Fränzel J.B. van Duijnhoven², Jody van den Ouweland³, Stéphanie O. Breukink⁴, Maryska L.G. Janssen-Heijnen^{1,5}, Eric T.P. Keulen⁶, and Matty P. Weijenberg¹

ABSTRACT

Background: Vitamin D status may be an important determinant of health-related quality of life of colorectal cancer survivors. The current study investigated longitudinal associations between serum 25-hydroxyvitamin D₃ (25OHD₃) concentrations and quality of life in stage I–III colorectal cancer survivors up to 2 years after treatment.

Methods: Patients with colorectal cancer ($n = 261$) were included upon diagnosis. Home visits (including blood sampling) were performed at diagnosis and at 6 weeks, 6 months, 1 year, and 2 years after treatment. Serum 25OHD₃ concentrations were measured using LC/MS-MS and adjusted for season. Validated questionnaires were used to assess global quality of life and cognitive functioning (EORTC-QLQ-C30), fatigue (EORTC-QLQ-C30 and Checklist Individual Strength, CIS), and depression and anxiety (Hospital Anxiety and Depression Scale). Statistical analyses were performed using linear mixed models and adjusted for sex, age, time since diagnosis, therapy, comorbidities, physical activity, and body mass index.

Results: At diagnosis, 45% of patients were vitamin D deficient (<50 nmol/L). After treatment, 25OHD₃ concentrations increased on average with 3.1 nmol/L every 6 months. In confounder-adjusted models, 20 nmol/L increments in 25OHD₃ were longitudinally associated with increased global quality of life [β 2.9; 95% confidence interval (CI), 1.5–4.3] and reduced fatigue (EORTC-QLQ-C30 subscale: β –3.5; 95% CI, –5.3 to –1.8 and CIS: β –2.8; 95% CI, –4.7 to –0.9). Observed associations were present both within and between individuals over time.

Conclusions: Higher concentrations of 25OHD₃ were longitudinally associated with better global quality of life and less fatigue in colorectal cancer survivors.

Impact: This study suggests that higher 25OHD₃ concentrations may be beneficial for colorectal cancer survivors. Future intervention studies are needed to corroborate these findings.

Introduction

Colorectal cancer survivors are susceptible to enduring physical and psychologic distress as a result of the tumor and therapy. A growing number of individuals are living with a history of colorectal cancer given the improving 5-year survival rate after colorectal cancer, currently 65% in the Netherlands (1, 2). Previous studies reported complaints of fatigue by more than one third of Dutch colorectal cancer survivors, especially within 5 years after diagnosis (3–5).

In addition, colorectal cancer survivors are at increased risk of impaired cognitive functioning and mental health problems such as depression (6–8).

Vitamin D is a potentially important determinant of health-related quality of life (HRQoL) of colorectal cancer survivors given its involvement in many cellular processes related to cognition (9, 10), depression (11, 12), and fatigue (13). Exposure of the skin to ultraviolet B radiation contributes to about two thirds of vitamin D supply, whereas about one third originates from vitamin D intake (14). Vitamin D deficiencies have become a concern of public health (15, 16), and the Dutch Health Council recommends women aged 50 to 70 (10 mcg) and men and women aged >70 (20 mcg) to daily supplement vitamin D (14). No specific guidelines for cancer survivors are available. To maintain adequate vitamin D concentrations, 15 to 30 minutes of daily sunlight exposure from March to November is generally sufficient for the general population (17). Colorectal cancer survivors are at risk of low vitamin D because of the negative impact of the cancer and chemotherapy on circulating concentrations (18–20). Moreover, the association between low vitamin D and increased colorectal cancer risk (21–24) makes colorectal cancer survivors particularly susceptible as low concentrations may be sustained after therapy. Inflammation may be an important confounder in the association between vitamin D and HRQoL. Inflammatory processes and vitamin D interact through 1,25-dihydroxyvitamin D, the biologically active metabolite also known as calcitriol, which is engaged in different intracellular inflammatory reactions (25). In addition, as inflammatory markers have been associated with increased fatigue

¹Department of Epidemiology, GROW School for Oncology and Developmental Biology, Maastricht University, Maastricht, the Netherlands. ²Division of Human Nutrition and Health, Wageningen University and Research, Wageningen, the Netherlands. ³Department of Clinical Chemistry, Canisius-Wilhelmina Hospital, Nijmegen, the Netherlands. ⁴Department of Surgery, GROW School for Oncology and Developmental Biology, Maastricht University Medical Centre+, Maastricht, the Netherlands. ⁵Department of Clinical Epidemiology, VieCuri Medical Centre, Venlo, the Netherlands. ⁶Department of Internal Medicine and Gastroenterology, Zuyderland Medical Centre, Sittard, the Netherlands.

Note: Supplementary data for this article are available at Cancer Epidemiology, Biomarkers & Prevention Online (<http://cebp.aacrjournals.org/>).

Corresponding Author: Janna L. Koole, Maastricht University, 6229HA Maastricht, the Netherlands. Phone: 314-3388-2043; Fax: 314-3388-4128; E-mail: annaleen.koole@maastrichtuniversity.nl

Cancer Epidemiol Biomarkers Prev 2020;29:1135–44

doi: 10.1158/1055-9965.EPI-19-1522

©2020 American Association for Cancer Research.

after cancer, a potential observed association between vitamin D and HRQoL may be altered by inflammation (26).

Few studies have investigated associations between vitamin D and HRQoL after colorectal cancer. An observational study among 453 stage II colorectal cancer survivors reported better symptom-related quality of life among participants using vitamin D supplements over 2 years of follow-up (27). Cancer survivors are known to be highly motivated to intentionally alter their lifestyle in order to make health improvements (28). Therefore, lifestyle recommendations that focus on the enhancement of vitamin D concentrations may be an opportunity to prevent or reduce problems of diminished HRQoL after colorectal cancer.

The objective of the present study was to investigate longitudinal associations of serum 25-hydroxyvitamin D₃ (25OHD₃) concentrations with HRQoL in colorectal cancer survivors from 6 weeks to 2 years after treatment, including global quality of life, fatigue, depression, anxiety, and cognitive functioning.

Materials and Methods

Study design and population

The EnCoRe study (Energy for Life after ColoRectal Cancer) is an ongoing prospective cohort study that was initiated in 2012 (Netherlands Trial Register no. NL6904; refs. 5, 29). The purpose of the EnCoRe study is to evaluate longitudinal associations between lifestyle factors and HRQoL, functioning, and prognosis after colorectal cancer. Patients with stage I–III colorectal cancer ≥18 years of age were recruited at diagnosis. Research dietitians performed home visits at diagnosis (prior to treatment) and at 6 weeks, 6 months, 1 year, and 2 years after the end of treatment. The study was approved by the Medical Ethics Committee of the University Hospital Maastricht and Maastricht University, the Netherlands. All participants signed informed consent. Exclusion criteria were: diagnosis with stage IV colorectal cancer, no home address in the Netherlands, inability to understand the Dutch language, and the presence of comorbidities obstructing successful study participation. Data used in the current analyses were based on the first 4.5 years of follow-up until November 1, 2016. Participants with at least 1 follow-up visit with available data on both 25OHD₃ and HRQoL were included in the current analyses. The final analyses contained 261 participants at diagnosis, 260 at 6 weeks, 213 at 6 months, 168 at 12 months, and 77 at 24 months after treatment. Response rate for inclusion was 46% and >90% for follow-up visits.

Measurement of 25OHD₃

At diagnosis, blood samples were drawn either during the first home visit or at the hospital, and were mostly nonfasting. At follow-up, fasting blood samples were drawn during home visits. Samples were collected in 8.5 mL serum tubes (BD Vacutainer SST II Advance) and pipetted into aliquots after centrifugation. Aliquots were stored at –80°C within 4 hours after blood draw until analysis. Serum 25OHD₃ concentrations were measured using LC/MS-MS at the Canisius-Wilhelmina Hospital in Nijmegen, the Netherlands (30). Interassay coefficients of variation were 5.3%, 3.1%, and 2.9% at 25OHD₃ concentrations of 39.0, 92.5, and 127.0 nmol/L, respectively. 25OHD₃ concentrations are a robust indicator and the most commonly used marker of vitamin D status (31).

Health-related quality of life

HRQoL was assessed during posttreatment time points. Global quality of life, cognitive functioning, and fatigue were assessed by the Quality of Life Questionnaire of the European Organisation for

Research and Treatment of Cancer (EORTC QLQ-C30, version 3.0; refs. 32, 33). Higher scores reflect better global quality of life, better cognitive functioning, and more fatigue (range, 0–100). Fatigue was also measured by the Checklist Individual Strength (CIS), a validated 20-item questionnaire that has been used in colorectal cancer survivors before (34, 35). Higher scores indicate more fatigue (range, 20–140). Depression and anxiety were assessed using the Hospital Anxiety and Depression Scale (HADS; refs. 36, 37). The HADS includes separate depression and anxiety scales, and higher scores indicate more depression and anxiety (range, 0–42).

Sociodemographic, lifestyle, and clinical data

Sociodemographic information was collected based on self-report. Cancer stage, type of therapy, and other clinical information were obtained from medical records. Information on comorbidities at diagnosis was retrieved from medical records by the Charlson Comorbidity Index (38) and during follow-up from the Self-Administered Comorbidity Questionnaire, which included heart condition; stroke; high blood pressure; asthma, chronic bronchitis, or chronic obstructive pulmonary disease; diabetes; stomach ulcer; kidney disease; liver disease; anemia or other disease of the blood; thyroid gland disease; depression; osteoarthritis; back pain; and rheumatoid arthritis (39).

Information on dietary supplement use, including vitamin D supplements, was collected during home visits. Information on (brand) name, dosage, frequency, duration, and ingredients was listed on standardized forms. Vitamin D intake from supplements was calculated in micrograms per day by multiplying daily frequency and dosage. At diagnosis, participants completed a semiquantitative 253-item food frequency questionnaire to retrospectively assess dietary vitamin D intake during the preceding year (40). Participants completed a 7-day dietary record as part of each posttreatment follow-up visit. Vitamin D levels in food products for both methods were obtained from the 2011 Dutch Food Composition Database (41). Methods were described in more detail previously (5, 42).

Measurements of height and weight were performed during home visits to determine body mass index (BMI). Physical activity was self-reported using the Short QUestionnaire to ASsess Health enhancing physical activity (SQUASH). Time spent in moderate-to-vigorous physical activity (MVPA; ≥3 metabolic equivalents of task) was calculated in hours/week (43). Sun exposure was self-reported at each time point by the number of days per week that participants had spent at least 15 minutes outside (in daylight), on average over the preceding month. The variable was dichotomized for the analyses in order to reflect adherence to the Dutch recommendations regarding sun exposure (spent 15 minutes outside on 7 or <7 days/week).

Plasma concentrations of inflammation markers IL6, IL8, IL10, and TNFα were measured for all time points. The selection of this set of inflammation markers was based on a review that evaluated inflammation markers that interact with vitamin D and that are specifically relevant in relation to colorectal cancer (25). Samples were collected in 6.0 mL EDTA plasma tubes (BD Vacutainer K2E) and pipetted into aliquots after centrifugation. Aliquots were stored at –80°C within 4 hours after blood draw until analysis. Measurements were performed using a custom-made multiplex assay and electrochemiluminescence (Meso Scale Diagnostics) at Wageningen University and Research, the Netherlands. Assay plates were analyzed on a QuickPlex SQ 120 plate reader (Meso Scale Diagnostics).

Statistical analyses

Descriptive analyses were performed to calculate means and SD for normally distributed variables and medians and interquartile ranges

Vitamin D and Quality of Life after Colorectal Cancer

Table 1. Sociodemographic and clinical characteristics of study participants at time of inclusion (colorectal cancer diagnosis), included in the EnCoRe study.

	Total population (<i>n</i> = 261) ^a	Vitamin D deficiency (<50 nmol/L) ^b	
		Yes (<i>n</i> = 111)	No (<i>n</i> = 136)
Age, mean (SD)	67 (9)	67 (9)	66 (9)
Sex, <i>n</i> (%)			
Men	179 (69)	78 (70)	91 (67)
Women	82 (31)	33 (30)	45 (33)
Education level, <i>n</i> (%)			
Low	66 (25)	26 (23)	35 (26)
Medium	105 (40)	48 (43)	53 (39)
High	89 (34)	37 (33)	47 (35)
Cancer type, <i>n</i> (%)			
Colon	159 (61)	72 (65)	79 (58)
Rectum	102 (39)	39 (35)	57 (42)
Cancer stage ^c , <i>n</i> (%)			
I	77 (30)	32 (30)	43 (34)
II	60 (23)	26 (24)	27 (21)
III	113 (43)	50 (46)	58 (45)
Treatment, <i>n</i> (%)			
Neither chemotherapy nor radiotherapy	143 (55)	60 (54)	77 (57)
Chemotherapy only	46 (18)	25 (23)	18 (13)
Radiotherapy only	20 (8)	6 (5)	10 (7)
Both chemotherapy and radiotherapy	52 (20)	20 (18)	31 (23)
Inflammatory markers, median (IQR)			
IL6 in pg/mL	1.1 (0.7–1.7)	1.2 (0.8–1.8)	1.0 (0.6–1.6)
IL8 in pg/mL	5.5 (4.4–8.0)	5.7 (4.4–8.0)	5.5 (4.5–7.9)
IL10 in pg/mL	0.3 (0.2–0.4)	0.3 (0.2–0.4)	0.3 (0.2–0.4)
TNF α in pg/mL	2.2 (1.8–2.8)	2.2 (1.9–2.7)	2.2 (1.7–2.8)
Summary z-score	–0.1 (–1.4 to 1.1)	–0.1 (–1.0 to 1.2)	–0.1 (–1.6 to 1.1)
Sun exposure ^d , <i>n</i> (%)			
<7 days	87 (33)	41 (37)	41 (30)
7 days	173 (67)	70 (63)	94 (70)
Number of comorbidities, <i>n</i> (%)			
None	49 (19)	18 (16)	29 (21)
1	57 (22)	21 (19)	36 (26)
≥ 2	155 (59)	72 (65)	71 (52)
BMI in kg/m ² , mean (SD)	28.4 (4.5)	29.2 (5.0)	27.6 (4.1)
Physical activity, median (IQR)			
LPA (hours/week)	11 (4–23)	11 (4–21)	12 (4–30)
MVPA (hours/week)	11 (5–20)	11 (4–19)	12 (6–21)
Current smoking, <i>n</i> (%)			
Yes	33 (13)	15 (14)	16 (12)
No	227 (87)	96 (86)	119 (88)

Abbreviations: BMI, body mass index; LPA, light physical activity; MVPA, moderate-to-vigorous physical activity.

^aThe number of participants may slightly vary for the different variables given a small percentage of missing data.^bFourteen participants had unknown 25OHD₃ status at time of inclusion.^cEleven participants had missing stage I, II, or III colorectal cancer.^dNumber of days per week spent at least 15 minutes outside (in daylight), on average over the preceding 4 weeks.

(IQR) for skewed variables. 25OHD₃ concentrations at each time point were adjusted for season using the week (1–52) of blood collection. Locally weighted regression analyses (LOWESS) were performed in order to account for the variation caused by seasonal differences (44, 45). For posttreatment time points, longitudinal associations between 25OHD₃ concentrations and HRQoL were investigated using linear mixed-model analyses. Each model contained a random intercept for subject. The addition of a random slope for vitamin D was evaluated in each model according to the change in log-restricted likelihood values. A separate model was used to disentangle inter- and intraindividual associations (46). Interindividual associations were

estimated by the centered person-mean value, indicating the difference between participants' mean 25OHD₃ concentrations over time. Intraindividual associations were estimated by the individual deviations from the person-mean, indicating individual changes in vitamin D concentration over time. Vitamin D was modeled as a continuous variable in units of 20 nmol/L and as a dichotomous variable (vitamin D concentration <50 nmol/L; yes/no; ref. 20). In addition, scoring on the outcomes was compared for 25OHD₃ concentrations of <30 nmol/L, 30–50 nmol/L, and >50 nmol/L. Vitamin D supplement use (yes/no), total vitamin D intake from diet and supplements (μ g/day), and vitamin D intake only from supplements (μ g/day) were

Table 2. Descriptive statistics of 25OHD₃ concentrations, vitamin D supplement use and dietary intake, HRQoL outcomes, and other important characteristics of patients with colorectal cancer from the EnCoRe study, from inclusion (colorectal cancer diagnosis) to 24 months after treatment.

	At diagnosis <i>n</i> = 261 ^a	Posttreatment follow-up measurements			
		6 weeks after treatment <i>n</i> = 260	6 months after treatment <i>n</i> = 213	12 months after treatment <i>n</i> = 168	24 months after treatment <i>n</i> = 77
25OHD ₃ concentration in nmol/L, median (IQR)	53.1 (40.6–67.6)	48.0 (35.2–66.1)	53.4 (41.8–69.4)	57.0 (42.5–70.5)	62.9 (46.5–82.4)
Vitamin D deficiency, <i>n</i> (%)					
Yes (<50 nmol/L)	111 (45)	133 (53)	87 (42)	60 (36)	22 (30)
No (≥50 nmol/L)	136 (55)	120 (47)	120 (58)	106 (64)	52 (70)
Vitamin D supplement use, <i>n</i> (%)					
Yes	62 (24)	49 (19)	44 (21)	38 (23)	17 (22)
No	194 (76)	207 (81)	163 (79)	129 (77)	60 (78)
Vitamin D intake from supplements in µg/d, median (IQR)	5 (5–10)	5 (5–10)	6 (5–15)	7 (5–20)	10 (5–20)
Dietary vitamin D intake ^b in µg/d, median (IQR)	3.5 (2.5–4.7)	4.1 (2.9–5.6)	3.7 (2.8–4.7)	3.8 (2.9–5.0)	3.7 (2.6–4.6)
EORTC QLQ-C30 ^c					
Global quality of life, mean (SD)		74.1 (18.3)	76.9 (19.0)	77.8 (18.0)	79.6 (18.5)
Range (min–max)		16.7–100	0–100	0–100	33.3–100
Cognitive functioning, mean (SD)		86.3 (20.8)	85.9 (19.1)	87.6 (19.4)	87.7 (15.1)
Range (min–max)		0–100	16.7–100	16.7–100	33.3–100
Fatigue, mean (SD)		29.1 (23.0)	23.6 (21.6)	21.6 (23.0)	19.3 (21.5)
Range (min–max)		0–100	0–100	0–100	0–88.9
CIS (incl. 4 subscales) ^d					
Total fatigue, mean (SD)		62.9 (26.4)	59.3 (27.4)	54.0 (25.9)	51.5 (25.2)
Range (min–max)		20–127	20–132	20–134	20–101
Subjective fatigue, mean (SD)		27.2 (13.3)	24.9 (12.8)	22.6 (12.3)	21.3 (12.8)
Range (min–max)		8–56	8–56	8–56	8–54
Reduced motivation, mean (SD)		12.3 (6.1)	12.0 (6.2)	10.9 (6.1)	10.5 (5.9)
Range (min–max)		4–28	4–27	4–28	4–27
Reduced physical activity, mean (SD)		10.6 (5.1)	9.6 (5.1)	8.6 (5.0)	8.1 (4.8)
Range (min–max)		3–21	3–21	3–21	3–21
Concentration problems, mean (SD)		12.9 (7.3)	12.8 (7.3)	11.9 (6.7)	11.6 (6.5)
Range (min–max)		5–34	5–33	5–31	5–28
HADS (incl. 2 subscales) ^e					
Total depression and anxiety, mean (SD)		7.4 (6.4)	7.4 (6.7)	6.8 (6.2)	5.6 (5.5)
Range (min–max)		0–30	0–32	0–33	0–28
Depression, mean (SD)		3.8 (3.6)	3.9 (3.9)	3.4 (3.4)	2.7 (3.1)
Range (min–max)		0–15	0–19	0–17	0–14
Anxiety, mean (SD)		3.6 (3.4)	3.5 (3.4)	3.4 (3.5)	2.9 (3.1)
Range (min–max)		0–16	0–15	0–16	0–16
Inflammatory markers, median (IQR)					
IL6 in pg/mL	1.1 (0.7–1.7)	1.5 (0.8–2.2)	1.2 (0.8–2.0)	0.9 (0.6–1.4)	0.9 (0.5–1.5)
IL8 in pg/mL	5.5 (4.4–8.0)	5.6 (4.4–7.3)	5.2 (4.4–7.0)	3.9 (3.1–4.8)	4.8 (3.8–6.2)
IL10 in pg/mL	0.3 (0.2–0.4)	0.4 (0.3–0.5)	0.4 (0.2–0.5)	0.2 (0.2–0.4)	0.2 (0.1–0.3)
TNFα in pg/mL	2.2 (1.8–2.8)	2.9 (2.4–3.8)	2.8 (2.3–3.6)	2.0 (1.6–2.5)	2.0 (1.6–2.9)
Summary z-score	−0.1 (−1.4 to 1.1)	−0.1 (−1.5 to 1.1)	−0.2 (−1.1 to 1.1)	−0.3 (−1.4 to 1.0)	0.1 (−1.5 to 1.3)
Sun exposure ^f , <i>n</i> (%)					
<7 days	87 (33)	119 (46)	87 (41)	65 (39)	32 (44)
7 days	173 (67)	141 (54)	124 (59)	102 (61)	40 (56)
Number of comorbidities, <i>n</i> (%)					
None	49 (19)	55 (21)	49 (21)	41 (24)	16 (21)
1	57 (22)	64 (25)	51 (24)	39 (23)	17 (22)
≥2	155 (59)	141 (54)	113 (53)	87 (52)	43 (57)
BMI in kg/m ² , mean (SD)	28.4 (4.5)	27.7 (4.4)	28.2 (4.4)	28.5 (4.5)	28.5 (5.0)
Physical activity, median (IQR)					
LPA (hours/week)	11 (4–23)	8 (2–16)	11 (4–21)	11 (3–22)	11 (5–24)
MVPA (hours/week)	11 (5–20)	7 (3–14)	9 (4–15)	9 (4–18)	8 (3–18)
Current smoking, <i>n</i> (%)					
Yes	33 (13)	24 (9)	17 (8)	17 (10)	6 (8)
No	227 (87)	235 (91)	194 (92)	150 (90)	66 (92)

Abbreviations: BMI, body mass index; LPA, light physical activity; MVPA, moderate-to-vigorous physical activity.

^aParticipants with at least 1 follow-up measurement with available data on both vitamin D and quality of life were included in the analyses. The numbers of included participants decrease during follow-up because data collected until November 1, 2016, were used and participants had not yet reached their follow-up measurement at that time. The number of participants may slightly vary for the different variables given a small percentage of missing data.^bDietary intake of vitamin D was measured by a semiquantitative 253-item food frequency questionnaire at diagnosis, and by a 7-day dietary record at each follow-up measurement.^cRanges EORTC QLQ-C30 subscales: 0–100.^dRanges CIS: total score, 20–140; subjective fatigue, 8–56; reduced motivation, 4–28; reduced physical activity, 3–21; concentration problems, 5–35.^eRanges HADS: total score, 0–42; depression, 0–21; anxiety, 0–21.^fNumber of days per week spent at least 15 minutes outside (in daylight), on average over the preceding 4 weeks.

Vitamin D and Quality of Life after Colorectal Cancer

modeled separately as the independent variable in multivariable adjusted models to obtain more insight into the role of vitamin D supplement use.

A summary score for the inflammatory markers was calculated by summing z-scores of natural-log transformed concentrations of IL6, IL8, and TNF α , and subtracting IL10, for each time point (47). Higher scores are indicative of higher inflammation.

Relevant confounders were preselected according to the literature and included sex, age at diagnosis, time since diagnosis (units of 6 months), cancer treatment (neither chemotherapy nor radiotherapy, chemotherapy only, radiotherapy only, or both), number of comorbidities (0, 1, or ≥ 2), MVPA (h/wk), and BMI (kg/m²). All time-dependent variables, including confounders, were included in the model as the repeated measurements. Other potential covariates that did not change the beta-coefficients for the relationship of main exposures with outcomes, and therefore not included in the models, were: inflammatory markers (z-score), 25OHD₃ at diagnosis, season, vitamin D intake from supplements ($\mu\text{g/d}$), and sun exposure (dichotomous: spent 15 minutes outside on 7 or <7 days/week). Subgroup analyses were performed for sex (men/women), age at diagnosis (<70 and ≥ 70 years), vitamin D supplement use (users/nonusers), and (neo-)adjuvant chemotherapy (yes/no).

Statistical analyses were performed using Stata15 (StataCorp.). *P* values <0.05 (two-sided) were considered as statistically significant.

Results

Participant characteristics

The 261 participants who enrolled in the study (31% women) had a mean $\pm$ SD age of 67 ± 9 years (Table 1). Median 25OHD₃ concentrations decreased after diagnosis and subsequently increased during follow-up (Table 2, Fig. 1). Almost half (45%) of participants were vitamin D deficient (<50 nmol/L) at diagnosis. Regarding the subgroups advised to use extra vitamin D, 24% of men and women aged ≥ 70 years ($n = 98$) were vitamin D supplement users at diagnosis. In addition, 33% of women aged 50–70 years ($n = 39$) used vitamin D supplements at diagnosis. Users of vitamin D supplements had higher serum concentrations compared with non-users (Fig. 1B). Participants who received chemotherapy had consistently lower concentrations compared with the group who did not receive chemotherapy (Fig. 1C).

Vitamin D and HRQoL

With every 6 months, global quality of life scores improved on average by 1.3 points [95% confidence interval (CI), 0.4–2.1], and fatigue levels decreased by 2.8 points (95% CI, -3.8 to -1.8 ; EORTC) and 3.3 points (95% CI, -4.3 to -2.2 ; CIS; Table 3, Supplementary Fig. S1).

25OHD₃ was longitudinally associated with better global quality of life (β 2.9; 95% CI, 1.5–4.3) and reduced fatigue (EORTC: β -3.5 ; 95% CI, -5.3 to -1.8 and CIS: β -2.8 ; 95% CI, -4.7 to -0.9). An increase of 20 nmol/L 25OHD₃ within an individual over time was associated with an average 3.7 point (95% CI, 1.7–5.6) higher global quality of life score. In addition, a 20 nmol/L higher mean 25OHD₃ concentration between individuals over time was associated with an average 2.1 point (95% CI, 0.2–4.1) higher global quality of life score. Intra- and interindividual associations of 25OHD₃ with fatigue were of similar degree when measured by the CIS (β 's -2.5 and -3.2 , respectively), whereas for fatigue as measured by the EORTC, the association for intraindividual changes was stronger compared with interindividual

Figure 1.

Median concentrations of 25OHD₃ over time from colorectal cancer diagnosis (study inclusion) to 24 months after treatment for patients with colorectal cancer included in the EnCoRe study, in the total population (with IQR) and stratified by sex (A), stratified by vitamin D supplement user type (B), and stratified by chemotherapy (C). The actual time between the measurement at diagnosis and the measurement 6 weeks after treatment can vary between individual patients due to differences in treatment duration.

differences (β 's -4.4 and -2.6 , respectively). Not being vitamin D deficient was longitudinally associated with better global quality of life (β 4.4; 95% CI, 1.6–7.2) and less fatigue (EORTC: β -5.2 ; 95% CI, -8.6 to -1.8 and CIS: β -5.4 ; 95% CI, -9.0 to -1.8), both within and between individuals (Table 3). Dose–response relations were observed for the outcomes global quality of life and fatigue when comparing the scoring of participants having severely deficient (<30 nmol/L) and

Table 3. Results of mixed-model analyses on changes of HRQoL outcomes over time and longitudinal associations of vitamin D with HRQoL outcomes in patients with colorectal cancer from the EnCoRe study followed up from diagnosis to 2 years after treatment.

	Change in quality of life outcome over time ^a β (95% CI)	Longitudinal associations of 25OHD ₃ concentrations (per 20 nmol/L)				Longitudinal associations of vitamin D deficiency (yes/no)			
		Adjusted model I ^b		Adjusted model II ^c		Adjusted model I ^b		Adjusted model II ^c	
		Overall β (95% CI)	Intraindividual β (95% CI)	Interindividual β (95% CI)	Overall β (95% CI)	Intraindividual β (95% CI)	Interindividual β (95% CI)	Overall β (95% CI)	Interindividual β (95% CI)
EORTC QLQ-C30 ^d									
Global quality of life	1.3 (0.4-2.1)	3.6 (2.2-5.0)	3.7 (1.7-5.6)	2.1 (0.2-4.1)	4.4 (1.6-7.2)	4.4 (0.9-7.9)	4.4 (-0.03 to 8.9)	4.4 (1.6-7.2)	4.4 (-0.03 to 8.9)
Cognitive function	-0.1 (-0.9 to 0.7)	0.3 (-1.1, 1.7)	0.1 (-1.9 to 2.0)	0.8 (-1.6 to 3.1)	0.9 (-1.9 to 3.6)	0.2 (-3.2 to 3.5)	3.2 (-2.0 to 8.5)	1.1 (-1.7 to 3.9)	3.2 (-2.0 to 8.5)
Fatigue	-2.8 (-3.8 to -1.8)	-4.8 (-6.5 to -3.1)	-4.4 (-6.9 to -2.0)	-2.6 (-5.1 to -0.2)	-7.4 (-10.8 to -4.1)	-5.0 (-9.3 to -0.7)	-5.7 (-11.2 to -0.1)	-5.2 (-8.6 to -1.8)	-5.7 (-11.2 to -0.1)
CIS (incl. 4 subscales) ^e									
Total fatigue	-3.3 (-4.3 to -2.2)	-4.5 (-6.4 to -2.7)	-2.5 (-4.9 to -0.1)	-3.2 (-6.1 to -0.2)	-8.1 (-11.7 to -4.5)	-4.3 (-8.6 to -0.1)	-7.9 (-14.6 to -1.3)	-5.4 (-9.0 to -1.8)	-7.9 (-14.6 to -1.3)
Subjective fatigue	-1.7 (-2.2 to -1.2)	-2.5 (-3.4 to -1.5)	-1.7 (-2.9 to -0.4)	-1.3 (-2.8 to 0.2)	-4.4 (-6.3 to -2.6)	-2.8 (-4.9 to -0.7)	-3.3 (-6.5 to -0.03)	-2.9 (-4.7 to -1.2)	-3.3 (-6.5 to -0.03)
Reduced motivation	-0.6 (-0.8 to -0.3)	-0.5 (-0.9 to -0.1)	0.0 (-0.6 to 0.6)	-0.4 (-1.0 to 0.3)	-1.5 (-2.3 to -0.6)	-0.8 (-1.9 to 0.3)	-1.4 (-2.9 to 0.1)	-1.0 (-1.9 to -0.1)	-1.4 (-2.9 to 0.1)
Reduced physical activity	-0.8 (-1.0 to -0.5)	-0.9 (-1.3 to -0.5)	-0.4 (-0.9 to 0.2)	-0.6 (-1.2 to -0.1)	-1.6 (-2.4 to -0.8)	-0.7 (-1.6 to 0.3)	-1.4 (-2.6 to -0.2)	-0.9 (-1.7 to -0.2)	-1.4 (-2.6 to -0.2)
Concentration problems	-0.3 (-0.6 to 0.1)	-0.7 (-1.2 to -0.1)	-0.5 (-1.2 to 0.3)	-0.8 (-1.7 to -0.1)	-0.8 (-1.8 to 0.2)	-0.1 (-1.4 to 1.2)	-1.9 (-3.7 to -0.1)	-0.7 (-1.7 to 0.4)	-1.9 (-3.7 to -0.1)
HADS (incl. 2 subscales) ^f									
Total anxiety/depression	-0.2 (-0.4 to 0.1)	-0.1 (-0.5 to 0.4)	0.1 (-0.5 to 0.6)	-0.2 (-1.0 to 0.5)	-0.1 (-0.9 to 0.7)	-0.1 (-0.9 to 1.0)	-0.6 (-2.3 to 1.1)	-0.1 (-0.9 to 0.7)	-0.6 (-2.3 to 1.1)
Depression	-0.1 (-0.3 to -0.0)	-0.1 (-0.4 to 0.1)	0.0 (-0.3 to 0.3)	-0.2 (-0.6 to 0.2)	-0.2 (-0.7 to 0.3)	-0.04 (-0.6 to 0.5)	-0.4 (-1.3 to 0.7)	-0.1 (-0.6 to 0.4)	-0.4 (-1.3 to 0.7)
Anxiety	-0.1 (-0.2 to 0.1)	0.1 (-0.2 to 0.3)	0.1 (-0.2 to 0.4)	0.0 (-0.4 to 0.4)	0.05 (-0.4 to 0.5)	0.1 (-0.4 to 0.7)	-0.2 (-1.1 to 0.7)	0.03 (-0.4 to 0.5)	-0.2 (-1.1 to 0.7)

Note: Bold content indicates statistically significant associations.

Abbreviation: β, beta-coefficient.

^aChanges in quality of life scores over time were measured from 6 weeks to 24 months after treatment, in units of 6 months.^bModel I: adjusted for sex, age at diagnosis.^cModel II: adjusted for sex, age at diagnosis, time since diagnosis, cancer treatment (neither chemotherapy nor radiotherapy, chemotherapy only, radiotherapy only, and both chemotherapy and radiotherapy), number of comorbidities (0, 1, and ≥2), MVPA (hours/week), BMI (kg/m²).^dRanges EORTC QLQ-C30 subscales: 0-100.^eRanges CIS: total score, 20-140; subjective fatigue, 8-56; reduced motivation, 4-28; reduced physical activity, 3-21; concentration problems, 5-35.^fRanges HADS: total score, 0-42; depression, 0-21; anxiety, 0-21.

Vitamin D and Quality of Life after Colorectal Cancer

Figure 2.

Comparison of scoring on quality of life outcomes over time for participants having severely deficient (<30 nmol/L) and deficient (30–50 nmol/L) 25OHD₃ concentrations, with participants having adequate concentrations (>50 nmol/L), from 6 weeks to 24 months after treatment in patients with colorectal cancer from the EnCoRe study.

deficient (30–50 nmol/L) 25OHD₃ concentrations, with participants having adequate concentrations (>50 nmol/L; **Fig. 2**).

Although not statistically significant, better global quality of life and less fatigue (EORTC QLQ-C30) were observed for individuals who changed from being nonuser to user of vitamin D supplements over time, whereas interindividual differences suggested lower global quality of life and more fatigue for supplement users compared with nonusers (**Table 4**). No longitudinal associations were found between the sum of dietary and supplemental vitamin D intake with global quality of life and fatigue, and between vitamin D intake from supplements only and these outcomes.

In subgroup analyses, longitudinal associations of 25OHD₃ concentrations with global quality of life and fatigue were only present in men (Supplementary Table S1).

Discussion

To our knowledge, the current study is the first to investigate longitudinal associations between serum 25OHD₃ concentrations and HRQoL outcomes in colorectal cancer survivors. Higher concentrations of 25OHD₃ were longitudinally associated with better global quality of life and less fatigue from 6 weeks up to 2 years after colorectal cancer treatment, both within and between individuals. In addition, having a vitamin D deficiency was associated with lower global quality of life and more fatigue. Intraindividual associations suggested better

global quality of life and less fatigue for individuals who started using vitamin D supplements during follow-up.

The prevalence of vitamin D deficiency increased from 45% at diagnosis to 53% 6 weeks after treatment (cutoff 50 nmol/L). The latter is slightly higher compared with the general Dutch population aged 60+ with prevalences of 34% to 51% (48). A study among 2,910 patients with stage I–IV colorectal cancer from Germany observed deficiencies in 84% of participants, yet samples were assessed approximately 3 weeks after diagnosis (49). Another study among 1,598 Scottish patients with stage I–III colorectal cancer found concentrations <25 nmol/L among 50% of participants approximately 15 weeks after treatment (50). This compares with only 10% of participants when using the cutoff point of 25 nmol/L in our population. Differences may be due to vitamin D supplement use, latitude, and time of sampling as concentrations in our study steadily increased after treatment. Nevertheless, the important health concern of vitamin D deficiency among this group of patients is indisputable.

Previous studies examining the potential relation between vitamin D and aspects of HRQoL are still inconclusive and scarce among population subgroups such as cancer survivors. A systematic literature review concluded a moderate positive effect of short-term vitamin D supplementation on HRQoL in clinical populations, not including cancer survivors (51). Despite the large heterogeneity between studies included in this review, results correspond to the observed associations

Table 4. Longitudinal associations of vitamin D supplement use, vitamin D intake from diet and supplements, and vitamin D intake from supplements only, with HRQoL outcomes in patients with colorectal cancer from the EnCoRe study followed up from diagnosis to 2 years after treatment.

	Vitamin D supplement use (no/yes) ^a			Sum vitamin D intake from diet and supplements (μg/d) ^b			Vitamin D intake from supplements (μg/d) ^c		
	Overall		Interindividual	Overall		Interindividual	Overall		Interindividual
	β (95% CI)	Intraindividual		β (95% CI)	Intraindividual		β (95% CI)	Intraindividual	
EORTC QLQ-C30									
Global quality of life	-1.5 (-6.2 to 3.2)	2.7 (-3.5 to 8.8)	-5.2 (-11.0 to 0.7)	-0.1 (-0.4 to 0.1)	0.2 (-0.1 to 0.6)	-0.4 (-0.7 to -0.1)	-0.05 (-0.4 to 0.3)	0.2 (-0.2 to 0.7)	-0.3 (-0.7 to 0.1)
Fatigue	0.3 (-5.5 to 6.0)	-5.6 (-12.9 to 1.8)	5.9 (-1.4 to 13.1)	0.1 (-0.2 to 0.4)	-0.2 (-0.6 to 0.2)	0.3 (-0.1 to 0.7)	-0.01 (-0.4 to 0.3)	-0.3 (-0.8 to 0.2)	0.2 (-0.3 to 0.7)
CIS									
Total fatigue	5.3 (-1.0 to 11.6)	4.2 (-3.2 to 11.7)	6.8 (-1.8 to 15.5)	0.2 (-0.1 to 0.5)	0.2 (-0.3 to 0.6)	0.3 (-0.2 to 0.7)	0.4 (-0.0 to 0.8)	0.5 (0.01 to 1.0)	0.2 (-0.4 to 0.8)

Note: Bold content indicates statistically significant associations.

^aAdjusted for sex, age at diagnosis, overall supplement use (y/n), time since diagnosis, cancer treatment (neither chemotherapy nor radiotherapy, chemotherapy only, radiotherapy only, or both chemotherapy and radiotherapy), number of comorbidities (0, 1, or ≥2), MVPA (hours/week), and BMI (kg/m²).

^bAdjusted for sex, age at diagnosis, overall supplement use (y/n), total energy intake (kcal/d), time since diagnosis, cancer treatment (neither chemotherapy nor radiotherapy, chemotherapy only, radiotherapy only, or both chemotherapy and radiotherapy), number of comorbidities (0, 1, or ≥2), MVPA (hours/week), and BMI (kg/m²).

^cAdjusted for sex, age at diagnosis, time since diagnosis, cancer treatment (neither chemotherapy nor radiotherapy, chemotherapy only, radiotherapy only, or both chemotherapy and radiotherapy), number of comorbidities (0, 1, or ≥2), MVPA (hours/week), and BMI (kg/m²).

of the current study. Regarding fatigue, contradicting results were found in two intervention studies. A randomized placebo-controlled trial among patients with chronic fatigue syndrome found no effect of vitamin D supplementation (52), whereas another trial for treating fatigue among otherwise healthy individuals observed lower fatigue among vitamin D supplement users compared with a placebo group (53). A retrospective study among 100 ambulatory advanced cancer patients, including patients with gastrointestinal cancer, found no association between vitamin D concentrations and symptoms of depression, anxiety, and fatigue (54). Our study neither found associations between serum 25OHD₃ concentrations and depression, anxiety, and cognitive functioning, potentially caused by the lack of variation over time in these outcomes. In summary, results from the current study are largely in line with the literature, yet comparing results is difficult given the large differences in methodology and study parameters between studies.

The observed effect sizes for associations with global quality of life and fatigue were small and could raise questions about clinical relevance. Minimally important differences for global quality of life and fatigue (EORTC QLQ-C30) were defined as 10 and 9 points (55, 56), respectively, and for overall fatigue (CIS) as 10 points (57). Increments of 20 nmol/L 25OHD₃ were used as clinically relevant contrast as it resembled 1 SD and represented realistic fluctuations in intraindividual 25OHD₃ concentrations. Although effect sizes were larger in dichotomous analyses (vitamin D deficiency yes/no), none of the effect sizes reached the level of clinical relevance. Nevertheless, observed beta-coefficients suggest the presence of a longitudinal association between higher 25OHD₃ and better HRQoL.

An important strength of the current study was the prospective character with repeated measurements of 25OHD₃ and HRQoL outcomes. Further, follow-up response rates were high, and the percentage of missing data was low. Another strength was the use of mixed-model analysis techniques that enabled disentangling of inter- and intraindividual associations, thereby providing additional insights into the nature of the associations. A limitation of the current study concerns the inability to draw conclusions on causality. Observed associations could be due to the fact that colorectal cancer survivors with poorer HRQoL spend more time indoors, resulting in lower 25OHD₃ concentrations. However, adjustment for sun exposure did not change the associations. In addition, despite adjustment for important variables such as therapy, residual confounding may have occurred, as observed associations could be caused by unmeasured underlying factors related to the cancer that also influenced biological mechanisms related to 25OHD₃ and HRQoL.

The use of vitamin D supplements was low despite the national recommendations that applied to about half of the study population. Although vitamin D supplement users on average had higher serum 25OHD₃ concentrations compared with nonusers, no direct beneficial association of vitamin D supplement use with HRQoL was found. In addition, associations of 25OHD₃ with global quality of life and fatigue were also present in participants not using vitamin D supplements. In fact, vitamin D supplement use itself seemed associated with poorer HRQoL and more fatigue in inter-individual associations. An alternative explanation, also hypothesized by Patterson and colleagues (58), is that supplement use may be a coping strategy rather than actually improving health of cancer survivors. Associations with supplement use should therefore be interpreted with caution, because individuals with poorer HRQoL and more fatigue may begin to use (vitamin D) supplements as a way to alleviate complaints (5). Intraindividual associations

contrarily suggested that individuals who started using supplements during follow-up were generally likely to report better global quality of life and less fatigue. Whether this association is causal needs to be addressed in intervention studies.

Our study does not provide strong enough evidence to formulate recommendations on the necessity of vitamin D supplementation for colorectal cancer survivors. However, higher 25OHD₃ concentrations seem beneficial, and colorectal cancer survivors are advised to follow the guidelines for the general Dutch population because no specific guidelines on vitamin D for cancer survivors are available. According to the national guidelines, women aged >50 and women and men aged >70 are recommended to use vitamin D supplements (10 and 20 mcg, respectively) to ensure adequate blood concentrations for the prevention of osteoporosis (14). In addition, all colorectal cancer survivors should spend the recommended time outdoors to sufficiently expose their skin to sunlight (17). In practice, however, many colorectal cancer survivors seem unaware of the prevailing recommendations, potentially as a consequence of the lack of proper information provided.

In conclusion, our results suggest that higher 25OHD₃ concentrations are longitudinally associated with better global quality of life and reduced fatigue in colorectal cancer survivors. It is important for colorectal cancer survivors to become aware of the national guidelines regarding sun exposure and vitamin D supplementation, and individual 25OHD₃ status should be monitored by medical professionals. Placebo-controlled randomized trials are needed to examine the potential advantage of the use of vitamin D supplements in colorectal cancer survivors for improvement of HRQoL and fatigue to clarify questions on cause and effect and to deepen the understanding of possible underlying mechanisms.

Disclosure of Potential Conflicts of Interest

No potential conflicts of interest were disclosed.

References

- Denlinger CS, Barsevick AM. The challenges of colorectal cancer survivorship. *J Natl Compr Canc Netw* 2009;7:883–93.
- Integraal Kankercentrum Nederland. Nederlandse Kankerregistratie; 2019. Available from: https://www.cijfersoverkanker.nl/selecties/Overleving_darm/img5c598055e0605.
- Thong MS, Mols F, Wang XS, Lemmens VE, Smilde TJ, van de Poll-Franse LV. Quantifying fatigue in (long-term) colorectal cancer survivors: a study from the population-based patient reported outcomes following initial treatment and long term evaluation of survivorship registry. *Eur J Cancer* 2013;49:1957–66.
- Husson O, Mols F, van de Poll-Franse L, de Vries J, Schep G, Thong MS. Variation in fatigue among 6011 (long-term) cancer survivors and a normative population: a study from the population-based PROFILES registry. *Support Care Cancer* 2015;23:2165–74.
- Koole JL, Bours MJL, Breedveld-Peters JLL, van Roekel EH, Breukink SO, Janssen-Heijnen MLG, et al. Is dietary supplement use longitudinally associated with fatigue in stage I-III colorectal cancer survivors? *Clin Nutr* 2020;39:234–41.
- Vardy J, Dhillon HM, Pond GR, Rourke SB, Xu W, Dodd A, et al. Cognitive function and fatigue after diagnosis of colorectal cancer. *Ann Oncol* 2014;25:2404–12.
- Cruzado JA, Lopez-Santiago S, Martinez-Marin V, Jose-Moreno G, Custodio AB, Feliu J. Longitudinal study of cognitive dysfunctions induced by adjuvant chemotherapy in colon cancer patients. *Support Care Cancer* 2014;22:1815–23.
- Lloyd S, Baraghoshi D, Tao R, Garrido-Laguna I, Gilcrease GW 3rd, Whisenant J, et al. Mental health disorders are more common in colorectal cancer survivors and associated with decreased overall survival. *Am J Clin Oncol* 2019;42:355–62.
- Krause D, Roupas P. Effect of vitamin intake on cognitive decline in older adults: evaluation of the evidence. *J Nutr Health Aging* 2015;19:745–53.
- Goodwill AM, Szoek C. A systematic review and meta-analysis of the effect of low vitamin D on cognition. *J Am Geriatr Soc* 2017;65:2161–8.
- Anglin RE, Samaan Z, Walter SD, McDonald SD. Vitamin D deficiency and depression in adults: systematic review and meta-analysis. *Br J Psychiatry* 2013;202:100–7.
- Parker GB, Brothie H, Graham RK. Vitamin D and depression. *J Affect Disord* 2017;208:56–61.
- Hock AD. Review: vitamin D3 deficiency results in dysfunctions of immunity with severe fatigue and depression in a variety of diseases. *In Vivo* 2014;28:133–45.
- Health Council of the Netherlands. Evaluation of the dietary reference values for vitamin D. The Hague (the Netherlands): Health Council of the Netherlands; 2012.
- Hossein-nezhad A, Holick MF. Optimize dietary intake of vitamin D: an epigenetic perspective. *Curr Opin Clin Nutr Metab Care* 2012;15:567–79.
- Cashman KD, Dowling KG, Skrabakova Z, Gonzalez-Gross M, Valtuena J, De Henauw S, et al. Vitamin D deficiency in Europe: pandemic? *Am J Clin Nutr* 2016;103:1033–44.
- Dutch Cancer Society. The relation between cancer, sun exposure and vitamin D. Amsterdam (the Netherlands): Dutch Cancer Society; 2010.
- Savoie MB, Paciorek A, Zhang L, Van Blarigan EL, Sommovilla N, Abrams D, et al. Vitamin D levels in patients with colorectal cancer before and after treatment initiation. *J Gastrointest Cancer* 2019;50:769–79.
- Fakih MG, Trump DL, Johnson CS, Tian L, Muindi J, Sunga AY. Chemotherapy is linked to severe vitamin D deficiency in patients with colorectal cancer. *Int J Colorectal Dis* 2009;24:219–24.
- Holick MF. Vitamin D deficiency. *N Engl J Med* 2007;357:266–81.

Authors' Contributions

Conception and design: J.L. Koole, M.J.L. Bours, M.P. Weijnenberg

Development of methodology: J.L. Koole, M.J.L. Bours, E.H. van Roekel, J.J.L. Breedveld-Peters, M.P. Weijnenberg

Acquisition of data (provided animals, acquired and managed patients, provided facilities, etc.): J.L. Koole, E.H. van Roekel, J.J.L. Breedveld-Peters, F.J.B. van Duijnhoven, E.T.P. Keulen

Analysis and interpretation of data (e.g., statistical analysis, biostatistics, computational analysis): J.L. Koole, F.J.B. van Duijnhoven, J. van den Ouweland, M.P. Weijnenberg

Writing, review, and/or revision of the manuscript: J.L. Koole, M.J.L. Bours, E.H. van Roekel, J.J.L. Breedveld-Peters, F.J.B. van Duijnhoven, J. van den Ouweland, S.O. Breukink, M.L.G. Janssen-Heijnen, E.T.P. Keulen, M.P. Weijnenberg

Administrative, technical, or material support (i.e., reporting or organizing data, constructing databases): J.L. Koole

Study supervision: M.J.L. Bours, M.P. Weijnenberg

Acknowledgments

The EnCoRe study was supported by grants from the Stichting Alpe d'HuZes within the research program "Leven met kanker" of the Dutch Cancer Society (Grant Nos. UM-2010-4867 and UM-2012-5653), by grants from Kankeronderzoeksfonds Limburg as part of Health Foundation Limburg (Grant No. 00005739), and by a grant from Wereld Kanker Onderzoek Fonds (WKOF), as part of the World Cancer Research Fund International grant program (grant number 2016/1620).

We would like to thank all participants of the EnCoRe study and the health professionals in the three hospitals involved in the recruitment of participants of the study: Maastricht University Medical Centre+, VieCuri Medical Centre, and Zuyderland Medical Centre. We would also like to thank the MEMIC center for data and information management for facilitating the logistic processes and data management of our study. We thank Michiel Balvers and Nhien Ly for their work on the inflammation markers. Finally, we would like to thank the research dietitians and research assistants who are responsible for patient inclusion and follow-up, performing home visits, as well as data collection and processing.

The costs of publication of this article were defrayed in part by the payment of page charges. This article must therefore be hereby marked *advertisement* in accordance with 18 U.S.C. Section 1734 solely to indicate this fact.

Received December 9, 2019; revised February 4, 2020; accepted March 31, 2020; published first April 3, 2020.

21. Ma Y, Zhang P, Wang F, Yang J, Liu Z, Qin H. Association between vitamin D and risk of colorectal cancer: a systematic review of prospective studies. *J Clin Oncol* 2011;29:3775–82.
22. Giovannucci E. Epidemiology of vitamin D and colorectal cancer. *Anticancer Agents Med Chem* 2013;13:11–9.
23. McCullough ML, Zoltick ES, Weinstein SJ, Fedirko V, Wang M, Cook NR, et al. Circulating vitamin D and colorectal cancer risk: an international pooling project of 17 cohorts. *J Natl Cancer Inst* 2019;111:158–69.
24. Lee JE, Li H, Chan AT, Hollis BW, Lee IM, Stampfer MJ, et al. Circulating levels of vitamin D and colon and rectal cancer: the Physicians' Health Study and a meta-analysis of prospective studies. *Cancer Prev Res* 2011;4:735–43.
25. van Harten-Gerritsen AS, Balvers MG, Witkamp RF, Kampman E, van Duin-hoven FJ. Vitamin D, inflammation, and colorectal cancer progression: a review of mechanistic studies and future directions for epidemiological studies. *Cancer Epidemiol Biomarkers Prev* 2015;24:1820–8.
26. Bower JE. Cancer-related fatigue—mechanisms, risk factors, and treatments. *Nat Rev Clin Oncol* 2014;11:597–609.
27. Lewis C, Xun P, He K. Vitamin D supplementation and quality of life following diagnosis in stage II colorectal cancer patients: a 24-month prospective study. *Support Care Cancer* 2016;24:1655–61.
28. Demark-Wahnefried W, Peterson B, McBride C, Lipkus I, Clipp E. Current health behaviors and readiness to pursue life-style changes among men and women diagnosed with early stage prostate and breast carcinomas. *Cancer* 2000;88:674–84.
29. van Roekel EH, Bours MJ, de Brouwer CP, Ten Napel H, Sanduleanu S, Beets GL, et al. The applicability of the international classification of functioning, disability, and health to study lifestyle and quality of life of colorectal cancer survivors. *Cancer Epidemiol Biomarkers Prev* 2014;23:1394–405.
30. van den Ouweland JM, Beijers AM, van Daal H. Overestimation of 25-hydroxyvitamin D3 by increased ionisation efficiency of 3-epi-25-hydroxyvitamin D3 in LC-MS/MS methods not separating both metabolites as determined by an LC-MS/MS method for separate quantification of 25-hydroxyvitamin D3, 3-epi-25-hydroxyvitamin D3 and 25-hydroxyvitamin D2 in human serum. *J Chromatogr B Analyt Technol Biomed Life Sci* 2014;967:195–202.
31. Seamans KM, Cashman KD. Existing and potentially novel functional markers of vitamin D status: a systematic review. *Am J Clin Nutr* 2009;89:1997S–2008S.
32. Aaronson NK, Ahmedzai S, Bergman B, Bullinger M, Cull A, Duez NJ, et al. The European Organization for Research and Treatment of Cancer QLQ-C30: a quality-of-life instrument for use in international clinical trials in oncology. *J Natl Cancer Inst* 1993;85:365–76.
33. Fayes PM, Aaronson NK, Bjordal K, Groenvold M, Curran D, Bottomley A; on behalf of the EORTC Quality of Life Group. The EORTC QLQ-C30 scoring manual. 3rd ed. Brussels (Belgium): European Organisation for Research and Treatment of Cancer; 2001.
34. Vercoulen JH, Swanink CM, Fennis JF, Galama JM, van der Meer JW, Bleijenberg G. Dimensional assessment of chronic fatigue syndrome. *J Psychosom Res* 1994;38:383–92.
35. Servaes P, Verhagen C, Bleijenberg G. Fatigue in cancer patients during and after treatment: prevalence, correlates and interventions. *Eur J Cancer* 2002;38:27–43.
36. Zigmond AS, Snaith RP. The hospital anxiety and depression scale. *Acta Psychiatr Scand* 1983;67:361–70.
37. Bjelland I, Dahl AA, Haug TT, Neckelmann D. The validity of the Hospital Anxiety and Depression Scale. An updated literature review. *J Psychosom Res* 2002;52:69–77.
38. Charlson ME, Pompei P, Ales KL, MacKenzie CR. A new method of classifying prognostic comorbidity in longitudinal studies: development and validation. *J Chronic Dis* 1987;40:373–83.
39. Sangha O, Stucki G, Liang MH, Fossel AH, Katz JN. The Self-Administered Comorbidity Questionnaire: a new method to assess comorbidity for clinical and health services research. *Arthritis Rheum* 2003;49:156–63.
40. Koole JL, Bours MJL, Breedveld-Peters JLL, van Roekel EH, van Dongen M, Eussen S, et al. Evaluating the validity of a food frequency questionnaire in comparison with a 7-day dietary record for measuring dietary intake in a population of survivors of colorectal cancer. *J Acad Nutr Diet* 2020;120:245–57.
41. RIVM/Voedingscentrum. Dutch Food Composition (NEVO) Table 2011. The Hague (the Netherlands): Netherlands Nutrition Center; 2011.
42. Breedveld-Peters JLL, Koole JL, Muller-Schulte E, van der Linden BWA, Windhausen C, Bours MJL, et al. Colorectal cancers survivors' adherence to lifestyle recommendations and cross-sectional associations with health-related quality of life. *Br J Nutr* 2018;120:188–97.
43. Wendel-Vos GC, Schuit AJ, Saris WH, Kromhout D. Reproducibility and relative validity of the short questionnaire to assess health-enhancing physical activity. *J Clin Epidemiol* 2003;56:1163–9.
44. Zhang H, Ahn J, Yu K. Comparing statistical methods for removing seasonal variation from vitamin D measurements in case-control studies. *Stat Interface* 2011;4:85–93.
45. Cleveland WS, Devlin SJ. Locally weighted regression: an approach to regression analysis by local fitting. *J Am Statist Assoc* 1988;83:596–610.
46. Twisk JWR, de Vente W. Hybrid models were found to be very elegant to disentangle longitudinal within- and between-subject relationships. *J Clin Epidemiol* 2019;107:66–70.
47. Hopkins MH, Owen J, Ahearn T, Fedirko V, Flanders WD, Jones DP, et al. Effects of supplemental vitamin D and calcium on biomarkers of inflammation in colorectal adenoma patients: a randomized, controlled clinical trial. *Cancer Prev Res* 2011;4:1645–54.
48. Health Council of the Netherlands. Towards an adequate intake of vitamin D. The Hague (the Netherlands): Health Council of the Netherlands, 2008.
49. Maalmi H, Walter V, Jansen L, Chang-Claude J, Owen RW, Ulrich A, et al. Relationship of very low serum 25-hydroxyvitamin D3 levels with long-term survival in a large cohort of colorectal cancer patients from Germany. *Eur J Epidemiol* 2017;32:961–71.
50. Zgaga L, Theodoratou E, Farrington SM, Din FV, Ooi LY, Glodzik D, et al. Plasma vitamin D concentration influences survival outcome after a diagnosis of colorectal cancer. *J Clin Oncol* 2014;32:2430–9.
51. Hoffmann MR, Senior PA, Mager DR. Vitamin D supplementation and health-related quality of life: a systematic review of the literature. *J Acad Nutr Diet* 2015;115:406–18.
52. Witham MD, Adams F, McSwiggan S, Kennedy G, Kabir G, Belch JJ, et al. Effect of intermittent vitamin D3 on vascular function and symptoms in chronic fatigue syndrome—a randomised controlled trial. *Nutr Metab Cardiovasc Dis* 2015;25:287–94.
53. Nowak A, Boesch L, Andres E, Battegay E, Hornemann T, Schmid C, et al. Effect of vitamin D3 on self-perceived fatigue: a double-blind randomized placebo-controlled trial. *Medicine* 2016;95:e5353.
54. Dev R, Del Fabbro E, Schwartz GG, Hui D, Palla SL, Gutierrez N, et al. Preliminary report: vitamin D deficiency in advanced cancer patients with symptoms of fatigue or anorexia. *Oncologist* 2011;16:1637–41.
55. Cocks K, King MT, Velikova G, Martyn St-James M, Fayes PM, Brown JM. Evidence-based guidelines for determination of sample size and interpretation of the European Organisation for the Research and Treatment of Cancer Quality of Life Questionnaire Core 30. *J Clin Oncol* 2011;29:89–96.
56. Jayadevappa R, Cook R, Chhatre S. Minimal important difference to infer changes in health-related quality of life—a systematic review. *J Clin Epidemiol* 2017;89:188–98.
57. Peters JB, Heijdra YF, Daudey L, Boer LM, Molema J, Dekhuijzen PN, et al. Course of normal and abnormal fatigue in patients with chronic obstructive pulmonary disease, and its relationship with domains of health status. *Patient Educ Couns* 2011;85:281–5.
58. Patterson RE, Neuhauser ML, Hedderson MM, Schwartz SM, Standish LJ, Bowen DJ. Changes in diet, physical activity, and supplement use among adults diagnosed with cancer. *J Am Diet Assoc* 2003;103:323–8.

Cancer Epidemiology, Biomarkers & Prevention

Higher Serum Vitamin D Concentrations Are Longitudinally Associated with Better Global Quality of Life and Less Fatigue in Colorectal Cancer Survivors up to 2 Years after Treatment

Janna L. Koole, Martijn J.L. Bours, Eline H. van Roekel, et al.

Cancer Epidemiol Biomarkers Prev 2020;29:1135-1144. Published OnlineFirst April 3, 2020.

Updated version Access the most recent version of this article at:
doi:[10.1158/1055-9965.EPI-19-1522](https://doi.org/10.1158/1055-9965.EPI-19-1522)

Supplementary Material Access the most recent supplemental material at:
<http://cebp.aacrjournals.org/content/suppl/2020/04/03/1055-9965.EPI-19-1522.DC1>

Cited articles This article cites 52 articles, 13 of which you can access for free at:
<http://cebp.aacrjournals.org/content/29/6/1135.full#ref-list-1>

E-mail alerts [Sign up to receive free email-alerts](#) related to this article or journal.

Reprints and Subscriptions To order reprints of this article or to subscribe to the journal, contact the AACR Publications Department at pubs@aacr.org.

Permissions To request permission to re-use all or part of this article, use this link
<http://cebp.aacrjournals.org/content/29/6/1135>.
Click on "Request Permissions" which will take you to the Copyright Clearance Center's (CCC) Rightslink site.