Augmented cognitive behavioral therapy for post stroke depressive symptoms

Citation for published version (APA):

Kootker, J. A., Rasquin, S. M., Lem, F. C., van Heugten, C. M., Fasotti, L., & Geurts, A. C. (2017). Augmented cognitive behavioral therapy for post stroke depressive symptoms: a randomized controlled trial. *Archives of Physical Medicine and Rehabilitation*, 98(4), 687–694. <https://doi.org/10.1016/j.apmr.2016.10.013>

Document status and date:

Published: 01/04/2017

DOI:

[10.1016/j.apmr.2016.10.013](https://doi.org/10.1016/j.apmr.2016.10.013)

Document Version:

Publisher's PDF, also known as Version of record

Document license:

Taverne

Please check the document version of this publication:

- A submitted manuscript is the version of the article upon submission and before peer-review. There can be important differences between the submitted version and the official published version of record. People interested in the research are advised to contact the author for the final version of the publication, or visit the DOI to the publisher's website.
- The final author version and the galley proof are versions of the publication after peer review.
- The final published version features the final layout of the paper including the volume, issue and page numbers.

[Link to publication](#)

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal.

If the publication is distributed under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license above, please follow below link for the End User Agreement:

www.umlib.nl/taverne-license

Take down policy

If you believe that this document breaches copyright please contact us at:

repository@maastrichtuniversity.nl

providing details and we will investigate your claim.

ORIGINAL RESEARCH

Augmented Cognitive Behavioral Therapy for Poststroke Depressive Symptoms: A Randomized Controlled Trial

Joyce A. Kootker, MSc,^a Sascha M.C. Rasquin, PhD,^{b,c} Frederik C. Lem, MD,^d
Caroline M. van Heugten, PhD,^{e,f} Luciano Fasotti, PhD,^{g,h} Alexander C.H. Geurts, PhD^{a,g}

From the ^aDepartment of Rehabilitation, Donders Institute for Brain, Cognition and Behaviour, Radboud University Medical Center, Nijmegen; ^bAdelante Rehabilitation Center, Hoensbroek; ^cCAPHRI, Department of Rehabilitation Medicine, Maastricht University Medical Center, Maastricht; ^dGroot Klimmendaal, Rehabilitation Center, Arnhem; ^eSchool for Mental Health and Neuroscience, Department of Psychiatry and Neuropsychology, Faculty of Health, Medicine and Life Sciences, Maastricht University, Maastricht; ^fDepartment of Neuropsychology and Psychopharmacology, Faculty of Psychology and Neuroscience, Maastricht University, Maastricht; ^gDonders Institute for Brain, Cognition and Behaviour, Radboud University, Nijmegen; and ^hKlimmendaal Rehabilitation Center, Arnhem, The Netherlands.

Abstract

Objective: To evaluate the effectiveness of individually tailored cognitive behavioral therapy (CBT) for reducing depressive symptoms with or without anxiety poststroke.

Design: Multicenter, assessor-blinded, randomized controlled trial.

Setting: Ambulatory rehabilitation setting.

Participants: Patients who had a Hospital Anxiety and Depression Scale-depression subscale (HADS-D) score >7 at least 3 months poststroke (N=61).

Interventions: Participants were randomly allocated to either augmented CBT or computerized cognitive training (CCT). The CBT intervention was based on the principles of recognizing, registering, and altering negative thoughts and cognitions. CBT was augmented with goal-directed real-life activity training given by an occupational or movement therapist.

Main Outcome Measures: HADS-D was the primary outcome, and measures of participation and quality of life were secondary outcomes. Outcome measurements were performed at baseline, immediately posttreatment, and at 4- and 8-month follow-up. Analysis was performed with linear mixed models using group (CBT vs CCT) as the between-subjects factor and time (4 assessments) as the within-subjects factor.

Results: Mixed model analyses showed a significant and persistent time effect for HADS-D (mean difference, -4.6; 95% confidence interval, -5.7 to -3.6; $P<.001$) and for participation and quality of life in both groups. There was no significant group $\times$ time effect for any of the outcome measures.

Conclusions: Our augmented CBT intervention was not superior to CCT for the treatment of mood disorders after stroke. Future studies should determine whether both interventions are better than natural history.

Archives of Physical Medicine and Rehabilitation 2017;98:687-94

© 2016 by the American Congress of Rehabilitation Medicine

Approximately one third of all stroke survivors experience depressive symptoms at some point in time,¹ with a negative effect on the outcome of rehabilitation and quality of life.² Hence,

adequate treatment of poststroke depressive symptoms is of the utmost importance. Typically, studies on the treatment of depressive symptoms after stroke have focused on pharmacologic interventions. However, drug treatment only produces small improvements in mood, and it is associated with a frequent occurrence of side effects,^{3,4} restricting routine prescription. Because poststroke depressive symptoms are strongly associated with individuals' perception of, and coping with the consequences

Supported by The Netherlands organization for health research and development, ZonMw (grant no. 89000004).

Dutch Trial Register Clinical Trial Registration No.: NTR2999.

Disclosures: A.C.H.G. declares financial relationships with Ipsen and Merz outside the submitted work. The other authors have nothing to disclose.

of, stroke,⁵ a psychological treatment approach seems warranted. Because cognitive behavioral therapy (CBT) is aimed at changing irrational cognitions and negative thoughts, its effects may endure after treatment, and chances of relapse of symptoms may be smaller compared with pharmacotherapy.⁶⁻⁸ In addition, the side-effect profile of psychological treatment is favorable compared with medication.⁶ However, psychological treatments of post-stroke mood disorders have not yet yielded convincing results.^{9,10}

To our knowledge, only 1 randomized controlled trial has been reported investigating psychological treatment of poststroke depressive symptoms.¹⁰ This study yielded inconclusive results regarding the effectiveness of CBT. As the authors acknowledged, this work was a pioneering study with several methodologic weaknesses that compromised its quality. For instance, cognitive impairments were not considered in the inclusion criteria, and the therapy content was insufficiently adjusted to the cognitive consequences of stroke. In addition, the training of therapists was limited. Other researchers argued that to optimize the effectiveness of CBT in patients with stroke, cognitive and emotional impairments and limited awareness of deficits should be taken into account.⁸ It was also recommended that CBT should be augmented with real-life activity training to help patients and their caregivers set and attain realistic goals aimed at social participation, taking into account their motor and cognitive impairments.⁸ In addition, anxiety symptoms should be recognized and addressed in CBT treatment because these are often associated with depressive symptoms after stroke.^{11,12}

In a pilot study, we already assessed the feasibility of a CBT intervention aimed at reducing depressive symptoms in stroke survivors.¹³ Three of 5 participants showed positive results on mood and quality of life immediately after treatment, and these positive results were retained at 3-month follow-up. These findings justified the conduct of a multicenter randomized controlled trial as part of the Restore4Stroke study in The Netherlands.^{14,15} In the present study, we hypothesized that patients treated with individually tailored augmented CBT would show a larger decrease in depressive symptoms and more improvement in social participation and quality of life than those receiving computerized cognitive training (CCT). CCT was selected as the comparator intervention to control for Hawthorne effects, in the expectation that CCT might improve cognition but not mood.

Methods

Participants

Participants were recruited over a period of 18 months from January 2012. They were screened for eligibility by their treating physicians and psychologists during regular outpatient visits in the following 7 participating rehabilitation centers or hospital rehabilitation departments in The Netherlands: Groot Klimmendaal, Arnhem; St Maartenskliniek, Nijmegen; Adelante, Hoensbroek; Roessingh, Enschede; VieCuri, Venlo; Tolbrug, 's Hertogenbosch;

and ViaReva, Apeldoorn. Participants were eligible when they met the following inclusion criteria: they should (1) have sustained any type of clinically confirmed stroke at least 3 months earlier; (2) a score >7 on the Hospital Anxiety and Depression Scale-depression subscale (HADS-D)^{16,17}; (3) be ≥ 18 years; (4) have only mild cognitive impairments (Mini-Mental State Examination score¹⁸ >27 out of 30) and score positively on the communication-related items of the National Institutes of Health Stroke Scale¹⁹; and (5) master the Dutch language. Exclusion criteria were (1) prestroke major depression requiring psychiatric care, (2) premorbid disability as reflected in a Barthel Index score <19 (out of 20),²⁰ (3) stay in an inpatient setting, (4) severe comorbidity that might affect mood (eg, cancer), and (5) poststroke major depression requiring a start with medication.

The study design and methods were previously published¹⁶ and approved by the Medical Ethical Committee of Nijmegen (The Netherlands) and by the executive boards of all participating rehabilitation institutes. The adjusted time points of assessment were approved post hoc by the same medical ethical committee.

Procedure

After referral to the primary investigator (J.A.K.), participants were enrolled based on the inclusion and exclusion criteria. Oral and written informed consent was obtained by the primary investigator (J.A.K.). Subsequently, patients were randomized to either 4 months of augmented CBT or an equal period of CCT. Interventions were given in the same 6 rehabilitation institutions where patients were treated.

All outcome measures were collected at 4 time points. Initially, we intended to measure at baseline, immediately posttreatment, and 6 and 12 months posttreatment.¹⁵ Because of a lower than expected inclusion rate, the timing of the assessments was adjusted to baseline (t0), immediately posttreatment (t1), and 4 (t2) and 8 months (t3) posttreatment (fig 1). All assessments took place in the rehabilitation institute where the patient was treated. Baseline assessments were performed by the primary investigator (J.A.K.). From t1, all outcome assessments were performed by research assistants who were not involved in the administration of the interventions and who were blinded to treatment allocation of the participants. When patients mentioned the content of their intervention to the assessor, this was reported, and the subsequent assessment was performed by a different, still blinded, assessor.

Randomization

Stratified block randomization (block size 4) was performed by a randomization program for each participating rehabilitation institution separately. Main factors that were expected to affect outcomes were selected for minimization (ie, rehabilitation institute, patients' anxiety level [Hospital Anxiety and Depression Scale-anxiety subscale score ≤ 7 vs >7]). As a result, for each participating institute, patients with high anxiety scores and those with low anxiety scores were equally allocated to either the experimental or nonexperimental group.

Interventions

Both interventions were administered during a 4-month time period, with a minimum of 13 and a maximum of 16 sessions. Each session consisted of two 20- to 25-minute blocks divided by a 10- to 15-minute break. Therefore, each session lasted

List of abbreviations:

CBT	Cognitive behavioral therapy
CCT	Computerized cognitive training
CI	Confidence interval
HADS-D	Hospital Anxiety and Depression Scale-depression subscale

Fig 1 Flowchart of the Restore4Stroke Trial for depressive symptoms after stroke (Consolidated Standards of Reporting Trials diagram). t0, baseline; t1, immediately posttreatment; t2, 4 months posttreatment; t3, 8 months posttreatment.

approximately 1 hour. The CBT intervention was administered by a certified health care psychologist (therapist) who had ample experience in treating depression and stroke rehabilitation in general. All therapists were additionally trained by the primary investigator (J.A.K.) to master the specific aspects of the current CBT intervention. Goals for attaining daily life activities were primarily set together by the patient and the therapist using pictures from the Activity Card Sort.²¹ Concurrently with the psychological sessions, the CBT intervention was augmented with 3 sessions of occupational therapy or movement therapy. During these sessions, an occupational or movement therapist helped patients with establishing and attaining goals aimed at meaningful activities and social participation. These goals were attuned to the content of the psychological sessions. In the case of a baseline Hospital Anxiety and Depression Scale-anxiety subscale score >7, the protocol was extended with an additional (fourth) occupational therapy/movement therapy session aimed at relaxation techniques, as provided by the Dutch Heart Foundation.²² To implement the recommendations made by Broomfield et al,¹⁰ therapists were continuously reminded to use concrete, repetitive, accessible, slow, and structured communication strategies, as proposed by Judd.²³ A detailed description of the applied CBT intervention has been published elsewhere.¹⁴

We selected CCT as the comparative intervention to control for nonspecific (eg, Hawthorne) effects. In a previous study, this type of training yielded high satisfaction scores.²⁴ In the current study, CCT was largely self-administered, but either cognitive trainers or psychological assistants were present to assist the participants during the training. They were instructed not to engage in any conversation with the patients about topics other than the cognitive training. A desktop was setup with headphones and a keyboard with colored patches attached to 2 keys. Patients could select any (or a combination) of 4 specific cognitive domains for training (ie, attention, memory, executive functioning, visual attention). As patients improved, the Cogniplus^a program adjusted the level of difficulty for each training task accordingly. In this way, each patient trained at his/her individual level and pace.

Cointerventions

During the study, patients were requested to minimize cointerventions by other therapists and to refrain from starting new medication. Any existing (psycho)pharmacologic treatment was continued. All cointerventions were registered in a booklet at each assessment.

Assessments

Baseline assessments consisted of the following characteristics: age (years), sex (male or female), employment (yes or no), time since stroke (months), stroke type (ischemic, hemorrhagic, or other), affected hemisphere (left, right, or other), mobility (Stroke Impact Scale mobility subscale²⁵), comorbidity (Cumulative Illness Rating Scale²⁶), level of independence (Barthel Index²⁰), and cognition (Mini-Mental State Examination¹⁸). Except for comorbidity, higher scores reflect better outcomes on the scales mentioned.

The primary outcome was the severity of depressive symptoms as assessed with the HADS-D. The Hospital Anxiety and Depression Scale has been specifically validated to assess depressive symptoms in patients with stroke.¹⁷ Secondary outcomes were the Hospital Anxiety and Depression Scale-anxiety subscale for symptoms of anxiety, the Post Stroke Depression Rating Scale for assessing the more qualitative aspects of mood,²⁷ coping assessed with the Utrecht Proactive Coping Competence scale,²⁸ quality of life assessed with the Stroke Specific Quality of Life scale,²⁹ social participation assessed with the Utrecht Scale for Evaluation of Rehabilitation-Participation,³⁰ and subjective well-being assessed with a life satisfaction questionnaire (Life Satisfaction questions).³¹ Except for the Hospital Anxiety and Depression Scale and Post Stroke Depression Rating Scale scores, higher scores reflect better outcomes.

Statistical analyses

This study was originally powered on the HADS-D at 4 months posttreatment (ie, t2), without accounting for improved precision

because of repeated measures (at t1 and t3). To obtain a power of 80% with an α level of 5%, we originally aimed to include a total of 106 participants.¹³ In this calculation, adjustments for baseline values (ie, t0) and posttreatment values (ie, at t2) and a dropout rate of 15% were taken into account. Because of the lower than expected inclusion rate, we investigated whether accounting for the additionally repeated measures at t1 and/or t3 could provide sufficient power. Using the method described by de Hoop,³² we calculated that including an extra posttreatment value (ie, t1) would provide sufficient power with half the originally planned subjects. This was discussed with and approved by the medical-ethical committee. As a consequence, a minimal number of 53 participants was needed.

Median and ranges or numbers and percentages were used for descriptive statistics in the case of continuous and categorical variables, respectively. After confirming normative distribution of the dependent continuous variables, we used linear mixed models for repeated measures to study the differences between groups for each outcome (dependent variable). The independent fixed variables were group (CBT vs CCT), baseline score, time point of measurement (t0–t3), the minimization factor Hospital Anxiety and Depression Scale-anxiety subscale score >7, and the interaction term between point of measurement and group. The participating institute was treated as a random variable. We present the baseline-adjusted mean differences between groups at each time point with 95% confidence intervals (CIs). The effects of the intervention on the primary and secondary outcome measures were analyzed according to the intention-to-treat principle using SAS 9.2 for Windows^b and IBM SPSS Statistics 20 for Windows.^c Data analysis was performed by an independent statistician who was blinded to group allocation.

Results

Participants

Figure 1 presents the patient flow throughout the study. Trial inclusion took place from January 2011 until August 2012, and ended according to prior agreements with the medical ethical committee. Out of 163 referred patients, 61 ultimately participated. Of these, 31 patients were assigned to the CBT intervention and 30 patients were assigned to the CCT intervention. Incomplete interventions occurred once in the CBT group (change in protocol by psychologist) and 4 times in the CCT group (related to suicidal thoughts: $n=1$; dissatisfaction with intervention: $n=2$; no reason/not specified: $n=1$). Fifty-two patients completed the posttreatment assessment, of whom 44 completed the last follow-up. Patient characteristics are listed in table 1.

Control of bias

More than 95% of all assessments were performed according to the protocol.¹⁵ Some exceptions were as follows: extra pauses (t0, $n=1$), missing page in assessment booklet (t1, $n=1$), Utrecht Proactive Coping Competence Scale questionnaire too difficult to understand (t1, $n=1$), and Utrecht Proactive Coping Competence Scale questionnaire and Life Satisfaction questions filled in at home because of time constraints (t1, $n=1$). Assessor unblinding by the patient occurred in 13% of the t1 assessments, 10% of the t2 assessments, and 9% of the t3 assessments. In the CBT group, 1

Table 1 Patient characteristics at baseline

Characteristic	CBT Patients		CCT Patients	
	n	Median (range) or n (%)	n	Median (range) or n (%)
Age (y)	31	61 (45–79)	30	61 (25–76)
TSS (mo)		26 (2–243)		21.5 (2–138)
Men	31	19 (61.3)	30	19 (63.3)
Employed	31	9 (29)	30	13 (43.3)
Stroke type	28		28	
Infarct		21 (75)		24 (85.7)
Hemorrhage/other		7 (25)		2 (7.1)
Unknown		0		2 (7.1)
Stroke hemisphere	26		28	
Left		15 (57.7)		9 (32.1)
Right		8 (30.8)		11 (39.3)
Other		3 (11.5)		8 (28.6)
SIS mobility	31	65 (13–98)	30	65 (25–100)
CIRS	31	3 (0–18)	30	5 (0–13)
Barthel Index	31	20 (17–20)	30	20 (12–20)
MMSE	31	30 (27–30)	30	30 (24–30)
HADS	31		30	
Depression		12 (8–20)		12 (8–20)
Anxiety		9 (4–17)		10 (1–18)

Abbreviations: CIRS, Cumulative Illness Rating Scale (range, 0–52); HADS, Hospital Anxiety and Depression Scale (subscale range, 0–21); MMSE, Mini-Mental State Examination (range, 0–30); SIS, Stroke Impact Scale (range, 10–50); TSS, time since stroke.

patient had sought contact with a psychologist outside the trial for additional anxiety therapy (t1) and 1 patient had started antidepressants (t2). In the CCT group, 1 patient commenced with antidepressants at t1.

Interaction effects and group differences

Table 2 presents the primary and secondary outcomes by point of measurement for each group. Changes in the HADS-D were never significant between groups (fig 2). No group differences after treatment were found for any of the primary or secondary outcomes. Therefore, a parallel line model is presented for all variables in table 3.

Time effects

Table 3 depicts the estimated mean differences between points of measurement, adjusted for baseline values, for all outcomes, and for both groups together. A posttreatment effect was found for the primary outcome, HADS-D (mean difference, -4.6 ; 95% CI, -5.7 to -3.6). The overall time effect of the HADS-D from posttreatment to 8-month follow-up was not significant. However, the HADS-D showed a significant increase from posttreatment to 4-month follow-up (mean difference, 1.1 ; 95% CI, 0.2 – 1.9), followed by a significant decline from 4-month to 8-month follow-up (mean difference, -1.4 ; 95% CI, -2.1 to -0.7) for both groups.

A posttreatment effect was also observed for the Hospital Anxiety and Depression Scale-anxiety subscale (mean difference, -2.6 ; 95% CI, -3.6 to -1.5), Post Stroke Depression Rating Scale (mean difference, -3.4 ; 95% CI, -4.9 to -2.0), Utrecht

Table 2 Observed median (range) of the primary and secondary outcomes by point of measurement by CBT and CCT groups

Outcome	Baseline		Posttreatment		4-mo Follow-Up		8-mo Follow-Up	
	n	Median (range)	n	Median (range)	n	Median (range)	n	Median (range)
HADS-D								
CBT	31	12.0 (8.0–20.0)	24	8.00 (0.0–15.0)	24	9.0 (0.0–19.0)	23	8.0 (0.0–19.0)
CCT	30	12.0 (8.0–20.0)	28	9.00 (1.0–14.0)	24	8.5 (0.0–16.0)	20	7.5 (0.0–12.0)
HADS-A								
CBT	31	9.0 (4.0–17.0)	24	6.5 (2.0–18.0)	24	6.0 (0.0–17.0)	23	5.0 (0.0–17.0)
CCT	30	10.0 (1.0–18.0)	28	7.5 (0.0–16)	24	7.0 (2.0–15.0)	20	6.0 (0.0–13.0)
PSDRS								
CBT	30	14.0 (3.0–24.0)	23	10.3 (2.0–21.0)	24	8.0 (2.0–22.3)	22	8.2 (1.0–18.0)
CCT	30	12.0 (4.0–25.3)	28	9.0 (1.0–21.0)	24	8.5 (0.0–23.3)	20	7.5 (2.0–18.0)
UPCC								
CBT	31	2.5 (1.5–3.2)	24	2.5 (1.4–3.8)	23	2.6 (1.7–3.5)	23	2.5 (1.7–3.6)
CCT	30	2.5 (1.5–3.8)	27	2.4 (1.7–4.0)	23	2.4 (1.5–3.8)	20	2.7 (1.8–4.0)
SSQoL								
CBT	31	3.2 (2.1–4.7)	24	3.8 (2.2–5.0)	24	3.5 (2.1–4.9)	24	3.7 (2.2–5.0)
CCT	30	3.1 (2.0–4.7)	28	3.5 (1.9–4.8)	24	3.3 (2.2–4.7)	20	3.7 (1.8–5.0)
USERP vocational activities								
CBT	31	12.6 (5.0–30.0)	24	14.0 (5.0–35.0)	24	11.5 (5.0–25.0)	24	12.7 (5.0–30.0)
CCT	30	14.5 (5.0–50.0)	28	13.4 (5.0–45.0)	24	13.1 (0.0–35.0)	20	14.3 (0.0–45.0)
USERP leisure activities								
CBT	31	45.0 (7.5–82.5)	24	43.8 (20.0–77.5)	24	51.3 (27.5–70.0)	24	47.5 (17.5–77.5)
CCT	30	42.5 (15.0–80.0)	28	45.0 (5.0–70.0)	24	46.3 (12.5–72.5)	20	50.0 (17.5–70.0)
USERP satisfaction								
CBT	31	56.3 (19.4–88.9)	24	66.3 (27.8–94.4)	24	68.1 (30.6–99.4)	24	63.9 (30.6–91.7)
CCT	31	51.4 (11.1–78.1)	28	62.5 (30.6–88.9)	24	62.5 (30.6–80.6)	20	63.9 (34.4–91.7)
USERP restriction								
CBT	31	72.5 (50.0–94.4)	24	75.0 (43.6–100.0)	24	77.6 (55.0–94.4)	24	73.8 (52.8–100.0)
CCT	30	71.4 (46.9–100.0)	28	75.0 (50.0–94.4)	24	73.8 (50.0–96.9)	20	81.2 (52.8–100.0)
LS2								
CBT	31	4.0 (2.0–11.0)	24	6.0 (3.0–13.0)	24	7.5 (2.0–13.0)	24	7.0 (0.0–13.0)
CCT	30	4.0 (2.0–12.0)	28	5.0 (3.0–11.0)	24	6.5 (2.0–13.0)	20	7.0 (3.0–13.0)

NOTE. Ranges are as follows: HADS-A (0–21); HADS-D (0–21); LS2 (2–13); PSDRS (0–45); SSQoL (1–5); UPCC (1–4); USERP (0–100, each subscale). Abbreviations: HADS-A, Hospital Anxiety and Depression Scale-anxiety subscale; HADS-D, Hospital Anxiety and Depression Scale-depression subscale; LS2, 2 Life Satisfaction questions; PSDRS, Post Stroke Depression Rating Scale; SSQoL, short Stroke Specific Quality of Life Scale; UPCC, Utrecht Proactive Coping Competence list; USERP, Utrecht Scale of Rehabilitation-Participation.

Scale for Evaluation of Rehabilitation-Participation satisfaction subscale (mean difference, 9.7; 95% CI, 5.4–14.1), Stroke Specific Quality of Life Scale (mean difference, 0.3; 95% CI, 0.1–0.5), and Life Satisfaction questions (mean difference, 1.3;

95% CI, 0.7–2.0), but not for the Utrecht Proactive Coping Competence Scale. However, the Utrecht Proactive Coping Competence Scale showed a significant change from posttreatment to 8-month follow-up (mean difference, 0.1; 95% CI, 0.0–0.3), as did the Post Stroke Depression Rating Scale (mean difference, –1.9; 95% CI, –3.1 to –0.7) and Life Satisfaction questions (mean difference, 0.6; 95% CI, 0.0–1.3).

Fig 2 Mean scores for CCT and CBT patient groups at all assessment time points.

Discussion

The aim of our study was to evaluate the effectiveness of individually tailored CBT for reducing depressive symptoms with or without anxiety after stroke. The results of our multicenter randomized controlled trial indicate that there was significant and persistent improvement of depressive and anxiety complaints after treatment; however, this was independent of the type of intervention. In addition, the subjective ratings of patients' participation level (Utrecht Scale for Evaluation of Rehabilitation-Participation satisfaction subscale) increased after treatment, as did quality of life (Stroke Specific Quality of Life Scale) and life satisfaction (Life Satisfaction questions).

Table 3 Estimated mean differences (95% CIs) between points of measurement for both the CBT and CCT groups, using linear mixed model for repeated measurements while adjusting for baseline values

Outcome	Time Differences				
	Posttreatment— Baseline	4-mo Follow-Up— Posttreatment	8-mo Follow-Up— 4-mo Follow-Up	8-mo Follow-Up— Posttreatment	Overall Group Difference After Treatment Period
HADS-D	-4.6 (-5.7 to -3.6)	1.1 (0.2 to 1.9)	-1.4 (-2.1 to -0.7)	-0.4 (-1.3 to 0.5)	0.5 (-1.3 to 2.3)
HADS-A	-2.6 (-3.6 to -1.5)	-0.3 (-1.0 to 0.4)	-0.4 (-1.3 to 0.5)	-0.7 (-1.6 to 0.2)	1.1 (-0.7 to 2.9)
PSDRS	-3.4 (-4.9 to -2.0)	-0.7 (-1.7 to 0.4)	-1.2 (-2.5 to 0.1)	-1.9 (-3.1 to -0.7)	0.7 (-1.6 to 3.0)
UPCC	0.0 (-0.1 to 0.1)	0.1 (-0.2 to 0.1)	0.1 (-0.0 to 0.2)	0.1 (0.0 to 0.3)	-0.1 (-0.3 to 0.1)
SSQoL	0.3 (0.1 to 0.5)	-0.1 (-0.2 to 0.0)	0.2 (0.0 to 0.4)	0.1 (-0.1 to 0.3)	-0.2 (-0.6 to 0.1)
USERP					
Vocational activities	0.3 (-1.8 to 2.4)	-1.6 (-3.7 to 0.5)	1.2 (-0.7 to 3.1)	-0.4 (-2.1 to 1.4)	0.5 (-2.5 to 3.6)
Leisure activities	0.0 (-4.0 to 4.0)	3.1 (-0.7 to 6.9)	-0.1 (-4.0 to 3.8)	3.0 (-1.4 to 7.5)	0.3 (-5.1 to 5.6)
Satisfaction	9.7 (5.4 to 14.1)	1.6 (-1.1 to 4.2)	0.4 (-2.8 to 3.5)	1.9 (-1.5 to 5.4)	-1.1 (-7.8 to 5.5)
Restriction	2.9 (-0.4 to 6.2)	-0.2 (-3.0 to 2.5)	3.0 (-0.4 to 6.4)	2.8 (-1.0 to 6.7)	0.4 (-4.4 to 5.3)
LS2	1.3 (0.7 to 2.0)	0.4 (-0.1 to 0.9)	0.2 (-0.3 to 0.8)	0.6 (0.0 to 1.3)	-0.5 (-1.6 to 0.5)

NOTE. Values are presented as mean (95% CI). The interaction between time and group never reached the level of statistical significance. Therefore, a parallel line model is presented for all variables. Ranges are as follows: HADS-A (0–21); HADS-D (0–21); LS2 (2–13); PSDRS (0–45); SSQoL (1–5); UPCC (1–5); USERP (0–100, each subscale).

Abbreviations: HADS-A, Hospital Anxiety and Depression Scale-anxiety subscale; HADS-D, Hospital Anxiety and Depression Scale-depression subscale; LS2, 2 Life Satisfaction questions; PSDRS, Post Stroke Depression Rating Scale; SSQoL, short Stroke Specific Quality of Life Scale; UPCC, Utrecht Proactive Coping Competence list; USERP, Utrecht Scale of Rehabilitation-Participation.

These results appear to be similar to those of Lincoln and Flannaghan,¹⁰ even though our CBT was of longer duration and higher intensity, was augmented with occupational or movement therapy, was provided by well-trained psychologists, and took cognitive impairments and anxiety complaints into account. Like our study, Lincoln and Flannaghan¹⁰ found no differences between intervention groups (ie, no intervention, attention placebo, CBT)³³; however, they did find a small decline (3–4 points) in median score on the Beck Depression Inventory (range, 0–63) for each group immediately after the intervention period. Notably, the relative improvements that were observed in the present study were larger (ie, 3–4 points on the HADS-D; range, 0–21), which suggests that both treatments may have caused a beneficial effect. As our study lacked a nonintervention group, it cannot be ruled out that any beneficial effect of either CBT or CCT was nonspecific or that HADS-D scores simply improved because of regression to the mean. However, the observed improvements were relatively large, whereas recent literature shows that depressive symptoms poststroke (measured with the Beck Depression Inventory) remain stable over the first 2 years.³⁴ In addition, our Restore4Stroke consortium recently reported nonsignificant fluctuations in poststroke depressive symptoms (maximally 2-point change in HADS-D) over the first 2 years.³⁵ These findings point toward the possibility that the observed decline in depressive symptoms after treatment in the present study may actually represent a beneficial effect of both CBT and CCT. This notion raises the question of what the effective component of CCT might have been because we took care that cognitive trainers and psychological assistants were instructed not to address issues other than the cognitive training. Interestingly, in a recent evaluation of CCT, Akerlund et al³⁶ showed that patients with acquired brain injury who suffered from depressive complaints demonstrated mood improvements after CCT, which suggests that cognitive training may improve mood through motivational mechanisms or perhaps through cognitive improvement. Irrespective of the underlying mechanism, it may be that such effects occurred in our study. Unfortunately, at the initiation of our study, beneficial effect results of CCT on mood problems were not yet known.

Quality of life scores and satisfaction with life and with participation equally and persistently improved in both groups after treatment, as did anxiety level. Proactive coping did not immediately respond after the intervention, but it improved during follow-up, together with life satisfaction and qualitative aspects of mood. In a recent study from our Restore4Stroke consortium,³⁷ change in depression scores was associated with subjective experience of participation, emphasizing that rehabilitation should focus on resuming occupational activities when treating depressive complaints after stroke. Although in the present study this approach was applied in the CBT intervention, it remains to be explained why we found no group differences for any of the secondary outcomes, and why CBT and CCT had similar effects.

Study limitations

Apart from the absence of a nonintervention group, the main limitation of this study is the relatively small patient sample and the recruitment through rehabilitation institutes, which limits the generalizability of our results. It is well known that recruiting patients with depressive symptoms for participation in research is notoriously difficult.³⁸ Patients who did participate may have been more motivated than those who did not, which may have led to selection bias. This too would limit the generalizability of our

results. Although we did not reach the inclusion number as originally planned, the risk of false-negative outcomes for group by time interactions seems to be small because no trends were observed toward significant interactions. We were not able to directly compare our results with those of Lincoln and Flannaghan¹⁰ because we used the HADS-D instead of the Beck Depression Inventory as a primary outcome. This choice was determined by the intention to prevent overestimation of depressive symptoms because of general symptoms (eg, fatigue) of stroke. Unlike the Beck Depression Inventory, the Hospital Anxiety and Depression Scale does not include fatigue as a symptom of depressed mood.

Conclusions

This randomized controlled trial showed persistent improvement of depressive symptoms, anxiety symptoms, quality of life, satisfaction with life, and participation after both CBT and CCT in patients with depressive complaints minimally 3 months after stroke. This finding implies that, for now, both types of interventions may be considered to improve depressive symptoms after stroke. Future research should determine whether both interventions are better than natural history and what underlying mechanisms are responsible for such effects.

Suppliers

- a. Cogniplus; SCHUHFRIED.
- b. SAS 9.2; SAS Institute.
- c. IBM SPSS Statistics 20; IBM.

Keywords

Anxiety; Depression; Psychology; Rehabilitation; Stroke

Corresponding author

Joyce A. Kootker, MSc, Reinier Postlaan 12, PO Box 9101, 6500 HB, Nijmegen, The Netherlands. *E-mail address:* joyce.kootker@radboudumc.nl.

Acknowledgments

We thank the participating centers: Groot Klimmendaal, Arnhem; St Maartenskliniek, Nijmegen; Adelante, Hoensbroek; Roessingh, Enschede; VieCuri, Venlo; Tolbrug, 's Hertogenbosch; and ViaReva, Apeldoorn. We also thank J. Hendriks, PhD, for assistance during the process of analyses and L. Vlutters, MSc, T. Vaessen, MSc, D. Mennen, MSc, and I. Kortland, MD, for their contribution to inclusion of participants.

References

1. Hackett ML, Pickles K. Part I: frequency of depression after stroke: an updated systematic review and meta-analysis of observational studies. *Int J Stroke* 2014;9:1017-25.
2. Ayerbe L, Ayis S, Crichton S, Wolfe CD, Rudd AG. The long-term outcomes of depression up to 10 years after stroke; the South London Stroke Register. *J Neurol Neurosurg Psychiatry* 2013;85:514-21.
3. Mottram P, Wilson K, Strobl J. Antidepressants for depressed elderly. *Cochrane Database Syst Rev* 2006;(1):CD003491.
4. Coupland CA, Dhiman P, Barton G, et al. A study of the safety and harms of antidepressant drugs for older people: a cohort study using a large primary care database. *Health Technol Assess* 2011;15:1-202. iii-iv.
5. Gainotti G, Marra C. Determinants and consequences of post-stroke depression. *Curr Opin Neurol* 2002;15:85-9.
6. Hackett ML, Anderson CS, House AO. Management of depression after stroke: a systematic review of pharmacological therapies. *Stroke* 2005;36:1098-103.
7. Cuijpers P. Prevention of depression in chronic medical disorders: a pilot study. *Psychol Rep* 1988;82:735-8.
8. Broomfield NM, Laidlaw K, Hickabottom E, et al. Post-stroke depression: the case for augmented, individually tailored cognitive behavioural therapy. *Clin Psychol Psychother* 2011;18:202-17.
9. Hackett ML, Anderson CS, House AO, Xia J. Interventions for treating depression after stroke. *Cochrane Database Syst Rev* 2008;(4):CD003437.
10. Lincoln NB, Flannaghan T. Cognitive behavioral psychotherapy for depression following stroke: a randomized controlled trial. *Stroke* 2003;34:111-5.
11. De Wit L, Putman K, Baert I, et al. Anxiety and depression in the first six months after stroke. A longitudinal multicentre study. *Disabil Rehabil* 2008;30:1858-66.
12. Wilkinson PR, Wolfe CD, Warburton FG, et al. A long-term follow-up of stroke patients. *Stroke* 1997;28:507-12.
13. Rasquin SM, Van De SP, Praamstra AJ, van Heugten CM. Cognitive-behavioural intervention for depression after stroke: five single case studies on effects and feasibility. *Neuropsychol Rehabil* 2009;19:208-22.
14. Kootker JA, Rasquin SM, Smits P, Geurts AC, van Heugten CM, Fasotti L. An augmented cognitive behavioural therapy for treating post-stroke depression: description of a treatment protocol. *Clin Rehabil* 2014;29:833-43.
15. Kootker JA, Fasotti L, Rasquin SM, van Heugten CM, Geurts A. The effectiveness of an augmented cognitive behavioural intervention for post-stroke depression with or without anxiety (PSDA): the Restore4Stroke-PSDA trial. *BMC Neurol* 2012;12:51.
16. Zigmond AS, Snaith RP. The hospital anxiety and depression scale. *Acta Psychiatr Scand* 1983;67:361-70.
17. Aben I, Verhey F, Lousberg R, Lodder J, Honig A. Validity of the beck depression inventory, hospital anxiety and depression scale, SCL-90, and hamilton depression rating scale as screening instruments for depression in stroke patients. *Psychosomatics* 2002;43:386-93.
18. Folstein MF, Folstein SE, McHugh PR. "Mini-mental state". A practical method for grading the cognitive state of patients for the clinician. *J Psychiatr Res* 1975;12:189-98.
19. Meyer BC, Lyden PD. The modified National Institutes of Health Stroke Scale: its time has come. *Int J Stroke* 2009;4:267-73.
20. Collin C, Wade DT, Davies S, Horne V. The Barthel ADL Index: a reliability study. *Int Disabil Stud* 1988;10:61-3.
21. Eriksson G, Chung J, Beng L, et al. Occupations of older adults: a cross cultural description. *OTJR (Thorofare N J)* 2011;31:182-92.
22. van Erp J. Relaxation exercises (ontspanningsoefeningen; bodyscan, aandacht voor uw lichaam). De hart&vaatgroep; 2010.
23. Judd T. Neuropsychotherapy and community integration; brain illness, emotions and behaviour. New York: Kluwer Academic/Plenum Publishers; 1999.
24. Spikman JM, Boelen DH, Lamberts KF, Brouwer WH, Fasotti L. Effects of a multifaceted treatment program for executive dysfunction after acquired brain injury on indications of executive functioning in daily life. *J Int Neuropsychol Soc* 2010;16:118-29.
25. Duncan PW, Wallace D, Lai SM, Johnson D, Embretson S, Laster LJ. The stroke impact scale version 2.0. Evaluation of reliability, validity, and sensitivity to change. *Stroke* 1999;30:2131-40.

26. Giaquinto S. Comorbidity in post-stroke rehabilitation. *Eur J Neurol* 2003;10:235-8.
27. Gainotti G, Azzoni A, Razzano C, Lanzillotta M, Marra C, Gasparini F. The Post-Stroke Depression Rating Scale: a test specifically devised to investigate affective disorders of stroke patients. *J Clin Exp Neuropsychol* 1997;19:340-56.
28. Bode C, Thoolen B, de Ridder D. [Measuring proactive coping. Psychometric characteristics of the Utrecht Proactive Coping Competence scale (UPCC)] [Dutch]. *Psychol Gezond* 2008;36:81-91.
29. Post MW, Boosman H, van Zandvoort MM, Passier PE, Rinkel GJ, Visser-Meily JM. Development and validation of a short version of the Stroke Specific Quality of Life Scale. *J Neurol Neurosurg Psychiatry* 2011;82:283-6.
30. Post MW, van der Zee CH, Hennink J, Schafrat CG, Visser-Meily JM, van Berlekom SB. Validity of the utrecht scale for evaluation of rehabilitation-participation. *Disabil Rehabil* 2012;34:478-85.
31. van Koppenhagen CF, Post MW, van der Woude LH, et al. Recovery of life satisfaction in persons with spinal cord injury during inpatient rehabilitation. *Am J Phys Med Rehabil* 2009;88:887-95.
32. Hoop de E. Efficient designs for cluster randomized trials with small numbers of clusters; stepped wedge and other repeated measurements designs [dissertation]. Nijmegen: Radboud University Medical Centre; 2014.
33. Lincoln NB, Flannaghan T, Sutcliffe L, Rother L. Evaluation of cognitive behavioural treatment for depression after stroke: a pilot study. *Clin Rehabil* 1997;11:114-22.
34. Eriksen S, Gay CL, Lerdal A. Acute phase factors associated with the course of depression during the first 18 months after first-ever stroke. *Disabil Rehabil* 2015;38:30-5.
35. van Mierlo ML, van Heugten CM, Post MW, Hajos TR, Kappelle LJ, Visser-Meily JM. Quality of life during the first two years post stroke: The Restore4Stroke Cohort Study. *Cerebrovasc Dis* 2016;41:19-26.
36. Akerlund E, Esbjornsson E, Sunnerhagen KS, Bjorkdahl A. Can computerized working memory training improve impaired working memory, cognition and psychological health? *Brain Inj* 2013;27:1649-57.
37. Blomer AM, van Mierlo ML, Visser-Meily JM, van Heugten CM, Post MW. Does the frequency of participation change after stroke and is this change associated with the subjective experience of participation? *Arch Phys Med Rehabil* 2015;96:456-63.
38. Hughes-Morley A, Young B, Waheed W, Small N, Bower P. Factors affecting recruitment into depression trials: systematic review, meta-synthesis and conceptual framework. *J Affect Disord* 2015;172:274-90.