New observations on the role of the A-V junction in tachycardias in man

Citation for published version (APA):

Brugada i Terradellas, P. (1982). *New observations on the role of the A-V junction in tachycardias in man*. [Doctoral Thesis, Maastricht University]. Rijksuniversiteit Limburg.
<https://doi.org/10.26481/dis.19821104pb>

Document status and date:

Published: 01/01/1982

DOI:

[10.26481/dis.19821104pb](https://doi.org/10.26481/dis.19821104pb)

Document Version:

Publisher's PDF, also known as Version of record

Please check the document version of this publication:

- A submitted manuscript is the version of the article upon submission and before peer-review. There can be important differences between the submitted version and the official published version of record. People interested in the research are advised to contact the author for the final version of the publication, or visit the DOI to the publisher's website.
- The final author version and the galley proof are versions of the publication after peer review.
- The final published version features the final layout of the paper including the volume, issue and page numbers.

[Link to publication](#)

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal.

If the publication is distributed under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license above, please follow below link for the End User Agreement:

www.umlib.nl/taverne-license

Take down policy

If you believe that this document breaches copyright please contact us at:

repository@maastrichtuniversity.nl

providing details and we will investigate your claim.

New observations on the role of the A-V junction in tachycardias in man

Proefschrift

Ter verkrijging van de graad van doctor in de geneeskunde aan de Rijksuniversiteit Limburg te Maastricht, op gezag van de Rector Magnificus Prof.Dr. H.C. Hemker, volgens besluit van het College van Dekanen in het openbaar te verdedigen in de aula van de universiteit op donderdag, 4 november 1982 des namiddags te vier uur.

door

Pere Brugada i Terradellas geboren in Banyoles, Catalunya (Spanje),
11 augustus 1952.

Promotor: Hein J.J. Wellens, M.D.

Referenten: Maurits Allessie, M.D., The Netherlands
Jerónimo Farré, M.D., Madrid, Spain
Mark E. Josephson, M.D., Philadelphia, U.S.A.

A la Mercé i a l'Isabel

A en Ramón i a la Pepita.

A Banyoles i a l'Estany.

*Als desmaïs, al Drac, i a l'aigüa
de La Puda que tot ho cura.*

Al pa amb tomate i pernil,

A un passeig per les Rambles

i a una Sardana una tarda d'estiu.

A una berenada a la muntanya,

i a un descans vora del mar.

Al soroll de les onades,

i a les barques dels pescadors.

A la frescor sota un pí en el despenyat,

i al sentiment del sol sobre el cos.

A Catalunya!

Contents

Foreword

1.	Introduction	1
2.	The method of programmed electrical stimulation of the heart used in Maastricht. With emphasis on the approach in patients with supraventricular tachycardia.	3
3.	Atrio-ventricular nodal tachycardia with and without discontinuous anterograde and retrograde atrio-ventricular nodal conduction curves: a reappraisal of the dual pathway concept.	31
4.	Observations on spontaneous termination of atrio-ventricular nodal re-entrant tachycardia.	41
5.	Initiation of atrio-ventricular nodal re-entrant tachycardia in patients with discontinuous anterograde atrio-ventricular nodal conduction curves with and without documented supraventricular tachycardia. Observations on the role of a discontinuous retrograde conduction curve.	47
6.	Dual atrio-ventricular nodal pathways and atrial fibrillation.	67
7.	Ventricular tachycardia with concealed retrograde conduction to two atrio-ventricular nodal pathways.	75
8.	Incessant reciprocating atrio-ventricular tachycardia. Factors playing a role in the mechanisms of the arrhythmia.	79
9.	Observations in patients showing A-V junctional echoes with a shorter P-R than R-P interval. Distinction between intranodal re-entry or re-entry using an accessory pathway with long conduction time.	87
10.	Observations in patients showing a supraventricular tachycardia with a shorter P-R than R-P interval. Distinction between atrial tachycardia or tachycardia using an accessory pathway with long conduction time.	99
11.	Summary	119

Foreword

During the fall of 1979 my family and I arrived in Maastricht. At that time Prof. H.J.J. Wellens gave me the opportunity to work as a granted Fellow in his laboratory of clinical electrophysiology. The problem was that I had no grant!

Survival would have been difficult without the support of this beautiful, tiny little girl Mercé to whom I am married. She agreed to stop her work as a Rheumatologist in our lovely Catalan country and abandoned any idea about a quiet, easy and comfortable position in the medical field in Barcelona.

Mercé is a genius. She started on the road of scientific investigation very early in life. As a child she questioned and replied to everybody. She even questioned her mother, when she told her the fairytale of Cinderella.

As you will remember Cinderella was a poor orphan girl abused by her stepmother and stepsisters. She worked all day in the ashes. She would have loved to go to the fancy ball which the Prince offered in the Palace to choose his wife. But this was prevented by the stepmother and stepsisters. Cinderella's fairy godmother, however, provided her with the necessary dress and carriage to get her to the Palace. She wore fancy cristal slippers. The condition was that at midnight the magic would stop and everything would be back in its original form. While dancing in the garden of the Palace the clock struck twelve, and Cinderella ran out of the Prince's arms. The magic was broken and everything regained its ashy look, except for the crystal shoe that Cinderella lost during her desperate escape. The Prince tried to find Cinderella, searching all over the kingdom.

Lovely, but..... at the end of the story Mercé will always ask the same question: "Why did the crystal shoe not turn back into a rough hemp sandal at midnight?" That is the principle of scientific investigation! The identification of the unusual and unexplained, the small detail that escapes the keen look of all but one!

I do not know what would have happened to us without Hein Wellens. I did not know that he by accepting me in Maastricht committed himself to the responsibility of taking care of my entire family. It did not take too long to realize this. He taught me electrophysiology. But I also found him to be an excellent teacher of clinical cardiology, the art of presenting in public but also English and Dutch! The best thing that I continuously learn from him, however, is to have an open and interested mind for all details, such as the crystal shoe of Cinderella. I realize that it was a tremendous privilege that Hein Wellens gave me the opportunity to stay in Maastricht.

To all dreamers that listen to stories and ask why the shoe of Cinderella did not turn back into a rough hemp sandal at midnight I would like to offer this piece of work which I did in Maastricht in the past years inspired and guided by Hein Wellens. Drs Maurits Allessie, Mark Josephson and Jerónimo Farré not only showed me how to perform scientific work but also kindly consented to review and discuss this thesis. Miss Moniek Vermeulen expertly typed the manuscript.

I expect that this thesis will be the first step on my road of research. I do not know yet why the shoe of Cinderella did not turn back into a rough hemp sandal, but hopefully we will find out!

1.

Introduction

Programmed electrical stimulation of the heart introduced in 1967 into clinical cardiology proved to be a useful technique for identification of the mechanism, the site of origin and the pathways involved in arrhythmias. It also showed to be valuable in the selection of appropriate therapy and improved our diagnostic ability using the 12-lead electrocardiogram.

The purpose of this thesis is to concentrate on findings obtained during programmed electrical stimulation in patients suffering from supraventricular tachycardia.

In chapter 2 the method of programmed electrical stimulation of the heart is described, and the ways to diagnose pathways used during the tachycardia are outlined.

The complexity of intra A-V nodal tachycardia is illustrated in chapters 3 to 5. Chapter 3 discusses modes of initiation of A-V nodal tachycardia. In chapter 4 unusual modes of termination of that arrhythmia are presented. It illustrates the interaction between different re-entrant circuits in the human heart, and also gives evidence for the presence of several A-V nodal pathways in the human A-V node. Chapter 5 describes the importance of the relation between the electrophysiological properties of the different A-V nodal pathways for initiation of A-V nodal tachycardia. Knowledge about the properties of these pathways is helpful to identify patients able to sustain AV nodal tachycardia. Chapter 6 describes the determinants of the ventricular response during atrial fibrillation in patients with dual A-V nodal pathways. It shows that the presence of dual A-V nodal pathways does not result in a different pattern of atrio-ventricular conduction when compared to the ventricular response during atrial fibrillation in patients without dual A-V nodal pathway curves. Chapter 7 illustrates how the presence of longitudinal dissociation in the A-V node can result in complex electrocardiographic patterns during ventricular tachycardia. The syndrome of incessant supraventricular tachycardia is discussed in chapters 8 to 10. The incessant tachycardias occurring in the setting of the Wolff-Parkinson-White syndrome or in the presence of a concealed "fast" accessory A-V pathway are not considered. The factors playing a role in the mechanisms of atrio-ventricular re-entrant incessant tachycardia are discussed in chapter 8. The distinction by means of programmed electrical stimulation of the heart between intranodal re-entry of the uncommon variety and re-entry using a slowly conducting accessory A-V pathway is given in chapter 9, and between atrial tachycardia and slowly conducting accessory A-V pathway in chapter 10. The thesis ends with a summary in the English, Dutch, Spanish and Catalan language.

2.

The method of programmed electrical stimulation of the heart used in Maastricht.

With emphasis on the approach in patients with supraventricular tachycardia

The method of programmed electrical stimulation of the heart (PES) presently used in Maastricht is an extension of the one introduced in 1967 by Durrer et al. (1). As described by Wellens (2,3) programmed stimulation of the heart has shown to be effective in studying the mechanisms, pathways and sites of origin of tachycardias. The importance of this type of investigation for treatment of patients with tachycardia has been extensively discussed by Wellens (2,3). In recent years our group made some changes in the stimulation protocol which increase the amount of information which can be obtained during the study. In this chapter we would like to review value and limitations of our present modes of stimulation with special emphasis on the study of mechanisms of supraventricular arrhythmias.

General

Patients are studied in the postabsorptive non-sedated state using local anaesthesia with 2% Lidocaine. Children below five years of age are studied under general anaesthesia which is maintained as light as possible once the catheters have been introduced. Unless the antiarrhythmic effects of drugs are being evaluated, patients are studied without medication.

In the absence of any contraindication full heparinization (100 IU/kg body weight) is used during the procedure. Electrode catheters are used to record intracavitary electrograms and to pace the heart at different locations. Catheters are passed through the femoral veins to the right atrium, right ventricle and to the tricuspid valve area to record activation of the bundle of His. Catheterization of the left atrium via a patent foramen ovale or of the coronary sinus is also done by way of the femoral approach. Catheterization of the coronary sinus is attempted even when a patent foramen ovale is present (fig. 1). Using the femoral route we are presently able to catheterize the coronary sinus in 95% of our patients. If during catheterization from the femoral vein the coronary vein cannot be entered the left cephalic vein is used. In this way the coronary sinus can be catheterized in 98% of patients where this is considered indicated. When a patent foramen ovale is crossed, direct recordings from the left atrium are made in addition to those from the coronary sinus. This results in the use of a total number of four to five venous catheters during PES, with most patients having four catheters. The minimal number of catheters used is three: right atrium, right ventricle and His bundle. Six has been the maximal number of catheters that we have used in a single patient: one or two quadripolar catheters in the right atrium, one or two bipolar or one bipolar and one quadripolar in the His bundle area, one quadripolar in the left atrium, one quadripolar in the coronary sinus, one quadripolar in the right ventricle, and one quadripolar in the left ventricle through a foramen ovale. In addition, retrograde arterial catheterization of the left ventricle using a bi or preferably quadripolar catheter is used when indicated (fig. 1). Catheters are moved to other locations during the procedure when required.

During the study 12-lead electrocardiograms are obtained during sinus rhythm and tachycardia. Leads I, II, III, V1 and V6 are recorded continuously simultaneous with 6 bipolar recordings from the different intracardiac positions described. It is also possible to record 6 surface leads with 5 endocavitary bipolar electrograms (fig. 2). A switch-box allows immediate changes in the electrodes used for stimulation or recording and the recorded surface leads.

A sixteen channel Siemens-Eléma Mingograph is used for real-time continuous recording and display and two 8 channel oscilloscopes for real-time monitoring (one at the patient's side and the other at the recording table). The remaining five channels are used for the coding system shown at the bottom of figure 2, to display 20 ms interval time-marks, to indicate the stimulus artifact, and

Figure 1
Catheter positions during study of supraventricular (top figure) and ventricular (bottom figure) tachycardia. Catheters are located in the right atrium (RA), left atrium (LA), coronary sinus (CS), His bundle area (His), right ventricular apex (RV) and left ventricle via a foramen ovale (LVfo) and retrogradely via the femoral artery (LV). V1 to V3: surface electrodes of leads V1 to V3.

Figure 2

A tracing as registered by a 16 channel recorder is shown. 6 Surface leads are recorded simultaneously with bipolar electrograms from the right atrium (RA), coronary sinus (CS), His bundle, right ventricle (RV) and left ventricle (LV). A one second time mark (mechanical) occurs in the top part of the tracing. Also the stimulus artifact, a 20 ms time-mark and data on pacing program, time and patient number are displayed.

the recording of arterial pressure. An additional one second mechanical time-mark is displayed in the top line of the registration. Intracardiac electrograms are preamplified using Janssen JSI 0387 preamplifiers directly connected to the catheters, and then amplified and filtered by Hewlett-Packard 8811A amplifiers. All data are continuously stored on tape (14 channel, Ampex PR2200). Patterns of A-V and V-A conduction and duration of refractory periods are determined using the extrastimulus technique during sinus rhythm, paced supraventricular and paced ventricular rhythms and tachycardia(s). Several (at least three) basic cycle lengths of pacing are used during atrial and ventricular stimulation. When needed a maximum of 4 extrastimuli are delivered during a spontaneous or a paced rhythm. Progressive incremental pacing is done from atrium and ventricle. We rarely perform pacing of the ventricles at rates faster than 300 beats/min, but we do not have a rate limit during atrial stimulation particularly when we try to induce atrial flutter or fibrillation. Several atrial sites (minimally right atrium and one site in the coronary sinus or left atrium) are paced. Pacing at different right or left ventricular sites is only used when insufficient

information is obtained during right ventricular apex stimulation or because of particular research protocols. Stimuli are given at twice diastolic threshold and have a 2 ms duration, except when the effect of changes in current or duration of the stimulus is studied.

All equipment is grounded and leakage current measured at regular intervals. Informed consent is obtained from each patient. The procedure is explained to the patient in the presence of the catheterizer, the electrophysiologist and the in-hospital attending physician.

Catheters used:

For pacing and recording from atrium, coronary sinus and ventricle bipolar (4 or 5F) or quadripolar (6F) electrodes are used. Quadripolar catheters allow pacing and recording from the same catheter. We do not use tripolar catheters to record electrograms from the His bundle area. They are helpful and probably necessary during special procedures like electrical ablation of the A-V node-His bundle, as recently described by Gallagher et al. (4). We have obtained stable good quality bipolar electrograms of the His bundle using a regular bipolar 1 cm interelectrode distance 4F catheter in most patients in whom it was attempted. This usually requires some rotation of the catheter to obtain a good contact of the tip against the His bundle.

A quadripolar right ventricular catheter allows pacing and recording from the ventricle. Ventricular signals are of value to study timing and pattern of ventricular activation during sinus rhythm, pacing or tachycardia, and invariably used in patients with ventricular arrhythmias. Mapping of endocardial activation is performed at our laboratory using a bi or quadripolar catheter and fluoroscopic control including angulated views (fig. 3). Gallagher et al. have designed a specially modified Brockenbrough catheter to map the right atrium (5). We have no experience with this catheter.

We recently had the opportunity to evaluate the quadripolar catheter designed by Mark E. Josephson (USCI). This catheter is a 6F quadripolar electrode catheter with an interelectrode distance of half a centimeter. The distal electrode is arbitrarily named 1 and the most proximal 4. The catheter is suitable for recording of uni or bipolar electrograms in any combination and for pacing. Because of the small interelectrode distance we have found this catheter particularly useful in two situations: recording of a proximal and distal (electrodes 3,4 and electrodes 1,2) bipolar electrograms of the His bundle, and recording of two one centimeter distance bipolar electrograms (electrodes 1,3 and electrodes 2,4) of the His bundle or at any other location in the heart.

Figure 4 shows the good quality of the recordings obtained with this catheter and how unusual forms of conduction disturbances within the His bundle can easily be identified. The two very close bipolar recordings are helpful in mapping procedures. Mapping is facilitated by the special connector and elongator of the catheter which avoids the limitations of a too short catheter. This also allows manipulation without losing torque control.

Using full heparinization the incidence of clinical venous thrombosis is low following PES. We have observed 3 cases in a consecutive series of 500 patients. Subclinical thrombosis seems to be high following any catheterization procedure, however. Heparin-coated catheters might be of help to reduce this incidence. They are not yet available for electrophysiologic investigations however.

The pulse generator

We are presently using a Janssen stimulator (Janssen Pharmaceutica, Belgium). The modular basis of this stimulator allows enlargement of the unit to any desired combination of stimuli and sensing during a driven rhythm or a spontaneous rhythm.

Apart from modes of stimulation universally applied during programmed stimulation of the heart (2,3), this stimulator allows some special modes of stimulation like:

Figure 3
 Example of atrial mapping during right atrial tachycardia using Josephson's catheter. 5 Surface leads are shown simultaneously with bipolar endocavitary electrograms of the right and left atrium at different locations, His bundle and coronary sinus (CS). Note that site 18 (low anterior right atrium) is the earliest site of endocardial atrial activation.

Josephson's catheter

Figure 4

Use of Josephson's catheter to record two close bipolar, 0.5 cm interelectrode distance, electrograms of the His bundle in a patient with intra-Hisian block. 5 Surface leads are recorded simultaneously with a bipolar electrogram of the high RA, and a proximal and distal His bundle bipolar electrogram. During sinus rhythm the Hp-Hd interval measures 10 ms, and prolongs to 80 ms after an atrial premature given after 330 ms (A1).

1. Pacing simultaneously at the supraventricular and ventricular level with the same basic rhythm followed by 1 to 4 stimuli at the desired (usually premature) interval. These premature beats can be given at one or both levels at any desired premature beat interval combination. An example of this mode of pacing is shown in fig. 5. We found it useful in studying the pattern of V-A conduction in patients with poor retrograde conduction during the basic driven rhythm. It is also helpful in the study of the mode of initiation of ventricular arrhythmias during ventricular stimulation in patients without V-A conduction during ventricular pacing by avoiding the occurrence of ventricular captures from sinus rhythm (fig. 6). It also allows analysis of the mode of initiation of A-V junctional tachycardia in patients having tachycardia initiated by any ventricular paced beat which is retrogradely conducted to the atria.

2. A continued paced rhythm in the atrium after a pre-set V-A interval. This mode of stimulation simulates a re-entrant tachycardia with a fixed V-A interval, as in the Wolff-Parkinson-White syndrome. Premature stimuli are delivered at the atrial or ventricular level during this simulated tachycardia and their effects studied. This mode of stimulation has been of interest to understand some mechanisms of re-entrant tachycardias like alternation of the tachycardia cycle due to alternation in anterograde A-V nodal conduction time (6-7).

3. Delivery of a train of stimuli at a pre-set rate during spontaneous rhythm with a pre-set coupling interval of the first paced beat. One of the most important drawbacks of the majority of commercially available stimulators is their inability to deliver the first paced beat of a train of stimuli at a choosable interval. Figures 7 and 8 show how depending upon the coupling interval of the first paced beat S3 is refractory or a 1-to-1 or a 2-to-1 capture of the ventricle occurs. The ability to set the interval of the first paced beat allows study of the effects of pacing under reproducible conditions.

4. Pacing of one cardiac chamber at a fixed or variable rate during an exactly determined number of stimuli. Most stimulators will pace the heart for eight beats followed by a certain number of extrastimuli (usually two or three). When accurate effects of a long train of paced beats are being studied these stimulators will pace the heart at a fixed or variable rate until the stimulation is switched off. The number of stimuli delivered will vary from stimulation to stimulation. We can select an exact number of stimuli with pacing rates that can be varied in steps of 1 beat/min. This has been of interest to study some electrophysiological mechanisms of re-entrant tachycardias like entrainment.

Figure 6
Use of simultaneous atrial and ventricular pacing to avoid ventricular captures by supraventricular impulses during ventricular pacing in patients with poor V-A conduction. In the top panel during pacing of the right ventricle with a basic cycle length of 600 ms captures occurred from conducted sinus impulses (arrow) and from reversed A-V nodal echoes. In the bottom panel simultaneous atrial and ventricular pacing prevents supraventricular captures of the ventricle. 5 Surface leads are shown simultaneously with two bipolar endocavitary electrograms.

Figure 7
Importance of the coupling interval of the first beat of a train of paced stimuli. All paced stimuli activate the ventricle in panel B. In panel A the first stimulus falls too early in the cycle resulting in ventricular activation by only 7 paced basic stimuli. Now S3 given after the same interval as in panel B finds the ventricle refractory. 6 Surface leads are shown.

Figure 8
In panel A 6 beats delivered during sinus rhythm with a CL of 240 ms result in 2-to-1 ventricular activation. In panel B initiation of a tachycardia using anterogradely a nodo-ventricular Mahaim fiber occurs after 1-to-1 ventricular capture. Although the same pacing cycle length is used 1-to-1 ventricular activation occurs because the first paced beat falls earlier in the sinus cycle.

5. Accurate selection of the amount of current delivered can only be accomplished when the selector of current intensity allows digital steps of at least 0.1 mA. This is particularly true when effects of drugs on the excitability properties of cell membranes (strength-duration and strength-interval curves) are being evaluated.

Future aims with the pulse generator

We believe that some modifications of the stimulator are of value. Firstly, an increase in the number of outputs. Like most stimulators ours has two outputs. Simultaneous pacing at several sites in the heart can be of interest to terminate arrhythmias. Simultaneous A-V pacing terminates A-V junctional re-entrant tachycardia and prevents re-initiation. We have not been able so far to terminate atrial flutter by delivering a single atrial premature beat at a single or simultaneously at two atrial sites. Delivery of a single premature beat simultaneously at three or four atrial sites might however prove to be useful in terminating the arrhythmia. The same holds for ventricular tachycardia. In those cases in which a single or multiple ventricular premature beats are ineffective to terminate ventricular tachycardia simultaneous depolarization of the ventricles at two or more sites by a single premature beat might prove helpful in terminating the arrhythmia.

Secondly, changes in the mode of current given. Our stimulator gives the desired intensity of current with the selected strength and duration of the impulse (both mA and ms duration are selected). Selection of mA is superior to selection of mV (voltage difference between electrodes) because in the last situation the intensity of current given is unknown. Intensity varies in relation to the resistance from the catheter and the tissues receiving the current. The ability to select either voltage difference or current intensity allows measurements of tissue and catheter resistance and might be desirable. For example when thresholds of stimulation are being compared between normal and abnormal areas of the heart. Accurate measurement of membrane properties by catheter techniques is questionable. Factors that cannot be easily controlled, like changes in catheter position during diastole and systole also play a role in the amount of current required to depolarize the pacing site. Control of current intensity in steps of 0.01 mA might be of interest. When given in the ventricles currents higher than 20 mA seem to be associated with an increased risk of ventricular fibrillation but they can be delivered to the atria without trouble and without discomfort to the patient if the catheter position is carefully checked. When the atrial refractory period approaches the cycle length of the tachycardia as in atrial flutter or reentrant tachycardia using an accessory pathway, atrial capture can only be obtained by delivering high intensity current. The possibility of delivering current intensities as high as 100 mA might be desirable in those situations. This can result in atrial capture when lower current intensities fail. Similarly, during pace-mapping for ventricular tachycardia current intensities of 35-50 mA are sometimes required to obtain capture in abnormal areas within an aneurysm. The benefit of increased intensity current in initiating ventricular tachycardia has recently been demonstrated (8).

There is no doubt that stimulation protocols are becoming more and more aggressive. Most patients with chronic recurrent ventricular tachycardia will require at least two extrastimuli during ventricular pacing or tachycardia to initiate or terminate the arrhythmia (9). Any stimulator should be able to deliver four extrastimuli. This can be done with our present stimulator. When research protocols are considered however, we would like to be able to vary the interval of each paced beat.

Defibrillation:

In our laboratory the incidence of defibrillation in patients with severe life-threatening ventricular arrhythmias has been 19% in a series of 105 patients. In 75% of ventricular tachycardias-ventricular flutter leading to severe hemodynamic problems we are able to terminate the arrhythmia by pacing within 15 seconds. Obviously one would like to terminate all these arrhythmias and earlier. Therefore delivery of extrastimuli or overdrive pacing should be instantaneous when required. This means that a separate pacing module should be available with pre-set premature beat intervals or overdrive rhythm. Part of our success rate in pacing control of severe ventricular arrhythmias is based upon the presence of a technician who only takes care of the stimulator while the physician is in exclusive control of the real-time recordings. Unfortunately the number of

people participating in a study is frequently less in most institutions. When defibrillation is required because of ventricular fibrillation the current given in the first shock is 200 w/s. Two out of 20 patients did not respond to the first shock with this amount of current but reverted to sinus rhythm using the same energy at a second attempt.

Occasionally, usually following intravenous administration of an antiarrhythmic drug, patients develop a ventricular tachycardia which does not result in serious hemodynamic depression to make the patient lose consciousness, but cannot be terminated by pacing. In this situation cardioversion may be required to terminate the arrhythmia. Our approach is to first sedate the patient with a rapid-acting short-lasting anaesthetic like etomidate. Of this drug 0.2 to 0.3 mg/kg body weight is given by intravenous route and followed by a synchronized countershock of 10-20 w/s. The action of etomidate lasts 4-5 min. The study can then be continued with the patient conscious and without the longer lasting effects of drugs like diazepam. When required, atrial fibrillation is converted to sinus rhythm using the same approach by delivering an initial shock of 25 to 100 w/s.

Acute intravenous testing of antiarrhythmic drugs:

Administration of antiarrhythmic drugs in patients undergoing electrophysiological investigations allows study of the effects of these drugs on the mechanisms of the arrhythmia. The aim of administration of an antiarrhythmic drug in a patient with tachycardias is to select the best therapy for that particular patient. The drug can be given by any route at the appropriate dose and the patient retested once the antiarrhythmic effect is supposed to be present. The acute intravenous dose of an antiarrhythmic is given the day of the first electrophysiological investigation. The effect of oral treatment is studied after the number of days required to obtain adequate loading with the antiarrhythmic.

During the acute study the drug is given intravenously during tachycardia, except when tachycardia is poorly tolerated. When the arrhythmia is terminated by the intravenous drug, programmed stimulation using the same basic cycle lengths and stimulation sites as before drug administration is immediately repeated and continued for a period of at least 30-45 min. When the intravenous injection does not terminate tachycardia a period of 5-10 min is allowed between the end of injection and reinitiation of programmed stimulation to terminate tachycardia. Thereafter the effect of the drug on echo and tachycardia-zones, rate of tachycardia, mode of termination and patterns of A-V and V-A conduction and refractory periods are studied. Oral therapy is initiated simultaneously with or four to six hours after the intravenous injection. This is continued until the patient is restudied. The electrocardiogram of the patient is continuously monitored during that time.

Restudies during oral drug therapy:

There are several ways to restudy patients during oral antiarrhythmic treatment. When the effect of the given drug can be evaluated within three days of therapy we usually leave one or two intracardiac catheters in place to use them for stimulation and recording. However, this approach has some disadvantages. As previously outlined, the catheters are placed using the femoral approach which results in immobilization of the patient during the days before restudy. Also when catheters are left in place, a continuous infusion of full dose heparine is used. We do not leave a catheter in place for restudy in patients showing signs of congestive failure, those who are overweight, have venous insufficiency or those with contraindications to heparine therapy. Patients in whom the restudy cannot be performed within three days will leave the catheterization laboratory without catheters. The only exception are patients with frequent severely symptomatic or life-threatening arrhythmias in whom the catheter might be of help to repeatedly terminate the arrhythmia while the full effect of the drug has not been reached. In these patients we sometimes place a new catheter by way of the left cephalic vein. Placement of a catheter using the arm has the advantage of easy mobilization of the patient. In these patients full heparinization by a continuous infusion pump is done by using an infusion holder on wheels. If the catheter is placed in the right ventricle by way of the arm the restudy does not require readmission to the

catheterization laboratory. After right ventricular stimulation has been performed withdrawal of the catheter under electrocardiographic or portable X-ray control allows atrial stimulation. The catheter used is a quadripolar electrode, which permits recording of a bipolar electrogram during pacing.

In general we prefer to insert new catheters on the day of restudy. The patient is then readmitted to the catheterization laboratory. This has the advantage of allowing recording of electrograms at the same location as during the initial study. It is more costly however and occupies the catheterisation laboratory. We have safely performed restudies at the patient's bed using a portable recording and stimulation unit with resuscitation requirements available. Patients have found this approach more comfortable. Patients with life-threatening arrhythmias are not discharged from hospital before control of the arrhythmia has been obtained. This might require testing of several antiarrhythmics, alone and in combinations.

Special ways of stimulation to study patterns of conduction and mechanisms of tachycardias

By using the method of programmed electrical stimulation of the heart one can study (1-3) sites of origin, pathways and mechanisms of most tachycardias. As will be demonstrated in chapters 9 and 10 the differential diagnosis of mechanism and pathways in patients with supraventricular tachycardia can be difficult, therefore the following special protocols have been developed:

Special pacing protocols during supraventricular tachycardia.

Incessant supraventricular tachycardia.

We have studied eleven patients with the syndrome of incessant A-V junctional supraventricular tachycardia. In these patients participation of an accessory pathway in retrograde direction having a long conduction time and A-V nodal-like properties could be demonstrated by using some uncommon modes of stimulation during sinus rhythm, ventricular pacing or tachycardia.

There is a lot of controversy about the mechanism responsible for the different types of incessant supraventricular tachycardia. Incessant supraventricular tachycardia can either originate in the atrium or be based on a concealed accessory pathway having A-V nodal like properties and a long ventriculo-atrial conduction time (10-12). Theoretically a tachycardia showing a similar electrocardiogram can also be based upon an A-V nodal re-entry mechanism with anterograde conduction over a fast A-V nodal pathway and retrograde conduction over a slow A-V nodal pathway (the so-called reversed A-V nodal tachycardia).

Incessant supraventricular tachycardia can only be considered to be the result of an intranodal reversed (fast-slow) re-entrant circuit when use of an accessory pathway has been excluded. The A-V nodal-like properties of the slowly conducting accessory pathway have been responsible for confusion. As will be shown, demonstration of the presence and use of an accessory pathway with long conduction times can sometimes be very difficult.

Demonstration of the use of an accessory pathway in supraventricular tachycardia is easily obtained when during tachycardia a single ventricular premature beat delivered at the time of refractoriness of the bundle of His is able to advance subsequent atrial activation with the same activation pattern as during tachycardia (13). The ability to produce this phenomenon depends on several factors: first, the ventricular premature beat must be able to invade the re-entrant circuit. This will not only depend on the distance between the site of the ventricular premature beat and the location of the re-entrant circuit but also on the conduction velocity of the tissues between the site of stimulation and the re-entrant circuit. The ventriculo-atrial conduction time of a right ventricular apical extrasystole (a common site of stimulation during PES) is 30-50 ms longer when conducted over a left lateral accessory pathway as compared to the ventriculo-atrial conduction time of an extrasystole delivered at the same site with V-A conduction over a septally located accessory pathway (14). In the former situation the increase in the V-A conduction time due to transeptal activation can prevent demonstration of pre-excitation of the atria at the time of refractoriness of the bundle of His. This is a not uncommonly observed phenomenon in patients

using retrogradely a left-sided accessory pathway (14).

Secondly, advancement of subsequent atrial activation will not only depend upon location but also upon the properties of the retrograde pathway in the tachycardia circuit. In the common type of accessory pathway, which is one having fast conduction velocity, decremental conduction properties are usually absent. Accessory pathways having long conduction times show decremental conduction (10-13, 15-16). In these cases even when the re-entry circuit is invaded by a ventricular premature beat delivered during tachycardia an increase in the conduction time over the accessory pathway can prevent demonstration of advancement of atrial activation. In 5 out of our 11 patients having a septally located accessory pathway with a long conduction time a single ventricular premature beat given repeatedly at the time of refractoriness of the bundle of His did not result in advancement of subsequent atrial activation during tachycardia. This is a major challenge to the electrophysiologist performing the stimulation study, and special modes of pacing must be designed "in situ" to try to prove or exclude the use of an accessory pathway during tachycardia. Some of these methods are described here.

1. Delivery of a single late ventricular premature beat during sinus rhythm. A ventricular premature beat is delivered during sinus rhythm at the time of refractoriness of the bundle of His. As observed in one of our patients having an accessory pathway with long conduction time ventriculo-atrial conduction can occur following the ventricular premature beat and result in initiation of tachycardia (12). This mode of initiation of the arrhythmia can best be explained by accepting the presence of an accessory pathway. Theoretically one might consider the presence of simultaneously occurring dissociation in two electrophysiologically different pathways within the His bundle and the A-V node. A distal connection within the His bundle should be present and the A-V nodal pathways must be able to conduct retrogradely. So far such a mechanism has not been demonstrated. Functionally, though not anatomically, this situation would be similar to an accessory pathway.

Demonstration of V-A conduction during sinus rhythm at the time of refractoriness of the bundle of His is not possible in patients having the common fastly conducting type of accessory pathway. The fast conduction velocity results in arrival of the impulse in the atrium at a time when this is still refractory.

2. Delivery of a single late ventricular premature beat during tachycardia. Sometimes a single ventricular premature beat delivered at the time of refractoriness of the bundle of His will not pre-excite the atria, but result in termination of the arrhythmia when the accessory pathway has a long retrograde refractory period during tachycardia (11). Termination of tachycardia without V-A conduction of a ventricular premature beat given when the His bundle is refractory excludes an atrial site of origin of the arrhythmia, and proves that the ventricles are a necessary link in this supraventricular tachycardia. Such a finding proves that an accessory A-V pathway is used in retrograde direction (figure 9).

3. Delivery of single ventricular premature beats at different ventricular sites. When an accessory pathway located on the left side is used retrogradely during tachycardia a left ventricular premature beat will reach more easily the re-entry circuit than a right ventricular premature beat. The reverse is true in case of a right sided accessory pathway. Use of single ventricular premature beats at the appropriate ventricular site will demonstrate advancement of subsequent atrial activation at the time of His bundle refractoriness while they fail to do so when given at another ventricular site (fig. 10-11). In patients using retrogradely an accessory pathway located in the septum left ventricular pacing is probably of no benefit to demonstrate atrial pre-excitation during His bundle refractoriness when right ventricular apical premature beats failed to show this. In these cases, as described Coumel et al (17) stimulation high in the right ventricular septum can sometimes demonstrate the phenomenon because of the closer location to the reentry circuit.

4. Delivery of single ventricular premature beats during intraventricular aberrant conduction. Though small, the risks of left ventricular stimulation are rarely justified in patients studied because of a supraventricular arrhythmia when they do not undergo simultaneous left heart catheterization because of coronary heart disease or other reasons. When left heart catheterization

Figure 9

Termination of a supraventricular tachycardia having a P-R interval shorter than R-P' interval by a ventricular premature beat given in the right ventricular apex. The ventricular premature beat occurs at the time when the His bundle is refractory. Tachycardia is terminated without V-A conduction of the ventricular premature beat. This mode of termination indicates that both atria and ventricles are incorporated in the tachycardia circuit and proves that the accessory pathway has a long retrograde refractory period. 5 Surface leads were simultaneously recorded with five bipolar endocavitary electrograms.

is not performed, delivery of single ventricular premature beats during intraventricular aberrant conduction can be of help in demonstrating the presence of an accessory pathway. In the patient using retrogradely a left sided accessory pathway a single ventricular premature beat delivered in the right ventricular apex may fail to advance subsequent atrial activation when the His bundle is refractory for reasons discussed already. During left bundle branch block however even a much later premature beat can advance atrial activation when the His bundle is refractory. This is the result of the increase in the dimensions of the re-entrant circuit during left bundle branch block, the stimulation site (the right ventricular apex) being now closer to or even within the re-entry circuit. When a right-sided accessory pathway is used retrogradely single right ventricular premature beats usually result in clear advancement of subsequent atrial activation during tachycardia at the time of refractoriness of the bundle of His. This may even lead to a so called "paradoxical" capture of the atria due to a shorter conduction time from the site of stimulation in the right ventricle to the atria than the ventriculo-atrial conduction time during tachycardia. Left bundle branch block does not modify the dimensions of the re-entry circuit in these cases and does not change the effects of right ventricular apical premature beats. "Paradoxical" captures by right ventricular apical premature beats are however most easily demonstrated when right bundle branch block develops and the circuit enlarges. In patients using a truly septally located accessory pathway, intraventricular aberrant conduction does not result in easier or more difficult demonstration of atrial pre-excitation at the time of His bundle refractoriness.

5. Use of two ventricular premature beats to demonstrate retrograde conduction over an accessory pathway during supraventricular tachycardia. When a single ventricular premature beat fails to terminate tachycardia without V-A conduction at the time of refractoriness of the bundle of His or

Figure 10

A ventricular premature beat is given during circusmovement tachycardia using retrogradely an accessory pathway located laterally in the left A-V groove at the time that the His bundle is refractory for conduction. This ventricular premature beat which is given in the apex of the right ventricle fails to advance subsequent atrial activation.

to advance subsequent atrial activation, other modes of stimulation have to be applied to exclude or prove the use of a retrogradely conducting accessory pathway. Intraventricular aberrant conduction lengthens only minimally (≤ 25 ms) the V-A conduction time during a tachycardia using a truly septally located accessory pathway (18). Therefore in these patients the delivery of a second ventricular premature beat during tachycardia can be of help to demonstrate that an accessory pathway is being used retrogradely. There are several ways of giving these two ventricular premature beats:

a) The first ventricular premature beat is given at the time of refractoriness of the His bundle but fails to advance subsequent atrial activation. Under these circumstances it is impossible to decide whether or not the ventricular premature beat has entered the re-entry circuit and failed to advance subsequent atrial activation because of transeptal activation time or because of decremental conduction properties in the V-A link of the tachycardia circuit. Giving a second ventricular premature beat late in the cycle several possible observations can be made. The first and second ventricular premature beat are given when the His bundle is refractory for conduction during two consecutive cycles, and the interval between the two ventricular premature beats (S1-S2) is adjusted to exactly the tachycardia cycle length or a shorter cycle. Pre-excitation of the atria by the second ventricular premature beat can sometimes be demonstrated at the time of refractoriness of the bundle of His (fig. 12). This suggests that the first ventricular premature beat indeed invaded the re-entry circuit, but failed to pre-excite the atria because of the transeptal activation time or because of decremental conduction in the retrograde limb of the tachycardia circuit. When

Figure 11

Same patient as in fig. 10. A left ventricular premature beat paradoxically advances subsequent atrial activation and terminates tachycardia. These observations are explained by the shorter distance between the re-entry circuit and the stimulus delivered in the left ventricle. 6 Surface leads are shown simultaneously with six endocavitary bipolar electrograms.

the first premature beat did not capture the re-entry circuit atrial pre-excitation is not demonstrated by a second, third or fourth ventricular premature beat (fig. 13).

b) The first ventricular premature beat is given at the time of refractoriness of the His bundle and fails to advance subsequent atrial activation. When the second ventricular premature beat is given early in the cycle before the expected anterograde His bundle depolarization this can be of help to demonstrate the presence of an accessory pathway. In figure 14 two examples are given. In the top panel the second ventricular premature beat advances subsequent atrial activation without modifying anterograde depolarization of the His bundle. In the bottom panel a 10 ms shortening of the second ventricular premature beat interval terminates tachycardia without V-A conduction and without influencing the expected anterograde depolarization of the His bundle.

c) The first ventricular premature beat has failed to pre-excite the atria at the time of refractoriness of the bundle of His, but when given early in the cycle results in termination of tachycardia without retrograde conduction to the atria. In this situation it is not possible to decide whether the tachycardia is located intranodally or uses an accessory pathway retrogradely, although an atrial site of origin of tachycardia is excluded. Interesting observations can be made by giving a second ventricular premature beat. When the first ventricular premature beat is given after the correct interval to terminate tachycardia and the second is given very late after the first (at least twice the tachycardia cycle length), this second ventricular premature beat can reinitiate tachycardia following V-A conduction. On increasing the prematurity of the second premature beat a zone of

Figure 12

During circus movement tachycardia using retrogradely an accessory left-sided A-V pathway a right ventricular apical premature beat given after 230 ms (panel A) fails to advance subsequent atrial activation. The ventricular premature beat is given just before the expected anterograde His bundle depolarization and does not affect the A-H interval. A premature beat given after the same interval advances subsequent atrial activation by 10 ms when it is preceded by another ventricular premature beat (panel B) and when given during tachycardia with left bundle branch block aberrancy (panel C). Five surface leads are simultaneously recorded with 3 endocavitary bipolar electrograms.

Figure 14

In panels A and B the first ventricular premature beat given at the time of refractoriness of the bundle of His fails to advance subsequent atrial activation. In panel A a second ventricular premature beat advances atrial activation without modifying anterograde depolarization of the bundle of His. A 10 ms shortening of the second ventricular premature beat interval terminates tachycardia. This occurs without V-A conduction and without modifying anterograde depolarization of the bundle of His (panel B). These observations indicate participation of an accessory A-V pathway as the retrograde arm of the tachycardia circuit. 5 Surface leads simultaneously recorded with four bipolar endocavitary electrograms.

complex). Under these circumstances, advancement of subsequent atrial activation can occur because of retrograde conduction over an accessory pathway or over an A-V nodal retrograde pathway prematurely engaged after retrograde depolarization of the His bundle. A second ventricular premature beat can advance subsequent atrial activation when the first one is given early in the cycle and the second one at the time of refractoriness of the bundle of His by the same principles previously discussed. When given earlier in the cycle it can be shown that the V-A conduction time of the first and second premature beat are the same and that no decremental conduction occurs. This finding is very suggestive for the use of an accessory pathway in ventriculo-atrial direction.

These special modes of stimulation during supraventricular tachycardia are not the only way to demonstrate use of an accessory pathway during tachycardia. There are other criteria useful to suggest that diagnosis like eccentric atrial activation during tachycardia when an atrial site of origin of the arrhythmia can be excluded, or an increase in V-A interval during tachycardia upon development of bundle branch block ipsilateral to the location of the accessory pathway. The special modes of stimulation which we discussed are especially useful when A-V nodal retrograde conduction simulates accessory pathway conduction (12,19) or when accessory pathway conduction simulates retrograde A-V nodal conduction (as in septally located accessory pathways).

In the literature it can be repeatedly observed that an A-V nodal re-entrant tachycardia is diagnosed when **no evidence** for the use of an accessory pathway during tachycardia is found. An intranodal tachycardia is diagnosed when a ventricular premature beat **does not** pre-excite the atria at the time of His bundle refractoriness, when the atrial activation sequence is **not** eccentric, when intraventricular aberrant conduction **does not** modify the V-A interval during tachycardia. These arguments do not prove the presence of an accessory pathway, but they do not exclude its use during tachycardia either. Negative observations do not **prove** that the tachycardia is located intranodally. Positive arguments have to be presented to exclude use of an accessory pathway. Positive arguments also have to prove an intranodal or atrial location of the re-entry circuit. We want to stress that the presence of discontinuous A-V nodal conduction curves suggesting the presence of dual A-V nodal pathways is not diagnostic of an intranodal location of the tachycardia circuit, even when tachycardia is initiated at the time of the shift in conduction from the fast to the slow A-V nodal pathway. Up to 10% of patients having an accessory pathway used retrogradely during tachycardia will demonstrate discontinuous A-V nodal conduction curves during atrial stimulation (20). Similarly, the presence of an accessory pathway does not mean that it is used during supraventricular tachycardia. Examples of intranodal tachycardias in patients with accessory pathways have been reported (21-23).

Positive arguments to exclude use of an accessory pathway during supraventricular tachycardia:

There are some very useful findings which positively exclude use of an accessory A-V pathway during supraventricular tachycardia, and also some observations which positively prove an intranodal location of the re-entry circuit:

- a) Demonstration of A-V block during tachycardia without termination of the arrhythmia: This phenomenon can sometimes be observed to occur spontaneously, positively excluding the use of an accessory A-V pathway during tachycardia. Single or multiple atrial or ventricular premature beats can result in A-V block of subsequent tachycardia P waves without terminating tachycardia (fig. 15). It can also be produced by vagal manoeuvres.
- b) The occurrence during tachycardia of atrial activation which precedes or occurs simultaneously with ventricular activation (fig. 16). In this situation there is no true "V-A" conduction therefore an accessory A-V pathway as participating retrogradely in the tachycardia circuit is excluded (13).

Positive arguments to suggest an intranodal location of the tachycardia circuit are:

- a) The demonstration that only the A-V node is required to initiate and perpetuate the arrhythmia. One has to show the simultaneous occurrence of H-A and H-V block during tachycardia without termination of the arrhythmia. Josephson et al. have suggested that by delivering atrial premature beats it can be shown that most of the atrial activation can be advanced without modifying the timing of activation of the His bundle (24). This phenomenon depends on the conduction

Figure 15

Demonstration of A-V block during supraventricular tachycardia by ventricular pacing without termination of the arrhythmia. These observations exclude a tachycardia using an accessory A-V pathway. 5 surface leads and one right atrial lead are shown.

properties of the A-V node during tachycardia. The atrial premature beat can pre-excite the atrium completely, penetrate the re-entry circuit, but because of delay in conduction over the slow A-V nodal pathway result in activation of the His bundle on time, suggesting that the re-entry circuit has not been penetrated. We believe that this observation cannot be taken as a positive argument to exclude participation of the atria in the tachycardia circuit (figure 17).

b) Termination of a supraventricular tachycardia showing simultaneous activation of atria and ventricles (common type of A-V nodal tachycardia) or a P'-R shorter than the R-P' (uncommon or "reversed" form of A-V nodal tachycardia) by a ventricular premature beat which is not retrogradely conducted to the atrium after an accessory pathway has been excluded. This mode of termination does exclude an atrial site of origin of tachycardia.

c) Initiation of the uncommon type of A-V nodal tachycardia by a ventricular premature beat in relation to delay in retrograde conduction (shift from retrograde fast to retrograde slow A-V nodal pathway) after an accessory pathway has been excluded by positive arguments (figure 18).

d) Demonstration of two zones of initiation of tachycardia of the uncommon variety during ventricular stimulation. The first zone is determined by the latest premature beat resulting in shift of conduction from the retrograde fast to the retrograde slow pathway up to the ventricular premature beat interval which reaches the retrograde refractory period of the slow pathway. The second zone occurs after a "gap" in initiation of tachycardia, and begins when sufficient subnodal delay of retrograde conduction of the ventricular premature beat allows arrival of the impulse to the A-V nodal slow pathway late enough after its refractory period. This observation proves that the slow pathway is located above the His bundle, and that there is no accessory A-V pathway with long conduction times (12).

We have observed only one case of long-lasting (60 beats) but non-sustained A-V nodal tachycardia of the reversed form. This arrhythmia was never documented clinically. Our patient demonstrated A-V block during tachycardia, without termination of the arrhythmia, thus excluding use of a retrogradely conducting accessory pathway with long conduction times.

Figure 16

Initiation of a supraventricular tachycardia during atrial pacing by a critically timed atrial premature beat during atrial pacing. At the time of initiation and during supraventricular tachycardia atrial activation occurs simultaneously with ventricular activation (panel B). This excludes incorporation of an accessory A-V pathway in the tachycardia circuit.

Special pacing protocols used during a wide qrs complex tachycardia with 1-to-1 relation between atrial and ventricular activation.

Several diagnostic possibilities must be considered during a wide QRS complex tachycardia showing a 1-to-1 A-V relation.

- Supraventricular tachycardia in a patient with a functional or a pre-existing bundle branch defect,
- Atrial tachycardia with anterograde conduction over an accessory A-V pathway,
- Circusmovement tachycardia using anterogradely an accessory pathway and retrogradely the normal A-V pathway or a second accessory pathway, and
- Ventricular tachycardia with 1-to-1 retrograde conduction to the atria.

In situation a) His bundle activation precedes and does not follow the QRS complex. This type of tachycardia will therefore not be considered. When confronted with possibilities b) to d) our first approach is to give an atrial premature beat with a progressively shorter interval. Several possibilities can occur:

1. Ventricular capture by the atrial premature beat with a QRS complex showing a different morphology as compared to the tachycardia QRS complex, without termination of tachycardia,

Figure 17

During a circusmovement tachycardia using retrogradely a left-sided accessory A-V pathway an atrial premature beat advances all atrial electrograms but does not modify the timing of subsequent anterograde activation of the His bundle. This observation cannot be used to exclude participation of the atria in the mechanisms of the arrhythmia. In tachycardias using an accessory A-V pathway, like the one shown in this figure, the atria are part of the re-entrant circuit.

followed by a fully compensatory pause at the ventricular level. This observation indicates that the pathway used anterogradely by the atrial premature beat is not used anterogradely during tachycardia. When a fully compensatory pause follows at the ventricular level anterograde use of an accessory A-V pathway during tachycardia is excluded. The observation of a fully compensatory pause without atrial activation in between indicates not only a ventricular site of origin of tachycardia, but suggests as pointed out by Josephson et al. (25) a relatively small and protected circuit of ventricular tachycardia. Ventricular capture by supraventricular stimuli can sometimes only occur by giving several atrial premature beats or by pacing the atria at rates faster than the tachycardia rate.

2. Exact ventricular capture (the V-Ve interval being identical to the A-Ac interval) with the same QRS morphology as during tachycardia with advancement of the tachycardia cycle. This observation (fig. 19) is the best way to demonstrate the use of anterograde conduction over an accessory A-V pathway during tachycardia. It indicates that the same pathway used to capture the ventricles by the atrial premature beat is used anterogradely during tachycardia. Exact advancement of the tachycardia cycle indicates absence of decremental conduction properties in the anterograde limb, and participation of the atria in the tachycardia circuit and of the ventricles, thus proving use of an accessory A-V pathway anterogradely.

3. Atrial premature beat does not result in ventricular capture but terminates tachycardia. This observation excludes a ventricular site of origin of tachycardia, the only exception being bundle branch re-entry tachycardia (26). If a supraventricular site of origin of the arrhythmia has been excluded tachycardia is the result of re-entry in the bundle branch system. When the atrial premature beat terminates the wide QRS tachycardia without being conducted to the bundle of His a ventricular site of origin and bundle branch re-entrant tachycardia are excluded. This observation indicates participation of the atria in the mechanisms of the arrhythmia. When during tachycardia activation of the His bundle follows the QRS complex, an A-V nodal site of origin of tachycardia with anterograde use of a nodo-ventricular accessory pathway cannot be excluded by this observation alone.

Figure 18

Initiation of a supraventricular tachycardia with a P-R shorter than R-P interval by a ventricular premature beat at the time of a change in conduction from a retrograde fast to a retrograde slow pathway. Following initiation A-V block occurs not resulting in termination of the arrhythmia. These observations suggest an intra A-V nodal location of the re-entry circuit.

Figure 19

Ventricular capture by an atrial premature beat showing the same QRS morphology as during tachycardia. This capture has at the ventricular level the same interval as the coupling interval of the atrial premature beat. The capture is followed by a shift in subsequent tachycardia cycles to the left. This observation proves that:

a. The same pathway used to activate the ventricles during tachycardia is used during anterograde conduction of the atrial premature beat.

b. the atria and ventricles are incorporated in the tachycardia circuit.

c. no decremental conduction occurs in the anterograde pathway.

They prove the use during tachycardia of an anterogradely conducting accessory A-V pathway.

4. The atrial premature beat does not result in ventricular capture or termination of tachycardia, but the subsequent tachycardia QRS complex shows delay or block in retrograde conduction to the atria without affecting the ventricular rate of the tachycardia. This observation excludes participation of the atria in the mechanisms of tachycardia.

5. The atrial premature beat fails to modify subsequent atrial and ventricular events. No conclusions can be made from this observation in regard to mechanism or pathway of tachycardia.

In patients with suspected or documented supraventricular tachycardia we start our stimulation study by pacing in the ventricle. We start by pacing in the atria in patients with suspected or documented ventricular tachycardia. The observation of V-A dissociation during ventricular pacing makes a tachycardia based upon re-entry using an accessory A-V pathway or an intranodal tachycardia unlikely and suggests the likelihood of an atrial tachycardia. A long refractory period of the fast retrograde A-V nodal pathway resulting in discontinuous retrograde A-V nodal conduction curves makes initiation of A-V nodal tachycardia of the common variety unlikely during atrial stimulation. As pointed out elsewhere A-V nodal tachycardia may then be initiated after abbreviation of the fast pathway refractory period by atropine.

Electrophysiologic studies are expensive (3). They may be dangerous in inexperienced hands. Adequate training resulting in understanding of the mechanisms of tachycardias and their expressions during a stimulation study is required before the start of a clinical electrophysiologic

program. Well performed electrophysiologic studies are of help in selecting the best therapy for a particular arrhythmia. Poor electrophysiological investigations, however, can result in wrong conclusions, the selection of inadequate therapy and risk for the patient.

Results of programmed electrical stimulation of the heart

Table I summarizes the results of programmed electrical stimulation of the heart in a series of 440 consecutive patients with supraventricular tachycardia. Only in 3 out of these 440 patients the exact mechanism of the arrhythmia could not be uncovered. All three patients suffered from A-V junctional supraventricular tachycardia, but the exact pathways could not be determined.

Table I

Incidence of different types of tachycardia in 440 patients with electrocardiographically documented regular paroxysmal supraventricular tachycardia (SVT), studied by programmed electrical stimulation of the heart.

<i>Type of SVT</i>	<i>Number</i>	<i>Percentage</i>
Atrial tachycardia	58	13.1
Atrial flutter	60	13.6
Intranodal tachycardia	107	24.3
Circusmovement tachycardia using a concealed accessory pathway	61	13.8
Unknown mechanism	3	0.6
WPW syndrome	151	34.3
	440	

References

1. Durrer D, Schöo L, Schuilenburg RM, Wellens HJJ: The role of premature beats in the initiation and termination of supraventricular tachycardia in the Wolff-Parkinson-White syndrome. *Circulation* 1967; 36:644-62.
2. Wellens HJJ: Electrical stimulation of the heart in the study and treatment of tachycardias. Leiden, Stenfert Kroese, 1971.
3. Wellens HJJ: Value and limitations of programmed electrical stimulation of the heart in the study and treatment of tachycardias. *Circulation* 1978; 57:845-52.
4. Gallagher JJ, Svenson RH, Kasell JK, et al: Catheter technique for closed-chest ablation of the atrio-ventricular conduction system. *N.Eng.J.Med.* 1982; 306:194-200.
5. Gallagher JJ, Pritchett ELC, Benditt DG et al: New catheter techniques for analysis of the sequence of retrograde atrial activation in man. *Eur.J.Cardiol.* 1977; 6: 1-14.
6. Dassen WRM, Ross DL, Brugada P, Bär F, Vanagt E, Wellens HJJ: A computer model for the study of cycle length alternation during tachycardia. *Proceedings of Computers in Cardiology*, 1980; 179-82.
7. Ross DL, Dassen WRM, Vanagt E, Brugada P, Bär FWHM, Wellens HJJ: Cycle length alternation in circus movement tachycardia using an atrio-ventricular accessory pathway. *Circulation* 1982; 65:862-8.
8. Mercer AB, Simpson RJ, Gettes LS, Foster JR: Strength-interval testing to enhance ventricular tachycardia induction during clinical electrophysiology study. *Am. J. Cardiol.* 1982; 49:958 (Abstr.)
9. Vandepol CJ, Farshidi A, Spielman SR, Greenspan AM, Horowitz LN, Josephson ME: Incidence and clinical significance of induced ventricular tachycardia. *Am. J. Cardiol.* 1980; 45:725-31.

10. **Farré J, Ross D, Wiener I, Bär F, Vanagt E, Wellens HJJ:** Reciprocal tachycardias using accessory pathways with long conduction times. *Am. J. Cardiol.* 1979; 44:1099-109.
11. **Brugada P, Vanagt EJ, Bär F, Wellens HJJ:** Incessant reciprocating atrio-ventricular tachycardia. *PACE* 1980; 3:670-7.
12. **Brugada P, Bär F, Vanagt EJ, Friedman PL, Wellens HJJ:** Observations in patients showing A-V junctional echoes with a P-R shorter than R-P interval. *Am. J. Cardiol.* 1981; 64:611-22.
13. **Wellens HJJ, Durrer D:** The role of an accessory atrio-ventricular pathway in reciprocal tachycardia. *Circulation* 1975; 52:58-72.
14. **Weiss J, Brugada P, Roy D, Bär FWHM, Wellens HJJ:** Localization of the accessory pathway in the WPW syndrome from the ventriculo-atrial conduction time of right ventricular apical extrasystoles. *PACE*, in press.
15. **Gallagher JJ, Sealy WC:** The permanent form of junctional reciprocating tachycardia: further elucidation of the underlying mechanism. *Eur J Cardiol.* 8:413, 1978.
16. **Denes P, Kehoe P, Rosen KM:** Multiple re-entrant tachycardias due to retrograde conduction of dual atrio-ventricular bundles with atrio-ventricular nodal-like properties. *Am J Cardiol.* 1979; 44:498.
17. **Coumel P, Attuel P, Mugica I:** Junctional reciprocal tachycardia. The permanent form. In HE Kulbertus (ed): *Re-entrant arrhythmias*. MTP Press, Lancaster 1977, p. 170.
18. **Kerr ChR, Gallagher JJ, German LD:** Effect of bundle branch block on ventriculo-atrial intervals during reciprocating tachycardia in patients with accessory atrio-ventricular pathways. *Am. J. Cardiol.* 1982, 49:994 (Abstr).
19. **Sung RJ, Waxman HL:** Physiological shift of retrograde atrio-ventricular nodal output mimicking retrograde accessory pathway conduction. *Circulation* 1979; 54-60: Suppl II:II-191 (Abstr).
20. **Wellens HJJ, Bär F, Farré J, Wiener I, Ross D, Gorgels AP, Vanagt E, Brugada P:** Diagnóstico e incidencia de vías accesorias ocultas en pacientes con taquicardias paroxísticas de la Unión AV. *Rev. Lat. Cardiol.* 1980, 3:60-71.
21. **Spurrell RAJ, Krikler D, Sowton E:** Two or more intra A-V nodal pathways in association with either a James or Kent extranodal bypass in three patients with paroxysmal supraventricular tachycardia. *Br. Heart J.* 1973, 35:113-9.
22. **Neuss M, Schlepper M:** Unusual re-entry mechanisms in patients with the Wolff-Parkinson-White syndrome. *Br. Heart J.* 1974; 36:880-6.
23. **Mandel WJ, Laks MM, Obayashi K:** Atrio-ventricular nodal re-entry in the Wolff-Parkinson-White syndrome. *Chest* 1975; 68:321-6.
24. **Josephson ME, Kastor JA:** Paroxysmal supraventricular tachycardia. Is the atrium a necessary link? *Circulation* 1976; 54:430-8.
25. **Josephson ME, Horowitz LN, Farshidi A, Kastor JA:** Recurrent sustained ventricular tachycardia. *Circulation* 1978; 57:431-40.
26. **Wellens HJJ, Lie KI, Durrer D:** Further observations on ventricular tachycardia as studied by electrical stimulation of the heart. *Circulation* 1974; 49:647-53.

3.

Atrioventricular nodal tachycardia with and without discontinuous anterograde and retrograde atrioventricular nodal conduction curves: a reappraisal of the dual pathway concept

P. Brugada, E. Vanagt, W.R. Dassen, A.P. Gorgels, F.W.H.M. Bär and H.J.J. Wellens.

Department of Cardiology, University of Limburg, Annadal Hospital, Maastricht, The Netherlands

KEY WORDS: A-V nodal tachycardia, dual pathways.

To illustrate the complexity of the electrophysiological behaviour of the human atrioventricular (A-V) node, two patients suffering from A-V nodal tachycardia are described. During tachycardia an A-V nodal slow pathway was used for anterograde conduction, and an A-V nodal fast pathway for retrograde conduction. Patient 1 showed smooth A-V nodal conduction curves in both the anterograde and the retrograde direction. Tachycardia could only be initiated by ventricular premature beats. No critical delay in ventriculo-atrial conduction time was required for initiation of tachycardia. Patient 2 showed smooth A-V nodal conduction curves at the lowest rate of pacing during atrial and ventricular stimulation. The curves became discontinuous in both directions when the basic driven-cycle length was decreased. Tachycardia could only be initiated by atrial premature beats. Ventricular premature beats induced non-sustained A-V nodal reentry that used an A-V nodal fast pathway for anterograde conduction, and an A-V nodal slow pathway for retrograde conduction. Accepting dual pathways in the anterograde and retrograde directions in the A-V node, means that depending upon their electrophysiological properties a large number of combinations of anterograde and retrograde conduction are possible. When more than two A-V nodal pathways are present, the number of possible combinations will increase markedly. These considerations are of help in understanding electrophysiological findings in patients with A-V nodal tachycardia. They are also useful to explain the importance of autonomic tone and the results of drug administration in these patients.

Most patients with atrioventricular (A-V) nodal reentrant tachycardia and their tachycardia circuit confined to the A-V node show discontinuous anterograde A-V nodal conduction curves during extrastimulus testing, suggesting the presence of dual A-V nodal pathways^[1]. In retrograde direction, A-V nodal conduction curves are usually continuous. Only a few examples have been reported of patients with intra A-V nodal reentrant tachycardia showing discontinuous anterograde and retrograde A-V nodal conduction curves, or discontinuous curves solely in the retrograde direction^[2]. The initiation of a supraventricular

tachycardia by a single timed ventricular premature beat without apparent increase in the V-A conduction time of the premature beat has been considered suggestive of a tachycardia either using an accessory pathway or located in the atrium^[3-5]. The initiation of intra A-V nodal reentrant tachycardia by an atrial premature beat in patients showing continuous anterograde A-V nodal conduction curves has been reported^[6-8], but the electrophysiological mechanisms involved have not been explained.

In this article, two patients are described in whom an A-V nodal reentrant tachycardia utilizing an A-V nodal slow pathway for anterograde conduction, and an A-V nodal fast pathway for retrograde conduction was initiated during the stimulation study. Patient 1 showed continuous anterograde and retrograde A-V nodal conduction curves,

Received for publication 23 June 1980.

Requests for reprints to: Hein J. J. Wellens, Department of Cardiology, University of Limburg, Annadal Hospital, Maastricht, The Netherlands.

and tachycardia could only be initiated with a single ventricular premature beat. Patient 2 showed continuous anterograde and retrograde A-V conduction curves at the lowest rates of pacing, but the curves became discontinuous in both directions on increasing the basic pacing rate. The examples were chosen to illustrate the wide spectrum of electrophysiological behaviour of the A-V node during stimulation studies in patients with A-V nodal tachycardia.

Patients and methods

Patient 1 was a 63 year old male. Nine years prior to the study he suffered from an uncomplicated inferior myocardial infarction. Palpitation started four years before our investigation. Spontaneous episodes of tachycardia were characterized by a narrow QRS complex without identifiable P waves. Oral digoxin in a dose of 0.25 mg/day did not prevent the arrhythmia.

Patient 2 was a 75 year old female. She complained of palpitations for three years before our study. No tachycardia had been documented electrocardiographically. Cardiovascular examination revealed no abnormalities. She was receiving no medication.

Electrophysiological studies

Following informed consent, an electrophysiological study was performed on both patients in the resting, postabsorptive, non-sedated state. In patient 1, digoxin had been discontinued 48 h prior to the investigation. Using the femoral approach, catheters were placed for both stimulation and recording of intracardiac electrograms, at the right ventricular apex, tricuspid valve ring and high lateral right atrium. In patient 1, a fourth catheter was placed in the left atrium via a patent foramen ovale. Our methods of recording, stimulation and analysis of tracings have been previously described⁹. After recording of basic conduction intervals during sinus rhythm, the right atrium was paced in patient 1 at basic cycle lengths of 665 and 600 ms, and in patient 2 at basic cycle lengths of 600, 545, 500, 460 and 430 ms. Refractory periods were determined using the extrastimulus technique. Ventriculo-atrial (V-A) conduction was studied during pacing of the right ventricular apex using the single test stimulus technique at the same basic cycle lengths as during atrial stimulation. The effect

of timed atrial and ventricular premature beats was tested during tachycardia.

Definitions

Conventional definitions were used for refractory periods. An A-V nodal conduction curve was considered to be discontinuous if the A-V nodal conduction time of a premature beat (atrial or ventricular) increased by 50 ms or more on increasing the prematurity of the test pulse by 10 ms⁴. Diagnosis of A-V nodal tachycardia using anterograde slow and retrograde fast pathway conduction was based upon:

- (1) Simultaneous activation of atria and ventricles during tachycardia⁴.
- (2) Inability to advance atrial activation by ventricular premature beats given during tachycardia at the time of refractoriness of the bundle of His.¹⁰
- (3) A similar pattern of atrial activation during tachycardia and following V-A conduction during ventricular pacing.

Results

PATIENT 1

His bundle recordings during sinus rhythm (rate 75 beats/min) revealed an A-H interval of 80 ms, and a prolonged H-V interval of 65 ms.

Single atrial test stimulation at a basic cycle length of 665 and 600 ms showed a continuous A_1A_2/H_1H_2 curve with an A-V nodal effective refractory period of 310 ms (Fig. 1). Neither echo beats nor tachycardia could be induced.

Ventricular stimulation at a basic cycle length of 665 ms revealed a continuous V_1V_2/A_1A_2 curve up to a V_1V_2 interval of 310 ms. Ventricular premature beats given at shorter V_1V_2 intervals showed an increase in V_2A_2 intervals. As shown in Fig. 2 this increase in V-A conduction time was subnodally located at very premature beat intervals, and it was assumed that the same site of delay accounted for the changes in V-A conduction time of ventricular premature beats given later in diastole. Without relation to the V-A conduction time of the ventricular test stimulus (V_2), tachycardia could be initiated by premature beats given with V_1V_2 intervals ranging from 460 to 250 ms, but only if the resulting A_1A_2 interval ranged from 460 to 360 ms. A premature beat given with a V_1V_2 interval of 310 ms did not induce tachycardia when the A_1A_2 interval was 350 ms, but only when the A_1A_2

Figure 1 Case 1. (a) Graphs showing the relation between the A_1A_2 and the H_1H_2 and A_2H_2 intervals during atrial pacing with basic cycle lengths of 665 and 600 ms. As shown the curves were continuous and no echo beats or tachycardia were induced. (b) The relation between the V_1V_2 and the A_1A_2 and V_2V_2 intervals during ventricular pacing with basic cycle lengths of 665 and 600 ms. Tachycardia (TC) was initiated without relation to the ventriculo-atrial conduction time of V_2 . Initiation of tachycardia was related to a critical range of A_1A_2 intervals. See text.

interval measured 400 ms. Similar observations were made during ventricular extrastimulus testing at a basic cycle length of 600 ms (Fig. 1).

During tachycardia atria and ventricles were simultaneously activated (Fig. 2). Onset of atrial activation was first recorded in the His bundle lead

and preceded ventricular activation. Subsequent atrial activation occurred in the proximal left atrial (LAp), the distal left atrial (LAd), and the high right atrial (HRA) recording. The R-R interval during tachycardia ranged from 410 to 480 ms. One single ventricular premature beat and one single HRA

Figure 2 Case 1. Six different ventricular premature beat intervals are shown during pacing of the right ventricle with a basic cycle length of 665 ms. Tachycardia is initiated without apparent relation to the ventriculoatrial conduction time of V_6 , when the A_1A_2 interval ranges from 360 to 460 ms. Note that with very early premature beats a retrograde His potential is recorded preceding atrial activation. Lead V_6 , a high right atrial (HRA) lead, a distal left atrial (LAAd) lead, a His bundle lead, and a proximal left atrial (LAP) lead were recorded simultaneously.

premature beat could reproducibly terminate the tachycardia.

PATIENT 2

His bundle recordings during sinus rhythm (rate 80 beats/min) revealed an A-H interval of 75 ms, and an H-V interval of 40 ms.

Single atrial test stimulation at basic cycle lengths of 600 and 545 ms showed continuous anterograde A-V nodal conduction curves, with A-V nodal conduction occurring up to the refractory period of the right atrium (Fig. 3). During pacing with basic cycle lengths of 500, 460 and 430 ms however the A-V nodal conduction curves became discontinuous, with an A-V nodal effective refractory period of the fast pathway of 280, 290 and 300 ms respectively. A-V nodal conduction over the slow pathway occurred up to the refractory period of the right atrium. Tachycardia could only be initiated by a single

atrial premature beat during pacing of the right atrium at a basic cycle length of 430 ms (Fig. 4).

Ventricular stimulation at a basic cycle length of 600 ms revealed a continuous retrograde conduction curve up to the refractory period of the right ventricle (Fig. 3). Decreasing the basic cycle length resulted in discontinuous retrograde conduction curves, with the induction of A-V nodal echo beats. As shown in the His bundle recordings, the discontinuity in the retrograde conduction curve was the result of retrograde A-V nodal conduction over the slow pathway (Fig. 4). A-V nodal echo beats following a ventricular test stimulus used the A-V nodal slow pathway for retrograde conduction and the A-V nodal fast pathway for anterograde conduction. No sustained tachycardia could be initiated.

During the tachycardia initiated by an atrial premature beat atria and ventricles were simultaneously activated (Fig. 4). The onset of atrial activation was first recorded in the His bundle lead,

Figure 4 Case 2. (a), (b) and (c) Show a ventricular premature beat given after 290 ms during ventricular pacing with a basic cycle length of 600, 500 and 430 ms respectively. Note that the V_2H_2 interval becomes shorter on decreasing the basic cycle length and is hidden into the QRS complex at a basic cycle length of 430 ms (c). In (a) V_2 is retrogradely conducted over the A-V nodal fast pathway, and no echo beat is induced. In (b) and (c) V_2 is retrogradely conducted over the slow pathway and an A-V nodal echo beat appears showing anterograde conduction over the fast pathway. (d) Shows the initiation of A-V nodal tachycardia by an atrial premature beat given after 300 ms during atrial stimulation at a basic cycle length of 430 ms. During tachycardia there is anterograde conduction over the slow pathway and retrograde conduction over the fast pathway.

and preceded the onset of ventricular activation. The R-R interval during tachycardia ranged from 370 to 410 ms. Tachycardia could reproducibly be terminated by a single atrial or ventricular premature beat.

Discussion

The initiation of a supraventricular tachycardia by a single ventricular premature beat in patient 1, without apparent increase in the V-A conduction time has been considered suggestive that the tachycardia is either using an accessory pathway or is located in the atrium¹³⁻¹⁵. In this patient the identical sequence of atrial activation during ventricular pacing and tachycardia, and the inability to pre-excite the atria when the His bundle was refractory, argues against the presence of an accessory pathway.

In 1975 one of us¹¹ reported a case of supraventricular tachycardia initiated by a single ventricular premature beat. No critical increase in the V-A conduction time of the ventricular premature

beat was required to initiate the tachycardia. When the A_1A_2 interval following V_1V_2 reached a certain value, tachycardia was initiated. The initiation of tachycardia without an increase in V-A conduction time was considered suggestive of an intra-atrial location of the tachycardia circuit. A limit of A_1A_2 intervals below which no tachycardia could be induced was not present in that patient. Similarly our patient 1 showed a critical range of A_1A_2 intervals for the initiation of tachycardia. Now also a lower limit was found of A_1A_2 intervals which were not followed by tachycardia. The most likely explanation for our patient from 1975 and our present patient 1 is an A-V nodal tachycardia in the presence of dual A-V nodal pathways with the fast pathway having the shortest effective refractory period in both the anterograde and the retrograde direction. Under these circumstances, discontinuities in the anterograde and retrograde A-V nodal conduction curve will not be found during atrial and ventricular pacing. The presence of dual A-V nodal pathways can only indirectly be demonstrated by inducing A-V nodal echoes or

tachycardia during the stimulation study. If with continuous anterograde and retrograde A-V nodal conduction curves, the effective refractory period of the slow pathway in the anterograde direction is shorter than the retrograde effective refractory period, only critically timed and conducted ventricular premature beats will be able to induce A-V nodal echoes or tachycardia, as shown in Fig. 5. The relation between the refractory periods of the two A-V nodal pathways and the mode of anterograde and retrograde A-V nodal conduction explains the absence of dual pathway curves during anterograde and retrograde A-V nodal conduction, and the initiation of tachycardia without change in the V-A conduction time.

In patient 2 decreasing the basic paced cycle length brought out discontinuities in the A-V nodal conduction curve that were not apparent at slower rates of pacing. Factors influencing demonstration of dual A-V nodal pathways during atrial stimulation are the refractory period of the atrium and

the properties of the A-V node itself. Both are known to be related to the basic cycle length^[11-14]. In our patient 2, the effective refractory period of the fast pathway in the anterograde direction became longer on shortening the basic cycle length. Also during ventricular stimulation the A-V nodal properties are expected to change with changes in basic cycle length. In patient 2, the retrograde effective refractory period of the fast pathway lengthened on decreasing the basic cycle length. When the ventricle was paced at different rates the His bundle electrogram of the basic beats was hidden in the QRS complex, so that plotting of the V_1V_2 and H_1H_2 intervals was not possible. However, as shown in Fig. 4, when the same V_1V_2 interval was studied at different basic cycle lengths, the V_1H_2 interval shortened on decreasing the cycle length of basic pacing. Figure 4 shows how depending upon the basic cycle length subnodal delay of V_2 can lead to failure to demonstrate discontinuous retrograde A-V nodal conduction curves. It illustrates that the

Figure 5 Case 1. Diagram showing the proposed mechanism of initiation of A-V nodal tachycardia by a single ventricular premature beat. A, atrium; AVN, atrioventricular node; V, ventricle; V_1 , A₁, last basic driven ventricular beat and its retrogradely conducted atrial activation; V_2 , A₂, ventricular premature beat and its retrogradely conducted atrial activation. a, b, c and d are different degrees of prematurity of V_2 . Top dashed area = anterograde effective refractory period of the slow pathway. Bottom dashed area = retrograde effective refractory period of the slow pathway. During basic pacing, ventriculo-atrial conduction occurs over the A-V nodal fast pathway (—). This impulse reenters the A-V nodal slow pathway (—) and collides with the impulse retrogradely conducted over the slow pathway. A late V_2 , a, is retrogradely conducted similar to the basic beat, and tachycardia is not initiated. A more premature V_2 , b, is retrogradely conducted to the atrial level over the fast pathway, but blocked retrogradely in the slow pathway. Now the impulse reentering the A-V nodal slow pathway is able to traverse the slow pathway and tachycardia is initiated. A more premature V_2 , c, will be conducted to the atrium over the fast pathway, and blocked retrogradely in the slow pathway. The reentering impulse, however, finds the slow pathway refractory in anterograde direction and no tachycardia will be initiated. A very premature V_2 , d, is retrogradely conducted to the atrium with subnodal delay. The later time of arrival at the distal A-V node will permit the same sequence of events as described for a premature beat given after interval b. Note that the tachycardia uses the A-V nodal fast pathway retrogradely and the A-V nodal slow pathway anterogradely.

late arrival of the impulse at the distal A-V node will result in an inability to attain retrograde H_1H_2 intervals short enough to reach the refractory period of the retrograde fast pathway. It stresses the necessity to use ventricular pacing at high rates to expose dual retrograde A-V nodal pathways.

If one accepts the concept of the presence of two different pathways in anterograde and retrograde directions, a large combination of possibilities can theoretically exist depending on the electrophysiological properties of these pathways. Table 1 shows that at least six combinations are possible showing discontinuous conduction curves in both anterograde and retrograde direction. There are at least six other possible combinations with continuous curves in both anterograde and retrograde direction. This illustrates the complexity of the A-V node, which is further complicated by the possibility that a certain combination of electrophysiological properties of the A-V nodal pathways may change into another depending upon autonomic tone, basic pacing rate, drugs, etc. The ability to demonstrate dual A-V nodal pathways seems also related to the stamina of the investigator! Using several basic cycle lengths and different atrial stimulation sites will increase the chance of showing the presence of dual pathways^[15]. As demonstrated in our patient 2, subnodal delay during ventricular pacing may

result in failure to prove the presence of two retrograde pathways. As shown, this subnodal delay can be overcome by increasing the basic pacing rate during ventricular stimulation. We have selected our two examples to show that in some patients with A-V nodal tachycardia (as in patient 1) dual A-V nodal pathways can only be demonstrated indirectly, while in others they are readily apparent both during atrial and ventricular pacing (patient 2). The concept that pathways with different electrophysiological properties are present in the A-V node is supported by observations from animal studies^[16]. The animal studies suggest that acceptance of a two pathway concept might be an oversimplification. Availability of several A-V nodal pathways in anterograde and retrograde direction makes the theoretical number of possible combinations for reentry in the A-V node almost infinite. We believe that this complex situation may account for the differences in findings between different groups of investigators^[1, 2, 6, 8, 13, 14, 17-20]. It also helps in understanding the importance of changes in autonomic tone for initiation of A-V nodal tachycardia^[1, 2, 4, 21, 22], and it might explain some of the sometimes conflicting results following drug administration^[1, 6, 7, 23-25].

References

- [1] Rosen KM, Denes P, Wu D, Dhingra RC. Electrophysiological diagnosis and manifestation of dual A-V nodal pathways. In: Wellens HJJ, Lie KI, Janse MJ, eds. The conduction system of the heart. Leiden: HE Stenfert Kroese BV, 1976: 453-66.
- [2] Wu D, Denes P, Amat-y-Leon F, Wyndham CRC, Dhingra R, Rosen KM. An unusual variety of atrioventricular nodal reentry due to retrograde dual atrioventricular nodal pathways. *Circulation* 1977; 56: 50-9.
- [3] Wellens HJJ. Unusual examples of supraventricular reentrant tachycardias. *Circulation* 1975; 51: 997-1002.
- [4] Wellens HJJ, Durrer D. The role of an accessory atrioventricular pathway in reciprocal tachycardia. *Circulation* 1975; 52: 58-72.
- [5] Wellens HJJ, Bär FW, Farré J, et al. Diagnosis and incidence of concealed accessory pathways in patients suffering from paroxysmal A-V junctional tachycardia. *Rev Lat Cardiol* 1980 (in press).
- [6] Wu D, Wyndham C, Amat-y-Leon F, Denes P, Dhingra R, Rosen KM. The effects of ouabain on induction of atrioventricular nodal reentrant paroxysmal supraventricular tachycardia. *Circulation* 1975; 52: 201-7.
- [7] Wellens HJJ, Tan SL, Bär FWH, Düren DR, Lie KI, Dohmen HM. Effect of verapamil studied by programmed electrical stimulation of the heart in patients with paroxysmal reentrant supraventricular tachycardia. *Br Heart J* 1977; 39: 1058-66.

Table 1 Theoretical approach to the possible combinations of dual A-V nodal pathways. The combinations are made by changing the relation in duration of the effective refractory periods of the two pathways in the anterograde and retrograde directions

ABCD+++	BACD++	CABD+++	DABC+++
ABDC++	BADC+	CADB+++	DACB+++
ACBD+++	BCAD++	CBAD++	DBAC+
ACDB+++	BCDA++	CBDA++	DBCA+
ADCB++	BDAC+	CDAB+++	DCAB+++
ADBC++	BDCA+	CDBA++	DCBA+

+, Discontinuous A-V nodal conduction curves in both anterograde and retrograde direction; ++, discontinuous A-V nodal conduction curves only anterogradely; +++, discontinuous A-V nodal conduction curves only retrogradely; +, continuous A-V nodal conduction curves in both anterograde and retrograde direction. A, Anterograde effective refractory period of the fast pathway; B, anterograde effective refractory period of the slow pathway; C, retrograde effective refractory period of the fast pathway; D, retrograde effective refractory period of the slow pathway. In each column the duration of the four effective refractory periods are given. The duration lengths going from left to right.

- [8] Akhtar M, Damato AN, Ruskin JN, *et al.* Anterograde and retrograde conduction characteristics in three patterns of paroxysmal atrioventricular junctional reentrant tachycardia. *Am Heart J* 1978; 95: 22-42.
- [9] Ross D, Farre J, Bär FWHM, *et al.* Comprehensive clinical electrophysiological studies in the investigation of documented or suspected tachycardias. *Circulation* 1980; 61: 1010-6.
- [10] Wu D, Denes P, Wyndham C, Amat-y-Leon F, Dhingra RC, Rosen KM. Demonstration of dual atrioventricular nodal pathways utilizing a ventricular extrastimulus in patients with atrioventricular nodal reentrant paroxysmal supraventricular tachycardia. *Circulation* 1975; 52: 789-98.
- [11] Cagin NA, Kunstadt D, Wolfish P, Levit B. The influence of heart rate on the refractory period of the atrium and A-V conducting system. *Am Heart J* 1973; 85: 358-66.
- [12] Denes P, Wu D, Dhingra R, Pietras RJ, Rosen KM. The effects of cycle length on cardiac refractory periods in man. *Circulation* 1974; 49: 32-41.
- [13] Denes P, Wu D, Dhingra R, Amat-y-Leon F, Wyndham C, Rosen KM. Dual atrioventricular nodal pathways. A common electrophysiological response. *Br Heart J* 1975; 37: 1069-76.
- [14] Neuss H, Schlepper M, Spies HF, De Mey CH. Effects of heart rate and atropine on 'dual AV conduction'. *Br Heart J* 1975; 37: 1216-27.
- [15] Ross D, Farre J, Wiener I, Bär F, Vanagt E, Wellens H. Comparison of right v. coronary sinus stimulation in AV nodal reentry. *Circulation* 1979; 60: 63.
- [16] Janse MJ, van Capelle FJL, Anderson RH, Touboul P, Billette J. Electrophysiology and structure of the atrioventricular node of the isolated rabbit heart. In: Wellens HJJ, Lie KI, Janse MJ, eds. *The conduction system of the Heart*. Leiden: HE Stenfert Kroese BV, 1976: 296-315.
- [17] Touboul P, Huerta F, Porte J, Delahaye JP. Reciprocal rhythm in patients with normal electrocardiogram: evidence for dual conduction pathways. *Am Heart J* 1976; 91: 3-10.
- [18] Denes P, Wu D, Amat-y-Leon F, Dhingra R, Wyndham CR, Rosen KM. The determinants of atrioventricular nodal reentry with premature atrial stimulation in patients with dual A-V nodal pathways. *Circulation* 1977; 56: 253-9.
- [19] Pritchett ELC, Gallagher JJ, Wallace AG. Reentry within the atrioventricular node in man: a reassessment. *Eur J Cardiol* 1978; 6/6: 437-47.
- [20] Akhtar M. Paroxysmal atrioventricular nodal reentrant tachycardia. In: Narula OS, ed. *Cardiac arrhythmias*. Baltimore: The Williams and Wilkins Co., 1979: 294-317.
- [21] Akhtar M, Damato AN, Batsford WP, *et al.* Induction of atrioventricular nodal reentrant tachycardia after atropine. *Am J Cardiol* 1975; 36: 286-93.
- [22] Wu D, Denes P, Bauernfeind R, Dhingra R, Wyndham C, Rosen KM. Effects of atropine on induction and maintenance of atrioventricular nodal reentrant tachycardia. *Circulation* 1979; 59: 779-88.
- [23] Wu D, Denes P, Dhingra R, Khan A, Rosen KM. The effects of propranolol on induction of A-V nodal reentrant tachycardia. *Circulation* 1974; 50: 665-7.
- [24] Wellens HJJ, Duren DR, Liem KL, Lie KI. Effect of digitalis in patients with paroxysmal atrioventricular nodal tachycardia. *Circulation* 1975; 52: 779-88.
- [25] Wu D, Denes P, Bauernfeind R, Amat-y-Leon F, Rosen KM. Effects of procainamide on A-V nodal reentrant paroxysmal tachycardia. *Circulation* 1978; 57: 1171-9.

4.

Observations on Spontaneous Termination of Atrioventricular Nodal Reentrant Tachycardia

Pedro Brugada, MD; David Ross, FRACP*; Frits W.H.M. Bär, MD; Willem R.M. Dassen, PhD; Hein J.J. Wellens, MD, FACC.

Maastricht, The Netherlands.

Unusual mechanisms of spontaneous termination of atrioventricular (A-V) nodal reentrant tachycardia were observed in two patients during programmed electrical stimulation of the heart. In both patients the mechanism of termination was based on the use of another reentrant pathway than the one used during tachycardia. This pathway was located extra-nodally in one patient and intranodally in the other. The observations illustrate some of the complexities of reentry in the human heart and how they can play a role in spontaneous termination of A-V nodal tachycardia.

Observations in the animal heart^{1,2} and data from programmed electrical stimulation in the human heart^{3,4} support reentry as the most common mechanism for atrioventricular (A-V) nodal tachycardia. Reentry requires the presence of two interconnected pathways having different electrophysiologic properties. Termination of reentry occurs when refractory tissue halts the circulating impulse somewhere in the reentrant pathways. In this study we report observations on spontaneous termination of tachycardia in two patients with A-V nodal tachycardia, emphasizing how a third pathway that is connected to the reentry circuit can result in refractoriness in the reentry circuit and termination of the arrhythmia.

Methods

Patient 1 was a 20 year old man with A-V nodal tachycardia at a rate of 200/min. Patient 2 was a 70 year old man with A-V nodal tachycardia at a rate of 150/min. After the patients gave informed consent, they underwent an electrophysiologic study in the resting postabsorptive, nonsedated state. Neither patient was receiving medication at the time of study. With use of the femoral approach, catheters were placed for both stimulation and recording of intracardiac electrograms in the apex of the right ventricle, the bundle of His region, the high right atrium and the coronary sinus. Our stimulation protocol has previously been described.⁵ Both patients underwent a complete study. Only relevant data will be presented.

Criteria for the diagnosis of A-V nodal reentrant tachycardia: The diagnosis of A-V nodal tachycardia was based largely on exclusion of the use of an accessory A-V pathway as the retrograde limb of the tachycardia circuit and exclusion of an atrial site of origin of the tachycardia. The exclusion criteria were as follows: (1) Supraventricular tachycardia inducible by atrial premature beats related to A-V nodal delay.^{3-4,6} (2) Simultaneous activation of atria and ventricles during tachycardia.⁶ (3) H-A interval during tachycardia independent of H-V interval or bundle branch block pattern. (4) Persistence of tachycardia during A-V block distal to the bundle of His area (Patient 1). (5) Inability to advance atrial activation during tachycardia by ventricular premature beats given at the time of refractoriness of the bundle of His.⁷ (This observation does not exclude the presence of an accessory A-V pathway, but the opposite finding would prove its presence.⁸) (6) Ventriculoatrial conduction during ventricular pacing and after ventricular premature beats related to depolarization of the bundle of His⁸ and subnodal delay of ventricular premature beats.⁹ (7) Similar atrial activation

From the Department of Cardiology, University of Limburg, Annadal Hospital, Maastricht, The Netherlands. Manuscript received August 13, 1980; revised manuscript received October 22, 1980; accepted October 24, 1980.

* Overseas clinical fellow of the National Heart Foundation of Australia, Melbourne, Australia.

Address for reprints: Hein J. J. Wellens, MD, Department of Cardiology, University of Limburg, Annadal Hospital, Maastricht, The Netherlands.

FIGURE 1. Case 1. Atrioventricular (A-V) nodal tachycardia originally showing 2:1 block distal to the bundle of His. After recovery of 1:1 A-V conduction over the right bundle branch (third QRS complex, with a left bundle branch block configuration) subsequent atrial activation is advanced. When this pattern is repeated (fifth QRS complex) tachycardia is terminated. Indicated are the tachycardia cycle length in ms, the H-V interval preceding the widened QRS complexes and the V-A interval following these widened QRS complexes as measured in the distal coronary sinus (CSd) lead for clarity. Arrows point to possible retrograde His bundle potentials labeled H*. See text and Figure 2 for explanation. Leads I, II, III, V₁, and V₆ were simultaneously recorded with a high right atrial (HRA), a distal coronary sinus and a His bundle electrogram.

pattern during tachycardia and after retrograde conduction during ventricular pacing, suggesting that in both circumstances the same pathway was used for retrograde conduction.

Conventional definitions were used for refractory periods. The A-V nodal conduction curve was considered discontinuous if increasing the prematurity of the test pulse (atrial or ventricular) by 10 ms resulted in a sudden increase in A-V nodal conduction time by 50 ms or more. This sudden unexpected increase in A-V nodal conduction time of a premature beat suggests a shift in conduction from a fast to a slow A-V nodal pathway.^{1-4,6,7}

Results

Patient 1: Termination of A-V nodal tachycardia by bundle branch reentry: A-V nodal tachycardia with a cycle length of 295 ms originally showed 2:1 block distal to the bundle of His (Fig. 1). During recovery of 1:1 A-V conduction the first QRS complex shows a prolonged H-V interval (140 ms) and a complete left bundle branch block configuration. This QRS complex is followed by atrial activation with the sequence of atrial activation similar to the one recorded during tachycardia and occurring 45 ms earlier than expected, indicating that the atrial complex is not the result of regular impulse formation during tachycardia. The fifth QRS complex during tachycardia again shows a left bundle branch block pattern, but with an H-V interval of 115 ms. This QRS complex is followed by atrial activation occurring earlier than expected. After this atrial complex the tachycardia terminates.

FIGURE 2. Case 1. Diagram showing the proposed mechanism of termination of tachycardia in Figure 1. A = atrium; AVN = atrioventricular node; BB = bundle branches; F = fast A-V nodal pathway; LB = left bundle branch; RB = right bundle branch; S = slow A-V nodal pathway; V = ventricle. See text.

The H-A interval stays the same in spite of changing H-V intervals during the aberrant QRS complexes, or in spite of block of conduction to the ventricles at a site distal to the bundle of His. We postulate (Fig. 2) that during left bundle branch block the impulse traveling to the ventricles over the right bundle branch is able to invade in retrograde fashion the left bundle branch and reenters the A-V node. Retrograde conduction over the fast A-V nodal pathway allows premature activation of the atrium.

In Figures 1 and 3 possible retrograde His bundle potentials following the aberrant QRS complexes are identified. It was considered that they should precede the atrial echo by 80 ms or more (the H-A interval during tachycardia). An inverse relation is observed between the H-V interval and the V-H interval after the aberrant QRS complexes, (the V-H interval measures 100 ms after an H-V interval of 140 ms, and 110 ms after an H-V interval of 115 ms). The V-A interval following the aberrant complexes (as measured from the ventriculogram in the distal coronary sinus lead to the onset of atrial activation in the same lead) changes in the absence of changes in the bundle branch block pattern. This observation suggests that a free wall bypass tract is not the pathway used for retrograde conduction. Tachycardia is terminated depending on the retrograde and anterograde input in the A-V nodal pathways as reflected by the H-H and A-A intervals. After the first aberrant QRS complex the H-H interval measures 240 ms (Fig. 1) and the A-A interval 250 ms. This results in some degree of concealed penetration of the impulse in the slow A-V nodal pathway (Fig. 2) and collision with the anterograde front within this pathway, whereas retrograde conduction to the atria takes place over the fast A-V nodal pathway. The atrial complex is conducted in anterograde manner with a transient increase in the slow pathway conduction time. However, after the second aberrant QRS complex both the H-H interval (225 ms) and the A-A interval (240 ms) are shorter (Fig. 1). This shortening results in a major degree of concealed penetration of the impulse in the slow pathway that, in combination with a shorter A-A interval, results in block of impulse conduction in the slow pathway and termination of tachycardia (Fig. 1 and 2).

Occasionally right bundle branch block was observed during the change from 2:1 to 1:1 A-V conduction. This also resulted in the echo phenomenon (Fig. 3). However, because of the retrograde and anterograde input in the A-V nodal pathways

(H-H interval 240 ms and A-A interval 260 ms after the aberrant QRS complex) a fully compensatory pause followed and tachycardia continued. In this patient atrial premature beats delivered during atrial pacing at a basic cycle length of 600 ms did not result in aberrant conduction to the ventricles, the shortest attainable H₁-H₂ interval measuring 330 ms.

Patient 2. Termination of A-V nodal tachycardia by a sudden change in another anterograde A-V nodal pathway: An A-V nodal tachycardia with anterograde conduction over a "slow" pathway and retrograde conduction over a "fast" pathway (Fig. 4) was terminated spontaneously by a cycle that (1) was 115 ms shorter than the cycles seen during tachycardia, and (2) was not followed by retrograde conduction to the atria. This mode of spontaneous termination was repeatedly observed and probably can be explained by the A-V nodal conduction curves shown in Figure 5. During single test stimulation of the right atrium at a basic cycle length of 500 ms, several discontinuities in the A₂-H₂ curve are seen. This was also during stimulation with a basic cycle length of 460 ms. These findings support the presence of either more than one "slow" pathway or delay in a final common pathway. The former hypothesis is supported by the fixed H to atrial echo interval at the time of initiation of tachycardia irrespective of the preceding A-H interval. If a final common pathway were responsible for delay in conduction to the His bundle and the "jumps" in the conduction curve, a changing relation between His bundle activation and atrial echo would have been observed.

We postulate that in this patient tachycardia terminates spontaneously because of conduction over a "slow" pathway that is not slow enough to overcome retrograde refractoriness of the "fast" retrograde pathway (Fig. 6). Although the anterograde and retrograde refractory periods of the fast pathway need not be equal, the anterograde refractory period of the fast pathway measured 290 ms and the H-H interval terminating tachycardia measured 280 ms (a lesser value). After initiation of tachycardia (Fig. 6), we postulate a decreasing concealment in the third pathway that is finally relieved, allowing anterograde conduction over this pathway, the impulse colliding in the slow pathway with the anterograde front and blocking in the fast pathway because it is entering at a time when the pathway is still refractory.

Discussion

A third pathway with bundle branch reentry: In these two patients the findings strongly suggest that reentry into the tachycardia circuit over a third pathway was responsible for termination of the arrhythmia. This third pathway was not involved in the reentrant pathways during initiation and perpetuation of tachycardia. Atrial echo beats based on reentry over the bundle branches have been demonstrated to occur.¹⁰⁻¹² As in those studies, the atrial echoes in our Patient 1 required the presence of block in one of the bundle branches and sufficient delay in conduction over the other bundle branch to complete a reentrant cycle. The time of arrival of the impulse in the A-V nodal pathways (in both anterograde and retrograde directions as reflected by the H-H and A-A intervals following the aberrant complex) was crucial for termination of the reentrant movement. It seems unlikely that the widened QRS complexes represented ventricular extrasystoles occurring late in diastole. Ventricular premature beats were not observed to occur spontaneously throughout the investigation and, as shown, advancement of subsequent atrial activation was observed during both left and right bundle

FIGURE 3. Case 1. A-V nodal tachycardia with 2:1 block distal to the bundle of His showing recovery of conduction over the left bundle branch (second QRS complex, with a right bundle branch block configuration). Subsequent atrial activation is advanced but tachycardia continues. Same leads and labeling as in Figure 1. See text.

branch block-shaped QRS complexes. The range of H-V intervals preceding these widened QRS complexes is in keeping with previous clinical observations¹⁰ and the results of experimental studies.¹²

An inverse relation between H-V and V-H intervals during the widened QRS complexes was demonstrated in our Patient 1, supporting the concept of bundle branch reentry.¹⁰ In this patient we could exclude the presence of an extranodal accessory pathway. Depolarization of the bundle of His occurred twice during H-H intervals of 240 to 225 ms (Fig. 1 and 3). Experimental studies¹² have shown that retrograde depolarization of the His bundle can occur as soon as 160 ms after its anterograde depolarization. The observation of bundle branch block with slow conduction over the contralateral branch with advancement of subsequent atrial activation makes bundle branch reentry the most plausible explanation for termination of tachycardia in this patient. Unfortunately, bundle branch block and atrial echoes were not observed during atrial premature beats delivered during pacing of the right atrium at a basic cycle length of 600 ms. Their absence might have occurred because the H-H interval of 300 ms during tachycardia could result in concealment in the bundle branch system, allowing development of bundle branch block patterns, whereas such concealed conduction presumably did not occur during atrial pacing at a basic cycle length of 600 ms.

Anterograde intranodal conduction over a third A-V nodal pathway: In Patient 2 a sudden shortening in the cycle length of reentry was followed by termina-

tion of tachycardia. The degree of shortening is too great to be accounted for by a sudden change in autonomic tone. The most attractive hypothesis to explain this phenomenon is anterograde intranodal conduction over a different A-V nodal pathway with slow conduction time. The finding of several discontinuities in the A-V nodal conduction curve might support this theory. It is of interest that the A-H interval after the first break in

the A-V nodal conduction curve is similar to the A-H interval before termination of tachycardia, measuring 230 and 215 ms, respectively (Fig. 4 and 5). We cannot exclude the possibility that a His bundle extrasystole was responsible for termination of tachycardia and the apparent shortening of the A-H interval. Also we do not know the validity of comparing the A-H interval terminating tachycardia with the A-H interval after the

FIGURE 4. Case 2. A-V nodal tachycardia with a cycle length of 395 ms. Tachycardia terminates after sudden shortening in the H-H interval by 115 ms, corresponding to 115 ms shortening in the A-H interval. This repeatedly observed phenomenon was not preceded by any changes in time relations during the tachycardia. Leads V₁ and V₆ were simultaneously recorded with four endocavitary electrograms. CS_p = Proximal coronary sinus electrodegram.

FIGURE 5. Case 2. Relation between the A₁-A₂ intervals (abscissa) during pacing of the high right atrium at basic cycle lengths of 500 and 460 ms. Three discontinuities in the A-V nodal conduction curve were observed during atrial pacing with a basic cycle length of 500 ms (left), and two when the basic cycle length of pacing was decreased to 460 ms (right). TC = atrial premature beats initiating tachycardia.

FIGURE 6. Case 2. Diagram illustrating the proposed mechanism of termination of tachycardia as observed in Figure 4. α , β and γ represent three A-V nodal pathways. In panel 1, an atrial premature beat (APB) is blocked in pathways β and γ and initiates tachycardia after anterograde conduction over pathway α (the slowest conducting pathway) and retrograde conduction over pathway β (the fastest pathway). Pathway γ is maintained refractory because of concealed anterograde and retrograde penetration of impulses during tachycardia. In panel 2, decreasing concealed penetration in pathway γ in the retrograde direction is postulated. In panel 3, after recovery of conduction in the anterograde direction in pathway γ the impulse is conducted to the ventricles with a shorter than expected A-H interval. The impulse blocks in pathway β because of refractoriness of this pathway, and collides in pathway α with the anterograde impulse, resulting in termination of tachycardia. Abbreviations as in Figure 2.

first break in the A-V nodal conduction curve of the left graph from Figure 5 when cycle lengths are different. However, we believe that the reproducibility of our observations, the several discontinuities in the A-V nodal conduction curve and the mode of termination of tachycardia suggest the availability of several A-V nodal pathways. The availability of several A-V nodal pathways has already been advanced from animal studies.² In Patient 2 a similar A-V nodal conduction time was observed not to initiate or to terminate tachycardia, thus suggesting the use of a third A-V nodal pathway that was not slow enough to overcome refractoriness of the retrograde A-V nodal pathway used during tachycardia.

Implications: In these two patients, sudden unexpected conduction over another reentrant pathway accounted for termination of the arrhythmia. Why these

pathways were suddenly able to conduct is difficult to explain. Although, the factors leading to the new reentrant pathway terminating tachycardia could be recognized in Patient 1, they remain unclear in Patient 2. One can speculate that differences in degree of concealed penetration of the impulse in potential reentrant circuits or changes in refractoriness and conduction velocity during tachycardia might have played a role. However, our present methods of investigation in the intact human heart are not refined enough to uncover most of these complex mechanisms.

Acknowledgment

We gratefully acknowledge the many helpful suggestions made by Mark E. Josephson, MD during the preparation of the manuscript.

References

1. Moe GK, Mendez C. The physiological basis of reciprocal rhythm. *Prog Cardiovasc Dis* 1966;8:461-82.
2. Janse MJ, van Capelle FJL, Anderson RH, Touboul P, Billewicz J. Electrophysiology and structure of the atrioventricular node of the isolated rabbit heart. In: Wellens HJJ, Lie KI, Janse MJ, eds. *The Conduction System of the Heart*. Philadelphia: Lea & Febiger, 1976:296-315.
3. Goldreyer BN, Damato AN. The essential role of atrioventricular conduction delay in the initiation of paroxysmal supraventricular tachycardia. *Circulation* 1971;43:679-87.
4. Rosen KM, Mehta A, Miller RA. Demonstration of dual atrioventricular nodal pathways in man. *Am J Cardiol* 1974;33:291-4.
5. Ross D, Farré J, Bar FWHM, et al. Comprehensive electrophysiological studies in the investigation of documented or suspected tachycardias. *Circulation* 1980;61:1010-4.
6. Wellens HJJ, Durrer D. The role of an accessory atrioventricular pathway in reciprocal tachycardia. *Circulation* 1975;52:58-71.
7. Wu D, Denes P, Wyndham C, Amat-y-Leon F, Dhingra RC, Rosen KM. Demonstration of dual atrioventricular nodal pathways utilizing a ventricular extrastimulus in patients with atrioventricular nodal reentrant paroxysmal supraventricular tachycardia. *Circulation* 1975;52:789-98.
8. Akhtar M, Damato AN, Batsford WP, Ruskin JN, Ogunkelu JB. A comparative analysis of antegrade and retrograde conduction patterns in man. *Circulation* 1975;52:766-78.
9. Brugada P, Vanagt EJ, Bar FWHM, Wellens HJJ. Incessant reciprocating atrioventricular tachycardia. *Pace* 1980;3:670-7.
10. Varghese PJ, Damato AN, Caracta AN, Gallagher JJ, Josephson ME, Lau SH. Intraventricular conduction delay as a determinant of atrial echo beats. *Circulation* 1974;49:805-10.
11. Castellanos A, Embl A, Aranda J, Befeler B. Retrograde His bundle deflection in bundle branch re-entry. *Br Heart J* 1976;38:301-3.
12. Moe GK, Mendez C, Han J. Aberrant A-V impulse propagation in the dog heart: a study of functional bundle branch block. *Circ Res* 1965;16:261-86.

5.

Initiation of atrio-ventricular nodal re-entrant tachycardia in patients with discontinuous anterograde atrio-ventricular nodal conduction curves with and without documented supraventricular tachycardia. Observations on the role of a discontinuous retrograde conduction curve.

Pedro Brugada, M.D.; Bill Heddle, M.B., B.S.; Frits W.H.M. Bär, M.D.; and Hein J.J. Wellens, M.D.

From the Department of Cardiology, University of Limburg, Annadal Hospital, Maastricht, The Netherlands.

Summary

To evaluate factors playing a role in initiation of atrioventricular (A-V) nodal re-entrant tachycardia utilizing anterogradely a slow and retrogradely a fast conducting A-V nodal pathway, thirty-eight patients having no accessory pathways and showing discontinuous anterograde A-V nodal conduction curves during atrial stimulation were studied. Twenty-two patients (group A) underwent an electrophysiologic investigation because of recurrent paroxysmal supraventricular tachycardia that had been electrocardiographically documented before the study. Sixteen patients (group B) underwent the study because of a history of palpitations (15 patients) or recurrent ventricular tachycardia (1 patient). In none of them a supraventricular tachycardia had been ever electrocardiographically documented before the investigation.

Twenty-one of the twenty-two patients of group A demonstrated continuous retrograde conduction curves during ventricular stimulation. In twenty tachycardia was initiated by either a single atrial premature beat (18 patients) or two atrial premature beats (2 patients). The remaining two patients received atropine and tachycardia could then be initiated by a single atrial premature beat.

Fifteen of the sixteen patients of group B had discontinuous retrograde conduction curves during ventricular stimulation, with a long refractory period of their retrograde fast pathway. Tachycardia was initiated by multiple atrial premature beats in one patient. Thirteen out of the remaining fifteen patients received atropine. Thereafter tachycardia could be initiated in three patients by a single atrial premature beat, by two atrial premature beats in one patient, and by incremental atrial pacing in another one. In the remaining eight patients tachycardia could not be initiated.

This study demonstrates that in patients having discontinuous anterograde A-V nodal conduction curves and clinical documentation of supraventricular tachycardia, the refractory period of their retrograde fast pathway is short, and tachycardia can easily be initiated during atrial stimulation. Only a few patients require administration of atropine to facilitate initiation of the arrhythmia. Patients without documentation of supraventricular tachycardia have a long refractory period of their retrograde fast pathway, and tachycardia is not likely to be initiated during atrial stimulation. Administration of atropine however can in some of these patients result in shortening of the refractory period of the retrograde fast pathway and initiation of tachycardia. Our observations indicate that the pattern of ventriculo-atrial conduction found during ventricular stimulation is a marker for ease of initiation of A-V nodal tachycardia in patients with discontinuous anterograde A-V nodal conduction curves.

Introduction

Experimental and clinical studies indicate that longitudinal dissociation of the atrio-ventricular (A-V) node leading to functionally different pathways plays a role in supraventricular tachycardia in man (1-8). In several patients with supraventricular tachycardia in whom the presence of an accessory pathway is excluded during the electrophysiological investigation a re-entry circuit confined to the A-V node can be demonstrated (9). These patients usually show discontinuous anterograde A-V nodal conduction curves during atrial stimulation suggesting the presence of dual A-V nodal pathways. Initiation of tachycardia is related to a shift in anterograde conduction from a fast to a slowly conducting A-V nodal pathway. Some studies have shown however absence of initiation of tachycardia in the presence of a discontinuous anterograde conduction curve (10-12). In these studies the possible role of a long refractory period of the retrograde fast pathway preventing initiation of tachycardia was not systematically studied. Also how a drug that induces changes in the properties of the A-V nodal pathways may facilitate initiation of tachycardia was not analysed.

In the present study the results of programmed electrical stimulation of the heart are reported in 38 patients showing discontinuous anterograde A-V nodal conduction curves during atrial stimulation in the absence of accessory atrio-ventricular pathways. Twenty-two patients had electrocardiographic documentation of supraventricular tachycardia before the investigation and sixteen patients not. The results of this study emphasize the role of the refractory period of the retrograde fast pathway in initiation of A-V nodal tachycardia during atrial stimulation. They also show why the administration of atropine can result in initiation of tachycardia.

Patients and methods

Consecutive patients were included in the present study when they met the following criteria:

- a. A complete electrophysiologic study using a previously described protocol of stimulation, recording and analysis of tracings (13). During the study leads I, II, III, V1 and V6 were simultaneously recorded with bipolar endocavitary electrograms from the right atrium, His bundle area, coronary sinus or left atrium.
- b. Demonstration of discontinuous anterograde A-V nodal conduction curves during single test stimulation of the atria at at least two of three different basic cycle lengths of pacing. Discontinuous anterograde conduction curves of the A-V node were diagnosed when a 50 ms or more increase in the A-V nodal conduction time occurred on increasing the prematurity of the atrial premature beat by 10 ms (4), and by the method described by Rosen et al (1).
- c. Presence of ventriculo-atrial conduction during ventricular pacing.
- d. No evidence of accessory atrio-ventricular pathways.

The extrastimulus technique was used during atrial and ventricular stimulation to determine the patterns and refractory periods of A-V and V-A conduction. Anterograde A-V nodal conduction was studied by plotting the atrial premature beat intervals (A1-A2) against their A-V nodal conduction time (the A2-H2 interval and the H1-H2 interval). The same cycle lengths used during atrial stimulation were used during ventricular stimulation. Retrograde conduction curves were analysed by plotting the ventricular premature intervals (V1-V2) against their ventriculo-atrial conduction time (V2-A2 and A1-A2 interval). Care was taken to identify when possible retrograde His bundle potentials both during basic paced ventricular beats and following ventricular premature beats. When both could be identified the retrograde H1-H2 intervals were taken as the premature beat interval of the retrograde input into the A-V node. An approximation of the retrograde input into the A-V node was obtained by measuring the S1-H2 interval when

retrograde activation of the His bundle was only observed following the ventricular premature beat (V2) (14). This interval results in an overestimation of the interval of the retrograde input into the A-V node. In all patients in whom a discontinuous retrograde conduction curve was diagnosed during ventricular stimulation the initiation of A-V nodal echoes using retrogradely a slow and anterogradely a fast pathway was required to support the A-V nodal site of the discontinuity in the retrograde conduction curve, including those in whom retrograde His bundle potentials were readily identified both during ventricular pacing and following the ventricular premature beat.

An accessory atrio-ventricular pathway was excluded, and the diagnosis of the A-V node as site of origin of tachycardia was made by using previously reported criteria (4,15).

Informed consent was obtained from each patient. All medication was stopped at least 72 hours before the study.

Results

a. Clinical data.

Thirty-eight out of 350 consecutive patients studied by programmed electrical stimulation of the heart in our laboratory fulfilled the criteria for inclusion in the study. Their clinical data are given in table I. In twenty-two patients a paroxysmal recurrent supraventricular tachycardia had been electrocardiographically documented before the investigation (group A). On analyzing the relation between P waves and QRS complexes during tachycardia (16,17) the documented arrhythmia was considered to be A-V nodal in origin in these 22 patients. Of the remaining 16 patients one was studied because of an electrocardiographically documented recurrent ventricular tachycardia on the basis of chronic coronary artery disease, but he never had an electrocardiogram showing a supraventricular arrhythmia. The remaining 15 patients were studied because of a history of recurrent palpitations. In all of them multiple electrocardiograms had failed to document a supraventricular tachycardia. In each patient a least one out of hospital 24 hours electrocardiographic recording had been made. Additional monitoring of variable duration (6-48 hours) was performed in hospital in all these sixteen patients (group B). Although supraventricular and ventricular premature beats were registered, in none of them a supraventricular tachycardia was documented. The history of complaints, ranged from 0.3 to 39 years in patients of group A and from 1 to 30 years in patients of group B. Excluding the patient of group B with ventricular tachycardia the mean duration of symptoms was longer in patients of group A (16.3 ± 12.7 years) than in patients of group B (9.6 ± 8.4 years). These results were statistically significant ($p < 0.05$) (Student's t test). There was a wide variation in frequency of complaints and in duration of episodes both inter and intraindividually, and these data were not analysed.

b. Electrophysiological studies (table II):

During single test stimulation of the atria during atrial pacing all 38 patients demonstrated the presence of discontinuous anterograde A-V nodal conduction curves at at least two different basic cycle lengths of pacing. In patients of group A the anterograde refractory period of the fast A-V nodal pathway ranged from 240 to 500 ms, and in group B from 250 to 560 ms. These results were not significantly different. Also the cycle lengths of pacing used in each group were not significantly different. In most patients, both in group A and B, the anterograde refractory period of the slow pathway could not be exactly determined, because conduction over this pathway was present up to the refractory period of the atrium. Statistical analysis of these values was not performed.

During single test stimulation of the ventricle during ventricular pacing twenty-one patients of group A demonstrated (fig. 1) a continuous retrograde conduction curve with conduction to the atrium over a fast A-V nodal pathway. In most instances ventriculo-atrial conduction occurred up to the refractory period of the right ventricle. There was one patient (patient 22) in whom a

Table I

Clinical data of thirty-eight patients studied

<i>Patient</i>	<i>Age(yr)/sex</i>	<i>Indication for study</i>	<i>ECG documentation of SVT</i>	<i>Heart Disease</i>
Group A:				
1	57 F	Paroxysmal SVT	+	-
2	54 F	Paroxysmal SVT	+	-
3	65 F	Paroxysmal SVT	+	-
4	46 M	Paroxysmal SVT	+	-
5	50 M	Paroxysmal SVT	+	-
6	23 M	Paroxysmal SVT	+	-
7	35 M	Paroxysmal SVT	+	-
8	30 F	Paroxysmal SVT	+	-
9	30 F	Paroxysmal SVT	+	-
10	52 F	Paroxysmal SVT	+	-
11	71 F	Paroxysmal SVT	+	-
12	43 M	Paroxysmal SVT	+	-
13	62 M	Paroxysmal SVT	+	-
14	49 F	Paroxysmal SVT	+	-
15	28 F	Paroxysmal SVT	+	-
16	78 F	Paroxysmal SVT	+	CAD
17	24 M	Paroxysmal SVT	+	-
18	80 F	Paroxysmal SVT	+	CAD
19	24 F	Paroxysmal SVT	+	-
20	49 F	Paroxysmal SVT	+	-
21	44 F	Paroxysmal SVT	+	RHD
22	21 M	Paroxysmal SVT	+	-
Group B:				
23	15 M	Palpitations	-	-
24	55 F	Palpitations	-	-
25	40 M	Palpitations	-	-
26	30 M	Palpitations	-	-
27	45 M	Palpitations	-	-
28	25 F	Palpitations	-	-
29	38 M	Palpitations	-	-
30	23 M	Palpitations	-	-
31	60 M	Palpitations	-	-
32	55 F	Palpitations	-	-
33	43 M	Palpitations	-	-
34	52 M	Recurrent VT	-	CAD
35	29 M	Palpitations	-	-
36	36 F	Palpitations	-	-
37	53 M	Palpitations	-	-
38	49 M	Palpitations	-	-

Abbreviations: SVT: Supraventricular Tachycardia; VT: Ventricular Tachycardia; F: Female; M: Male; CAD: Coronary Artery Disease; RHD: Rheumatic Heart Disease. CCM: Congestive cardiomyopathy; yr: years; -: absent, +: present.

Figure 1
 Patient 12. Representative example of findings in group A patients during ventricular stimulation studies. During ventricular pacing with a basic cycle length of 600 ms ventriculo-atrial conduction over a retrograde fast A-V nodal pathway is present up to the refractory period of the ventricle. The top panel shows a ventricular premature beat given after 240 ms which is retrogradely conducted to the atria with a normal retrograde atrial activation sequence and preceded by a retrograde His bundle depolarization. The H2A2 interval measures 70 ms, indicating conduction over a retrograde fast pathway. In the bottom panel a premature ventricular stimulus given 10 ms earlier reaches the refractory period of the ventricle. Five surface leads are shown simultaneously with four endocavitary electrograms of the right atrium (RA), the proximal (p) and distal (d) coronary sinus (CS) and the His bundle (His).

Figure 2

Patient 30. Typical example of findings in group B patients during ventricular stimulation before administration of atropine. In panel A pacing of the right ventricular apex with a basic cycle length of 600 ms results in retrograde conduction to the atria over a slow A-V nodal pathway. (Note the retrograde His bundle potential at the end of the ventricular electrogram in the His bundle lead (H1)). A Wenckebach sequence of conduction over the slow pathway, with prolongation of the S1A1 interval is observed after the second paced ventricular complex. The first atrial complex shown is of sinus node origin. These findings indicate a long retrograde refractory period of the fast pathway. In panel B the right atrium and the right ventricle are paced simultaneously with the same basic cycle length. A premature beat is given in the ventricle but not in the atrium. This allows retrograde conduction of the ventricular premature beat to the atria with marked subnodal delay over a slow

A-V nodal pathway. An A-V nodal echo using anterogradely a fast A-V nodal pathway is observed. During retrograde conduction to the atria over the slow pathway the typical sequence of atrial activation is observed (18,19), with atrial activation recorded first in the proximal coronary sinus lead. Five surface leads are shown recorded simultaneously with 5 endocavitary bipolar electrograms (RV = right ventricular lead).

discontinuity in the retrograde conduction curve was present. In him the A-V node as the only retrograde pathway could be demonstrated by the initiation of the "reversed" form of A-V nodal echoes (fast-slow) after the occurrence of the discontinuity in the retrograde conduction curve. In these twenty-one patients the retrograde refractory period of the fast pathway varied individually from less than 190 ms to 360 ms. In the remaining patient of group A having discontinuous retrograde conduction curves the retrograde fast pathway refractory period measured ≥ 750 ms.

During single test stimulation of the ventricle during ventricular pacing a discontinuous retrograde conduction curve in the A-V node followed by initiation of "reversed" A-V nodal echoes was demonstrated in 14 patients of group B. This occurred at the same cycle lengths as the ones used during atrial stimulation (fig. 2). Patient 25 had discontinuous retrograde conduction curves with initiation of "reversed" A-V nodal echoes only at one basic cycle length of pacing, and patient 37 demonstrated continuous retrograde conduction curves at all basic cycle lengths tested.

In all patients the "reversed" form of re-entry in the A-V node terminated spontaneously after 1 to 3 cycles, except for one patient in whom once a maximum of six cycles was observed. Using previously reported criteria (18) we excluded the presence of an accessory pathway with a long conduction time in all patients showing discontinuous retrograde conduction curves.

Initiation of the common type of re-entry in the A-V node (using anterogradely a slow and retrogradely a fast A-V nodal pathway) was possible in the majority of patients of group A by giving a single atrial premature beat during atrial stimulation (table II, fig. 3). Two patients (patient 5 and patient 13) required two atrial premature beats. In two patients (patients 10 and 22) single or multiple atrial premature beats, and incremental atrial pacing failed to initiate re-entry in the A-V node in the basal condition. In all patients of group A in whom the retrograde refractory period of the fast pathway could be measured during ventricular stimulation, the A-H interval over the slow pathway at the time of initiation of re-entry in the A-V node during atrial stimulation was similar to or longer than the retrograde refractory period of the fast pathway (table II). In all patients of group A where re-entry in the A-V node could be initiated during atrial stimulation a sustained intra-A-V nodal tachycardia followed (fig. 1 and 3).

In patients of group B a sustained A-V nodal tachycardia using anterogradely a slow and retrogradely a fast pathway was initiated in the basal state during atrial stimulation in one patient (patient 28) by delivering three consecutive atrial premature beats during atrial stimulation. In 4 patients (patients 25, 26, 28 and 37) only a single atrial echo of A-V nodal origin was initiated by a single atrial premature beat during atrial stimulation. This echo was usually followed by anterograde conduction to the ventricles. Thereafter re-entry in the A-V node terminated. Except for patient 28, multiple atrial premature beats and incremental atrial pacing did not result in initiation of sustained A-V nodal tachycardia in the basal condition. The maximal duration of the A-H intervals over the slow pathway during atrial stimulation in the patients of group B showing no A-V nodal re-entry was similar to those in patients from group A and B in whom re-entry in the A-V node could be initiated. However, as shown in table II, in patients in whom re-entry in the A-V node could not be initiated during atrial stimulation, the values of the maximal A-H intervals over the slow pathway were usually far below the value of the retrograde refractory period of the fast pathway when measured during ventricular stimulation (fig. 2 and fig. 4). This finding suggests that conduction over the slow pathway was not slow enough to overcome refractoriness of the retrograde pathway.

c. Administration of atropine:

In two patients of group A, and in thirteen patients of group B 1 mg of atropine was given intravenously. In the two patients of group A (table III) a sustained A-V nodal tachycardia using

Figure 3

Same patient as in figure 1. During pacing of the right atrium with a basic cycle length of 600 ms an atrial premature beat given after 330 ms is conducted to the ventricles with an A2H2 interval of 30 ms and an H1H2 interval of 380 ms (panel A). Shortening of the atrial premature beat interval by 10 ms (panel B) results in marked prolongation of both the A2H2 (340 ms) and H1H2 (580 ms) interval. No echo follows, however. In

panel C further shortening of the atrial premature beat interval to 270 ms results in initiation of A-V nodal tachycardia when a critical conduction time over the slow A-V nodal pathway is reached (A2H2 = 380 ms). Three surface leads are shown simultaneously with three endocavitary electrograms.

Table II

Summary of electrophysiologic data on thirty-eight patients studied, during basal condition

Pt	BCL	ERPfa	ERPsa	ERPfr	shortest A-H over Sa at time of echoes')	Mode of initiation of tachycardia
Group A						
1	600	350	< AFRP	≤ 250	445	1 APB
	500	360	< AFRP	320	450	
2	600	300	< AFRP	≤ 220	320	1 APB
	500	320	< AFRP	≤ 210	330	
3	600	430	340	≤ 270	260	1 APB
	500	450	340	≤ 260	230	
4	600	250	< AFRP	≤ 220	300	1 APB
	500	260	< AFRP	≤ 210	290	
5	600	260	< AFRP	≤ 240	200	2 APB's
	500	250	< AFRP	≤ 230	200	
6	600	290	< AFRP	≤ 230	200	1 APB
	500	310	< AFRP	≤ 230	200	
7	600	340	< AFRP	250	269	1 APB
	500	360	< AFRP	≤ 240	290	
8	460	260	< AFRP	≤ 190	220	1 APB
	400	240	< AFRP	≤ 190	230	
9	600	270	220	≤ 250	400	1 APB
	500	300	< AFRP	240	290	
10	600	310	< AFRP	250	—	—
	500	280	< AFRP	≤ 220	—	—
11	600	500	< AFRP	≤ 260	200	1 APB
	500	470	< AFRP	≤ 230	270	
12	600	320	< AFRP	≤ 320	380	1 APB
	500	360	300	≤ 330	360	
13	550	280	240	≤ 210	260	2 APB's
	500	270	260	≤ 200	260	
14	600	390	200	≤ 270	250	1 APB
	500	390	< AFRP	≤ 260	250	
15	600	350	< AFRP	300	350	1 APB
	500	340	< AFRP	250	300	
16	600	380	330	≤ 250	250	1 APB
	500	430	350	≤ 240	400	
17	750	340	320	≤ 240	410	1 APB
	600	420	340	≤ 220	350	
18	680	280	< AFRP	≤ 220	360	1 APB
	600	290	> AFRP	≤ 210	290	
19	600	320	280	≤ 220	—	1 APB
	500	290	240	360	390	
20	500	310	< AFRP	≤ 210	250	1 APB
	450	320	270	≤ 200	270	
21	600	290	< AFRP	≤ 230	300	1 APB
	500	340	< AFRP	≤ 210	320	
22	750	320	< AFRP	≥ 750	—	—
	600	350	< AFRP	≥ 600	—	

Table II (Cont)

<i>Pt</i>	<i>BCL</i>	<i>ERPfa</i>	<i>ERPsa</i>	<i>ERPfr</i>	<i>longest possible A-H over Sa</i>	<i>echoes') initiation</i>	<i>Mode of of tachycardia</i>
23	680	250	< AFRP	500	220	-	-
	600	270	< AFRP	500	280	-	
24	680	310	< AFRP	630	340	-	-
	600	290	< AFRP	> 600	320	-	
25	600	320	< AFRP	≤ 250	530	-	-
	500	360	300	370	500	+	
26	750	370	290	410	260	-	-
	600	430	280	290	270	+	
27	600	400	< AFRP	340	330	-	-
	500	420	< AFRP	390	310	-	
28	750	320	< AFRP	350	310	-	M APB
	600	350	< AFRP	300	320	+	
29	600	430	< AFRP	370	320	-	-
	500	420	< 310	340	330	-	
30	600	430	< AFRP	> 600	440	-	-
	500	400	< AFRP	> 500	410	-	
31	600	300	< AFRP	350	300	-	-
	500	280	< AFRP	360	310	-	
32	600	410	< AFRP	350	290	-	-
	450	420	< AFRP	360	290	-	
33	600	300	< AFRP	380	380	-	-
	500	310	270	410	400	-	
34	680	560	320	≥ 680	330	-	-
	600	560	280	≥ 600	470	-	
35	600	460	340	550	380	-	-
	500	450	330	≥ 500	400	-	
36	600	310	< AFRP	460	300	-	-
	430	250	< AFRP	260	260	-	
37	600	280	< AFRP	≤ 220	240	+	-
	500	290	< AFRP	≤ 220	290	+	
38	900	350	< AFRP	580	300	-	-
	750	370	< AFRP	440	300	-	

Abbreviations: ERP: Effective Refractory Period; fa: anterograde fast A-V nodal pathway; sa: anterograde slow A-V nodal pathway; fr: retrograde fast A-V nodal pathway; AFRP: Functional Refractory Period of the Atrium; APB: Atrial Premature Beat; '): A-V nodal echoes slow-fast; MAPB: Multiple Atrial Premature Beats. -: not initiated; +: initiated.

Figure 4

Same patient as in figure 2. During pacing of the right atrium with a basic cycle length of 600 ms an atrial premature beat given after 440 ms is conducted to the ventricles with an A2H2 interval of 170 ms and a H1H2 interval of 500 ms (panel A). Shortening of the atrial premature beat interval by 10 ms to 430 ms (panel B) results in an increase of the A2H2 interval by 50 ms to 220 ms, the H1H2 interval now measuring 540 ms. In panel C the atrial premature beat interval has been shortened to 260 ms, the A2H2 interval measuring 440 ms and the H1H2 interval 590 ms. In spite of this very long A2H2 interval, no A-V nodal echo follows, presumably because it is unable to overcome refractoriness of the retrograde pathway. In panel D an atrial premature stimulus given after 240 ms reaches the refractory period of the right atrium. Three surface leads are shown simultaneously with 5 endocavitary bipolar electrograms.

anterogradely a slow and retrogradely a fast pathway could then be initiated by a single atrial premature beat given during atrial stimulation. In both patients the A-V nodal conduction curves were still discontinuous in the anterograde direction. In patient 22 the retrograde conduction curves which were discontinuous before atropine administration became continuous following administration of the drug. The A-H intervals over the slow pathway at the time of initiation of tachycardia were not longer after atropine as compared to the A-H intervals over the slow pathway before atropine. These data indicate that atropine facilitated initiation of A-V nodal tachycardia by shortening of the retrograde refractory period of the fast pathway.

In 5 patients of group B initiation of a sustained A-V nodal tachycardia became possible after administration of atropine (table III, fig. 5-6). In three patients following a single atrial premature beat during atrial stimulation, in 1 patient after two consecutive atrial premature beats and in the remaining patient during incremental atrial pacing. In these 5 patients, similar to the two patients of group A in whom tachycardia could be initiated after atropine, the A-H intervals over the slow pathway at the time of initiation of tachycardia did not become longer after atropine (fig. 5-6).

Table III

Effects of atropine on the A-V nodal conduction curves and on initiation of A-V nodal tachycardia using anterogradely a slow and retrogradely a fast pathway

<i>Patient</i>	<i>Anterograde A-V nodal conduction curves</i>	<i>Retrograde A-V nodal conduction curves</i>	<i>Mode of initiation of tachycardia</i>
Group A			
10	discontinuous	continuous	1 APB
22	discontinuous	continuous	1 APB
Group B			
23	continuous	continuous	—
24	continuous	absent V-A conduction	—
25	discontinuous	continuous	IAP
26	discontinuous	continuous	-')
27	discontinuous	continuous	2 APB's
29	discontinuous	continuous	1 APB
30	continuous	continuous	—
31	discontinuous	continuous	1 APB
33	continuous	continuous	—
35	discontinuous	?	—
36	discontinuous	continuous	1 APB
37	continuous	continuous	—
38	continuous	continuous	—

Abbreviations: Same as table II. ? : Unknown; —: not initiated; ') Maximal A-H interval over slow pathway shortened from 400 ms to 200 ms after atropine; IAP: Incremental Atrial Pacing.

In these patients initiation of tachycardia was facilitated by shortening of the retrograde refractory period of the retrograde fast pathway. In all 5 patients the discontinuity in the retrograde conduction curve had disappeared while the characteristic discontinuous anterograde A-V nodal conduction curve was maintained. All 5 patients had initiation of the "common" form of A-V nodal tachycardia (slow-fast) after atropine. In the remaining eight patients any attempt to initiate an A-V nodal tachycardia failed. In none of them we were able to initiate re-entry in the A-V node of the "common" or "reversed" variety. Analysis of anterograde and retrograde conduction curves of the A-V node in these 8 patients showed a continuous retrograde conduction curve both in the anterograde and retrograde direction in 5 patients (fig. 7-8), with inability to shift conduction from the fast to the slow A-V nodal pathway during atrial or ventricular stimulation. This was due to a shortening of the anterograde and retrograde fast pathway refractory period after administration of atropine. In patient 24 the conduction curves were discontinuous in anterograde direction, but V-A dissociation was observed during ventricular pacing using a basic cycle length of 375 ms. In patient 26 the conduction curves were discontinuous in anterograde direction but the A-H intervals obtained over the slow pathway during atrial stimulation shortened from a maximal basal value of 400 ms to a maximal value of 200 ms following atropine. In patient 35 the anterograde conduction curves were discontinuous but the exact mechanism of failure to initiate tachycardia remained unknown because ventricular stimulation could not be performed following atropine.

Discussion

Although we frequently cannot assess the correct intervals of retrograde input into the A-V node during ventricular stimulation, our results are of interest because they suggest that:

Figure 5

A representative example of findings during ventricular stimulation of group B patients in whom tachycardia could be initiated after administration of atropine. Patient 29. Before atropine a discontinuity in the retrograde conduction curve was present when during pacing with a basic cycle length of 500 ms a ventricular premature beat was given after 340 ms. This was followed by the initiation of a single "reversed" A-V nodal echo (panels A and B). Atropine resulted in shortening of the retrograde refractory period of the fast pathway, retrograde conduction over this pathway being now present up to the refractory period of the right ventricle. In contrast (panels C and D) to panels A and B where His bundle electrograms were not recorded, following atropine administration in panels C and D the retrograde His bundle potential precedes the ventriculogram during basic paced beats and follows the ventricular premature beat in panel C. The retrograde H1H2 interval shown in panel C measures 360 ms indicating that the refractory period of the retrograde pathway must have shortened to at least 360 msec after atropine administration. MRA = mid right atrium.

- a. the presence of a discontinuous retrograde conduction curve during ventricular stimulation, suggesting a long refractory period of the retrograde fast A-V nodal pathway, hampers initiation of A-V nodal tachycardia in patients having discontinuous anterograde A-V nodal conduction curves;
- b. in patients without documented tachycardia initiation of a sustained tachycardia in the A-V node after atropine is mainly the result of shortening of the refractory period of the retrograde fast pathway;

Figure 6

Same patient as in figure 5. Atrial stimulation before (panels A and B) and after (panels C and D) atropine. Before atropine pacing in the coronary sinus with a basic cycle length of 500 ms demonstrated a discontinuity in the anterograde A-V nodal conduction curve when an atrial premature beat was given after 420 ms. The longest possible A2H2 interval of 330 ms was obtained after an A1A2 interval in the coronary sinus of 260 ms (panel A). This A2H2 interval was not followed by initiation of A-V nodal echoes of the "common" type (slowfast). In panel B on further shortening of the premature stimulus interval from 210 to 200 ms the refractory period of the coronary sinus was reached. After atropine the refractory period of the anterograde A-V nodal fast pathway shortened to 260 ms. In panel C an atrial premature beat given after 270 ms is conducted to the ventricles with an A2H2 interval of 120 ms and a H1H2 interval of 340 ms. In panel D 10 ms shortening of the atrial premature beat interval to 260 ms results in demonstration of a discontinuity in the anterograde conduction curve and in initiation of tachycardia. That shortening of the retrograde refractory period of the fast pathway by atropine as shown in fig. 5 during ventricular stimulation was responsible for initiation of tachycardia is supported by observing that in panel D initiation of tachycardia occurs with an A2H2 interval of 290 ms, which is 40 ms shorter than the A2H2 interval in panel A when no echoes were initiated. Note that in panels C and D the electrograms from the HRA and CSp are reversed as compared to panels A and B.

c. patients with electrocardiographic documentation of sustained supraventricular tachycardia have a shorter retrograde refractory period of their fast pathway, as compared to those with anterograde dual A-V nodal pathways but no documentation of supraventricular tachycardia.

This study confirms previous observations by other authors (10-12) that the presence of a discontinuous A-V nodal conduction curve, either anterograde, or retrograde, is by no means a marker of the clinical occurrence of A-V nodal tachycardia. By manipulating the autonomic tone in patients with dual A-V nodal pathways but no documented tachycardia, thereby changing their A-V nodal properties, a sustained A-V nodal tachycardia can be initiated in some of these patients. The clinical significance of this finding is not clear at the present time.

Figure 7

Patient 33. A representative example of patients of group B in whom no tachycardia could be initiated after atropine. Before atropine (panels A and B) during pacing of the right ventricular apex with a basic cycle length of 500 ms a ventricular premature beat given after 420 ms was conducted retrogradely to the atria with an S2A2 interval of 270 ms and an A1A2 interval of 440 ms (panel A). A premature beat given in the ventricle 10 ms earlier (panel B) results in a marked increase in the retrograde conduction time (S2A2) to 350 ms and a corresponding A1A2 interval of 510 ms. A "reversed" (fast-slow) A-V nodal echo follows. This indicates that the refractory period of the fast pathway is approximately 410 ms (no retrograde His is observed during basic paced beats or following the ventricular premature beat). After atropine (panels C and D) pacing at the right ventricle with a shorter basic cycle length (440 ms) than in panels A and B shows retrograde conduction to the atria with a shorter S1A1 interval (250 ms in panels A and B and 120 ms in panels C and D). A ventricular premature beat given after 210 ms is still retrogradely conducted to the atria over the fast A-V nodal pathway, the H2A2 interval measuring 160 ms (panel C). On further shortening of the interval of the ventricular premature test pulse to 200 ms is reached the refractory period of the ventricle. Atropine therefore resulted in shortening of retrograde conduction time during basic pacing and in shortening of the refractory period of the retrograde pathway with disappearance of A-V nodal echoes during ventricular stimulation. 5 surface leads were simultaneously recorded with one endocavitary bipolar electrogram from the RA and three electrograms from the distal (d), mid (m) and proximal (p) His bundle area.

Figure 8
Same patient as fig. 7. Atrial stimulation studies before and after atropine. Before atropine (panels A and B) a discontinuity in the conduction curve was present when during pacing the right atrium with a basic cycle length of 500 ms an atrial premature beat was given after 310 ms. Further shortening of the atrial premature beat interval to 280 ms resulted in an A2H2 interval of 400 ms, the longest observed during atrial stimulation (panel A). It was not followed by initiation of re-entry in the A-V node most likely because the anterograde conduction time in the slow pathway was not long enough to overcome refractoriness of the retrograde fast pathway (which as shown in fig. 7 panel B measured 410 ms). On further shortening of the interval of the atrial test pulse to 270 ms (panel B) the refractory period of the slow A-V nodal pathway was reached. After atropine the anterograde conduction curves of the A-V node became continuous. Pacing of the right atrium at a basic cycle length of 440 ms resulted in anterograde conduction to the ventricles with an A1H1 interval of 80 ms (50 ms shorter than in panel A and B during pacing with a cycle length of 50 ms. An atrial premature beat given after 230 ms is conducted to the ventricles with an A2H2 interval of 190 ms and a H1H2 interval of 340 ms. On further shortening of the test stimulus interval by 10 ms the refractory period of the right atrium was reached. After atropine no re-entry in the A-V node could be initiated after atropine both during ventricular (fig. 7) and atrial stimulation. Three surface leads are shown simultaneously with four endocavitary bipolar electrograms.

The determinants of initiation of A-V nodal tachycardia in patients having electrocardiographic documentation of the arrhythmia have been studied by several authors (1-8). They all stressed the importance of the classical conditions required to initiate and sustain a re-entrant loop:

- a. presence of two electrophysiologically different pathways,
- b. unidirectional block,
- c. conduction which is slow enough to overcome refractoriness of the formerly blocked pathway and
- d. ability of both pathways to sustain sequential conduction.

Different electrophysiological methods have been used to assess the properties of the A-V nodal pathways, a major problem being the assessment of the retrograde properties of the fast pathway. In the present study we have used an approximation of the retrograde properties of the A-V node by measuring the V1-V2, H1-H2 or the S1-H2 interval during ventricular stimulation. In this way we have studied the refractoriness of the retrograde fast pathway. We realize that this method is not accurate but the only way to obtain in patients of group B an idea of the refractory period of the retrograde fast pathway. In all of them the initiation of "reversed" A-V nodal echoes at the time of the discontinuity in the retrograde conduction curve proved the A-V node to be the site of origin of the discontinuity in the conduction curve. In patients with documented supraventricular tachycardia (group A) we could show the presence of a continuous retrograde conduction curve, in most of them suggesting a short retrograde refractory period of the fast pathway. This was in contrast with the results obtained in patients of group B having no documentation of supraventricular tachycardia. In most of them we could demonstrate the presence of a discontinuous retrograde conduction curve suggesting a rather long refractory period of the retrograde fast pathway. By obtaining an approximate value of the refractory period of the retrograde fast pathway we were able to show in patients of group A that the A-H intervals over the slow pathway were sufficiently prolonged to overcome retrograde refractoriness of the fast pathway. In contrast, in patients of group B, even when the A-H intervals over the slow anterograde A-V nodal pathway were similar to those of group A patients, they were not long enough to overcome refractoriness of the retrograde fast A-V nodal pathway. These findings indicate that one can rapidly get information about the likelihood of initiation of A-V nodal tachycardia at the start of the study by looking for the presence or absence of a discontinuity in the retrograde conduction curves during single test stimulation of the ventricle. Our data demonstrate that a long refractory period of the retrograde fast pathway hampers ease of initiation of tachycardia.

The majority of the patients receiving atropine showed marked shortening in the refractory period of the retrograde fast pathway during ventricular stimulation. Initiation of tachycardia was facilitated by atropine-induced shortening of the refractory period of the retrograde fast pathway. This mechanism of facilitation of initiation of A-V nodal tachycardia has been described by other authors however in patients with electrocardiographic documentation of the arrhythmia (20-22). Our study extends these observations to patients without documentation of supraventricular tachycardia.

In 5 of the patients of group B (table II) we were able to initiate a sustained A-V nodal tachycardia after atropine administration. The mechanism by which initiation of tachycardia was facilitated was the same as in patients having documented recurrent paroxysmal A-V nodal tachycardia. In eight patients of group B, however, the conditions required to initiate and maintain re-entry in the A-V node were not induced by atropine, and tachycardia could not be initiated. Six out of 7 patients (in one patient ventricular stimulation was not performed after atropine) showed shortening of the refractory period of the retrograde fast pathway. Two mechanisms prevented initiation of reentry in the A-V node following atropine in these patients:

In five the anterograde conduction curve became continuous after atropine because of shortening in the anterograde refractory period of the fast pathway. In the remaining patient showing anterograde discontinuous curves following atropine marked shortening of the A-H interval over the slow pathway during atrial stimulation, probably prevented retrograde activation of the fast pathway and initiation of reentry. One patient showed no V-A conduction following atropine.

The exact level of V-A block after atropine could not be determined in this patient. These observations indicate that in patients without electrocardiographic documentation of A-V nodal tachycardia in whom the arrhythmia can be initiated following administration of atropine the mechanism of facilitation of initiation of the arrhythmia was shortening of the retrograde refractory period of the fast pathway. Although initiation of tachycardia occurred in some patients of group B, atropine-induced changes in the anterograde conduction curves, leading to continuous anterograde A-V nodal conduction prevented initiation of tachycardia in the majority of them.

Analysis of the results of programmed electrical stimulation in the two groups of patients studied show that initiation of A-V nodal tachycardia by this technique is easier in patients with electrocardiographic documentation of the arrhythmia (group A) as compared to patients with dual A-V nodal pathways but without documented tachycardia. These data are in agreement with previous experiences by other authors (9). In most of our patients with documented supraventricular tachycardia we could initiate the arrhythmia in the basal state by a single atrial premature beat during atrial pacing. In these patients the conditions for re-entry in the A-V node were already present in the basal state except for unidirectional block which was created by the premature impulse. In patients without documented tachycardia, however, a long retrograde refractory period of the fast pathway prevented initiation and/or perpetuation of re-entry of the common type in the A-V node. A minority of patients of group A required atropine to initiate the arrhythmia, in contrast to a majority of group B patients. The presence of a discontinuous conduction curve in the A-V node, even when retrograde, has been considered a possible marker of A-V nodal tachycardia in patients without documented arrhythmias suffering from palpitations (23). This is not the opinion of all authors (11-12). It has been shown that in arrhythmia-free children discontinuous A-V nodal conduction curves during atrial stimulation are far from rare (11-12). We have shown that in patients with a discontinuous retrograde conduction curve initiation of a sustained "reversed" (fast-slow) intra-A-V nodal tachycardia is unlikely (18). All our patients having a sustained A-V junctional tachycardia with a P-R interval shorter than the R-P interval in the presence of discontinuity in the curves of ventriculo-atrial conduction suffered from a tachycardia with a slowly conducting accessory A-V pathway with decremental conduction properties incorporated in the tachycardia circuit.

We conclude that electrophysiologic studies of the A-V node in most patients clinically suffering from A-V nodal tachycardia of the common (slow-fast) type reveal discontinuous anterograde conduction curves and continuous retrograde conduction curves. In patients with dual A-V nodal pathways but no documented tachycardia manipulation of the autonomic system will facilitate induction of A-V nodal tachycardia when this results in abbreviation of the duration of the refractory period of the retrograde fast pathway to values below the conduction time of the anterograde slow A-V nodal pathway. The clinical significance of this arrhythmia is presently unknown.

References

1. Rosen KM, Mehta A, Miller RA: Demonstration of dual atrioventricular nodal pathways in man. *Am J Cardiol* 33:291, 1974.
2. Wu D, Denes P, Wyndham C, Amat-y-Leon F, Dhingra RC, Rosen KM: Demonstration of dual A-V nodal pathways utilizing a ventricular extrastimulus in patients with atrioventricular nodal re-entrant paroxysmal supraventricular tachycardia. *Circulation* 52:789, 1975.
3. Rosen KM, Denes P, Wu D, Dhingra RC. Electrophysiological diagnosis and manifestations of dual A-V nodal pathways. In: *The conduction system of the heart*, edited by Wellens HJJ, Lie KI, Janse MJ. Leiden: Stenfert-Kroese BV; 1976; 453-66.
4. Wellens HJJ, Durrer D: The role of an accessory atrioventricular pathway in reciprocal tachycardia. *Circulation* 51:997, 1975.

5. Touboul P, Huerta F, Porte J, Delahaye JP. Reciprocal rhythm in patients with normal electrocardiogram: evidence for dual conduction pathways. *Am Heart J* 91:3, 1976.
6. Denes P, Wu D, Amat-Y-Leon F, Dhingra RC, Wyndham CR, Rosen KM: The determinants of atrioventricular nodal re-entry with premature atrial stimulation in patients with dual A-V nodal pathways. *Circulation* 56:253, 1977.
7. Pritchett ELC, Gallagher JJ, Wallace AG: Re-entry within the atrioventricular node in man: a reassessment. *Eur J Cardiol* 6/6:437, 1978.
8. Akhtar M: Paroxysmal atrioventricular nodal re-entrant tachycardia. In: *Cardiac arrhythmias*, edited by Narula OS. Baltimore: The Williams and Wilkins CO., 1979; 294-317.
9. Wellens HJJ: Value and limitations of programmed electrical stimulation of the heart in the study and treatment of tachycardias. *Circulation* 57: 845, 1977.
10. Denes P, Wu D, Dhingra R, Amat-y-Leon F, Wyndham C, Rosen KM: Dual atrio-ventricular nodal pathways. A common electrophysiological response. *Br. Heart J* 37:1069, 1975.
11. Thapar MK, Gillette PC. Dual atrioventricular nodal pathways: A common electrophysiologic response in children. *Circulation* 60:1369, 1979.
12. Casta A, Wolff GS, Mehta AV, Tamer D, Garcia OK, Pickoff A, Ferrer PL, Sung RJ, Gelband H: Dual atrioventricular nodal pathways: A benign finding in arrhythmia-free children with heart disease. *Am J Cardiol* 46:1013, 1980.
13. Ross D, Farré J, Bär FWHM, Dassen WRM, Wiener I, Wellens HJJ: Comprehensive clinical electrophysiological studies in the investigation of documented or suspected tachycardias. *Circulation* 61:1010, 1980.
14. Josephson ME, Seides SF: *Clinical cardiac electrophysiology*. Philadelphia: Lea & Febiger, 1979; 23-59.
15. Brugada P, Ross D, Bär WHM, Vanagt EJ, Dassen WRM, Wellens HJJ: Observations on spontaneous termination of atrioventricular nodal re-entrant tachycardia. *Am J Cardiol* 47:703, 1981.
16. Akhtar M, Damato AN, Ruskin JN, Batsford WP, Reddy CP, Ticzon AR, Dhatt MS, Gomes JAC, Calon AH: Anterograde and retrograde conduction characteristics in three patterns of paroxysmal atrioventricular junctional re-entrant tachycardia. *Am Heart J* 95:22, 1978.
17. Farré J, Wellens HJJ. The value of the electrocardiogram in diagnosing the mechanism and site of origin of supraventricular tachycardia. In: *What's new in electrocardiography*, edited by Wellens HJJ and Kulbertus HE. The Hague: Martinus Nijhoff Publishers, 1981; 131-71.
18. Brugada P, Bär FWHM, Vanagt EJ, Friedman PL, Wellens HJJ: Observations in patients showing A-V junctional echoes with a shorter P-R than R-P interval. *Am J Cardiol* 48:611, 1981.
19. Sung RJ, Waxman HL: Physiological shift of retrograde atrioventricular nodal output mimicking retrograde accessory pathway conduction *Circulation* 59,60: (Suppl. II) II-191, 1979.
20. Neuss H, Schlepper M, Spies HF, De Mey CH. Effects of heart rate and atropine on "dual AV conduction". *Br Heart J* 37:1216, 1975.
21. Akhtar M, Damato AN, Batsford WP, Caracta AR, Ruskin JN, Weisfogel GM, Lau SH: Induction of atrioventricular nodal re-entrant tachycardia after atropine. *Am J Cardiol* 36:286, 1975.
22. Wu D, Denes P, Bauernfeind R, Dhingra R, Wyndham C, Rosen KM: Effects of atropine on induction and maintenance of atrioventricular nodal re-entrant tachycardia. *Circulation* 59:779, 1979.
23. Strasberg B, Swiryn S, Bauernfeind R, Palileo E, Scagliotti D, Elise Duffy C, Rosen KM: Retrograde dual atrioventricular nodal pathways. *Am J Cardiol* 48:639, 1981.

6.

Dual atrio-ventricular nodal pathways and atrial fibrillation

Pedro Brugada, M.D.; Denis Roy, M.D.; Jim Weiss, M.D.; Willem R.M. Dassen, Ph.D.; Frits W.H.M. Bär, M.D.; and Hein J.J. Wellens, M.D.

From the Department of Cardiology, University of Limburg, Annadal Hospital, Maastricht, The Netherlands.

Summary

The determinants of the ventricular rate during atrial fibrillation were studied in a group of eleven patients demonstrating dual A-V nodal pathways during atrial stimulation. The shortest R-R interval and the mean ventricular cycle length during at least 1 min of pacing induced atrial fibrillation were compared a) to the effective and functional refractory period of the fast pathway, b) to the effective refractory period of the slow pathway determined during atrial stimulation, at two or more different basic cycle lengths of pacing and c) to the shortest cycle length during atrial stimulation followed by 1-to-1 A-V conduction. A group of 8 patients not demonstrating dual A-V nodal pathway-curves during atrial stimulation was used as control.

In both groups the shortest R-R interval during atrial fibrillation was best predicted by the shortest cycle length followed by 1-to-1 A-V conduction during atrial stimulation. The mean ventricular cycle length during atrial fibrillation was not accurately predicted by any of the variables studied. The similar results in patients with and without dual A-V nodal pathways suggest that **retrograde** concealed conduction in the fast pathway does not play a role in determining the ventricular response during atrial fibrillation in patients with dual A-V nodal pathways.

Introduction

The electrophysiological properties of the normal A-V node protect the ventricle against supraventricular rhythms with excessively fast rates (1-2). The decremental conduction properties of the A-V node and its rather long refractory period at rest result in ventricular rates that rarely exceed 180 beats/min in the normal individual (2). Concealed conduction in the A-V node is known to play an important role in the ventricular response during atrial fibrillation (1). The technique of programmed electrical stimulation of the heart has been of help to demonstrate longitudinal dissociation of the A-V node into different electrophysiological pathways (3). The determinants of the ventricular rate during atrial fibrillation in the presence of dual A-V nodal pathways are not known. In patients with dual A-V nodal pathways retrograde concealed conduction from one to another pathway might result in a different pattern of A-V conduction during atrial fibrillation than in patients not having dual A-V nodal pathways. We therefore compared a group of eleven patients with dual A-V nodal pathways and a control group of eight patients without dual A-V nodal pathways. In both groups of patients a similar pacing protocol was used during atrial stimulation to assess the properties and patterns of A-V conduction. The results of this study are reported here.

Patients and methods:

Patients were included in this study when the following criteria were fulfilled:

- a. Sinus rhythm at the time of study.
- b. A complete electrophysiological investigation using our previously described protocol of stimulation, recording, and analysis of tracings (4).
- c. No accessory pathways.

- d. A duration of atrial refractoriness not preventing determination of the exact values for the refractory periods of the A-V nodal pathway(s).
- e. At least one episode of pacing induced atrial fibrillation lasting at least 1 minute.
- f. No antiarrhythmic medication.

Patients having discontinuous A-V nodal conduction curves but having a refractory period of the slow A-V nodal pathway which was less than the refractory period of the atrium were not included in this study. This markedly limited the number of patients considered for inclusion.

Nineteen patients out of a series of 370 consecutive patients studied by programmed electrical stimulation of the heart were included. Following informed consent they were studied because of a variety of reasons (table 1). There were 15 males and 4 females. Aged ranged from 15 to 70 years. Eleven patients (group A) demonstrated the presence of dual A-V nodal pathway and eight patients (group B) not. Catheters were introduced in the femoral veins under fluoroscopic control and placed at the right atrium, right ventricular apex, His bundle area and in the coronary sinus in all but one patient (patient 7) of group A and in three patients of group B (patients 13-15). Simultaneous recordings were made of the surface leads I, II, III, V1 and V6 and bipolar endocavitary electrograms in the right atrium (RA), proximal and distal coronary sinus (CSp, CSd) when available, and His bundle. Atrial stimulation was performed using at least two different basic cycle lengths of pacing. Five patients in group A and 2 patients in group B had three and 2 patients in group A and 1 in group B four cycle lengths tested. The extrastimulus technique was used to determine patterns of A-V conduction and refractory periods. The prematurity of the test pulse was increased by 10 ms steps. Incremental atrial pacing was performed starting with a rate just above the spontaneous sinus rate and every 10-15 seconds increased in steps of 10 beats/min. This was done without interrupting pacing, to the pacing rate where the 1-to-1 A-V relation was lost, and usually thereafter continued until 2-to-1 A-V conduction developed. The Wenckebach point indicated in our patients failure of the anterograde fast A-V nodal pathway.

The presence of dual A-V nodal pathways was diagnosed when a discontinuous A-V nodal conduction curve was present. A discontinuity in the anterograde (A1A2/H1H2; A1A2/A2H2) A-V nodal conduction curve was defined as a reproducible increase by 40 ms or more in the A-V nodal conduction time of a premature beat when its prematurity was increased by 10 ms (5), and by the curve fitting method described by Rosen et al (3).

All patients in group A, and none in group B demonstrated the atypical form of Wenckebach sequence described by Rosen et al. (3) during incremental atrial pacing. These patients showed sudden increases in A-H interval at the Wenckebach point, a finding consistent with the presence of dual A-V nodal pathways. The effective refractory period (ERP) of the fast (f) A-V nodal pathway was defined as the longest A1-A2 interval not conducted over this pathway. The ERP of the slow (s) A-V nodal pathway was defined as the longest A1-A2 interval not conducted over this pathway. The functional refractory period (ERP) of the fast A-V nodal pathway was defined as the shortest H1H2 interval resulting from any A1A2 anterogradely conducted over the fast pathway. We have previously shown that the atrial stimulation site modifies the A-H intervals but not refractoriness of the A-V node in patients with dual A-V nodal pathways (6). However, in view of the observations by Batsford et al. (7) who found significant differences in A-V nodal properties depending on the atrial stimulation site in patients without dual A-V nodal pathways, only results of right atrial stimulation were used. After studies were performed during sinus rhythm atrial fibrillation was induced in these patients by high rate (300-500 beats/min) atrial stimulation. The shortest R-R interval during atrial fibrillation and the mean ventricular cycle length were assessed by measuring them over a continuous 1 min period of atrial fibrillation. Measurements were made at a paper speed of 100 mm/s.

The ERPF, ERPs, FRPF and shortest cycle length followed by 1-to-1 A-V conduction were compared to the shortest R-R interval and mean ventricular cycle length during atrial fibrillation, looking for the best fit with the line of identity.

Results

A summary of the electrophysiologic studies is given in table I.

Table I

Summary of clinical and electrophysiological data of patients studied

<i>Pt</i>	<i>Age/sex</i>	<i>clinical diagnosis</i>	<i>BCL</i>	<i>ERPf</i>	<i>ERPs</i>	<i>FRPf</i>	<i>Sh CL 1-to-1</i>	<i>Sh RR Afib</i>	<i>MVCL Afib</i>
Group A									
1	22 M	SVT/AoS	750 600	300 310	230 330	450 430	400	400	600
2	15 M	SVT	600 500	290 290	280 280	430 430	370	360	468
3	52 F	SVT	680 500	360 360	220 250	440 440	300	290	375
4	49 F	SVT	500 450	280 330	250 270	380 400	300	280	384
5	62 M	SVT	600 550	280 270	230 260	390 370	333	310	441
6	40 M	SVT	600 500	370 370	270 270	470 500	400	380	500
7	53 M	LBBB/ Asympt	850 600	410 400	350 340	500 500	450	490	714
8	69 M	SVT	600 500	370 370	300 320	450 450	400	380	600
9	45 M	Palp.	600 500	440 400	230 260	500 430	353	300	340
10	19 M	SVT	600 550	290 290	280 280	590 590	300	310	500
11	70 M	SVT	600 500	290 270	270 250	350 330	333	320	435

In patients of group A the ERPf ranged from 270 to 440 ms, the ERPs from 220 to 350 ms, and the FRPf from 330 to 590 ms. In patients of group B the ERP of the A-V node ranged from 220 to 450 ms, and the FRP of the A-V node from 310 to 550 ms. The shortest cycle length during atrial stimulation followed by 1-to-1 A-V conduction ranged from 300 to 450 ms in patients of group A, and from 300 to 500 in group B.

Using the unpaired student's T-test group A and B did not differ significantly in any measured parameter.

Shortest R-R interval during atrial fibrillation ranged from 280 to 490 ms in patients of group A. Those values were not accurately predicted from the data on ERPf, FRPf and ERPs (fig. 1). The shortest R-R interval during atrial fibrillation was best predicted from the shortest cycle length during atrial pacing followed by 1-to-1 A-V conduction (fig. 1).

Similar results were obtained in patients of group B, where the shortest cycle length with 1-to-1 A-

Table I (contd.)

<i>Pt</i>	<i>Age/sex</i>	<i>Clinical diagnosis</i>	<i>BCL</i>	<i>ERPf</i>	<i>FRPf</i>	<i>Sh CL 1-to-1</i>	<i>Sh RR Afib</i>	<i>MVCL Afib</i>
Group B								
12	62 M	APB's	800 760	230 310	400 390	370	380	490
13	46 F	VPB's	670 550	280 300	490 480	428	420	517
14	52 M	VPB's	750 600	430 430	550 550	500	410	625
15	64 F	Dizzy spells	600 500	280 280	390 390	300	300	353
16	20 M	Parox. Afib	600 500	310 300	310 320	410	340	545
17	69 M	Atrial tachyc	600 500	300 280	365 350	300	300	500
18	50 M	Parox. Afib	750 500	310 300	390 370	333	260	361
19	35 M	Palp.	600 500	250 220	400 360	300	330	405

Age in years. Abbreviations: SVT: supraventricular tachycardia. Aos: aortic stenosis. LBBB/asympt: asymptomatic patient with left bundle branch block. Palp: palpitations. APB's: atrial premature beats. VPB's: ventricular premature beats. BCL: basic cycle length of pacing. ERPf: Effective refractory period of the fast A-V nodal pathway. ERPs: effective refractory period of the slow A-V nodal pathway. FRPf: Functional refractory period of the fast A-V nodal pathway. Sh CL 1-to-1 AV conduction during atrial stimulation at increasing rates. Sh RR Afib: Shortest RR interval during atrial fibrillation. MVCL Afib: Mean ventricular cycle length during atrial fibrillation. All values in milliseconds.

V conduction during atrial pacing best predicted the shortest R-R observed during atrial fibrillation. But this was not predicted accurately by any other of the variables studied (ERPf, FRPf, ERPs) (Fig. 1).

Mean ventricular cycle length during atrial fibrillation ranged from 340 to 714 ms in patients of group A, and from 353 to 625 ms in patients of group B. In both groups the ERPf, FRPf, ERPs and shortest cycle length with 1-to-1 A-V conduction did not accurately predict the mean ventricular cycle length during atrial fibrillation (Fig. 2).

Distribution of R-R intervals.

Most of our patients had short-lasting (1 to 3 min.) atrial fibrillation and analysis of the distribution of R-R intervals was hampered because of the short observation period. In four patients, however, more than 500 cycles during atrial fibrillation were available. Two patients were in group A and two patients in group B. R-R intervals in these four patients followed a unimodal distribution curve during atrial fibrillation (Fig. 3). None of these 4 patients showed a bimodal distribution curve which may be observed in the presence of two A-V pathways. The ventricular response was irregular in all patients of group A and group B. The mean difference between shortest R-R interval and mean ventricular cycle length during atrial fibrillation was 137.9 ms in patients of group A and 117.3 ms in patients of group B. These results were not significantly different.

Figure 1

Relationships between the shortest R-R interval during atrial fibrillation and the ERP_f, ERP_s, FRP_f and shortest cycle length (CL) followed by 1-to-1 AV conduction during atrial pacing.

The shortest CL followed by 1-to-1 AV conduction was the best predictor of the shortest R-R interval during atrial fibrillation in both groups. Only the longest cycle length of pacing tested during right atrial stimulation is depicted in this figure and figure 2. Results were similar when the shortest cycle length of pacing used or combinations of the shortest and the longest were studied.

Discussion

The results of this study indicate that the determinants of the ventricular rate during atrial fibrillation (shortest R-R interval and mean ventricular cycle length) are not affected by the presence or absence of dual A-V nodal pathways as demonstrated by atrial stimulation.

Some limitations of this study are obvious. First, we cannot exclude that patients of group B also had longitudinal dissociation of the A-V node. When the anterograde fast pathway has a refractory period which is shorter than the refractory period of the slow anterograde pathway discontinuities in the A-V nodal conduction curve will not be present, and a diagnosis of dual A-V nodal pathways cannot be made. Patients of group A and B underwent a similar stimulation protocol and the A-V nodal conduction curves behaved differently during stimulation. The ventricular response during atrial fibrillation was however similar. Only the shortest cycle length followed by 1-to-1 A-V conduction during incremental atrial pacing was able to predict in both groups the shortest R-R interval during atrial fibrillation. None of the variables studied could predict the mean ventricular cycle length during atrial fibrillation in any group. Second, the possibility exists that the similarity in ventricular response during atrial fibrillation in both groups

Figure 2
Relationships between the mean ventricular cycle length (MVCL) during atrial fibrillation and the ERP_f, ERPs, FRP_f and shortest CL with 1-to-1 AV conduction during atrial pacing.

Figure 3
Distribution of R-R intervals during atrial fibrillation in two patients in whom 600 ventricular cycles were measured. Distribution of R-R intervals was unimodal in the patient of group A (solid line - closed circles) and in the patient of group B (broken line - open circles).

was the result of a similar behaviour of the distal common pathway (the His bundle) while the A-V nodal output might be different. His bundle recordings were available in all patients during atrial fibrillation. A-V block distal to the His bundle was never observed. Third, these were selected patients from a population demonstrating dual A-V nodal pathways, and it is difficult to generalize the conclusions reached here to the whole population of patients showing dual A-V nodal pathways.

Concealed conduction in the A-V junction is known to play a role in the ventricular response during atrial fibrillation (1). Impulses penetrating the A-V node not being anterogradely conducted are able to delay or block conduction of subsequent impulses. It has also been shown that concealed re-entry in the AV node occurs during atrial pacing (8-11). The question arises therefore whether in the presence of dual A-V nodal pathways additional concealed retrograde conduction in the fast A-V nodal pathway can play a role in the behaviour of A-V conduction during atrial fibrillation. The similar results obtained in this limited number of observations in patients of group A and group B suggests however, that in patients with dual A-V nodal pathways concealed retrograde conduction into the fast A-V nodal pathway does not play a role in the shortest R-R interval and mean ventricular cycle length during atrial fibrillation.

The studies of Meijler et al. (12) are pertinent to our observations. Investigations performed by these authors in rat and dog hearts, and during electrophysiological investigations in patients, showed small differences in A-V nodal memory (time required to accommodate A-V nodal conduction to changes in atrial rate) when compared to the differences in size of the A-V nodes of these species. Human A-V nodal memory was found to be relatively short compared to the average R-R intervals during atrial fibrillation. They concluded that the ventricular irregularity during atrial fibrillation is caused primarily by the random excitatory process of the atria. The role of the A-V node in the ventricular response is limited to scaling down the atrial impulses, a result of the interplay between refractoriness and concealed conduction. A-V nodal memory was too short to affect conduction of subsequent impulses. Our data, with similar results in patients of group A and B would suggest that A-V nodal memory plays no role in the ventricular response in patients with dual A-V nodal pathways. Unfortunately we did not specifically study A-V nodal memory in our patients.

In conclusion, the ventricular response during atrial fibrillation in patients with dual A-V nodal pathways does not seem to differ from the ventricular response during atrial fibrillation in patients not demonstrating dual A-V nodal pathways during atrial stimulation studies.

References

1. **Langendorf R, Pick A, Katz LN:** Ventricular response in atrial fibrillation. Role of concealed conduction in the A-V junction. *Circulation* 32:69, 1965
2. **Godtfredsen J:** Atrial fibrillation. Etiology, course and prognosis. A follow-up study of 1212 cases. University of Copenhagen press. 1974, p. 107-9.
3. **Rosen KM, Denes P, Wu D, Dhingra RC:** Electrophysiological diagnosis and manifestations of dual A-V nodal pathways. In: *The conduction system of the heart*. Edited by Wellens H.J.J. Lie KI, Janse MJ. HE Stenfert Kroese BV. Leiden 1976, p. 453-66.
4. **Ross DL, Farré J, Bär FWHM, Vanagt EJ, Dassen WRM, Wiener I, Wellens HJJ:** Comprehensive electrophysiological studies in the investigation of documented or suspected tachycardias: times, staff, problems and costs involved. *Circulation* 61:1010, 1980
5. **Wellens HJJ, Durrer D:** The role of an accessory atrioventricular pathway in reciprocal tachycardia. *Circulation* 52:58, 1974
6. **Ross DL, Brugada P, Bär FWHM, Vanagt EJ, Wiener I, Farré, Wellens HJJ:** Comparison of right and left atrial stimulation in demonstration of dual atrioventricular nodal pathways and induction of intranodal re-entry. *Circulation* 64:1051, 1981
7. **Batsford WP, Akhtar M, Caracta AR, Josephson ME, Seides SF, Damato AN:** Effect of atrial stimulation site on the electrophysiological properties of the atrio-ventricular node in man. *Circulation* 50: 283, 1974
8. **Damato AM, Varghese PJ, Lau SH, Gallagher JJ, Bobb GA:** Manifest and concealed re-entry. A mechanism of AV nodal Wenckebach phenomenon. *Circ. Res.* 30:283, 1972.
9. **Gallagher JJ, Damato AN, Varghese PJ, Lau SH:** Manifest and concealed re-entry. A mechanism of AV nodal Wenckebach in man. *Circulation* 47: 752, 1973.
10. **Simson MB, Spear JF, Moore EW:** Electrophysiologic studies in atrio-ventricular nodal Wenckebach cycles. *Am. J. Cardiol.* 41: 244, 1978.
11. **Littmann L, Svenson RH:** Concealed re-entry: A mechanism of atrioventricular nodal alternating Wenckebach periodicity. *Circulation* 65: 1269, 1982.
12. **Meijler FL, Heethaar RM, Harms FMA, Robles de Medina EO, Prystowsky EN, Zipes DP, Fisch C:** Comparative atrioventricular conduction and its consequences for atrial fibrillation in man. In: *Atrial Fibrillation*. Edited by Kulbertus HE, Olsson SB, Schlepper M. A Lindgren and Söner AB. Mölndal, Sweden 1982, p. 72-80.

Ventricular Tachycardia with Concealed Retrograde Conduction to Two Atrio-ventricular Nodal Pathways

Pedro Brugada, Erik W. Andries, and Hein J.J. Wellens.

From the Departments of Cardiology, University of Limburg, Annadal Hospital, Maastricht, The Netherlands, and O.L. Vrouw Hospital, Aalst, Belgium.

In patients with ventricular tachycardia, analysis of the relation between atrial and ventricular activations during arrhythmia demonstrates some form of ventriculo-atrial (V-A) conduction in approximately 50% of the cases.¹ In 5% of the patients with ventricular tachycardia, V-A conduction is a concealed phenomenon, only recognized by its effects on conduction of subsequent cardiac impulses. Concealed conduction usually results in delay of conduction or block of subsequent impulses. This can be transient or persistent.² When two A-V nodal pathways are present, concealed retrograde conduction during ventricular tachycardia can lead to very complex electrocardiographic patterns. An example is reported here.

Discussion of the Electrocardiogram

The electrocardiogram shown in Figure 1 was recorded the sixth day after an acute infero-lateral wall myocardial infarction in a 60-year-old male. That day the patient developed frequent short runs of ventricular tachycardia. He was not receiving antiarrhythmic or cardiotonic drugs at the time of electrocardiogram recording. A central venous catheter having a unipolar electrode permitted the recording of a unipolar right atrial electrogram during the arrhythmia. In the electrocardiogram two types of QRS complexes can be identified. One type has a width of

0.08 sec. These QRS complexes are in the right part of the Figure preceded by P waves of sinus origin. A second type of QRS complex having a width of 0.16 sec and a left bundle branch block morphology in lead V₁ and QS in lead II is also observed. These wide QRS complexes occur both in runs of three-to-four beats and as isolated premature beats. Measurement of the P-P interval starting in the right part of the Figure and going back to the left part of the tracing reveals a sinus tachycardia at the atrial level. There is dissociation between the wide QRS complexes and atrial activity. This suggests a ventricular origin of the wide QRS complexes. Following termination of a short run of ventricular tachycardia two different patterns of A-V conduction of the sinus impulses are observed. After the first run of ventricular tachycardia lasting three beats, the next sinus P wave is conducted with a very long P-R interval (0.44 sec). The time interval between the last tachycardia beat and the next P wave measures 0.36 sec. This pattern of conduction with a very long P-R interval persists until a new episode of ventricular tachycardia develops. After four beats the ventricular tachycardia terminates. The P wave following the last beat of this run of ventricular tachycardia is not conducted to the ventricles. The next sinus impulse is conducted to the ventricle with a P-R interval of 0.16 sec. The interval between the last tachycardia beat and the blocked P wave measures 0.24 sec. The sinus P wave following the last episode of ventricular tachycardia (right part of the Figure) is conducted anterogradely to the ventricles with a P-R interval of 0.16 sec. The interval between

Address for reprints: Hein J.J. Wellens, M.D., Department of Cardiology, University of Limburg, Annadal Hospital, Maastricht, The Netherlands.

Received August 11, 1981; accepted August 18, 1981.

Figure 1. Leads II and V₁ are simultaneously recorded with a unipolar electrogram of the right atrium at a paper speed of 25 mm/sec. The broken line in the ladder diagram represents conduction over a slow A-V nodal pathway, the solid line conduction over a fast A-V nodal pathway. A = atrium, AVN = atrioventricular node, V = ventricle. In the retrograde direction the line indicates retrograde concealed penetration of the A-V nodal pathways. See text.

the last tachycardia beat and this P wave measures 0.44 sec.

The most likely explanation for the differences in P-R interval during anterograde A-V conduction is given in the ladder diagram (Fig. 1). It is postulated that the patient has two pathways for anterograde conduction in the A-V node: a fast conducting pathway with a long refractory period, and a slow pathway with a short refractory period.³ The last beat of the first run of ventricular tachycardia results in concealed retrograde penetration of the A-V node. The next sinus P wave, occurring 0.36 sec after the last tachycardia beat, finds the fast A-V nodal pathway refractory for conduction, but can be conducted over the slow A-V nodal pathway. Sustained slow pathway conduction with a P-R interval of 0.44 sec develops. The sustained nature of this slow pathway conduction leading to a long P-R interval can be explained by two

mechanisms: a. persistence of the effect of concealed conduction during tachycardia in the fast pathway prolonging its refractory period to a value longer than the sinus cycle length, and b. concealed re-entry in the A-V node due to retrograde penetration into the fast pathway of the impulse which is conducted anterogradely over the slow pathway. The last explanation is considered to be the most likely one and is shown in the diagram. The second episode of ventricular tachycardia observed in the Figure is followed by a P wave which is not conducted to the ventricle. Following the third run of ventricular tachycardia the next P wave occurs late enough to find the fast A-V nodal pathway recovered to conduct the impulse. This and subsequent P waves are then conducted with a normal P-R interval.

As beautifully discussed by Pick and Langendorf,⁴ concealed conduction and concealed re-

entry are not uncommon mechanisms for arrhythmias in the A-V junction. To our knowledge, concealed re-entry leading to persistent P-R prolongation is a rare phenomenon, however. In our patient no His bundle electrogram was recorded. This could have facilitated the location of the area of concealed re-entry. Concealed conduction and concealed re-entry in the A-V node is a more likely explanation for the phe-

nomena observed than concealed re-entry located subnodally, considering the high prevalence of dual A-V nodal pathways in man.^{3,4}

The patient was treated with intravenous procainamide followed by oral doses of the same drug. Ventricular tachycardia did not recur during continuous electrocardiographic monitoring for ten days. A-V nodal tachycardia was never observed.

References

1. Wellens, H.J.J., Bar, F.W.H.M., and Lie, K.I.: The value of the electrocardiogram in the differential diagnosis of a tachycardia with a wide QRS complex. *Am. J. Med.*, **64**:27, 1978.
2. Langendorf, R.: Concealed A-V conduction: The effect of blocked impulses on the formation and conduction of subsequent impulses. *Am. Heart J.*, **35**:542, 1948.
3. Rosen, K.M., Denes, P., Wu, D., et al.: Electro-physiological diagnosis and manifestation of dual A-V nodal pathways. In, H.J.J. Wellens, K.I. Lie, and M.J. Janse, (Eds.): *The Conduction System of the Heart*. HE Stenfert Kroese BV, Leiden, 1976, p. 453-66.
4. Pick, A., and Langendorf, R.: *Interpretation of Complex Arrhythmias*. Philadelphia, Lea and Febiger, 1979, p. 471-533.
5. Denes, P., Wu, D., Dhingra, R., et al.: Dual atrio-ventricular nodal pathways. A common electrophysiological response. *Br. Heart J.*, **37**:1069, 1975.

Incessant Reciprocating Atrioventricular Tachycardia Factors Playing a Role in the Mechanism of the Arrhythmia

Pedro Brugada, Eddy J. Vanagt, Frits W.H.M. Bär, and Hein J.J. Wellens

From the Department of Cardiology, University of Limburg, Annadal Hospital, Maastricht, The Netherlands.

BRUGADA, P., ET AL.: Incessant reciprocating atrioventricular tachycardia. Factors playing a role in the mechanism of the arrhythmia. The case of a patient suffering from incessant supraventricular tachycardia is presented. The electrophysiological study showed the presence of an accessory atrioventricular (A-V) bundle with nodal-like properties and long conduction times. This structure was used as the retrograde arm of the tachycardia circuit. Tachycardia was intermittent at rest, but had a sustained character during slight exercise. Administration of atropine and isoproterenol failed to sustain the arrhythmia and spontaneous initiation during sinus rhythm was no longer observed. During handgrip exercise a sustained tachycardia developed immediately. During ventricular stimulation a dual atrial response to a single paced ventricular premature beat was repeatedly observed, proving the availability of two separate A-V pathways for retrograde conduction. The case illustrates the labile nature of this type of accessory pathway, and suggests that autonomic changes can play an important role in the initiation, maintenance, termination, or even spontaneous cure of tachycardia in patients with this anomaly. (*PACE*, Vol. 3, November-December, 1980)

programmed electrical stimulation, incessant supraventricular tachycardia, autonomic tone, accessory pathway, dual atrial response, single ventricular premature beat

Recently, evidence has been published for the existence of accessory atrioventricular (A-V) pathways having conduction times longer than those over the normal A-V nodal-His-Purkinje axis.¹⁻⁵ In patients with this anomaly, supraventricular tachycardias presented with a so-called "permanent" or "incessant" character.¹ Usually in these patients the accessory pathway is used

as the retrograde limb of the tachycardia circuit. Characteristically, these patients show a supraventricular tachycardia with negative P waves in II, III, AVF, and the precordial leads and a PR interval which is shorter than the RP interval.¹ It has been pointed out that the accessory pathway in these patients might be composed of nodal tissue, as suggested³⁻⁵ by the behavior of this structure during incremental pacing and following the administration of drugs, which is similar to the response shown by normal A-V nodal tissue. The anomaly is known to be congenital, but little is known about the mechanisms leading to initiation, maintenance, termination,

Submitted 4/22/80; revised 6/16/80; accepted 6/20/80.

Address for reprints: Hein J.J. Wellens, M.D., Department of Cardiology, University of Limburg, Annadal Hospital, Maastricht, The Netherlands.

or even spontaneous cure³ of the episodes of tachycardia in these cases.

We had the opportunity to study one patient with an incessant supraventricular tachycardia with some unusual features. The case illustrates the importance of autonomic changes on the genesis, maintenance, and termination of tachycardia in patients having this type of accessory pathway. A dual atrial response to a single paced ventricular premature beat, supporting the existence of two separate A-V pathways, was repeatedly observed.

Patient and Methods

The patient was a 50-year-old male who had noticed palpitations starting 11 years prior to this investigation. The electrocardiogram during tachycardia, having a rate of 120-160 beats/min, showed a negative P wave in leads I, II, III, and in the precordial leads V₁ to V₆, followed 140 ms later by a normal QRS complex. The tachycardia, which was intermittent in rest, became incessant during slight exercise, such as elevating the legs (Fig. 1). Cardiovascular examination revealed no abnormalities other than tachycardia.

Electrophysiological Study

Following informed consent an electrophysiological study was performed in the resting, post-absorptive, nonsedated state. Using the femoral approach, catheters were placed at the right ventricular apex, tricuspid valve ring, and high lateral right atrium. A fourth quadripolar electrode catheter was placed in the coronary sinus. Our methods of recording, stimulation and analysis of tracings have been previously described.⁶ After recording of basic conduction intervals during sinus rhythm, the right ventricular apex was paced at basic cycle lengths of 690 and 520 ms. Ventriculo-atrial (V-A) conduction and retrograde refractory periods were determined using the extrastimulus technique. Atrial test stimuli during sinus rhythm and atrial pacing immediately resulted in a brief period of tachycardia (2 to 3 beats) making it impossible to study the role of atrial premature beats in the initiation of tachycardia. Following these obser-

Figure 1. 6-lead electrocardiogram showing a supraventricular tachycardia with a rate of 160 beats/min. Note the negative P wave preceding the QRS complex with a PR interval shorter than the RP interval, which is characteristic of an incessant A-V tachycardia.¹

ations, 0.5 mg of atropine was administered intravenously and a single atrial test stimulation during sinus rhythm was performed from the coronary sinus. Forty-five minutes later, isoproterenol was given by continuous infusion intravenously in a dose of 1.5 μ g/min, and the ventricular stimulation study, at a basic cycle length of 520 ms, was repeated. This was followed by single atrial test stimulation from the coronary sinus during sinus rhythm. One hour after atropine administration, and 10 min after discontinuation of isoproterenol, handgrip exercise was performed. During tachycardia the effect of timed ventricular premature beats was tested.

Results

Resting State

His bundle recordings during sinus rhythm (cycle length 800 ms) revealed an A-H interval of 90 ms and an H-V interval of 40 ms. The QRS complex showed no pre-excitation, and was normal and narrow. Single beats or short runs of tachycardia were observed to occur spontaneously during sinus rhythm and during atrial pacing. Their initiation could not be related to a shortening or lengthening of the sinus cycle length. During tachycardia (cycle length 500-540 ms) the A-H interval ranged from 90 to 120 ms. Atrial activation was first recorded in the proximal coronary sinus (CSp) lead, and was closely thereafter followed by activation in the His bundle, distal coronary sinus (CSd), and high right atrial (HRA) lead. Tachycardia was never sustained; it terminated spontaneously without a definite pattern of shortening or lengthening of its cycle length, A-H or V-A intervals. Tachycardia always terminated by a QRS complex not followed by a P wave.

During ventricular stimulation with a basic cycle length of 690 ms the following observations were made: the basic ventricular beats were retrogradely conducted to the atria, and the onset of atrial activation was recorded first in the His bundle lead, followed by atrial activation in the CSp, HRA, and CSd leads. A ventricular premature beat given with a V_1V_2 interval of 440 ms was retrogradely conducted to the atria with the same atrial activation sequence as the basic paced ventricular beats, and a 20 ms increase in retrograde conduction time. This was not followed by initiation of tachycardia (Fig. 2, panel 1). A decrease of the premature beat interval by 10 ms showed retrograde conduction to the atrium with the same activation sequence as during basic ventricular pacing. However, following atrial activation, a second P wave was recorded showing an atrial activation pattern in the endocavitary leads similar to the atrial activation sequence during tachycardia. This was followed by initiation of tachycardia (Fig. 2, panel 2 and Fig. 5). A further decrease in the V_1V_2 interval revealed that the first atrial activation (showing the same sequence as during basic pac-

Figure 2. Ventricular stimulation study before drug administration. Lead II was simultaneously recorded with four bipolar endocavitary electrograms from the high right atrium (HRA), distal coronary sinus (CSd), His bundle, and proximal coronary sinus (CSp). Indicated are the basic pacing cycle length and the premature beat intervals in milliseconds between bars. V-A = ventriculo-atrial conduction time of the premature beat in milliseconds. See text.

ing) was not necessary for initiation of tachycardia (Fig. 3). A shortening of the coupling interval of the test pulse showed subnodal delay in retrograde conduction of V_2 again followed by retrograde conduction to the atria with the type of atrial activation sequence similar to basic

Figure 4. See legend for Figure 2.

paced ventricular beats; however, the second P wave and initiation of tachycardia could still be observed (Fig. 4, panel 1). At V_1V_2 intervals from 290 ms to the refractory period of the ventricle the first P wave appeared, but the second P wave was no longer observed and tachycardia no longer initiated (Fig. 4, panel 2).

During ventricular stimulation with a basic cycle length of 520 ms retrograde conduction to the atria with an atrial activation sequence similar to basic paced beats was seen up to the refractory period of the right ventricle. There was a gap in V-A conduction of the first atrial response at V_1V_2 intervals from 380 to 360 ms. This gap terminated after sufficient subnodal delay in retrograde conduction had occurred. During this gap in V-A conduction tachycardia was initiated.

During pacing with a basic cycle length of 690 ms and with a basic cycle length of 520 ms, the V-A interval as measured from the onset of the QRS complex of V_2 to the onset of atrial activation of the first tachycardia P wave was observed to increase on decreasing the V_1V_2 interval (Figs. 2-5).

Atropine Administration

Because of the unsustained nature of the tachycardia observed, atropine, 0.5 mg, was administered intravenously. The sinus cycle length shortened to 700 ms, and this was accompanied by disappearance of the spontaneously occurring echo beats or short runs of tachycardia which were previously present. Single atrial test stimulation from the coronary sinus during sinus rhythm with A_1A_2 intervals of 490 to 350 ms initiated echo beats, but no sustained tachycardia (Fig. 6).

Isoproterenol Administration

Administration of isoproterenol by continuous infusion in a dose of $1.5 \mu\text{g}/\text{min}$ resulted in shortening of the sinus cycle length to 580 ms. Spontaneous initiation of tachycardia was not observed during sinus rhythm.

Right ventricular stimulation at a basic cycle length of 520 ms showed initiation of tachycar-

Figure 5. A ventricular premature beat given after 300 ms during pacing of the right ventricular apex with a basic cycle length of 690 ms. Leads I, II, III, V₁, and V₆ were simultaneously recorded with the same four endocavitary leads as in Figures 2-4. The first retrogradely conducted atrial activation shows the same sequence of atrial activation as during atrial activation following the paced beats of the basic rhythm, and is preceded by a retrograde His bundle electrogram. After this P wave a second one, showing an atrial activation sequence similar to the one recorded during tachycardia, is seen. Same abbreviations as in Figures 2-4.

dia only when a ventricular premature beat was given after 350 ms. As compared to similar V₁V₂ intervals during ventricular stimulation with the same basic cycle length prior to drug administration, the V-A interval at the time of initiation of tachycardia shortened [V-A interval measured 640 ms after a V₁V₂ interval of 360 ms prior to drug administration, and measured 550 ms after a V₁V₂ interval of 350 ms during isoproterenol administration). The R-R interval during tachycardia shortened to 440 ms during isoproterenol administration with an A-H interval of 80 ms and a V-A interval of 320 ms (as compared to an R-R interval of 520 ms, an A-H inter-

val of 90 ms, and a V-A interval of 390 ms prior to drug administration) (Table I).

Single atrial test stimulation from the coronary sinus during sinus rhythm initiated echo beats with A₁A₂ intervals from 410 to 360 ms. A short run of tachycardia could only be initiated by an atrial premature beat given after 350 ms.

Spontaneous Termination of Tachycardia

Under all conditions tachycardia terminated spontaneously by a QRS complex which was not followed by retrograde conduction to the atrium.

Figure 6. Four atrial premature beat intervals delivered from the coronary sinus during sinus rhythm after atropine administration. Lead II was simultaneously recorded with three bipolar endocavitary electrograms. Note initiation of single echos after atrial premature beats given after 490 to 350 ms, showing shorter V-A conduction times as compared to Figures 2-5, and terminating by a QRS complex not followed by next P wave. Same abbreviations as in Figures 2-5.

Handgrip Exercise

One hour after atropine administration and 10 min after discontinuation of isoproterenol, handgrip exercise was performed. This was immediately followed by sustained tachycardia. Ventricular premature beats given during tachycardia terminated the arrhythmia. These ventricular premature beats were never followed by a retrograde P wave (Fig. 7). This finding indicates that the accessory pathway had a relatively long refractory period in ventriculo-atrial direction. The R-R interval during tachycardia

measured 440 ms, with an A-H interval of 80 ms and a V-A interval of 320 ms [Table I].

Discussion

In our patient, two A-V pathways were available for retrograde conduction, the normal His-A-V nodal axis, and an A-V accessory pathway. During basic ventricular pacing ventriculo-atrial conduction occurred over the normal A-V pathway, as suggested by the atrial activation pattern of the first P wave;⁷ the gap in retrograde

Table 1.
Time Intervals During Tachycardia Under Different Conditions

	Basal	Isoproterenol	Handgrip Exercise
R-R (ms)	520	440	440
A-H (ms)	90	80	80
V-A (ms)	390	320	320

conduction which was terminated after sufficient subnodal delay of V_2 was obtained on increasing the prematurity of V_2 .⁸ The P wave initiating tachycardia showed the same atrial activation sequence as atrial activation during tachycardia. This sequence of atrial activation was the result of retrograde conduction to the atria over the accessory pathway, showing a different entrance point into the atria as compared to the A-V node (Figs. 2-5). Of importance for the determination of the ventricular end of the accessory pathway was the observation that subnodal delay of V_2 has no effect on the retrograde conduction time over the accessory pathway, excluding a ventricular origin of this structure in the bundle of His or the A-V node. The retrograde conduction time over the accessory pathway increased on decreasing the V_1V_2 interval, demonstrating that this structure had decremental conduction properties.⁸ A dual atrial response to a single ventricular premature beat was possible because of a very long retrograde conduction time over the accessory pathway (up to 660 ms in Fig. 5), resulting in arrival over the latter structure in the atrium after the end of the atrial refractory period which resulted from the first retrogradely conducted impulse. During basic ventricular pacing with a cycle length of 690 ms the basic paced beats showed no retrograde conduction to the atria over the accessory pathway. This can be explained on the basis of concealed antegrade conduction in the accessory pathway of the atrial impulses resulting from retrograde conduction over the normal A-V axis. Of importance to support this hypothesis is the previously reported observation that ante-

grade conduction over such an accessory pathway is possible.⁹

While our observations demonstrate ventriculo-atrial conduction over a slowly conducting accessory pathway, an atrial site of origin of the tachycardia must be excluded. This can be done by noting: 1) the similar sequence of atrial activation during tachycardia and during ventricular premature beats retrogradely conducted over the accessory pathway; and 2) the reproducible termination of tachycardia after ventricular premature beats not conducted retrogradely to the atria.

The previously reported shortening in V-A conduction time over the accessory bundle following atropine administration⁸ was also seen in this patient, again suggesting A-V nodal-like tissue as the functional make-up of the accessory pathway.¹⁻³ Interesting observations were made during isoproterenol administration and during handgrip exercise. Both maneuvers are known to increase sympathetic tone, but in a different manner. Isoproterenol administration produces beta-adrenergic stimulation,¹⁰ while handgrip exercise mainly results in alpha-adrenergic

Figure 7. Tachycardia initiated during handgrip exercise. The H-H and V-A intervals during tachycardia are indicated. A ventricular premature beat given after 400 ms terminates tachycardia. This ventricular premature beat is not conducted to the atrium. Leads V_1 and V_6 and three endocavitary leads were recorded simultaneously.

stimulation.¹¹ During isoproterenol administration, spontaneous initiation of tachycardia during sinus rhythm was no longer observed and during single atrial test stimulation the echo zone was observed to narrow. As shown, the R-R, A-H, and V-A intervals were similar in tachycardia while isoproterenol was given and during handgrip exercise (Table I). Handgrip exercise, however, readily initiated and sustained tachycardia. These facts suggest that the electrophysiological changes induced in the accessory bundle were different depending upon the type of sympathetic stimulation produced.

These findings demonstrate that a delicate and labile balance in electrophysiological prop-

erties of the tachycardia circuit was required to initiate and maintain tachycardia. This balance could easily be disrupted by changing autonomic tone. Our observations suggest that autonomic changes might play an important role in the mechanisms of tachycardia in patients with this type of accessory pathway with A-V nodal-like properties.

In our patient, and also in the ones which we have reported previously,³ the weak link in the tachycardia circuit was the accessory pathway. Both during spontaneous termination and termination by induced ventricular premature beats, block always occurred in this pathway.

References

1. Coumel, P., Attuel, P., and Mugica, J.: Junctional reciprocal tachycardia. The permanent form. In H.E. Kulbertus (Ed.): *Re-entrant Arrhythmias. Mechanisms and Treatment*. MTP Press, Lancaster, England, 1977, p. 170.
2. Gallagher, J.J., and Sealy, W.C.: The permanent form of junctional reciprocating tachycardia: further elucidation of the underlying mechanism. *Eur. J. Cardiol.*, **6**:413, 1978.
3. Denes, P., Kehoe, R., and Rosen, K.M.: Multiple reentrant tachycardias due to retrograde conduction of dual atrioventricular bundles with atrioventricular nodal-like properties. *Am. J. Cardiol.*, **44**:162, 1979.
4. Epstein, M.L., Stone, F.M., and Benditt, D.G.: Incessant atrial tachycardia in childhood: association with rate-dependent conduction in an accessory atrioventricular pathway. *Am. J. Cardiol.*, **44**:498, 1979.
5. Farré, J., Ross, D., Wiener, I., Bär, F.W.H.M., Vanagt, E.J., and Wellens, H.J.J.: Reciprocal tachycardias using accessory pathways with long conduction times. *Am. J. Cardiol.*, **44**:1099, 1979.
6. Ross, D., Farré, J., Bär, F.W.H.M., Vanagt, E.J., Dassen, W.R.M., Wiener, I., and Wellens, H.J.J.: Comprehensive electrophysiological studies in the investigation of documented or suspected tachycardias: times, staff, problems and costs involved. *Circulation*, **61**:1010, 1980.
7. Gallagher, J.J., Pritchett, E.L.C., Benditt, D.G., Tonkin, A.M., Campbell, R.W.F., Dugan, F.A., Bashore, T.M., Tower, A., and Wallace, A.G.: New catheter techniques for analysis of the sequence of retrograde atrial activation in man. *Eur. J. Cardiol.*, **6**:1, 1977.
8. Akhtar, M., Damato, A.N., Caracta, A.R., Bastford, W.P., and Lau, S.H.: The gap phenomena during retrograde conduction in man. *Circulation*, **49**:811, 1974.
9. Hoffman, B.F., and Cranefield, P.F.: *Electrophysiology of the Heart*. Futura Publishing Company, Mount Kisco, New York, 1976, p. 132.
10. Lands, A.M., Arnold, A., MacAuliff, J.P., Ludvena, F.P., and Brown, T.L.: Differentiation of receptor systems activated by sympathomimetic amines. *Nature*, **214**:597, 1967.
11. Donald, K.W., Lind, A.R., McNicol, G.W., Humphreys, P.W., Taylor, S.H., and Stauton, H.P.: Cardiovascular responses to sustained (static) contractions. *Circ. Res.*, **20** & **21** (Suppl I):15, 1967.

Observations in Patients Showing A-V Junctional Echoes With a Shorter P-R Than R-P Interval

Distinction Between Intranodal Reentry or Reentry Using an Accessory Pathway With a Long Conduction Time

Pedro Brugada, MD; Frits W.H.M. Bär, MD; Eddy J. Vanagt, MD; Peter L. Friedman, MD, PhD; Hein J.J. Wellens, MD, FACC

Maastricht, The Netherlands

Single test stimulation of the ventricle revealed initiation of echoes with a supraventricular QRS complex with a shorter P-R than R-P interval in 28 of 300 patients consecutively studied with programmed electrical stimulation of the heart because of documented or suspected tachycardias. In all 28 the initiation of echoes was related to a discontinuity in the retrograde conduction curve. In 10 patients a different atrial activation sequence in the endocavitary leads was present before and after the discontinuity in the retrograde conduction curve. In five of these a sustained tachycardia with a shorter P-R than R-P interval could be initiated, and in all five patients an accessory pathway with a long conduction time as the retrograde arm of the tachycardia circuit could be demonstrated. In these five patients spontaneous initiation of tachycardia was observed during sinus rhythm or after atrial premature beats. Tachycardia accelerated after the administration of atropine. In the remaining 23 patients the initiation of echoes showing a shorter P-R than R-P interval was nonsustained. In these patients spontaneous initiation of such echoes during sinus rhythm or initiation by atrial premature beats was not observed, and echoes with this relation of the P-R and R-P intervals systematically disappeared after administration of atropine.

It is postulated that in these patients a slow atrioventricular (A-V) nodal pathway is used in the retrograde direction during echoes showing a shorter P-R than R-P interval. Sustained A-V junctional tachycardia showing this relation between P-R and R-P intervals favors incorporation of an accessory pathway with slow retrograde conduction in the tachycardia circuit.

When single test stimulation of the ventricle results in initiation of echoes with a supraventricular QRS complex in relation to a critical delay in retrograde conduction to the atria, the presence of dual pathways with ventriculoatrial (V-A) conduction over a slowly conducting intranodal pathway is considered the causal mechanism.^{1,2} These echoes present electrocardiographically with a retrograde P wave and a P-R interval that is shorter than the R-P interval. A few examples of sustained tachycardia of this type have been reported.^{2,3} Recently the existence of accessory pathways whose conduction time in the retrograde direction is longer than conduction time over the normal atrioventricular (A-V) pathway has been documented.⁴⁻⁸ Patients with this anomaly present with a supraventricular tachycardia showing a retrograde P wave and a shorter P-R than R-P interval, because a slowly conducting accessory pathway in the retrograde direction is being used. The electrocardiographic distinction between V-A conduction over a slowly conducting accessory pathway or a slowly conducting A-V nodal pathway can be extremely difficult or even impossible. Knowledge of the pathways involved in supraventricular tachycardia is of therapeutic interest, particularly in those patients whose arrhythmia is poorly controlled by drug

From the Department of Cardiology, University of Limburg, Annadal Hospital, Maastricht, The Netherlands. Manuscript received February 17, 1981; revised manuscript received April 14, 1981, accepted April 17, 1981.

Address for reprints: Hein J. J. Wellens, MD, FACC, Department of Cardiology, University of Limburg, Annadal Hospital, Maastricht, The Netherlands.

therapy and who might benefit from surgical procedures. The present study was undertaken to further elucidate the pathways involved during reentry in the human heart with a retrograde P wave and a shorter P-R than R-P interval.

Methods

Study patients: Patients were included in this investigation if they fulfilled the following criteria: (1) supraventricular echoes or tachycardia was initiated by ventricular premature beats or ventricular pacing related to a critical delay in V-A conduction. These echoes showed in the surface electrocardiogram with a normal QRS or typical bundle branch block configuration and a shorter P-R than R-P interval. The H-V interval of the His bundle electrogram preceding the echo had to be 35 ms or more. (2) A complete electrophysiologic investigation was performed because of documented or suspected tachycardia, using a previously described⁹ protocol of stimulation, recording and analysis of tracings. In all patients a coronary sinus or left atrial electrogram was recorded simultaneously with the surface leads I, II, III, V₁ and V₆, and endocavitary recordings were obtained from the right atrium and His bundle. (3) Anterograde conduction over an accessory pathway was excluded during the stimulation study.

Informed consent was obtained from every patient. No patient was receiving medication at the time of study, which was performed in the resting, postabsorptive, nonsedated state. In 12 patients in whom only echoes or nonsustained tachycardias were initiated, the effect of 1 mg of atropine given intravenously was studied. The same amount of atropine was given to four of the five patients in whom sustained tachycardia could be induced.

Electrophysiologic studies: Anterograde and retrograde refractory periods and patterns of A-V and V-A conduction were studied as previously described⁹ using the extrastimulus technique. The atrial activation sequence and the P wave configuration in the surface leads were carefully analyzed both during retrograde conduction to the atria during ventricular pacing and at the time of initiation of echoes or tachycardia. Anterograde conduction curves were analyzed by plotting the atrial premature beat intervals (A₁-A₂) against their A-V nodal conduction times (as measured by the A-H intervals) (A₂-H₂). Retrograde conduction curves were studied by plotting the ventricular premature beat intervals (V₁-V₂) against their retrograde conduction times to the atrium (V₂-A₂). A conduction curve was considered discontinuous when shortening of the premature beat interval (atrial or ventricular) by 10 ms resulted in an increase in the conduction time of the premature beat by 40 ms or more. This sudden unexpected increase in conduction time of the premature beat on slight shortening of the premature beat interval was considered suggestive of a shift in conduction from one pathway to another with a longer conduction time.^{2,10}

In all 28 patients included in this study this concept of a shift in conduction from one to another pathway was supported by the initiation of echoes after the discontinuity in the retrograde conduction curve. Special care was taken to identify possible retrograde His bundle potentials during retrograde conduction to the atria during ventricular stimulation. When the retrograde His bundle potential could not be identified during basic ventricular pacing, presumably because it was hidden in the QRS complex in the His bundle lead, the S₁-H₂ interval was taken as an approximation of the retrograde input to the His bundle and A-V node.¹¹ S₁ represents the stimulus artifact of the last basic paced ventricular

beat, and H₂ the retrograde His bundle potential resulting from the ventricular premature beat V₂.

A diagnosis of A-V nodal echoes or tachycardia using in the anterograde direction a slow and in the retrograde direction a fast A-V nodal pathway—the so-called usual type of A-V nodal tachycardia—was made using previously reported criteria.¹⁰

Results

In 28 (9.3 percent) of 300 consecutive patients undergoing electrophysiologic investigation because of documented or suspected tachycardias, echoes with a shorter P-R than R-P interval were initiated during ventricular stimulation. Clinical and electrophysiologic data on these patients are summarized in Table I. Data on 3 of these 28 patients (Patients 1, 2 and 3) have previously been reported in detail.^{7,8}

Mode of initiation of echoes or tachycardia: In 28 patients it was possible to initiate echoes or tachycardia with a shorter P-R than R-P interval during ventricular stimulation. However, only five patients (Patients 1 to 5) showed initiation of the same type of echoes spontaneously during sinus rhythm or after induced atrial premature beats.

Initiation of echoes or tachycardia during ventricular stimulation (Fig. 1 to 4): Initiation of echoes or tachycardia was related to all patients to delay in retrograde conduction to the atria during ventricular stimulation. In 27 patients retrograde conduction curves were available, and initiation of echoes in all was related to a discontinuity in the retrograde conduction curve. In Patient 2 the retrograde conduction curve could not be analyzed because of the presence of almost incessant tachycardia. However, two ranges of V-A conduction times with two different atrial activation sequences were identified, and have previously been reported.⁷

Initiation of echoes or tachycardia by atrial premature beats or spontaneously during sinus rhythm: In 5 of the 28 patients (group A) the same type of echoes initiated during ventricular stimulation, with the same pattern of atrial activation in the endocavitary leads, relation of P-R and R-P intervals and supraventricular QRS complex could be initiated by atrial premature beats or occurred spontaneously during sinus rhythm. In these patients, as previously reported,^{7,8} we could not identify a critical sinus cycle length or a critical A-V conduction time at the time of initiation of tachycardia. The remaining 23 patients not showing these phenomena were classified in group B.

Anterograde conduction curves: Complete anterograde conduction curves were available for two patients in group A. In both patients the conduction curves were continuous. In the remaining three patients in group A spontaneous initiation of tachycardia prevented complete study of anterograde conduction curves. Anterograde conduction curves were available for all 23 patients of group B (Table I). In 15 patients a discontinuity in the anterograde conduction curve was present, which was followed in 4 patients by initiation of single echoes and in 2 patients by initiation of A-V nodal tachycardia utilizing in the anterograde direction

a slow and in the retrograde direction a fast A-V nodal pathway.

Observations on the atrial activation sequence in the endocavitary leads and P wave configuration in the surface leads during retrograde conduction to the atria: During retrograde conduction to the atria two different atrial activation sequences were observed in all five patients in group A and in five patients in group B (Table I). The change in the atrial activation sequence was related in nine of these patients to a critical delay in retrograde conduction to the atria during ventricular stimulation and marked by a discontinuity in the retrograde conduction curve. As previously emphasized, this could not be observed in Patient 2. In all 10 patients retrograde activation of the atria during fast pathway conduction started in the His bundle lead and was followed by activation in the remaining endocavitary leads (Fig. 1, 3 and 4). In the remaining 18 patients in group B the same pattern of atrial activation in the

endocavitary leads was recorded before and after the discontinuity in retrograde conduction (Fig. 2).

We attempted a comparative analysis of the configuration of the P wave in the surface leads before and after the discontinuity in the retrograde conduction curve. This was facilitated by the very long retrograde conduction time at the time of initiation of echoes (Fig. 1 to 4). A negative P wave was recorded in surface leads II, III and V_6 during retrograde conduction to the atria at the time of initiation of echoes or tachycardia. The P wave was flat with a diphasic negative-positive component in lead I and V_1 .

A completely positive P wave in lead V_1 during retrograde conduction to the atria at the time of initiation of echoes was recorded only in those patients not showing a change in the atrial activation sequence before and after the discontinuity in the retrograde conduction curve (Fig. 2). In patients in group A the P wave was positive in lead V_1 before the discontinuity in the

TABLE I
Summary of Clinical and Electrophysiologic Data on 28 Patients Studied

Case	Age (yr) & Sex	Indication for Study	Anterograde A-V Nodal Conduction	Atrial Sequence*	Tachycardia With P-R < R-P Initiated by APBs or Sinus Rhythm	Relation Between Subnodal Delay of VPBs and Retrograde Slow Pathway Conduction
Group A						
1	9M	SVT	...	Different	Yes	...
2	6F	SVT	...	Different	Yes	...
3	50M	SVT	Continuous	Different	Yes	-
4	54F	SVT	...	Different	Yes	...
5	26M	SVT	Continuous	Different	Yes	-
Group B						
6	11F	SVT	Continuous	Same	No	+
7	52M	Palp	Discontinuous	Same	No	...
8	30M	Palp	Discontinuous	Same	No†	...
9	42M	Palp	Continuous	Same	No	+
10	38M	Palp	Discontinuous	Same	No	+
11	35M	Palp	Discontinuous	Same	No†	...
12	25F	VT	Continuous	Same	No	+
13	46F	VT	Discontinuous	Same	No	...
14	15M	SVT	Discontinuous	Same	No†	...
15	33M	VT	Continuous	Same	No	...
16	54M	VT	Continuous	Same	No	...
17	55F	SVT	Discontinuous	Same	No	...
18	55F	SVT	Discontinuous	Same	No†	+
19	69M	VT	Discontinuous	Same	No	+
20	45M	Palp	Discontinuous	Same	No	+
21	32M	Palp	Continuous	Same	No	+
22	35M	Palp	Discontinuous	Same	No	...
23	75F	SVT	Discontinuous	Same	No†	+
24	41M	SVT	Continuous	Different	No	+
25	41F	Palp	Discontinuous	Different	No†	+
26	43M	Palp	Discontinuous	Different	No	+
27	40M	Palp	Continuous	Different	No	+
28	24F	VT	Continuous	Different	No	+

* Before and after discontinuity in retrograde conduction curves. † Atrial premature beats initiated single A-V nodal echoes using in the anterograde direction a slow and in the retrograde direction a fast pathway. ‡ Atrial premature beats initiated sustained A-V nodal tachycardia using in the anterograde direction a slow and in the retrograde direction a fast pathway.

APBs = atrial premature beats; Palp = palpitations; SVT = supraventricular tachycardia; VPBs = ventricular premature beats; VT = ventricular tachycardia; + = present; - = absent; ... = not measurable.

FIGURE 1. Patient 3 (group A). Two ventricular premature beat intervals are shown during pacing of the right ventricular apex with a basic cycle length of 600 ms. In **A**, a ventricular premature beat given after 440 ms is followed by retrograde conduction to the atrium, the first atrial activation being recorded in the His bundle lead (thick arrow) and thereafter followed by activation in the remaining endocavitary leads (thin arrows). Note the negative P wave in leads II and III, and a positive P wave in lead V₁. In **B**, shortening of the premature beat interval by 10 ms results in prolongation of the ventriculoatrial conduction time of the premature beat (V₂-A₂ interval) by 390 ms, and the initiation of echo beats. Note the change in sequence of atrial activation as well as in the P wave configuration in lead V₁ at the time of initiation of echoes. Five surface leads were simultaneously recorded with four endocavitary electrograms from the high right atrium (HRA), distal (d) and proximal (p) coronary sinus (CS), and His bundle area.

FIGURE 2. Patient 18 (group B). Two ventricular premature beat intervals are shown during pacing of the right ventricular apex with a basic cycle length of 870 ms. In A, a ventricular premature beat given after 660 ms is followed by retrograde conduction to the atria. Atrial activation starts in the His bundle area (thick arrow), thereafter followed by activation in the mid right atrial (MRA), high right atrial (HRA) and distal coronary sinus (CSd) lead. In B, a marked increase in the V₂-A₂ interval occurs on shortening the V₁-V₂ interval by 10 ms, and echoes are initiated. Note, in contrast to Figure 1, there is no change in the sequence of atrial activation and P wave configuration in the surface leads at the time of initiation of echoes. The P wave remains positive in lead V₁.

retrograde conduction curve, changing into a flat P wave after the discontinuity (Fig. 1).

Relation between subnodal delay of ventricular premature beats and conduction over the slow pathway during ventricular stimulation: The relation between retrograde conduction to the atria over the slow pathway and subnodal delay of ventricular premature beats could be studied in 2 patients in group A and in 14 patients in group B (Table I). In patients in group A (Fig. 5) retrograde conduction to the atria over the slow pathway was not related to a critical S₁-H₂ interval, suggesting that retrograde conduction in these patients occurred over a slow pathway that was not related to retrograde input into the His bundle. This observation therefore indicates that the distal end of the slow retrograde pathway did not communicate with the His bundle.

In 14 patients in group B (Fig. 3 and 4), a gap was observed in retrograde conduction over the slow pathway, with resumption of conduction and echoes when sufficient Purkinje-His delay had occurred to overcome refractoriness of this area. This finding indicates that conduction over the slow pathway was related to retrograde input into the bundle of His. This observation was made in all patients in group B showing a change in the atrial activation sequence after the discontinuity in the retrograde conduction curve, and in nine patients in group B not showing such a change (Table I).

Duration and mode of termination of reentry showing a shorter P-R than R-P interval: All patients in group A had sustained episodes of tachycardia showing this relation of the P-R and R-P intervals. In patients in group B echoes lasted only one to three cycles except in one patient with a maximum of six con-

FIGURE 3. Patient 24 (group B). A ventricular premature beat given after 370 ms during basic pacing of the right ventricular apex with a basic cycle length of 680 ms is conducted in retrograde manner to the atria showing a change in the sequence of atrial activation as compared with the basic paced beats and to ventricular premature beats given before the discontinuity in the retrograde conduction curve (not shown). Following retrograde conduction to the atria an echo is observed. Note the diphasic negative-positive P wave in lead V₁.

FIGURE 4. Patient 24 (same patient as in Fig. 3). **B,** a ventricular premature beat given after 340 ms is not followed by retrograde conduction to the atria, and no echo is initiated. **C,** on further shortening of the V₁-V₂ interval to 320 ms, delay occurs at the subnodal level during retrograde conduction of V₂. This is shown by the retrograde His bundle potential (H₂) following V₂. This delay allows resumption of retrograde conduction over the His bundle followed by retrograde conduction over the slow A-V nodal pathway, and resumption of initiation of echoes.

FIGURE 5. Patient 5, group A. Two ventricular premature beats are shown during pacing of the right ventricular apex with a basic cycle length of 500 ms. In **A**, a ventricular premature beat given after 310 ms is followed by retrograde conduction to the atrium; the first atrial activation is recorded in the CSp lead. A retrograde His bundle potential (H₂) is recorded preceding retrograde atrial activation. The S₁-H₂ interval measures 430 ms. Note that during basic paced beats the earliest atrial activation is recorded in the His bundle lead. **B**, shortening of the ventricular premature beat interval by 10 ms results in failure of retrograde conduction to the atria in spite of prolongation of the S₁-H₂ interval, indicating that ventriculoatrial conduction in **A** occurred over a slowly conducting accessory pathway and that the refractory period of this pathway measured 300 ms (**B**).

secutive echo beats. In no patient in group B was a sustained tachycardia of this type observed.

Spontaneous termination of tachycardia was rarely observed in four patients in group A, but frequently occurred in Patient 3. Termination always occurred by a QRS complex not followed by subsequent atrial activation. Similarly, in patients in group B with repeated spontaneous termination, block always occurred in the retrograde slow pathway.

Effect of ventricular premature beats during reentry: In three patients in group A (with rates of tachycardia varying from 120 to 180 beats/min) atrial activation during tachycardia showing the same atrial activation in the endocavitary leads could be advanced by a single ventricular premature beat delivered at a time when the His bundle was refractory (Fig. 1 of reference 7). In Patient 3 tachycardia was terminated by a single ventricular premature beat that was not followed by retrograde conduction to the atria and that was delivered at the time of refractoriness of the bundle of His (Fig. 6 of reference 8). In Patient 5 in group A, with a rate of tachycardia of 215 beats/min, a ventricular premature beat delivered during tachycardia at the time

of refractoriness of the bundle of His did not advance subsequent atrial activation. However, when the first ventricular premature beat was followed by a second one, subsequent atrial activation could be advanced without modifying the activation of the His bundle. Tachycardia could also be terminated by this mode of pacing. Because of the short duration of reentry, the effect of ventricular premature beats could not be systematically studied in patients in group B.

Effect of ventricular premature beats during sinus rhythm: In Patient 3 in group A it was observed that during sinus rhythm, when a ventricular premature beat was delivered when the His bundle was expected to be refractory for V-A conduction, retrograde conduction to the atrium was possible (Fig. 6). In six patients in group B ventricular premature beats occurring at the time of refractoriness of the bundle of His were never followed by retrograde conduction to the atria.

Effect of administration of atropine (Table II): Four patients in group A and 12 patients in group B received atropine. As previously described,^{7,8} this resulted in acceleration and persistence of tachycardia in patients in group A, with shortening of the A-H and V-A

FIGURE 6. Patient 3. When ventricular pacing is initiated during sinus rhythm, the first paced ventricular complex occurs at a time when the His bundle is refractory from anterograde conduction of the sinus impulse (arrow). However, ventriculoatrial conduction occurs and is followed by an echo beat. This mode of ventriculoatrial conduction and initiation of an echo beat can be explained only by accepting the presence of a slowly conducting accessory pathway. Five surface leads were simultaneously recorded with four endocavitary electrograms.

TABLE II

Effect of Atropine Administration on Anterograde and Retrograde Conduction and on Initiation of Tachycardias

Conduction Curves			Tachycardias Initiated
Case	Anterograde	Retrograde	
Group A			
1	...	...	SVT, P-R < R-P
3	Continuous	...	SVT, P-R < R-P
4	Continuous	Discontinuous	SVT, P-R < R-P
5	Continuous	Discontinuous	SVT, P-R < R-P
Group B			
6	Continuous	...	No
7	Continuous	Continuous	No
8	Discontinuous	Continuous	No
10	Discontinuous	...	AVN (s-f)
12	...	...	No
14	Continuous	Continuous	No
17	Discontinuous	...	AVN (s-f)
20	Discontinuous	...	AVN (s-f)
22	Continuous	Continuous	No
24	Continuous	Continuous	No
26	Continuous	Continuous	No
27	Discontinuous	...	AVN (s-f)

AVN (s-f) = atrioventricular nodal tachycardia with use of in anterograde direction a slow and in retrograde direction a fast pathway; SVT, P-R < R-P = supraventricular tachycardia with a shorter P-R than R-P interval.

intervals during tachycardia in all but Patient 5, who showed only shortening of the A-H interval. In two patients of this group retrograde conduction curves remained discontinuous after administration of atropine.

In patients in group B administration of atropine gave completely different results. Echoes or tachycardia with a shorter P-R than R-P interval could not be initiated after the administration of atropine (Table II) in any patient in this group. The retrograde conduction curves became continuous in six patients, including two in whom a change in the atrial activation sequence was observed after the discontinuity in the retrograde conduction curve before administration of atropine. In four patients retrograde conduction was not studied after the administration of atropine because atrial premature beats resulted in initiation of A-V nodal tachycardia utilizing a slow pathway in the anterograde direction and a fast pathway in the retrograde direction. In the remaining two patients the retrograde conduction curves were not studied after administration of atropine.

Discussion

Characteristics of the dual A-V transmission system: In 1966 Moe and Mendez¹² postulated a dual A-V transmission system to explain echoes like those observed in our patients. Characteristically their occurrence is related to a critical delay in retrograde conduction to the atria during ventricular stimulation. Since then, the site of longitudinal dissociation has been considered to be located in the A-V node.^{1,2,12-14} Our

results demonstrate that the presence of an accessory A-V pathway with a long conduction time can give rise to an electrocardiographically similar pattern (echoes with a retrograde P wave, a supraventricular QRS complex and a shorter P-R than R-P interval). However, our patients were characterized by the sustained nature of the arrhythmia, which could also be initiated by atrial premature beats or occur spontaneously during sinus rhythm. Furthermore, the mechanism persisted with acceleration of tachycardia after administration of atropine. In contrast, in patients showing the "unusual" form of A-V nodal reentry, the arrhythmia is not sustained, is not initiated by atrial premature beats, does not occur spontaneously during sinus rhythm and cannot be initiated after the administration of atropine.

Characteristics of the retrograde accessory A-V pathway: In all patients in group A we found definite evidence of the presence of an accessory A-V pathway, which was concealed in the surface electrocardiogram but was used in retrograde manner during tachycardia. The accessory pathway had a retrograde conduction time that was longer than the retrograde conduction time over the normal A-V pathway. The normal A-V pathway had a retrograde refractory period that was longer than the refractory period of the accessory pathway. These properties resulted in the observation of discontinuous retrograde conduction curves with two different atrial activation sequences depending on the pathway used for retrograde conduction. The diagnosis of an accessory A-V pathway used in retrograde manner during tachycardia in patients in group A was based on the following criteria: (1) advancement of subsequent atrial activation during tachycardia with the same atrial activation sequence by a ventricular premature beat delivered during tachycardia at the time of refractoriness of the bundle of His^{7,10}; and (2) termination of tachycardia by a ventricular premature beat that was not conducted in retrograde manner to the atria, occurring at the time of refractoriness of the bundle of His.⁸ Occasionally, as in Patient 5, these features were demonstrated only by giving two premature ventricular beats consecutively.

New observations demonstrating V-A conduction over an accessory pathway with a long conduction time were: (1) the occurrence of V-A conduction after a ventricular premature beat given during sinus rhythm, at a time when the His bundle was refractory for retrograde conduction (Fig. 6); and (2) the finding that V-A conduction over a slow pathway is not related to the timing of retrograde activation of the bundle of His, indicating that the slow pathway is functionally dissociated from the His-A-V nodal pathway (Fig. 5).

Features of atrial activation in presence of accessory pathway: All patients with evidence of the presence of an accessory pathway with a long conduction time manifested two different atrial activation sequences on careful study with multiple endocavitary recordings. The location of the atrial end of these accessory pathways with a long conduction time, as well

as the criteria to exclude a nodoventricular accessory pathway, have been discussed previously.⁷ We and others⁴⁻⁶ have shown that this type of accessory pathway has its atrial end close to the mouth of the coronary sinus. As Sung and Waxman¹⁵ observed in a preliminary report, in some patients in whom the presence of an accessory pathway can be excluded, a change in the sequence of atrial activation can be found depending on the use of the fast or the slow A-V nodal pathway. These findings suggest that in some patients there are different atrial entry points for the fast and the slow A-V nodal pathway in the atrium. This observation was made in five of our patients in group B. Janse et al.¹⁶ previously described the existence of a dual input to the A-V node in the rabbit heart during sinus rhythm and atrial stimulation: a posterior input by way of the crista terminalis entering the node beneath the coronary sinus ostium, and an anterior input entering the node as a broad wave front anterior to the ostium of the coronary sinus. They also reported the presence of two exits to the atrium during retrograde conduction over the A-V node. A similar situation may exist in our five patients in group B who showed a change in atrial activation pattern after the discontinuity in the retrograde conduction curve.

In retrospect, we consider it unfortunate that we did not repeat in our patients the experiments performed by Schuilenburg and Durrer.¹⁷ They showed that in patients having the type of echo described in our report a part of the reentrant cycle could be preexcited by stimulating the atrium before the expected arrival at the atrium of the retrograde impulse in the course of the echo sequence. A similar study in our patients with comparison of the effect of coronary sinus stimulation versus low right atrial stimulation on the time relations during the reentrant cycle would probably have given us more information on the atrial entry sites during reentry and the role of the atrium in the reentrant circuit. Additional work is required to know whether the absence of a change in retrograde atrial activation pattern during reentry after the discontinuity in the retrograde conduction curve indicates an intra-A-V nodal reentrant circuit and whether a change in the retrograde atrial activation sequence suggests incorporation of the atrium in the reentrant circuit.

The five patients in group B showing the change in the sequence of atrial activation after the discontinuity in the retrograde conduction curve could not be distinguished from patients in group A by their pattern of atrial activation in the endocavitary leads or the configuration of the P waves in the surface leads. Patients in group B who manifested the same pattern of atrial activation before and after the discontinuity in the retrograde conduction curve showed (1) first onset of atrial activation in the His bundle electrogram, and (2) a positive P wave in the surface lead V₁ before and after the discontinuity. That the changes in the atrial activation pattern are real and not a recording artifact is supported by our observation that in patients showing this phenomenon the P wave changed from positive in

lead V₁ before the discontinuity into a flat diphasic P wave in the same lead after the discontinuity (Fig. 1).

Location of the slowly conducting pathway in the A-V node: In 14 patients in group B, including all five patients showing the change in retrograde atrial activation pattern after the discontinuity in the retrograde conduction curve (Patients 24 to 28), we could demonstrate that the slowly conducting pathway was located above the bundle of His. In all 14 the occurrence of retrograde conduction over the slow pathway was related to critical timing of the retrograde input into the His bundle-A-V node. This could be beautifully demonstrated by observing the phenomenon of a gap in retrograde conduction over the slow pathway, which was overcome when sufficient subnodal delay permitted resumption of retrograde conduction over the slow pathway (Fig. 3 and 4).

In seven of the remaining nine patients in group B our assumption that retrograde longitudinal dissociation most likely occurred in the A-V node was based on the lack of positive criteria for the presence of an accessory pathway, and the finding of anterograde longitudinal dissociation in the A-V node. The latter was shown by (1) presence of a discontinuous anterograde conduction curve, (2) initiation of the "usual" type of A-V nodal echoes (anterograde slow, retrograde fast) in Patients 8, 11 and 14, and (3) initiation of the "usual" type of A-V nodal tachycardia in Patient 17 after administration of atropine. In two patients in group B (Patients 15 and 16) no evidence could be obtained to support the presence of an accessory pathway with a long conduction time. These two patients also showed no signs of longitudinal dissociation of the A-V node in the anterograde direction. We realize that in these nine patients the presence of an accessory pathway in the retrograde direction cannot be totally excluded.

Eighteen patients in group B showed only one sequence of atrial activation in the endocavitary leads, and this did not change after the discontinuity in the retrograde conduction curve. In 9 of these 18 patients the presence of a retrograde gap in conduction of the type discussed proved that the distal end of the slow pathway was located above the bundle of His. As previously discussed, the similar atrial activation sequence before and after the discontinuity suggests the presence of an upper common pathway in the A-V node in these patients.

Diagnostic considerations: Of diagnostic importance is the observation of initiation of tachycardia or echoes with a shorter P-R than R-P interval spontaneously during sinus rhythm or after atrial premature beats in patients in whom an accessory pathway with a long conduction time is used in the retrograde direction. This was never observed in patients in whom a retrograde slow A-V nodal pathway was used. Although it is theoretically possible to observe the initiation of the "unusual" type of echoes during sinus rhythm or atrial stimulation in patients having dual A-V nodal pathways in the anterograde direction, this was never observed in our patients. The presence of a slow pathway with a

FIGURE 7. Diagram showing the mechanism of initiation during atrial pacing of the "unusual" type of intranodal atrioventricular echo (top panel), and of supraventricular tachycardia using in retrograde manner an accessory pathway with a long conduction time (bottom panel). Although theoretically possible, the mode of initiation depicted in the top panel was never observed during this study. See text for possible explanations. A = atrium; ap = accessory pathway with a long conduction time; AVN = atrioventricular node; BB = bundle branches; F = fast A-V nodal pathway; H = His bundle; S = slow A-V nodal pathway; V = ventricle.

longer refractory period than the fast pathway is essential for the initiation of the "unusual" type of intranodal reentry during atrial pacing (Fig. 7). In 15 of our 23 patients in group B we could demonstrate during atrial pacing that the reverse was true, thus explaining why the "unusual" type of intranodal reentry could not be initiated. In the remaining seven patients one can speculate that factors like depth of penetration into the slow pathway or early arrival of the impulse over the fast pathway at the distal common pathway prevented completion of an intranodal reentrant cycle with an-

terograde conduction over the fast and retrograde conduction over the slow pathway. It is understandable that reentry is much more easily initiated during atrial stimulation or even in sinus rhythm in the presence of a slowly conducting accessory pathway, simply by the combination of anterograde block in the accessory pathway and an echo circuit with much larger dimensions than those of the A-V node.

Effect of atropine: The effect of administration of atropine was very different in patients in group A and group B. Tachycardia persisted and accelerated in those patients in whom an accessory pathway with a long conduction time was used. However, in no patient in group B could echoes with a shorter P-R than R-P interval be initiated after the administration of atropine. In all patients in group B in whom this could be tested, atropine resulted in shortening of the retrograde refractory period of the fast pathway, and thus in disappearance of the discontinuity in conduction in the retrograde direction. Wu et al.² reported a case of "unusual" type of A-V nodal tachycardia initiated after administration of atropine. In that patient, and also in later observations by the same group,³ the use of an accessory pathway with a long conduction time as the retrograde arm of the tachycardia circuit was not excluded.

Implications: Our data suggest that in patients with A-V junctional tachycardia whose P-R interval is shorter than the R-P interval during the arrhythmia, an intranodal location of the reentrant circuit is extremely rare. For all practical purposes, including medical and surgical treatment, a tachycardia circuit incorporating a slowly conducting accessory pathway should be considered most likely.

References

- Mendez C, Moe GK. Demonstration of dual A-V nodal conduction system in the isolated rabbit heart. *Circ Res* 1966;19:378-93.
- Wu D, Dones P, Amat-y-Leon F, Wyndham CRC, Dhinra R, Rosen KM. An unusual variety of atrioventricular nodal re-entry due to retrograde dual atrioventricular nodal pathways. *Circulation* 1977;56:50-9.
- Straeßberg B, Bauernfeld R, Swiryn S, Wyndham C, Rosen K. Retrograde dual A-V nodal pathways (abstr). *Circulation* 1980; 62:Suppl III:43-47.
- Gallagher JJ, Sealy WC. The permanent form of junctional reciprocating tachycardia: further elucidation of the underlying mechanism. *Eur J Cardiol* 1978;8:413-30.
- Dones P, Kehoe P, Rosen KM. Multiple reentrant tachycardias due to retrograde conduction of dual atrioventricular bundles with atrioventricular nodal-like properties. *Am J Cardiol* 1979;44: 162-70.
- Epstein ML, Stone FM, Benditt DG. Incessant atrial tachycardia in childhood: association with rate-dependent conduction in an accessory atrioventricular pathway. *Am J Cardiol* 1979;44: 498-504.
- Farré J, Ross D, Wiener I, Barr FW, Vanagt EJ, Wellens HJJ. Reciprocal tachycardias using accessory pathways with long conduction times. *Am J Cardiol* 1979;44:1099-109.
- Brugada P, Vanagt EJ, Barr FWHM, Wellens HJJ. Incessant reciprocating atrioventricular tachycardia. Factors playing a role in the mechanisms of the arrhythmia. *Pace* 1980;3:670-7.
- Ross D, Farré J, Barr FWHM, et al. Comprehensive electrophysiological studies in the investigation of documented or suspected tachycardias: times, staff, problems and costs involved. *Circulation* 1980;61:1010-8.
- Wellens HJJ, Durrer D. The role of an accessory atrioventricular pathway in reciprocal tachycardia. *Circulation* 1975;52:58-72.
- Josephson ME, Seides SF. Clinical cardiac electrophysiology. Philadelphia: Lea & Febiger, 1979:23-59.
- Moe GK, Mendez C. The physiologic basis of reciprocal rhythm. *Prog Cardiovasc Dis* 1966;8:461-82.
- Mignone RJ, Wallace AG. Evidence for dissociation of conduction and re-entry within the AV node. *Circ Res* 1966;19:638-49.
- Schulenburg RM, Durrer D. Ventricular echo beats in the human heart elicited by induced ventricular premature beats. *Circulation* 1969;40:337-47.
- Sung RJ, Waxman HL. Physiological shift of retrograde atrioventricular nodal output mimicking retrograde accessory pathway conduction (abstr). *Circulation* 1979;59,60:Suppl II:191.
- Janse MJ, Van Capelle FJL, Anderson RH, Touboul P, Billewicz J. Electrophysiology and structure of the atrioventricular node of the isolated rabbit heart. In: Wellens HJJ, Lie KI, Janse MJ, eds. *The Conduction System of the Heart*. Leiden: Stenfert Kroese, 1978: 296-315.
- Schulenburg RM, Durrer D. Further observations of the ventricular echo phenomena elicited in the human heart. Is the atrium part of the echo pathway? *Circulation* 1972;45:629-38.

10.

Observations in patients with supraventricular tachycardia having a P-R interval shorter than the R-P interval.

The differentiation between atrial tachycardia and reciprocating atrio-ventricular tachycardia using an accessory pathway with long conduction times.

Pedro Brugada, M.D.; Jerónimo Farré, M.D.; Bill Heddle, M.B., B.S.; Denis Roy, M.D.; and Hein J.J. Wellens, M.D., F.A.C.C.

From the Department of Cardiology, University of Limburg, Annadal Hospital, Maastricht, The Netherlands.

Summary

In twenty-five consecutive patients with supraventricular tachycardia (16 incessant and 9 recurrent paroxysmal) having a P-R interval shorter than the R-P interval during the arrhythmia, the mechanism and site of origin of tachycardia was studied by programmed electrical stimulation of the heart. No patient had anterograde conduction over an accessory pathway. Fourteen patients with incessant tachycardia underwent an exercise test during the arrhythmia, and in sixteen patients the effect of 1 mg of atropine given intravenously on the time intervals during tachycardia was studied. Vagal manoeuvres were performed in all patients. In 7 patients (group A) all with incessant tachycardia, the presence of an accessory pathway with long conduction time used retrogradely during tachycardia was demonstrated. All patients had V-A conduction during ventricular pacing. Tachycardia initiated spontaneously during sinus rhythm, but was also initiated by atrial and ventricular premature beats. Exercise accelerated the rate of tachycardia by shortening both the P-R and the R-P intervals. The same effect was obtained by atropine administration. A continuous 1-to-1 A-V relation was required in these patients to perpetuate the arrhythmia. In 10 patients (group B) V-A conduction during ventricular pacing could not be demonstrated. Seven patients had incessant tachycardia. The arrhythmia initiated spontaneously during sinus rhythm and was also initiated by atrial premature beats. In two patients the mechanism of initiation remained unknown because incessant tachycardia could never be terminated. In no patient tachycardia could be initiated or terminated by ventricular premature beats or ventricular overdrive pacing.

In the remaining 8 patients (group C) V-A conduction was present. Tachycardia initiated spontaneously during sinus rhythm and was also initiated by atrial premature beats. In one patient it could also be initiated by a ventricular premature beat, but in contrast to patients of group A, without relation to a critical delay in retrograde conduction.

In group B and C atropine accelerated the rate of tachycardia in 9 of 11. In seven patients with incessant tachycardia (4 of group B and 3 of group C) the rate of tachycardia increased during exercise. Continuation of exercise resulted in the appearance of sinus tachycardia in three of them. In all patients of group B and C a 1-to-1 A-V relation was not required to perpetuate the arrhythmia. Vagal manoeuvres were of help to demonstrate perpetuation of tachycardia when conduction to the ventricles did not occur. An atrial site of origin of tachycardia could be demonstrated in these patients.

All patients of group A had during tachycardia the characteristic configuration of the P wave

described by Coumel. No patient of group B and C had this type of P wave. In them five different P wave configurations were observed.

Our study in patients with incessant or recurrent paroxysmal supraventricular tachycardia having a P-R interval shorter than the R-P interval during the arrhythmia shows that the differentiation between an atrial site of origin of tachycardia and a circus movement tachycardia using an accessory pathway with long conduction time in retrograde direction can be made by careful analysis of the 12-lead electrocardiogram and use of simple manoeuvres like carotid sinus massage, atropine administration and exercise.

Introduction

Programmed electrical stimulation of the heart has improved our knowledge of the site of origin, mechanisms and pathways of supraventricular tachycardia in man (1-2). This has resulted in a more rational therapeutical approach to the different types of supraventricular tachycardia, and the development of refined surgical techniques for some of them (3-4). It also increased our abilities of making the correct diagnosis on the 12-lead electrocardiogram in patients with supraventricular tachycardia (5). We have recently reported on patients with sustained A-V junctional tachycardia showing a P-R interval shorter than the R-P interval (6). It was demonstrated that an intranodal location of the re-entry circuit (the so-called "uncommon type" of A-V nodal tachycardia) during this arrhythmia is unlikely. These patients have a circusmovement tachycardia using in atrio-ventricular direction the normal A-V nodal-His-Purkinje pathway and in ventriculo-atrial direction an accessory pathway with long conduction times and A-V nodal-like properties. Usually these patients present clinically with a tachycardia having an almost incessant or "permanent" character (7).

Atrial tachycardia can present with a similar electrocardiographic and clinical picture. Also in view of the therapeutical implications we would like to discuss how to differentiate between both types of supraventricular tachycardia. Special emphasis will be placed upon the use of the 12-lead ECG and simple non-invasive manoeuvres (like carotid sinus massage, the administration of atropine, and the exercise test) in making this distinction.

Patients and methods

Twenty-five consecutive patients (table I) fulfilled the following criteria:

- a. Permanent or recurrent paroxysmal supraventricular tachycardia continuously or intermittently showing a 1-to-1 A-V relation with a P-R interval shorter than the R-P interval.
- b. A complete electrophysiological investigation to delineate the mechanisms, pathways and site of origin of the arrhythmia using our previously described protocol of stimulation, recording, and analysis of tracings (8). The study was performed in the resting, postabsorptive nonsedated state, or under light sedation with oral diazepam except in patient 25 who was studied under general anaesthesia. Medication was discontinued at least 48 hours before the study. No patient had received antiarrhythmic drugs with cumulative effects (like Amiodarone) before the investigation. Catheters were placed for both stimulation and recording of intracardiac bipolar electrograms in the right atrium, coronary sinus and/or left atrium, His bundle area, and the apex of the right ventricle. With the single test stimulus technique at different basic cycle lengths of pacing (at least two) patterns of A-V and V-A conduction and refractory periods were measured during atrial and ventricular stimulation. Complete studies could rarely be performed during sinus rhythm in patients having incessant tachycardia. During tachycardia the effect of timed atrial and ventricular premature beats was tested. V-A conduction over a slowly conducting accessory A-V pathway was diagnosed according to previously reported criteria (6, 9-12).
- c. Patients with anterograde conduction over an accessory pathway and patients with atrial flutter were excluded.

Fourteen patients having incessant tachycardia underwent a limited exercise test to assess its effects on tachycardia time-intervals. Sixteen patients received 1 mg of atropine intravenously to

Summary of clinical and electrophysiological data on the 25 patients studied

Patient	Age(yrs)/sex	Indication for study	Tachycardia rate (beats/min)	A-V relation	Initiation	Termination	V-A conduction
Group A Tachycardia initiated by a ventricular premature in relation to delay in retrograde conduction							
1	9 M	Incessant SVT	130	1:1	SR, APB, VPB	APB, VPB, SPT(	+
2	6 F	Incessant SVT	130	1:1	SR, APB, VPB	APB, VPB, SPT(	+
3	50 M	Incessant SVT	175	1:1	SR, APB, VPB	APB, VPB, SPT	+
4	54 F	Incessant SVT	145	1:1	SR, APB, VPB	APB, VPB(	+
5	49 F	Incessant SVT	160	1:1	SR, APB, VPB	APB, VPB(	+
6	56 F	Incessant SVT	120	1:1	SR, APB, VPB	APB, VPB	+
7	32 M	Incessant SVT	140	1:1	SR, APB, VPB	APB, VPB(	+
Group B Absent VA conduction							
8	42 F	Paroxysmal SVT	170	1:1, W, 2:1	APB	SPT	-
9	57 F	Paroxysmal SVT	140-180	1:1, W	SR	APB	-
10	35 F	Paroxysmal SVT	210	1:1, W, 2:1	SR	SPT	-
11	17 M	Incessant SVT	160	1:1, W, 2:1	SR	- (	-
12	65 F	Incessant SVT	150	1:1, W	SR, APB	SPT(	-
13	41 F	Incessant SVT	130	1:1	SR, APB	SPT(	-
14	10 F	Incessant SVT	160	1:1, W	SR, APB	SPT(	-
15	42 M	Incessant SVT	200	1:1, W, 2:1	SR	SPT(	-
16	10 F	Incessant SVT	170	1:1, W, 2:1	?	-	-
17	10 M	Incessant SVT	140	1:1, W	?	-	-
Group C V-A conduction present, but tachycardia not initiated by a ventricular premature in relation to delay in retrograde conduction							
18	73 M	Incessant SVT	140	1:1	SR, APB	APB(	+
19	19 M	Incessant SVT	120	1:1	?	-	+
20	4 M	Incessant SVT	160	1:1	SR, APB	APB	+
21	57 F	Paroxysmal SVT	130	1:1, W, 2:1	SR, APB	APB	+
22	60 F	Paroxysmal SVT	160	1:1, W, 2:1	APB	APB, SPT	+
23	69 F	Paroxysmal SVT	110	1:1, W	APB, VPB	SPT	+
24	63 M	Paroxysmal SVT	150	1:1	APB	APB, SPT	+
25	52 F	Paroxysmal SVT	180	1:1, W, 2:1	APB	APB, SPT	+

Abbreviations: A-V relation: atrio-ventricular relation during supraventricular tachycardia. APB: atrial premature beat. Initiation: Mode of initiation of tachycardia. Termination: Mode of termination of tachycardia. SR: during sinus rhythm. VPB: Ventricular premature beat. SPT: Spontaneous termination. SVT: Supraventricular tachycardia. W: Different degrees of Wenckebach pattern of conduction. +: Present. -: Absent. (?): Short-lived termination of tachycardia (?): terminated by carotid

Figure 1

Patient 1. Spontaneous termination (left) and re-initiation (right) of a tachycardia using retrogradely an accessory pathway with long conduction times. Tachycardia terminates by a QRS complex not followed by a retrograde P wave. After five sinus cycles showing a progressive increase in rate tachycardia starts spontaneously. Note the deep negative P wave in leads II, III and in the precordial leads (7). 5 Surface leads were recorded simultaneously.

determine its effects on tachycardia (table II). Apart from 12-lead electrocardiograms recorded before the stimulation study showing simultaneous recordings of three extremity or precordial leads or six extremity or precordial leads, leads I, II, III, V1 and V6 were simultaneously recorded during tachycardia in the catheterization laboratory in all patients. These tracings were used to analyse the P wave morphology during the arrhythmia.

Results

Seventeen of the 25 patients presented clinically with a permanent or almost continuous supraventricular tachycardia (table I). Eight patients had a recurrent paroxysmal form of supraventricular tachycardia. There were 15 females and 10 males. Measured at rest during stable basal conditions, rates during tachycardia individually ranged from 110 to 210 beats/min.

1. Electrophysiological studies:

a. Initiation of tachycardia:

In twenty-two patients tachycardia initiated **spontaneously during sinus rhythm** or could be initiated by **atrial premature beats**. In the remaining three patients with continuous tachycardia in whom the arrhythmia could not be terminated by pacing (patients 16, 17 and 19) the mode of initiation of tachycardia could not be documented. In patients with spontaneous initiation of tachycardia during sinus rhythm no definite relation was found between initiation of tachycardia and sinus cycle length although two patients (patient 1 and 12) had a trend towards facilitation of tachycardia by increasing the sinus rate (fig. 1). In patients in whom tachycardia could be initiated by an electrically induced atrial premature beat a spontaneously occurring atrial premature beat was sometimes recorded as the mechanism of initiation of tachycardia (fig. 2).

Figure 2

Patient 13. Spontaneous initiation of atrial tachycardia during sinus rhythm. 5 Surface leads were recorded simultaneously with four endocavitary bipolar electrograms from the right atrium (RA 1 to RA 4), the His bundle area and the coronary sinus (CS). The first P wave of the tachycardia shows the same pattern of atrial activation as subsequent tachycardia P waves (broken vertical lines). Compare the P wave morphology in the surface leads during tachycardia in this patient with the one shown in figure 1 and subsequent figures.

In eight patients a tachycardia with the same P wave morphology, pattern of atrial activation in the endocavitary leads, and P-R/R-P relation could be initiated by a ventricular premature beat. In seven of the eight patients initiation of tachycardia by a ventricular premature beat **was related** to a critical delay in retrograde conduction to the atria. This was not the case in the remaining patient (patient 23).

Based upon their behaviour during ventricular stimulation our patients were divided into three groups: Group A consisted of the seven patients in whom tachycardia could be initiated by a ventricular premature beat on reaching a critical delay in ventriculo-atrial conduction. Group B was composed of patients in whom V-A conduction during ventricular pacing was absent. Group C contained seven patients having ventriculoatrial conduction but no induction of tachycardia during ventricular pacing plus the one patient in whom tachycardia could be initiated by a ventricular premature beat without relation to a critical delay in ventriculo-atrial conduction of the ventricular premature beat.

In patients of group A initiation of tachycardia by an atrial premature beat was never possible unless the premature beat was conducted to the ventricle. In patients of group B and C atrial premature beats initiating tachycardia did not require conduction over the A-V node (documented by recording a His bundle electrogram following the atrial premature beat). When tachycardia initiated spontaneously during sinus rhythm (table I) a critical delay in A-V conduction could not

be identified in any patient of any group. These observations suggest that in patients of group A structures below the atria (A-V node, His bundle, bundle branches, ventricle) were required to initiate the arrhythmia, whereas this was not the case in patients of group B and C.

b. Atrio-ventricular relation during tachycardia

Patients of group A always had a 1-to-1 A-V relation during tachycardia. The arrhythmia terminated whenever a QRS complex was not followed by a P wave, or whenever a P wave was not anterogradely conducted to the ventricles. This phenomenon could be observed spontaneously (fig. 3) or could be provoked by carotid sinus massage in all of them and also by other vagal manoeuvres (fig. 4). Characteristically in this group of patients a Wenckebach type of conduction preceded termination of the arrhythmia in A-V or V-A direction (fig. 3 and 4).

Figure 3
 Patient 2. Spontaneous initiation of circusmovement tachycardia using retrogradely an accessory A-V pathway (top panel) and spontaneous termination (continuous tracing bottom panel). Termination of tachycardia occurs after progressive prolongation in the V-A conduction time during tachycardia from 310 to 370 ms. There are no changes in the A-H interval. Four endocavitary electrograms were simultaneously recorded with four endocavitary bipolar electrograms.

CAROTID SINUS MASSAGE

Figure 4

Patient 6. Termination of circummovement tachycardia incorporating an accessory pathway with long conduction times retrogradely by carotid sinus massage (CSM). CSM induced cycle length alternation due to alternation in the RP' interval. Note that the last RR interval is the shortest and is not followed by a retrograde P' wave. Surface leads V1 + V2 were recorded simultaneously with an oesophageal lead (OE).

In patients of group B and C tachycardia continued in spite of A-V nodal block. When a change in relation between atrial and ventricular events did not occur spontaneously, it could be provoked by carotid sinus massage or other vagal manoeuvres (fig. 5-6), or by inducing ventricular premature beats that changed the atrio-ventricular relation but not the atrial rate during

Figure 5

Patient 11. 2-to-1 A-V block and short-lasting termination of atrial tachycardia during carotid sinus massage in a patient with incessant atrial tachycardia. Tachycardia continues in the left part of the figure in spite of the absence of A-V conduction of every second P wave. Following termination of tachycardia and a single sinus beat (right part) tachycardia reinitiates spontaneously. 4 Surface leads were recorded simultaneously.

Figure 6
Patient 20. A-V block during atrial tachycardia without termination of the arrhythmia produced by a single deep inspiration. Six surface leads were recorded simultaneously.

tachycardia. All patients of group B and C showed episodes with a 1-to-1 A-V relation during tachycardia with a P-R shorter than R-P, but characteristically also changes in the relation between atrial and ventricular complexes (table I). This A-V relation could change from 1-to-1 with a P-R interval shorter than R-P interval to 1-to-1 with P-R interval longer than R-P, and thereafter to a Wenckebach type of conduction or higher degrees of A-V block without changes in the atrial rate. This could occur spontaneously, after drug administration (fig. 7), or during exercise or excitement.

It is of interest that patients of group A on changing tachycardia rate by atropine, exercise or vagal manoeuvres, always maintained a P-R interval shorter than the R-P interval during tachycardia (fig. 3, 4 and 8).

c. Termination of tachycardia

In five patients of group A (table I) tachycardia could not be long-lastingly terminated by programmed stimulation of the heart. In these patients (patients 1, 2, 4, 5 and 7) tachycardia terminated for a few sinus cycles but reinitiated immediately thereafter. In the remaining two patients tachycardia could be terminated for longer periods. **Spontaneous termination** of tachycardia was observed in three patients. This could occur either by a P wave which was not conducted to the ventricles, or more commonly by a QRS complex. In the latter situation a Wenckebach sequence of conduction in V-A direction was observed (fig. 3). Termination of tachycardia was possible in all patients of group A by **atrial premature beats**. Termination of tachycardia occurred whenever the atrial premature beat produced block in atrio-ventricular conduction. In all seven patients termination of tachycardia was possible by a **ventricular premature beat which was not conducted to the atria** (fig. 9).

Figure 7

Patient 11. Effect of 1 mg of atropine given intravenously on time intervals during atrial tachycardia. In panel A (before atropine) the tachycardia cycle length measures 400 ms and the P-R interval is only slightly shorter than the R-P interval. The A-H interval is 180 ms. Following atropine (panel B) there is marked shortening of the A-H interval to 70 ms with no increase in rate of the atrial tachycardia (cycle length 430 ms). In the extremity leads marked shortening of P-R interval and prolongation of the R-P interval can be observed. This was never seen after atropine in patients of group A.

In six group B patients tachycardia terminated **spontaneously** by a QRS complex which was not followed by a P wave. In one patient (patient 9) tachycardia could reproducibly be terminated by an induced atrial premature beat. In three patients tachycardia could not be terminated by any mode of stimulation. In this group of patients having no V-A conduction tachycardia could never be terminated by ventricular premature beats or ventricular overdrive pacing.

Group C. In four patients of group C **spontaneous termination** of tachycardia was observed. In five of them induced **atrial premature beats** were also able to terminate tachycardia. In one patient tachycardia could not be terminated by pacing.

Even when ventriculo-atrial conduction could be demonstrated in this group of patients, **ventricular premature beats** never terminated tachycardia. Following ventriculo-atrial conduction of the ventricular premature beat atrial activation did not occur prematurely enough to terminate the arrhythmia (fig. 10).

2. Effect of atropine and exercise during tachycardia

a. Atropine: (table II)

Five patients of group A, 6 of group B, and 5 of group C received 1 mg of atropine intravenously during tachycardia. The tachycardia rate accelerated in all patients of group A, in 4 patients of group B, and in 5 of group C. It did not change in one patient of group B and slowed slightly in another patient of group B (fig. 7). The A-H interval shortened in all patients following atropine. When tachycardia rate accelerated patients of all three groups showed shortening both of the P-R

Figure 8

Patient 4. Effect of exercise on the time intervals of a tachycardia using retrogradely an accessory pathway with long conduction times. Before exercise the tachycardia cycle length is 400 ms. The P-R interval is shorter than the R-P interval (140 and 260 ms respectively). Immediately after exercise the tachycardia has accelerated markedly to a cycle length of 280 ms. Both the P-R and the R-P interval have shortened as compared to the tracing in the left panel.

Figure 9

Patient 4. Termination of circummovement tachycardia using retrogradely an accessory pathway with long conduction times by a ventricular premature beat (S1). As shown this ventricular premature beat is not followed by retrograde conduction to the atria. Following termination of tachycardia the arrhythmia reinitiates after two sinus beats (right part). 5 Surface leads were recorded simultaneously with 4 endocavitary electrograms. STIM = stimulation.

Figure 10

Patient 19. Demonstration of V-A conduction during incessant atrial tachycardia having an interval between two successive atrial complexes ranging from 510 to 530 msec. During atrial tachycardia pacing is performed in the right ventricular apex. Note that following a ventricular premature beat given after 310 ms occurs earlier than expected atrial activation (arrow). The pattern of atrial activation of this complex differs from those of the atrial tachycardia (particularly at the RA and CSp electrograms) indicating that retrograde conduction following the ventricular premature beat results in a different atrial activation pattern as during atrial tachycardia. Such an observation indicates an atrial origin of the tachycardia. 5 Surface leads were recorded simultaneously with 4 endocavitary bipolar electrograms.

and R-P intervals. Therefore the response to atropine administration did not distinguish between them. In the two patients of group B with no or minimal change in tachycardia rate a marked change was observed in the P-R/R-P relation following atropine (fig. 7). This response was not observed in any patient of group A.

b. Exercise: (table II)

Exercise accelerated the rate of tachycardia in all 7 patients of group A and in 6 patients of group B and C having incessant tachycardia (table II). In all patients the P-R and R-P intervals shortened during tachycardia (figs. 8 and 11). Therefore patients of group A, B and C could not be distinguished on the basis of their behaviour during exercise. It is of interest however to observe that in three patients (patients 11, 16 and 19) maximal exercise resulted in sinus P waves replacing the tachycardia P wave. When exercise was terminated progressive slowing in sinus rate was followed by a transition from a sinus P wave back to the atrial tachycardia P wave. During this period fusion P waves were observed in these patients (fig. 12). This phenomenon was not observed during exercise in any patient of group A.

3. Carotid sinus massage and other vagal manoeuvres

Carotid sinus massage, deep inspiration or the valsalva manoeuvre produced short-lived termination of tachycardia in all patients of group A. Perpetuation of the arrhythmia when A-V or V-A block occurred was never observed (fig. 4) in this group of patients.

Table II

Effect of atropine and exercise on tachycardia time-intervals

Patient	Atropine				Exercise			
	P-P (ms)		P-R	R-P	P-P (ms)		P-R	R-P
	before	- after	after		before	- during	during	
Group A								
1	460	380	S	S	430	315	S	S
2	N.G.		N.G.	N.G.	460	350	S	S
3	520	400	S	S	500	440	S	S
4	430	330	S	S	400	280	S	S
5	N.G.		N.G.	N.G.	375	310	S	S
6	500	380	S	S	500	390	S	S
7	450	360	S	S	450	350	S	S
Group B								
8	N.G.		N.G.	N.G.	N.P.		N.P.	N.P.
9	N.G.		N.G.	N.G.	N.P.		N.P.	N.P.
10	310	240	S	S	N.P.		N.P.	N.P.
11	400	430	S	I	450	400')	S	S
12	360	350	S	I	N.P.		N.P.	N.P.
13	480	400	S	S	500	320	S	S
14	N.G.		N.G.	N.G.	400	380	S	S
15	N.G.		N.G.	N.G.	N.P.		N.P.	N.P.
16	350	300	S	S	350	300')	S	S
17	440	400	S	S	N.P.		N.P.	N.P.
Group C								
18	460	430	S	S	460	400	S	S
19	500	400	S	S	500	380')	S	S
20	N.G.		N.G.	N.G.	380	240	N.P.	N.P.
21	N.G.		N.G.	N.G.	N.P.		N.P.	N.P.
22	470	430	S	I	N.P.		N.P.	N.P.
23	N.G.		N.G.	N.G.	N.P.		N.P.	N.P.
24	480	340	S	S	N.P.		N.P.	N.P.
25	380	335	S	S	N.P.		N.P.	N.P.

Abbreviations: P-P: interval between subsequent P waves during tachycardia. P-R: P-R interval during tachycardia. R-P: R-P interval during tachycardia. N.G.: Not given. N.P.: Not performed. S: shortening. I: increase or prolongation. '): Sinus tachycardia at this cycle length appeared during exercise.

Vagal manoeuvres produced A-V block during tachycardia in all patients of group B and C (fig. 5-6) with perpetuation of tachycardia. In patient 11 (fig. 5) tachycardia was also terminated by carotid sinus massage after a short period of perpetuation in spite of A-V block. Vagal manoeuvres were of help to identify those patients in whom the arrhythmia perpetuated when a 1-to-1 A-V relation was not present, excluding participation of an accessory A-V pathway in the mechanism of tachycardia.

Figure 11
Patient 11. Effect of exercise in atrial tachycardia. Before exercise the tachycardia cycle length is 450 ms (right panel) and shortens to 400 ms during exercise (left panel) with shortening of both the P-R and R-P intervals. 6 Surface leads were simultaneously recorded.

Figure 12
Same patient as in figure 11. Immediately after termination of exercise there is a change from sinus tachycardia back to atrial tachycardia. During maximal exercise the rate of the sinus node exceeded the rate of the atrial tachycardia. During the change from sinus rhythm to atrial tachycardia fusion P waves are observed. 6 Surface leads were simultaneously recorded.

4. P wave morphology in the surface leads and pattern of endocavitary activation during tachycardia

Patients of group A showed a characteristic P wave configuration in the 12-lead electrocardiogram. As observed in figures 1, 3 and 9, the P wave was deeply negative in leads II, III and in leads V3 to V6. The P wave was diphasic or flat in leads I and V1.

In patients of group B and C five morphologically different types of P waves were recorded (fig. 6 and fig. 13). Ten patients had a cranio-caudal P wave during tachycardia which was not always clearly different on the 12-lead electrocardiogram from the P wave during sinus rhythm (panel B of fig. 13). A different pattern of atrial activation was observed, however, in the endocavitary leads when compared to the sinus P wave in all 10 patients. One patient (patient 17) had a negative P wave in lead I (panel C of fig. 13). In this patient endocardial atrial mapping demonstrated a site of origin of tachycardia in the left atrium. Three patients showed during tachycardia a caudo-cranial P wave (panel D of fig. 13) which was negative in the left precordial leads. A comparison between these three patients and patients of group A revealed that patients of group A had a much deeper and wider P wave in the inferior leads (compare fig. 13 panel D with the tachycardia shown in figs. 1 and 3). The remaining patient (patient 20) showed a right-to-left P wave which was diphasic and deeply negative in lead V1. Atrial mapping demonstrated a site of origin of tachycardia in the low anterior right atrium in this patient.

Figure 13

Four panels showing 5 surface leads in four different patients of group B and C to illustrate five different types of P waves observed during atrial tachycardia. Panel A and B show two different sorts of cranio-caudal P waves. Patients having P waves like the ones shown in panel B required endocavitary electrograms to differentiate them from sinus P waves. Panel C shows our only patient with P wave indicating activation from left to right. Panel D illustrates the caudo-cranial P wave observed in three patients of group B. A right-to-left P wave observed in patient 20 is illustrated in fig. 6.

Discussion

Site of origin and mechanisms of supraventricular tachycardia in patients of group A, B and C

Twenty-five consecutive patients with paroxysmal or incessant supraventricular tachycardia showing intermittently or continuously a 1-to-1 A-V relation with a P-R interval shorter than the R-P interval were included in the present study.

In the seven patients of group A a continuous 1-to-1 A-V relation was required to perpetuate tachycardia. In all seven patients we were able to terminate reproducibly the tachycardia for shorter or longer periods whenever this 1-to-1 A-V relation was interrupted. This could be accomplished by giving an atrial premature beat which was not conducted to the ventricles, or by a

ventricular premature beat which was not followed by retrograde activation of the atria. This last mode of termination of tachycardia excludes an atrial site of origin of the arrhythmia and indicates the incorporation of A-V junctional structures in the tachycardia circuit. In all 7 patients of group A, using previously reported criteria (6, 9-12), a slowly conducting accessory A-V pathway was demonstrated to be the retrograde pathway of a circusmovement using anterogradely the normal A-V nodal-His-Purkinje pathway. The use of this slowly conducting accessory A-V pathway in retrograde direction resulted in the electrocardiographic pattern observed (a P-R interval shorter than the R-P interval). As expected all group A patients had ventriculo-atrial conduction during ventricular pacing.

Ten patients of group B had no ventriculo-atrial conduction during stimulation of the ventricles. This observation excludes the possibility of a tachycardia using retrogradely an accessory A-V pathway. Six of the ten patients had a configuration of their P waves during tachycardia indicating cranio-caudal atrial activation suggesting that neither an accessory A-V pathway with an atrial end in the A-V groove, nor an intranodal re-entry circuit was responsible for the arrhythmia (13-15). One patient (patient 17) had a left-to right P wave, and the remaining three patients (patients 11, 15 and 16) had a caudo-cranial P wave. Definitive proof in these patients, as well as in the six patients with craniocaudal P wave, that an accessory A-V pathway was not playing a role in the mechanism of their arrhythmia was obtained by observing changes in the pattern of A-V conduction from 1-to-1 to different degrees of A-V block without termination of the arrhythmia. This observation demonstrates that activation of the ventricles is not necessary to sustain the tachycardia. When A-V block did not occur spontaneously, it could be provoked by carotid massage or other vagal manoeuvres. With the aid of endocavitary recordings it could be shown that A-V nodal conduction (as assessed from recording a His bundle electrogram following the atrial complex) was not necessary to perpetuate tachycardia. This observation is of importance in the three patients having a caudo-cranial P wave during tachycardia, because it suggests that the arrhythmia is not originating in the A-V node. Theoretically a "reversed" type of A-V nodal tachycardia with a distal common A-V nodal pathway located above the bundle of His cannot be completely excluded (16).

Patients of group C also showed variations in A-V relation without termination of tachycardia either spontaneously or following vagal manoeuvres. In contrast to patients of group B however, V-A conduction was possible in all eight patients of group C. An A-V nodal origin of the tachycardia could be excluded however by observing:

- a. A cranio-caudal or eccentric P wave during the arrhythmia and
- b. A pattern of endocardial atrial activation during tachycardia which was different from the pattern observed in the endocavitary leads following retrograde conduction during ventricular pacing.

In some patients fusion between atrial tachycardia P wave and retrograde conduction could be recorded. In one patient of this group (patient 23) tachycardia could be initiated by a ventricular premature beat not showing a critical delay in retrograde conduction to the atria. In this patient tachycardia could also easily be initiated by a single atrial premature beat. Apparently retrograde conduction following the ventricular premature beat resulted in atrial activation early enough to initiate the arrhythmia.

Effect of atropine and exercise on time intervals during tachycardia

In patients of group A, B and C atropine administration could increase the rate of the tachycardia. Both the P-R, R-P and P-P intervals shortened during tachycardia in these patients. It was not possible to distinguish between those patients having an atrial site of origin of their arrhythmia and those using retrogradely an accessory A-V pathway with long conduction time (group A). It is of interest however, that while tachycardia rate increased in all patients of group A, two patients of group B showed no change or even slight slowing in rate of tachycardia (fig. 7). In these patients the R-P interval prolonged because of shortening of the P-R interval after atropine.

In all patients with incessant tachycardia exercise accelerated the rate of tachycardia whatever the mechanism. This observation therefore was of no help to differentiate the mechanism of tachycardia because all patients showed 1-to-1 A-V relation during exercise. In three patients however exercise accelerated sinus rate to such an extent that sinus tachycardia replaced atrial tachycardia (fig. 12). Atrial tachycardia regained control of the heart immediately after a decrease in sinus rate in the postexercise period. There was a trend for patients with incessant reciprocating atrio-ventricular tachycardia to show a more marked increase in tachycardia rate during exercise as compared to patients with incessant atrial tachycardia (table II). Accurate analysis of this aspect was however not possible because of the sinus tachycardia appearing during exercise in three patients with incessant atrial tachycardia. In patients having reciprocating atrio-ventricular tachycardia using retrogradely an accessory pathway with long conduction times acceleration of the sinus rate and overdriving of the tachycardia can theoretically occur in the presence of sinus node entrance block. This phenomenon was not observed however in any patient of group A.

Differential diagnosis of a supraventricular tachycardia with a P-R interval shorter than the R-P interval

In patients presenting with a sustained supraventricular tachycardia having a 1-to-1 A-V relation and a P-R interval shorter than the R-P interval four different mechanisms for the arrhythmia have to be considered:

- a. atrial tachycardia,
- b. a "reversed type" of A-V nodal tachycardia,
- c. reciprocating A-V tachycardia using an accessory pathway with long conduction times,
- d. reciprocating A-V tachycardia retrogradely using an accessory pathway with fast conduction velocity in the presence of enhanced anterograde A-V nodal conduction, or, as recently pointed out by Gallagher et al. (17) in the presence of delay in retrograde conduction in the atrialized portion of the right ventricle in patients with Ebstein's anomaly of the tricuspid valve, and
- e. A-V nodal tachycardia with 2-to-1 A-V block (16)

Recently we have shown that a sustained supraventricular tachycardia having a P-R interval shorter than the R-P interval is unlikely to be the result of a "reversed" (fast/slow) re-entry loop in the A-V node (6). In this study of twenty-five consecutive patients no one was included having a retrogradely conducting accessory pathway of the fast type in the presence of enhanced anterograde A-V nodal conduction nor was any patient with "common" type of A-V nodal tachycardia having 2-to-1 A-V block included. The former patients usually present with a fast tachycardia because the R-P interval is in the order of 90 to 160 msec and the P-R interval of similar magnitude. In childhood enhanced anterograde atrio-ventricular conduction is more common and probably accounts for the rather poor discriminating value of the P-R/R-P relation during supraventricular tachycardia in localizing the site of origin of the arrhythmia in that age group (18). The majority of adult patients using an accessory pathway of the fast type have a tachycardia with a P-R interval longer than the R-P interval (1-2, 5). Therefore, in the adult patient when confronted with a supraventricular tachycardia showing a P-R interval shorter than the R-P interval two major possibilities have to be considered:

- a. atrial tachycardia, and
- b. reciprocating atrio-ventricular tachycardia using an accessory pathway with long conduction time in retrograde direction.

The correct diagnosis can be made by using the following approach (fig. 14):

- a. Perpetuation of tachycardia during A-V block of the P waves. This definitely excludes the possibility of a tachycardia incorporating an accessory A-V pathway. As shown in figs. 5-6 if A-V block does not occur spontaneously vagal manoeuvres is of great help to demonstrate this phenomenon.

Diagnostic approach to supraventricular tachycardia with PR < RP

Figure 14
Diagnostic approach to a supraventricular tachycardia presenting with a P'-R shorter than the R-P' interval and a 1-to-1 A-V relation. Abbreviations: AccP LCT: accessory pathway with long conduction time.

- b.** A P wave configuration indicating cranio-caudal activation of the atria during tachycardia. This makes retrograde activation of the atria from the A-V node or over an accessory pathway with its atrial end located in the A-V groove extremely unlikely. Such a configuration of the P wave argues in favour of atrial tachycardia.
- c.** Termination of tachycardia by a ventricular premature beat (spontaneous or induced) which is not retrogradely conducted to the atria. This observation indicates that structures other than the atria are required to perpetuate the arrhythmia. When in addition the ventricular premature beat occurs at the time when the His bundle is refractory for retrograde conduction and the tachycardia is terminated without V-A conduction of the ventricular premature beat, participation of an accessory pathway in retrograde direction is certain (6).
- d.** Initiation of the tachycardia by a ventricular premature beat on reaching a critical delay in retrograde conduction to the atria. As previously reported (6), in patients using retrogradely an accessory pathway with long conduction time tachycardia is initiated when retrograde conduction to the atria is blocked over the A-V node and exclusively occurs over the accessory pathway.
- e.** Disappearance of the abnormal P waves during tachycardia and the appearance of sinus P waves during exercise. This argues for atrial tachycardia.
- f.** Marked shortening in the P-R interval during tachycardia without change in the P-P interval during atropine administration indicates an atrial origin of the arrhythmia. In patients with an accessory pathway with long conduction time the R-P interval shortens following atropine because of the A-V nodal-like properties of the accessory pathway.
- g.** As described by Coumel (7) the P wave observed in patients of group A has a characteristic configuration due to the location of the atrial end of the accessory pathway close to the mouth of the coronary sinus. Theoretically an atrial tachycardia arising in this area can show the same P wave morphology.

Conclusions

Our study demonstrates that differentiation between atrial tachycardia and a tachycardia using an accessory pathway with long conduction time in retrograde direction can be made on the 12-lead ECG alone or by using simple manoeuvres like carotid sinus massage. Observations on the mode of initiation and termination of the arrhythmia (particularly by ventricular premature beats) facilitates making the correct diagnosis.

References

- 1. Wellens, H.J.J.; Durrer, D.:** The role of an accessory atrioventricular pathway in reciprocal tachycardia: observations in patients with and without the Wolff-Parkinson-White syndrome. *Circulation* 52:58, 1975.
- 2. Akhtar, M. Damato, A.N.; Ruskin, I.N.; et al.:** Antegrade and retrograde conduction characteristics in three patterns of paroxysmal atrio-ventricular junctional re-entrant tachycardia. *Am. Heart J.*, 95:22, 1978.
- 3. Gallagher, J.J.; Gilbert M.; Svenson, R.H.; Sealy, W.C.; Kasell, J.; Vallau, A.G.:** Wolff-Parkinson-White syndrome. The problem, evaluation and surgical correction. *Circulation* 51:767, 1975.
- 4. Harrison, L.; Gallagher, J.J.; Kasell, J., et al.:** Cryosurgical ablation of the atrio-ventricular node-His bundle: a new method for producing atrio-ventricular block. *Circulation* 55:463, 1977.
- 5. Farré, J. and Wellens, H.J.J.:** The value of the electrocardiogram in diagnosing site of origin and mechanism of supraventricular tachycardia. In: Wellens H.J.J. and Kulbertus H.E., eds.: "What's new in electrocardiography". Martinus Nijhoff Publishers, The Hague 1981; p. 131.
- 6. Brugada, P.; Bär, F.W.H.M.; Vanagt, E.J.; Friedman, P.L.; Wellens, H.J.J.:** Observations in patients showing A-V junctional echoes with a shorter P-R than R-P interval. *Am. J. Cardiol.* 48:611, 1981.
- 7. Coumel, P.; Attuel, P.; Leclercq, J.F.:** Permanent form of junctional reciprocating tachycardia: mechanism, clinical and therapeutical implications. In: Narula O.S. (Ed).: *Cardiac arrhythmias, electrophysiology, diagnosis and management.* Williams and Wilkins, Baltimore 1979, p. 347.

8. Ross, D.; Farré, J.; Bär, F.W.H.M. et al.: Comprehensive electrophysiological studies in the investigation of documented or suspected tachycardias: times, staff, problems and costs involved. *Circulation* 61:1010, 1980.
9. Gallagher, J.J.; Sealy, W.C.: The permanent form of reciprocating tachycardia: further elucidation of the underlying mechanism. *Eur. J. Cardiol.* 8:413, 1978.
10. Denes, P.; Kehoe, P.; Rosen, K.M.: Multiple re-entrant tachycardias due to retrograde conduction of dual atrio-ventricular bundles with atrio-ventricular nodal-like properties. *Am. J. Cardiol.* 44:162, 1979.
11. Epstein, M.L.; Stone, F.M.; Benditt, D.G.: Incessant atrial tachycardia in childhood: association with rate-dependent conduction in an accessory atrio-ventricular pathway. *Am. J. Cardiol.* 44:498, 1979.
12. Farré, J.; Ross, D.; Wiener, I.; Bär, F.W.H.M.; Vanagt, E.J.; Wellens, H.J.J.: Reciprocal tachycardias using accessory pathways with long conduction times. *Am. J. Cardiol.* 44:1099, 1979.
13. Amat-y-Leon, F.; Dhingra, R.C.; Wu, D.; Denes, P.; Wyndham, C.; Rosen, K.M.: Catheter mapping of retrograde atrial activation. Observations during ventricular pacing and A-V nodal re-entrant paroxysmal tachycardia. *British Heart J.* 38:355, 1976.
14. Maclean, W.A.H.; Karp, R.; Kouchoukos, N.T.; James, T.N.; Waldo, A.L.: P-waves during ectopic atrial rhythms in man. A study utilizing atrial pacing with fixed electrodes. *Circulation* 52:426, 1975.
15. Garcia-Civera, R.; Ferrero, J.A.; Sanjuan, R. et al.: Retrograde P-wave polarity in reciprocating tachycardias utilizing lateral bypass tracts. *Eur. Heart J.* 1:137, 1980.
16. Wellens H.J.J.; Wesdorp, H.; Düren, D.R.; Lie, K.I.: Second degree block during reciprocal atrio-ventricular nodal tachycardia. *Circulation*, 53:595, 1976.
17. Gallagher, J.J.; Smith, W.M.; Kerr, C.R.; German, L.D.; Sterba, R.; Sealy, W.C.: Etiology of long R-P' tachycardia in 33 cases of supraventricular tachycardia. *Circulation* 64 (IV), 145, 1981.
18. Garson Jr., J. and Gillette, P.C.: Electrophysiologic studies of supraventricular tachycardia in children. II. Prediction of specific mechanisms by noninvasive features. *Am. Heart J.* 102:383, 1981.

Summary

In this study new observations are described on the role of the A-V junction in tachycardias in man using the method of programmed electrical stimulation of the heart. In chapter III the complexity of the human A-V node during stimulation is illustrated by some unusual modes of initiation of A-V nodal tachycardia. As shown in chapter IV different re-entry circuits can interact in the human heart and result in termination of arrhythmias. The observation of a discontinuous A-V nodal conduction curve in anterograde direction has not the same significance in patients with and without documented tachycardia. Chapter V showed that in the last group of patients a long retrograde refractory period of the fast pathway prevents initiation of re-entry in the A-V node. In some of these patients, however, a sustained tachycardia can be initiated after atropine administration. Atropine results in shortening of the refractory period of the retrograde fast pathway. Chapter VI describes that during atrial fibrillation the pattern of atrio-ventricular conduction in the presence of dual A-V nodal pathways does not differ from the pattern observed during A-V conduction over an A-V node without discontinuous curves. Complex electrocardiographic patterns can occur during ventricular tachycardia in patients having dual A-V nodal pathways as shown in chapter VII. Chapter VIII reports that in the incessant form of reciprocating atrio-ventricular tachycardia autonomic tone plays an important role. As pointed out in chapter IX and X the electrocardiogram, effect of drug administration, exercise test, vagal manoeuvres and observations during electrophysiologic studies help us to differentiate between the different mechanisms of incessant supraventricular tachycardia.

Samenvatting

In deze studie worden nieuwe waarnemingen beschreven over de rol van de AV knoop bij tachycardiën. Deze waarnemingen zijn gedaan tijdens geprogrammeerde elektrische stimulatie van het hart bij de mens.

In hoofdstuk III wordt de complexiteit van A-V nodale geleiding gedemonstreerd aan de hand van enkele voorbeelden, die de verschillende manieren laten zien waarop A-V nodale tachycardiën opgewekt kunnen worden. In hoofdstuk IV wordt aangegeven hoe verschillende re-entry circuits in het hart elkaar kunnen beïnvloeden en dat deze wisselwerking kan leiden tot het beëindigen van de tachycardie. Wanneer men een of twee wegen in de anterograde A-V nodale geleidingscurve is de betekenis verschillend bij patiënten met en patiënten zonder gedokumenteerde tachycardiën.

Hoofdstuk V laat zien, dat in deze laatste groep patiënten een lange retrograde refractaire periode van het snel geleidende AV knoop pad het ontstaan van re-entry tachycardiën voorkomt. Bij sommigen van deze patiënten kan een stabiele tachycardie wel worden gestart na atropine toediening. Atropine veroorzaakt dan een verkorting van de refractaire periode van het in retrograde richting snel geleidende pad in de A-V knoop.

In hoofdstuk VI wordt beschreven dat de aan- of afwezigheid van twee intra A-V nodale paden het A-V geleidingspatroon bij boezemfibrilleren niet beïnvloedt. Dat de aanwezigheid van twee A-V nodale paden kan leiden tot complexe electrocardiografische beelden bij een kamertachycardie wordt gedemonstreerd in hoofdstuk VIII.

In hoofdstuk VIII wordt geïllustreerd dat het autonome zenuwstelsel een belangrijke rol speelt bij A-V nodale re-entry tachycardiën.

In de hoofdstukken IX en X wordt beschreven hoe men moet differentiëren tussen de verschillende soorten van supraventriculaire tachycardiën door op een juiste wijze gebruik te maken van het electrocardiogram, resultaten van geneesmiddeltoediening, inspanningselectrocardiografie, vagale manoeuvres en geprogrammeerde prikkeling van het hart.

Resumen

Usando la técnica de estimulación eléctrica programada del corazón se han descrito en este estudio nuevas observaciones sobre el papel que la unión A-V tiene en las taquicardias humanas. En el capítulo III se ha ilustrado la complejidad del nodo A-V analizando modos poco comunes de inicio de la taquicardia intranodal. Tal como se ha ilustrado en el capítulo IV cuando mas de un posible circuito de reentrada se halla presente en el corazón, pueden interferir entre sí y resultar en terminación de una taquicardia en uno de los circuitos. Una curva de doble paso intranodal observada durante la estimulación auricular no tiene el mismo significado en pacientes con taquicardia supraventricular documentada que en pacientes sin documentación electrocardiográfica de taquicardia supraventricular. En el capítulo V se describió que en el último grupo de pacientes el período refractario de la vía rápida intranodal es largo en dirección retrógrada e impide iniciar la taquicardia nodal. En alguno de estos pacientes sin embargo, la taquicardia puede iniciarse después de atropina, debido al acortamiento del período refractario retrógrado de la vía rápida. El significado de esta arritmia en estos pacientes es hasta ahora desconocido. En el capítulo VI se ha mostrado que el patrón de conducción atrio-ventricular durante la fibrilación auricular no difiere entre pacientes con o sin curvas de doble paso intranodal. El capítulo VII ha ilustrado como patrones electrocardiográficos complejos pueden aparecer durante una taquicardia ventricular cuando existe un doble paso intranodal. El capítulo VIII demuestra que el tono autonómico tiene un papel importante en el síndrome de la taquicardia incesante supraventricular debida a una vía accesoria oculta con conducción lenta. En los capítulos IX y X se ha descrito como usando el electrocardiograma, efectos de administración de drogas, prueba de esfuerzo, maniobras vagales y las observaciones hechas durante el estudio electrofisiológico, podemos llegar al diagnóstico del mecanismo de las taquicardias incesantes supraventriculares.

Resum

En aquest estudi es presenten noves observacions sobre el paper de l'unió A-V en els mecanismes de les taquicardies en el home. Aquestes observacions s'han fet utilitzant la tècnica de estimulació programada del cor. En el capítol III s'il·lustra la complexitat del nodo A-V humà per mitjà de modes poc comuns d'inici de la taquicardia. El capítol IV demostra la possibilitat d'interacció entre dos circuits per re-entrada. Una corba discontinua del nodo A-V no té el mateix significat en pacients que s'estudien degut a taquicardia supraventricular documentada que en pacients sense taquicardia documentada. El capítol V ensenya que en l'últim grup de pacients al període refractari de la via nodal ràpida es llarg en direcció retrògrada i impedeix iniciar la taquicardia. En algun d'aquests pacients es pot iniciar una taquicardia després de donar atropina; que disminueix la duració del període refractari de la via ràpida retrògrada. El capítol VI descriu que el patró de conducció A-V durant la fibril·lació auricular no varia degut a la presència de dos passos intranodals. El capítol VII demostra un patró electrocardiogràfic complex degut a una taquicardia ventricular en un pacient amb doble pas intranodal. Al capítol VIII s'explica com el tò autonòmic té un paper important en els mecanismes de la taquicardia incessant A-V per re-entrada. Als capítols IX i X s'ensenya com es pot arribar al diagnòstic correcte del mecanisme de les taquicardies incessants supraventriculars fent servir l'electrocardiograma, els resultats d'administració de drogues, la prova d'esfors, maniobres vagals i l'estudi electrofisiològic.