Dealing with a life changing event

Citation for published version (APA):

Wachelder, E. M., Moulaert, V. R. M. P., van Heugten, C., Gorgels, A., Wade, D. T., & Verbunt, J. A. (2016). Dealing with a life changing event: The influence of spirituality and coping style on quality of life after survival of a cardiac arrest or myocardial infarction. *Resuscitation*, 109, 81-86.
<https://doi.org/10.1016/j.resuscitation.2016.09.025>

Document status and date:

Published: 01/12/2016

DOI:

[10.1016/j.resuscitation.2016.09.025](https://doi.org/10.1016/j.resuscitation.2016.09.025)

Document Version:

Publisher's PDF, also known as Version of record

Document license:

Taverne

Please check the document version of this publication:

- A submitted manuscript is the version of the article upon submission and before peer-review. There can be important differences between the submitted version and the official published version of record. People interested in the research are advised to contact the author for the final version of the publication, or visit the DOI to the publisher's website.
- The final author version and the galley proof are versions of the publication after peer review.
- The final published version features the final layout of the paper including the volume, issue and page numbers.

[Link to publication](#)

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal.

If the publication is distributed under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license above, please follow below link for the End User Agreement:

www.umlib.nl/taverne-license

Take down policy

If you believe that this document breaches copyright please contact us at:

repository@maastrichtuniversity.nl

providing details and we will investigate your claim.

Clinical paper

Dealing with a life changing event: The influence of spirituality and coping style on quality of life after survival of a cardiac arrest or myocardial infarction[☆]

E.M. Wachelder^{a,*}, V.R.M.P. Moolaert^{a,b}, C. van Heugten^{c,d}, T. Gorgels^e, D.T. Wade^f, J.A. Verbunt^{a,b}

^a Adelante Centre of Expertise in Rehabilitation and Audiology, Hoensbroek, The Netherlands

^b Department of Rehabilitation Medicine, Care and Public Health Research Institute, Maastricht University Medical Centre, Maastricht, The Netherlands

^c School for Mental Health and Neuroscience, Maastricht University, Maastricht, The Netherlands

^d Department of Neuropsychology and Psychopharmacology, Maastricht University, Maastricht, The Netherlands

^e Maastricht University Hospital, Department of Cardiology, Maastricht, The Netherlands

^f Oxford Centre for Enablement, Oxford, United Kingdom

ARTICLE INFO

Article history:

Received 11 February 2016

Received in revised form 29 June 2016

Accepted 16 September 2016

Keywords:

Heart arrest
Quality of life
Spirituality

ABSTRACT

Background: Survivors of a cardiac arrest often have cognitive and emotional problems. As a cardiac arrest is also an obvious life-threatening event, other psychological sequelae associated with surviving such as spirituality may also affect quality of life.

Objectives: To determine the relationship between spirituality, coping and quality of life in cardiac patients both with and without a cardiac arrest.

Methods: In this retrospective cohort study, participants received a questionnaire by post. The primary outcome measure was quality of life (LiSat-9). Secondary outcome measures were spiritual well-being (FACIT-Sp12), coping style (UPCC), emotional well-being (HADS, IES), fatigue (FSS) and daily activities (FAI). Statistical analyses included multiple regression analyses.

Results: Data were available from 72 (60% response rate) cardiac arrest survivors and 98 (47%) patients with a myocardial infarction. Against our hypothesis, there were no differences in spirituality or other variables between the groups, with the exception of more depressive symptoms in patients with myocardial infarction *without* arrest. Analysis of the total data set (170 participants) found that a better quality of life was associated with higher levels of meaning and peace in life, higher levels of social and leisure activities, and lower levels of fatigue.

Conclusions: Quality of life after a cardiac arrest and after a myocardial infarction without arrest are not different; fatigue, a sense of meaning and peace, and level of extended daily activities are factors related to higher life satisfaction.

© 2016 Published by Elsevier Ireland Ltd.

Introduction

In the Netherlands, the overall incidence of cardiac arrest is 0.6–0.9 per 1000 persons per year.^{1,2} During a cardiac arrest, the brain suffers from hypoxia, which may cause diffuse hypoxic–ischemic injury and often results in long-term cognitive impairments.^{3–5} Furthermore, cardiac arrest survivors often report severe fatigue,⁶ feelings of anxiety and/or depression.^{6,7}

and posttraumatic stress reactions.⁶ Fortunately, but also remarkably, quality of life (QoL) after cardiac arrest seems only slightly decreased compared to the general population.⁸

A closer look at QoL after cardiac arrest reveals that it is less associated with medical variables, but more with cognitive complaints, anxiety/depression, posttraumatic stress reactions and difficulties in daily activities.^{6,9,10} This may implicate that differences in responding to the traumatic event of a cardiac arrest may account for variances in quality of life. Two examples of influence are spirituality and coping style.

Spirituality can be defined as a framework that provides people with a sense of ultimate purpose and meaning in life, offering people stability, support and direction in critical times.¹¹ Increasing interest has been shown in the association between spiritual

[☆] A Spanish translated version of the abstract of this article appears as Appendix in the final online version at <http://dx.doi.org/10.1016/j.resuscitation.2016.09.025>.

* Corresponding author at: Adelante Centre of Expertise in Rehabilitation and Audiology, P.O. Box 88, 6430 AB Hoensbroek, The Netherlands.

E-mail address: j.verbunt@mumc.nl (E.M. Wachelder).

well-being and QoL.^{12–14} Patients with terminal cancer, HIV and heart failure and a higher level of spiritual well-being seemed less depressed.^{15,16} Furthermore, higher spiritual well-being may result in better life satisfaction.^{11,17} As surviving a cardiac arrest can be a life changing event, it may also change a patient's outlook on life and spirituality.¹⁸

Coping has been defined as 'the person's cognitive and behavioural efforts to manage the internal and external demands of the person-environment transaction that is appraised as taxing or exceeding the person's resources'.¹⁹ In heart failure patients coping appeared to influence emotional state.²⁰ In patients with subarachnoid haemorrhage, a passive coping style influenced health-related QoL negatively²¹ and in brain injury patients, both increases in active problem-focused coping and decreases in passive emotion-focused coping predicted a higher QoL.²²

So far, the relationship between spirituality, coping style and QoL has not been studied in cardiac arrest survivors. The aim of the present study was to investigate how spiritual well-being and coping are related to QoL two years after a cardiac arrest.

We hypothesized that cardiac arrest survivors, as a consequence of their brain injury, would have a lower level of functioning regarding emotional well-being and fatigue than cardiac patients without a cardiac arrest. In addition, we hypothesized that cardiac arrest survivors would report higher spiritual well-being, as we considered that a cardiac arrest would influence life more as compared to a myocardial infarction. Moreover, we hypothesized a positive contribution of higher spiritual well-being and active coping style on QoL for both cardiac patients with and without a cardiac arrest.

Methods

For this study, data were collected from two groups: cardiac arrest survivors and patients who suffered from a myocardial infarction without cardiac arrest. Patients with a myocardial infarction without cardiac arrest were chosen as a control group, as they were considered most comparable in baseline characteristics such as age, gender and underlying cardiac pathology, and they also were likely to perceive a risk of sudden death. By comparing both groups, the additional impact on life of a cardiac arrest will be studied.

Potential participants with a cardiac arrest were derived from a database of a prospective cohort study called Activity and Life after Survival of a Cardiac Arrest (ALASCA).²³ The inclusion criteria for the ALASCA study were: (1) survivor of a cardiac arrest, (2) living within 50 km of a participating hospital, (3) age 18 years or older, and (4) sufficient knowledge of the Dutch language. Patients were recruited from coronary care units and intensive care units of Dutch hospitals from April 2007 till 2010 and followed for one year. For the present study, ALASCA-participants with a cardiac arrest before April 2009 who had given consent to be approached for follow-up studies were invited to participate two years after the event.

Participants with a myocardial infarction without cardiac arrest were identified out of patient files of the Department of Cardiology of the University Hospital Maastricht, The Netherlands. Patients admitted to the department between July 2008 and January 2009 with myocardial infarction, aged 18 years or older and with sufficient knowledge of the Dutch language were identified as potential participants and were also invited to participate two years after the event.

Data collection

Two years after their cardiac event, all potential participants were invited to participate in the study by sending them an information letter, a set of questionnaires and an informed consent form by post. One reminder was sent. The study protocol was approved

by the Medical Ethics Committee of the University Hospital Maastricht.

Measurement instruments

Socio-demographic variables regarding marital status, living situation and work situation were collected by self-report.

Quality of life, the main outcome of interest, was measured using the **Life Satisfaction Questionnaire (LiSat-9)**, designed to assess the level of satisfaction with the present living situation. It consists of nine items (satisfaction with life in general and with eight life domains), scoring from 1 (very dissatisfied) to 6 (very satisfied). Its internal consistency is satisfactory (Cronbach alpha 0.74).²⁴

Spiritual well being was measured using the **Functional Assessment of Chronic Illness Therapy–Spiritual Well Being (FACIT-Sp12)** scale, a 12-item questionnaire measuring overall spiritual well-being divided over 2 subscales (meaning/peace and faith) with item scores ranging from 0 (not at all) to 4 (very much). Both subscales have a high internal consistency, and show good reliability and validity.^{15,25}

Coping style was measured using the **Utrecht Proactive Coping Competence scale (UPCC)**, a questionnaire assessing an individual's competency with regard to proactive coping, using 21 items with scores ranging from 1 (not at all capable) to 4 (capable). Cronbach alpha ranges from 0.83 to 0.95.²⁶

Emotional well-being was measured using the **Hospital Anxiety and Depression Scale (HADS)**, a questionnaire designed to assess mood disorders in non-psychiatric hospital outpatients. It consists of 14 items divided in two subscales, directed at measuring signs of anxiety and depression. Validity, internal consistency and reliability are good.²⁷

The **Impact of Event Scale (IES)**, a 15-item questionnaire assessing stress reactions after traumatic events was used to measure the significance of the event to the particular patient. Validity and reliability are good.²⁸

Fatigue was assessed using the **Fatigue Severity Scale (FSS)**, a 9-item questionnaire assessing the impact of fatigue on patients' functioning. Each item is scored on a 7-point scale and the total score is the overall mean (range 1–7). The scale has good psychometric properties.²⁹

Extended activities of daily living were measured using the **Frenchay Activities Index (FAI)**, a 15-item questionnaire for instrumental activities of daily living (ADL) with item scores ranging from 0 to 3. It is valid and reliable measure as assessed in stroke.^{30,31}

Statistical analysis

Possible age- or gender related differences between responders and non-responders in both groups were investigated using *t*-test or Mann–Whitney testing (depending on parametric distribution) for age and chi-square testing for gender. To detect baseline differences between groups, similar tests were performed for age and gender, Mann–Whitney testing for time since event and chi-square testing for current marital status and living situation.

In case of <15% missing values, missing data were imputed by the mean score per instrument. In case of ≥ 15% missing values in a questionnaire, a total score was not calculated. In the cardiac arrest group, scores on all questionnaires, except the FAI, showed a non-parametric distribution.

To detect statistical significant differences between variables in both groups, Mann–Whitney 2 testing was performed. A *p*-value of ≤0.05 was considered statistically significant.

Multiple linear regression analyses (ENTER method) were carried with independent variables spiritual well-being (subscales 'meaning & peace', 'faith' (FACIT-Sp12), coping style (UPCC), emo-

Fig. 1. Flow chart of patient inclusion.

tional well being (HADS and IES), fatigue (FSS), daily functioning (FAI) and group (cardiac arrest versus myocardial infarction)) and the dependent variable QoL (LiSat-9).

To rule out different pathways or relationships within both groups, potential interactions of the variable “study group” with the variables “spiritual well-being”, “emotional well-being”, “age” and “gender” were analysed. A similar check was performed for interaction between spiritual and emotional well-being. Age and gender were introduced in the model as potential confounders. A limit value for elimination in the regression analyses was set at $p < 0.10$ and a p -value ≤ 0.05 was considered statistically significant. Regression models were checked for co-linearity and outliers. Analyses were performed using SPSS software version 20.0 (SPSS Inc., Chicago, Illinois, USA).

Results

Fig. 1 shows that 72 patients (response rate 60%) with a cardiac arrest and 98 patients (response rate 47%) with a myocardial infarction participated.

Table 1 shows the characteristics of all participants. There were neither statistically significant gender-related nor age-related differences between responders and non-responders in the cardiac arrest group and the myocardial infarction group. Participants with a myocardial infarction were significantly older and more often female as compared to the cardiac arrest group.

Current level of functioning

Table 2 shows scores for basic functioning, activities in daily living and QoL (following the ICF-domains³²).

In general, the level of functioning on different domains is good for both groups, except for fatigue, which is present in over half of all participants. Also, the level of functioning is comparable between both groups. Only the total score of the HADS and its subscale depression were higher in the myocardial infarction group compared with the cardiac arrest group.

Participants were also asked if they suffered from any other medical problems influencing their functioning in daily life. This was the case in 19 (26%) cardiac arrest survivors and 37 (38%) patients in the control group (chi-square = 2.505, $p = 0.113$).

Variables related to quality of life

Regression analysis did not confirm different pathways or relationships between variables within groups, as all analyses to test interactions and confounders were negative (results not shown). Therefore, the group of both cardiac arrest and myocardial infarction patients were analysed together to study the contribution of spiritual well-being and coping, emotional well-being, fatigue and daily functioning on QoL.

Table 3 shows that the subscale ‘meaning & peace’ of the FACIT-Sp12 contributed positively to QoL; a higher sense of meaning and peace was associated with higher life satisfaction. Also, less fatigue and a more independent level of daily functioning were associated with a higher quality of life, explaining 59% of the variance in the model. Coping style and emotional well-being did not have any significant effect on QoL.

Discussion

This study found that cardiac arrest does not have a specific impact on quality of life in people with cardiac disease, and that the factors associated with higher life satisfaction two years after an acute cardiac event are a higher level of social and leisure activities, a lower level of fatigue and a higher level of a sense of meaning and peace. Coping style had no influence.

In this study, cardiac arrest survivors were compared with myocardial infarction patients, and we expected that cardiac arrest survivors would experience more emotional problems and fatigue (as a consequence of hypoxic brain injury), and would have a higher level of spiritual well-being (due to the higher impact of the event). In fact we found a quite good and also comparable level of functioning in both groups. *Fatigue* was notably a frequently reported problem in both groups, and a lower level of *emotional well-being* (especially depressive symptoms) was found in the group who had a myocardial infarction without cardiac arrest. The presence of emotional problems (30% in the cardiac arrest group and 40% in the myocardial infarction group) is comparable with numbers found in stroke patients.^{33,34}

The lower level of *emotional well-being* in the myocardial infarction group may, to some extent, be explained by differences in age and gender. Previous studies described that older age and female gender were negatively related to the patient’s functioning and

Table 1
Demographic characteristics of the participants.

	Cardiac arrest (N = 72) Current situation	Myocardial infarction (N = 98) Current situation	p-Value
Time since cardiac event (days)			0.70
Median	744	748	
Range	698–932	712–783	
Age			<0.001
Mean (SD)	59.8 (11.5)	66.6 (11.7)	
Range	29–82	37–90	
Gender			0.002
Male	66 (92%)	71 (72%)	
Female	6 (8%)	27 (28%)	
Marital status			0.08
Married/living with partner	57 (79%)	73 (74%)	
Unmarried/divorced/widowed	3 (4%)	20 (20%)	
Unknown	12 (17%)	5 (5%)	
Living situation			0.03
Home (independent)	61 (85%)	69 (70%)	
Home (with help from others)	6 (8%)	24 (25%)	
Sheltered housing	–	1 (1%)	
Nursing home	–	1 (1%)	
Unknown	5 (7%)	3 (3%)	

Table 2
Basic functioning, activities in daily living and quality of life of the participants.

ICF domain	Range of questionnaire	Cut-offscore	Cardiac arrest (N = 72)		Myocardial infarction (N = 98)		p-Value
			Mean (SD)/median (range)	No (%) impaired	Mean (SD)/median (range)	No (%) impaired	
Basic functioning							
Spiritual well-being (FACIT-Sp12)	0–48	–	32 (4–46)	–	31 (8–48)	–	0.41
Emotional well-being (HADS)	0–42	≥11	5 (0–31)	21 (30%)	9 (0–36)	38 (40%)	0.02
– Subscale anxiety	0–21	≥9	3 (0–15)	16 (23%)	4 (0–20)	17 (18%)	0.17
– Subscale depression	0–21	≥9	2 (0–20)	14 (20%)	4 (0–19)	19 (20%)	0.01
Stress reactions (IES)	0–75	≥26	11 (0–63)	19 (27%)	11 (0–59)	22 (23%)	0.51
Fatigue (FSS)	1–7	≥4	4.0 (1–7)	35 (51%)	4.3 (1–7)	57 (60%)	0.07
Activities of daily living							
Instrumental daily activities (FAI)	0–45	≤16	27.3 (7.4)	6 (9%)	28.0 (1–44)	10 (11%)	0.63
Quality of life							
Life satisfaction (LiSat-9)	1–6	≤4	5.0 (2.8–6.0)	15 (22%)	4.8 (1.9–6.0)	12 (14%)	0.35
Other							
Coping style (UPCC)	1–4	–	3.1 (1–4)	–	3.1 (1.5–4)	–	0.98

Table 3
Final model from multiple linear regression analyses (ENTER method) N = 144.

Dependent variable	Independent variables	R ²	Adjusted R ²	Standardized beta	p-Value
LiSat-9		0.59	0.57		0.00
	Spiritual well-being; meaning/peace (FACIT-Sp 12)			0.50	0.00
	Spiritual well-being; faith (FACIT-Sp12)			–0.05	0.40
	Coping style (UPCC)			0.05	0.50
	Emotional well-being (HADS)			0.03	0.80
	Stress reactions (IES)			–0.11	0.15
	Fatigue (FSS)			–0.25	0.00
	Instrumental daily activities (FAI)			0.14	0.04
	Group (cardiac arrest versus myocardial infarction)			–0.04	0.49

Dependent variable: LiSat-9, Life Satisfaction Questionnaire.

Independent variables: FACIT-Sp12, Functional Assessment of Chronic Illness Therapy—Spiritual Well-being scale, score on subscale meaning/peace and faith; total score on UPCC, Utrecht Proactive Coping Competence scale; HADS, Hospital Anxiety and Depression Scale; IES, Impact of Event Scale; FSS, Fatigue Severity Scale; FAI, Frenchay Activities Index; and research group (cardiac arrest group or myocardial infarction).

QoL.^{35,36} In addition, differences in medical condition may also explain some of the findings: patients with a myocardial infarction can have negative illness perceptions which increase over time and can result in increased feelings of *fatigue*.³⁷ Also, use of medication may have affected fatigue and emotional wellbeing.

Finally, the analysis of the cohort was done irrespective of treatment. However, about one third of the cardiac arrest survivors

received a new early intervention service during their participation in the ALASCA study and this intervention appeared to be effective in improving mood and quality of life.³⁸ This therefore may have led to better overall levels of functioning in the cardiac arrest group.

The absence of higher levels of *spirituality* in the cardiac arrest group may be due to underestimation of the impact on life of a myocardial infarction. Based on our results, it could be hypothe-

sized that the impact of an event such as a myocardial infarction is experienced as equally life threatening, and may therefore be equally life changing with similar effects on spiritual well-being. This theory seems supported by comparable mean scores on the Impact of Event Scale for both groups. Although the mean score is relatively low, the range is large indicating that a subgroup of patients within both groups is still experiencing significant stress reactions related to the cardiac event two years later.

The construct of spirituality is complex and still under much debate. Some suggest it might be regarded as a construct, which is present in the whole being of a person to a more or less extent but will not change much over time. Then, serving as a coping resource and helping people dealing better with adversity, the level of spiritual well-being within a person might eventually define the impact of an event.³⁹ So a more spiritual person might be more capable dealing with a cardiac event and as a result be less depressed and experience a higher QoL. Others however suggest that spirituality is a developmental process within a person,⁴⁰ and that posttraumatic growth of spirituality can be seen.^{41,42} More research is needed in this field.

Spirituality, fatigue and daily activities contributed significantly to QoL, whereas coping style and emotional well-being did not. We checked whether age, gender, or group were possible interacting or confounding factors, but this was not the case.

A previous study showed that QoL was related to emotional problems, fatigue, daily activities and cognitive complaints.¹⁰ In the present study cognition was not taken into account. The frequency of emotional problems was relatively high. However, since emotional problems did not contribute to QoL, spirituality may have overpowered the impact of emotional problems. The fact that coping style was not related to QoL was unexpected, as in other cardiovascular diseases an influence was shown.²⁰

There are several limitations to our study. First, the recruitment of patients with a cardiac arrest and with a myocardial infarction without cardiac arrest differed. Patients with a cardiac arrest were originally asked to participate in the ALASCA study. Baseline data were collected immediately after the event. Patients with a myocardial infarction were invited two years after the incident and for baseline data we had to rely on patient's files. Selection bias is possible, due to the fact that more patient with cardiac arrest responded than those with a myocardial infarction (60% versus 47%). Cardiac arrest survivors were already part of an on-going follow up study (the ALASCA study) which may have made them more committed to respond. One would expect a smaller percentage of the myocardial infarction group to have neurocognitive consequences as the brain is most likely not involved in the event. Selection bias is also possible due to the fact that there is no further background information available regarding the non-responders.

Secondly, even though there is increasing interest in spirituality and coping, and their effect on functioning and QoL, these are still concepts difficult to understand and study.

Last, many other factors, not assessed in our study, may show associations with QoL. For instance, Brands et al. showed that self-efficacy predicts long-term QoL in brain injured patients.⁴³ Further research is necessary.

Further research based on prospective studies with larger samples is needed. Both on the subject of spirituality and coping and the influence on QoL. This way, spirituality and coping can be taken more into account when treating patients after a cardiac event. In a recent qualitative study nurses experiences in spiritual care were examined and the authors concluded that spirituality could be of relevance to all areas of nursing care.⁴⁴ As a clinical implication of the current findings can be mentioned that for both patients with a cardiac arrest and a cardiac infarction, it seems important to focus on spirituality as it positively contributed to QoL, whereas fatigue influenced it in a negative way. Learning patients how to

cope with fatigue could probably add to their QoL. Whether, in addition, spirituality can also be influenced in a positive way needs further study.

Conclusion

This study investigated the influence of spirituality and coping style on QoL after a cardiac arrest. Spirituality positively contributed to QoL, whereas coping style did not have an additional effect. Furthermore, fatigue and daily functioning also contributed significantly. Spirituality and fatigue are both factors that should be addressed after a cardiac event.

Conflict of interest statement

We hereby declare that none of the participating authors of the manuscript "Dealing with a life changing event: the influence of spirituality and coping style on quality of life after survival of a cardiac arrest" had any potential conflicts regarding financial, personal or other relationships that might have biased their work.

Potential conflict of interest: none declared.

Appendix A. Supplementary data

Supplementary data associated with this article can be found, in the online version, at <http://dx.doi.org/10.1016/j.resuscitation.2016.09.025>.

References

- Gorgels AP, Gijsbers C, de Vreede Swagemakers J, Lousberg A, Wellens HJ. Out-of-hospital cardiac arrest—the relevance of heart failure. The Maastricht Circulatory Arrest Registry. *Eur Heart J* 2003;24:1204–9.
- Waalwijk RA, de Vos R, Koster RW. Out-of-hospital cardiac arrests in Amsterdam and its surrounding areas: results from the Amsterdam resuscitation study (ARREST) in 'Utstein' style. *Resuscitation* 1998;38:157–67.
- Lim C, Alexander MP, LaFleche G, Schnyer DM, Verfaellie M. The neurological and cognitive sequelae of cardiac arrest. *Neurology* 2004;63:1774–8.
- Sauve MJ, Doolittle N, Walker JA, Paul SM, Scheinman MM. Factors associated with cognitive recovery after cardiopulmonary resuscitation. *Am J Crit Care* 1996;5:127–39.
- Moulaert VR, Verbunt JA, van Heugten CM, Wade DT. Cognitive impairments in survivors of out-of-hospital cardiac arrest: a systematic review. *Resuscitation* 2009;80:297–305. Epub 2009/01/02.
- Wachelder EM, Moulaert VR, van Heugten C, Verbunt JA, Bekkers SC, Wade DT. Life after survival: long-term daily functioning and quality of life after an out-of-hospital cardiac arrest. *Resuscitation* 2009;80:517–22. Epub 2009/03/14.
- de Vos R, de Haes HC, Koster RW, de Haan RJ. Quality of survival after cardiopulmonary resuscitation. *Arch Intern Med* 1999;159:249–54.
- Elliott VJ, Rodgers DL, Brett SJ. Systematic review of quality of life and other patient-centred outcomes after cardiac arrest survival. *Resuscitation* 2011;82:247–56. Epub 2011/01/11.
- Stiell I, Nichol G, Wells G, et al. Health-related quality of life is better for cardiac arrest survivors who received citizen cardiopulmonary resuscitation. *Circulation* 2003;108:1939–44. Epub 2003/10/08.
- Moulaert VR, Wachelder EM, Verbunt JA, Wade DT, van Heugten CM. Determinants of quality of life in survivors of cardiac arrest. *J Rehabil Med* 2010;42:553–8. Epub 2010/06/16.
- Hill PC, Pargament KI. Advances in the conceptualization and measurement of religion and spirituality. Implications for physical and mental health research. *Am Psychol* 2003;58:64–74.
- Powell LH, Shahabi L, Thoresen CE. Religion and spirituality. Linkages to physical health. *Am Psychol* 2003;58:36–52.
- Miller WR, Thoresen CE. Spirituality, religion, and health. An emerging research field. *Am Psychol* 2003;58:24–35.
- Seeman TE, Dubin LF, Seeman M. Religiosity/spirituality and health. A critical review of the evidence for biological pathways. *Am Psychol* 2003;58:53–63.
- Bekelman DB, Dy SM, Becker DM, et al. Spiritual well-being and depression in patients with heart failure. *J Gen Intern Med* 2007;22:470–7.
- McClain CS, Rosenfeld B, Breitbart W. Effect of spiritual well-being on end-of-life despair in terminally-ill cancer patients. *Lancet* 2003;361:1603–7. Epub 2003/05/16.
- Kennedy JE, Abbott RA, Rosenberg BS. Changes in spirituality and well-being in a retreat program for cardiac patients. *Altern Ther Health Med* 2002;8:64–6, 8–70, 2–3. Epub 2002/07/20.

18. Parnia S, Spearpoint K, Fenwick PB. Near death experiences, cognitive function and psychological outcomes of surviving cardiac arrest. *Resuscitation* 2007;74:215–21.
19. Folkman S, Lazarus RS, Gruen RJ, DeLongis A. Appraisal, coping, health status, and psychological symptoms. *J Pers Soc Psychol* 1986;50:571–9. Epub 1986/03/01.
20. Doering LV, Dracup K, Caldwell MA, et al. Is coping style linked to emotional states in heart failure patients? *J Card Fail* 2004;10:344–9. Epub 2004/08/17.
21. Passier PE, Visser-Meily JM, van Zandvoort MJ, Rinkel GJ, Lindeman E, Post MW. Predictors of long-term health-related quality of life in patients with aneurysmal subarachnoid hemorrhage. *NeuroRehabilitation* 2012;30:137–45. Epub 2012/03/21.
22. Wolters G, Stapert S, Brands I, Van Heugten C. Coping styles in relation to cognitive rehabilitation and quality of life after brain injury. *Neuropsychol Rehabil* 2010;20:587–600. Epub 2010/05/07.
23. Moolaert VR, Verbunt JA, van Heugten CM, et al. Activity and Life After Survival of a Cardiac Arrest (ALASCA) and the effectiveness of an early intervention service: design of a randomised controlled trial. *BMC Cardiovasc Disord* 2007;7:26. Epub 2007/08/29.
24. Fugl Meyer AR, Melin R, Fugl Meyer KS. Life satisfaction in 18- to 64-year-old Swedes: in relation to gender, age, partner and immigrant status. *J Rehabil Med* 2002;34:239–46.
25. Peterman AH, Fitchett G, Brady MJ, Hernandez L, Cella D. Measuring spiritual well-being in people with cancer: the functional assessment of chronic illness therapy—Spiritual Well-being Scale (FACIT-Sp). *Ann Behav Med* 2002;24:49–58.
26. Bode C, Thoolen B, de Ridder D. Measuring proactive coping. Psychometric characteristics of the Utrecht Proactive Competence scale (UPCC). *Psychologie en Gezondheid* 2008;32:81–91.
27. Bjelland I, Dahl AA, Haug TT, Neckelmann D. The validity of the hospital anxiety and depression scale: an updated literature review. *J Psychosom Res* 2002;52:69–77.
28. Sundin EC, Horowitz MJ. Impact of Event Scale: psychometric properties. *Br J Psychiatry* 2002;180:205–9.
29. Krupp LB, LaRocca NG, Muir Nash J, Steinberg AD. The fatigue severity scale. Application to patients with multiple sclerosis and systemic lupus erythematosus. *Arch Neurol* 1989;46:1121–3.
30. Schuling J, de Haan R, Limburg M, Groenier KH. The Frenchay Activities Index. Assessment of functional status in stroke patients. *Stroke* 1993;24:1173–7.
31. Wade DT, Legh Smith J, Langton Hewer R. Social activities after stroke: measurement and natural history using the Frenchay Activities Index. *Int Rehabil Med* 1985;7:176–81.
32. World Health Organization. International Classification of Functioning, Disability and Health (ICF). WHO; 2001.
33. Bour A, Rasquin S, Aben I, Boreas A, Limburg M, Verhey F. A one-year follow-up study into the course of depression after stroke. *J Nutr Health Aging* 2010;14:488–93. Epub 2010/07/10.
34. van Mierlo ML, van Heugten CM, Post MW, de Kort PL, Visser-Meily JM. Psychological factors determine depressive symptomatology after stroke. *Arch Phys Med Rehabil* 2015;96:1064–70. Epub 2015/02/15.
35. Bunch TJ, White RD, Khan AH, Packer DL. Impact of age on long-term survival and quality of life following out-of-hospital cardiac arrest. *Crit Care Med* 2004;32:963–7.
36. van Alem AP, Waalewijn RA, Koster RW, de Vos R. Assessment of quality of life and cognitive function after out-of-hospital cardiac arrest with successful resuscitation. *Am J Cardiol* 2004;93:131–5.
37. Alsen P, Brink E, Persson LO, Brandstrom Y, Karlson BW. Illness perceptions after myocardial infarction: relations to fatigue, emotional distress, and health-related quality of life. *J Cardiovasc Nurs* 2010;25:E1–10. Epub 2010/02/20.
38. Moolaert VR, van Heugten CM, Winkens B, et al. Early neurologically-focused follow-up after cardiac arrest improves quality of life at one year: a randomised controlled trial. *Int J Cardiol* 2015;193:8–16. Epub 2015/05/26.
39. Koenig HG. Religion, spirituality, and health: the research and clinical implications. *ISRN Psychiatry* 2012;2012:278730. Epub 2012/01/01.
40. Shek DT. Spirituality as a positive youth development construct: a conceptual review. *Sci World J* 2012;2012:458953. Epub 2012/06/02.
41. Picoraro JA, Womer JW, Kazak AE, Feudtner C. Posttraumatic growth in parents and pediatric patients. *J Palliat Med* 2014;17:209–18. Epub 2014/01/22.
42. Schaefer FC, Blazer DG, Koenig HG. Religious and spiritual factors and the consequences of trauma: a review and model of the interrelationship. *Int J Psychiatry Med* 2008;38:507–24. Epub 2008/01/01.
43. Brands I, Kohler S, Stapert S, Wade D, van Heugten C. Influence of self-efficacy and coping on quality of life and social participation after acquired brain injury: a 1-year follow-up study. *Arch Phys Med Rehabil* 2014;95:2327–34. Epub 2014/06/29.
44. Giske T, Cone PH. Discerning the healing path—how nurses assist patient spirituality in diverse health care settings. *J Clin Nurs* 2015;24:2926–2935. Epub 2015/07/29.