Subgrouping patients with chronic low back pain

Citation for published version (APA):

Huijnen, I. P. J., Schasfoort, F. C., Smeets, R. J. E. M., Sneekes, E., Verbunt, J. A., & Busmann, J. B. J. (2020). Subgrouping patients with chronic low back pain: What are the differences in actual daily life behavior between patients classified as avoider or persister? *Journal of Back and Musculoskeletal Rehabilitation*, 33(2), 303-311. <https://doi.org/10.3233/BMR-171048>

Document status and date:

Published: 01/01/2020

DOI:

[10.3233/BMR-171048](https://doi.org/10.3233/BMR-171048)

Document Version:

Publisher's PDF, also known as Version of record

Document license:

Taverne

Please check the document version of this publication:

- A submitted manuscript is the version of the article upon submission and before peer-review. There can be important differences between the submitted version and the official published version of record. People interested in the research are advised to contact the author for the final version of the publication, or visit the DOI to the publisher's website.
- The final author version and the galley proof are versions of the publication after peer review.
- The final published version features the final layout of the paper including the volume, issue and page numbers.

[Link to publication](#)

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal.

If the publication is distributed under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license above, please follow below link for the End User Agreement:

www.umlib.nl/taverne-license

Take down policy

If you believe that this document breaches copyright please contact us at:

repository@maastrichtuniversity.nl

providing details and we will investigate your claim.

Subgrouping patients with chronic low back pain: What are the differences in actual daily life behavior between patients classified as avoider or persister?

Ivan P.J. Huijnen^{a,b,1,*}, Fabienne C. Schasfoort^{c,d,1}, Rob J.E.M. Smeets^{a,e}, Emiel Sneekes^{c,d}, Jeanine A. Verbunt^{a,b} and Johannes B.J. Bussmann^c

^aDepartment of Rehabilitation Medicine, Research Program Functioning and Rehabilitation, Research School CAPHRI, Maastricht University, Maastricht, The Netherlands

^bAdelante, Centre of Expertise in Rehabilitation and Audiology, Hoensbroek, The Netherlands

^cDepartment of Rehabilitation Medicine, Erasmus MC University Medical Center Rotterdam, Rotterdam, The Netherlands

^dRijndam Rehabilitation, Rotterdam, The Netherlands

^eCIR Revalidatie Location Eindhoven, Eindhoven, The Netherlands

Abstract.

BACKGROUND AND OBJECTIVE: The purpose of this study is to determine whether patients, classified by their treating consultant in rehabilitation medicine as avoider or persister, show differences in a large set of detailed outcomes of actual, objectively measured daily physical behaviour.

METHODS: In this explorative cross-sectional study, 16 patients were included; 9 patients were categorized as avoider and 7 patients as persister. Subjects wore the VitaMove activity monitor, a high-end accelerometry-based device that allowed automatic detection of a large set of body postures and motions. Physical behaviour was assessed in detail by total duration of body postures and motions as percentages of 24 hours, as well as by the number of sit-to-stand transfers, overall activity level, walking speed, and the distribution of bouts of physical activity and sedentary behaviour. Differences between groups were tested with the Mann Whitney U test.

RESULTS: There were no significant differences between groups in any of the physical behaviour outcomes.

CONCLUSIONS: Our study showed that activity-related behavioural style categorization by consultants in rehabilitation medicine is not expressed in objectively measured detailed outcomes of daily physical behaviour.

Keywords: Activity monitoring, activity-related behaviour, chronic low back pain, excessive persistence, avoidance

1. Introduction

Chronic low back pain (CLBP) is a major public health problem with a high impact on daily life functioning [1]. Interdisciplinary rehabilitation treatment therefore generally aims at reducing limitations in daily life functioning of this patient group [2]. Since not all patients seem to benefit from the same ap-

¹Both authors equally contributed to the organization and execution of the study presented in this manuscript.

*Corresponding author: Ivan P.J. Huijnen, Department of Rehabilitation Medicine, Maastricht University, P.O. Box 616, 6200 MD Maastricht, The Netherlands. Tel.: +31 43 3882162; E-mail: ivan.huijnen@maastrichtuniversity.nl.

proach [2], tailoring treatment to individual patient's coping strategies may increase effectiveness [3]. Valid classification systems to create subgroups lack, however.

In recent years, research has increasingly focused on avoidance and persistence behaviour as two maladaptive activity-related behavioural styles in patients with CLBP. Avoidance behaviour is assumed to be a result of catastrophizing thoughts about pain and the development of fear of movement. Eventually, this will lead to lower levels of daily life functioning. Such avoidance behaviour has been confirmed in patients with pain [4]. In persistence behaviour, other mechanisms are thought to explain a patient's disability level. Although persons with a persistence behavioural style perceive themselves as disabled, it is assumed that they continue their activities despite their pain. These patients will therefore have a daily life activity level similar to people without pain or other impairments [5,6].

A relatively small number of studies has focused on (disentangling the nature of) the relationship between the constructs avoidance and/or persistence on the one hand, and physical activity levels/daily functioning outcome measures on the other hand. For example, Huijnen and colleagues found that patients' self-reported daily life activity levels were significantly higher in 'persisters' compared to 'avoiders', whereas such a difference was not present when daily life activity levels were objectively measured with a simple accelerometer device [7]. It was also found that persisters, in comparison to avoiders, reported a longer daily uptime (defined as the period between getting up and going to sleep), using an objective as well as a self-reported outcome measure [7,8]. Andrews et al. have studied the relationship between an individual's self-reported activity-related behavioural style and patterns of both pain and objectively measured physical activity [9]. They did not find a significant association between avoidance behaviour and lower daily life activity levels, nor was there a significant association between persistence behaviour and higher daily life activity levels. There was some evidence, however, that individuals who were classified as 'excessive' persister (i.e. high self-reported level of persistence combined with low level of avoidance) are more likely to engage in daily life activity in a way that exacerbates pain complaints. Overall, the existing evidence appears to be inconclusive.

This may be related to discrepancies between self-reported/perceived activity levels and actual daily life activity levels: simply because it is difficult to assess

one's own or another person's physical activity levels findings are not easily interpretable [10]. It has been shown that experienced physicians both under- and overestimated the level of objectively measured daily activity levels in several rehabilitation patient groups, including a significant underestimation in patients with chronic benign pain [11].

The inconclusiveness may also be related to the questionable ability of simple monitoring devices to validly measure avoidance and (excessive) persistence behaviour, as pointed out by Andrews et al. [12]. It might be that maladaptive styles of patients are reflected in other aspects of people's physical behaviour than the overall daily life activity level, which is the main outcome of most simple monitoring devices [6]. For example, avoiders might avoid specific feared activities or postures, but may continue to perform other non-feared activities. As a result, maladaptive styles may not be expressed in e.g. a decline in the overall daily life activity level, but might become overt in other physical behaviour outcomes, such as the distribution of activities and postures over time. So far, actual physical behaviour of patients with CLBP has not been analysed in more detail in studies aiming to identify differences between avoiders and persisters; for this, a more advanced and high-end accelerometer device is required. The high-end monitoring device VitaMove provides detailed outcomes on types, intensity and patterns of physical behaviour, and therefore allows a more in depth evaluation of whether avoiders and persisters differ in their actual daily life behaviour [13–15].

Therefore, the aim of this explorative study was to evaluate whether patients with CLBP categorized as avoider or persister by consultants in rehabilitation medicine show differences in detailed outcomes of actual, objectively measured daily physical behaviour.

2. Methods

2.1. Participants

Consultants in rehabilitation medicine with ample expertise in chronic pain treatment from Maastricht University Medical Centre, Rijndam Rehabilitation Centre Rotterdam, and Top-Care Health Services/CIRAN assessed patients diagnosed with CLBP for eligibility to participate in this cross-sectional study. Inclusion criteria were: a) low back pain: pain localized below the scapulae and above the gluteal

folds for longer than six months [16]; b) age between 18 and 65 years. Exclusion criteria were: a) low back pain attributable to a recognizable, known specific pathology (e.g. infection, tumour, osteoporosis, fracture, structural deformity, inflammatory disorder, radicular syndrome or cauda equina syndrome) [17]; b) pregnancy; c) wearing a pacemaker or ICD device (i.e. safety measure related to activity monitoring); d) serious psychiatric disorders; e) non-fluency in Dutch. The Medical Ethics Committee of Rotterdam, The Netherlands, approved the study (METC-2011-378).

2.2. Study procedure

Prior to enrolment, all subjects received verbal and written information about the study. After the subjects agreed to participate, written informed consent was obtained, and subjects completed a set of self-reported measures on disability, activity-related strategies, fear of movement, habitual physical activity level, pain intensity level, and depressed mood. Also, an appointment was made to visit the hospital's human performance laboratory to have a short training session on using the activity monitoring device. Participants left the university hospital with the monitoring device, and wore it for preferably 5 days during daytime. After the measurement period, the device was returned to the university hospital for automated analyses.

Classification of patients as 'avoider' or 'persister' regarding their activity related behaviour style was done by a consultant in rehabilitation medicine. The consultant discussed a number of specific persistence and avoidance statements with participants. These statements were based on different dimensions of avoidance and persistence behaviour and were derived from a study of Kindermans et al. [18]. In that study, it was found that activity avoidance and excessive persistence (i.e. doing too much, not respecting one's physical limits and experiencing a rebound effect of over-activity) were the only dimensions of avoidance and persistence behaviour associated with a higher disability level. The 13 items from the Behavioral Responses to Illness Questionnaire (BRIQ) and 20 items from the Patterns of Activity Measure-Pain (POAM-P) loading on these factors in the study of Kindermans et al., were used as statements during the consultation to support clinical judgement of the consultant in rehabilitation medicine to classify patients as avoider or persister. Not all of these 33 statements had to be discussed, however. Classification was finished as

Fig. 1. Adult person wearing a VitaMove activity monitor (2M Engineering, Veldhoven, The Netherlands).

soon as the expert consultant in rehabilitation medicine was convinced about which activity-related behaviour style the patient used.

2.3. Main outcome measure: Actual physical behaviour

Actual physical behaviour was measured using the advanced high-end VitaMove activity monitor (2M Engineering, Veldhoven, The Netherlands; see Fig. 1). This Activity monitor consists of three body-fixed accelerometer units (Freescale MMA7260Q, Denver, USA). The VitaMove sensor units (one attached to each thigh, one to the sternum part of the trunk, using elastic belts) are wirelessly connected. Acceleration signals were continuously sampled with 128 Hz, and stored on micro SD memory cards. The recorders were ideally worn during waking hours on at least 5 consecutive days (and preferably including 1 weekend day). Patients were instructed to continue their normal daily life activities, but recorders had to be de-attached during swimming, bathing and sleeping.

After the measurement period, data were uploaded to a PC, for analysis with VitaScore Software (VitaScore BV, Gemert, The Netherlands). This analysis software allowed a one-second detection of a large set of body postures (e.g. lying, sitting, standing), body motions (e.g. walking, running, cycling, non-cyclic movements), and body transitions (e.g. sit-to-stand). Body postures and motions with a duration < 5 s were disregarded. Additionally, a motility signal was created, expressing the intensity of body postures and motions, based on the variability of all mea-

sured accelerometer signals. A detailed description of the configuration and analysis has been described elsewhere [19]. The VitaMove system is the successor of the Vitaport Activity Monitor, which is extensively validated to quantify body postures and motions, and often applied in research [11,19–23].

From the raw signals the start and end of the wear period was determined. Subsequently, from the VitaScore analyses the following physical behaviour outcomes were calculated over all days and averaged:

1. The total duration – expressed as % of 24 hours – of the composite measure of the duration of all body motions and standing (expressing physical activity)
2. The total duration (as % of 24 hours) of the composite measure of the duration of sitting and lying (expressing daytime sedentary behaviour)
3. The duration – expressed as % of 24 hours – of each body posture and motion
4. Number of sit-to-stand transitions
5. Mean motility of all activities expressed in g ($1\text{ g} = 9.81\text{ m/s}^2$), as an overall measure of the amount of physical activity
6. The mean motility during walking periods, expressed in g ($1\text{ g} = 9.81\text{ m/s}^2$), as an indicator of walking speed [24,25]

Subsequently, the VitaScore body posture and motion output signal was exported for distribution analyses using a custom-made MatLab algorithm. Binary (0/1) 1-second time resolution signals were created, expressing bouts (i.e. a time period of a defined activity category) of the physical behaviour domains physical activity (body motions and standing), and sedentary behaviour (sitting and lying). Based on these bout signals several distribution outcomes were calculated [13,14,26]. The analyses resulted in the following outcome variables, calculated for both physical behaviour domains separately:

1. Number of bouts; The number of bouts of uninterrupted physical activity or sedentary behaviour
2. Median bout length; The median duration (sec) of bouts
3. Covariance of Variation of bout length; Expresses the variance in bout length
4. Fragmentation; The number of (physical activity or sedentary) bouts are divided by the total duration. A higher score represents a more fragmented behaviour, i.e. consisting of more bouts of a shorter length
5. W-index; Is the total time of bouts above the me-

dian bout length divided by the total duration of physical activity or sedentary behaviour. The W-index will be higher in case there is a larger contribution of longer bouts in the total time

2.4. Descriptive assessment: Demographic and pain-related information

First, demographic (gender, age, education level, ethnic information, work status) and pain-related information (duration of complaints) was obtained, after which fear of movement/(re)injury was assessed with the Dutch version of the Tampa Scale for Kinesiophobia (TSK). The Dutch version has been reported to be reliable and valid [27,28]. Patients were asked to rate their mean pain level perceived in the past week on a 0 to 10 point scale. The level of depression was determined by the Beck Depression Inventory II (BDI-II) [29]. The questionnaire has good psychometric properties and is a valid questionnaire to measure the severity of depression in patients with chronic pain [30,31].

2.5. Self-reported habitual approach to activity

2.5.1. Patterns of Activity Measure-Pain (POAM-P)

The patients' degree of avoidance and persistence behaviour was measured with the POAM-P [32]. The POAM-P is a 30-item self-report questionnaire, developed to measure three activity patterns in patients with chronic pain, namely avoidance (10 items), overdoing (10 items) and pacing (10 items) [18,32]. Since the focus of the current study is on an avoidant versus persistent activity-related strategy, only the results of the subscales avoidance and overdoing were used.

2.5.2. Behavioral Responses to Illness Questionnaire (BRIQ)

The BRIQ is a 21-item self-report questionnaire originally developed to measure behavioural responses of patients in an acute phase of illness [33]. The questionnaire comprises four subscales, but in the present study, only 'all-or-nothing behaviour' and 'limiting behaviour' are selected, as these subscales measure behavioural persistence and avoidance behaviour, respectively. Psychometric properties of the BRIQ are satisfactory [33]. For the study of Kindermans et al., the BRIQ was translated into Dutch and the back translation was approved by the original authors [34].

2.5.3. Disability

Low back pain related disability was assessed using

Table 1
Characteristics of the study population of patients with CLBP

	1. Avoiders (<i>N</i> = 9)	2. Persisters (<i>N</i> = 7)
Male/female	5/4	3/4
Age	50 (37.5–55.0)	46 (45.0–59.0)
Duration of pain		
6–12 months	1	1 ^a
1–5 yrs	6	4 ^a
> 5 yrs	2	0 ^a
Born in The Netherlands (<i>N</i> (%))	8 (88.9%)	6 (85.7%)
Work status (<i>N</i> (%))		
Paid job	5 (55.6%)	4 (57.1%)
Disability payment	2 (22.2%)	2 (28.6%)
Fear of movement (TSK)	42.0 (34.0–47.0)	36.0 (31.0–48.0)
Depressed mood (BDI-II)	15.0 (10.5–22.0)	14.0 (5.0–24.0)
Pain intensity (VAS)	6.00 (5.17–6.83)	6.83 (5.83–7.67) ^a
Avoidance (POAM-P)	25.0 (15.5–31.5)	16.0 (9.0–25.0)
Overdoing (POAM-P)	22.0 (19.5–29.0)	30.0 (25.0–31.0)
BRIQ all or nothing behaviour	18.0 (12.5–18.5)	19.0 (17.0–23.0)
BRIQ limiting behaviour	19.0 (15.5–22.0)	22.0 (13.0–27.0)
Disability (QBPDS)	45.0 (38.0–57.5)	59.5 (30.3–72.5) ^a
Habitual activity level (BPAQ)	8.5 (5.86–8.92)	8.79 (6.01–9.08)

Data are represented by medians (interquartile ranges). ^a*N* = 6. TSK = Tampa Scale for Kinesiophobia. BDI-II = Beck Depression Inventory II. Avoidance = Score on the avoidance subscale of the POAM-P. Overdoing = Score on the overdoing subscale of the POAM-P. BRIQ = Behavioural Responses to Illness Questionnaire. QBPDS = Quebec Back Pain Disability Scale. BPAQ = Baecke Physical Activity Questionnaire.

the Quebec Back Pain Disability Scale (QBPDS). The original as well as the Dutch version of the QBPDS are valid and reliable [35,36].

2.5.4. Habitual physical activity in daily life

The level of daily life activities during the last year was measured with the Baecke Physical Activity Questionnaire (BPAQ) [37]. The reliability of the BPAQ in patients with LBP appears to be sufficient [38].

2.6. Statistical analysis

Descriptive statistics were used for the relevant characteristics of the activity-related styles and the outcome variables used. To evaluate whether the actual, objectively assessed, characteristics of activity-related behaviour are different between avoiders and persisters the Mann Whitney U test was used. In this analysis, alpha was set at 0.05. The analysis was performed using IBM SPSS Statistics for Windows, version 22 (IBM Corp, Armonk, NY, USA).

3. Results

Sixteen patients with CLBP (8 females) participated in this study. Eleven patients were included from

Maastricht University Medical Centre, one from Rijn-dam Rehabilitation, and four patients from CIRAN. No significant differences in activity-related behavioural style, gender and age were present between the inclusion locations. However, patients included in Maastricht felt significantly more disabled measured with the QBPDS ($p < 0.05$).

Overall, 70 measurement days were included in the VitaScore analyses. Due to the criteria used for the analyses, the number of days analysed for one patient was limited to two days. For 2 participants 3 days were analysed, for 3 patients 4 days were analysed, for 10 persons 5 days were analysed, and 22 out of the 70 days were weekend days. The mean length of the measurement day was 14.01 hours (SD = 1.7).

3.1. General characteristics of avoiders and persisters

Table 1 presents characteristics of the avoiders and persisters. Nine patients were classified as avoider and seven patients were classified as persister. No significant differences were found on age, gender and work status between avoiders and persisters. Similarly, no significant differences between these groups were found on other parameters, although avoiders, com-

Table 2
Actual physical behaviour outcomes for the two groups (as measured with VitaMove)

		1. Avoiders ($N = 9$)	2. Persisters ($N = 7$)
Quantity	Daytime (h)	14.00 (12.90–15.25)	13.93 (11.30–16.19)
	Total duration body motions & standing (% of 24 h)	21.02 (15.94–30.43)	20.73 (18.48–25.32)
	Total duration sitting & lying (% of 24 h)	36.72 (31.45–38.15)	32.70 (27.72–46.73)
	Total duration (% of 24 h)		
	Lying	2.46 (0.77–8.27)	2.64 (0.46–8.75)
	Sitting	31.10 (27.88–35.26)	28.42 (25.08–33.73)
	Standing	14.30 (10.90–19.63)	11.65 (11.54–15.86)
	Walking	5.17 (2.59–7.85)	4.44 (2.98–4.80)
	Cycling	0.02 (0.00–0.16)	0.08 (0.00–0.82)
	Non-cyclic movement	2.26 (1.42–3.92)	2.28 (1.68–4.50)
Quality	Number of sit-to-stand transitions	27.67 (17.80–35.25)	22.80 (16.25–31.00)
	Mean motility (g)	8.37 (6.12–13.30)	7.98 (6.58–10.66)
	Mean motility during walking (g)	46.20 (44.34–58.30)	49.00 (45.50–52.20)
	Number of bouts	92.67 (59.23–127.40)	93.50 (63.50–129.00)
	Median bout length (sec)	77.70 (53.75–115.53)	52.13 (29.40–82.13)
	CoV bout length (sec)	40.60 (35.55–42.74)	43.58 (41.97–44.97)
	Fragmentation	0.0039 (0.0033–0.0069)	0.0045 (0.0041–0.0073)
	W-index	0.94 (0.93–0.96)	0.95 (0.93–0.96)
	Number of bouts	86.33 (57.57–123.00)	89.00 (60.50–127.75)
	Median bout length (sec)	78.50 (52.52–199.75)	82.63 (69.90–136.60)
Distribution outcomes physical activity	CoV bout length (sec)	40.17 (37.37–44.33)	40.15 (38.53–45.38)
	Fragmentation	0.0030 (0.0019–0.0043)	0.0029 (0.0025–0.0032)
	W-index	0.96 (0.95–0.97)	0.96 (0.95–0.96)
	W-index		
Distribution outcomes sedentary behaviour	Number of bouts	86.33 (57.57–123.00)	89.00 (60.50–127.75)
	Median bout length (sec)	78.50 (52.52–199.75)	82.63 (69.90–136.60)
	CoV bout length (sec)	40.17 (37.37–44.33)	40.15 (38.53–45.38)
	Fragmentation	0.0030 (0.0019–0.0043)	0.0029 (0.0025–0.0032)
	W-index	0.96 (0.95–0.97)	0.96 (0.95–0.96)

pared to the persister group, showed a higher median score on fear of movement (42.0 vs 36.0), more avoidance behaviour as measured with the POAM-P (25.0 vs 16.0) and a lower score on overdoing (22.0 vs 30.0). Also no significant differences were found in the disability level or the self-reported daily life activity level.

3.2. Actual daily life activities of avoiders and persisters

Table 2 presents the median scores and interquartile ranges of the activity monitor data for avoiders and persisters. No significant differences in percentages sedentary or body motions were found between avoiders and persisters. Also no significant differences in distribution outcomes for the two different constructs of physical behaviour were found.

To check the validity of the results, two types of post-hoc analyses were performed in which the Mann Whitney U test was repeated. The first post-hoc analyses focused on differences between avoiders and persisters when data of specific days were analysed; weekend days and week days (avoiders $n = 9$, persisters $n = 7$), and data of work days and non-work days (avoiders $n = 5$; persisters $n = 4$). These post-hoc analyses also did not show a significant difference between avoiders and persisters for any of the outcomes. Secondly, to check for the potential effect of our strategy of patient categorization, patients scoring most ex-

treme on activity avoidance ($n = 5$) and excessive persistence ($n = 5$) – based on their scores on the relevant questions of the POAM-P and BRIQ – were selected. Again, for none of the outcomes significant differences between groups were found.

4. Discussion

This study had an explorative character, and aimed to evaluate whether patients displaying avoidance and persistence behavioural styles present differences in objectively measured actual physical behaviour. We were not able to confirm that patients classified as avoider or persister differ in their actual physical behaviour. Patients scored similarly on all VitaMove quantitative and qualitative outcomes, as well as on all distribution outcomes. The innovative aspect of this study was that physical behaviour was measured in detail and was expressed in a broad set of outcome measures for actual physical behaviour. Nevertheless, no differences were found. This is in line with other studies evaluating actual daily activity levels with relatively simple accelerometer devices [7,9]. The finding that persisters have a longer daily uptime – both reported and objectively measured [7,8] – was not confirmed by the wear time data in our study, however.

More detailed and in depth analysis of the quantity and distribution of body postures and movements

thus also did not reveal differences between patients categorized as avoider or persister. Literature indicates that differences between activity-related behavioural styles might indeed be more strongly expressed in self-reported outcomes [7,8,39]. However, in the present study also no significant differences in self-reported physical activities in daily life or the patient's disability level were found between both subgroups. This was also the case in the study of Hasenbring et al., where no differences in self-reported daily life activity levels between both styles were found in 24 patients [39]. This may be related to insufficient statistical power, because in a previous study in a larger group of 79 patients, a significant difference for self-reported daily life activity level between avoiders and persisters has been found [7].

Another explanation for the absence of differences between groups might be related to the categorization of patients. In our study, the classification was done by a consultant in rehabilitation medicine. The consultant discussed a number of specific persistence and avoidance statements derived from the POAM-P and BRIQ questionnaires. Selection of statements was based on a study of Kindermans [18]. That study showed that although two types of avoidance behaviour (pain avoidance and activity avoidance) and three types of persistence behaviour (task-contingent persistence, pain-contingent persistence, and excessive persistence) could be distinguished, only excessive persistence and activity avoidance appeared to be related to a higher level of disability [18]. Only questions of excessive persistence and activity avoidance were used as statement and discussed by the consultant in rehabilitation medicine to decide whether a patient was either persister or avoider. Patients classified as persister can therefore be seen as an excessive persister, whereas patients classified as avoider can be seen as an activity avoider. It should, however, be taken into account that the validity and reliability of this method is still unknown. In a study of Esteve et al. the types of activity-related behaviour distinguished by Kindermans et al. to develop the Activity Pattern Scale (APS) were also used and provided evidence that this instrument is valid and reliable to assess activity patterns [40]. This finding supports the classification method used in our study.

Although the samples of avoiders and persisters were relatively small and the median scores seemed different between both styles, it is remarkable that the subscale of overdoing and avoidance of the POAM-P as well as the BRIQ subscales "all or nothing be-

haviour" and "limiting behaviour" showed no significant differences between both subgroups in our study. Several other studies used the POAM-P [7], the Physical Activity Relations Questionnaire (PARQ) [8,9] and the Avoidance Endurance Questionnaire (AEQ) [41] to classify patients in subgroups. The activity-related styles measured in these questionnaires measure different constructs of persistence behaviour. However, excessive persistence, as identified in the study of Kindermans et al., as a maladaptive style, was not considered as a unique and only style. The different questionnaires contain questions of the different types of persistence behaviour. Therefore, in the present study only the statements of the maladaptive avoidance and persistence styles were used to classify patients in avoider or persister. However, as discussed earlier, no differences between avoiders and persisters could be found in neither detailed outcomes of actual physical behaviour nor in the self-reported daily life activity level. Differences in results between studies may depend on the method or instruments used to categorize patients in subgroups based on their activity-related behavioural style.

Besides the issue of categorizing patients, other explanations for not confirming our assumption that avoiders and persisters differ in detailed objectively measured physical behaviour may be related to the way the detailed VitaMove outcome measures were measured and calculated. For instance, several factors, such as weather and seasonality, will influence the physical behaviour outcomes. These might have led to increased variability in outcomes, but given the random order of measurements, no systematic effect on the data can be expected. Furthermore, differences might become overt by including data of specific days, such as weekend and weekdays, and work and non-work days. The post-hoc analyses we performed (comparing smaller persister and avoider subgroups), however, did not indicate that the effect of these factors was significant and did not result in other conclusions. We agree with Andrews et al. [12] that objective monitoring devices should be used in concordance with self-reported outcome for now. Both types of outcome measures have advantages and disadvantages related to validity and bias.

The most important limitation of our study was that only sixteen patients were included. Therefore, we have to be careful with interpreting the results because of the small sample and the impact on statistical power. More research is needed with a larger sample to replicate our findings and to make individually tailored treatment based on individual patient's coping strategies possible.

5. Conclusion

Based on the results presented here it can be concluded that patients classified by consultants in rehabilitation medicine as avoider or persister do not differ in detailed aspects of actual physical behaviour as assessed by objective accelerometry. For clinical practice, it seems important to realize that although patients present a certain activity-related behavioural style, this may not necessarily be reflected in their actual daily life activities.

Acknowledgments

This study was financially supported by Erasmus MC Pijnfonds. The authors also would like to thank Loes Swaan (Rijndam Rehabilitation), Malou Fanchamps (Erasmus MC), Bart Pepels (Maastricht University), Marco van Woensel, Jan Joosten, Joost Volmer, Jan Jochijms and Fons van Bergen (Top-Care Health Services/CIRAN).

Conflict of interest

The authors have no conflicts of interest relevant to this work.

References

- [1] Global Burden of Disease Study C. Global, regional, and national incidence, prevalence, and years lived with disability for 301 acute and chronic diseases and injuries in 188 countries, 1990–2013: a systematic analysis for the Global Burden of Disease Study 2013. *Lancet*. 2015; 386(9995): 743–800.
- [2] Kamper SJ, Apeldoorn AT, Chiarotto A, Smeets RJ, Ostelo RW, Guzman J, et al. Multidisciplinary biopsychosocial rehabilitation for chronic low back pain: cochrane systematic review and meta-analysis. *BMJ*. 2015; 350: h444.
- [3] Murphy SL. Overactivity in daily life: a crucial step in understanding how to tailor treatments. *Pain*. 2015; 156(10): 1831–2.
- [4] Leeuw M, Goossens ME, Linton SJ, Crombez G, Boersma K, Vlaeyen JW. The fear-avoidance model of musculoskeletal pain: current state of scientific evidence. *J Behav Med*. 2007; 30(1): 77–94.
- [5] van den Berg-Emons RJ, Schasfoort FC, de Vos LA, Bussmann JB, Stam HJ. Impact of chronic pain on everyday physical activity. *Eur J Pain*. 2007; 11(5): 587–93.
- [6] Verbunt JA, Westerterp KR, van der Heijden GJ, Seelen HA, Vlaeyen JW, Knottnerus JA. Physical activity in daily life in patients with chronic low back pain. *Arch Phys Med Rehabil*. 2001; 82(6): 726–30.
- [7] Huijnen IP, Verbunt JA, Peters ML, Smeets RJ, Kindermans HP, Roelofs J, et al. Differences in activity-related behaviour among patients with chronic low back pain. *Eur J Pain*. 2011; 15(7): 748–55.
- [8] McCracken LM, Samuel VM. The role of avoidance, pacing, and other activity patterns in chronic pain. *Pain*. 2007; 130(1–2): 119–25.
- [9] Andrews NE, Strong J, Meredith PJ. Overactivity in chronic pain: is it a valid construct? *Pain*. 2015; 156(10): 1991–2000.
- [10] Huijnen IP, Verbunt JA, Peters ML, Delespaul P, Kindermans HP, Roelofs J, et al. Do depression and pain intensity interfere with physical activity in daily life in patients with Chronic Low Back Pain? *Pain*. 2010; 150(1): 161–6.
- [11] van den Berg-Emons RJ, Bussmann JB, Stam HJ. Accelerometry-based activity spectrum in persons with chronic physical conditions. *Arch Phys Med Rehabil*. 2010; 91(12): 1856–61.
- [12] Andrews NE, Strong J, Meredith PJ. Avoidance or incapacitation: a discussion on definition and validity of objective measures of avoidance, persistence, and overactivity. *Clin J Pain*. 2015; 31(7): 670–2.
- [13] Chastin SF, Granat MH. Methods for objective measure, quantification and analysis of sedentary behaviour and inactivity. *Gait Posture*. 2010; 31(1): 82–6.
- [14] Lord S, Chastin SF, McInnes L, Little L, Briggs P, Rochester L. Exploring patterns of daily physical and sedentary behaviour in community-dwelling older adults. *Age Ageing*. 2011; 40(2): 205–10.
- [15] Sanchez MC, Bussmann J, Janssen W, Horemans H, Chastin S, Heijenbrok M, et al. Accelerometric assessment of different dimensions of natural walking during the first year after stroke: recovery of amount, distribution, quality and speed of walking. *J Rehabil Med*. 2015; 47(8): 714–21.
- [16] Merskey H, Bogduk N. Classification of chronic pain. edition S, editor. Seattle: IASP press; 1994.
- [17] Airaksinen O, Brox JJ, Cedraschi C, Hildebrandt J, Klaber-Moffett J, Kovacs F, et al. Chapter 4. European guidelines for the management of chronic nonspecific low back pain. *Eur Spine J*. 2006; 15(Suppl 2): S192–300.
- [18] Kindermans HP, Roelofs J, Goossens ME, Huijnen IP, Verbunt JA, Vlaeyen JW. Activity patterns in chronic pain: underlying dimensions and associations with disability and depressed mood. *J Pain*. 2011; 12(10): 1049–58.
- [19] Bussmann JB, Martens WL, Tulen JH, Schasfoort FC, van den Berg-Emons HJ, Stam HJ. Measuring daily behavior using ambulatory accelerometry: the Activity Monitor. *Behav Res Methods Instrum Comput*. 2001; 33(3): 349–56.
- [20] Bussmann HB, Reuvekamp PJ, Veltink PH, Martens WL, Stam HJ. Validity and reliability of measurements obtained with an “activity monitor” in people with and without a transtibial amputation. *Phys Ther*. 1998; 78(9): 989–98.
- [21] Bussmann JB, Tulen JH, van Herel EC, Stam HJ. Quantification of physical activities by means of ambulatory accelerometry: a validation study. *Psychophysiology*. 1998; 35(5): 488–96.
- [22] Bussmann JB, van de Laar YM, Neeleman MP, Stam HJ. Ambulatory accelerometry to quantify motor behaviour in patients after failed back surgery: a validation study. *Pain*. 1998; 74(2–3): 153–61.
- [23] van den Berg-Emons HJ, Bussmann JB, Balk AH, Stam HJ. Validity of ambulatory accelerometry to quantify physical activity in heart failure. *Scand J Rehabil Med*. 2000; 32(4): 187–92.

- [24] Bussmann JB, Hartgerink I, van der Woude LH, Stam HJ. Measuring physical strain during ambulation with accelerometry. *Med Sci Sports Exerc.* 2000; 32(8): 1462-71.
- [25] Bussmann JB, van den Berg-Emons HJ, Angulo SM, Stijnen T, Stam HJ. Sensitivity and reproducibility of accelerometry and heart rate in physical strain assessment during prosthetic gait. *Eur J Appl Physiol.* 2004; 91(1): 71-8.
- [26] Tieges Z, Mead G, Allerhand M, Duncan F, van Wijck F, Fitzsimons C, et al. Sedentary behavior in the first year after stroke: a longitudinal cohort study with objective measures. *Arch Phys Med Rehabil.* 2015; 96(1): 15-23.
- [27] Goubert L, Crombez G, Vlaeyen J, Van Damme S, Van Den Broeck A, Van Houdenhove B. De tampa schaal voor kinesiofobie: psychometrische karakteristieken en normering (the tampa scale kinesiophobia: psychometric properties and norms). *Gedrag en Gezondheid.* 2000; 28: 54-62.
- [28] Vlaeyen JW, Kole-Snijders AM, Boeren RG, van Eek H. Fear of movement/(re)injury in chronic low back pain and its relation to behavioral performance. *Pain.* 1995; 62(3): 363-72.
- [29] Beck AT, Steer RA, Brown GK, Van der Does AJW. BDI-II-NL Handleiding (BDI-II-Dutch Manual). Lisse: The Netherlands: Psychological Corporation; 2002.
- [30] Gretebeck RJ, Montoye HJ. Variability of some objective measures of physical activity. *Med Sci Sports Exerc.* 1992; 24(10): 1167-72.
- [31] Harris CA, D'Eon JL. Psychometric properties of the Beck Depression Inventory-second edition (BDI-II) in individuals with chronic pain. *Pain.* 2008; 137(3): 609-22.
- [32] Cane D, Nielson WR, McCarthy M, Mazmanian D. Pain-related activity patterns: measurement, interrelationships, and associations with psychosocial functioning. *The Clinical Journal of Pain.* 2013; 29(5): 435-42.
- [33] Spence M, Moss-Morris R, Chalder T. The Behavioural Responses to Illness Questionnaire (BRIQ): a new predictive measure of medically unexplained symptoms following acute infection. *Psychol Med.* 2005; 35(4): 583-93.
- [34] Kindermans HP, Huijnen IP, Goossens ME, Roelofs J, Verbunt JA, Vlaeyen JW. "Being" in pain: the role of self-discrepancies in the emotional experience and activity patterns of patients with chronic low back pain. *Pain.* 2011; 152(2): 403-9.
- [35] Kopec JA, Esdaile JM, Abrahamowicz M, Abenhaim L, Wood-Dauphinee S, Lamping DL, et al. The quebec back pain disability scale. Measurement properties. *Spine.* 1995; 20(3): 341-52.
- [36] Schoppink LE, van Tulder MW, Koes BW, Beurskens SA, de Bie RA. Reliability and validity of the dutch adaptation of the quebec back pain disability scale. *Phys Ther.* 1996; 76(3): 268-75.
- [37] Baecke JA, Burema J, Frijters JE. A short questionnaire for the measurement of habitual physical activity in epidemiological studies. *Am J Clin Nutr.* 1982; 36(5): 936-42.
- [38] Jacob T, Baras M, Zeev A, Epstein L. Low back pain: reliability of a set of pain measurement tools. *Arch Phys Med Rehabil.* 2001; 82(6): 735-42.
- [39] Hasenbring MI, Plaas H, Fischbein B, Willburger R. The relationship between activity and pain in patients 6 months after lumbar disc surgery: do pain-related coping modes act as moderator variables? *Eur J Pain.* 2006; 10(8): 701-9.
- [40] Esteve R, Ramirez-Maestre C, Peters ML, Serrano-Ibanez ER, Ruiz-Parraga GT, Lopez-Martinez AE. Development and initial validation of the activity patterns scale in patients with chronic pain. *J Pain.* 2016; 17(4): 451-61.
- [41] Hasenbring MI, Hallner D, Klasen B, Streitlein-Bohme I, Willburger R, Rusche H. Pain-related avoidance versus endurance in primary care patients with subacute back pain: psychological characteristics and outcome at a 6-month follow-up. *Pain.* 2012; 153(1): 211-7.