Social role functioning in Parkinson's disease

Citation for published version (APA):

Perepezko, K., Hinkle, J. T., Shepard, M. D., Fischer, N., Broen, M. P. G., Leentjens, A. F. G., Gallo, J. J., & Pontone, G. M. (2019). Social role functioning in Parkinson's disease: A mixed-methods systematic review. *International Journal of Geriatric Psychiatry*, 34(8), 1128-1138. <https://doi.org/10.1002/gps.5137>

Document status and date:

Published: 01/08/2019

DOI:

[10.1002/gps.5137](https://doi.org/10.1002/gps.5137)

Document Version:

Publisher's PDF, also known as Version of record

Document license:

Taverne

Please check the document version of this publication:

- A submitted manuscript is the version of the article upon submission and before peer-review. There can be important differences between the submitted version and the official published version of record. People interested in the research are advised to contact the author for the final version of the publication, or visit the DOI to the publisher's website.
- The final author version and the galley proof are versions of the publication after peer review.
- The final published version features the final layout of the paper including the volume, issue and page numbers.

[Link to publication](#)

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal.

If the publication is distributed under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license above, please follow below link for the End User Agreement:

www.umlib.nl/taverne-license

Take down policy

If you believe that this document breaches copyright please contact us at:

repository@maastrichtuniversity.nl

providing details and we will investigate your claim.

Social role functioning in Parkinson's disease: A mixed-methods systematic review

Kate Perepezko^{1,5} | Jared T. Hinkle^{1,2} | Melissa D. Shepard¹ | Nicole Fischer¹ | Martinus P.G. Broen³ | Albert F.G. Leentjens⁴ | Joseph J. Gallo⁵ | Gregory M. Pontone¹

¹Department of Psychiatry and Behavioral Sciences, Johns Hopkins University School of Medicine, Baltimore, Maryland

²Johns Hopkins School of Medicine, Medical Scientist Training Program, Baltimore, Maryland

³Department of Neurology, Maastricht University Medical Center, Maastricht, the Netherlands

⁴Department of Psychiatry, Maastricht University Medical Center, Maastricht, the Netherlands

⁵Department of Mental Health, Johns Hopkins Bloomberg School of Public Health, Baltimore, Maryland

Correspondence

Kate Perepezko, Department of Psychiatry and Behavioral Sciences, Johns Hopkins University School of Medicine, Baltimore, MD.
Email: kperepe1@jhu.edu

Objectives: Parkinson's disease (PD) is a progressive neurodegenerative disease that often impedes activities of daily living (ADL) and social functioning. Impairment in these areas can alter social roles by interfering with employment status, household management, friendships, and other relationships. Understanding how PD affects social functioning can help clinicians choose management strategies that mitigate these changes.

Methods: We conducted a mixed-methods systematic review of existing literature on social roles and social functioning in PD. A tailored search strategy in five databases identified 51 full-text reports that fulfilled the inclusion criteria and passed the quality appraisal. We aggregated and analyzed the results from these studies and then created a narrative summary.

Results: Our review demonstrates how PD causes many people to withdraw from their accustomed social roles and experience deficits in corresponding activities. We describe how PD symptoms (eg, tremor, facial masking, and neuropsychiatric symptoms) interfere with relationships (eg, couple, friends, and family) and precipitate earlier departure from the workforce. Additionally, several studies demonstrated that conventional PD therapy has little positive effect on social role functioning.

Conclusions: Our report presents critical insight into how PD affects social functioning and gives direction to future studies and interventions (eg, couple counseling and recreational activities).

KEYWORDS

caregiver, Parkinson's disease, quality of life, social roles

1 | INTRODUCTION

Parkinson's disease (PD) is a progressive neurodegenerative disease characterized by motor and nonmotor symptoms that impair functioning in activities of daily living (ADL) and cause changes in social functioning.^{1,2} Social functioning encompasses performance in specific social roles, or "expected ways of behaving," which are established by both an individual's personal goals and societal norms.³ In chronic illnesses, symptoms impact the ability to fulfill social roles such as employment, household management,

friendships, and other relationships.⁴ Changes in social role functioning are particularly troublesome because satisfaction with social role performance is related to overall happiness and quality of life (QoL).⁵ Additionally, social isolation and loneliness is a risk factor for depression, cognitive decline, increased health-care costs, and overall mortality.⁶⁻⁹

QoL has become a frequent outcome of interest in clinical trials and is one of the most important factors for determining clinical care for people with chronic diseases.^{10,11} In the PD field, research on QoL has increased, and several systematic reviews describe the

determinants of QoL, economic impact of decreased QoL, how to measure QoL, and prognostic factors related to QoL.¹²⁻¹⁵ Van Uem et al recently published a review on health-related QoL in patients with PD using the International Classification of Functioning, Disability and Health (ICF) developed by the World Health Organization as a framework. The ICF model was designed to represent health and disability at individual and population levels, using the following domains: body functions and structures, ADL, participation in social roles, personal features, and environmental factors.¹⁶ This review found that poor health-related QoL was most strongly associated with social role functioning in PD.¹⁷ Because no review has examined specific aspects of social functioning in PD and social functioning contributes greatly to QoL, we conducted a systematic review to identify what is currently known about this subject.

2 | METHODS

In order to capture the breadth of research on social role functioning, we incorporated studies estimating the association of social function with PD employing statistical methods and qualitative studies seeking to understand the construct of social functioning in PD. Inclusion criteria for the screening were as follows: (a) the PD participants and (b) at least one outcome of the paper focused on social functioning or a social role.

2.1 | Search strategy and study selection

A literature search was performed in March 2018, using several MeSH terms related to social roles (Table S1). All articles published before March 1, 2018, were included in the search, yielding 7129 articles from Pubmed (2257), Embase (3300), PsychINFO (1078), CINAHL (373), and Cochrane (121). After removing duplicates ($n = 1872$), the titles and abstracts of the remaining 5257 articles were screened by two reviewers (J.H. and K.P.). Articles were excluded from the review if they did not meet inclusion criteria, were not available in English, or if they did not present original research. A third reviewer (M.S.) served as an adjudicator when there was disagreement about inclusion in the full-text evaluation. In order to provide the most comprehensive review of papers investigating social functioning, we included studies employing a variety of instruments and data collection methods. However, because of the heterogeneity of instruments used, we were unable to conduct a meta-analysis.

After the initial screening, 4843 articles were excluded, leaving 414 articles in the full text evaluation (Figure 1). The full texts were then read to confirm they met inclusion criteria. After full text review, 344 articles were excluded, leaving 70 articles for quality appraisal and data extraction. We also searched the reference sections of the articles that were included after full-text review for any additional papers that may have been missed in the database search (two additional articles were identified). Table S2 describes the reasons for excluding these articles after full-text review.

Key points

- PD can affect performance in different social roles or may cause withdrawal from these roles.
- Standard pharmacologic and surgical interventions have little positive effect on social role functioning in PD.
- Our review identified several correlates of impaired social role performance in PD, including disease severity, anxiety, depression, and cognitive impairment.
- A few studies found that activities like dancing or support groups improved social role functioning and quality of life. Future research should focus on a broader range of interventions to improve social role functioning.

2.2 | Data extraction and quality appraisal of included studies

One reviewer (K.P.) used a quality appraisal checklist to independently evaluate the quality of the 70 articles remaining after the full-text assessment. We used the Critical Appraisal Skills Program checklist for the quality appraisal.¹⁸ The checklist type was selected based on the study design. The appraisal resulted in the inclusion of 51 articles and the exclusion of 19 articles. One reviewer (K.P.) extracted data from the 51 articles that remained after the quality appraisal.

2.3 | Data synthesis

Our mixed-methods approach facilitated a synthesis of published results and enabled us to produce a narrative summary of existing work. Using the Cochrane procedure for narrative summary, one reviewer (K.P.) read through each paper and first developed a preliminary composition of the results (Tables S3 to S8).¹⁹ A combination of inductive and deductive reasoning was applied for the analysis. We employed deductive synthesis to first divide articles based on the type of social role described: couple relationship, parent/family role, work role, friendship role, social/leisure role, or grouped into a general social functioning category if no specific role was described. Some papers discussed multiple roles; the information relevant to each role was extracted from the paper and categorized in the appropriate section. Then, within each group, we applied an inductive synthesis to identify patterns of topics discussed and further divide papers into a sub-theme. The data from studies were then translated using thematic analysis to describe common conclusions across different papers (within the previously defined social roles) (Table 1).

3 | RESULTS

We identified 51 papers to include in our analysis, all published between 1973 and 2018. Of these, 24 papers employed primarily quantitative

FIGURE 1 Consort diagram of the review process [Colour figure can be viewed at wileyonlinelibrary.com]

methods, 23 were qualitative, and four used a mixed-methods approach. Most study designs were cross sectional ($n = 44$), and the remaining articles were longitudinal ($n = 7$). The papers included in this review either covered general social functioning ($n = 16$) or discussed a specific social role, including the couple relationship ($n = 36$), parent/family role ($n = 14$), friendship role ($n = 9$), work role ($n = 19$), and/or social/leisure role ($n = 6$). Figure 2 depicts the distribution of articles based on methodology used and social role addressed.

3.1 | Quantitative studies: Instruments

A wide variety of instruments were used to measure social role functioning. Some studies included full questionnaires such as the Marital Adjustment Test. Other studies included questionnaires that assessed social functioning in a subsection of the full scale, for example, the Nottingham Health Profile.

3.2 | Qualitative studies: Data collection methods

Most of the qualitative studies employed in-depth interviews as the main data collection method and some employed focus group discussions to supplement the interviews. Questions that were included in the interviews were generally open ended (eg, "Can you tell me what your life is like with Parkinson's disease?"), some focused on specific social roles (eg, "Can you describe how the disease has affected your

relationship with family, friends, and your community?"), while others were broad (eg, "How has your 'usual state of health' changed after diagnosis?").

3.3 | Narrative analysis of papers by social role category

3.3.1 | General social functioning

General social functioning included papers ($n = 16$) that described overall socialization or social functioning but did not discuss a specific social role. These articles were categorized into three subthemes—clinical observations/PD symptoms ($n = 4$), treatment/intervention effect ($n = 9$), and QoL/life changes after PD ($n = 3$). Gunnery et al described that facial masking was associated with social functioning problems such as social rejection, although this finding was attenuated after controlling for depression.²⁰ Miller et al described how communication problems (ie, voice problems) were associated with changes in socialization such as social withdrawal.²¹ People living with Parkinson's felt that PD restricted their activity, decreased their socialization, and limited their ability to have a "meaningful" social contribution.²¹⁻²³ Papers also described treatments and interventions to improve Parkinson's symptoms including "typical" PD treatments (ie, medication and deep brain stimulation [DBS] surgery), exercise interventions (eg, tango class), and educational programs (eg, psychoeducation). Quantitative studies describing DBS and levodopa

TABLE 1 Summary of results from review papers

Social Role	Subtheme	Summary of Results
General social functioning	Clinical observations/PD symptoms	<ul style="list-style-type: none"> •Facial masking is correlated with social rejection. •Communication problems (eg, voice problems) can lead to social withdrawal. •PD can limit socialization and ability to feel capable of making a meaningful contribution.
	Treatment/intervention effect	<ul style="list-style-type: none"> •A tango class helped increase social support. •There is little evidence that traditional therapies (medication and DBS) improve social role functioning. •For some people, social functioning can decline after DBS surgery; however, psychoeducation can address this decline.
	QoL/life changes after PD	<ul style="list-style-type: none"> •Cognitive functioning is related to social integration. •Females report worse social support than males.
Primary relationships/couple	Clinical/PD symptoms	<ul style="list-style-type: none"> •Facial masking and disease severity were associated with worse partner interaction. •The added responsibility of taking care of a spouse with PD can strain relationships. •Ability to cope with changes predicted psychosocial functioning.
	Relationship satisfaction (nonsexual aspects)	<ul style="list-style-type: none"> •Nonsexual relationship aspects (eg, talking and sharing activities) can become more important after PD diagnosis. •Female participants still valued being perceived as a wife and mother. •Greater relationship mutuality predicted better QoL.
	Sexual satisfaction	<ul style="list-style-type: none"> •Sexual satisfaction was lower for younger onset People with PD. •Sex can become problematic because of decreased drive, fatigue, and rigidity. •Sexual satisfaction was associated with marital satisfaction. •Predictors of quality of sexual life include male gender, better motor functioning, and quality of sexual life for partner.
	Treatment effect	<ul style="list-style-type: none"> •Higher levodopa doses were associated with increased thoughts of breaking up. •People with PD who participated in a dance class with their spouse found it increased moral support. •The effect of DBS on the couple role is mixed.
Parent/family role	Treatment/intervention	<ul style="list-style-type: none"> •A tango intervention improved family role functioning. •After DBS people with PD typically had improved relations with their children.
	Sharing disease/communication	<ul style="list-style-type: none"> •Family relationships are affected early in the disease. •Some people with PD feel the need to "protect" their families from the diagnosis. •Some people with PD feel sharing their diagnosis with their families helps them better manage.
	Relationship satisfaction	<ul style="list-style-type: none"> •Sometimes there is no change in family relationship quality after diagnosis. •For some parenting becomes more important after diagnosis.
Friendship role	Number of social contacts/social connectedness	<ul style="list-style-type: none"> •People with PD are less likely to have many close friends. •The number of friends may stay the same, but typically people with PD initiate fewer visits to friends. •Some people conceal themselves from others, which disrupts social connectedness.
	Quality of life Relationship quality	<ul style="list-style-type: none"> •Changes in social contacts also substantially influence quality of life. •Some friendships were strengthened while others lost friends because of the disease. •Identification and involvement with the larger PD community was a meaningful source of friendship and support.
Work role	Clinical/PD symptoms/predictors	<ul style="list-style-type: none"> •Age, anxiety, fatigue, motor symptoms, apathy, and cognition interfere with work performance.
	Housework Work unavailability/leaving the workforce	<ul style="list-style-type: none"> •People with PD were less likely to participate in household management. •People with PD are less likely to work. •Some people try to keep their jobs as long as they can and worry about losing social connections from work. •Reasons for leaving work include being unable to meet demands. •People who continue to work try to hide their symptoms and try to maintain job performance rather than working toward a promotion.
	Treatment/intervention	<ul style="list-style-type: none"> •No evidence that levodopa influences work role. •Work activity often became worse after DBS.
Social and leisure role	Activity type	<ul style="list-style-type: none"> •More likely to engage in solitary and/or sedentary activities.
	Treatment/intervention effect	<ul style="list-style-type: none"> •No evidence that levodopa helps with social activities. •Participation in physical activities (eg, dance class) helped with social interaction. •DBS had no effect on social and leisure role.

(Continues)

TABLE 1 (Continued)

Social Role	Subtheme	Summary of Results
	Clinical/PD symptoms	<ul style="list-style-type: none"> •Hoehn and Yahr stage, fatigue, mobility, and lack of symptom control were related to leisure role functioning. •Some people with PD are embarrassed by symptoms that makes them pull away from activities.

Abbreviations: DBS, deep brain stimulation; PD, Parkinson's disease; QoL, quality of life.

FIGURE 2 Distribution of articles based on social role examined and analysis performed

frequently reported no change in social functioning following surgery or in some cases a detrimental change.²⁴⁻²⁶ It is unclear why social functioning inconsistently improves post-DBS; however, Schüpbach et al revealed that expectations of surgery outcomes influenced relationships (couple and work) after surgery.²⁷ A psychoeducation class for DBS participants was found to significantly improve postsurgical social adjustment.²⁸ The Stanford Chronic Disease Self-Management Program, which aimed to improve social functioning, did not significantly improve social support scores; however, there were some positive correlations between changes in social support and changes in self-management outcomes after program participation.²⁹

Another class of PD therapy was behavioral interventions, including a dance class and a self-management training program. These interventions had a positive effect on social functioning, by increasing social engagement directly through the activity.²⁹⁻³⁴ Papers describing QoL or life changes after PD found that cognitive functioning related to participation.^{35,36} Additionally, Farhadi et al found that females reported worse psychosocial functioning and social support.³⁷

3.3.2 | Couple relationship

The couple relationship has been measured by concordance with partner, dependence on partner, feelings about partner, relationship satisfaction, and sexual adjustment (ie, frequency and enjoyment of intercourse).³⁸ Articles discussing this social role focused on the effect

of clinical/PD symptoms ($n = 7$), relationship satisfaction (nonsexual) ($n = 6$), sexual satisfaction ($n = 5$), and the effect of treatment ($n = 5$). The quantitative papers that described clinical/PD symptoms affecting the couple relationship reported that facial masking and Hoehn and Yahr stage were correlated to an impaired couple relationship, while speech problems had no significant impact.^{20,39} Older couples and those who were able to better cope with the disease reported better relationships.⁴⁰ The qualitative studies revealed positive and negative impacts on the couple relationship. A positive change that people living with Parkinson's and their spouses reported was the affirmation of their commitment to each other after the diagnosis.^{41,42} Some negative changes reported included shifting relational roles (with more responsibility falling on the care partner), changes in sexual intimacy, engaging in fewer activities together, and financial burden.^{39,41-44} Ten papers described the impact of PD on nonsexual aspects (eg, communication, attention, and shared activities) of relationship satisfaction. Buhmann et al found that people living with Parkinson's, especially women, believed these nonsexual aspects of their relationship became more important after PD diagnosis.⁴⁵ Eleanor Singer compared marriage satisfaction between people living with Parkinson's and age-matched controls and found no significant difference.⁴⁶ Three studies found that depression, anxiety, negative social exchanges, and alexithymia were associated with reduced relationship satisfaction.⁴⁷⁻⁴⁹ Mavandadi et al measured relationship satisfaction from the care partner perspective and found that satisfaction was

related to the care partner's "benefit finding," or ability to experience positive change when faced with a stressor like PD.⁴⁹ Karlstedt et al investigated relationship mutuality, "the positive quality of a relationship," and found that having a male care partner was associated with higher mutuality score for the people living with Parkinson's. Care partner mutuality score was associated with the person living with Parkinson's mutuality score and cognitive ability.⁵⁰ Qualitative papers in this subtheme reported that uncertainty about the future and role changes that placed a greater burden on the care partner were strongly associated with relationship quality.^{39,41,51}

Five papers described sexual functioning in PD and how it affects the couple relationship. Lower sexual satisfaction was more common among people with a younger disease onset, males, and people with worse motor scores (Movement Disorders Society Unified Parkinson's disease Rating Scale (MDS-UPDRS)), fatigue, and rigidity.^{52,53} Papers also reported that sex life satisfaction was significantly associated with marital satisfaction.^{44,47} Fleming et al elucidated this issue, listing dramatic increases or decreases in libido, as well as a shift in relationship roles from partner to carer as the main reasons for sex life dissatisfaction.⁴¹ Five papers described how different interventions or therapies impacted the couple relationship. DBS was found to diminish sexual desire and, in some cases, worsen marital satisfaction.^{26,46,54} Agid et al provided an explanation for this worsened marital quality following surgery: either people living with Parkinson's rejected their spouse after they felt "cured" or they were rejected by their spouse who expected them to be able to return to their premorbid level of functioning following surgery.²⁶ Similarly, higher doses of levodopa were associated with more frequent thoughts about breaking up with a partner and with relationship termination.⁴⁵ However, support group attendance had a positive influence on the relationship by providing a way for couples to have a shared social activity.⁵⁵

3.3.3 | Family role

The family role has been defined by feelings about family interactions, the ability to handle family financial needs, the frequency and quality of interactions with family members, interest in a child's activities and quality of interactions with children.³⁸ Two mixed-methods papers described how treatments/interventions influenced the family relationship.^{27,31} People living with Parkinson's who participated in a tango intervention reported improved family role functioning after the classes.³¹ Schüpbach et al reported improved family relationships after DBS surgery were more common than strained relationships.²⁷ Another common theme related to the family role was discussing or sharing the disease with family members. Five papers described that communication early in the disease was crucial for facilitating understanding of the disease among family members and reducing its burden.^{40,41,53,56,57} Fleming et al provided more information about why communication was a struggle for some people living with Parkinson's. People living with Parkinson's reported the need to "protect" their families and did not want their children to "miss out" on anything because of the diagnosis.⁴¹ Navarta-Sanchez et al described how health-care providers and family members influenced the way people

living with Parkinson's handled their disease.⁴⁰ Receiving support from their family helped make people living with Parkinson's feel more secure and motivated to maintain their treatment regimen.⁴⁰

Four papers described family relationship satisfaction in PD. Eleanor Singer compared people living with Parkinson's to age-matched controls and found no difference in parental role satisfaction.²⁴ There were three qualitative papers that discussed the importance of family relationships for QoL in people living with Parkinson's.^{23,42,58} Additionally, for some people living with Parkinson's who were unable to work, parenting or family relationships became a higher priority.

3.3.4 | Friendship role

The friendship role has been measured by how frequently contact (eg, telephone, email, and in person) is initiated with friends and the quality of these interactions.³⁸ Most people living with Parkinson's reported the number of social contacts they had remained stable after diagnosis and following some treatments for PD (eg, DBS surgery).^{4,54,59,60} Rubenstein et al noted that although the number of visits friends made to people living with Parkinson's remained the same, people living with Parkinson's were less likely to initiate social outings with friends.⁵⁹ Soleimani et al revealed that people living with Parkinson's were concerned about losing social connectedness because of their disease (preventing them from leaving the house or increasing their desire to remain isolated to conceal symptoms).⁶⁰ Reduced social outings were found to greatly influence QoL and functioning in other relationships (eg, couple and family).^{23,61} Fleming et al reported a divide in how friendships changed with PD. Some friendships were strengthened while others ended.⁴¹ There was no consensus on factors that predicted relationship outcome following diagnosis.

3.3.5 | Work role

The work role has previously been measured by incorporating both paid and unpaid work (eg, house work). Work role is defined by the duration of work, any changes to employment (eg, full time to part time), respondent's feelings about the quality of their work, and relationships with coworkers.³⁸ Nineteen papers described the work role and how it was impaired by PD. Three quantitative papers found factors that contributed to leaving the workforce included anxiety, older age, longer disease duration, female sex, cognitive performance, depression, and ability to perform ADL (eg, dressing and hygiene).^{36,62,63} Several papers described that people living with Parkinson's decreased their work outside of the home and at home. Decreased employment was more evident in the young-onset group, with reasons for leaving the workforce often tied to the inability to meet job demands.^{39,42,53,54,59,64,65} Barbara Habermann reported that people living with Parkinson's who remained in the workforce described goal adjustment, eg, changing their focus from career advancement to maintaining their current position.³⁹ Qualitative studies also revealed that leaving the workforce impacted other social roles and overall QoL due to perceived loss of societal contribution and social contacts from work.^{23,51,60} Two papers described the

impact of DBS and levodopa on work roles. There was no evidence that levodopa led people to rejoin the workforce.²⁴ Professional activity following DBS was more often worsened than improved.²⁷

3.3.6 | Social and leisure role

The social and leisure role has been measured by the frequency, duration, and quality of social activities (eg, hobbies and membership in organizations).³⁸ Four papers described that the types of activities in which people living with Parkinson's engaged tended to be more solitary and sedentary, such as reading or watching TV.^{4,24,41,66} Two qualitative papers provided reasons for this shift to more sedentary activities, including people living with Parkinson's giving up more physically demanding hobbies because of the disease or favoring more solitary activities because of the unpredictability of symptoms and embarrassment about symptoms.^{41,66} People living with Parkinson's mentioned that planning ahead was crucial for maintaining social activities and navigating symptom demands.⁶⁶ Six papers described the impact of treatment/interventions on social activities. Social and leisure role performance was improved for people living with Parkinson's who participated in activities with other people living with Parkinson's (eg, tango class).⁶⁷⁻⁶⁹ These classes naturally provided an opportunity for socialization, as well as the ability to meet people with similar challenges. Two papers evaluated the effect of DBS on social and leisure roles. Boel et al described no change in membership to organizations following DBS.⁵⁴ Liddle et al found that people living with Parkinson's reported improved leisure performance after DBS.⁷⁰

4 | DISCUSSION

4.1 | Summary of findings

The aim of this review was to describe the effect of PD on social role functioning. Our work integrates research on how PD affects various social roles, including the couple relationship, family, friendship, work, and social/leisure roles. Our analysis uncovered three central findings: (a) PD can affect performance in different social roles or may cause withdrawal from these roles; (b) standard pharmacologic and surgical interventions have little positive effect on social role functioning in PD; and (c) a wide variety of instruments and data collection methods were used in the reviewed studies, demonstrating a pressing need for a more uniform method to evaluate social role functioning in PD.

4.2 | Strengths and limitations

Our systematic review is the first to summarize social functioning in PD, using a mixed-methods approach. We applied the Cochrane robust methodological procedures during the review process and had more than one reviewer at each stage to reduce bias. We also acknowledge that we could have missed some potentially relevant articles in other databases or the grey literature. However, we do feel

that the five databases we have selected provide relevant sources for papers that would be included in our review.

Furthermore, our review only addressed social functioning from the perspective of the person with Parkinson disease. However, the patient's functioning is also dependent on how partners or caregivers cope with the disease of their loved one and the extent to which this influences their own lives. Some studies focusing on this issue have been published, but we considered caregiver outcomes to be beyond the scope of this review.^{71,72}

4.3 | Implications for research, policy, and practice

Social role performance is a priority for people with chronic diseases and the clinicians who treat them. Hammarlund et al researched which outcome measures were most important in PD trials from the perspectives of health-care professionals and people living with Parkinson's.⁶⁷ Study participants (people living with Parkinson's and health-care professionals) most frequently ranked aspects related to social involvement (ie, QoL, control of disease processes, ability to be on visiting terms, socializing, and participating in society) as the most important outcome. Additionally, the World Health Organization (WHO) emphasizes participation in social roles and social role performance among people with chronic disease or disability as a crucial method in which to prevent physical and mental health problems.¹⁶

Our review demonstrates that social functioning is an expanding area of research in clinical practice. Although some papers were published in more academic research-oriented journals ($n = 20$), there were several papers in neurology journals ($n = 8$), nursing journals ($n = 7$), rehabilitation journals ($n = 12$), and occupational therapy journals ($n = 4$). Additionally, publications related to social functioning have been increasing in popularity, with most of the papers included in this review published in the last 4 years.

Our review identified several correlates of impaired social role performance in PD, including disease severity, anxiety, depression, and cognitive impairment. However, less attention has been paid to interventions or methods of preserving or improving social role functioning. In fact, standard therapies for the motor symptoms of PD (DBS and levodopa) have not been shown to reliably improve social functioning and in some cases even worsen social role performance. Therefore, other approaches, such as nonpharmacologic therapies, should be investigated. A few studies found that activities like dancing or support groups improved social role functioning and QoL.^{31,32,39,73} Future research should focus on a broader range of interventions to improve social role functioning in PD.

This review provides several targets related to social role functioning, which could be used to develop interventions. Poor general social functioning was associated with facial masking, communication problems, and cognitive impairment.^{20,21,30,35,74-78} Impaired couple relationships were related to facial masking, higher Hoehn and Yahr stage, lack of coping responses (ie, spouse "benefit finding"), and sexual dysfunction.^{20,40,79} Earlier departure from the workforce was associated with female sex, older age, longer disease duration,

anxiety, ADL performance, depression, and cognitive impairment.^{33,42,53,54,59,64,65} The breadth of factors that contribute to impaired functioning signals a variety of plausible intervention targets. Although some factors are fixed, such as sex, age of disease onset, disease duration, age, and disease severity, there are many that can be targeted with treatment, including neuropsychiatric symptoms, sexual dysfunction, and motor function. Additionally, fixed factors can serve as markers to identify people living with Parkinson's at greater risk for problems in social role functioning. Certain current interventions and treatments address some of the modifiable factors associated with impaired social functioning. A variety of medications, therapies, and complementary strategies exist to address the neuropsychiatric and motor symptoms in PD. However, research on how these approaches affect social functioning is limited. Additionally, treatments for modifiable symptoms, such as sexual dysfunction, have been less thoroughly explored despite the strong association with QoL in many patients with chronic diseases and the couple relationship in PD.⁸⁰ One review of managing sexual dysfunction in PD promoted several methods, including discussing sexual function in regular neurologist visits, managing medications with sexual side effects, sexual counseling, and timing medications to improve sexual function.⁸¹ Incorporating these methods into regular care could greatly improve the couple relationship and overall QoL for people living with Parkinson's and their spouses. Additionally, understanding how these treatments affect the couple relationship would be beneficial to people living with Parkinson's, care partners, and clinicians. Of note, within the couple relationship role, dysfunction is not only exclusively related to patient variables or patient coping but also depends on caregiver coping and support. This stresses the importance of involving caregivers in (nonpharmacological) supportive treatments.

Social support is another potential target for interventions to improve social functioning in PD. Previous studies have found that positive affect in PD is increased by the number of social contacts maintained.⁸² Our summary revealed that some friendships are maintained while others are terminated after people living with Parkinson's received their diagnosis.⁴¹ However, the friendship role did not have specific factors associated with changes. Future research could further explore if clinical factors can predict changes in these roles. Beyond friendships, this review described activities that provided socialization, resulting in an increase in social support for people living with Parkinson's (eg, support groups and tango classes). This approach to socialization has been supported in other diseases as well, such as dementia and serious mental illness.^{83,84} Several other exercise classes have been researched in PD, including Tai Chi, boxing, aerobic exercise, and Qigong.^{4,61,85,86} Exercise classes have been shown to improve motor symptoms and QoL; however, direct impact on social functioning requires further investigation. Despite the improvement in health status that social support can provide, many patients view social support utilization as an indicator of functional decline and described actively avoiding the need for social support by adopting behaviors to maintain their independence.²⁹ It is important to consider stigma as a potential barrier to participating in interventions to increase social support.

In addition to informing directions for future interventions to improve social role functioning, this review also revealed the lack of PD-specific social functioning measures available to researchers. The quantitative studies included 24 different questionnaires or activities to assess aspects of social role functioning; however, only three questionnaires were used in more than one study: the 39-Item Parkinson's Disease Questionnaire ($n = 3$), Social Adjustment Scale ($n = 2$), and Golombok Rust Inventory of Marital State ($n = 2$). The 39-Item Parkinson's Disease Questionnaire is a PD specific questionnaire; however, it does not assess social functioning exclusively and includes only three questions that address social function. The Social Adjustment Scale focuses on social roles; however, it is not validated in people living with Parkinson's.³⁸ Finally, the Golombok Rust Inventory only measures the couple relationship and has not been validated in PD.⁸⁷ There are some factors that could be specific to social functioning in PD (eg, embarrassment caused by symptoms leading to isolation) that may not be adequately assessed in general questionnaires. A benefit of conducting this mixed-methods review was that the qualitative studies addressed some questions that arose from the quantitative assessments. For example, several studies of DBS revealed that social functioning in the couple relationship was often worsened following surgery; however, family role functioning was improved. This discrepancy could not be clarified by these questionnaires alone, but qualitative interviews uncovered that expectations from patients and spouses play a role in relationship satisfaction following surgery. In order to better measure social role functioning in PD, it would be important to develop assessments that are validated in this population and can capture the nuances of the disease that are not currently ascertained from existing questionnaires. Furthermore, new studies should consider incorporating a mixed methods approach to better understand individual experiences with PD.

5 | CONCLUSION

Successful aging involves engagement in social and productive activities.⁸⁸ Furthermore, reduced social participation is a risk factor for depression, cognitive decline, increased health-care costs, and overall mortality.⁶⁻⁹ Our review reveals how PD impairs general social functioning and the ability to fulfill specific roles. A number of symptoms associated with reduced social functioning can be targeted in order to improve function and QoL. However, current PD treatments and interventions have not been shown to adequately improve social functioning. Patients' social participation should be considered as soon as minor losses or changes are detected to prevent isolation and promote successful aging.

CONFLICT OF INTEREST

None declared.

ORCID

Kate Perepezko <https://orcid.org/0000-0003-3762-563X>

Jared T. Hinkle <https://orcid.org/0000-0001-6770-3067>

Martinus P.G. Broen <https://orcid.org/0000-0002-8834-2202>

REFERENCES

- Jankovic J. Parkinson's disease: clinical features and diagnosis. *J Neurol Neurosurg Psychiatry*. 2008;79(4):368-376.
- Schrag A, Jahanshahi M, Quinn N. How does Parkinson's disease affect quality of life? A comparison with quality of life in the general population. *Mov Disord*. 2000;15(6):1112-1118.
- Cohen F, Lazarus RS. Coping with the stresses of illness. *Heal Psychol A Handb*. 1979;1979:217-254.
- Singer E. Social costs of Parkinson's disease. *J Chronic Dis*. 1973;26(4):243-254.
- Bettencourt B, Sheldon K. Social roles as mechanism for psychological need satisfaction within social groups. *J Pers Soc Psychol*. 2001;81(6):1131.
- Holt-Lunstad J, Smith TB, Baker M, Harris T, Stephenson D. Loneliness and social isolation as risk factors for mortality. *Perspect Psychol Sci*. 2015;10(2):227-237. <https://doi.org/10.1177/1745691614568352>
- Shaw JG, Farid M, Noel-Miller C, et al. Social isolation and medicare spending: among older adults, objective isolation increases expenditures while loneliness does not. *J Aging Health*. 2017;29(7):1119-1143. <https://doi.org/10.1177/0898264317703559>
- Cornwell EY, Waite LJ. Social disconnectedness, perceived isolation, and health among older adults. *J Health Soc Behav*. 2009;50(1):31-48. <https://doi.org/10.1177/002214650905000103>
- Cacioppo J, Hawkey L, Twisted R. Perceived social isolation makes me sad. *Psychol Aging*. 2010;25(2):453-463.
- Wood-Dauphinee S. Assessing quality of life in clinical research: from where have we come and where are we going? *J Clin Epidemiol*. 1999;52(4):355-363.
- Deuschl G, Schade-Brittinger C, Krack P, et al. A randomized trial of deep-brain stimulation for Parkinson's disease. *N Engl J Med*. 2006;355(9):896-908.
- Den Ouden BL, Van Heck GL, De Vries J. Quality of life and related concepts in Parkinson's disease: a systematic review. *Mov Disord*. 2007;22(11):1528-1537. <https://doi.org/10.1002/mds.21567>
- Dowding CH, Shenton CL, Salek SS. A review of the health-related quality of life and economic impact of Parkinson's disease. *Drugs Aging*. 2006;23(9):693-721. <https://doi.org/10.2165/00002512-200623090-00001>
- Post B, Merkus MP, De Haan RJ, Speelman JD. Prognostic factors for the progression of Parkinson's disease: a systematic review. *Mov Disord*. 2007;22(13):1839-1851. <https://doi.org/10.1002/mds.21537>
- Soh S-E, Morris ME, McGinley JL. Determinants of health-related quality of life in Parkinson's disease: a systematic review. *Parkinsonism Relat Disord*. 2011;17(1):1-9.
- World Health Organization (WHO). (2016). *International classification of functioning, disability and health (ICF)*. Geneva, Switzerland: WHO; 2001.
- van Uem JMT, Marinus J, Canning C, et al. Health-related quality of life in patients with Parkinson's disease—a systematic review based on the ICF model. *Neurosci Biobehav Rev*. 2016;61:26-34.
- Critical Appraisal Skills Programme. 2018.
- Fox S, Cashell A, Kernohan WG, et al. Palliative care for Parkinson's disease: patient and carer's perspectives explored through qualitative interview. *Palliat Med*. 2017;31(7):634-641. <https://doi.org/10.1177/0269216316669922>
- Gunnery SD, Habermann B, Saint-Hilaire M, Thomas CA, Tickle-Degnen L. The relationship between the experience of hypomimia and social wellbeing in people with Parkinson's disease and their care partner. *S J Park Dis*. 2016;6(3):625-630.
- Miller N, Noble E, Jones D, Burn D. Life with communication changes in Parkinson's disease. *Age Ageing*. 2006;35(3):235-239. <https://doi.org/10.1093/ageing/afj053>
- Wressle E, Engstrand C, Granérus A. Living with Parkinson's disease: elderly patients' and relatives' perspective on daily living. *Aust Occup Ther J*. 2007;54(2):131-139. <http://search.ebscohost.com/login.aspx?direct=true&db=rzh&AN=105743000&site=ehost-live&scope=siteNS->
- Dauwerse L, Hendriks A, Schipper K, Struiksma C, Abma TA. Quality-of-life of patients with Parkinson's disease. *Brain Inj*. 2014;28(10):1342-1352. <https://doi.org/10.3109/02699052.2014.916417>
- Singer E. The effect of treatment with levodopa on Parkinson patients' social functioning and outlook on life. *J Clin Epidemiol*. 1974;27(11):581-594.
- Volkman J, Albanese A, Kulisevsky J, et al. Long-term effects of pallidal or subthalamic deep brain stimulation on quality of life in Parkinson's disease. *Mov Disord*. 2009;24(8):1154-1161.
- Agid Y, Schüpbach M, Gargiulo M, et al. Neurosurgery in Parkinson's disease: the doctor is happy, the patient less so? *J Neural Transm*. 2006. https://doi.org/10.1007/978-3-211-45295-0_61
- Schüpbach M, Gargiulo M, Welter ML, et al. Neurosurgery in Parkinson disease: A distressed mind in a repaired body? *Neurol*. 2006;66(12):1811-1816. <https://doi.org/10.1212/01.wnl.0000234880.51322.16>
- Dos Santos JFA, Du Montcel ST, Gargiulo M, et al. Tackling psychosocial maladjustment in Parkinson's disease patients following subthalamic deep-brain stimulation: a randomised clinical trial. *PLoS ONE*. 2017;12(4):e0174512. <https://doi.org/10.1371/journal.pone.0174512>
- Pappa K, Doty T, Taff SD, Knipmann K, Foster ER. Self-management program participation and social support in Parkinson's disease: mixed methods evaluation. *Phys Occup Ther Geriatr*. 2017;35(2):81-98. <https://doi.org/10.1080/02703181.2017.1288673>
- Fox S, Cashell A, Kernohan WG, et al. Palliative care for Parkinson's disease: patient and carer's perspectives explored through qualitative interview. *Palliat Med*. 2017;31(7):634-641. <https://doi.org/10.1177/0269216316669922>
- Zafar M, Bozzorg A, Hackney ME. Adapted tango improves aspects of participation in older adults versus individuals with Parkinson's disease. *Disabil Rehabil*. 2017;39(22):2294-2301. <https://doi.org/10.1080/09638288.2016.1226405>
- Rocha PA, Slade SC, McClelland J, Morris ME. Dance is more than therapy: qualitative analysis on therapeutic dancing classes for Parkinson's. *Complement Ther Med*. 2017;34:1-9.
- Habermann B, Shin JY. Preferences and concerns for care needs in advanced Parkinson's disease: a qualitative study of couples. *J Clin Nurs*. 2017;26(11-12):1650-1656.
- Choudhury TK, Harris C, Crist K, Satterwhite TK, York MK. Comparative patient satisfaction and feasibility of a pilot Parkinson's disease enrichment program. *J Geriatr Psychiatry Neurol*. 2017;30(5):253-260. <https://doi.org/10.1177/0891988717720299>
- Vlaspolder TT, Koerts J, Tucha O, et al. Objective versus subjective measures of executive functions: predictors of participation and quality of life in Parkinson disease? *Arch Phys Med Rehabil*. 2017;98(11):2181-2187. <https://doi.org/10.1016/j.apmr.2017.03.016>
- Meligrana L, Sgaramella Maria T, Bartolomei L, et al. The role of cognitive, functional and neuropsychiatric symptoms on community integration in Parkinson's disease. *Int J Child Heal Hum Dev*. 2011.

37. Farhadi F, Vosoughi K, Shahidi GA, Delbari A, Lökk J, Fereshtehnejad SM. Sexual dimorphism in Parkinson's disease: differences in clinical manifestations, quality of life and psychosocial functioning between males and females. *Neuropsychiatr Dis Treat*. 2017. <https://doi.org/10.2147/NDT.S124984>
38. Weissman M, Bothwell S. Self-report version of the Social Adjustment Scale. *Arch Gen Psychiatry*. 1976;33(9):1111-1115.
39. Habermann B. Day-to-day demands of Parkinson's disease. *West J Nurs Res*. 1996;18(4):397-413. <https://doi.org/10.1177/019394599601800403>
40. Navarta-Sánchez MV, Senosiain García JM, Riverol M, et al. Factors influencing psychosocial adjustment and quality of life in Parkinson patients and informal caregivers. *Qual Life Res An Int J Qual Life Asp Treat Care Rehabil*. 2016;(Journal Article PG-. <https://doi.org/10.1007/s11136-015-1220-3>
41. Fleming V, Tolson D, Schartau E. Changing perceptions of womanhood: living with Parkinson's disease. *Int J Nurs Stud*. 2004;41(5):515-524. <https://doi.org/10.1016/j.ijnurstu.2003.12.004>
42. Hodgson JH, Garcia K, Tyndall L. Parkinson's disease and the couple relationship: a qualitative analysis. *Fam Syst Health*. 2004;22(1):101-118. <https://doi.org/10.1037/1091-7527.22.1.101>
43. Martin SC. Relational issues within couples coping with Parkinson's disease: implications and ideas for family-focused care. *J Fam Nurs*. 2016;22(2):224-251.
44. O'Connor EJ, McCabe MP, Firth L. The impact of neurological illness on marital relationships. *J Sex Marital Ther*. 2008;(2):115-132. <https://doi.org/10.1080/00926230701636189>
45. Buhmann C, Dogac S, Vettorazzi E, Hidding U, Gerloff C, Jürgens TP. The impact of Parkinson disease on patients' sexuality and relationship. *J Neural Transm*. 2017;124(8):983-996. <https://doi.org/10.1007/s00702-016-1649-8>
46. Singer E. Premature social aging: the social-psychological consequences of a chronic illness. *Soc Sci Med*. 1974;8(3):143-151. [https://doi.org/10.1016/0037-7856\(74\)90020-1](https://doi.org/10.1016/0037-7856(74)90020-1)
47. Wielinski CL, Varpness SC, Erickson-Davis C, Paraschos AJ, Parashos SA. Sexual and relationship satisfaction among persons with young-onset Parkinson's disease. *J Sex Med*. 2010;7(4):1438-1444. <https://doi.org/10.1111/j.1743-6109.2009.01408.x>
48. Ricciardi L, Pomponi M, Demartini B, et al. Emotional awareness, relationship quality, and satisfaction in patients with Parkinson's disease and their spousal caregivers. *J Nerv Ment Dis*. 2015;203(8):646-649.
49. Mavandadi S, Dobkin R, Mamikonyan E, Sayers S, Ten Have T, Weintraub D. Benefit finding and relationship quality in Parkinson's disease: a pilot dyadic analysis of husbands and wives. *J Fam Psychol*. 2014;28(5):728-734. <https://doi.org/10.1037/a0037847>
50. Karlstedt M, Fereshtehnejad S-M, Aarsland D, Lökk J. Determinants of dyadic relationship and its psychosocial impact in patients with Parkinson's disease and their spouses. *Parkinsons Dis*. 2017;2017:1-9. <https://doi.org/10.1155/2017/4697052>
51. Ravenek M, Rudman DL, Jenkins ME, Spaulding S. Understanding uncertainty in young-onset Parkinson disease. *Chronic Illn*. 2017;13(4):288-298. <https://doi.org/10.1177/1742395317694699>
52. Bronner G, Cohen OS, Yahalom G, et al. Correlates of quality of sexual life in male and female patients with Parkinson disease and their partners. *Parkinsonism Relat Disord*. 2014;20(10):1085-1088. <https://doi.org/10.1016/j.parkreldis.2014.07.003>
53. Schrag A, Hovris A, Morley D, Quinn N, Jahanshahi M. Young- versus older-onset Parkinson's disease: impact of disease and psychosocial consequences. *Mov Disord*. 2003;18(11):1250-1256. <https://doi.org/10.1002/mds.10527>
54. Boel JA, Odekerken VJJ, Geurtsen GJ, et al. Psychiatric and social outcome after deep brain stimulation for advanced Parkinson's disease. *Mov Disord*. 2016;31(3):409-413. <https://doi.org/10.1002/mds.26468>
55. Smith LJ, Shaw RL. Learning to live with Parkinson's disease in the family unit: an interpretative phenomenological analysis of well-being. *Med Health Care Philos*. 2017;20(1):13-21. <https://doi.org/10.1007/s11019-016-9716-3>
56. Whetten-Goldstein K, Sloan F, Kulas E, Cutson T, Schenkman M. The burden of Parkinson's disease on society, family, and the individual. *J Am Geriatr Soc*. 1997;45(7):844-849. <https://doi.org/10.1111/j.1532-5415.1997.tb01512.x>
57. Kang MY, Ellis-Hill C. How do people live life successfully with Parkinson's disease? *J Clin Nurs*. 2015;24(15-16):2314-2322. <https://doi.org/10.1111/jocn.12819>
58. Den Oudsten BL, Lucas-Carrasco R, Green AM, Whoqol-Dis Group T. Perceptions of persons with Parkinson's disease, family and professionals on quality of life: an international focus group study. *Disabil Rehabil*. 2011;33(25/26):2490-2508. <https://doi.org/10.3109/09638288.2011.575527>
59. Rubenstein LM, Chrischilles EA, Voelker MD. The impact of Parkinson's disease on health status, health expenditures, and productivity. *Pharmacoeconomics*. 1997;12(4):486-498.
60. Soleimani MA, Bastani F, Negarandeh R, Greysen R. Perceptions of people living with Parkinson's disease: a qualitative study in Iran. *Br J Community Nurs*. 2016;21(4):188-195. <https://doi.org/10.12968/bjcn.2016.21.4.188>
61. Lee KS, Merriman A, Owen A, Chew B, Tan TC. The medical, social, and functional profile of Parkinson's disease patients. *Singapore Med J*. 1994.
62. Timpka J, Svensson J, Nilsson MH, Pålhagen S, Hagell P, Odin P. Workforce unavailability in Parkinson's disease. *Acta Neurol Scand*. 2017;135(3):332-338.
63. Ray J, Das SK, Gangopadhyaya PK, Roy T. Quality of life in Parkinson's disease-Indian scenario. *Japi*. 2006;54:17-21.
64. Wong SL, Gilmour H, Ramage-Morin PL. Parkinson's disease: prevalence, diagnosis and impact. *Heal Rep*. 2014;25(11):10.
65. Valcarengi RV. The daily lives of people with Parkinson's disease. *Rev Bras Enferm*. 2018;71(2):272-279. <https://doi.org/10.1590/0034-7167-2016-0577>
66. Thordardottir B, Nilsson MH, Iwarsson S, Haak M. "You plan, but you never know"—participation among people with different levels of severity of Parkinson's disease. *Disabil Rehabil*. 2014;36(26):2216-2224. <https://doi.org/10.3109/09638288.2014.898807>
67. Hammarlund CS, Nilsson MH, Hagell P. Measuring outcomes in Parkinson's disease: a multi-perspective concept mapping study. *Qual Life Res*. 2012;21(3):453-463.
68. Sabari JS, Ortiz D, Pallatto K, Yagerman J, Glazman S, Bodis-Wollner I. Activity engagement and health quality of life in people with Parkinson's disease. *Disabil Rehabil*. 2015;37(16):1411-1415.
69. Bogner S, DeFaria AM, O'Dwyer C, et al. More than just dancing: experiences of people with Parkinson's disease in a therapeutic dance program. *Disabil Rehabil*. 2016;39(11):1-6. <https://doi.org/10.1080/09638288.2016.1175037>
70. Liddle J, Phillips J, Gustafsson L, Silburn P. Understanding the lived experiences of Parkinson's disease and deep brain stimulation (DBS) through occupational changes. *Aust Occup Ther J*. 2018;65(1):45-53. <https://doi.org/10.1111/1440-1630.12437>

71. Tessitore A, Marano P, Modugno N, et al. Caregiver burden and its related factors in advanced Parkinson's disease: data from the PRE-DICT study. *J Neurol*. 2018;1-14.
72. Berger S, Chen T, Eldridge J, Thomas CA, Habermann B, Tickle-Degnen L. The self-management balancing act of spousal care partners in the case of Parkinson's disease. *Disabil Rehabil*. 2017;41(8):1-9. <https://doi.org/10.1080/09638288.2017.1413427>
73. Vann-Ward T, Morse JM, Charmaz K. Preserving self: theorizing the social and psychological processes of living with Parkinson disease. *Qual Health Res*. 2017;27(7):964-982.
74. Kalf JG, Smit AM, Bloem BR, Zwarts MJ, Munneke M. Impact of drooling in Parkinson's disease. *J Neurol*. 2007;254(9):1227-1232. <https://doi.org/10.1007/s00415-007-0508-9>
75. Anderson RJ, Simpson AC, Channon S, Samuel M, Brown RG. Social problem solving, social cognition, and mild cognitive impairment in Parkinson's disease. *Behav Neurosci*. 2013;127(2):184-192.
76. Shin JY, Pohlig RT, Habermann B. Self-reported symptoms of Parkinson's disease by sex and disease duration. *West J Nurs Res*. 2016;39(11):1412-1428. <https://doi.org/10.1177/0193945916670904>
77. Marr JA. The experience of living with Parkinson's disease. *J Neurosci Nurs*. 1991;23(5):325-329.
78. Shaw ST, Vivekananda-Schmidt P. Challenges to ethically managing Parkinson disease. *J Patient Exp*. 2017;4(4):191-196. <https://doi.org/10.1177/2374373517706836>
79. Greene SM, Griffin WA. Symptom study in context: effects of marital quality on signs of Parkinson's disease during patient-spouse interaction. *Psychiatry*. 1998;61(1):35-45. <https://doi.org/10.1080/00332747.1998.11024817>
80. Clayton A, Ramamurthy S. The impact of physical illness on sexual dysfunction. *Sexual Dysfunction*. 2008. <https://doi.org/10.1159/000126625>
81. Meco G, Rubino A, Caravona N, Valente M. Sexual dysfunction in Parkinson's disease. *Parkinsonism Relat Disord*. 2008;14(6):451-456.
82. Simpson J, Haines K, Lekwuwa G, Wardle J, Crawford T. Social support and psychological outcome in people with Parkinson's disease: evidence for a specific pattern of associations. *Br J Clin Psychol*. 2006;45(4):585-590. <https://doi.org/10.1348/014466506X96490>
83. Palo-Bengtsson L, Ekman S-L. Social dancing in the care of persons with dementia in a nursing home setting: a phenomenological study. *Sch Inq Nurs Pract*. 1997;11(2):23.
84. Hackney ME, Earhart GM. Social partnered dance for people with serious and persistent mental illness: a pilot study. *J Nerv Ment Dis*. 2010;198(1):76-78.
85. Bridgewater KJ, Sharpe MH. Trunk muscle training and early Parkinson's disease. *Physiother Theory Pract*. 1997;13(2):139-153.
86. Schmitz-Hübsch T, Pyfer D, Kielwein K, Fimmers R, Klockgether T, Wüllner U. Qigong exercise for the symptoms of Parkinson's disease: a randomized, controlled pilot study. *Mov Disord*. 2006;21(4):543-548.
87. Rust J, Golombok S. The Golombok-Rust Inventory of Sexual Satisfaction (GRISS). *Br J Clin Psychol*. 1985;24(1):63-64. <https://doi.org/10.1111/j.2044-8260.1985.tb01314.x>
88. Rowe JW, Kahn RL. Successful aging. *Gerontologist*. 1997;44(1):155-169. <https://doi.org/10.5054/tq.2010.215250>

SUPPORTING INFORMATION

Additional supporting information may be found online in the Supporting Information section at the end of the article.

How to cite this article: Perepezko K, Hinkle JT, Shepard MD, et al. Social role functioning in Parkinson's disease: A mixed-methods systematic review. *Int J Geriatr Psychiatry*. 2019;34:1128-1138. <https://doi.org/10.1002/gps.5137>