Left ventricular lead positioning in cardiac resynchronization therapy

Citation for published version (APA):

Nguyen, U. C., Prinzen, F. W., & Vernoooy, K. (2017). Left ventricular lead positioning in cardiac resynchronization therapy: Mission accomplished? *Heart Rhythm*, 14(9), 1373-1374.
<https://doi.org/10.1016/j.hrthm.2017.05.030>

Document status and date:

Published: 01/09/2017

DOI:

[10.1016/j.hrthm.2017.05.030](https://doi.org/10.1016/j.hrthm.2017.05.030)

Document Version:

Publisher's PDF, also known as Version of record

Document license:

Taverne

Please check the document version of this publication:

- A submitted manuscript is the version of the article upon submission and before peer-review. There can be important differences between the submitted version and the official published version of record. People interested in the research are advised to contact the author for the final version of the publication, or visit the DOI to the publisher's website.
- The final author version and the galley proof are versions of the publication after peer review.
- The final published version features the final layout of the paper including the volume, issue and page numbers.

[Link to publication](#)

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal.

If the publication is distributed under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license above, please follow below link for the End User Agreement:

www.umlib.nl/taverne-license

Take down policy

If you believe that this document breaches copyright please contact us at:

repository@maastrichtuniversity.nl

providing details and we will investigate your claim.

Left ventricular lead positioning in cardiac resynchronization therapy: Mission accomplished?

Uyên Châu Nguyễn, MD, MSc,^{*†} Frits W. Prinzen, PhD,^{*} Kevin Vernooy, MD, PhD^{*†}

From the ^{*}Department of Physiology, Maastricht University Medical Center, Maastricht, The Netherlands, and [†]Department of Cardiology, Maastricht University Medical Center, Maastricht, The Netherlands.

Cardiac resynchronization therapy (CRT) is an effective treatment for patients with symptomatic heart failure, reduced systolic left ventricular (LV) function, and QRS delay.¹ Nevertheless, evidence continues to increase that the benefit of CRT can be improved by positioning the LV lead remote from scar in an area of late electrical or mechanical activation.^{2,3} Numerous studies have emerged attempting to tackle this issue using variable modalities to guide LV lead placement to a patient-specific site of late activation remote from scar.^{3–8} Delayed enhancement cardiac magnetic resonance imaging (DE-CMR) remains the criterion standard for myocardial scar assessment. Accordingly, this modality is predominantly used to guide LV lead placement outside scar in most of the aforementioned studies.

In this issue of *HeartRhythm*, Behar et al⁹ report on an evaluation of the use of computed tomography (CT) for selection of the optimal region for LV lead placement in a segment of late mechanical activation outside scar. This study was performed in 18 patients with existing pacing systems and therefore contraindications for CMR. Segmental mechanical activation was calculated as the time to peak strain, derived from the stretch of the endocardial surface throughout a heartbeat (stretch quantifier of endocardial engraved zones [SQUEEZE] using CT data sets together with electrical delay. Excluded as target segments were septal segments and segments with low-amplitude CT-SQUEEZE strain curves, the latter being regarded as nonviable. LV lead positioning was guided by acute hemodynamic response (AHR). After implantation, CT-SQUEEZE target segments were compared with AHR-target segments^{7,8} and 70% concordance in target segments between both methods was found. AHR during pacing was not significantly lower in the CT-SQUEEZE target segments in comparison with the best achievable AHR and similar to pacing segments with latest electrical delay (“Q-LV”). Conversely, LV pacing in scar

led to a considerable reduction in AHR. The principal finding was that CT-SQUEEZE may be a valuable tool to guide LV lead placement.

The authors are the first to implement an algorithm that computes mechanical activation from CT data sets for LV lead guidance in patients undergoing CRT and should therefore be congratulated for the novelty of their work. Moreover, they took the initiative to provide alternatives for DE-CMR scar assessment in patients with preexisting pacing systems, a study population that has been underexposed in previous studies about LV lead placement. Nonetheless, several limitations should be acknowledged, the majority of them already pointed out by the authors. First, CT assessment of scar was done by contrast enhancement ($n = 1$) or subjective judgment of wall thinning and hypoperfusion. However, in the end only low-amplitude strain was used to exclude segments as target. Second, CT-SQUEEZE circumferential strain has been validated against CMR tagging in an animal model of ischemia, but the reliability of associating low-amplitude strains from this metric as scar yet remains to be investigated.¹⁰ Finally, although the sample size is sufficient for a feasibility study, the generalization of concordance between the AHR and the CT-SQUEEZE target segment remains limited.

The hypothesis that pacing at a site of delayed activation leads to more resynchronization, and subsequently improved clinical outcome seems plausible from an electrophysiological point of view. Nevertheless, data supporting this notion are limited to single-center studies using different imaging modalities and algorithms for mechanical activation computation. In the TARGET study, the mortality rate was indeed lower when the LV lead was placed in the latest activated region defined by speckle-tracking echocardiography.³ However, even more mortality risk reduction was observed when the LV lead was placed remote from scar, suggesting that avoiding scar is more important than targeting latest activated regions. Another concern about the concept of positioning the LV lead in the segment with latest activation at baseline is that during right ventricular pacing, which is commonly used in CRT as part of biventricular pacing, the location of the latest activation differs from that of intrinsic activation in a large proportion of patients.^{11,12} In addition,

Dr Prinzen has received research grants from Medtronic, St. Jude Medical, Sorin, MSD, and Biotronik. Dr Vernooy has received research grants from Medtronic and St. Jude Medical. **Address reprint requests and correspondence:** Dr Kevin Vernooy, Department of Cardiology, Maastricht University Medical Center, P. Debyelaan 25, PO Box 5800, 6202 AZ Maastricht, The Netherlands. E-mail address: kevin.vernooy@mumc.nl.

computer simulations showed that the best electrical resynchronization and AHR were achieved when the LV lead was positioned mid-basolateral, as this site induced fast activation of the LV, regardless of underlying conduction disturbance.¹³

Taken collectively, avoiding scar appears to be the priority for optimal LV lead placement. Accurate scar assessment in this context is therefore relevant, and thus validating imaging modalities other than DE-CMR for this purpose is desired. An additional advantage of CT-based scar assessment over DE-CMR may be the ability to evaluate the anatomy of the coronary veins, as it appears that target segments based on scar anatomy or activation are irrelevant in the absence of a coronary vein in that region.

In conclusion, positioning the LV lead remote from scar is relevant for optimal CRT response and may even more important than targeting regions of late activation. Evaluating scar by CT seems a promising method for LV lead guidance in patients with contraindication for CMR and may be even more useful for CRT road mapping when coronary venous anatomy is assessed preprocedurally as well.

References

1. European Heart Rhythm Association, European Society of Cardiology, Heart Rhythm Society, Heart Failure Society of Association, American Society of Echocardiography, Heart Failure Association, Daubert JC, Saxon L, Adamson PB, et al. 2012 EHRA/HRS expert consensus statement on cardiac resynchronization therapy in heart failure: implant and follow-up recommendations and management. *Heart Rhythm* 2012;9:1524–1576.
2. Kandala J, Upadhyay GA, Altman RK, et al. QRS morphology, left ventricular lead location, and clinical outcome in patients receiving cardiac resynchronization therapy. *Eur Heart J* 2013;34:2252–2262.
3. Khan FZ, Virdee MS, Palmer CR, et al. Targeted left ventricular lead placement to guide cardiac resynchronization therapy: the TARGET study: a randomized, controlled trial. *J Am Coll Cardiol* 2012;59:1509–1518.
4. Nguyen UC, Mafi-Rad M, Aben JP, et al. A novel approach for left ventricular lead placement in cardiac resynchronization therapy: intraprocedural integration of coronary venous electroanatomic mapping with delayed enhancement cardiac magnetic resonance imaging. *Heart Rhythm* 2017;14:110–119.
5. Bakos Z, Markstad H, Ostenfeld E, Carlsson M, Roijer A, Borgquist R. Combined preoperative information using a bullseye plot from speckle tracking echocardiography, cardiac CT scan, and MRI scan: targeted left ventricular lead implantation in patients receiving cardiac resynchronization therapy. *Eur Heart J Cardiovasc Imaging* 2014;15:523–531.
6. Shetty AK, Duckett SG, Ginks MR, et al. Cardiac magnetic resonance-derived anatomy, scar, and dyssynchrony fused with fluoroscopy to guide LV lead placement in cardiac resynchronization therapy: a comparison with acute haemodynamic measures and echocardiographic reverse remodelling. *Eur Heart J Cardiovasc Imaging* 2013;14:692–699.
7. Rad MM, Blaauw Y, Dinh T, et al. Left ventricular lead placement in the latest activated region guided by coronary venous electroanatomic mapping. *Europace* 2015;17:84–93.
8. Laksman Z, Yee R, Stirrat J, et al. Model-based navigation of left and right ventricular leads to optimal targets for cardiac resynchronization therapy: a single-center feasibility study. *Circ Arrhythm Electrophysiol* 2014;7:1040–1047.
9. Behar JM, Rajani R, Pourmorteza A, et al. Comprehensive use of cardiac computed tomography to guide left ventricular lead placement in cardiac resynchronization therapy. *Heart Rhythm* 2017;14:1364–1372.
10. Pourmorteza A, Chen MY, van der Pals J, Arai AE, McVeigh ER. Correlation of CT-based regional cardiac function (SQUEEZ) with myocardial strain calculated from tagged MRI: an experimental study. *Int J Cardiovasc Imaging* 2016; 32:817–823.
11. Mafi Rad M, Blaauw Y, Dinh T, et al. Different regions of latest electrical activation during left bundle-branch block and right ventricular pacing in cardiac resynchronization therapy patients determined by coronary venous electroanatomic mapping. *Eur J Heart Fail* 2014;16:1214–1222.
12. Ludwig DR, Tanaka H, Friehling M, Gorcsan J, Schwartzman D. Further deterioration of LV ejection fraction and mechanical synchrony during RV apical pacing in patients with heart failure and LBBB. *J Cardiovasc Transl Res* 2013; 6:425–429.
13. Pluijmer M, Bovendeerd PH, Lumens J, Vernooij K, Prinzen FW, Delhaas T. New insights from a computational model on the relation between pacing site and CRT response. *Europace* 2016;18:iv94–iv103.