

Nonpain goal pursuit inhibits attentional bias to pain

Citation for published version (APA):

Schrooten, M. G. S., van Damme, S., Crombez, G., Peters, M. L., Vogt, J., & Vlaeyen, J. W. S. (2012). Nonpain goal pursuit inhibits attentional bias to pain. *Pain*, 153(6), 1180-1186. <https://doi.org/10.1016/j.pain.2012.01.025>

Document status and date:

Published: 01/06/2012

DOI:

[10.1016/j.pain.2012.01.025](https://doi.org/10.1016/j.pain.2012.01.025)

Document Version:

Publisher's PDF, also known as Version of record

Document license:

Taverne

Please check the document version of this publication:

- A submitted manuscript is the version of the article upon submission and before peer-review. There can be important differences between the submitted version and the official published version of record. People interested in the research are advised to contact the author for the final version of the publication, or visit the DOI to the publisher's website.
- The final author version and the galley proof are versions of the publication after peer review.
- The final published version features the final layout of the paper including the volume, issue and page numbers.

[Link to publication](#)

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal.

If the publication is distributed under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license above, please follow below link for the End User Agreement:

www.umlib.nl/taverne-license

Take down policy

If you believe that this document breaches copyright please contact us at:

repository@maastrichtuniversity.nl

providing details and we will investigate your claim.

Nonpain goal pursuit inhibits attentional bias to pain

Martien G.S. Schrooten^{a,b,*}, Stefaan Van Damme^c, Geert Crombez^c, Madelon L. Peters^a, Julia Vogt^c, Johan W.S. Vlaeyen^{a,b}

^a Department of Clinical Psychological Science, Maastricht University, Maastricht, The Netherlands

^b Department of Psychology, University of Leuven, Leuven, Belgium

^c Department of Experimental-Clinical and Health Psychology, Ghent University, Belgium

Sponsorships or competing interests that may be relevant to content are disclosed at the end of this article.

ARTICLE INFO

Article history:

Received 7 November 2011

Received in revised form 14 January 2012

Accepted 26 January 2012

Keywords:

Pain

Fear

Motivation

Goal pursuit

ABSTRACT

Although dealing with pain is a vital goal to pursue, most individuals are also engaged in the pursuit of other goals. The aim of the present experiment was to investigate whether attentional bias to pain signals is inhibited when one is pursuing a concurrent salient but nonpain task goal. Attentional bias to pain signals was measured in pain-free volunteers ($n = 63$) using a spatial cueing task with pain cues and neutral cues. The pursuit of a concurrent goal was manipulated by including additional trials in which a digit appeared at the middle of the screen. Half of the participants (goal group) were instructed to name these additional stimuli aloud. In order to increase the affective-motivational value of this non-pain-related goal, monetary reward and punishment were made contingent upon the performance of this task. Participants of the control group did not perform the additional task. As predicted, the results show attentional bias to pain signals in the control group, but not in the goal group. This indicates that attentional bias to signals of impending pain is inhibited when one is engaged in the pursuit of another salient but nonpain goal. The results of this study underscore a motivational view on attention to pain, in which the pursuit of multiple goals, including nonpain goals, is taken into account.

© 2012 International Association for the Study of Pain. Published by Elsevier B.V. All rights reserved.

1. Introduction

An influential idea is that attention to pain and somatic threat plays an important role in acute and chronic pain [7,11,28,47,52]. Consistent with this idea, evidence from behavioural and neurobiological studies suggests attentional bias favouring the processing of pain-related information in both healthy persons and chronic pain patients [35,47]. However, some seemingly inconsistent and inconclusive findings also have been reported regarding attentional biases to pain-related information, particularly for pain patients [35,47]. These inconsistencies might be resolved by assuming that, in contrast to what is often thought, attention to pain-related information is not a stable mechanism [47]. It has, for example, recently been argued that attention to (signals of impending) pain, as well as affective information processing in general, depends on the motivational context, or in other words, the pursuit of concurrent goals [31,47]. Indeed, pain and pain-related fear mostly occur within a context in which individuals also face pain-unrelated tasks and challenges.

* Corresponding author. Address: Faculty of Psychology and Neuroscience, Department of Clinical Psychological Science, Maastricht University, P.O. Box 616, Maastricht, MD 6200, The Netherlands. Tel.: +31 43 388 1484; fax: +31 43 388 4155.

E-mail address: martien.schrooten@maastrichtuniversity.nl (M.G.S. Schrooten).

When confronted with multiple goals, the pursuit of one goal often impairs the likelihood of achieving other goals [30]. Commitment to one goal might result in reduced accessibility of alternative goals [12,40,41] and attentional priority to information related to the focal goal [17,22,27,53,54]. Within such a goal-pursuit perspective, the extent to which pain captures attention is not merely dependent upon characteristics related to the painful stimulus, but also upon characteristics of the focal goal [11,47]. In that respect, one could expect attention to pain being attenuated by top-down processing induced by nonpain goals [21].

Supporting this reasoning, previous research has shown that engagement in a pain-unrelated task modulates the processing of pain and pain-related information. For instance, cognitively demanding distraction tasks have been associated with lower pain ratings and reduced pain-related brain activity [1,2,39,49]. Moreover, there are indications that working memory load during nociceptive stimulation [20] and motivational relevance of the task [49] modulate the disruptive effect of pain on task performance. In the present experiment, we advance this work by directly measuring the effects of a concurrent nonpain goal on attentional bias to pain-related information.

More specifically, the present experiment tests the hypothesis that biasing attention to pain signals, as previously observed [43,45,46,48], is inhibited when a concurrent nonpain goal is

pursued. Attentional bias was assessed with a spatial cueing task in otherwise pain-free participants [43,45,46,48]. Nonpain goal pursuit was manipulated independently of the cueing task: in half of the participants, the cueing task was combined with an independent goal task (cf. [32,33]). To enhance motivational salience of the goal task, participants were led to believe that they could win or lose money depending on their performance on this task. Because the goal task was presented concurrently (intermixed) with the cueing task, motivational context and nonpain goal pursuit remained salient during the cueing task and thus, during assessment of attentional bias for pain signals. We predicted that in participants who were motivated to engage in the goal task, attentional bias to pain signals would be smaller compared to participants who only performed the cueing task.

2. Method

2.1. Participants

Sixty-three undergraduates (11 men) with a mean age of 20.7 years (SD = 2.4) participated. Candidate participants were recruited through advertisements at Maastricht University. Exclusion criteria included pain complaints (acute/chronic), pregnancy, electronic implant, attention deficit disorder, and insufficient knowledge of the Dutch language. All participants reported themselves to be healthy, to have (corrected to) normal vision and colour vision, and were naïve as to the purpose of the experiment. They gave informed consent and received €10. The department's Ethics Committee approved the study.

2.2. Apparatus and task materials

2.2.1. Electrocutaneous stimuli

Electrocutaneous stimuli (bipolar sinus waveform; 300-ms duration) were administered by a direct current stimulator with 50-Hz internal frequency (Instrument Development Engineering & Evaluation, Maastricht University). Stimuli were applied to the external side of the left ankle using 2 8-mm stainless steel electrodes, vertically placed with 1-cm inter-electrode distance and filled with hypertonic gel. Stimulus intensity was individually determined using a work-up procedure. More specifically, a series of electrocutaneous stimuli with gradually increasing intensity was delivered. After each stimulus, participants rated its unpleasantness on an 11-point Likert scale (0 = "not unpleasant at all"; 10 = "very unpleasant") and indicated whether the stimulus was tolerable or not. Intensity was increased to a rating of 9. Then the series was delivered again. The stimulus that was perceived as very unpleasant (rating 9) and difficult to tolerate during the second series was presented during the computer task. Participants were not informed about these procedural details.

After the work-up procedure, in order to increase the threat value of the electrocutaneous stimuli, participants were led to believe that stimuli of a higher intensity than selected would be occasionally presented during the test phase [4]. Participants had no prior experience with such a high-intensity stimulus. In fact, no stimuli of higher intensity were delivered. The threat value of the electrocutaneous stimuli was increased because attentional bias has been especially found for stimuli associated with relatively high levels of threat (cf. [4,5,26,42]).

2.2.2. Spatial cueing task

Throughout the task, a black fixation cross (7 mm × 7 mm) was presented on a light grey background at the centre of the computer screen. Two black rectangular frames (6.5 cm high × 4.8 cm wide), serving as position markers for spatial cues, were continuously

displayed at the left and right of fixation (distance from screen centre to the centre of each frame = 9.8 cm). Participants were encouraged to fixate the cross consistently.

The typical trial configuration of the cueing task was as follows. After 1000 ms, a spatial cue (ie, coloured rectangle) appeared for 200 ms within the left or right frame, completely filling the frame. On half of the trials, the cue was pink; on the other half, green (eg, [43,45,46,48]). Each cue colour appeared equally often at either position. Thirty milliseconds after cue offset, a small target ("/" or "\"; 4 mm) appeared at the centre of either the left or the right frame. Following each cue colour, each target identity appeared equally often. Within each combination of cue colour and target identity, on half of the trials, the target appeared at the position previously occupied by the spatial cue (*valid trials*); on the other half of the trials, the target appeared at the other position (*invalid trials*). Participants' task was to press on each trial, as quickly and accurately as possible, the top key on a response box with the right index finger to "/" and the bottom key with the left index finger to "\". Targets were displayed until a response was made or for a maximum of 2000 ms. Intertrial interval randomly varied from 1000 ms to 1500 ms.

Attention to the visual cue is reflected in faster responses to targets at cue location than to targets at the other location (ie, *cue validity effect*) (eg, [29,55]).

2.2.3. Differential conditioning

During the test phase (see Section 2.3.4.), spatial cues were conditioned stimuli, with one of the colours (*pain cue*) followed by an aversive electrocutaneous stimulus, whereas the other colour was never followed by electrocutaneous stimulation (*no-pain cue*). On one-third of the trials in which the pain cue appeared, the electrocutaneous stimulus was delivered at cue offset (acquisition by partial reinforcement). The test phase was immediately preceded by a short acquisition phase (see Section 2.3.3.) during which the electrocutaneous stimulus was delivered at cue offset each time the pain cue appeared (acquisition by full reinforcement). Colour of pain cue and no-pain cue was counterbalanced between participants. Participants were informed that one of the colours (but not which colour) would predict the occurrence of the electrocutaneous stimulus.

Biases in attention to pain signals are reflected in larger cue validity effects for pain cues than for no-pain cues (eg, [13,46]). The size of attentional bias effects as measured with the spatial cueing task is in the order of milliseconds.

2.2.4. Nonpain goal task

Spatial cueing task trials were intermixed with digit trials. Digit trial configuration was similar to that for cueing task trials, except that a black digit (7 mm) also appeared for 50 ms at central fixation, replacing the cross. Digits from 1 to 9 were randomly selected and appeared throughout the trial (but not simultaneously with the target or with responses to the target, for technical reasons) as well as during the intertrial interval. Participants' task with regard to targets ("/" or "\") was the same as on cueing task trials. Participants' task with regard to digits depended on the group they were assigned to.

The *goal group* was instructed to read aloud each digit as quickly and accurately as possible. To enhance motivational salience of this *nonpain goal task*, the goal group was told that digit-naming performance would influence monetary compensation for their participation. It was explained that they would gain one point for each fast and accurate response, but lose one point for each slow, inaccurate, or missed response. Digit-naming responses were categorized as fast or slow based on a criterion that was adjusted after each response. Using this floating criterion, an equivalent proportion of relatively fast and slow responses was established

throughout the task. Intermediate scores were not presented to the participant following each digit, but during regular task breaks (see Section 2.3.). Instead of the individual's end score, an end score equal to 0 was given to all participants. The task started with a score of 0 (at start of baseline phase; see paragraph 2.3.2.). At the start of the session, participants assigned to the goal group received 2 €5 vouchers and were told that their compensation would depend on their end score (at the end of the test phase). They would receive €15 compensation (ie, €5 gain) with a positive end score, €5 compensation (ie, €5 loss) with a negative end score, and €10 compensation (ie, no gain, no loss) with an end score of 0. Because cueing task and goal task required different responses to different stimuli, the tasks were independent. So, the nonpain goal manipulation had no direct effect on cueing task performance, from which attentional bias to pain signals was derived. This is important in order to allow firm conclusions about effects of competing goals on pain-related attentional bias.

The *control group* received no instruction to respond to digits. They were informed that the digits were presented as an additional aid to focus on central fixation. Pilot testing indicated that undergraduates considered this a plausible explanation. Because digit trials were also presented to the control group, perceptual load of the task was equal for both groups.

2.2.5. Apparatus

Task presentation and response registration (latency and accuracy) were controlled by a Dell Optiplex GX 620 (Dell, Round Rock, TX, USA) computer that was running Presentation software (Neurobehavioral Systems, Albany, CA, USA) and that was connected to 2 19-inch Samsung Syncmaster 940 BF LCD (Samsung, Ridgefield Park, NJ, USA) monitors (one for the participant and one for the experimenter). In the goal group, verbal response latency was registered via a Sennheiser HMD/HME 25-1 (Sennheiser Electronic Corporation, Old Lyme, CT, USA) microphone/headphone combination connected to a voice key. In order to record verbal response accuracy, the experimenter manually entered the corresponding digit (or 0 in case of missing) at the end of each digit trial (numeric keypad). In order to establish comparable testing conditions for both groups, the control group also wore the microphone/headphone combination (supposedly as part of the intercom system and to attenuate distracting noise).

2.3. Procedure

Participants were tested individually at a viewing distance of about 60 cm from the computer screen in a dimly lit, sound-attenuated room designed for psycho-physiological experiments. They were video-monitored and could communicate with the experimenter (in a separate room) through an intercom. They were informed that the study investigated the relationship between concentration and performance and that during the computer task, sensory stimuli would be delivered through electrodes on their ankle; that these stimuli feel like pinpricks, stimulate pain nerves, and are perceived by the majority as unpleasant. The true purpose was explained after all participants had been tested.

Upon arrival in the laboratory, participants were randomly assigned to the goal group or the control group. Following electrocutaneous stimulus selection, both groups performed the computer task consisting of a mixture of cueing task trials and digit trials. Task trials were the same for both groups, but instructions differed. Whereas the goal group was instructed to respond manually to targets (“/” or “\”) on every trial and verbally to digits that appeared on 25% of the trials, the control group had only to respond to targets. Because the configuration of cueing task trials and digit trials was the same, except for the occurrence of digits,

digits were expected to appear throughout cueing task trials. So, for the goal group, stimuli for the additional goal task were also expected on trials on which attentional bias to pain signals was measured. The goal group received 2 vouchers worth €5 each before the computer task performance. These vouchers remained visible throughout the session. For both groups, the computer task consisted of the following phases.

2.3.1. Practice phase

The goal group practiced the cueing task, first without (32 cueing task trials) and then in combination with the digit-naming task (16 cueing task trials; 16 digit trials). The control group practiced the cueing task (2 × [16 cueing task trials; 16 digit trials]). No electrocutaneous stimuli were delivered and participants were informed about this. Following practice, all participants assigned to the goal group were able to repeat the rules for gaining/losing points and money.

2.3.2. Baseline phase

The baseline phase consisted of 96 cueing task trials and 32 digit trials. The goal group performed the cueing task in combination with the digit naming task; the control group performed only the cueing task. No electrocutaneous stimuli were delivered, and participants were informed about this.

2.3.3. Acquisition phase

The acquisition phase consisted of 8 cueing task trials (no digit trials) and was immediately followed by the test phase (no break in between). The goal group performed the cueing task in combination with the digit-naming task; the control group performed only the cueing task. Spatial cues were differentially conditioned. In all 4 trials in which a pain cue appeared, the electrocutaneous stimulus was delivered; in the other 4 trials, a no-pain cue was presented, never followed by the electrocutaneous stimulus.

2.3.4. Test phase

The test phase consisted of 144 cueing task trials and 48 digit trials. The goal group performed the cueing task in combination with the digit-naming task; the control group performed only the cueing task. Spatial cues were differentially conditioned. On one-third of the trials in which a pain cue appeared (24 cueing task trials; 8 digit trials), the electrocutaneous stimulus was delivered; on the other trials, no electrocutaneous stimuli were delivered.

During practice, baseline, and test phase, cueing task trials and digit trials were presented in a different random order for each participant. During the whole task, incorrect and premature responses to targets (“/” or “\”) were indicated by a short beep along with the display of an error message at screen centre for 500 ms (+1000-ms pause). Missed responses to these targets were also followed by a visual message lasting 500 ms (+1000-ms pause). Performance feedback was given (at screen centre) every 32 trials during short breaks terminated by the participant: mean reaction time (RT) and number of incorrect responses with regard to manual responses to targets and, to the goal group, intermediate score on the digit-naming task.

Following computer task performance, electrodes were detached and participants indicated on 11-point Likert rating scales (0 = “no at all”; 10 = “to a very large extent” or “extremely”) to what extent they expected green and pink cues to be followed by sensory stimulation, how fearful they were when green and pink cues were presented, and how painful and unpleasant they perceived the sensory stimulation during task performance. Finally, all participants received 2 vouchers worth €5 each. Each session took about 55 minutes to complete.

2.4. Design

This experiment employed a 2 (valid cueing vs invalid cueing) \times 2 (pain cue vs no-pain cue) \times 2 (goal group vs control group) factorial design with RT to targets as main dependent variable. Attentional bias for pain signals is reflected in differential cue validity effects for pain cues and no-pain cues during the test phase. The test phase was preceded by a baseline phase, without differential conditioning of the spatial cues, in order to assess whether cue validity effects do not differ as a function of distinctive visual features of the cues, independent of their conditioned signal value.

2.5. Data analysis

Attention effects were derived from performance on cueing task trials on which no electrocutaneous stimulus was delivered. Outlier participants who were slow or inaccurate on these trials during baseline and/or test phase were identified separately for each group (>2.5 SD above group mean) and were excluded from the analyses.

To examine whether the final groups differed in demographic variables and debriefing scores, χ^2 tests and independent *t*-tests were conducted. To determine whether differential conditioning had occurred, analyses of variance (ANOVAs) were conducted with cue identity (pain cue vs no-pain cue) as within-subjects factor, group (goal group vs control group) as between-subjects factor, and self-reported fear when the cue was presented or expectancy of electrocutaneous stimulation following the cue as dependent variable.

The reported RT analysis was based on *median* correct RTs to reduce the impact of outlier values. (The same pattern of results was obtained with *mean* correct RTs, also after exclusion of responses deviating more than 2.5 SDs from the mean latency per condition). All reported *P*-values are 2-tailed.

In order to assess whether cue validity effects differ as a function of distinctive features of the cues, median RTs on cueing task trials during baseline phase were subjected to a mixed ANOVA with cue validity (valid vs invalid) and cue identity (pain cue vs no-pain cue) as within-subjects factors, and group (goal group vs control group) as between-subjects factor. In order to assess group differences in attentional bias for pain signals (ie, cue validity \times cue identity), median RTs on cueing task trials during test phase were subjected to a mixed ANOVA with cue validity, cue identity, and group as factors. Significant interactions were broken down by using 2 \times 2 ANOVAs and *t*-tests, when appropriate. Magnitude of cue validity effects was calculated by subtracting median RTs on valid trials from median RTs on invalid trials; magnitude of differential cue validity effects for pain cues and no-pain cues was calculated by subtracting cue validity effects for no-pain cues from cue validity effects for pain cues.

Accuracy data (the log of percentage correct; [37]) were analysed in the same way as was done for RTs.

3. Results

3.1. Group characteristics

In total, 31 participants were assigned to the goal group and 32 to the control group. Five outlier participants who responded slowly or inaccurately were excluded, leaving 29 participants in the goal condition and 29 in the control condition. The final groups had the same gender ratio and did not differ in age (Table 1). The goal group perceived the sensory stimulation as somewhat less unpleasant and less painful (Table 1; Cohen's *d* = .64 and .72, respectively).

Self-reported fear and expectancy of electrocutaneous stimulation indicated that differential conditioning had occurred. That is, as can be seen in Table 1, participants were more fearful when pain cues were presented as compared to the presentation of no-pain cues; $F(1,56) = 90.3$, $P < 0.001$, $\eta_p^2 = .62$ ($d = 1.73$); and electrocutaneous stimulation was more often expected after pain cues than after no-pain cues; $F(1,56) = 91.4$, $P < 0.001$, $\eta_p^2 = .62$ ($d = 1.72$). These differences in fear and expectancy ratings between pain cues and no-pain cues, reflecting differential conditioning, were smaller in the goal group [fear: $t(28) = 5.0$, $P < 0.001$, $d = 1.33$; expectancy: $t(28) = 4.5$, $P < 0.001$, $d = 1.19$] than in the control group [fear: $t(28) = 8.6$, $P < 0.001$, $d = 2.40$; expectancy: $t(28) = 10.3$, $P < 0.001$, $d = 2.70$], as indicated by significant interaction effects of cue identity and group on fear; $F(1,56) = 4.0$, $P = 0.05$, $\eta_p^2 = .07$; and expectancy of electrocutaneous stimulation; $F(1,56) = 4.2$, $P < 0.05$, $\eta_p^2 = .07$. The goal group, as compared to the control group, reported a somewhat higher expectation of electrocutaneous stimulation following no-pain cues ($d = .59$) and, though nonsignificantly, a somewhat higher level of fear when no-pain cues were presented ($d = .48$), whereas the groups did not differ in those ratings for pain cues (Table 1). Note that fear and expectancy ratings for no-pain cues were very low in either group.

3.2. Spatial cueing task: RTs

3.2.1. Baseline phase

In the goal group, an average of 3.1% (SD = 2.1) of the responses on cueing task trials was incorrect; in the control group, 2.4% (SD = 1.8). Median correct RTs on cueing task trials (Table 2, left-hand columns) were subjected to an ANOVA with cue validity, cue identity, and group as factors. Responses were faster following valid cues than following invalid cues; $F(1,56) = 111.2$, $P < 0.001$, $\eta_p^2 = .67$ ($d = 2.02$). As expected, this cue validity effect (Table 2, left-hand columns) did not depend on cue identity: cue validity \times cue identity: $F(1,56) = 1.3$, $P = 0.25$, $\eta_p^2 = .02$; cue validity \times group: $F < 1.0$; for cue colour was not yet differentially conditioned during the baseline phase. Overall, the goal group responded somewhat faster and showed a smaller cue validity effect as compared to the control group [group: $F(1,56) = 7.1$, $P = 0.01$, $\eta_p^2 = .11$ ($d = .82$); cue validity \times group: $F(1,56) = 5.9$, $P < 0.05$, $\eta_p^2 = .10$].

3.2.2. Test phase

In the goal group, an average of 2.2% (SD = 1.7) of the responses on cueing task trials during which no electrocutaneous stimulus was delivered was incorrect; in the control group, 2.9% (SD = 1.9). Median correct RTs on cueing task trials during which no electrocutaneous stimulus was delivered (Table 2, right-hand columns) were subjected to an ANOVA with cue validity, cue identity, and group as factors. Responses were again faster following valid cues than following invalid cues; $F(1,56) = 69.6$, $P < 0.001$, $\eta_p^2 = .55$ ($d = 1.60$). In line with our hypotheses, the magnitude of this cue validity effect (Table 2, right-hand columns; Fig. 1) depended on cue identity as well as on group (cue validity \times cue identity \times group: $F(1,56) = 8.2$, $P < 0.01$, $\eta_p^2 = .13$; cue validity \times cue identity: $F < 1.0$; cue validity \times group: $F(1,56) = 5.3$, $P < 0.05$, $\eta_p^2 = .09$), indicating a group difference in attentional bias to pain signals. There were no other significant results from the ANOVA.

For the control group, the cue validity effect was larger for pain cues than for no-pain cues (mean difference = 11.1 ms, SD = 23.0), indicating attentional bias for pain signals [cue validity \times cue identity: $F(1,28) = 6.8$, $P < 0.05$, $\eta_p^2 = .20$; cue validity: $F(1,28) = 37.6$, $P < 0.001$, $\eta_p^2 = .57$ ($d = 1.77$); cue identity: $F(1,28) = 3.2$, $0.05 < P \leq 0.10$, $\eta_p^2 = .10$ ($d = .50$)]. In contrast, for the goal group, the cue validity effect seemed to be smaller for pain cues than

Table 1
Group sizes, mean age, and mean total scores (SD in parentheses) on debriefing questions (11-point Likert scales) at the end of the session, immediately after the computer task.

	Goal group n = 29 (5 men)	Control group n = 29 (5 men)	<i>t</i> ^a	<i>df</i>
Age (in years)	20.1 (1.6)	21.1 (2.5)	1.8 [#]	56
Unpleasantness sensory stimulation?	5.0 (1.8)	6.2 (2.1)	2.4 [*]	56
Painfulness sensory stimulation?	3.6 (1.9)	5.2 (2.5)	2.7 ^{**}	56
Expectancy sensory stimulation after pain cue?	5.6 (3.1)	6.3 (2.1)	1.1	49.33 ^b
Expectancy sensory stimulation after no-pain cue?	2.0 (2.3)	.9 (1.5)	-2.2 [*]	48.21 ^b
How fearful when pain cue?	3.8 (2.9)	4.8 (2.9)	1.4	56
How fearful when no-pain cue?	.8 (1.5)	.3 (.6)	-1.8 [#]	36.36 ^b

^a Independent *t*-test.

^b Equality of variances could not be assumed (Levene's test).

^{*} $P \leq 0.05$.

^{**} $P \leq 0.01$.

[#] $0.05 < P \leq 0.10$.

Table 2
Median correct RTs (in ms; SD in brackets) on cueing task trials during which no electrocutaneous stimulus was delivered, as a function of cue validity, cue identity, and group (baseline phase and test phase).

		Baseline phase			Test phase		
		Valid	Invalid	Cue validity effect	Valid	Invalid	Cue validity effect
Goal group	Pain cue	411.4 (46.7)	442.9 (50.6)	31.5 (32.3)	404.7 (43.1)	425.6 (44.5)	21.0 (22.2)
	No-pain cue	411.4 (44.6)	437.4 (47.9)	26.0 (22.3)	397.4 (41.8)	427.6 (50.0)	30.2 (31.7)
Control group	Pain cue	429.3 (31.3)	477.8 (50.2)	48.5 (36.4)	406.0 (32.8)	456.5 (56.2)	50.5 (45.3)
	No-pain cue	434.2 (36.9)	477.5 (49.4)	43.3 (35.7)	406.5 (29.2)	445.9 (47.0)	39.4 (36.4)

RT, reaction time.

Note: Magnitude of cue validity effects was calculated by subtracting median RTs on valid trials from median RTs on invalid trials.




Fig. 1. Mean cue validity effects of the goal group ($n = 29$) and the control group ($n = 29$) for pain cues and no-pain cues during the test phase. Magnitude of cue validity effects was calculated by subtracting median reaction times (RTs) on valid trials from median RTs on invalid trials. Error bars indicate the SE of the group average of cue validity effects in each condition.

for no-pain cues (mean difference = -9.2 ms, $SD = 30.6$), but this difference was not significant [cue validity \times cue identity: $F(1,28) = 2.6$, $P = 0.1$, $\eta_p^2 = .09$; cue validity: $F(1,28) = 36.9$, $P < 0.001$, $\eta_p^2 = .57$ ($d = 1.62$); cue identity: $F < 1.0$].

For pain cues, the cue validity effect was smaller in the goal group as compared to the control group [cue validity \times group: $F(1,56) = 9.9$, $P < 0.01$, $\eta_p^2 = .15$; cue validity: $F(1,56) = 58.1$, $P < 0.001$, $\eta_p^2 = .51$ ($d = 1.39$); group: $F(1,56) = 2.2$, $P = 0.14$, $\eta_p^2 = .04$ ($d = .39$)], but for no-pain cues, there was no significant group difference [cue validity \times group: $F(1,56) = 1.1$, $P = 0.31$, $\eta_p^2 = .02$; cue validity: $F(1,56) = 60.4$, $P < 0.001$, $\eta_p^2 = .52$ ($d = 1.52$); group: $F(1,56) = 1.8$, $P = 0.19$, $\eta_p^2 = .03$ ($d = .35$)].

3.3. Spatial cueing task: accuracy

ANOVAs of log percentage correct with cue validity, cue identity, and group as factors revealed a significant cue validity effect (ie, more accurate on valid than invalid trials) for the baseline phase; $F(1,56) = 17.7$, $P < 0.001$, $\eta_p^2 = .24$ ($d = .80$), and for the test phase; $F(1,56) = 14.1$, $P < 0.001$, $\eta_p^2 = .20$ ($d = .70$). There were no other significant effects.

4. Discussion

The present experiment investigated whether attentional bias to pain signals is reduced when a pain-unrelated but salient goal is pursued. Biased attention to pain signals was assessed within a spatial cueing task whilst half of the participants were also motivated to perform well on an independent goal task. As predicted, in participants who were motivated to engage in the goal task (goal group), attentional bias to pain signals, substantiated in a larger RT difference between valid and invalid conditions for pain cues than for no-pain cues, was smaller as compared to participants who only performed the cueing task (control group).

Recently, researchers have called for a motivational view on attention to pain, in which the role of a person's goals is taken into account [47]. Pain and fear of pain do not occur in a motivational vacuum, but in a context of different, possibly conflicting goals (eg, [6,16,24,51]). Efficient goal pursuit and self-regulation require that information processing can be flexibly adjusted, with attention oriented correspondingly (eg, [9,15,25]). It would, for example, be adaptive when the activation of one's focal goal inhibits the accessibility of alternative goals and other distracting information [14,18,41]. Attention has been considered as a central process of such "goal shielding" [15,40]. Our finding that attentional bias to pain (signals) can be influenced by the motivational context and by current goal pursuit underscores a dynamic and motivational view on attention to pain.

The observed reduction of attentional bias in the goal group as compared to the control group adds to previous findings outside

the field of pain that support the modulation of attentional bias by top-down engagement (eg, [22,27,31,32]). In addition, the present experiment extends in several ways previous studies within the field of pain that indicate that the disruption of task performance by experimentally induced pain can be modulated by the motivational salience of the task [49] and by working memory load during painful stimulation [20]. First, in the present experiment, attention to signals of impending pain, rather than pain itself, was studied and attentional processing was derived from spatial cueing effects, rather than only indirectly measured via interference effects. Second, goal pursuit was manipulated within a secondary task that was completely independent from the cueing task, measuring pain-related attentional bias. That is, the stimuli as well as response options for the goal (task) were not related to the stimuli and response options for the cueing task. By doing so, the goal manipulation had no influence on the way the cueing task was performed, and group differences in attentional bias could be solely interpreted in terms of differences in pain-unrelated goal pursuit.

Besides the reduced attentional bias to pain signals in the context of nonpain goal pursuit, supporting the main hypothesis of the present study, some further aspects of the data warrant discussion. *First*, the observed reduction of cue validity effects in the goal group, during baseline and test phase, adds to previous findings showing that attention to peripheral cues that do not predict target position (so-called exogenous cues) is suppressed when attentional resources are already focused elsewhere, for example, on a demanding task or perceptual display [34]. Santangelo et al. [34] observed a suppression of exogenous cueing effects (ie, no significant difference between performance on valid and invalid trials) when during cueing task performance participants' voluntary attention was engaged at the centre of the display – by presenting target digits (and distractor letters) at central fixation – independent of whether a response to these additional stimuli was required. These results suggest that reflexive orienting of attention is not truly automatic. In the present experiment, contrasting Santangelo et al.'s observation [34], cue validity effects for unconditioned exogenous cues (baseline phase) depended on whether a response to central digits was required (goal group) or not (control group). An important difference between the current task and the one applied by Santangelo et al. [34] is that our focused attention task was motivationally enhanced. Note that because digit task trials were also presented to the control group without requiring any response to them, the group difference in cueing effects cannot be due to a mere difference in perceptual load [19,34]. It might be valuable for future experiments to also avoid differences in attentional focus between goal conditions. Finally, Santangelo et al.'s study does not allow any conclusions about the attentional processing of affective cues in a motivational context [34]. The current finding of a reduced cue validity effect for conditioned spatial cues in the goal group (test phase) suggests that attention to pain signals is not entirely automatic, in the sense that it can be suppressed by focusing attentional resources on a salient, demanding, and potentially rewarding task.

Second, immediately after the computer task, participants in either group reported a higher expectation of electrocutaneous stimulation following pain cues than following no-pain cues, as well as a higher level of fear when pain cues were presented, indicating that the cues were indeed, as expected, differentially conditioned. These differential conditioning effects were somewhat smaller for the goal group as compared to the control group. Importantly, the group difference in attentional bias to pain signals was due to a group difference in cue validity effect following pain cues for which the groups did not differ in their rating of expectancy or fear. Therefore, the group difference in attentional bias

to pain signals cannot be explained in terms of the slight group difference in differential conditioning.

After the computer task, the goal group also evaluated the sensory stimulation during task performance as somewhat less unpleasant and less painful as compared to the control group. This finding is in line with previous studies showing that engagement in a more cognitively demanding task reduces pain ratings [10,38,47,49,50]. Exactly the same procedure was followed in both groups with regard to electrocutaneous stimulus selection and the above-mentioned pain and cue ratings. Hence, it can be concluded that the significant group differences in retrospective self-report ratings were due to the goal pursuit manipulation and are therefore not controlled for in the attentional bias analysis [23]. This not only holds for the pain and cue ratings, but also for the expectancy and fear ratings discussed before. Furthermore, from the present data, it is not possible to discern the relative effects of dual-task performance and bias in visuospatial orienting on these ratings.

Our finding that attentional bias to pain signals can be reduced in the context of nonpain goal pursuit may have clinical implications. Pain management programs that focus on the pursuit of valued, but pain-unrelated, goals may help reduce attention to pain and bodily threat, consequently improving daily functioning. It has been suggested that cognitive-behavioural therapies that promote the pursuit of daily life goals and the engagement in valued activities would be effective for treatment of chronic pain as well (eg, [3,36,44]). Without further research, our findings with pain-free volunteers and experimentally induced pain cannot simply be generalized to chronic pain (or other populations). Indeed, chronic pain, as well as various forms of psychopathology, have been thought to be characterized by inflexible, maladaptive goal pursuit, associated with a rigid attentional bias and weakened executive functioning (eg, [3,8,20,33,44]). If so, within a context of multiple-goal pursuit, chronic pain might be associated with a reduced inhibition of attentional bias to pain.

Some study limitations should be acknowledged. First, this experiment is the first of its kind and replication is needed. Second, clinical variables or variables affecting overall response speed and/or general motivation (eg, fatigue, alcohol, caffeine) were not systematically assessed. However, participants were randomly assigned to groups, which helped to ensure that there were no systematic group differences regarding such additional variables at the onset of the experiment. Future studies may benefit from more explicit consideration of these variables. Third, in order to address the main objective of this experiment, we induced concurrent nonpain goal pursuit by requiring participants to perform an additional task, the affective-motivational value of which was increased by associating task performance with monetary incentives; the control group was not instructed to perform an additional task. Therefore, the specific effects of additional task performance (cf. working memory load) and motivation manipulation cannot be disentangled with the present experimental design. It would be interesting for further research on pain-related attention to further examine the differential effect of affective-motivational and cognitive aspects or to focus on the impact of different goal orientations and goal task characteristics (cf. [32,33,47]). Fourth, no cues were included predicting the occurrence of non-painful somatosensory stimulation or the occurrence of another aversive outcome. So, it is not clear whether the observed effects are specific to pain. Prior research with a similar paradigm as the one applied in the present experiment to measure attentional orienting, but also including cues predicting nonpainful vibrotactile stimulation, has suggested that the modulation of attentional disengagement is pain-specific [43]. An interesting avenue would be to examine attentional bias to pain cues, relative to cues that are associated with nonpain goals (cf. [53,54]).

Conflict of interest statement

The authors have no conflicts of interest with regard to the study described in this manuscript.

Acknowledgments

The contribution of Martien Schrooten and Johan Vlaeyen was supported by the NWO Social Sciences Research Council of The Netherlands, Grant No. 453-04-003. Stefaan van Damme is supported by an IASP Early Career Research Grant and by the Research Foundation – Flanders (FWO – Vlaanderen). Julia Vogt is a Postdoctoral Fellow of the Research Foundation – Flanders (FWO – Vlaanderen). The authors thank Ron Hellenbrand for technical support and Rena Gatzounis for help in data acquisition.

References

- Bantick SJ, Wise RG, Ploughs A, Clare S, Smith SM, Tracey I. Imaging how attention modulates pain in humans using functional MRI. *Brain* 2002;125:310–9.
- Bingel U, Rose M, Gläscher J, Büchel C. fMRI reveals how pain modulates visual object processing in the ventral stream. *Neuron* 2007;55:157–67.
- Christiansen S, Oettingen G, Dahme B, Klinger R. A short goal-pursuit intervention to improve physical capacity: a randomized clinical trial in chronic back pain patients. *Pain* 2010;149:444–52.
- Crombez G, Eccleston C, Baeyens F, Eelen P. When somatic information threatens, catastrophic thinking enhances attentional interference. *Pain* 1998;75:187–98.
- Crombez G, Eccleston C, Baeyens F, Van Houdenhove B, Van den Broeck A. Attention to chronic pain is dependent upon pain-related fear. *J Psychosom Res* 1999;47:403–10.
- Crombez G, Eccleston C, Van Damme S, Vlaeyen JWS, Karoly P. Fear avoidance model of chronic pain: the next generation. *Clin J Pain*; in press.
- Crombez G, Van Damme S, Eccleston C. Hypervigilance to pain: an experimental and clinical analysis. *Pain* 2005;116:4–7.
- Derryberry D, Reed M. Anxiety-related attentional biases and their regulation by attentional control. *J Abnorm Psychol* 2002;111:225–36.
- Dijksterhuis A, Aarts H. Goals, attention, and (un)consciousness. *Annu Rev Psychol* 2010;61:467–90.
- Eccleston C. The attentional control of pain: methodological and theoretical concerns. *Pain* 1995;63:3–10.
- Eccleston C, Crombez G. Pain demands attention: a cognitive-affective model of the interruptive function of pain. *Psychol Bull* 1999;125:356–66.
- Fishbach A, Ferguson MF. The goal construct in social psychology. In: Kruglanski AW, Higgins ET, editors. *Social psychology: handbook of basic principles*. New York: Guilford Press; 2007. p. 490–515.
- Fox E, Russo R, Bowles R, Dutton K. Do threatening stimuli draw or hold visual attention in subclinical anxiety? *J Exp Psychol Gen* 2001;130:681–700.
- Goschke T, Dreisbach G. Conflict-triggered goal shielding: response conflicts attenuate background monitoring for prospective memory cues. *Psychol Sci* 2008;19:25–32.
- Johnson RE, Chang CD, Lord RG. Moving from cognition to behavior: what the research says. *Psychol Bull* 2006;132:381–415.
- Karoly P, Okun MA, Ruelhman LS, Pugliese JA. The impact of goal cognition and pain severity on disability and depression in adults with chronic pain: an examination of direct effects and mediated effects via pain-induced fear. *Cogn Ther Res* 2008;32:418–33.
- Kiss M, Driver J, Eimer M. Reward priority of visual target singletons modulates event-related potential signatures of attentional selection. *Psychol Sci* 2009;20:245–51.
- Kruglanski A, Kopetz C. The role of goal systems in self-regulation. In: Morsella E, Bargh JA, Gollwitzer PM, editors. *Oxford handbook of human action*. New York: Oxford University Press; 2009. p. 350–67.
- Lavie N. Distracted and confused? Selective attention under load. *Trends Cogn Sci* 2005;9:75–82.
- Legrain V, Crombez G, Verhoeven K, Mouraux A. The role of working memory in the attentional control of pain. *Pain* 2011;152:453–9.
- Legrain V, Van Damme S, Eccleston C, Davis KD, Seminowicz DA, Crombez G. A neurocognitive model of attention to pain: behavioral and neuroimaging evidence. *Pain* 2009;144:230–2.
- Maner JK, Gailliot MT, Miller SL. The implicit cognition of relationship maintenance: inattention to attractive alternatives. *J Exp Soc Psychol* 2009;45:174–9.
- Miller GA, Chapman JP. Misunderstanding analysis of covariance. *J Abnorm Psychol* 2001;110:40–8.
- Morley S, Eccleston C. The object of fear in pain. In: Asmundson GJ, Vlaeyen JW, Crombez G, editors. *Understanding and treating fear of pain*. Oxford, UK: Oxford University Press; 2004. p. 163–88.
- Norman DA, Shallice T. Attention to action: willed and automatic control of behaviour. In: Davidson RJ, Schwarz GE, Shapiro D, editors. *Consciousness and self-regulation: advances in research and theory*, vol. 4. London: Plenum Press; 1986. p. 1–18.
- Notebaert L, Crombez G, Vogt J, De Houwer J, Van Damme S, Theeuwes J. Attempts to control pain prioritize attention towards signals of pain: an experimental study. *Pain* 2011;152:1068–73.
- Papies EK, Stroebe W, Aarts H. The allure of forbidden food: on the role of attention in self-regulation. *J Exp Soc Psychol* 2008;44:1283–92.
- Pincus T, Morley S. Cognitive-processing bias in chronic pain: a review and integration. *Psychol Bull* 2001;127:599–617.
- Posner MI, Snyder CR, Davidson BJ. Attention and the detection of signals. *J Exp Psychol Gen* 1980;109:160–74.
- Riediger M, Freund AM. Interference and facilitation among personal goals: differential associations with subjective well-being and persistent goal pursuit. *Pers Soc Psychol Bull* 2004;30:1511–23.
- Rothermund K. Counter-regulation and control-dependency: affective processing biases in the service of action regulation. *Soc Psychol* 2011;42:56–66.
- Rothermund K, Voss A, Wentura D. Counter-regulation in affective attentional biases: a basic mechanism, that warrants flexibility in emotion and motivation. *Emotion* 2008;8:34–46.
- Rothermund K, Wentura D, Bak PM. Automatic attention to stimuli signalling chances and dangers: moderating effects of positive and negative goal and action contexts. *Cogn Emot* 2001;15:231–48.
- Santangelo V, Belardinelli MO, Spence C. The suppression of reflexive visual and auditory orienting when attention is otherwise engaged. *J Exp Psychol Hum Percept Perform* 2007;33:137–48.
- Schoth DE, Delgado Nunes V, Lioffi C. Attentional bias towards pain-related information in chronic pain; a meta-analysis of visual-probe investigations. *Clin Psychol Rev* 2012;32:13–25.
- Schrooten MGS, Vlaeyen JWS. Becoming active again? Further thoughts on goal pursuit in chronic pain. *Pain* 2010;149:422–3.
- Schweickert R. Separable effects of factors on speed and accuracy: memory scanning, lexical decision, and choice tasks. *Psychol Bull* 1985;97:530–46.
- Seminowicz DA, Davis KD. A re-examination of pain-cognition interactions: implications for neuroimaging. *Pain* 2007;130:8–13.
- Seminowicz DA, Davis KD. Interactions of pain intensity and cognitive load: the brain stays on task. *Cereb Cortex* 2007;17:1412–22.
- Shah JY. The automatic pursuit and management of goals. *Curr Direct Psychol Sci* 2005;14:10–3.
- Shah JY, Friedman R, Kruglanski AW. Forgetting all else: on the antecedents and consequences of goal shielding. *J Person Soc Psychol* 2002;83:1261–80.
- Van Damme S, Crombez G, Eccleston C. Retarded disengagement from pain cues: the effects of pain catastrophizing and pain expectancy. *Pain* 2002;100:111–8.
- Van Damme S, Crombez G, Eccleston C. The anticipation of pain modulates spatial attention: evidence for pain-specificity in high-pain catastrophizers. *Pain* 2004;111:392–9.
- Van Damme S, Crombez G, Eccleston C. Coping with pain: a motivational perspective. *Pain* 2008;139:1–4.
- Van Damme S, Crombez G, Eccleston C, Koster EHW. Hypervigilance to learned pain signals: a componential analysis. *J Pain* 2006;7:346–57.
- Van Damme S, Crombez G, Hermans D, Koster EHW, Eccleston C. The role of extinction and reinstatement in attentional bias to threat: a conditioning approach. *Behav Res Ther* 2006;44:1555–63.
- Van Damme S, Legrain V, Vogt J, Crombez G. Keeping pain in mind: a motivational account of attention to pain. *Neurosc Biobehav Rev* 2010;34:204–13.
- Van Damme S, Lorenz J, Eccleston C, Koster EHW, De Clerq A, Crombez G. Fear-conditioned cues of impending pain facilitate attentional engagement. *Clin Neurophysiol* 2004;34:33–9.
- Verhoeven K, Crombez G, Eccleston C, Van Ryckeghem DML, Morley S, Van Damme S. The role of motivation in distracting attention away from pain: an experimental study. *Pain* 2010;149:229–34.
- Villemure C, Bushnell MC. Cognitive modulation of pain: how do attention and emotion influence pain processing? *Pain* 2002;95:195–9.
- Vlaeyen JWS, Crombez G, Linton SJ. The fear-avoidance model of pain: we are not there yet. Comment on Wideman et al. "A prospective sequential analysis of the fear-avoidance model of pain" [Pain, 2009] and Nicholas "First things first: reduction in catastrophizing before fear of movement" [Pain, 2009]. *Pain* 2009;146:222–3.
- Vlaeyen JWS, Linton SJ. Fear-avoidance and its consequences in chronic musculoskeletal pain: a state of the art. *Pain* 2000;85:317–32.
- Vogt J, De Houwer J, Crombez G. Multiple goal management starts with attention: goal prioritizing affects the allocation of spatial attention to goal-relevant events. *Exp Psychol* 2011;58:55–61.
- Vogt J, De Houwer J, Moors A, Van Damme S, Crombez G. The automatic orienting of attention to goal-relevant stimuli. *Acta Psychol* 2010;134:61–9.
- Yantis S. Control of visual attention. In: Pashler H, editor. *Attention*. San Diego: Psychology Press; 1996. p. 223–56.