Response to "Vitamin D status is negatively correlated with retinal nerve fiber layer thickness in relapsing-remitting MS patients without acute optic neuritis" by Knippenberg et al.

Citation for published version (APA):

Knippenberg, S., Smolders, J., van den Maegdenbergh, M., Porro, G., Damoiseaux, J., & Hupperts, R. (2017). Response to "Vitamin D status is negatively correlated with retinal nerve fiber layer thickness in relapsing-remitting MS patients without acute optic neuritis" by Knippenberg et al. *Multiple Sclerosis Journal*, 23(1), 129-129. <https://doi.org/10.1177/1352458516652951>

Document status and date:

Published: 01/01/2017

DOI:

[10.1177/1352458516652951](https://doi.org/10.1177/1352458516652951)

Document Version:

Publisher's PDF, also known as Version of record

Document license:

Taverne

Please check the document version of this publication:

- A submitted manuscript is the version of the article upon submission and before peer-review. There can be important differences between the submitted version and the official published version of record. People interested in the research are advised to contact the author for the final version of the publication, or visit the DOI to the publisher's website.
- The final author version and the galley proof are versions of the publication after peer review.
- The final published version features the final layout of the paper including the volume, issue and page numbers.

[Link to publication](#)

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal.

If the publication is distributed under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license above, please follow below link for the End User Agreement:

www.umlib.nl/taverne-license

Take down policy

If you believe that this document breaches copyright please contact us at:

repository@maastrichtuniversity.nl

providing details and we will investigate your claim.

Download date: 26 Apr. 2023

A total of 10 subjects had a history of ON. Patient characteristics were not significantly different between MS subjects with ON or without ON. The worst performing eye was included in the analysis.

Like Burton et al., we observed a negative correlation between vitamin D status and RNFL thickness at baseline (Figure 1(a)). This correlation was stronger when only subjects without previous ON were analyzed ($r = -0.601$, $p = 0.011$). We investigated other outcomes of optic nerve function. Vitamin D status did not correlate with visual acuity, VEP P100 latency, or VEP amplitude (Figure 1(b)–(d), respectively). Stratification for a history of ON did not affect the results (data not shown). Finally, we assessed the correlation between vitamin D status at baseline and the 12-month change of RNFL thickness, VEP P100 latency, and VEP amplitude. We found no correlation between vitamin D status and change of RNFL thickness, change of VEP P100 latency, and change of VEP amplitude (Figure 1(e)–(g), respectively).

Although the negative correlation between 25(OH)D levels and RNFL thickness reported by Burton et al. was hypothesized to be due to inflammation of the optic nerve, we replicated this finding in a cohort without active ON. Therefore, an effect of inflammation is unlikely. Alternatively, a detrimental effect of 25(OH)D on RNFL thickness could be hypothesized but is not supported by the longitudinal data of Burton et al., and by the longitudinal OCT and VEP data in this letter. Providing alternative explanations is challenging. A higher ultraviolet (UV) exposure has been associated with a higher risk of macular degeneration.² Therefore, the lower vitamin D status in patients with a reduced RNFL thickness could be a biomarker for detrimental effects of UV-light exposure on the retina. Nevertheless, although the interpretation of the correlation between vitamin D status and RNFL thickness is uncertain, the beneficial effect of vitamin D status on ON recovery demonstrated by Burton et al. is exciting and warrants further investigation.

Declaration of Conflicting Interests

The author(s) declared no potential conflicts of interest with respect to the research, authorship, and/or publication of this article.

Response to “Vitamin D status is negatively correlated with retinal nerve fiber layer thickness in relapsing-remitting MS patients without acute optic neuritis” by Knippenberg et al.

Date received: 7 May 2016; accepted: 9 May 2016

Funding

The author(s) disclosed receipt of the following financial support for the research, authorship, and/or publication of this article: This project was sponsored by an unrestricted grant by TEVA.

References

1. Burton JM, Eliasziw M, Trufyn J, et al. A prospective cohort study of vitamin D in optic neuritis recovery. *Mult Scler*. Epub ahead of print 1 April 2016. DOI: 10.1177/1352458516642315.
2. Schick T, Ersoy L, Lechanteur YT, et al. History of sunlight exposure is a risk factor for age-related macular degeneration. *Retina* 2016; 36(4): 787–790.

**Stephanie Knippenberg^{1,2,3}, Joost Smolders^{1,4},
Marianne van den Maegdenbergh⁵,
Giorgio Porro⁶, Jan Damoiseaux⁷ and
Raymond Hupperts^{1,2}**

¹Academic MS Center Limburg, Department of Neurology, Zuyderland Medical Center, Sittard, The Netherlands

²School for Mental Health and Neuroscience, Maastricht University Medical Center, Maastricht, the Netherlands

³Department of Neurology, Zuyderland Medical Center, Heerlen, The Netherlands

⁴Department of Neurology, Canisius Wilhelmina Hospital, Nijmegen, The Netherlands

⁵Department of ophthalmology, Zuyderland-Eyescan, Sittard, the Netherlands

⁶Department of ophthalmology, Amphia Hospital, Breda, the Netherlands

⁷Central diagnostic laboratory, Maastricht University Medical Center, Maastricht, The Netherlands

Correspondence to:

S Knippenberg

Department of Neurology, Zuyderland Medical Center, Henri Dunantstraat 5, 6419 PC Heerlen, The Netherlands.

S.Knippenberg@maastrichtuniversity.nl

Dear *Multiple Sclerosis Journal* Editors and Dr Knippenberg and colleagues,

Thank you to Dr Knippenberg and colleagues for your letter.¹ Dr Knippenberg et al. report that in their study of 27 patients with relapsing-remitting multiple

Visit SAGE journals online
<http://msj.sagepub.com>

SAGE journals

Multiple Sclerosis Journal
2017, Vol. 23(1) 129–130

DOI: 10.1177/
1352458516652951

© The Author(s), 2016.
Reprints and permissions:
[http://www.sagepub.co.uk/
journalsPermissions.nav](http://www.sagepub.co.uk/journalsPermissions.nav)

sclerosis (RRMS) (with or without a history of optic neuritis), 25(OH)D values did negatively correlate with the initial retinal nerve fiber layer (RNFL) thickness measurement, regardless of optic neuritis history. In our prospective longitudinal study of acute optic neuritis of 36 patients, all patients had acute optic neuritis, and so the 25(OH)D measurements we obtained at presentation were temporally associated with all baseline optical coherence tomography (OCT) results including RNFL thickness, which was only elevated in those with a 25(OH)D <80 nmol/L (RNFL of 134.3 vs 95.2 μ m in sufficient patients, $p=0.003$). As well, we distinguished groups based on vitamin D status using a threshold 25(OH)D above or below 80 nmol/L, consistent with the biochemical definition of vitamin D sufficiency. The definition of vitamin D sufficiency may vary between different studies and thus could be a factor.

With respect to subsequent ophthalmological measures over time (6 months in our study, 12 months in Knippenberg et al.), we agree that we did not see significant differences in thinning in RNFL at 6 months between our vitamin D cohorts (79.3 vs 76.3 μ m in insufficient vs sufficient patients, respectively). However, in vitamin D insufficient patients, we did find evidence of significant neuronal loss in the ganglion cell layer (GCL) on OCT, as well as poorer visual recovery on high-contrast visual acuity testing. In our study, the GCL inter-eye difference, which uses the unaffected eye of the patient as another level of control (unaffected eye value minus affected eye value) differed significantly between vitamin D cohorts. In vitamin D insufficient patients, the GCL inter-eye difference was 14.2 μ m, while it was only 4 μ m in the sufficient group ($p=0.008$), indicating relatively greater thinning in insufficient patients. The use of the GCL measure is highly practical as it is not impacted by inflammation (as is the RNFL) and correlates with brain atrophy in MS patients.²

Admittedly, both our study and that by Knippenberg et al. are affected by a small sample size. Furthermore, differences in the patient cohorts between studies could be contributing to the different results found by the two studies. Knippenberg et al.'s inclusion of patients without a history of optic neuritis, and those

without acute optic neuritis, may present a challenge in elucidating the apparent impact of vitamin D on retinal structure measurements by OCT, as the timing of vitamin D sufficiency in the context of optic nerve injury itself could be an important variable. Additionally, we did not study ultraviolet (UV) exposure in our patients, so there is the possibility that this variable may be a contributing factor. Both studies have shown some very interesting correlations between OCT measurements of optic nerve integrity and vitamin D status, suggesting that further, larger scale studies of the relationship between vitamin D and optic nerve integrity are warranted.

Declaration of Conflicting Interests

The author(s) declared no potential conflicts of interest with respect to the research, authorship, and/or publication of this article.

Funding

The author(s) received no financial support for the research, authorship, and/or publication of this article.

References

1. Knippenberg S, Smolders J, van den Maegdenbergh M, Porro G, Damoiseaux J, & Hupperts R. Vitamin D status is negatively correlated with retinal nerve fiber layer thickness in relapsing-remitting MS patients without acute optic neuritis. *Mult Scler*. Epub ahead of print. DOI: 10.1177/1352458516652950
2. Saidha S, Al-Louzi O, Ratchford JN, et al. Optical coherence tomography reflects brain atrophy in multiple sclerosis: A four-year study. *Ann Neurol* 2015; 78: 801–813.

Jodie M Burton

Departments of Clinical Neurosciences and Community Health Sciences, Hotchkiss Brain Institute, University of Calgary, Calgary, AB, Canada

Correspondence to:

JM Burton

Departments of Clinical Neurosciences and Community Health Sciences, Hotchkiss Brain Institute, University of Calgary, Calgary, AB T2N 2T9, Canada.
jodie.burton@albertahealthservices.ca