Can we really conclude that treatment delays are associated with poorer outcome in patients with colorectal cancer awaiting elective surgery?

Citation for published version (APA):

Franssen, R. F. W., Strous, M. T. A., Vogelaar, F. J., & Janssen-Heijnen, M. L. G. (2021). Can we really conclude that treatment delays are associated with poorer outcome in patients with colorectal cancer awaiting elective surgery? *Colorectal Disease*, 23(8), 2197-2198. <https://doi.org/10.1111/codi.15731>

Document status and date:

Published: 01/08/2021

DOI:

[10.1111/codi.15731](https://doi.org/10.1111/codi.15731)

Document Version:

Publisher's PDF, also known as Version of record

Document license:

Taverne

Please check the document version of this publication:

- A submitted manuscript is the version of the article upon submission and before peer-review. There can be important differences between the submitted version and the official published version of record. People interested in the research are advised to contact the author for the final version of the publication, or visit the DOI to the publisher's website.
- The final author version and the galley proof are versions of the publication after peer review.
- The final published version features the final layout of the paper including the volume, issue and page numbers.

[Link to publication](#)

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal.

If the publication is distributed under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license above, please follow below link for the End User Agreement:

www.umlib.nl/taverne-license

Take down policy

If you believe that this document breaches copyright please contact us at:

repository@maastrichtuniversity.nl

providing details and we will investigate your claim.

Download date: 26 Apr. 2023

DATA AVAILABILITY STATEMENT

Data sharing not applicable to this article as no datasets were generated or analysed during the current study

John Bunni

Department of Colorectal Surgery, Royal United Hospitals Bath
NHS Foundation Trust, Bath, UK
Email: johnbunni@nhs.net

ORCID

John Bunni <https://orcid.org/0000-0003-2251-9201>

REFERENCES

1. Cecil T, Bunni J, Mehta A, editors. A Practical Guide to Peritoneal Malignancy. Boca Raton, FL: The PMI Manual. CRC Press; 2019.
2. Bonomi AM, Colombo F, Frontali A, Ferrario L, Danelli P. Laparoscopic appendectomy for a large appendiceal mucinous neoplasm - A video vignette. *Colorectal Dis.* 2021;23(5):1293

Received: 15 April 2021 | Accepted: 18 April 2021

DOI: 10.1111/codi.15731

Can we really conclude that treatment delays are associated with poorer outcome in patients with colorectal cancer awaiting elective surgery?

Dear Editor,

We read with great interest the systematic review by Whittaker et al. [1] published in *Colorectal Disease*. Their study aimed to explore the association between treatment delay (TD) and overall survival (OS) or disease-free survival (DFS) in patients undergoing elective surgery for colorectal cancer (CRC). The authors concluded that elective surgery for CRC should not be postponed for longer than 4 weeks. Although we endorse the importance of the aim of the study, we have some methodological concerns regarding this systematic review and meta-analysis.

Considering the wide heterogeneity in the definition of TD, it is questionable whether it is valid to perform a meta-analysis of the included studies. Although the end of the TD is generally clearly defined, the time point used as diagnosis, which is mostly taken as the starting point of TD, is defined less consistently. Definitions of diagnosis can vary from first investigation for defining malignancy to diagnosis confirmed by a multidisciplinary team meeting [2]. From experience in our own hospital, these time-points can be as far as 7–14 days apart. In some studies, such as that of Kucejko et al. [3], diagnosis was not specified at all. Heterogeneity in the definition of TD makes it questionable whether it is valid to pool hazard ratios from the included studies in a meta-analysis.

Besides our concern about the methodology, we believe that the conclusion is not sufficiently supported by the data presented. The authors state that elective surgery for CRC should not be postponed for longer than 4 weeks. This interval seems to be chosen arbitrarily, as this 4-week interval is not consistent with the intervals used in the included studies. For example, the study of Strous et al. [4] used

a cut-off point of >5 weeks. In addition, with regard to the inconsistencies in definition of diagnosis, it is unclear how this maximum TD of 4 weeks (or any TD) should be interpreted (i.e. 4 weeks starting when?).

The conclusion seems to be based on the association between TD and OS, while the association between TD and DFS is far more interesting with regard to treatment delays and preparation for surgery. It is well known that OS is influenced by many factors. Some studies show that frailer patients often have a longer TD, which might at least partially explain a possible association between a longer TD and OS [3]. As also acknowledged by the authors, not all included studies adjusted their analysis for patient factors associated with frailty [5]. As the main concern with longer treatment delays is tumour growth and risk for metastasis, DFS should be the main outcome of this systematic review.

We agree with the authors that TDs other than those that are needed for pretreatment work-up or optimization of health should be avoided in order to provide optimal care and minimize patient distress. However, especially in patients at high risk for complications, interventions aimed at reducing complications might be more instrumental than aiming for short treatment delays. Perhaps a shift in thinking from 'treatment delay' to 'optimization period' would allow for more individualized 'treatment intervals'.

Ruud F. W. Franssen^{1,2}

Maud T. A. Strous^{2,3}

F. Jeroen Vogelaar³

Maryska L. G. Janssen-Heijnen^{2,4}

¹Department of Clinical Physical Therapy, VieCuri Medical Center, Venlo, The Netherlands

²Department of Epidemiology, GROW School for Oncology and Developmental Biology, Faculty of Health, Medicine and Life Sciences, Maastricht University, Maastricht, The Netherlands

³Department of Surgery, VieCuri Medical Center, Venlo, The Netherlands

⁴Department of Epidemiology, VieCuri Medical Center, Venlo, The Netherlands

ORCID

Ruud F. W. Franssen <https://orcid.org/0000-0003-2911-4020>

REFERENCES

1. Whittaker TM, Abdelrazek MEG, Fitzpatrick AJ, Froud JLJ, Kelly JR, Williamson JS, et al. Delay to elective colorectal cancer surgery and

implications for survival: a systematic review and meta-analysis. *Colorectal Dis.* 2021;23(7):1699–1711. <https://doi.org/10.1111/codi.15625>

2. Molenaar CJL, Janssen L, van der Peet DL, Winter DC, Roumen RMH, Slooter GD. Conflicting guidelines: a systematic review on the proper interval for colorectal cancer treatment. *World J Surg.* 2021. [Epub ahead of print].
3. Kucejko RJ, Holleran TJ, Stein DE, Poggio JL. How soon should patients with colon cancer undergo definitive resection? *Dis Colon Rectum.* 2020;63(2):172–82.
4. Strous MTA, Janssen-Heijnen MLG, Vogelaar FJ. Impact of therapeutic delay in colorectal cancer on overall survival and cancer recurrence – is there a safe timeframe for prehabilitation? *Eur J Surg Oncol.* 2019;45(12):2295–301.
5. Grass F, Behm KT, Duchalais E, Crippa J, Spears GM, Harmsen WS, et al. Impact of delay to surgery on survival in stage I–III colon cancer. *Eur J Surg Oncol.* 2020;46(3):455–61.

Received: 4 May 2021 | Accepted: 8 May 2021

DOI: 10.1111/codi.15730

Reply to: Franssen, Ruud; Strous, Maud; Vogelaar, F Jeroen; Janssen-Heijnen, Maryska. Can we really conclude that treatment delays are associated with poorer outcome in patients with colorectal cancer awaiting elective surgery?

Dear Sir,

The authors of the original systematic review would like to thank Franssen et al. for their correspondence with regard to our recent work. The feedback and constructive criticism provided is greatly appreciated.

Given the lack of a recent meta-analysis on this topic [1] we attempted to quantify the delay from diagnosis to surgical treatment in elective colorectal cancer from the available literature. The evidence base available to the authors was of poor quality as acknowledged by the ROBINS-I score which declared three papers at serious risk of bias. The lack of uniformity in delay time at 4 weeks was acknowledged; however, it was the most reported interval and national guidelines recommend this interval. Thus, we felt it was the most appropriate approximation for a short delay interval.

As there is no consensus in the evidence base on the categorization of delay, we still believe future work should report the association between delay and mortality as a continuous variable. This will allow uniformity in statistical analyses so that future work can compare correlations directly.

We would like to highlight that a significant conclusion from this meta-analysis was the lack of consistency in the literature. From our work, a statistically significant correlation between delay and overall survival at 4 and 12 weeks justified a cautious recommendation for clinicians counselling patients. This is particularly prescient given the worldwide pandemic and current delays to surgical waiting times [2].

The categorization for 12 weeks was consistently reported in the cited studies. Therefore, a meta-analysis at this delay point was warranted and important to assess for patient safety. We do agree that even with this time interval the three studies adjusted for varying confounding variables. However, the error was reduced by using a random inverse model as suggested by the Cochrane Handbook [3].

We agree with the final point made by [4] that the interval between diagnosis and treatment could be an opportunity for optimization and should be utilized in the most efficient manner possible. Perhaps future research could address the optimum point for treatment delay and the benefits of prehabilitation. As highlighted by our results, the majority of studies did not describe the reason for