Men and women have an equal oropharyngeal and anorectal *Chlamydia trachomatis* bacterial load

Citation for published version (APA):

Wijers, J. N. A. P., Dukers-Muijrers, N. H. T. M., van Liere, G. A. F. S., Dirks, J. A. M. C., Wolffs, P. F. G., & Hoebe, C. J. P. A. (2021). Men and women have an equal oropharyngeal and anorectal *Chlamydia trachomatis* bacterial load: A Comparison of 3 Anatomic Sites. *Journal of Infectious Diseases*, 223(9), 1582-1589. <https://doi.org/10.1093/infdis/jiz668>

Document status and date:

Published: 01/05/2021

DOI:

[10.1093/infdis/jiz668](https://doi.org/10.1093/infdis/jiz668)

Document Version:

Publisher's PDF, also known as Version of record

Document license:

Taverne

Please check the document version of this publication:

- A submitted manuscript is the version of the article upon submission and before peer-review. There can be important differences between the submitted version and the official published version of record. People interested in the research are advised to contact the author for the final version of the publication, or visit the DOI to the publisher's website.
- The final author version and the galley proof are versions of the publication after peer review.
- The final published version features the final layout of the paper including the volume, issue and page numbers.

[Link to publication](#)

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal.

If the publication is distributed under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license above, please follow below link for the End User Agreement:

www.umlib.nl/taverne-license

Take down policy

If you believe that this document breaches copyright please contact us at:

repository@maastrichtuniversity.nl

providing details and we will investigate your claim.

Download date: 26 Apr. 2023

Men and Women Have an Equal Oropharyngeal and Anorectal *Chlamydia trachomatis* Bacterial Load: A Comparison of 3 Anatomic Sites

Juliën N. A. P. Wijers,^{1,2} Nicole H. T. M. Dukers-Muijers,^{1,2} Geneviève A. F. S. van Liere,^{1,2} Jeanne A. M. C. Dirks,¹ Petra F. G. Wolffs,¹ and Christian J. P. A. Hoebe^{1,2}

¹Department of Social Medicine and Medical Microbiology, Care and Public Health Research Institute Maastricht University Medical Center, Maastricht, the Netherlands, and ²Department of Sexual Health, Infectious Diseases, and Environmental Health, South Limburg Public Health Service, Heerlen, the Netherlands

Background. The *Chlamydia trachomatis* bacterial load could have impact on transmission and sequelae. This is the first study providing comparison of *C. trachomatis* load at 3 anatomic sites estimated by cycle quantification (Cq) values.

Methods. Data from 7900 *C. trachomatis*-positive samples were included (2012–2018). Cq value was used as an inversely proportional measure for *C. trachomatis* load. Multivariable linear regression analyses assessed differences in mean Cq values.

Results. Vaginal swabs had the lowest Cq values (31.0) followed by urine (32.5), anorectal swabs (34.0), and oropharyngeal swabs (36.8) ($P < .001$). Men and women had similar oropharyngeal (36.4 vs 37.3; $P = .13$) and anorectal (34.2 vs 33.9; $P = .19$) Cq values. Men (32.2) and women (30.7) aged <25 years had lower urogenital Cq values than men (32.8) and women (31.9) aged ≥ 25 years ($P < .001$). HIV-positive patients had higher urogenital Cq values than HIV-negative patients (33.8 vs 32.6; $P < .03$).

Conclusions. Men and women have a similar *C. trachomatis* load at extragenital locations arguing for similar transmission potential and clinical relevance. Older patients and HIV-coinfected patients had lower *C. trachomatis* load, suggesting exposure to previous *C. trachomatis* infections potentially leading to partial immunity reducing load.

Keywords. *Chlamydia*; bacterial load; urogenital; anorectal; oropharyngeal; extragenital.

Chlamydia trachomatis is the most reported bacterial sexually transmitted infection (STI) worldwide [1]. *C. trachomatis* is associated with reproductive sequelae in women, such as pelvic inflammatory disease, ectopic pregnancy, infertility, and chronic lower abdominal pain [2].

The *C. trachomatis* bacterial load, frequently expressed as the number of bacteria/mL, could potentially affect transmission of the disease and sequelae [3, 4]. Currently, it is not clear what determines a high *C. trachomatis* load in a patient. Although symptoms may be associated with a higher *C. trachomatis* load, studies to date show a weak association [3, 5, 6]. According to a systematic review, only a few nucleic acid amplification test (NAAT)-based studies on associations with *C. trachomatis* load have been conducted [3, 5, 7–10]. Sample sizes of these studies were small and most studies had a small number of cases within categories of independent determinants [3, 5, 7–10]. The

assessment of determinants for a high *C. trachomatis* load could be relevant. For example, viral STIs like human immunodeficiency virus (HIV) and herpes simplex virus have shown increased transmission potential with higher viral load [3]. Likely the transmission potential of *C. trachomatis* is also dependent on *C. trachomatis* load [3].

Extragenital infections, mainly anorectal infections, are common among men who have sex with men (MSM) and women [11]. Several studies have assessed the *C. trachomatis* load of different anatomic locations and sample types [3, 5, 7, 12, 13]. However, those studies used different methods to quantify and report *C. trachomatis* load making them less comparable. Therefore, standardization of *C. trachomatis* load measurements is needed [14]. In the current study, we compared the *C. trachomatis* load of all relevant urogenital and extragenital sites using the same methods.

Coinfections with *Neisseria gonorrhoeae* are prevalent among *C. trachomatis*-positive high-risk individuals such as MSM, young people (aged <25 years), and some ethnic groups, and these coinfections might influence *C. trachomatis* load [15–17]. According to a systematic review, only 6 studies to date have assessed the association between coinfections and *C. trachomatis* load [3]. One such study observed a higher anorectal *C. trachomatis* load among *C. trachomatis*-positive men coinfecting with anorectal *N. gonorrhoeae* compared to anorectal *N. gonorrhoeae*-negative men [18]. Another study

Received 11 October 2019; editorial decision 10 December 2019; accepted 13 December 2019; published online December 16, 2019.

Presented in part: STI & HIV 2019 World Congress, 14–17 July 2019, Vancouver, Canada.

Correspondence: Juliën Wijers, Msc, Department of Sexual Health, Infectious Diseases and Environmental Health, South Limburg Public Health Service, PO Box 33, 6400 AA, Heerlen, The Netherlands (julien.wijers@ggdz.nl).

The Journal of Infectious Diseases® 2021;223:1582–9

© The Author(s) 2019. Published by Oxford University Press for the Infectious Diseases Society of America. All rights reserved. For permissions, e-mail: journals.permissions@oup.com. DOI: 10.1093/infdis/jiz668

suggested a lower urogenital *C. trachomatis* load among women concurrently infected with *C. trachomatis* and *N. gonorrhoeae*; however, this was not statistically significant ($P = .06$) [19]. Associations between *C. trachomatis* load and coinfections with other STIs like HIV and syphilis were not observed and this is still unclear due to lack of research evidences [3, 12, 18, 20]. Nevertheless, coinfection with other STIs might have impact on acquiring and transmitting *C. trachomatis*.

To our knowledge, this is the first study to date providing a comparison of *C. trachomatis* load of all anatomic sites in both men and women estimated by cycle quantification (Cq) values. Our objectives were to assess whether different sample types have a different *C. trachomatis* load and whether age and coinfection with *N. gonorrhoeae*, HIV, or syphilis were associated.

METHODS

Study Population

Data of *C. trachomatis*-positive patients (aged ≥ 16 years) from January 2012 to June 2018 were used in this cross-sectional study. Data derived from 7224 consultations (with 1 or more *C. trachomatis*-positive samples available) originating from 6170 *C. trachomatis*-positive patients (from a total of $n = 62\,306$ *C. trachomatis* test consultations; 11.6% *C. trachomatis* positive) were obtained from the laboratory registry of the Medical Microbiology Laboratory of Maastricht University Medical Center. All samples were tested for *C. trachomatis* with the same NAAT for both plasmid and chromosomal DNA (Cobas 4800, Roche Diagnostics), as per the manufacturer's protocol [21]. We used the NAAT-derived Cq value as a proxy for *C. trachomatis*

load. The Cq value reflects the number of amplification cycles that occur before a positive *C. trachomatis* signal is detected. Thus Cq values show inverse relationships, that is a low Cq value indicated a high load and vice versa [22–24].

Patients were tested by taking samples from different anatomic locations (urogenital, anorectal, and oropharyngeal) according to testing guidelines and indications [25]. The study included data from all *C. trachomatis*-positive samples for which the Cq value could be retrieved (99.2%; 7900/7965; Figure 1). For men, data from first void urine samples (further referred to as urine samples) and urethral swabs were available. Mean Cq values from urine samples ($n = 2601$, mean = 32.49 [SD 3.18]) and urethral swabs ($n = 10$, mean = 33.47 [SD 4.89]) were similar ($P = .54$), therefore urethral swabs of men were excluded from the analysis. For women, urogenital data were available from urine samples, vaginal swabs, and cervical swabs. The mean Cq values from vaginal swabs ($n = 3273$, mean = 31.00 [SD 3.72]) and cervical swabs ($n = 147$, mean = 30.49 [SD 3.94]) were comparable in women ($P = .11$), therefore values for cervical swabs were excluded from the analysis.

Cycle Quantification Validation for *C. trachomatis* Load

In order to ensure that the Cq value can indeed be used as a proxy for *C. trachomatis* load in the different sample types, we compared the Cq values obtained from the Cobas system with previously determined quantitative in-house polymerase chain reaction (PCR) [12, 21, 26]. Absolute *C. trachomatis* load values (bacteria/mL) were available from a subset of the clinical STI clinic population: 129 urine samples and 69 anorectal swabs from men, and 403 vaginal swabs and 101 anorectal

Figure 1. Flowchart, including urogenital and extragenital *Chlamydia trachomatis* samples from men and women. Abbreviation: Cq, cycle quantification.

samples from women. In brief, we quantified *C. trachomatis* load by an in-house TaqMan real-time PCR to quantify *C. trachomatis* *OmpA*-gene copies/mL [21]. A full description of the *C. trachomatis* load quantification has been described elsewhere [21].

For men, moderate correlation between Cq values of urine samples and urine *C. trachomatis* load (Pearson r , -0.61 ; $n = 129$; $P \leq .001$), and high correlation between Cq values of anorectal samples and anorectal *C. trachomatis* load (Pearson r , -0.93 ; $n = 69$; $P \leq .001$) were observed. For women, high correlation between Cq values of vaginal swabs and vaginal *C. trachomatis* load (Pearson r , -0.88 ; $n = 403$; $P \leq .001$), and high correlation between Cq values of anorectal swabs and anorectal *C. trachomatis* load (Pearson r , -0.96 ; $n = 101$; $P \leq .001$) were observed. In addition, Cobas Cq values are highly correlated (Pearson $R^2 > 0.98$) with known concentrations of *C. trachomatis* (inclusion forming units/mL) (personal communication, B. van der Veer, unpublished results). Therefore, we argue that Cq values are a valid proxy for *C. trachomatis* load in urogenital and extragenital samples used in this study.

Statistical Analyses

In the main analyses, the association between sample type and the outcome measure Cq values, as a proxy for *C. trachomatis* load, was assessed using univariable and multivariable linear regression analyses adjusting for sex and age (<25 years, ≥ 25 years). Anorectal samples were used as the reference group.

In the secondary analyses, association of sex (women, men), age (<25 years, ≥ 25 years), urogenital *N. gonorrhoeae* coinfection, anorectal *N. gonorrhoeae* coinfection, oropharyngeal *N. gonorrhoeae* coinfection, HIV coinfection, and a positive syphilis screening test (*Treponema pallidum* hemagglutination assay/*T. pallidum* particle agglutination assay [TPHA/TPPA]) were assessed with the *C. trachomatis* load per sample type, that is vaginal swab, urine sample, oropharyngeal swab, and anorectal swab.

In the linear regression analyses, determinants with $P < .05$ in the univariable model were included in the multivariable model. A P value of $< .05$ was considered statistically significant. Mean values, standard deviations, (adjusted) mean differences, and 95% confidence intervals (CI) were calculated. Mean differences represented the size of the associations found.

As Cq values represent values on a log scale, this is a less accessible and readily measure. To make it more accessible a “factor *C. trachomatis* load” that represents the number of times the *C. trachomatis* load was higher between studied groups was calculated for the statistically significant determinants using the formula: $2^{(\text{adjusted mean difference of Cq values})}$. For example, the adjusted mean difference of Cq values between vaginal swabs and anorectal swabs was -2.87 Cq. Therefore, factor load was $2^{2.87} = 7.3$. Thus, vaginal swabs have a 7.3-times higher *C. trachomatis* load compared to anorectal swabs.

All analyses were performed using SPSS Version 24 (IBM SPSS Statistics for Windows, IBM Corporation).

Ethical Clearance

The Medical Ethics Committee of the Maastricht University Medical Center (Maastricht, the Netherlands) approved this study (METC 2017-0251) and waived the need for patient consent. Because the retrospective data originated from regular care and were analyzed anonymously, no further informed consent for data analysis was obtained.

RESULTS

C. trachomatis Load Per Sample Type, Estimated by Cycle Quantification Values

In multivariable analyses, vaginal swabs (-2.87 Cq) and urine samples (-1.33 Cq) had lower Cq values than anorectal swabs, which represented 7.3- and 2.5-times higher *C. trachomatis* load, respectively. Oropharyngeal Cq values were higher (2.73 Cq) compared to anorectal swabs, indicating a 6.6-times lower *C. trachomatis* load (Table 1).

The range of Cq values per sample type were comparable for men and women (Figure 2A). Men and women had similar urine ($P = .15$), anorectal ($P = .19$), and oropharyngeal ($P = .13$) Cq values indicative of similar *C. trachomatis* loads (Figure 2B). The mean Cq values from vaginal swabs ($n = 3273$, mean = 31.00 [SD 3.7]) and urine samples ($n = 290$, mean = 32.82 [SD 3.7]) in women were significantly different ($P \leq .001$).

Determinants Associated With *C. trachomatis* Load Stratified Per Sample Type

In multivariable analyses, women aged <25 years had lower vaginal Cq values (-0.69 Cq) compared to women aged ≥ 25 years, resulting in a 1.6-times higher *C. trachomatis* load in the former (Table 2). HIV-positive women had higher vaginal Cq values (6.86 Cq) compared to HIV-negative women, which represented a 116.2-times lower *C. trachomatis* load.

We additionally assessed whether or not the 3 HIV-infected women were repeatedly infected with *C. trachomatis*. All were not repeatedly infected with *C. trachomatis*. However, because the analyses included only 3 HIV-infected women this result was unreliable.

In multivariable analyses, patients aged <25 years had lower urine Cq values (-0.52 Cq) compared to patients aged ≥ 25 years, indicating a 1.4-times higher *C. trachomatis* load (Table 2). Patients coinfecting with urogenital *N. gonorrhoeae* had higher urine Cq values (0.96 Cq) compared to urogenital *N. gonorrhoeae*-negative patients, representing a 1.9-times lower *C. trachomatis* load. HIV-positive patients had higher urine Cq values (0.95 Cq) compared to HIV-negative patients, which represented a 1.9-times lower *C. trachomatis* load.

We additionally assessed whether urogenital *N. gonorrhoeae* and HIV-coinfecting patients with urine samples had repeat *C. trachomatis* infections (≥ 2 infections) or reported to have

Table 1. Main Analyses Including Different Sample Types and Associations With Sex and Age for *Chlamydia trachomatis* Cycle Quantification Values as a Proxy for *C. trachomatis* Load

Cq Values					
Determinants	% (n)	Mean Cq Value (SD)	Mean Difference (95% CI)	Adjusted Mean Difference (95% CI)	Factor CT Load
Overall	100 (7743)				
Sample type					
Vaginal swab ^a	42.3 (3273)	31.0 (3.7)	−3.02 (−3.24 to −2.80)	−2.87 (−3.13 to −2.62)^b	7.3
Urine sample	37.3 (2891)	32.5 (3.2)	−1.50 (−1.74 to −1.28)	−1.33 (−1.58 to −1.08)^c	2.5
Oropharyngeal swab	1.6 (124)	36.8 (3.1)	2.79 (2.14 to 3.44)	2.73 (2.08 to 3.38)	6.6
Anorectal swab	18.8 (1455)	34.0 (3.8)	Ref	Ref	
Sex ^d					
Women	56.2 (4354)	31.6 (4.0)	−1.24 (−1.40 to −1.07)	0.21 (−.06 to .49)	
Men	43.8 (3389)	32.9 (3.4)	Ref	Ref	
Age ^d					
<25 y	60.3 (4666)	31.7 (3.7)	−1.38 (−1.55 to −1.21)	−0.76 (−.93 to −.59)	1.7
≥25 y	39.7 (3077)	33.1 (3.7)	Ref	Ref	

Statistically significant associations ($P < .05$) are depicted in bold.

Abbreviations: CI, confidence interval; Cq, cycle quantification; CT, *Chlamydia trachomatis*; Ref, reference.

^aOnly measured among women.

^bWhen using oropharyngeal samples as the reference group the adjusted mean difference for vaginal swabs was −5.60 (95% CI, −6.26 to 4.95), factor CT load, 48.5.

^cWhen using oropharyngeal samples as the reference group the adjusted mean difference for urine was −4.06 (95% CI, −4.70 to 3.41), factor CT load, 16.7.

^dIncluding all sample types, that is vaginal swabs, urine samples, oropharyngeal swabs, and anorectal swabs.

had an earlier STI (only among STI clinic patients). HIV-coinfected patients more often had repeat *C. trachomatis* infections compared to HIV-negative men (24.4% vs 6.4%; $P < .001$). Patients coinfecting with HIV reported to have had an earlier STI more often compared to HIV-negative men (35.7% vs 16.0%; $P < .001$). Patients coinfecting with *N. gonorrhoeae* had similar repeat *C. trachomatis* infections compared to urogenital *N. gonorrhoeae*-negative patients (11.2% vs 9.1%; $P = .69$). Patients coinfecting with *N. gonorrhoeae* frequently reported to have had an earlier STI compared to *N. gonorrhoeae*-negative patients (31.7% vs 19.5%; $P = .07$).

No determinants were associated with oropharyngeal or anorectal *C. trachomatis* load in multivariable analyses (Supplementary Table 1).

DISCUSSION

This is the first and largest study to date providing a comparison of *C. trachomatis* load, estimated by Cq values, in urogenital and extragenital locations of men and women. The mean *C. trachomatis* load differed significantly per sample type and subsequently per anatomic location. Notably, no difference ($P > .05$) in *C. trachomatis* load was observed for men and

Figure 2. Boxplots and error bars showing the distribution of Cq values per sample type for men and women. A, Boxplots showing the distribution of Cq values as a proxy for *Chlamydia trachomatis* load per sample type for men (grey) and women (white). Middle lines are used to indicate median, boxes are used to indicate 25th and 75th percentile and whiskers indicate lowest and highest value. Outliers are indicated by circles. B, Mean Cq values and 95% CI (error bars) as a proxy for *C. trachomatis* load per sample type for men (dotted line) and women (solid line). Univariable regression analyses were performed to test unadjusted mean Cq value differences between men and women. Men and women had similar Cq values: urine (mean [SD] = 32.4 Cq [3.2] vs mean = 32.8 Cq [3.7]; $P = .15$), oropharyngeal (mean = 36.4 Cq [3.4] vs mean = 37.3 Cq [2.7]; $P = .13$), and anorectal (mean = 34.2 Cq [3.6] vs mean = 33.9 Cq [3.9]; $P = .19$). Abbreviations: CI, confidence interval; Cq, cycle quantification.

Table 2. Additional Analyses Including Potential Determinants Associated With Vaginal and Urine *Chlamydia trachomatis* Cycle Quantification Values as a Proxy for *C. trachomatis* Load

	Vaginal Cq Values				Urine Cq Values				Factor CT Load
	% (n)	Mean Cq Value (SD)	Mean Difference (95% CI)	Adjusted Mean Difference (95% CI)	Factor CT Load	% (n)	Mean Cq Value (SD)	Mean Difference (95% CI)	Adjusted Mean Difference (95% CI)
Overall	100 (3273)					100 (2891)			
Sex									
Women	100 (3273)					10.0 (290)	32.8 (3.7)	0.33 (−0.06 to .72)	
Men						90.0 (2601)	32.5 (3.2)	Ref	
Age									
<25 y	75.4 (2468)	30.7 (3.6)	−1.13 (−1.42 to −.84)	−0.69 (−1.00 to −.38)	1.6	53.3 (1542)	32.2 (3.2)	0.60 (−.83 to −.36)	−0.52 (−0.78 to −.27)
≥25 y	24.6 (805)	31.9 (3.9)	Ref	Ref		46.7 (1349)	32.8 (3.3)	Ref	Ref
Urogenital NG coinfection ^a									
Yes	2.9 (95)	31.8 (3.3)	0.78 (.02 to 1.54)		4.0 (116)		33.7 (3.2)	1.20 (.60 to 1.81)	0.96 (0.32 to 1.59)
No	95.0 (3108)	31.0 (3.7)	Ref		92.5 (2675)		32.5 (3.2)	Ref	Ref
Anorectal NG coinfection ^b									
Yes	0.7 (22)	33.9 (2.2)	2.28 (.71 to 3.85)	1.36 (−0.33 to 3.04)	1.6 (45)		34.4 (3.5)	1.27 (.26 to 2.28)	0.87 (−0.34 to 2.09)
No	22.3 (731)	31.6 (3.8)	Ref	Ref	10.8 (311)		33.1 (3.4)	Ref	Ref
Oropharyngeal NG coinfection ^c									
Yes	0.5 (18)	33.9 (3.4)	1.04 (−0.71 to 2.80)	0.31 (−1.57 to 2.19)	0.9 (27)		34.0 (3.0)	0.98 (−0.29 to 2.24)	0.01 (−1.45 to 1.48)
No	8.4 (275)	32.8 (3.8)	Ref	Ref	12.6 (364)		33.0 (3.4)	Ref	Ref
HIV coinfection ^{d,e}									
Yes	0.1 (3)	378 (0.25)	6.45 (2.26 to 10.64)	6.86 (2.72 to 10.99)	116.2	1.4 (41)	33.8 (3.1)	1.23 (.23 to 2.23)	0.95 (0.07 to 1.96)
No	41.0 (1389)	31.4 (3.8)	Ref	Ref	57.5 (1663)		32.6 (3.3)	Ref	Ref
TPHA/TPPA positive (syphilis screening test) ^f									
Yes	0.1 (4)	31.6 (15)	0.74 (−2.92 to 4.40)		0.7 (21)		33.3 (3.2)	0.48 (−0.92 to 1.88)	−0.19 (−1.62 to 1.24)
No	26.3 (860)	30.9 (3.9)	Ref		32.8 (949)		32.8 (3.3)	Ref	Ref

Statistically significant associations ($P < .05$) are depicted in bold.Abbreviations: CI, confidence interval; Cq, cycle quantification; CT, *Chlamydia trachomatis*; HIV, human immunodeficiency virus; NG, *Neisseria gonorrhoeae*; Ref, reference; TPHA, *Treponema pallidum* hemagglutination assay; TPPA, *Treponema pallidum* particle agglutination assay.^an = 70 patients with vaginal samples and n = 100 patients with urine samples were not tested for urogenital NG.^bn = 2520 patients with vaginal samples and n = 2535 patients with urine samples were not tested for anorectal *N. gonorrhoeae*.^cn = 2980 patients with vaginal samples and n = 2500 patients with urine samples were not tested for oropharyngeal *N. gonorrhoeae*.^dn = 1878 patients with vaginal samples and n = 1178 patients with urine samples were not tested for HIV.^eHIV result was missing for n = 3 patients with vaginal swabs and n = 9 patients with urine samples.^fn = 2409 patients with vaginal swabs and n = 1921 patients with urine samples were not tested for TPHA/TPPA (syphilis screening test).

women on extragenital locations, arguing for similar transmission potential, clinical relevance, and NAAT detection limits. Moreover, in women vaginal swabs had a higher *C. trachomatis* load compared to urine samples, confirming that vaginal swabs rather than urine samples should be used. A higher urogenital *C. trachomatis* load was observed in younger patients, whereas a lower urogenital *C. trachomatis* load was found in patients coinfecting with HIV and *N. gonorrhoeae*.

The strength of the current study is the comparison of *C. trachomatis* load of different sample types of both men and women in 1 large study using the same diagnostic test (NAAT), allowing for the best comparison possible. *C. trachomatis* load can be quite variable in different patients depending on various factors, including time of diagnosis since acquiring the infection. However, our study group showed that the majority of patients had a stable *C. trachomatis* load in the time interval between diagnosis and treatment (66.3%, 73.1%, and 48.6% for vaginal swabs, urine samples, and anorectal swabs, respectively) [26]. The *C. trachomatis* load for each sample type might also vary as patients could apply the sampling instructions differently. Nevertheless, mean *C. trachomatis* load at a population level was quite stable, as seen in our results of anorectal and oropharyngeal samples across both sexes [6]. In earlier work we discussed the limitations of comparing different sampling methods (ie, urine vs swabs) and different sampling locations (vaginal swab vs anorectal or oropharyngeal swabs). For example, different sites have different epithelial cells, which might influence the *C. trachomatis* load. Some previous studies included the number of epithelial cells in assessment of the *C. trachomatis* load count but as anatomical sample sites differ in the number of cells in the samples, we argue that this hampers accurate comparison [14]. The Cq values of urine samples of men were moderately, but significantly, correlated with urine *C. trachomatis* load (expressed as copies/mL). Therefore, the results of urine Cq values could be a less stable indication of *C. trachomatis* load. This is an expected finding as we know concentration in urine can differ inter- and intraindividually and with time because infected cells from the male urethra may get washed out by first void urine. Another reason could be that the *C. trachomatis* major outer membrane protein breaks down faster in urine samples than in vaginal swabs leading to higher Cq values in the in-house qPCR (Taqman) compared to lower Cq values in the NAAT (Cobas 4800). Furthermore, we were unable to assess whether the amplified DNA was from viable or nonviable *C. trachomatis*. Previously, our study group developed a new viability technique (viability PCR) to discriminate between viable and nonviable *C. trachomatis* and showed that a substantial amount of *C. trachomatis* DNA originated from nonviable cells [27]. The nonviable *C. trachomatis* DNA would have no impact on *C. trachomatis* transmission and sequelae.

There is an ongoing discussion about what difference in *C. trachomatis* load is clinically or microbiologically relevant.

Earlier, our group used a difference of 1 log load (3.3 Cq) as microbiology relevant to overcome potential technical variations when measuring the load within the same patient over time. However, in the current study, we averaged the Cq values over an entire group (eg, men, women, or patients aged <25 year). Therefore, potential technical differences between individuals were flattened out and had little impact on the results because these variations are randomly distributed over all samples tested within the particular group [26]. The precise clinical or microbiological relevance of the differences in *C. trachomatis* load between groups remains unknown. However, the fact that men and women had a similar extragenital *C. trachomatis* load indicates similar clinical relevance.

Nevertheless, of all sample types included, vaginal swabs had the highest *C. trachomatis* load, which could relate to the highest impact on transmission. Recently, our group showed a borderline significant association ($P = .54$) between a higher urogenital *C. trachomatis* load and microbiological *C. trachomatis* detection after treatment with azithromycin in women [28]. Our current study confirmed the presence of high urogenital *C. trachomatis* load in some patients, which might be clinically relevant when it hampers the treatment of a *C. trachomatis* infection with azithromycin.

Our research team and others have shown that the majority of women (summary estimate of 68.1%) diagnosed with urogenital *C. trachomatis* also have a concurrent anorectal *C. trachomatis* infection irrespective of reported anal sex or anal symptoms [29, 30]. The clinical relevance (risk of complications) and public health implications (transmission potential) associated with anorectal *C. trachomatis* detection in women are, however, under debate [31]. Anorectal infections in women could result from autoinoculation due to the close proximity between the vagina and anorectum [19, 32]. In our previous study, in a sample ($n = 105$) of concurrently infected STI clinic women showed that in the majority of cases (in 79% and 56% of women reporting anorectal intercourse and no anorectal intercourse, respectively), the anorectal *C. trachomatis* load was in the same range as the urogenital *C. trachomatis* load, suggesting similar clinical relevance [33]. Moreover, in the current study, we confirmed this finding and also showed that the anorectal *C. trachomatis* load in men equals that of women. We saw as many low as high anorectal loads in both men and women, which was suggested in an earlier smaller study based on *C. trachomatis* qPCR [12]. While diagnosis and treatment for anorectal *C. trachomatis* in MSM is recommended, whether this should apply in women is still under debate. However, the finding of a similar load in men and women suggests a similar clinical relevance. Relevance in women might be even more substantial as they bear the burden of reproductive morbidity caused by *C. trachomatis* infections [2]. Moreover, women concurrently infected with urogenital and anorectal *C. trachomatis* have an 8.5-times higher urogenital *C. trachomatis* load compared to anorectal-negative women [6]. Therefore,

women concurrently infected with urogenital and anorectal *C. trachomatis* might have higher chances of sequelae such as reproductive morbidity. Although, anorectal *C. trachomatis* infections in women might be coincidentally treated with a urogenital infections due to the high concurrency, a recent study from our group suggested that the frequently used drug azithromycin may be less effective than doxycycline in clearing anorectal *C. trachomatis* infections in women [28]. Nevertheless, the question remains, whether and how substantial autoinoculation or other mechanisms for transmission from the female anorectal site to the female urogenital site, and vice versa, occur.

Studies have shown a lower *C. trachomatis* load among older patients and patients with repeat infections potentially due to development of partial immunity against *C. trachomatis* [8, 9, 34, 35]. Especially in HIV-infected men (a high-risk group) who were repeatedly infected with *C. trachomatis*, development of partial immunity might explain a lower load.

In conclusion, the mean *C. trachomatis* load estimated by Cq values differed per sample type and thus per anatomic location. Vaginal swabs had the highest *C. trachomatis* load, whereas oropharyngeal swabs had the lowest load. Notably, men and women had a similar *C. trachomatis* load on extragenital locations, arguing for similar transmission potential and clinical relevance. A lower load was associated with older age (>25) and coinfections with *N. gonorrhoeae* and HIV, which was suggestive of exposure to previous *C. trachomatis* and potential development of some kind of partial immunity reducing *C. trachomatis* load.

Supplementary Data

Supplementary materials are available at *The Journal of Infectious Diseases* online. Consisting of data provided by the authors to benefit the reader, the posted materials are not copyedited and are the sole responsibility of the authors, so questions or comments should be addressed to the corresponding author.

Notes

Acknowledgments. The authors acknowledge the Medical Microbiology Laboratory of Maastricht University Medical Center (MUMC+) for providing the laboratory data, particularly Brian van der Veer, MUMC+ for providing the *Chlamydia trachomatis* cycle quantification data.

Potential conflicts of interest. All authors: No reported conflicts. All authors have submitted the ICMJE Form for Disclosure of Potential Conflicts of Interest. Conflicts that the editors consider relevant to the content of the manuscript have been disclosed.

References

1. Newman L, Rowley J, Vander Hoorn S, et al. Global estimates of the prevalence and incidence of four curable sexually transmitted infections in 2012 based on systematic review and global reporting. *PLoS One* **2015**; 10:e0143304.
2. Haggerty CL, Gottlieb SL, Taylor BD, Low N, Xu F, Ness RB. Risk of sequelae after *Chlamydia trachomatis* genital infection in women. *J Infect Dis* **2010**; 201(suppl 2):S134–55.
3. Vodstrcil LA, McIver R, Huston WM, Tabrizi SN, Timms P, Hocking JS. The epidemiology of *Chlamydia trachomatis* organism load during genital infection: a systematic review. *J Infect Dis* **2015**; 211:1628–45.
4. Rogers SM, Miller WC, Turner CF, et al. Concordance of *Chlamydia trachomatis* infections within sexual partnerships. *Sex Transm Infect* **2008**; 84:23–8.
5. Jalal H, Verlander NQ, Kumar N, Bentley N, Carne C, Sonnex C. Genital chlamydial infection: association between clinical features, organism genotype and load. *J Med Microbiol* **2011**; 60:881–8.
6. Wijers JNAP, Hoebe CJPA, van Liere GAFS, Wolffs PFG, Dukers-Muijers NHTM. *Chlamydia trachomatis* bacterial load, estimated by Cq values, in urogenital samples from men and women visiting the general practice, hospital or STI clinic. *PLoS One* **2019**; 14:e0215606.
7. Michel CE, Sonnex C, Carne CA, et al. *Chlamydia trachomatis* load at matched anatomic sites: implications for screening strategies. *J Clin Microbiol* **2007**; 45:1395–402.
8. Gomes JP, Borrego MJ, Atik B, et al. Correlating *Chlamydia trachomatis* infectious load with urogenital ecological success and disease pathogenesis. *Microbes Infect* **2006**; 8:16–26.
9. Wiggins R, Graf S, Low N, Horner PJ; Chlamydia Screening Studies (ClasS) Study Group. Real-time quantitative PCR to determine chlamydial load in men and women in a community setting. *J Clin Microbiol* **2009**; 47:1824–9.
10. Walker J, Fairley CK, Bradshaw CS, et al. The difference in determinants of *Chlamydia trachomatis* and *Mycoplasma genitalium* in a sample of young Australian women. *BMC Infect Dis* **2011**; 11:35.
11. Chan PA, Robinette A, Montgomery M, et al. Extragenital infections caused by *Chlamydia trachomatis* and *Neisseria gonorrhoeae*: a review of the literature. *Infect Dis Obstet Gynecol* **2016**; 2016:5758387.
12. van Liere GA, Dirks JA, Hoebe CJ, Wolffs PF, Dukers-Muijers NH. Anorectal *Chlamydia trachomatis* load is similar in men who have sex with men and women reporting anal sex. *PLoS One* **2015**; 10:e0134991.
13. van Rooijen MS, van der Loeff MF, Morré SA, van Dam AP, Speksnijder AG, de Vries HJ. Spontaneous pharyngeal *Chlamydia trachomatis* RNA clearance. A cross-sectional study followed by a cohort study of untreated STI clinic patients in Amsterdam, The Netherlands. *Sex Transm Infect* **2015**; 91:157–64.
14. Dirks JAMC, Hoebe CJPA, van Liere GAFS, Dukers-Muijers NHTM, Wolffs PFG. Standardisation is necessary in urogenital and extragenital *Chlamydia trachomatis* bacterial load determination by quantitative PCR: a review of

- literature and retrospective study. *Sex Transm Infect* **2019**; 95:562–8.
15. van Veen MG, Koedijk FD, van der Sande MA; Dutch STD Centres. STD coinfections in The Netherlands: specific sexual networks at highest risk. *Sex Transm Dis* **2010**; 37:416–22.
  16. Tongtoyai J, Todd CS, Chonwattana W, et al. Prevalence and correlates of *Chlamydia trachomatis* and *Neisseria gonorrhoeae* by anatomic site among urban Thai men who have sex with men. *Sex Transm Dis* **2015**; 42:440–9.
  17. Lim RB, Wong ML, Cook AR, et al. Determinants of chlamydia, gonorrhea, and coinfection in heterosexual adolescents attending the national public sexually transmitted infection clinic in Singapore. *Sex Transm Dis* **2015**; 42:450–6.
  18. Boisvert JE, Koutsky LA, Suchland RJ, Stamm WE. Clinical features of *Chlamydia trachomatis* rectal infection by serovar among homosexually active men. *Sex Transm Dis* **1999**; 26:392–8.
  19. Dubbink JH, de Waaij DJ, Bos M, et al. Microbiological characteristics of *Chlamydia trachomatis* and *Neisseria gonorrhoeae* infections in South African Women. *J Clin Microbiol* **2016**; 54:200–3.
  20. Barnes RC, Katz BP, Rolfs RT, Batteiger B, Caine V, Jones RB. Quantitative culture of endocervical *Chlamydia trachomatis*. *J Clin Microbiol* **1990**; 28:774–80.
  21. Dirks JA, Wolffs PF, Dukers-Muijrs NH, Brink AA, Speksnijder AG, Hoebe CJ. *Chlamydia trachomatis* load in population-based screening and STI-clinics: implications for screening policy. *PLoS One* **2015**; 10:e0121433.
  22. Bolotin S, Deeks SL, Marchand-Austin A, et al. Correlation of real time PCR cycle threshold cut-off with *Bordetella pertussis* clinical severity. *PLoS One* **2015**; 10:e0133209.
  23. Dionne LL, Raymond F, Corbeil J, Longtin J, Gervais P, Longtin Y. Correlation between *Clostridium difficile* bacterial load, commercial real-time PCR cycle thresholds, and results of diagnostic tests based on enzyme immunoassay and cell culture cytotoxicity assay. *J Clin Microbiol* **2013**; 51:3624–30.
  24. Ott SJ, Musfeldt M, Ullmann U, Hampe J, Schreiber S. Quantification of intestinal bacterial populations by real-time PCR with a universal primer set and minor groove binder probes: a global approach to the enteric flora. *J Clin Microbiol* **2004**; 42:2566–72.
  25. De Vries H, van Dam A, Bax C. Seksueel Overdraagbare Aandoeningen Multidisciplinaire Richtlijn. Update 2018. www.soa aids.nl/nl/professionals/beroepsgroep/arts/professioneel-handelen/de-specialist. Accessed 3 September 2018.
  26. Dirks JA, van Lier GA, Bogers S, Dukers-Muijrs NH, Wolffs PF, Hoebe CJ. Natural course of *Chlamydia trachomatis* bacterial load in the time interval between screening and treatment in anogenital samples. *PLoS One* **2015**; 10:e0145693.
  27. Janssen KJ, Hoebe CJ, Dukers-Muijrs NH, Eppings L, Lucchesi M, Wolffs PF. Viability-PCR shows that NAAT detects a high proportion of DNA from non-viable *Chlamydia trachomatis*. *PLoS One* **2016**; 11:e0165920.
  28. Dukers-Muijrs NHTM, Wolffs PFG, De Vries H, et al. Treatment effectiveness of azithromycin and doxycycline in uncomplicated rectal and vaginal *Chlamydia trachomatis* infections in women: a multicenter observational study (FemCure). *Clin Infect Dis* **2019**; 69:1946–54.
  29. van Lier GAFS, Dukers-Muijrs NHTM, Levels L, Hoebe CJPA. High proportion of anorectal *Chlamydia trachomatis* and *Neisseria gonorrhoeae* after routine universal urogenital and anorectal screening in women visiting the sexually transmitted infection clinic. *Clin Infect Dis* **2017**; 64:1705–10.
  30. Chandra NL, Broad C, Folkard K, et al. Detection of *Chlamydia trachomatis* in rectal specimens in women and its association with anal intercourse: a systematic review and meta-analysis. *Sex Transm Infect* **2018**; 94:320–6.
  31. Ding A, Challenor R. Rectal *Chlamydia* in heterosexual women: more questions than answers. *Int J STD AIDS* **2014**; 25:587–92.
  32. Heijne JCM, van Lier GAFS, Hoebe CJPA, Bogaards JA, van Benthem BHB, Dukers-Muijrs NHTM. What explains anorectal *Chlamydia* infection in women? Implications of a mathematical model for test and treatment strategies. *Sex Transm Infect* **2017**; 93:270–5.
  33. Dirks JAMC, van Lier GAFS, Hoebe CJPA, Wolffs P, Dukers-Muijrs NHTM. Genital and anal *Chlamydia trachomatis* bacterial load in concurrently infected women: a cross-sectional study. *Sex Transm Infect* **2019**; 95:317–21.
  34. Gupta K, Bakshi RK, Van Der Pol B, et al. Repeated *Chlamydia trachomatis* infections are associated with lower bacterial loads [published online ahead of print 4 October 2018]. *Epidemiol Infect* doi: 10.1017/S0950268818002704.
  35. Walker J, Tabrizi SN, Fairley CK, et al. *Chlamydia trachomatis* incidence and re-infection among young women—behavioural and microbiological characteristics. *PLoS One* **2012**; 7:e37778.