Re

Citation for published version (APA):

Wesselius, A., & Zeegers, M. (2017). Re: Lifestyle and bladder cancer prevention: no consistent evidence from cohort studies. *European Journal of Epidemiology*, 32(11), 1037-1038.
<https://doi.org/10.1007/s10654-017-0307-2>

Document status and date:

Published: 01/11/2017

DOI:

[10.1007/s10654-017-0307-2](https://doi.org/10.1007/s10654-017-0307-2)

Document Version:

Publisher's PDF, also known as Version of record

Document license:

Taverne

Please check the document version of this publication:

- A submitted manuscript is the version of the article upon submission and before peer-review. There can be important differences between the submitted version and the official published version of record. People interested in the research are advised to contact the author for the final version of the publication, or visit the DOI to the publisher's website.
- The final author version and the galley proof are versions of the publication after peer review.
- The final published version features the final layout of the paper including the volume, issue and page numbers.

[Link to publication](#)

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal.

If the publication is distributed under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license above, please follow below link for the End User Agreement:

www.umlib.nl/taverne-license

Take down policy

If you believe that this document breaches copyright please contact us at:

repository@maastrichtuniversity.nl

providing details and we will investigate your claim.

CORRESPONDENCE

Re: Lifestyle and bladder cancer prevention: no consistent evidence from cohort studies

Anke Wesselius¹ · Maurice Zeegers¹

Received: 29 August 2017 / Accepted: 30 August 2017 / Published online: 19 September 2017
© Springer Science+Business Media B.V. 2017

We thank Dr. Vrieling for her knowledgeable comments on our study [1]. Her concern was the use of estimates based on case–control and cohort studies combined.

In our systematic review of meta-analyses we choose to review the most comprehensive meta-analyses for all modifiable risk factors for bladder cancer, published up to September 2015. In our criteria list we prioritized adjustment for confounding, including adjustment for smoking, over study design.

Both case–control and cohort studies are susceptible to confounding bias, which can influence the magnitude of the relationship between the independent variable and outcome drastically. Therefore, unadjusted estimates are often considered incorrect.

Regarding the modifiable risk factor “physical activity” this criterion has led to report an estimate based on case–control and cohort studies separately, rather than cohort studies only. Although Keimling et al. [2], did report an estimate for physical activity based on cohort studies only, this estimate was not restricted to primary studies that adjusted for important confounders, e.g. four cohort studies were not adjusted for the most important risk factor for bladder cancer; smoking [3–6]. We, therefore, believe that our reported “more adjusted estimated” risk estimate, including both case–control and cohort studies, is more

correct than reporting the risk estimate based on cohort studies only.

In nutritional epidemiology, most individual studies have insufficient sample sizes and thus lacked adequate statistical power to find all true associations. Therefore, meta-analyses, in which results of independent studies are integrated into a more precise estimate, play an important role in this area of research. However, the examination of variability or heterogeneity in methodological diversity is crucial to this approach. In our review, the reported significant risk estimates for vegetables, vitamin E, vitamin C, antioxidant supplementation, obesity and folate showed low levels of heterogeneity. Although low levels of heterogeneity do not appear to be predictive of the accuracy of the meta-analysis result, they do suggest low methodological diversity. The role of recall bias is therefore likely to be present but with minimal influence on the results.

Regarding the remaining reported risk estimates that showed a significant association (i.e. processed meat, selenium, cruciferous vegetables, fruit and citrus) we do agree with Vrieling that results should be interpreted with caution, since methodological diversity among the included individual studies may play a significant role. However, for consistent reporting, we opted to use similar criteria/use a similar approach for all investigated modifiable risk factors. In addition, although the form of bias addressed by Vrieling (i.e. recall bias) has also been addressed and analyzed for its consequences in many epidemiological/methodological papers [7–10], no clear answer on the magnitude of the effect of this specific type of bias could be drawn. Furthermore, the magnitude and direction of recall bias might differ for different exposure factors, and concern are especially expected for those exposures that attracted public attention, since the knowledge of an

✉ Anke Wesselius
anke.wesselius@maastrichtuniversity.nl
Maurice Zeegers
m.zeegers@maastrichtuniversity.nl

¹ Department of Complex Genetics, School of Nutrition, and Translational Research in Metabolism (NUTRIM), Maastricht University Medical Centre, P. O. Box 616, 6200 MS Maastricht, The Netherlands

unhealthy or healthy effect of such an exposure is a prerequisite for study subjects to change their reporting behavior of individual exposure. In this regard, it is expected that especially fruit and vegetables intake would be effected by recall bias. However, the positive effect of fruit remains significant even after restricting to cohort studies only. Apart from methodological considerations and consistent real-life observational data, it may be worth pointing that the results on processed meat, cruciferous vegetables and (citrus)fruit are supported by strong biological evidence, which are reported in our discussion section.

References

1. AL-Zalabani AH, Stewart KF, Wesselius A, Schols AM, Zeegers MP. Modifiable risk factors for prevention of bladder cancer: a systematic review of meta-analyses. *Eur J Epidemiol.* 2016;31(9): 811–51
2. Keimling M, Behrens G, Schmid D, Jochem C, Leitzmann MF. The association between physical activity and bladder cancer: systematic review and meta-analysis. *Br J Cancer.* 2014;110:1862–70.
3. Paffenbarger RS Jr, Lee IM, Wing AL. The influence of physical activity on the incidence of site-specific cancers in college alumni. *Adv Exp Med Biol.* 1992;322:7–15.
4. Soll-Johanning H, Bach E. Occupational exposure to air pollution and cancer risk among Danish urban mail carriers. *Int Arch Occup Environ Health.* 2004;77:351–6.
5. Sormunen J, Backmand HM, Sarna S, Kujala UM, Kaprio J, Dyba T, Pukkala E. Lifetime physical activity and cancer incidence—A cohort study of male former elite athletes in Finland. *J Sci Med Sport.* 2013;S1440–2440:00470–2.
6. Wilson RT, Donahue M, Gridley G, Adami J, El Ghormli L, Dosemeci M. Shared occupational risks for transitional cell cancer of the bladder and renal pelvis among men and women in Sweden. *Am J Ind Med.* 2008;51:83–99.
7. Barry D. Differential recall bias and spurious associations in case/control studies. *Stat Med.* 1996;15(23):2603–16.
8. Barry D, Livingstone V. The investigation and correction of recall bias for an ordinal response in a case-control study. *Stat Med.* 2006;25(6):965–75.
9. Chouinard E, Walter S. Recall bias in case-control studies: an empirical analysis and theoretical framework. *J Clin Epidemiol.* 1995;48(2):245–54.
10. Gefeller O. Invited commentary: recall bias in melanoma—much ado about almost nothing? *Am J Epidemiol.* 2009;169:267–70.