Intimal hyperplasia in human vascular grafts : a study in peripheral bypasses and arteriovenous fistulas

Citation for published version (APA):

Hofstra, L. (1995). *Intimal hyperplasia in human vascular grafts : a study in peripheral bypasses and arteriovenous fistulas*. [Doctoral Thesis, Maastricht University]. Datawyse / Universitaire Pers Maastricht. <https://doi.org/10.26481/dis.19950602lh>

Document status and date:

Published: 01/01/1995

DOI:

[10.26481/dis.19950602lh](https://doi.org/10.26481/dis.19950602lh)

Document Version:

Publisher's PDF, also known as Version of record

Please check the document version of this publication:

- A submitted manuscript is the version of the article upon submission and before peer-review. There can be important differences between the submitted version and the official published version of record. People interested in the research are advised to contact the author for the final version of the publication, or visit the DOI to the publisher's website.
- The final author version and the galley proof are versions of the publication after peer review.
- The final published version features the final layout of the paper including the volume, issue and page numbers.

[Link to publication](#)

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal.

If the publication is distributed under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license above, please follow below link for the End User Agreement:

www.umlib.nl/taverne-license

Take down policy

If you believe that this document breaches copyright please contact us at:

repository@maastrichtuniversity.nl

providing details and we will investigate your claim.

Download date: 26 Apr. 2023

INTIMAL HYPERPLASIA IN HUMAN VASCULAR GRAFTS
A STUDY IN PERIPHERAL BYPASSES AND ARTERIOVENOUS FISTULAS

Supported by a grant of the Netherlands Heart Foundation (NHS 90.055)

INTIMAL HYPERPLASIA IN HUMAN VASCULAR GRAFTS

A study in peripheral bypasses and arteriovenous fistulas

PROEFSCHRIFT

ter verkrijging van de graad van doctor
aan de Rijksuniversiteit Limburg te Maastricht,
op het gezag van de Rector Magnificus, Prof. mr. M.J. Cohen,
volgens het besluit van het College van Dekanen,
in het openbaar te verdedigen
op vrijdag 2 juni 1995 om 14.00 uur

door

Leonard Hofstra

geboren te Emmen op 25 juni 1963

UNIVERSITAIRE PERS MAASTRICHT

Promotores

Prof. dr. P.J.E.H.M. Kitslaar

Prof. dr. A.P.G. Hoeks

Co-promotores

Dr. J.H.M. Tordoir

Dr. M.J.A.P. Daemen

Beoordelingscommissie

Prof. dr. R.S. Reneman (voorzitter)

Prof. dr. J.W. Arends

Prof. dr. B. Eykelboom (Rijksuniversiteit Utrecht)

Prof. dr. N. Westerhof (Vrije Universiteit Amsterdam)

Prof. dr. H.J.J. Wellens

CIP-DATA KONINKLIJKE BIBLIOTHEEK, DEN HAAG

Hofstra, Leonard

Intimal hyperplasia in human vascular grafts; a study in
peripheral bypasses and arteriovenous fistulas / Leonard
Hofstra. - Maastricht: Universitaire Pers Maastricht. - Ill.
Thesis Rijksuniversiteit Limburg Maastricht. - With ref. -
With summary in Dutch.

ISBN 90-5278-190-7

Subject headings: intimal hyperplasia / human vascular grafts / fistulas.

Layout: Karin Scheele, A Point, Maastricht

Production: Datawyse, Maastricht

Financial support by the Netherlands Heart Foundation and the Netherlands Kidney Foundation for the publication of this thesis is gratefully acknowledged.

CONTENTS

	page
Preface	7
INTRODUCTION	
Chapter 1	General introduction 9
Chapter 2	The development of intimal hyperplasia after vascular reconstruction in humans 19
MATERIALS AND METHODS	
Chapter 3	Vessel Wall Doppler Tracking: A new technique to assess vessel wall elastic properties 31
Chapter 4	Short-term variations in elastic properties of a muscular artery in humans 41
ELASTIC PROFILES OF PERIPHERAL BYPASSES AND GRAFT ARTERIO- VENOUS FISTULAS IN HUMANS	
Chapter 5	Assessment of inhomogeneities in elastic properties of in situ and reversed saphenous vein grafts in humans 57
Chapter 6	Elastic properties of PTFE and saphenous vein bypass grafts in humans 71
Chapter 7	Mismatch in elastic properties around anastomoses of interposition grafts for hemodialysis access 85
Chapter 8	The dimensions and elastic properties of human saphenous vein grafted into the arterial circulation are stable with time 101

MECHANICAL INFLUENCES ON THE OCCURRENCE OF INTIMAL HYPERPLASIA IN PERIPHERAL BYPASSES AND GRAFT ARTERIO-VEINOUS FISTULAS IN HUMANS

Chapter 9	Anastomotic intimal hyperplasia in prosthetic arterio-venous fistulas for hemodialysis is associated with high shear rate and not with mismatch in elastic properties	115
Chapter 10	The influence of mismatch in elastic properties and flow on the occurrence of stenotic intimal thickening in peripheral bypass grafts in humans; a prospective analysis	131
Chapter 11	High shear rate mediated denudation inhibits proliferation in human stenotic tissue derived from AV fistulas and peripheral bypass grafts	145

GENERAL DISCUSSION

Chapter 12	Mechanical and flow characteristics affecting intimal hyperplasia in peripheral bypasses and graft arteriovenous fistulas in humans	165
Summary		173
Samenvatting		177
Dankwoord		181
Curriculum vitae		183
List of publications		185

PREFACE

The studies presented in this thesis are part of a larger research program, conducted in the Cardiovascular Research Institute Maastricht (CARIM). In that program attention is focused on studying the elastic properties of medium sized arteries in patients with a high risk of developing atherosclerosis, like hypertensive and diabetic patients, to gain more insight in the pathogenesis of vascular disease. Vessel Wall Doppler Tracking, a non-invasive method to assess the elastic properties of large vessels, is one of the techniques used and has been developed in the Department of Biophysics of our University.

We have used this system to investigate a different vascular problem, that is the occurrence of stenoses in peripheral bypass grafts and arterio-venous fistulas used for hemodialysis. Patency and function of these vascular conduits are jeopardized by the development of a stenosis due to intimal hyperplasia, mainly occurring at the anastomoses. A difference in elastic properties between graft and native vessels ('compliance mismatch') and unphysiologic flow conditions (shear rate) are possible contributing factors in the development of the stenotic lesions. In this thesis we have attempted to clarify the role of mismatch in elastic properties around anastomoses in the development of intimal hyperplasia in peripheral bypasses and graft arteriovenous fistulas in humans in a prospective way. Furthermore, the influence of flow characteristics on intimal thickening was studied.

When we started our studies in 1991, quantitative mechanical information on bypass grafts and AV fistulas was scarce. Therefore, we first studied the mechanical profiles of different graft materials in both types of reconstructions. The elastic properties and flow characteristics were assessed two weeks after construction at defined sites in the graft and in the inflow and outflow vascular segments in patients who underwent femorodistal bypass surgery for stage II-IV occlusive arterial disease of the legs and in patients with a graft arteriovenous fistula for hemodialysis access. The initial mechanical and flow profiles were correlated with the occurrence of intimal hyperplasia during follow-up. This allowed a prospective analysis of the influence of mismatch in elastic properties and flow characteristics on intimal hyperplasia. Finally, the extent and distribution of cellular proliferation in stenotic lesions, surgically derived from these patients was studied in relation to local flow characteristics, in order to obtain insight in the direct coupling between flow and intimal proliferation.

Research on this subject has recently been continued, by investigating the velocity profiles in arteries with a new ultrasound technique. For the first time, in vivo shear rate measurements in humans can be performed, allowing investigation of the effect of shear rate on the development of intimal hyperplasia in humans. Investigations with this new system may further unravel the influence of mechanical factors on the development of intimal hyperplastic lesions in vascular reconstructions in humans.

CHAPTER 1

GENERAL INTRODUCTION

The treatment of lower limb ischaemia is one of the major clinical problems encountered in vascular surgery. After the first implantation of a saphenous vein graft as a bypass to reestablish distal perfusion of the lower limb in 1949, major technical progress has been made in bypassing peripheral occlusive atherosclerotic lesions¹. Nowadays, femoro-pedal bypasses are performed with the use of the in situ technique, and femoro-crural bypasses are constructed using prosthetic graft material, to achieve limb salvage in patients previously considered inoperable²⁻¹⁸. Despite these technical improvements¹⁹⁻²¹, peripheral bypasses are often complicated by the development of stenoses, mainly due to intimal hyperplasia. When a saphenous vein is grafted into the arterial circulation, generalized intimal thickening occurs, but to a limited extent²²⁻²⁹. However, sometimes, locally, the process of intimal thickening is uncontrolled and leads to the formation of focal stenotic lesions^{16,17,30-34}. Intimal hyperplasia often occurs at the anastomoses or in the areas adjacent to the anastomoses³³⁻³⁹ and leads to deterioration of graft flow and eventually to occlusion in about 20-30% of cases in the first year after operation^{2,3,16,17,36,37,40}. Graft occlusion often necessitates surgical repair of the bypass graft and may result in loss of the graft. Another type of vascular reconstruction, the arteriovenous fistula for hemodialysis⁴¹, is even more susceptible to develop intimal hyperplasia, with a 50-60% incidence of stenosis in the first postoperative year⁴²⁻⁴⁵. Since hemodialysis patients depend on a proper functioning of the fistula, hemodynamically important stenoses always necessitate surgical correction or transluminal angioplasty. These interventions result in a mean hospital stay of one month per patient each year, illustrating the financial consequences and the impact of this clinical problem on the daily life of hemodialysis patients^{42,45}. Another major clinical problem related to intimal hyperplasia is restenosis after percutaneous transluminal coronary angioplasty (PTCA) in patients with severe coronary artery disease. Restenosis due to intimal thickening occurs within three months in approximately 30-40% of patients after PTCA, results in recurrence of symptoms, and often necessitates reintervention⁴⁶⁻⁵⁴.

Because of the profound clinical impact of intimal hyperplasia after vascular reconstruction, extensive research has been performed in the last two decades to increase our understanding of the biology of this vascular response. Immunologic reactions⁵⁵⁻⁶⁹, platelet aggregation after vascular injury^{59,60,70-72}, a large variety of growth factors^{30,51} and different mechanical factors^{73,74} have been implicated as playing a role in the development of stenoses after vascular reconstruction.

The studies presented in this thesis concentrate on the role of different mechanical factors, such as compliance mismatch and unphysiological shear rate, on the occurrence of intimal hyperplasia. Our hypothesis that compliance mismatch between graft and native vessels is one of the factors contributing to the development of intimal hyperplasia is based on the observation, that intimal thickening very often develops at the site of the anastomoses^{34,35,38,42,45,75-77}. A possible mechanism may be a difference in compliance between graft and host vessel, occurring within a very short vessel segment, resulting in excessive mechanical stretching at the site of the anastomosis. Mechanical stretch may induce vessel wall injury, initiating a reparative response which results in intimal thickening^{73,78}. Another physical factor affecting intimal growth is shear stress, i.e. shear rate times local viscosity. In one of the first studies evaluating the effect of varying levels of shear rate on the vessel wall, endothelial cell denudation was observed when the vessel wall was exposed to excessive high shear rate levels^{79,80}. Therefore, it was suggested that high shear rate induces a hemodynamic injury of the vessel wall, initiating a vascular response to injury. This hypothesis was supported by observations in an animal study, investigating the influence of varying flow velocities on the occurrence of intimal hyperplasia at the venous anastomosis and the efferent vein of graft AV fistulas. A higher incidence of intimal hyperplasia was observed in areas with the highest flow velocities^{81,82}. On the other hand, numerous investigations in different animal models demonstrated increased intimal thickening in low shear rate regions^{74,83-89}. Therefore, the current opinion is that the development of intimal thickening occurs in areas with low shear rate conditions.

AIM OF THE STUDY

In this thesis we have attempted to investigate prospectively the influence of mechanical factors, mismatch in elastic properties and shear rate, on the occurrence of intimal hyperplasia in peripheral bypass grafts and graft AV fistulas. One of the main advantages of studying the influence of mechanical factors in these 'models' is that peripheral bypass grafts and AV fistulas are easily accessible to the techniques used in this thesis. Moreover, the inclusion of the two different types of vascular reconstruction allows the investigation of mechanical factors on intimal hyperplasia in conditions of normal physiologic arterial flow (peripheral bypass) and very high flow (arteriovenous fistula). Our hypothesis is that (sub)-occlusive intimal hyperplasia in peripheral bypasses or AV fistulas occurs in areas of low shear rates. In addition, we hypothesize that an increased mismatch in elastic properties between graft and native vessels is associated with the occurrence of peri-anastomotic intimal hyperplasia. To test these hypotheses, a stepwise approach is used, which is reflected by the structure of the thesis.

OUTLINE OF THE THESIS

After an introduction of the problem of intimal hyperplasia following vascular reconstructions in humans (chapter 2), the thesis is divided in three different sections:

1. MATERIALS AND METHODS

In this section Vessel Wall Doppler Tracking (VWDT), the method used to measure diameter and distension of vessel segments is described and evaluated (chapter 3). Furthermore, the results of repeated measurements of the elastic properties of the common femoral artery in humans are reported (chapter 4). Especially, intra-individual variation in elastic behavior in relation to blood pressure was evaluated.

2. ELASTIC PROFILES OF PERIPHERAL BYPASSES AND GRAFT AV FISTULAS IN HUMANS

Since quantitative information on the in vivo mechanical profiles of peripheral bypass grafts and AV fistulas in humans was not available, we started to measure the elastic properties of different graft materials and the adjacent native vessels. The goal of these studies was to quantify mismatch in elastic properties around anastomoses and to obtain mechanical profiles along the graft. In addition, we tried to identify clues for the predisposition of intimal hyperplasia to develop at the site of the anastomoses. In chapter 5 the assessment of mechanical profiles of 'in situ' and 'reversed' saphenous vein bypass grafts are reported. The goal of this study was to compare the mechanical profiles of the in situ and the reversed saphenous vein peripheral bypass graft and to obtain insight in possible inhomogeneities in elastic properties around anastomoses. In chapter 6 a comparison is made between elastic profiles of poly-tetra-fluoro-ethylene (PTFE) prosthesis and reversed saphenous vein bypass grafts in the femorodistal position. In chapter 7 the differences in mismatch in elastic properties around the venous and arterial anastomoses of graft arteriovenous fistulas for hemodialysis are reported. Chapter 8 describes the dimensions and elastic properties of saphenous vein femorodistal grafts during the first 6 months after implantation. The goal of this study was to investigate possible functional or structural adaptation with time of the saphenous vein when grafted into the arterial circulation.

3. MECHANICAL INFLUENCES ON THE DEVELOPMENT OF INTIMAL HYPERPLASIA IN PERIPHERAL BYPASSES AND GRAFT AV FISTULAS IN HUMANS

The occurrence of stenoses due to intimal hyperplasia in the peripheral bypass grafts and the arteriovenous fistulas, respectively, was correlated with the initial mechanical profiles and flow characteristics as reported in chapter 5-7. The goal of these studies was to prospectively analyse the influence of mechanical factors, flow and perianastomotic mismatch in elastic properties, on the occurrence of intimal thickening.

The results of these studies are reported in chapter 9 (AV fistulas) and 10 (peripheral bypasses). In chapter 11 the assessment of intimal DNA synthesis in surgically harvested stenotic lesions in relation to flow and endothelial cell coverage of the lumen is reported. The goal of that study was to investigate a possible link between peak shear rate and cell proliferation in human stenotic lesions. In addition, the influence of the presence of endothelial cells on intimal proliferation was studied.

REFERENCES

1. Kunlin J. Le traitement de l'arterite oblitterante par la greffe veineuse. *Arch Chir Mal Coeur* 1949; 42:371-372.
2. Harris PL, How TV, Jones DR. Prospectively randomized clinical trial to compare in situ and reversed saphenous vein grafts for femoropopliteal bypass. *Br J Surg* 1987; 74:252-255.
3. Harris RW, Andros G, Dulawa LB, Oblath RW, Apyan R, Salles-Cunha S. The transition to "in situ" vein bypass grafts. *Surg Gynecol Obstet* 1986; 163:21-28.
4. Harris PL, Veith FJ, Shanik GD, Nott D, Wengerter KR, Moore DJ. Prospective randomized comparison of in situ and reversed infrapopliteal vein grafts. *Br J Surg* 1993; 80:173-176.
5. Acher CW, Turnipseed WD. In situ distal saphenous vein bypass using the intraluminal valve-disruption technique. *Arch Surg* 1985; 120:933-936.
6. Baird RJ, Tutassaura H, Hiyagishima RT. Saphenous vein bypass graft to arteries of the ankle and foot. *Ann Surg* 1970; 172:1059-1063.
7. Batson RC, Sottiurai VS. Nonreversed and in situ grafts. Clinical and experimental observations. *Ann Surg* 1985; 201:771-779.
8. Buchbinder D, Rolins DL, Verta MJ, et al. Early experience with in situ saphenous vein bypass for distal arterial reconstruction. *Surgery* 1986; 99:350-356.
9. Buchbinder D, Singh JK, Karmody AM, Leather RP, Shah DM. Comparison of patency rate and structural changes of in situ and reversed vein arterial bypass. *J Surg Res* 1981; 30:213-222.
10. Hurley JJ, Auer AI, Binnington HB, et al. Comparison of initial limb salvage in 98 consecutive patients with either reversed autogenous or in situ vein bypass graft procedures. *Am J Surg* 1985; 150:777-781.
11. Leather RP, Shah DM, Chang BB, Kaufman JL. Resurrection of the in situ saphenous vein bypass. 1000 cases later. *Ann Surg* 1988; 208:435-442.
12. Leather RP, Shah DM, Karmody AM. Infrapopliteal arterial bypass for limb salvage: Increased patency and utilization of the saphenous vein used "in situ". *Surgery* 1981; 90:1000-1008.
13. Najmaldin A, Clifford PC, Farrands PA, Campbell MJ, Chant AD, Webster JH. The success of in situ vein grafting: Early results in comparison with the reversed vein technique. *Eur J Vasc Surg* 1987; 1:165-168.
14. Prendiville EJ, Yeager A, O'Donnell TF, et al. Long-term results with the above-knee popliteal expanded polytetrafluoroethylene graft. *J Vasc Surg* 1990; 11:517-524.
15. Shah DM, Buchbinder D. Modified technique to produce valvular incompetence in in situ saphenous vein arterial bypass. *Arch Surg* 1981; 116:356-358.
16. Taylor LM, Edwards JM, Phinney ES, Porter JM. Reversed vein bypass to infrapopliteal arteries. *Ann Surg* 1987; 205:90-97.
17. Taylor LM, Edwards JM, Porter JM. Present status of reversed vein bypass grafting: Five-year results of a modern series. *J Vasc Surg* 1990; 11:193-206.
18. Taylor RS, Loh A, McFarland RJ, Cox M, Chester JF. Improved technique for polytetrafluoroethylene bypass grafting: long-term results using anastomotic vein patches. *Br J Surg* 1992; 79:348-354.
19. Tyrrell MR, Chester JF, Vipond MN, Clarke GH, Taylor RS, Wolfe JHN. Experimental evidence to support the use of interposition vein collars/patches in distal PTFE anastomoses. *Eur J Vasc Surg* 1990; 4:95-101.
20. Tyrrell MR, Wolfe JHN. New prosthetic venous collar anastomotic technique: combining the best of other procedures. *Br J Surg* 1991; 78:1016-1017.
21. Veith FJ, Gupta SK, Ascer E, et al. Six year prospective multicenter randomized comparison of autologous saphenous vein and polytetrafluoroethylene grafts in infrainguinal arterial reconstructions. *J Vasc Surg* 1986; 3:104-114.

22. Cambria RP, Brewster DC, Hasson J, Megerman J, Warnock DF, Abbott WM. The evolution of morphologic and biomechanical changes in reversed and in situ vein grafts. *Ann Surg* 1987; 205:167-174.
23. Dilley RJ, McGeachie JK, Prendergast FJ. A review of the histologic changes in vein-to-artery grafts, with particular reference to intimal hyperplasia. *Arch Surg* 1988; 123:691-696.
24. Fuchs JCA, Mitchener JS, Hagen PO. Postoperative changes in autologous vein grafts. *Ann Surg* 1978; 188:1-15.
25. Dilley RJ, McGeachie JK, Tennant M. Vein to artery grafts: a morphological and histochemical study of the histogenesis of intimal hyperplasia. *Aust N Z J Surg* 1992; 62:297-303.
26. Adcock GD, Adcock OT, Wheeler JR, et al. Arterialization of reversed autogenous vein grafts: Quantitative light and electron microscopy of canine jugular vein grafts harvested and implanted by standard or improved techniques. *J Vasc Surg* 1987; 6:283-295.
27. Dobrin PB, Littoo FN, Golan J, Blakeman B, Fareed J. Mechanical and histological changes in canine vein grafts. *J Surg Res* 1988; 44:259-265.
28. Milroy CM, Scott DJA, Beardt JD, Horrocks M, Bradfield JWB. Histological appearances of the long saphenous vein. *J Pathology* 1989; 159:311-316.
29. Zacharias RK, Kirkman TR, Clowes AW. Mechanisms of healing in synthetic grafts. *J Vasc Surg* 1987; 6:429-436.
30. Clowes AW, Reidy MA. Prevention of stenosis after vascular reconstruction: Pharmacologic control of intimal hyperplasia - A review. *J Vasc Surg* 1991; 6:885-891.
31. Szilagy DE, Elliot JP, Hageman JH, Smith RF, Daall'olmo CA. Biologic fate of autogenous vein implants as arterial substitutes. *Ann Surg* 1973; 178:232-246.
32. Echave V, Koornick AR, Haimov M, Jacobson JH. Intimal hyperplasia as a complication of the use of the polytetrafluoroethylene graft for femoral-popliteal bypass. *Surgery* 1979; 86:791-798.
33. Madras PN, Johnson WR, Ward CA. Enhancement of thrombus formation by vascular prostheses. *Can J Surg* 1980; 23:544-546.
34. Madras PN, Ward CA, Johnson WR, Singh PI. Anastomic hyperplasia. *Surgery* 1981; 90:922-923.
35. Imparato AM, Bracco A, Eun Kim G, Zeff R. Intimal and neointimal fibrous proliferation causing failure of arterial reconstructions. *Surgery* 1972; 72:1007-1017.
36. Berkowitz HD, Fox AD, Deaton DH. Reversed vein graft stenosis: Early diagnosis and management. *J Vasc Surg* 1992; 15:130-142.
37. Berkowitz HD, Greenstein S, Barker CF, Perloff LJ. Late failure of reversed vein bypass grafts. *Ann Surg* 1989; 210:782-786.
38. LoGerfo FW, Quist WC, Nowak MD, Crawshaw HM, Haudenschild CC. Downstream anastomotic hyperplasia. A mechanism of failure in Dacron arterial grafts. *Ann Surg* 1983; 197:479-483.
39. Mills JL, Harris EJ, Taylor LM, Beckett WC, Porter JM. The importance of routine surveillance of distal bypass grafts with duplex scanning: A study of 379 reversed vein grafts. *J Vasc Surg* 1990; 12:379-389.
40. Buth J, Disselhoff B, Sommeling C, Stam L. Color-flow duplex criteria for grading stenosis in infrainguinal vein grafts. *J Vasc Surg* 1991; 14:716-728.
41. May J, Tiller D, Johnson J, Stewart J, Ross Sheil AG. Saphenous-vein arteriovenous fistula in regular dialysis treatment. *New Eng J Med* 1969; 280:770.
42. Swedberg SH, Brown BG, Sigley R, Wight TN, Gordon D, Nicholls SC. Intimal fibromuscular hyperplasia at the venous anastomosis of PTFE grafts in hemodialysis patients. *Circulation* 1989; 80:1726-1736.

43. Tordoir JHM, Herman JM, Kwan TS, Diderich PM. Long-term follow-up of the polytetrafluoroethylene (PTFE) prosthesis as an arteriovenous fistula for hemodialysis. *Eur J Vasc Surg* 1988; 2:3-7.
44. Windus DW. Permanent vascular access: a nephrologist's view. *Am J Kidney Dis* 1993; 21:457-471.
45. Jenkins AMcL, Buist TAS, Glover SD. Medium-term follow-up of forty autogenous vein and forty polytetrafluoroethylene (Gore-Tex) grafts for vascular access. *Surgery* 1980; 88:667-672.
46. Bertrand ME, Lablanche JM, Fourrier JL, Gommeaux A, Ruel M. Relation to restenosis after percutaneous transluminal coronary angioplasty to vasomotion of the dilated coronary arterial segment. *Am J Cardiol* 1989; 63:277-281.
47. Vlietstra RE, Holmes DR, Rodeheffer RJ, Bailly KR. Consequences of restenosis after coronary angioplasty. *Int J Cardiol* 1991; 31:143-148.
48. Waller BF, Pinkerton CA, Orr CM, Slack JD, VanTassel JW, Peters T. Restenosis 1 to 24 months after clinically successful coronary balloon angioplasty: A necropsy study of 20 patients. *J Am Coll Cardiol* 1991; 17:58b-70b.
49. Serruys PW, Foley DP, Kirkheide RL, King III SB. Restenosis revisited: Insights provided by quantitative coronary angiography. *Am Heart J* 1993; 126:1243-1267.
50. Serruys PW, Breeman A. Coronary angioplasty - Long-term follow-up results and detection of restenosis: guidelines for aviation cardiology. *Eur Heart J* 1992; 13:76h-88h.
51. Liu MW, Roubin GS, King III SB. Restenosis after coronary angioplasty. Potential biologic determinants and role of intimal hyperplasia. *Circulation* 1989; 79:1374-1387.
52. Liu MW, Berk BC. Restenosis following coronary balloon angioplasty. Role of smooth muscle cell proliferation. *Trends Cardiovasc Med* 1991; 1:107-111.
53. Serruys PW, Luyten HE, Beatt KJ, et al. Incidence of restenosis after successful coronary angioplasty: a time-related phenomenon. *Circulation* 1988; 77:361-371.
54. McBride W, Lange RA, Hillis LD. Restenosis after successful coronary angioplasty. Pathophysiology and prevention. *New Eng J Med* 1988; 318:1734-1737.
55. Bevilacqua MP, Nelson RM. Selectins. *J Clin Invest* 1993; 91:379-387.
56. Badimon JJ, Fuster V, Chesebro JH, Badimon L. Coronary atherosclerosis. A multifactorial disease. *Circulation* 1993; 87:II3-II16.
57. Farber HW, Beer DJ. Restricted secretion of a T-lymphocyte chemotactic cytokine by serotonin-stimulated cultured aortic endothelial cells. *Circ Res* 1991; 69:257-265.
58. Fitzmaurice M, Ratliff NB. Immunoglobulin deposition in atherosclerotic aortocoronary saphenous vein grafts. *Arch Pathol Lab Med* 1990; 114:388-393.
59. Fuster V, Badimon L, Badimon JJ, Chesebro JH. Mechanisms of disease: The pathogenesis of coronary artery disease and the acute coronary syndromes (First of two parts). *New Eng J Med* 1992; 326:242-250.
60. Fuster V, Badimon L, Badimon JJ, Chesebro JH. Mechanisms of disease: The pathogenesis of coronary artery disease and the acute coronary syndromes (Second of two parts). *New Eng J Med* 1992; 326:310-318.
61. Garratt KN, Edwards WD, Kaufmann URS, Vlietstra RE, Holmes DR. Differential histopathology of primary atherosclerotic and restenotic lesions in coronary arteries and saphenous vein bypass grafts: Analysis of tissue obtained from 73 patients by directional atherectomy. *J Am Coll Cardiol* 1991; 17:442-448.
62. Hansson GK, Holm J, Holm S, Fotev Z, Hedrich HJ, Fingerle J. T lymphocytes inhibit the vascular response to injury. *Proc Natl Acad Sci U S A* 1991; 88:10530-10534.
63. Mantovani A, Dejana E. Cytokines as communication signals between leukocytes and endothelial cells. *Immunol Today* 1989; 10:370-375.
64. McEver RP. Selectins: Novel receptors that mediate leukocyte adhesion during inflammation. *Thrombosis Haemostasis* 1991; 65:223-228.

65. Nelken NA, Coughlin SR, Gordon D, Wilcox JN. Monocyte chemoattractant protein-1 in human atheromatous plaques. *J Clin Invest* 1991; 88:1121-1127.
66. Pan XM, Nelken NA, Colyvas N, Rapp JH. Inhibition of injury induced intimal hyperplasia by saralasin in rats. *J Vasc Surg* 1992; 15:693-698.
67. Ricevuti G, Mazzone A, Pasotti D, de Servi S, Specchia G. Role of granulocytes in endothelial injury in coronary heart disease in humans. *Atherosclerosis* 1991; 91:1-14.
68. Chan P, Munro E, Patel M, et al. Cellular biology of human intimal hyperplastic stenosis. *Eur J Vasc Surg* 1993; 7:129-135.
69. Ross R. The pathogenesis of atherosclerosis: a perspective for the 1990s. *Nature* 1993; 362:801-809.
70. Okadome K, Yukizane T, Mii S, Fukuda A, Sugimachi K. Correlation of longterm results of femoropopliteal autogenous vein grafts with shear stress variation. In situ or reversed graft? *J Cardiovasc Surg Torino* 1989; 30:932-935.
71. Karino T, Goldsmith HL. Role of blood cell-wall interactions in thrombogenesis and atherogenesis: a microrheological study. *Biorheology* 1984; 21:587-601.
72. Yukizane T, Okadome K, Eguchi H, Muto Y, Sugimachi K. Isotopic study of the effects of platelets on development of intimal thickening in autologous vein grafts in dogs. *Br J Surg* 1991; 78:297-302.
73. Abbott WM, Megerman J, Hasson JE, L'Italien G, Warnock DF. Effect of compliance mismatch on vascular graft patency. *J Vasc Surg* 1987; 5:376-382.
74. Kohler TR, Jawien A. Flow affects development of intimal hyperplasia after arterial injury in rats. *Arterioscler Thromb* 1992; 12:963-971.
75. Baird RN, Abbott WM. Pulsatile blood-flow in arterial grafts. *Lancet* 1976; October 30:948-950.
76. Ark van der PD, Smakman JGJ. De rol van plaatjesaggregatie bij hart- en vaatziekten. Deel 3: Welke winst kan men verwachten van een plaatjesaggregatie-remmende therapie? *Janssen Nieuws* 1990; 5:47-50.
77. Mills JL, Fujitani RM, Taylor SM. The characteristics and anatomic distribution of lesions that cause reversed vein graft failure: a five-year prospective study. *J Vasc Surg* 1993; 17:195-204.
78. Abbott WM, Megerman J. Adaptive responses of arteries to grafting. *J Vasc Surg* 1989; 9:377-379.
79. Fry DL. Acute vascular endothelial changes are associated with increased blood velocity gradients. *Circ Res* 1968; 22:165-197.
80. Fry DL. Certain chemorheologic considerations regarding the blood vascular interface with particular reference to coronary artery disease. *Circulation* 1969; XXXIX and XI:IV38-IV59.
81. Fillinger MF, Reinitz ER, Schwartz RA, Resetarits DE, Paskanik AM, Bredenberg CE. Beneficial effects of banding on venous intimal-medial hyperplasia in arteriovenous loop grafts. *Am J Surg* 1989; 158:87-94.
82. Fillinger MF, Reinitz ER, Schwartz RA, et al. Graft geometry and venous intimal-medial hyperplasia in arteriovenous loop grafts. *J Vasc Surg* 1990; 11:556-566.
83. Kohler TR, Kirkman TR, Kraiss LW, Zierler BK, Clowes AW. Increased blood flow inhibits neointimal hyperplasia in endothelialized vascular grafts. *Circ Res* 1991; 69:1557-1565.
84. Asakura T, Karino T. Flow patterns and spatial distribution of atherosclerotic lesions in human coronary arteries. *Circ Res* 1990; 66:1045-1066.
85. Bassiouny HS, White S, Glagov S, Choi E, Giddens DP, Zarins CK. Anastomotic intimal hyperplasia: Mechanical injury or flow induced. *J Vasc Surg* 1992; 15:708-717.
86. Friedman MH. Some atherosclerosis may be a consequence of the normal adaptive response to shear. *Atherosclerosis* 1990; 82:193-196.
87. Friedman MH, Barger CB, Deters OJ, Hutchins GM, Mark FF. Correlation between wall shear and intimal thickness at a coronary artery branch. *Atherosclerosis* 1987; 68:27-33.

88. Kamiya A, Ando J, Shibata M, Masuda H. Roles of fluid shear stress in physiological regulation of vascular structure and function. *Biorheology* 1988; 25:271-277.
89. Kamiya A, Togawa T. Adaptive regulation of wall shear stress to flow change in the canine carotid artery. *Am J Physiol* 1980; 239:H14-H21.

CHAPTER 2

THE DEVELOPMENT OF INTIMAL HYPERPLASIA AFTER VASCULAR RECONSTRUCTION IN HUMANS

INTRODUCTION

Intimal hyperplasia has first been described in 1906 by Carrell¹, who transplanted a vein into the arterial circulation and observed intimal thickening at the site of the anastomoses. Nowadays, intimal hyperplasia is considered as the response of a vessel wall to injury. It is the main cause for obstructive complications in the first year following almost every type of vascular intervention. Stenoses due to intimal hyperplasia may develop after PTCA (30-40%)²⁻⁹, after coronary artery bypass grafting (25%)¹⁰, in peripheral artery bypass surgery (25%)¹¹⁻²⁶ and after the construction of a vascular access for hemodialysis (50%)²⁷⁻³¹. Knowledge on the biology of this process may help in developing therapeutic strategies to prevent this frequent complication after vascular intervention.

MORPHOLOGY

The development of an intimal hyperplastic lesion has been very well described in animal models, especially in the balloon injury model in the rat³²⁻³⁴. In this model, a large artery is denudated by passage of an inflated balloon, introduced by a catheter. In the response to this injury which leads to intimal thickening, three different stadia are recognized³⁴. The first stage starts as soon as 24 hours after injury and is characterized by the replication of vascular smooth muscle cells (VSMC) in the media of the damaged arterial wall. In the second stage, these smooth muscle cells migrate from the media to the intima, with a peak between the third and fourth day. In the final stage of the process, VSMC replicate in the intima with a maximum at day 4-7. Furthermore, the intimal VSMC start producing matrix material like collagen, elastin and proteoglycans³⁵. The thickening of the intima proceeds up to the 12th week after injury, due to the production of extracellular matrix and volume increase of the intimal VSMC. After this period, the lesion remains stable. Histologic studies evaluating the morphology of intimal hyperplasia in humans showed that VSMC and extracellular matrix are also prominent in the lesions. This suggests that the biology of intimal hyperplasia in humans and animals may be similar^{8,36-38}. However, some observations suggest that differences may be present between the development of intimal thickening in humans and in animals. First of all, evaluation of cell proliferation in human

stenotic lesions, regarded as the key event in the development of intimal thickening in animals³⁹⁻⁴³, provided conflicting results. In studies investigating proliferation in restenotic tissue after PTCA, both high and low proliferation indices were observed⁴⁴⁻⁴⁶. In stenotic lesions, surgically derived from arterio-venous fistulas of hemodialysis patients high proliferation rates have been observed⁴⁶. A second point is that administration of agents, like heparin or ACE-inhibitors^{27,47-56}, decrease the extent of intimal thickening in animals. The effect of these agents in humans when administered locally or systemically is currently under investigation. Systemic administration has shown no beneficial effect so far^{57,58}.

FACTORS INFLUENCING INTIMAL HYPERPLASIA

Peri-anastomotic mismatch in elastic properties and intimal thickening

When two vessels with different elastic properties are anastomosed, an abrupt change in elastic properties occurs within a very short vessel segment. The peri-anastomotic difference in elastic properties is often called compliance mismatch. The suggested link between compliance mismatch and anastomotic intimal hyperplasia stems from many observations that intimal hyperplasia in peripheral arterial bypass grafts and graft AV fistulas develops at the site of the anastomoses^{29,30,59-64}. A mismatch in elastic properties may affect intimal hyperplasia in at least 3 different ways^{61,65-72}. First, a mismatch in elastic properties leads to a point of excessive mechanical stretching, which may result in vessel wall injury and initiate a vascular response resulting in intimal thickening⁷³. A second mechanism may be the positive influence of excessive cyclic stretching on replication of VSMC and production of extra-cellular matrix (ECM)^{33,74-79}. In vitro studies demonstrated that VSMC start to produce ECM and to replicate when they are subjected to high levels of distension. This phenomenon seems to be more pronounced in VSMC derived from veins as compared to arterial VSMC⁷⁴. Animal studies demonstrated that intimal thickening in vein grafts is associated with increased tangential stress⁸⁰⁻⁸³. A third mechanism may be the influence of compliance on flow characteristics^{84,85}. Flow studies demonstrated that a sudden increase in compliance is associated with enhanced particle residence time, flow stasis and separation, suggesting low shear conditions. A sudden decrease in compliance was not associated with these unfavourable flow conditions. In humans a sudden increase in compliance is observed in the carotid bulb, where also enhanced particle residence time, reversal of flow and flow separation have been observed⁸⁶. Since the carotid bulb is a predilection site for the occurrence of atherosclerosis, one might postulate that disturbed flow and increased particle residence time are associated with the development of an atherosclerotic plaque.

In animal studies evaluating the effect of compliance mismatch, lower patency rates in grafts with a low compliance were observed⁶⁵. However, a direct effect of compliance mismatch on intimal thickening at the anastomoses of grafts has never been demonstrated. This may be due to the absence of occlusive intimal hyperplasia in animal models⁸⁷⁻⁸⁹. In humans, lower patency rates are found in stiff prosthetic grafts

as compared to more compliant venous grafts. However, a prospective study investigating the effect of compliance mismatch around anastomoses on the occurrence of intimal thickening has never been conducted due to the absence of techniques able to quantify elastic properties around anastomoses in vivo in humans^{90,91}. With the introduction of Vessel Wall Doppler Tracking⁹² the assessment of elastic properties at defined sites in the graft and the native arterial system has become possible. This allows the calculation of compliance mismatch around the anastomoses of human grafts^{93,94}.

Blood flow characteristics and intimal thickening

Flow is an important modulator of vascular structure and function, probably by affecting the function of the endothelium⁹⁵⁻¹⁰². Due to their location at the interface of blood and arterial wall, endothelial cells are exposed to pulsatile pressure and flow, generated by the beating heart. Although the magnitude of the tangential acting shear forces is smaller than the transmural forces imposed by variations in pressure, it is recognized that oscillations in shear forces have an important effect on the structure and function of the vessel wall. The shape and structure of endothelial cells can be modified by changing shear rate levels¹⁰³⁻¹⁰⁸. Furthermore, the release of vaso-active substances and growth factors by the endothelium seems to be affected by changes in the level of shear rate¹⁰⁹⁻¹¹⁹. The crucial role of the endothelial cells in mediating structural changes in response to modulation of flow levels was demonstrated by Langille, who found a 21% decrease in rat carotid artery diameter after reduction of flow in arteries with a normal endothelial cell lining^{120,121}. In denudated arteries no structural adaptations to reduced flow were observed. However, shear rate dependent functional and structural changes in the vessel wall have also been observed in the absence of endothelium^{95,122}, suggesting that the VSMC's beneath the endothelium also have the ability to sense differences in flow.

The effect of varying levels of shear rate on intimal thickening has been studied extensively in animals¹²³⁻¹²⁵. The first studies conducted in this field suggested that high shear rate levels have an injuring effect on the endothelial cell (EC) layer, thereby inducing a vascular response^{126,127}. High shear rate levels resulted in endothelial cell edema, deformation and eventually in EC erosion. These early observations are supported by a study investigating the occurrence of intimal hyperplasia in graft AV fistulas in dogs in relation to flow velocities. More intimal thickening was observed in areas with the highest flow velocities^{128,129}. However, the current opinion is that increased intimal thickening occurs in areas of low shear rates^{86,97,99,122,130-133}. This has been observed in at least two different models. In the balloon catheter injury model, increased flow velocities are associated with a decrease in intimal-medial thickening¹²⁴. Also in models based on the placement of a prosthetic graft in the arterial system, increased flow velocities resulted in decreased intimal thickening¹²³. Studies on flow models based of the human carotid bifurcation showed that in the carotid bulb, just opposite to the flow divider, low shear rate levels are present, illustrated by flow separation, enhanced particle residence time and reversal of flow. This part of the

carotid artery is predisposed to the development of atherosclerotic disease in contrast to the relatively disease free flow divider, where a high shear rate is present. This suggests an inhibiting effect of high shear rate on intimal thickening⁸⁶.

Influence of flow on growth factors release

The pathways by which flow acts on the vessel wall are still unknown, but an essential role in signal transduction is attributed to EC^{68,115,134-142}. In a flow model in rabbit iliac arteries it was shown that an increase in flow resulted in release of NO and vasodilatation¹¹⁰. The administration of a specific potassium channel antagonist blocked the flow mediated vasodilatation and NO release, suggesting that the signal transduction pathway underlying flow mediated vasodilatation may be the activation of a potassium channel on the endothelium cell membrane. The observation that increased flow results in a release of NO by endothelial cells was confirmed in a cell culture study, using bovine aortic EC¹⁰⁹. Besides a vasodilatory effect of an increase in NO production with increasing shear rate, increased NO release may also result in a reduction of platelet adherence and aggregation^{143,144}. In a similar model it was shown that fluid shear stress modulates intracellular calcium concentration in vascular endothelial cells. A transient increase in free calcium concentration was observed after increasing shear rates to a certain level¹⁴⁵. The increase in free calcium concentration was proportional to the increase in shear rate, suggesting an important role of calcium in the signal transduction pathway, underlying flow mediated responses of the vessel wall. The shear rate dependent release of growth factors by endothelial cells is reflected by a varying expression of multiple growth factor related genes when wall shear rate is changed. A possible mechanism underlying the increased expression of growth factor genes may be a recently described shear rate induced transcription factor¹⁴⁶, which may not only regulate the expression of platelet derived growth factor (PDGF) genes, but also of tissue plasminogen activator, intracellular adhesion molecule 1 (ICAM-1) and transforming growth factor β 1 (TGF- β 1). For instance, a shear dependent release of PDGF β mRNA has been observed in cultured human umbilical vein endothelial cells (HUVEC)¹¹⁴. Of interest is that the release of PDGF β mRNA increased when shear stress increased from 0 to 6 dyn/cm², then gradually declined to a minimum at a shear stress level of 31 dyn/cm² and increased again when shear stress rose to 50 dyn/cm². Since PDGF is a mitogen and a potent vasoconstrictor, one may speculate that a certain level of shear stress is associated with a minimum release of PDGF by endothelial cells, resulting in vasodilatation and inhibition of VSMC growth. In addition, these data indicate that in both low and high shear stress conditions, increased expression of PDGF may occur, possibly resulting in promotion of VSMC growth^{147,148}. The stimulatory effects of both high and low flow on growth factor release like PDGF may explain the apparent discrepancy in the literature that both high and low shear rates are associated with increased vascular growth.

So far, it is unclear whether the signal transduction pathways of shear rate on vascular function are more susceptible to mean shear rate or to pulsatile variations in shear rate. In most models the effect of mean shear rate on intimal thickening was studied, since

this can be calculated from the Haagen-Poiseuille formula^{123,124}. However, some studies show evidence that the temporal shear rate gradient (value between the minimum and maximum value of shear rate within a cardiac cycle) is of more importance than the mean level of shear. First, in an in-vitro study it was shown that in regions of low mean shear rate but with large temporal gradients, the endothelial cells structure is heavily affected in that region¹⁴⁹. Endothelial cells migrated away from these sites with the remaining cells starting to divide at a high rate. This suggests that this flow mediated response of endothelial cells is determined by the temporal gradient in wall shear rate rather than the mean level of shear. In a study evaluating the possible mechanisms leading to post-stenotic dilatation, it was shown that the mean level of shear was very low in these areas, but the temporal gradient was high¹⁰². This also suggests that the shear rate gradient may be more important in modifying vascular function and structure than the mean level of shear rate. A recently published study showed that the development of anastomotic intimal hyperplasia decreases temporal gradients of shear rate at the site of the stenosis¹⁵⁰. This suggests that the development of intimal thickening in the anastomotic area may be at least in part an adaptive response in order to decrease high wall shear rate gradients and supports the concept that the gradient in shear rate may be the regulatory factor rather than a certain mean level of shear rate.

REFERENCES

1. Carrel A. Resultats eloignes de la transplantation des veines sur les arteres. *Rev De Chir* 1910; 41:987.
2. Bertrand ME, Lablanche JM, Fourrier JL, Gommeaux A, Ruel M. Relation to restenosis after percutaneous transluminal coronary angioplasty to vasomotion of the dilated coronary arterial segment. *Am J Cardiol* 1989; 63:277-281.
3. Vlietstra RE, Holmes DR, Rodeheffer RJ, Baily KR. Consequences of restenosis after coronary angioplasty. *Int J Cardiol* 1991; 31:143-148.
4. Waller BF, Pinkerton CA, Orr CM, Slack JD, VanTassel JW, Peters T. Restenosis 1 to 24 months after clinically succesful coronary ballon angioplasty: A necropsy study of 20 patients. *J Am Coll Cardiol* 1991; 17:58b-70b.
5. Serruys PW, Foley DP, Kirkheide RL, King III SB. Restenosis revisited: Insights provided by quantitative coronary angiography. *Am Heart J* 1993; 126:1243-1267.
6. Serruys PW, Breeman A. Coronary angioplasty - Long-term follow-up results and detection of restenosis: guidelines for aviation cardiology. *Eur Heart J* 1992; 13:76h-88h.
7. Liu MW, Roubin GS, King III SB. Restenosis after coronary angioplasty. Potential biologic determinants and role of intimal hyperplasia. *Circulation* 1989; 79:1374-1387.
8. Liu MW, Berk BC. Restenosis following coronary balloon angioplasty. Role of smooth muscle cell proliferation. *Trends Cardiovasc Med* 1991; 1:107-111.
9. Serruys PW, Luyten HE, Beatt KJ, et al. Incidence of restenosis after successful coronary angioplasty: a time-related phenomenon. *Circulation* 1988; 77:361-371.
10. The Veterans Administration Coronary Bypass Surgery Cooperative Study Group. Eleven-year survival in the veterans administration randomized trial of coronary bypass surgery for stable angina. *New Eng J Med* 1984; 311:1333-1339.
11. Harris PL, How TV, Jones DR. Prospectively randomized clinical trial to compare in situ and reversed saphenous vein grafts for femoropopliteal bypass. *Br J Surg* 1987; 74:252-255.
12. Harris RW, Andros G, Dulawa LB, Oblath RW, Apyan R, Salles-Cunha S. The transition to "in situ" vein bypass grafts. *Surg Gynecol Obstet* 1986; 163:21-28.
13. Harris PL, Veith FJ, Shanik GD, Nott D, Wengerter KR, Moore DJ. Prospective randomized comparison of in situ and reversed infrapopliteal vein grafts. *Br J Surg* 1993; 80:173-176.
14. Acher CW, Turnipseed WD. In situ distal saphenous vein bypass using the intraluminal valve-disruption technique. *Arch Surg* 1985; 120:933-936.
15. Baird RJ, Tutassaura H, Hiyagishima RT. Saphenous vein bypass graft to arteries of the ankle and foot. *Ann Surg* 1970; 172:1059-1063.
16. Batson RC, Sottiurai VS. Nonreversed and in situ grafts. Clinical and experimental observations. *Ann Surg* 1985; 201:771-779.
17. Buchbinder D, Rolins DL, Verta MJ, et al. Early experiance with in situ saphenous vein bypass for distal arterial reconstruction. *Surgery* 1986; 99:350-356.
18. Buchbinder D, Singh JK, Karmody AM, Leather RP, Shah DM. Comparison of patency rate and structural changes of in situ and reversed vein arterial bypass. *J Surg Res* 1981; 30:213-222.
19. Hurley JJ, Auer AI, Binnington HB, et al. Comparison of initial limb salvage in 98 consecutive patients with either reversed autogenous or in situ vein bypass graft procedures. *Am J Surg* 1985; 150:777-781.
20. Najmaldin A, Clifford PC, Farrands PA, Campbell MJ, Chant AD, Webster JH. The succes of in situ vein grafting: Early results in comparison with the reversed vein technique. *Eur J Vasc Surg* 1987; 1:165-168.
21. Leather RP, Shah DM, Karmody AM. Infrapopliteal arterial bypass for limb salvage: Increased patency and utilization of the saphenous vein used "in situ". *Surgery* 1981; 90:1000-1008.

22. Prendiville EJ, Yeager A, O'Donnell TF, et al. Long-term results with the above-knee popliteal expanded polytetrafluoroethylene graft. *J Vasc Surg* 1990; 11:517-524.
23. Shah DM, Buchbinder D. Modified technique to produce valvular incompetence in in situ saphenous vein arterial bypass. *Arch Surg* 1981; 116:356-358.
24. Taylor LM, Edwards JM, Phinney ES, Porter JM. Reversed vein bypass to infrapopliteal arteries. *Ann Surg* 1987; 205:90-97.
25. Taylor LM, Edwards JM, Porter JM. Present status of reversed vein bypass grafting: Five-year results of a modern series. *J Vasc Surg* 1990; 11:193-206.
26. Taylor RS, Loh A, McFarland RJ, Cox M, Chester JF. Improved technique for polytetrafluoroethylene bypass grafting: long-term results using anastomotic vein patches. *Br J Surg* 1992; 79:348-354.
27. Endean ED, Kispert JF, Martin KW, O'Conner W. Intimal hyperplasia is reduced by ornithine decarboxylase inhibition. *J Surg Res* 1991; 50:634-637.
28. Tordoir JHM, Herman JM, Kwan TS, Diderich PM. Long-term follow-up of the polytetrafluoroethylene (PTFE) prosthesis as an arteriovenous fistula for hemodialysis. *Eur J Vasc Surg* 1988; 2:3-7.
29. Swedberg SH, Brown BG, Sigley R, Wight TN, Gordon D, Nicholls SC. Intimal fibromuscular hyperplasia at the venous anastomosis of PTFE grafts in hemodialysis patients. *Circulation* 1989; 80:1726-1736.
30. Jenkins AMcL, Buist TAS, Glover SD. Medium-term follow-up of forty autogenous vein and forty polytetrafluoroethylene (Gore-Tex) grafts for vascular access. *Surgery* 1980; 88:667-672.
31. Windus DW. Permanent vascular access: a nephrologist's view. *Am J Kidney Dis* 1993; 21:457-471.
32. Boonen HCM, Schiffers PMH, Fazzi GE, Janssen GMJ, Daemen MJAP, Mey de JGR. DNA synthesis in isolated arteries. Kinetics and structural consequences. *Am J Physiol* 1991; 260:H1-H8.
33. Clowes AW, Clowes MM, Fingerle J, Reidy MA. Kinetics of cellular proliferation after arterial injury. V. Role of acute distension in the induction of smooth muscle cell proliferation. *Lab Invest* 1989; 60:360-364.
34. Clowes AW, Reidy MA, Clowes MM. Mechanisms of stenoses after arterial injury. *Lab Invest* 1983; 49:208-215.
35. Nikkari TN, Jarvelainen HT, Wight TN, Ferguson M, Clowes AW. Smooth muscle cell expression of extracellular matrix genes after arterial injury. *Am J Pathol* 1994; 144:1348-1356.
36. Garratt KN, Edwards WD, Kaufmann URS, Vlietstra RE, Holmes DR. Differential histopathology of primary atherosclerotic and restenotic lesions in coronary arteries and saphenous vein bypass grafts: Analysis of tissue obtained from 73 patients by directional atherectomy. *J Am Coll Cardiol* 1991; 17:442-448.
37. Ueda M, Becker AE, Fujimoto T, Tsukada T. The early phenomena of restenosis following percutaneous transluminal coronary angioplasty. *Eur Heart J* 1991; 12:937-945.
38. Sayers RD, Jones L, Varty K, et al. The histopathology of infrainguinal vein graft stenoses. *Eur J Vasc Surg* 1993; 7:16-20.
39. Schwartz RM, Campbell GR, Campbell JH. Replication of smooth muscle cells in vascular disease. *Circ Res* 1986; 58:427-444.
40. Schwartz SM. Common mechanisms of cell proliferation in hypertension and atherosclerosis. In: Zanchetti A, Tarazi RC, eds. 7th ed. New York: Elsevier Science Publishers, 1986:454-464.
41. Schwartz SM, Liaw L. Growth control and morphogenesis in the development and pathology of arteries. *J Cardiovasc Pharmacol* 1993; 21:s31-s49.
42. Gordon D, Schwartz SM. Cell proliferation in human atherosclerosis. *Trends Cardiovasc Med* 1991; January/February:24-28.

43. Schwartz CJ, Valente AJ, Sprague EA, Kelly JL, Nerem RM. The pathogenesis of atherosclerosis: An overview. *Clin Cardiol* 1991; 14:1-16.
44. O'Brien ER, Alpers CE, Stewart DK, et al. Proliferation in primary restenotic coronary atherectomy tissue. Implications for antiproliferative therapy. *Circ Res* 1993; 73:223-231.
45. Pickering JG, Weir L, Jekanowski J, Kearny MA, Isner JM. Proliferative activity in peripheral and coronary atherosclerotic plaque among patients undergoing percutaneous revascularisation. *J Clin Invest* 1993; 91:1469-1480.
46. Rehker M, Nicholls S, Ferguson M, Gordon D. Cell proliferation in human arteriovenous fistulas use for hemodialysis. *Arterioscler Thromb* 1993; 13:609-617.
47. Kouchi Y, Esato K, O-Hara M, Zempo N. Effect of prostaglandin I₂ analogue TRK-100 on the suppression of intimal fibrous proliferation. *J Vasc Surg* 1992; 16:232-238.
48. Kranzhöfer R, Schirmer J, Schömig A, et al. Suppression of neointimal thickening and smooth muscle cell proliferation after arterial injury in the rat by inhibitors of Na⁺-H⁺ exchange. *Circ Res* 1993; 73:264-268.
49. Mitsuka M, Nagae M, Berk BC. Na⁺-H⁺ exchange inhibitors decrease neointimal formation after rat carotid injury. Effects on smooth muscle cell migration and proliferation. *Circ Res* 1993; 73:269-275.
50. Powell JS, Clozel JP, Muller RKM, et al. Inhibitors of angiotensin-converting enzyme prevent myointimal proliferation after vascular injury. *Science* 1989; 245:186-188.
51. Powell JS, Rouge M, Muller RK, Baumgartner HR. Cilazapril suppresses myointimal proliferation after vascular injury: effects on growth factor induction in vascular smooth muscle cells. *Basic Res Cardiol* 1991; 86:65-74.
52. Vashiht R, Sian M, Franks PJ, O'Malley MK. Long-term reduction of intimal hyperplasia by the selective alpha-1 adrenergic antagonist doxazosin. *Br J Surg* 1992; 79:1285-1288.
53. Wilson NV, Salisbury JR, Kakkar VV. Effect of low molecular weight heparin on intimal hyperplasia. *Br J Surg* 1991; 78:1381-1383.
54. Clowes AW, Clowes MM. Kinetics of cellular proliferation after arterial injury. IV. Heparin inhibits rat smooth muscle mitogenesis and migration. *Circ Res* 1986; 58:839-845.
55. Dryjski M, Mikat E, Björnsson TD. Inhibition of intimal hyperplasia after arterial injury by heparins and heparinoid. *J Vasc Surg* 1988; 8:623-633.
56. Clowes AW, Clowes MM, Vergel SC, et al. Heparin and cilazapril together inhibit injury-induced intimal hyperplasia. *Hypertension* 1991; 18:II65-II69.
57. The MERCATOR Study Group. Does the new angiotensin converting enzyme inhibitor cilazapril prevent restenosis after percutaneous transluminal coronary angioplasty? *Circulation* 1992; 86:100-110.
58. Jackson CL. Animal models of restenosis. *Trends Cardiovasc Med* 1994; 4:122-130.
59. Imparato AM, Bracco A, Eun Kim G, Zeff R. Intimal and neointimal fibrous proliferation causing failure of arterial reconstructions. *Surgery* 1972; 72:1007-1017.
60. Baird RN, Abbott WM. Pulsatile blood-flow in arterial grafts. *Lancet* 1976; October 30:948-950.
61. Kidson IG, Abbott WM. Low compliance and arterial graft occlusion. *Circulation* 1978; 58:II-14.
62. LoGerfo FW, Quist WC, Nowak MD, Crawshaw HM, Haudenschild CC. Downstream anastomotic hyperplasia. A mechanism of failure in Dacron arterial grafts. *Ann Surg* 1983; 197:479-483.
63. Madras PN, Ward CA, Johnson WR, Singh PI. Anastomotic hyperplasia. *Surgery* 1981; 90:922-923.
64. Mills JL, Fujitani RM, Taylor SM. The characteristics and anatomic distribution of lesions that cause reversed vein graft failure: a five-year prospective study. *J Vasc Surg* 1993; 17:195-204.
65. Abbott WM, Megerman J. Adaptive responses of arteries to grafting. *J Vasc Surg* 1989; 9:377-379.

66. Kinley CE, Marble AE, Eng P. Compliance: A continuing problem with vascular grafts. *J Cardiovas Surg* 1980; 21:163-170.
67. Laogun AA, Gosling RG. In vivo arterial compliance in man. *Clin Phys Physiol Meas* 1982; 3:201-212.
68. Levenson J, Simon AC, Bouthier JD, Benetos A, Safar ME. Post-synaptic alpha-blockade and brachial artery compliance in essential hypertension. *J Hypertens* 1984; 2:37-41.
69. Schmitz-Rixen T, Megerman J, Anderson JM, et al. Longterm study of a compliant biological vascular graft. *Eur J Vasc Surg* 1991; 5:149-158.
70. Okuhn SP, Connelly DP, Calakos N, Ferrell L, Man-Xiang P, Goldstone J. Does compliance mismatch alone cause neointimal hyperplasia? *J Vasc Surg* 1989; 9:35-45.
71. Bassiouny HS, White S, Glagov S, Choi E, Giddens DP, Zarins CK. Anastomotic intimal hyperplasia: Mechanical injury or flow induced. *J Vasc Surg* 1992; 15:708-717.
72. Pourdeyhimi B, Wagner D. On the correlation between the failure of vascular grafts and their structural and material properties: A critical analysis. *J Biomed Mat Res* 1986; 20:375-409.
73. Pevec WC, L'Italien GJ, Megerman J, Cambria RP, Abbott WM. Abnormal wall strain at distal end-to-side anastomoses. *Ann Vasc Surg* 1993; 7:14-20.
74. Predel HG, Yang Z, Von Segesser L, Turina M, Buhler FR, Luscher TF. Implications of pulsatile stretch on growth of saphenous vein and mammary artery smooth muscle. *Lancet* 1992; 340:878-879.
75. Sottiurai VS, Kollross P, Glagov S, Zarins CK, Mathews MB. Morphologic alteration of cultured arterial smooth muscle cells by cyclic stretching. *J Surg Res* 1983; 35:490-497.
76. Sottiurai VS, Sue SL, Feinberg II EL, Bringaze WL, Tran AT, Batson RC. Distal anastomotic intimal hyperplasia: Biogenesis and etiology. *Eur J Vasc Surg* 1988; 2:245-256.
77. Leung DYM, Glagov S, Mathews MB. Cyclic stretching stimulates synthesis of matrix components by arterial smooth muscle cells in vitro. *Science* 1976; 191:475-477.
78. Leung DYM, Glagov S, Mathews MB. Elastin and collagen accumulation in rabbit ascending aorta and pulmonary trunk during postnatal growth. Correlation of cellular synthetic response with medial tension. *Circ Res* 1977; 41:316-323.
79. Sumpio BE, Banas AJ, Levin LG, Johnson G. Mechanical stress stimulates aortic endothelial cells to proliferate. *J Vasc Surg* 1987; 6:252-256.
80. Zwolak RM, Adams MC, Clowes AW. Kinetics of vein graft hyperplasia: Association with tangential stress. *J Vasc Surg* 1987; 5:126-136.
81. Dobrin PB, Littooy FN, Golan J, Blakeman B, Fareed J. Mechanical and histological changes in canine vein grafts. *J Surg Res* 1988; 44:259-265.
82. Dobrin PB, Littooy FN, Endean ED. Mechanical factors predisposing to intimal hyperplasia and medial thickening in autogenous vein grafts. *Surgery* 1989; 105:393-400.
83. Dobrin PB. On the roles of deformation, tension, and wall stress as critical stimuli eliciting myointimal/medial hyperplasia. *J Vasc Surg* 1991; 581-582.
84. Stewart SFC, Lyman DJ. Effects of a vascular graft/natural artery compliance mismatch on pulsatile flow. *J Biomech* 1992; 25:297-310.
85. Ojha M, Cobbold RS, Johnston KW. Hemodynamics of a side-to-end proximal arterial anastomosis model. *J Vasc Surg* 1993; 17:646-655.
86. Zarins CK, Giddens DP, Bharadvaj BK, Sottiurai VS, Mabon RF, Glagov S. Carotid bifurcation atherosclerosis. Quantitative correlation of plaque localization with flow velocity profiles and wall shear stress. *Circ Res* 1983; 53:502-514.
87. McBride W, Lange RA, Hillis LD. Restenosis after successful coronary angioplasty. Pathophysiology and prevention. *New Eng J Med* 1988; 318:1734-1737.
88. Wu MH, Shi Q, Sauvage LR, et al. The direct effect of graft compliance mismatch per se on development of host arterial intimal hyperplasia at the anastomotic interface. *Ann Vasc Surg* 1993; 7:156-168.

89. Ferrel M, Fuster V, Gold HK, Chesebro JH. Choosing appropriate experimental animal model for the prevention of restenosis. *Circulation* 1992; 85:1630-1631.
90. Mozersky DJ, Sumner DS, Hokanson DE, Strandness DE. Transcutaneous measurement of the elastic properties of the human femoral artery. *Circulation* 1972; XLIV:948-955.
91. Lye CR, Sumner DS, Hokanson DE, Strandness DE. The transcutaneous measurement of the elastic properties of the human saphenous vein femoropopliteal bypass graft. *Surg Gynecol Obstet* 1975; 141:891-895.
92. Hoeks APG, Brands PJ, Smeets FAM, Reneman RS. Assessment of distensibility of superficial arteries. *Ultrasound Med Biol* 1990; 16:121-128.
93. Hofstra L, Bergmans DCJJ, Hoeks APG, Tordoir JHM, Kitslaar PJEHM. Assessment of inhomogeneities in elastic properties of in situ and reversed saphenous vein grafts in humans. *Eur J Vasc Surg* 1994; 100:100-105.
94. Hofstra L, Bergmans DCCJ, Hoeks APG, Kitslaar PJEHM, Leunissen KML, Tordoir JHM. Mismatch in elastic properties around anastomoses of interposition grafts for hemodialysis access. *JASN* 1994; 45:45-55.
95. Bevan JA, Joyce EH, Wellman GC. Flow-dependent dilation in a resistance artery still occurs after endothelium removal. *Circ Res* 1988; 63:980-985.
96. Kamiya A, Togawa T. Adaptive regulation of wall shear stress to flow change in the canine carotid artery. *Am J Physiol* 1980; 239:H14-H21.
97. Zarins CK, Bomberger RA, Glagov S. Local effects of stenoses: Increased flow velocity inhibits atherogenesis. *Circulation* 1981; 64:II221-II227.
98. Cashin WL, Sanmarco ME, Nessim SA, Blankenhorn DH. Accelerated progression of atherosclerosis in coronary vessels with minimal lesions that are bypassed. *New Eng J Med* 1984; 311:824-828.
99. Asakura T, Karino T. Flow patterns and spatial distribution of atherosclerotic lesions in human coronary arteries. *Circ Res* 1990; 66:1045-1066.
100. Kamiya A, Ando J, Shibata M, Masuda H. Roles of fluid shear stress in physiological regulation of vascular structure and function. *Biorheology* 1988; 25:271-277.
101. LaBarbera M. Principles of design of fluid transport systems in zoology. *Science* 1990; 249:992-999.
102. Ojha M, Johnston KW, Cobbold RS. Evidence of a possible link between poststenotic dilatation and wall shear stress. *J Vasc Surg* 1990; 11:127-135.
103. Nerem RM. Vascular fluid mechanics, the arterial wall, and atherosclerosis. *J Biomech Eng* 1992; 114:274-282.
104. Nerem RM, Girard PR. Hemodynamic influences on vascular endothelial biology. *Toxicologic Pathology* 1990; 18:572-582.
105. Cambria RP, Megerman J, Abbott WM. Endothelial preservation in reversed and in situ autogenous vein grafts. *Ann Surg* 1985; 202:50-55.
106. Nerem RM, Harrison DG, Taylor WR, Alexander RW. Hemodynamics and vascular endothelial biology. *J Cardiovasc Pharmacol* 1993; 21:s6-s10.
107. Langille BL, Graham JJ, Kim D, Gotlieb AI. Dynamics of shear induced redistribution of F-actin in endothelial cells in vivo. *Arterioscler Thromb* 1991; 6:1814-1820.
108. Langille BL, Reidy M, Kline R. Injury and repair of endothelium at sites of flow disturbances near abdominal aortic coarctations in rabbits. *Arterioscler Thromb* 1988; 6:146-154.
109. Buga GM, Gold ME, Fukuto JM, Ignarro LJ. Shear stress-induced release of nitric oxide from endothelial cells grown on beads. *Hypertension* 1991; 17:187-193.
110. Cooke JP, Rossitch E, Andon NA, Loscalzo J, Dzau VJ. Flow activates an endothelial potassium channel to release an endogenous nitrovasodilator. *J Clin Invest* 1991; 88:1663-1671.

111. Ralevic V, Milner P, Hudlicka O, Kristek F, Burnstock G. Substance P is released from the endothelium of normal and capsaicin-treated rat hind-limb vasculature, in vivo, by increased flow. *Circ Res* 1990; 66:1178-1183.
112. Rubanyi GM, Freay AD, Kauser K, Johns A, Harder DR. Mechanoreception by the endothelium: Mediators and mechanisms of pressure- and flow-induced vascular responses. *Blood Vessels* 1990; 27:246-257.
113. Tesfamariam B, Cohen RA. Inhibition of adrenergic vasoconstriction by endothelial cell shear stress. *Circ Res* 1988; 63:720-725.
114. Hsieh HJ, Li NQ, Frangos JA. Shear stress increases endothelial platelet-derived growth factor mRNA levels. *Am J Physiol* 1991; 260:H642-H646.
115. Lüscher TF. Endothelial control of vascular tone and growth. *Clin Exp Hypertens* 1990; 12:897-902.
116. Melkumyants AM, Balashov SA, Khayutin VM. Endothelium dependent control of arterial diameter by blood viscosity. *Cardiovasc Res* 1989; 23:741-747.
117. Olesen SP, Clapham DE, Davies PF. Hemodynamic shear stress activates a K⁺ current in vascular endothelial cells. *Nature* 1988; 331:168-170.
118. Werner GS, Wiegand V, Kreuzer H. Effect of acetylcholine on arterial and venous grafts and coronary arteries in patients with coronary artery disease. *Eur Heart J* 1990; 11:127-137.
119. Werner GS, Wiegand V, Tebbe U, Kreuzer H. Contrary effects of acetylcholine on coronary arteries and aortocoronary venous grafts in man. *Eur Heart J* 1989; 10:86-91.
120. Langille BL, O'Donnell F. Reductions in arterial diameter produced by chronic decreases in blood flow are endothelium-dependent. *Science* 1986; 231:405-407.
121. Langille BL, Bendeck MP, Keeley FW. Adaptations of carotid arteries of young and mature rabbits to reduced carotid blood flow. *Am J Physiol* 1989; 256:H931-H939.
122. Sterpetti AV, Cucina A, Santoro L, Cardillo B, Cavallaro A. Modulation of arterial smooth muscle cell growth by hemodynamic forces. *Eur J Vasc Surg* 1992; 6:16-20.
123. Kohler TR, Kirkman TR, Kraiss LW, Zierler BK, Clowes AW. Increased blood flow inhibits neointimal hyperplasia in endothelialized vascular grafts. *Circ Res* 1991; 69:1557-1565.
124. Kohler TR, Jawien A. Flow affects development of intimal hyperplasia after arterial injury in rats. *Arterioscler Thromb* 1992; 12:963-971.
125. Morinaga K, Eguchi H, Miyazaki T, Okadome K, Sugimachi K. Development and regression of intimal thickening of arterially transplanted autologous vein grafts in dogs. *J Vasc Surg* 1987; 5:719-730.
126. Fry DL. Acute vascular endothelial changes are associated with increased blood velocity gradients. *Circ Res* 1968; 22:165-197.
127. Fry DL. Certain chemorheologic considerations regarding the blood vascular interface with particular reference to coronary artery disease. *Circulation* 1969; XXXIX and XI:IV38-IV59.
128. Fillinger MF, Reinitz ER, Schwartz RA, Resetarits DE, Paskanik AM, Bredenberg CE. Beneficial effects of banding on venous intimal-medial hyperplasia in arteriovenous loop grafts. *Am J Surg* 1989; 158:87-94.
129. Fillinger MF, Reinitz ER, Schwartz RA, et al. Graft geometry and venous intimal-medial hyperplasia in arteriovenous loop grafts. *J Vasc Surg* 1990; 11:556-566.
130. Rittgers SE, Karayannacos PE, Guy JF, et al. Velocity distribution and intimal proliferation in autologous vein grafts in dogs. *Circ Res* 1978; 42:792-801.
131. Zarins CK, Zatina MA, Giddens DP, Ku DN, Glagov S. Shear stress regulation of artery lumen diameter in experimental atherogenesis. *J Vasc Surg* 1987; 5:413-420.
132. Friedman MH. Some atherosclerosis may be a consequence of the normal adaptive response to shear. *Atherosclerosis* 1990; 82:193-196.
133. Friedman MH, Barger CB, Deters OJ, Hutchins GM, Mark FF. Correlation between wall shear and intimal thickness at a coronary artery branch. *Atherosclerosis* 1987; 68:27-33.

134. Koo EWY, Gotlieb AI. Endothelial stimulation of intimal cell proliferation in a porcine aortic organ culture. *Am J Pathol* 1989; 134:497-503.
135. Lüscher TF. Imbalance of endothelium-derived relaxing and contracting factors. A new concept in hypertension? *Am J Hypertens* 1990; 3:317-330.
136. Luscher TF, Diederich D, Siebeenman R, Lehman K, Stulz P, Von Segesser L, Yang Z, Turina M, Weber E, Buhler FR. Difference between endothelium-dependent relaxation in arterial and venous coronary bypass grafts. *New Eng J Med* 1988; 319:462-467
137. Ludmer PL, Selwyn AP, Shook TL, et al. Paradoxical vasoconstriction induced by acetylcholine in atherosclerotic coronary arteries. *New Eng J Med* 1986; 315:1046-1051.
138. Mey de JGR, Dijkstra E, Fazzi G, Janssen G. Growth responses in the arterial wall. *Basic Res Cardiol* 1991; 86:13-17.
139. Mey de JGR, Dijkstra EH, Vrijdag MJF. Endothelium reduces DNA synthesis in isolated arteries. *Am J Physiol* 1991; 260:H1-H7.
140. Mey de JGR, Uitendaal MP, Boonen HCM, Schiffers PMH, Fazzi GE. Growth responses in isolated elastic, muscular and resistance-sized arterial segments of the rat. *Blood Vessels* 1991; 28:372-385.
141. Vanhoutte PM, Shimokawa H. Endothelium-derived relaxing factor and coronary vasospasm. *Circulation* 1989; 80:1-9.
142. Vanhoutte PM, Verbeuren TJ, Webb RC. Local modulation of adrenergic neuroeffector interaction in the blood vessel wall. *Physiol Rev* 1981; 61:151-215.
143. Furchgott RF, Vanhoutte PM. Endothelium-derived relaxing and contracting factors. *FASEB J* 1989; 3:2007-2018.
144. Linder L, Kiowski W, Bühler FR, Lüscher TF. Indirect evidence for release of endothelium-derived relaxing factor in human forearm circulation in vivo. Blunted response in essential hypertension. *Circulation* 1990; 81:1762-1767.
145. Shen J, Luscinskas FW, Conolly A, Dewey CF, Gimbrone MA. Fluid shear stress modulates cytosolic free calcium in vascular endothelial cells. *Am J Cell Physiol* 1992; 31:c384-c390.
146. Resnick N, Collins T, Atkinson W, Bonthron DT, Dewey CF, Gimbrone MA. Platelet-derived growth factor β chain contains a cis-acting fluid shear-stress-responsive element. *Proc Natl Acad Sci U S A* 1993; 90:4591-4595.
147. Nabel EG, Yang Z, Liptay S, et al. Recombinant platelet-derived growth factor β gene expression in porcine arteries induce intimal hyperplasia in vivo. *J Clin Invest* 1993; 91:1822-1829.
148. Taubman MB, Rollins BJ, Poon M, et al. JE mRNA accumulates rapidly in aortic injury and in platelet-derived growth factor-stimulated vascular smooth muscle cells. *Circ Res* 1992; 70:314-325.
149. DePaola N, Gimbrone MA, Davies PF, Dewey CF. Vascular endothelium responds to fluid shear stress gradients. *Arterioscler Thromb* 1992; 12:1254-1257.
150. Ojha M. Wall shear stress temporal gradient and anastomotic intimal hyperplasia. *Circ Res* 1994; 74:1227-1231.

CHAPTER 3

VESSEL WALL DOPPLER TRACKING: A NEW TECHNIQUE TO ASSESS VESSEL WALL COMPLIANCE

L Hofstra, EJM Ermers, APM Langeveldt, APG Hoeks, PJEHM Kitslaar
Adapted from the Journal of Vascular Technology, 1993; 17:65-68

ABSTRACT

Background. To obtain insight in the pathogenesis of atherosclerosis and intimal thickening, information on mechanical arterial vessel wall properties is necessary.

Methods. A Vessel Wall Doppler Tracking (VWDT) system, based on ultrasound M-mode detection in combination with computerized data-acquisition on a PC, was developed in our laboratory. This system transcutaneously measures vessel diameter and diameter change during a successive number of heartbeats. From these parameters and simultaneously recorded pulse pressure using Dinamap, relative distension (RD), distensibility coefficient (DC= relative distension per unit of pressure) and cross-sectional compliance (CC = area increase per unit of pressure) can be calculated. RD and DC are measures of mechanical stress exerted on the vessel wall, CC is a measure of the capacity of the vessel to store blood volume during cardiac cycle.

Results. We studied the vessel wall characteristics in the right common femoral artery in 41 healthy subjects (mean age: 39, range: 24-70 years, 19 male and 22 female). Diameter was significantly higher ($p<0.01$) in males (9.95 ± 0.40 mm) than in females (8.49 ± 0.22 mm). CC was also significantly higher ($p<0.05$) in the male group (0.837 ± 0.096 mm²/kPa) than in the female group (0.604 ± 0.044 mm²/kPa). RD did not differ significantly between males ($3.4\pm0.3\%$) and females ($3.8\pm0.3\%$). In DC no significant differences between males ($7.4\pm0.8 \cdot 10^{-3}$ kPa⁻¹) and females ($7.4\pm0.8 \cdot 10^{-3}$ kPa⁻¹) was found. Diameter, RD and CC did not change with age.

Conclusion. We conclude that VWDT is a valuable tool for establishing arterial vessel wall characteristics. The capacity of the common femoral artery to transport blood is higher in males than in females but the mechanical stress on the arterial vessel wall does not differ. These data suggest that VWDT might be of clinical relevance e.g. in bypass surveillance and early detection of atherosclerosis.

INTRODUCTION

The compliance of large conduit arteries, like the carotid and femoral arteries, play an important role in buffering phasic changes in pressure and flow during the cardiac cycle. This results in a gradual transition from phasic flow close to the heart to a more continuous flow in peripheral arteries.

From a pathogenetic and possibly diagnostic point of view it is important to study elastic properties of arteries. For instance hypertension and diabetes mellitus, which are major risk factors in the development of atherosclerosis^{1,2}, are associated with a decrease in compliance and distensibility of arteries³⁻¹⁵. An established technique to assess compliance of arteries, is the measurement of pulse wave velocity (PWV)^{3,4,8,15}. Stiffening of arteries leads to an increase in pulse wave velocity^{3,8}. This provides information on the global compliance in a defined segment of an artery. However, an increased PWV allows no distinction between plaque formation and general stiffening of the arterial segment investigated.

Vessel Wall Doppler Tracking (VWDT) is a new system to assess local elastic properties of superficial arteries¹⁶. This ultrasound system combines B-mode and processing of radio frequency (RF) signals to measure arterial diameter and diameter change during a cardiac cycle. In this article a description of the technique is presented. Furthermore, the preliminary results of measurements of compliance and distensibility of the common femoral artery, obtained in 41 presumed healthy volunteers, are reported.

METHODS

The VWDT-system consists of an echo-dopplersystem (ATL ULTRAMARK IV), a memory module and a Personal Computer. The vessel of interest is visualized in B-mode, to obtain anatomical and structural information of the vessel. A cursor line is positioned perpendicular to the arterial walls (Figure 1). A number (2048) of sequential lines, based on the received and non-processed radiofrequency (RF) signals, are digitized and temporarily stored in a memory¹⁶. Each line is transferred to a personal computer. The first digitized RF-line is displayed on the computer screen, allowing the operator to identify the anterior and posterior wall of the vessel. Two sample volumes, indicated by boxes, are placed on the anterior and posterior wall, respectively (Figure 2). The distance between the boxes corresponds with the diameter of the artery. The selected signal segments are subjected to Doppler-signal processing yielding a displacement signal with a high resolution. Based on the detected displacement of the walls, the positions of the sample volumes, representing the anterior and posterior wall, are updated for each subsequent line. Thus the position of the sample volume is linked with the vessel wall structure and follows the position of the walls (tracking). The process is triggered by an ECG-signal, which facilitates the detection of the maximum (systolic) and minimum (diastolic) arterial diameter within a cardiac cycle. The minimum diameter after an ECG trigger is considered as the initial diameter. The

difference between maximum and minimum diameter gives the change in diameter during a cardiac cycle. The result of a typical recording is shown in figure 3. The values of the parameters obtained during 3-5 heartcycles are averaged as shown in table I. During the measurements pulse pressure recordings (i.e. the difference between systolic and diastolic pressure) were obtained at the left arm with the use of an automated cuff device (Dinamap^R).

Figure 1: B-mode image of the common femoral artery (left panel). The cursor line is positioned perpendicular to the vessel. In M-mode the displacement of the artery wall can be observed. Arrows indicate the anterior (top) and posterior (bottom) wall of the common femoral artery.

Figure 2: The digitized RF-signal of the first acquired line. The anterior (left arrow) and posterior wall (right arrow) are identified and covered by sample-volumes indicated by the boxes.

Figure 3: Displacement waveforms of the anterior (top) and posterior (middle) wall of the common femoral artery. This results in the distension waveform (bottom), where a cross indicates the occurrence of an ECG-trigger.

Table I: Parameters derived from the distension waveform (see fig 3). The parameter values obtained during 4 successive cardiac cycles are averaged.

Beat	Diameter(mm)	Distension(μ m)	Relative Distension (%)
0	9.25	543	5.9
1	9.23	520	5.6
2	9.21	565	6.1
3	9.29	481	5.2
mean	9.25	527	5.7
std dev	0.04	36	0.41

From diameter, diameter change and simultaneously recorded pulse pressure the cross-sectional compliance (CC, mm²/kPa), distensibility coefficient (DC, 1/kPa) and relative distension (RD, %) can be calculated, using the following equations:

$$CC = \pi \frac{\Delta D * D}{2 \Delta P} \quad (1)$$

$$DC = \frac{2 \Delta D / D}{\Delta P} \quad (2)$$

$$RD = \Delta D / D * 100 \quad (3)$$

(D = diameter, ΔD = diameter change and ΔP = pulse pressure). The cross-sectional compliance is the cross-sectional area increase per unit of pressure, which is a measure of the capacity of the vessel to store blood volume during the cardiac cycle. The distensibility coefficient is the fractional diameter increase per unit of pressure, representing the stress-strain relationship of the arterial wall. Relative distension is the percentage diameter increase during the cardiac cycle.

The subjects, 19 male and 22 female, were measured in the supine position after an acclimatization period of 20 minutes. The measurements took place in a climatized laboratory, with a constant temperature of 24°C. There were no significant differences in age and smoking habits between the male and female volunteers (Student t-test).

The right common femoral artery was measured three times in each subject within 20 minutes. From these measurements, the median value of each parameter was used in this study to diminish the influence of outliers.

RESULTS

In all subjects, visualization of the common femoral artery was easily possible. A summary of the results is given in table II and III. Diameter and CC were significantly higher in the male group as compared to the female group (Student t-test, $p < 0.05$). In DC no difference could be found between the groups. RD was slightly higher in the female group as compared to the male group. Linear regression analysis revealed a significant negative correlation for CC ($r = -0.52$, $p < 0.05$) and DC ($r = -0.56$, $p < 0.05$) with age in the female group. RD in the female volunteers and all parameters in the male group lacked a significant correlation with age.

Table II: Arterial diameter, relative distension (RD), distensibility coefficient (DC), cross-sectional compliance (CC) and pulse pressure for the male and the female group (mean $\pm$ SE). Diameter and CC are significantly higher in the male group as compared to the female group (Student T-test).

Parameter	Male	Female	P-value
Diameter (mm)	9.95 (0.41)	8.49 (0.22)	0.01
RD (%)	3.42 (0.31)	3.83 (0.32)	0.36
DC (10^{-3} /kPa)	7.4 (0.76)	7.4 (0.76)	0.98
CC (mm^2/kPa)	0.837 (0.097)	0.604 (0.064)	0.05
Pulsepressure (mmHg)	51 (3.1)	56 (2.4)	0.25

Table III: Correlations of DC, CC and RD with age for the male and the female group, calculated with the use of linear regression analysis. (* = significant correlation, $p<0.05$).

	Male	Female
DC	-0.06	-0.56*
CC	0.25	-0.52*
RD	-0.12	-0.38

DISCUSSION

VWDT is a new technique, where visualization in B-mode is combined with off-line data processing. Based on B-mode imaging structural information about arteries, like plaque formation and distance to the bifurcations, can be obtained. Plaque formation in the vessel wall, probably influencing the elastic properties can be identified. B-mode imaging is also very helpful in longitudinal studies, where it is of importance to define the exact site of measurement in the subsequent sessions. Using landmarks, like the bifurcations and anastomoses, this can be achieved¹⁶.

Off-line data-processing has the advantage to identify carefully the position of the vessel wall reflections representing the anterior and posterior wall¹⁶. Doppler tracking provides the required high resolution necessary to assess compliance and distensibility accurately.

The preliminary data obtained from the common femoral artery in 41 healthy volunteers, show differences between male and female. Cross-sectional Compliance (CC), which is a measure of the capacity of the vessel to store blood volume during the cardiac cycle, is larger in the male group than in the female group. This is due to a difference in diameter rather than a difference in local elastic properties of the vessel, which is represented by RD and DC.

In the female group a significant decrease with age in compliance and distensibility of the common femoral artery was found. In the male group no significant relation with age was shown. Some authors demonstrated age-related changes in elastic properties of large arteries^{8,15,17}. In most studies a decrease in compliance with increasing age was reported. Laogun¹⁵ reported a difference in age related changes in aortic compliance between male and female. Male aortic compliance was found to decrease steadily between 20 and 50 yrs of age. On the other hand, compliance of the aorta in females showed hardly any decrease with age between 20 and 45 yrs., but a sharp fall was found between 45 and 50 yrs.. Our data also show a difference in age-related changes in compliance between male and female. This could be the effect of the difference in hormonal status between male and female before menopause.

VWDT is a promising new technique to assess accurately the local vessel wall properties of arteries. This system could become of great value to study compliance and distensibility in hypertensive and diabetic subjects, and to obtain more insight in the pathogenetic mechanisms resulting in the development of atherosclerosis. Furthermore, attention can be paid to vessel wall characteristics of venous grafts used as a bypass conduit in peripheral revascularisation, to get a better understanding in the occurrence of stenotic lesions, often leading to failure of bypasses.

APPLICATION OF VWDT IN VASCULAR SURGERY

In vascular surgery, saphenous vein grafts are often used in the construction of a bypass in order to reestablish flow to the lower limb. The occurrence of stenotic lesions in peripheral bypass grafts is a major threat to the function of the vascular conduit and often necessitates reintervention, leading to morbidity and hospitalisation of patients undergoing this kind of surgery. One of the causes for the occurrence of stenosis may be a difference in elastic properties between the venous graft and the native arterial system (mismatch). Quantitative data on the peri-anastomotic differences in elastic properties are scarce, which may be attributed to the lack of a suitable system to assess elastic properties at defined sites in the vascular system. With the

introduction of VWDT, measurements of elastic properties at pre-defined sites are possible, due to the help of the B-mode image. This facilitates the positioning of the cursor line perpendicular to the vessel walls. Since the anastomoses of peripheral bypasses provide a useful landmark on the B-mode image, assessment of elastic properties at pre-defined sites in the graft and native arterial system is possible. Therefore, in theory VWDT is a suitable system to assess spatial changes in vessel wall elastic properties around anastomoses of vascular reconstructions. Also the calculation of mismatch in elastic properties around anastomoses may be possible. This provides insight to the possible contribution of mismatch in elastic properties to the development of stenosis in vascular reconstructions such as peripheral bypasses.

REFERENCES

1. Brand FN, Abbott RD, Kannel WB. Diabetes, intermittent claudication, and risk of cardiovascular events. The Framingham study. *Diabetes* 1989; 38:504-509.
2. Rutan GH, Kuller LH, Neaton JD, Wentworth DN, McDonald RH, McFate-Smith W. Mortality associated with diastolic hypertension and isolated systolic hypertension among men screened for Multiple Risk Intervention Trial. *Circulation* 1988; 77:504-514.
3. Avolio AP, Deng FQ, Li WQ, et al. Effects of aging on arterial distensibility in populations with high and low prevalence of hypertension: comparison between urban and rural communities in China. *Circulation* 1985; 71:202-210.
4. Ting CT, Chang MS, Wang SP, Chiang BN, Yin FCP. Regional pulse wave velocities in hypertensive and normotensive humans. *Cardiovasc Res* 1990; 24:865-872.
5. Laurent S, London GM, Safar ME. Increased response of the brachial arterial diameter to norepinephrine in essential hypertension: Preliminary report. *J Hypertens* 1986; 4:237-239.
6. Levenson J, Pithois-Merli I, Simon A. Mechanical factors in large artery disease and antihypertensive drugs. *Am J Cardiol* 1990; 66:39C-42C.
7. Levy BI, Benessiano J, Poitevin P, Safar ME. Endothelium-dependent mechanical properties of the carotid artery in Wistar-Kyoto and spontaneously hypertensive rats. *J Hypertens* 1989; 7:S116-S119.
8. Avolio AP, Chen SG, Wang RP, Zhang CL, Li MF, O'Rourke MF. Effects of aging on changing arterial compliance and left ventricular load in a northern chinese urban community. *Circulation* 1983; 68:50-58.
9. O'Rourke MF. Arterial stiffness, systolic blood pressure, and logical treatment of arterial hypertension. *Hypertension* 1990; 15:339-347.
10. Asmar AG, Brunel PC, Pannier BM, Lacolley PJ, Safar ME. Arterial distensibility and ambulatory blood pressure monitoring in essential hypertension. *Am J Cardiol* 1988; 61:1066-1070.
11. Simon AC, Levenson J, Bouthier JD, Maarek B. Haemodynamic basis of early modifications of the large arteries in borderline hypertension. *J Hypertens* 1987; 5:179-184.
12. Scarpello JHB, Martin TRP, Ward JD. Ultrasound measurements of pulse-wave velocity in the peripheral arteries of diabetic subjects. *Clin Sci* 1980; 58:53-57.
13. Oxlund H, Rasmussen LM, Andreassen TT, Heickendorff L. Increased aortic stiffness in patients with type I (insulin-dependent) diabetes mellitus. *Diabetologia* 1989; 32:748-752.
14. Cairns SA, Woodcock JP, Marshall AJ. Early arterial lesions in maturity onset diabetes mellitus detected by an ultrasound technique. *Diabetologia* 1978; 14:107-111.
15. Laogun AA, Gosling RG. In vivo arterial compliance in man. *Clin Phys Physiol Meas* 1982; 3:201-212.
16. Hoeks APG, Brands PJ, Smeets FAM, Reneman RS. Assessment of distensibility of superficial arteries. *Ultrasound Med Biol* 1990; 16:121-128.
17. Reneman RS, Merode van T, Hick P, Muytjens AMM, Hoeks APG. Age-related changes in carotid artery wall properties in men. *Ultrasound Med Biol* 1986; 12:465-471.

CHAPTER 4

SHORT-TERM VARIATION IN ELASTIC PROPERTIES OF A MUSCULAR ARTERY IN HUMANS

L Hofstra, JM Willigers, NC Schaper, FC Huvers, ADM Kester, PJEHM Kitslaar,
APG Hoeks

Adapted from Clinical Science 1994;86:567-574

ABSTRACT

Background. Information on the interaction between mechanical strain and elastic properties of large arteries could give insight in the origin of hemodynamic vascular injury, possibly resulting in formation of atherosclerotic plaques. Intra-individual variation in elastic properties of muscular arteries, measured using non-invasive techniques, has been observed. This variation could be due to changes in blood pressure, since elastic properties of muscular arteries are strongly non-linear.

Methods. The objective of the present study was to evaluate variability in elastic properties of a muscular artery over a time period up to one hour. Furthermore the relation between changes in elastic properties and alterations in blood pressure was studied. With a non-invasive high resolution ultrasound technique diameter and distension of the femoral artery were assessed in repeated measurements in 10 healthy volunteers (mean age: 27 ± 2.9 years) in resting conditions during 40-60 minutes. Simultaneously intra-arterial blood pressure of the contralateral common femoral artery and heart rate were recorded. From distension and pulse pressure, distension pressure ratio (DPR) was calculated, a measure for the compliance. For each parameter the variation coefficient (VC) was computed.

Results. In all subjects a considerable variation was found in distension (mean VC 14.8%) but not in pulse pressure (mean VC 3.7%). The expected positive correlation between pulse pressure and distension was absent in 8 subjects. As a consequence, large intra-individual variation was found in DPR (mean VC 15.3%). When plotted as function of time, gradual fluctuations in DPR were observed. In the majority of the subjects no correlation between diastolic pressure and diameter was observed. Therefore the non-linear relation between diastolic diameter and distension cannot explain the observed variation in DPR.

Conclusions. In resting conditions femoral artery elastic properties in humans show large intra-individual variation with time. This variation cannot be explained by alterations in diastolic diameter, suggesting active vessel wall behavior to be responsible

for the variability. Due to the variability of DPR in this large artery, a single measurement cannot give a proper characterization of the elastic properties.

INTRODUCTION

Due to their visco-elastic properties, large conduit arteries play an important role in buffering phasic changes in pressure and flow during the cardiac cycle¹⁻³. Comprehension of the interaction between mechanical strain and elastic characteristics of the arterial wall is essential to get more insight in the aetiology of arterial degenerative disease due to hemodynamic vascular injury⁴⁻⁸. Variability in elastic behavior of large muscular arteries measured with the use of non-invasive techniques has been demonstrated. A wide range of individual values for stiffness has been reported for both the brachial and femoral artery⁹. Moreover, significant intra-individual variations in elasticity of the aorta and the brachial artery, calculated from pulse-wave velocity, have been reported^{10,11}. These observed variations can be explained by the non-linear relationship between blood pressure and elastic properties¹²⁻¹⁴. An elevation of mean blood pressure leads to a decrease in compliance of an artery. Intra-arterial measurements of blood pressure in the brachial artery demonstrated considerable variation, even in resting conditions¹⁵⁻¹⁷, and could possibly explain the observed variation in elastic behavior. Another explanation could be the modification of elastic properties by vaso-active substances or the central nerve system^{13,18}. Recent research revealed that the dimensions of large muscular arteries, like the common femoral artery, are influenced by vaso-active substances such as angiotensin, norepinephrin and atrial natriuretic factor¹⁹⁻²⁴. Furthermore, there is evidence of the existence of a local renin-angiotensin system (RAS) in large arteries^{19,25,26}. An increase in compliance of the carotid artery was found after the local application of ACE-inhibitors²⁷. Different pressure compliance curves were found for the same segment of the femoral artery in one dog measured several times during one month under the same conditions¹³. This indicates that spontaneous changes in elastic behavior of a large muscular vessel can occur. So far it is unclear whether the reported variation in elastic properties of large arteries can be attributed solely to alterations in blood pressure or is the result of changes in vasomotor tone.

The present study was designed to evaluate variability in elastic behavior of a large muscular artery in resting condition in man. Furthermore, the effect of alterations in blood pressure on elastic properties was studied. Repeated measurements of diameter and distension of the femoral artery were performed during 45 minutes and simultaneously blood pressure was recorded in a beat to beat fashion. In this period considerable spontaneous variation in pressure can be expected¹⁵⁻¹⁷. The femoral artery was chosen, because this vessel is prone to develop atherosclerotic lesions. Moreover, this part of the arterial tree is easily accessible for as well the ultrasound technique used as the intra-arterial measurement of blood pressure.

MATERIALS AND METHODS

Vessel Wall Doppler Tracking (VWDT)^{28,29} was used to measure diameter and distension. The VWDT-system consists of an echo Doppler system (Ultramark IV, Advanced Technology Laboratories) with a 7.5 MHz sector probe, a memory module and a Personal Computer. The vessel of interest is visualized in B-mode to obtain anatomical and structural information. A cursor line, representing the angle of investigation is positioned perpendicular to the arterial walls. A number (2048) of sequential non-processed RF lines are digitized and temporarily stored in the memory module for processing afterwards. The signal as function of depth of the first digitized RF line is displayed on the computer screen. From the shape and the position in depth of the reflections the operator can identify the anterior and posterior walls. Two sample volumes, indicated by boxes, are placed on the anterior and posterior wall, respectively. The distance between both boxes reflects the initial diameter of the artery (Figure 1).

Figure 1: The digitized RF signal of the first line acquired. From the shape and position in depth the operator can identify the position of the anterior and posterior wall. The walls are marked with (white) boxes representing the sample volumes used by the tracking program. The distance between the boxes reflects the initial diameter.

Analysis of the signal behavior within the sample volumes as function of time yields the change in vessel wall position (displacement waveform). The position of the sample volume is updated according to the detected displacement to ensure that for processing the same signal segment will be evaluated. The difference between the displacement waveforms of the anterior and posterior walls gives the change in diameter (distension) as function of time. The process is triggered by an electrocardiograph (ECG) signal, which facilitates the detection of the maximum (systolic) and minimum (diastolic) arterial diameter within a cardiac cycle. The difference between maximum and minimum diameter is considered as distension. The result of a typical recording of a distension waveform is shown in figure 2. In the present study the averaged values of diameter and distension values during a recording time of 4 seconds were used.

Figure 2: Displacement waveforms of the anterior (top tracing) and posterior (middle tracing) wall of the common femoral artery. This results in the distension waveform (bottom tracing), where a cross indicates the occurrence of an ECG trigger.

Intra-arterially blood pressure data were obtained simultaneously in the contralateral common femoral artery at a site corresponding with the position of the ultrasound measurements. We decided not to perform these measurements at the ipsilateral side, since introduction of a catheter could influence the elastic properties of the artery. A 20 Gauge rigid catheter (Arrow, Reading, USA) was introduced in the contralateral femoral artery after local anesthesia with 2% lidocaine. Via a fluid filled line, the catheter was connected with a pressure transducer (Baxter Pressure Set Uniflow 43-600F, Uden, The Netherlands). The resonance frequency of this system ranges from

18-20 Hz, which is sufficient to provide acceptable dynamic responses. During the measurements of the distension, pressure data were simultaneously recorded and digitized at a sample frequency of 250 Hz.. Pulse pressure was defined as the difference between the maximum and minimum value of pressure within 300 ms of the ECG-trigger (i.e. difference between peak systolic and end diastolic pressure). From distension (ΔD in μm) and simultaneously recorded pulse pressure (ΔP in mmHg), the distension pressure ratio (DPR) was calculated using the following equation:

$$\text{Distension Pressure Ratio (DPR)} = \frac{\Delta D}{\Delta P} \quad (1)$$

This simple equation gives a measure for the compliance of the vessel.

The study was performed on 10 healthy volunteers (9 male, 1 female, age 27 (± 2.9) (mean ($\pm$ SD)) without a history of cardiovascular disease. All subjects were normotensive and clinically free of vascular symptoms. None of them used medication affecting cardiovascular behavior. The study was approved by the medical ethical committee of our hospital and each subject gave written informed consent.

STUDY DESIGN

All subjects were investigated at 14.00 am after they had used a light meal. They abstained from nicotine and caffeine before the investigation. The ultrasound investigations were carried out in a climatized room (24°C) with the subjects in the supine position. After an acclimatization phase of 20 minutes, the subjects were measured during 38-70 minutes (mean: 52 $\pm$ 10 min.) with a mean sample interval of 77 $\pm$ 15 seconds. During the measurements, the echo transducer was constantly kept in place by the operator to avoid changes in measured values due to local inhomogeneities in compliance³⁰. The common femoral artery dimensions were measured 2-3 centimeters proximal to the femoral bifurcation. All measurements were performed by the same operator.

ANALYSIS

To calculate the correlation between the different parameters, the Pearson correlation coefficient was used. A p-value < 0.05 was accepted as level of statistical significance. The values for pulse pressure and distension for each subject are presented as mean and standard deviation. As a measure of variation, the variation coefficient of the parameters was calculated using the following equation:

$$\text{Variation Coefficient (VC)} = 100 * \text{Standard Deviation} / \text{Mean} \quad (\%) \quad (2)$$

RESULTS

In all subjects a wide range of values for distension was found. (Table I). The large variation in distension of the common femoral artery is illustrated by a mean variation coefficient of 14.8% for the whole group. In most subjects the variation coefficient of distension was approximately four times the variation coefficient of pulse pressure (mean 3.7%).

Table I

Subj.	n	Pulse pressure (mmHg)			Distension (µm)			DPR (µm/mmHg)		
		mean	sd	vc(%)	mean	sd	vc(%)	mean	sd	vc(%)
1	35	73.5	2.4	3	391	51.6	13	5.32	0.68	13
2	39	63.2	1.6	3	221	18.9	9	3.50	0.32	9
3	31	55.6	2.9	5	540	112.9	21	9.70	1.88	19
4	45	60.9	3.0	5	396	49.3	13	6.51	0.79	12
5	55	72.3	1.4	2	354	50.6	14	4.88	0.72	15
6	54	73.0	1.3	2	536	53.3	10	7.35	0.81	11
7	40	60.3	1.5	2	361	60.1	17	6.00	1.09	18
8	34	61.9	4.9	8	484	83.6	17	7.90	1.71	22
9	48	66.8	2.1	3	364	58.7	16	5.45	0.89	16
10	29	49.4	1.9	4	209	38.9	19	4.23	0.77	18
MEAN		63.7	2.3	3.7	386	57.8	14.8	6.14	1.04	15.3

As an example, the relation between pulse pressure and distension of subject 1 is displayed in figure 3. As is illustrated in figure 3, a single value of pulse pressure can lead to different values of distension. No correlation ($r = 0.25$, 95% confidence interval: -0.09 - 0.54) was found between pulse pressure and distension in this subject. Only in 2 of the remaining subjects a positive correlation between pulse pressure and distension was found. In 3 subjects a negative correlation and in 5 subjects no correlation was found (Table II).

To evaluate a possible shift in the pressure diameter relation due to changes in diastolic pressure, diastolic pressure and diastolic diameter were correlated. In only 2 subjects could a positive correlation between these parameters be observed. In 1 subject a negative correlation was found and in 7 no correlation was observed (Table II). This indicates that an increase in diastolic pressure does not necessarily lead to an increase in diastolic diameter. To evaluate the effect of changes in diastolic pressure on elastic properties, the correlation between diastolic pressure and distension was

calculated. In 7 subjects a negative correlation between diastolic pressure and distension was observed and in 3 subjects no correlation between these parameters was demonstrated.

Table II

Subj.	ρ (ΔP , dist)			ρ (Pdia, diam)			ρ (Pdia, dist)			ρ (diam, dist)		
	ρ	95% conf. interval		ρ	95% conf. interval		ρ	95% conf. interval		ρ	95% conf. interval	
1	0.25	-0.09	0.54	0.15	-0.20	0.46	-0.45	-0.68	-0.13	-0.60	-0.78	-0.33
2	-0.09	-0.39	0.23	-0.29	-0.55	0.03	-0.26	-0.53	0.06	0.03	0.29	-0.34
3	0.41	0.06	0.67	-0.21	-0.53	0.16	-0.25	-0.55	0.12	-0.35	-0.62	0.01
4	0.30	0.15	0.43	0.09	-0.07	0.24	-0.46	-0.57	-0.33	0.12	-0.04	0.27
5	-0.14	-0.28	0.00	0.62	0.52	0.70	-0.51	-0.61	-0.39	-0.28	-0.41	-0.14
6	-0.57	-0.73	-0.35	-0.30	-0.52	-0.03	-0.77	-0.86	-0.63	0.32	0.05	0.54
7	-0.26	-0.41	-0.10	0.16	0.00	0.32	-0.53	-0.64	-0.40	-0.35	-0.48	-0.19
8	-0.24	-0.40	-0.07	0.33	0.16	0.48	-0.48	-0.60	-0.33	-0.71	-0.79	-0.61
9	0.01	-0.14	0.15	-0.15	-0.29	0.00	-0.14	-0.28	0.01	-0.60	-0.69	-0.49
10	0.16	-0.22	0.50	0.07	-0.30	0.43	-0.41	-0.76	-0.05	-0.61	-0.80	-0.31

Figure 3: The relation between pulse pressure and distension of subject 1. This graph shows that an equal pulse pressure can lead to different distensions. The correlation coefficient between the parameters is not significant for this subject ($r = 0.25$, 95%confidence-interval: -0.09 - 0.54).

Correcting alterations in distension for pulse pressure changes (distension pressure ratio, equation 1) did not result in the expected reduction of the variation. For the whole group a mean variation coefficient for DPR of 15.3% was found. A typical graph of DPR as a function of time of subject 5 is shown in figure 4. The graph suggests that the variation is not at random but is characterized by a gradual fluctuation with time. Between heart rate, a marker for sympathetic activity, and distension no correlation was found (data not shown).

Figure 4: The distension pressure ratio (DPR, eq. 1) values of the femoral artery of subject 5 plotted as function of time. The pattern suggests a gradual fluctuation in time.

DISCUSSION

The goal of this study was to evaluate variability in elastic behavior of the common femoral artery in relation to changes in blood pressure. Spontaneous intra-individual variation in distension of this artery was found in all subjects, illustrated by an average variation coefficient of 14.8%, while the variation in pulse pressure was considerably lower (mean VC 3.7%).

Part of the variation in distension possibly originated from intra-observer variation or changes in experimentation circumstances. To eliminate this variation as much as possible, we optimized operator as well as subject conditions. The measurements were performed by one and the same experienced operator. During the whole length of the experiment, the transducer was constantly positioned at the same segment of the common femoral artery, 2-3 centimeter proximal from the femoral bifurcation. This measure was necessary to avoid changes in compliance due to local inhomogeneities

in elastic properties³⁰. The use of B-mode imaging facilitated the positioning of the cursor line using the bifurcation as a landmark. The measurements were performed in a quiet and climatized room (24°C) after an acclimatization period of at least 20 minutes. It is most likely that the variation in DPR, as reported in this study, originates from alterations in local mechanical behavior of the vessel wall rather than changes in circumstances or intra-observer variation.

One of the limitations of this study might be the correlation of blood pressure of the contralateral femoral artery with distension values of the ipsilateral femoral artery. However, it is likely that the values of pressure and the configuration of the pressure signal are equal at both sites, since intensity and timing of pulse wave reflection will have the same effect at both sites.

In previous reports the change in cross-section per unit of pressure, the cross-sectional compliance, was used as a measure for compliance. Diameter is included in this equation³⁰. In our data hardly any variation was observed in diameter (VC: 3%, data not shown), indicating that the changes in elastic properties of the femoral artery result from changes in distension over cardiac cycles rather than diastolic diameter changes. Therefore, it was decided to use a simple equation to represent the compliance, the distension pressure ratio.

Since pulse pressure is the force leading to the pulsatile distension of the arterial wall a positive correlation is expected between these two parameters at equal diameter values. In only 2 of the 10 subjects a positive correlation between pulse pressure and distension was found and in 3 subjects even a negative correlation was observed. Consequently, a large intra-individual variation was found in the distension pressure ratio (DPR) calculated from distension and pulse pressure, as illustrated in figures 3 and 4. This observation can be explained by the non-linear relationship between pressure and distension of an artery^{13,14,31,32} as is diagrammatically represented in figure 5. This figure shows that an equal pulse pressure occurring at different levels of diastolic pressure can lead to quite different values of distension. In our data, only for a minority of the subjects investigated, could a positive correlation between diastolic diameter and diastolic pressure be observed. Thus, an increase in diastolic pressure is not necessarily followed by an increase in diastolic diameter, indicating that a shift in the pressure distension curve does not occur. This suggests that passive changes in elastic behavior due to changes in blood pressure, cannot explain the observed lack of correlation between pulse pressure and distension. Therefore, the large observed variation in DPR cannot be attributed to changes in blood pressure.

Figure 5: Diagrammatic representation of the relation between pressure and diameter. Since this relation is non-linear an equal pulse pressure may lead to a different distension (d_1 , d_2) for a varying diastolic pressure (PD_1 , PD_2).

Figure 6: Hypothetical pressure-diameter curves of the same vessel at two different moments. Due to a different elastic behavior, equal pulse and diastolic pressure values can lead to different distension values.

Figure 7: Pressure-distension curves obtained for two different heart beats of subject 1. Only the leading part of the curve (i.e. the early systolic phase) is presented in the graph. Despite equal diastolic pressure, diastolic diameter and pulse pressure, distinct curves are found, indicating variation in elastic behavior of the vessel at different moments.

Differences in pressure-compliance curves for a femoral artery, obtained by repeated measurements in a single dog under established conditions, have been reported by Megerman¹³. The consequence of a change in the pressure distension relation is illustrated in figure 6. This figure shows that for two different heartbeats unchanged diastolic and pulse pressure values can lead to different distension values due to a change in the pressure distension relation. Changes in pressure compliance curves, indicating alterations in vascular tone, can fully explain the lack of a positive correlation between pulse pressure and distension. Two instantaneous pressure distension curves of subject 1 at two different moments are depicted in figure 7, indicating that the pressure distension relation of the femoral changes in a relatively short period of time (interval between the curves: 5 minutes).

Alterations in vascular tone³² can be the effect of a varying influence of vaso-active substances^{13,33,34} and/or the central nervous system^{18,24}. The endothelial cell layer plays an important role in local regulation of vascular tone^{19,21,35-40}. Increase in shear forces, exerted on the endothelium, leads to EDRF (NO) release^{41,42} and to an inhibition of adrenergic constriction⁴³. Both mechanisms lead to vasodilation. It has been hypothesized, and it is now widely accepted, that arteries tend to maintain the shear forces that are sensed by the endothelial layer, at a constant value^{44,45}. An increase in shear stress leads to a structural increase in artery diameter resulting in a return of the shear forces to base line values. The alterations in elastic properties, as found in our study, might

be a functional regulation mechanism to maintain shear stress values at a constant level. This is supported by our finding that in diameter hardly any variation was found. According to the Hagen-Poiseuille formula, mean shear stress is strongly diameter dependent ($\tau \sim 1/d^4$), and therefore slight changes in diameter have a profound effect on shear stress. Of interest in this respect is the negative correlation between diastolic pressure and distension found in all subjects, while no correlation was found between diastolic pressure and diastolic diameter. This indicates, that in response to diastolic pressure increase a contraction of smooth muscle cells occurs, to prevent an increase in diameter. At the same time, this contraction leads to stiffening of the vessel explaining the negative correlation between diastolic pressure and distension.

Another important control mechanism is the local renin-angiotensin system (RAS)^{19,46}. The central RAS has a short term effect on changes in blood pressure and vascular tone. The local RAS probably is of importance in long term regulation of pressure and vasomotor tone²⁵. The gradual fluctuation of DPR as depicted in figure 4, suggests a short term regulation mechanism.

The extensive influence of the sympathetic nerve system on brachial artery distensibility has been demonstrated in a group (n=10) of normal subjects⁴⁷. A decrease in distension of the brachial artery was found after applying contra-lateral cooling (cold pressor). However, an increase in pulse pressure was observed, measured at the middle finger of the ipsilateral hand, indicating a decrease in compliance. Kawasaki also found an increase in stiffness of the brachial artery after the application of cold at the contra-lateral hand⁹. Heart rate can be used as a measure of sympathetic activity⁴⁸. In the present data we found no correlation between heart rate and DPR, suggesting that the sympathetic nerve system does not play a crucial role in the variation of the elastic properties of the common femoral artery in resting conditions.

A recent study reported a 40% increase in compliance of the radial artery in hypertensive subjects after the infusion of papaverine⁴⁹. These data support our observation, that in a muscular artery compliance values can undergo considerable changes. The new information of the present study is that elastic properties of the common femoral artery, represented by the DPR, change considerably in resting conditions, without the administration of vaso-active substances. The implication of this finding is that this large muscular artery in a healthy subject cannot be characterized by a single measurement of distension or DPR. Another way to arrive at a better characterization of the elastic properties of this type of artery might be by the use of provocation tests, like the administration of nitroglycerine, the cold pressor test, or a reactive hyperaemia test⁴⁹. This might give a better insight in the mechanical response of a large muscular artery and probably results in a better characterization of the vessel.

In conclusion, our data demonstrate that the elastic properties of the common femoral artery cannot be considered constant in a young population investigated under resting conditions. Changes in blood pressure cannot explain the observed variation. This suggests that active mechanisms play a role in these changes rather than passive mechanisms. The physiologic implications of the observed variation in elastic properties are unclear, and cannot be concluded from this study. More research in this field is necessary to elicit the function of active mechanical behavior⁵⁰ of large muscular arteries.

ACKNOWLEDGEMENT

We thank Dr G. Ramsay, Department of General Surgery, Academic Hospital Maastricht, for providing us with advice in writing the manuscript.

REFERENCES

1. Cox RH. Blood flow and pressure propagation in the canine femoral artery. *J Biomech* 1970; 3:131-149.
2. Klanchar M, Tarbell JM, Wang DM. In vitro study of the influence of radial wall motion on shear stress in an elastic tube model of the aorta. *Circ Res* 1990; 66:1624-1635.
3. O'Rourke MF. Steady and pulsatile energy losses in the systemic circulation under normal conditions and in simulated arterial disease. *Cardiovasc Res* 1967; 1:313-326.
4. Nerem RM. Vascular fluid mechanics, the arterial wall, and atherosclerosis. *J Biomech Eng* 1992; 114:274-282.
5. Nerem RM, Girard PR. Hemodynamic influences on vascular endothelial biology. *Toxicologic Pathology* 1990; 18:572-582.
6. Bondjers G, Glukhova M, Hansson GK, Postnov YV, Reidy MA, Schwartz SM. Hypertension and atherosclerosis. Cause and effect, or two effects with one unknown cause? *Circulation* 1991; 84:VI2-VI6.
7. Karino T, Goldsmith HL. Role of blood cell-wall interactions in thrombogenesis and atherogenesis: a microrheological study. *Biorheology* 1984; 21:587-601.
8. Ross R. The pathogenesis of atherosclerosis - An update. *New Eng J Med* 1986; 314:488-499.
9. Kawasaki T, Sasayama S, Yagi SI, Asakawa T, Hirai T. Non-invasive assessment of the age related changes in stiffness of major branches of the human arteries. *Cardiovasc Res* 1987; 21:678-687.
10. Wright JS, Cruickshank JK, Kontis S, Dore C, Gosling RG. Aortic compliance measured by non-invasive doppler ultrasound: description of a method and its reproducibility. *Clin Sci* 1990; 78:463-468.
11. Kontis S, Gosling RG. On-line doppler ultrasound measurement of aortic compliance and its repeatability in normal subjects. *Clin Phys Physiol Meas* 1989; 10:127-135.
12. O'Rourke MF. Arterial stiffness, systolic blood pressure, and logical treatment of arterial hypertension. *Hypertension* 1990; 15:339-347.
13. Megerman J, Hasson JE, Warnock DF, L'Italien GJ, Abbott WM. Noninvasive measurements of nonlinear arterial elasticity. *Am J Physiol* 1986; 250:H181-H188.
14. Hayoz D, Rutschmann B, Perret F, et al. Conduit artery compliance and distensibility are not necessarily reduced in hypertension. *Hypertension* 1992; 20:1-6.
15. Imholz BPM, Settels JJ, Meiracker van der AH, Wesseling KH, Wieling W. Non-invasive continuous finger blood pressure measurement during orthostatic stress compared to intra-arterial pressure. *Cardiovasc Res* 1990; 24:214-221.
16. Idema RN, Meiracker van den AH, Imholz BPM, et al. Comparison of Finapres non-invasive beat-to-beat finger blood pressure with intrabrachial artery pressure during and after bicycle ergometry. *J Hypertens* 1989; 7:S58-S59.
17. Parati G, Casadei R, Gropelli A, Di Rienzo M, Mancina G. Comparison of finger and intra-arterial blood pressure monitoring at rest and during laboratory testing. *Hypertension* 1989; 13:647-655.
18. Vanhoutte PM, Verbeuren TJ, Webb RC. Local modulation of adrenergic neuroeffector interaction in the blood vessel wall. *Physiol Rev* 1981; 61:151-215.
19. Dzau VJ, Safar ME. Large conduit arteries in hypertension: role of the vascular renin-angiotensin system. *Circulation* 1988; 77:947-954.
20. Simon AC, Levenson JA, Levy BY, Bouthier JE, Peronneau PP, Safar ME. Effect of nitroglycerin on peripheral large arteries in hypertension. *Br J Clin Pharmac* 1982; 14:241-246.
21. Levy BI, Benessiano J, Poitevin P, Safar ME. Endothelium-dependent mechanical properties of the carotid artery in Wistar-Kyoto and spontaneously hypertensive rats. *J Hypertens* 1989; 7:S116-S119.

22. Levenson J, Simon AC, Bouthier JD, Benetos A, Safar ME. Post-synaptic alpha-blockade and brachial artery compliance in essential hypertension. *J Hypertens* 1984; 2:37-41.
23. Laurent S, London GM, Safar ME. Increased response of the brachial arterial diameter to norepinephrine in essential hypertension: Preliminary report. *J Hypertens* 1986; 4:237-239.
24. Bolton TB. Mechanisms of action of transmitters and other substances on smooth muscle. *Physiol Rev* 1979; 59:606-695.
25. Dzau VJ. Short- and long-term determinants of cardiovascular function and therapy: contribution of circulating and tissue renin-angiotensin systems. *J Cardiovasc Pharmacol* 1989; 14:S1-S5.
26. Krieger JE, Dzau VJ. Molecular biology of hypertension. *Hypertension* 1991; 18:13-17.
27. Levy BI, Benessiano J, Poitevin P, Safar ME. Angiotensin-converting enzyme inhibition and compliance of the carotid artery in normotensive and hypertensive rats. *J Hum Hypertens* 1989; 3:S57-S62.
28. Hoeks APG, Brands PJ, Smeets FAM, Reneman RS. Assessment of distensibility of superficial arteries. *Ultrasound Med Biol* 1990; 16:121-128.
29. Hofstra L, Ermers EJM, Langeveld APM, Hoeks APG, Kitslaar PJEHM. Vessel Wall Doppler Tracking: A new technique to assess vessel wall compliance. *J Vasc Tech* 1993; 17:65-68.
30. Reneman RS, Merode van T, Brands PJ, Hoeks APG. Inhomogeneities in arterial wall properties under normal and pathological conditions. *J Hypertens* 1992; 10:S35-S39.
31. Armentano R, Simon A, Levenson J, Chau NP, Megnien JL, Pichel R. Mechanical pressure versus intrinsic effects of hypertension on large arteries in humans. *Hypertension* 1991; 18:657-664.
32. Dobrin PB. Mechanical properties of arteries. *Physiol Rev* 1987; 58:397-460.
33. Dobrin PB, Rovick AA. Influence of vascular smooth muscle on contractile mechanics and elasticity of arteries. *Am J Physiol* 1969; 217:1644-1651.
34. Vatner SF, Pagani M, Manders WT, Pasipoularides AD. Alpha adrenergic vasoconstriction and nitroglycerin vasodilation of large coronary arteries in the conscious dog. *J Clin Invest* 1980; 65:5-14.
35. Rubanyi GM, Freay AD, Kauser K, Johns A, Harder DR. Mechanoreception by the endothelium: Mediators and mechanisms of pressure- and flow-induced vascular responses. *Blood Vessels* 1990; 27:246-257.
36. Olesen SP, Clapham DE, Davies PF. Hemodynamic shear stress activates a K⁺ current in vascular endothelial cells. *Nature* 1988; 331:168-170.
37. Furchgott RF, Vanhoutte PM. Endothelium-derived relaxing and contracting factors. *FASEB J* 1989; 3:2007-2018.
38. Asmar AG, Brunel PC, Pannier BM, Lacolley PJ, Safar ME. Arterial distensibility and ambulatory blood pressure monitoring in essential hypertension. *Am J Cardiol* 1988; 61:1066-1070.
39. Lüscher TF. Imbalance of endothelium-derived relaxing and contracting factors. A new concept in hypertension? *Am J Hypertens* 1990; 3:317-330.
40. Ralevic V, Milner P, Hudlicka O, Kristek F, Burnstock G. Substance P is released from the endothelium of normal and capsaicin-treated rat hind-limb vasculature, in vivo, by increased flow. *Circ Res* 1990; 66:1178-1183.
41. Cooke JP, Rossitch E, Andon NA, Loscalzo J, Dzau VJ. Flow activates an endothelial potassium channel to release an endogenous nitrovasodilator. *J Clin Invest* 1991; 88:1663-1671.
42. Buga GM, Gold ME, Fukuto JM, Ignarro LJ. Shear stress-induced release of nitric oxide from endothelial cells grown on beads. *Hypertension* 1991; 17:187-193.
43. Tesfamariam B, Cohen RA. Inhibition of adrenergic vasoconstriction by endothelial cell shear stress. *Circ Res* 1988; 63:720-725.

44. Langille BL, O'Donnell F. Reductions in arterial diameter produced by chronic decreases in blood flow are endothelium-dependent. *Science* 1986; 231:405-407.
45. Zarins CK, Zatina MA, Giddens DP, Ku DN, Glagov S. Shear stress regulation of artery lumen diameter in experimental atherogenesis. *J Vasc Surg* 1987; 5:413-420.
46. Dzau VJ, Gibbons GH, Pratt RE. Molecular mechanisms of vascular renin-angiotensin system in myointimal hyperplasia. *Hypertension* 1991; 18:III100-III105.
47. Kurvers HAJM, Hofstra L, Hoeks APG, Wildenberg Van De FAJM, Kitslaar PJEHM, Jacobs MJHM. Sympathetic dysfunction in Sudeck's dystrophy assessed by Vessel Wall Echo Tracking. *Eur J Clin Invest* 1992; 22:A8.
48. Flückiger JP, Gremaud G, Waeber B, et al. Measurement of sympathetic nerve activity in the unanesthetized rat. *J Appl Physiol* 1989; 67:250-255.
49. Trazzi S, Omboni S, Santucci C, Parati G, Mancina G. Variability in arterial diameter and compliance: compliance modulation reserve. *J Hypertens* 1992; 10:S41-S43.
50. Kon V, Harris RC, Ichikawa I. A regulatory role for large vessels in organ circulation. *J Clin Invest* 1990; 85:1728-1733.

CHAPTER 5

ASSESSMENT OF INHOMOGENEITIES IN ELASTIC PROPERTIES OF IN SITU AND REVERSED SAPHENOUS VEIN GRAFTS IN HUMANS

L Hofstra, DCJJ Bergmans, APG Hoeks, JHM Tordoir, PJEHM Kitslaar
Adapted from the European Journal of Vascular Surgery 1994; 8: 670-676

Winner of the European Society of Vascular Surgery prize 1993
for the best clinical paper

ABSTRACT

Background. In vascular surgery controversy remains concerning the optimal technique for the implantation of the saphenous vein graft to bypass peripheral atherosclerotic lesions. In comparison with the reversed graft, the in situ graft provides theoretical hemodynamic advantages due to its natural taper and the preservation of endothelial cell function.

Methods. Elastic properties of newly implanted in situ (IS, n=11) and reversed (RV, n=21) saphenous grafts were studied with the use of Vessel Wall Doppler Tracking. From diameter, diameter change and simultaneously recorded pulse pressure, distensibility coefficient (DC) representing the intrinsic elastic properties and compliance coefficient (CC), a hemodynamic parameter, were calculated. In order to obtain a mechanical profile, the parameters were assessed in defined sites in the graft and native arterial system.

Results. In the RV group the reversal of the grafts resulted in a difference in diameter around the proximal anastomoses (7.4 vs 4.3 mm, $p<0.01$) and from proximal to distal (4.3 vs 5.9 mm, $p<0.01$) in the grafts; around the distal anastomoses no differences in diameter were observed. Due to the natural taper of the in situ grafts, diameter decreased from proximal to distal in the grafts (4.5 vs 3.2 mm, $p<0.05$) and no size differences were found around the anastomoses. In the RV group a decrease was observed in DC from proximal to distal in the grafts. Around the anastomoses no differences in DC were apparent. In the IS group no change in DC was found from proximal to distal in the grafts; around the distal anastomoses a decrease in DC was observed. In the RV group a decrease in CC, representing the hemodynamic capacity, was observed in the proximal anastomoses (0.25 vs 0.09 mm²/kPa, $p<0.01$). From proximal to distal in the graft and around the distal anastomoses no differences were

found in hemodynamic capacity. In the IS group no change in hemodynamic capacity was observed around the proximal anastomoses and distal anastomoses. However, a decrease in the value of this parameter from proximal to distal in the grafts was found (0.07 vs 0.04 mm²/kPa, $p < 0.05$).

Conclusions. These results demonstrate that the elastic properties of newly inserted human saphenous vein grafts are comparable with native artery elastic properties. In contrast to the in situ technique, a mismatch in hemodynamic capacity is apparent around the proximal anastomoses using the reversed technique, indicating hemodynamic benefit in proximal anastomoses of in situ grafts. However, an important decrease in hemodynamic capacity is present in the in situ graft, whereas this parameter remains stable in the reversed grafts, suggesting intrinsic hemodynamic advantage in reversed grafts.

INTRODUCTION

The autologous saphenous vein is the material of choice to bypass atherosclerotic lesions in the superficial femoral, popliteal and crural arteries. Knowledge on the mechanical characteristic of grafts could be useful for various reasons. Controversy exists concerning the optimum technique for the implantation of these grafts¹⁻⁵. In theory, the in situ technique has advantages over the reversed technique. Due to its natural taper, the in situ graft offers a better diameter match around both the proximal and distal anastomoses as compared to the reversed graft, probably resulting in a better management of pulsatile flow. Total graft performance might therefore be better in in situ grafts leading to an improvement of distal perfusion⁶. Another important advantage of the in situ technique may be the protection of endothelial cell function due to the preservation of the vasa vasorum⁷. An intact endothelium inhibits platelet adherence and could maintain better mechanical control due to the release of relaxing factors⁸⁻¹⁰. Investigation of the mechanical characteristics of in situ and reversed grafts may verify these claims in favour of the in situ technique.

A second motive to study graft mechanical characteristics is the alleged relation between intimal hyperplasia and compliance mismatch around the anastomoses. Compliance mismatch between artery and graft leads to a region of excessive mechanical stretch, possibly resulting in wall injury^{11,12}, a major event in the initiation of intimal hyperplasia¹³. In addition, a mismatch in compliance leads to flow disturbance and flow separation, both associated with increased intimal thickening^{14,15}. Abbot demonstrated a decrease in patency rates in bypasses with low compliance in a canine model¹¹. However, occlusions of these grafts were not related to the presence of intimal thickening.

Measurements of compliance of saphenous vein grafts in humans have been performed in the past with the use of an ultra-sonic phase locked echo-tracking system. In a comparison of in situ and reversed saphenous vein grafts, slightly higher compliance values were found in in situ grafts¹. However, in that study as well as in previous reports no information was provided on the definition of the sites of measurements of

compliance¹⁶⁻¹⁸. This could possibly be the result of the limitations of the technique used. Furthermore, no information was provided on the differences in compliance around the anastomoses. This information is essential to get more insight in the influence of compliance mismatch around the anastomoses on the development of intimal hyperplasia. The values for compliance in saphenous vein grafts as found in these studies, were considerably lower as compared to native artery compliance values, implying a poor hemodynamic performance of the saphenous vein graft when subjected to arterial pressure.

Recently a new technique, Vessel Wall Doppler Tracking (VWDT), was developed to assess compliance and distensibility of large vessels and grafts¹⁹. An important advantage of VWDT over other techniques is that anatomical information can be obtained, allowing exact definition of the site of measurement. In addition, measurements of elastic properties of vessels situated as deep as 6 cm below the skin surface can be performed, permitting measurements of reversed bypass grafts in the anatomical positions. The aim of the present study was to assess the mechanical properties of newly inserted *in situ* and reversed saphenous vein grafts at defined sites in the graft and native arteries in order to obtain a mechanical profile. Furthermore, differences in elastic properties around the anastomoses were studied.

MATERIALS AND METHODS

Patients

Eligible for this study were patients undergoing a femoro-distal bypass operation with the use of autologous saphenous vein. Indications for operation were disabling claudication, restpain and/or ulceration. Before the elastic properties of the graft were measured, a Duplex scan was performed to test the proper functioning of the graft. Patients with evidence of graft related stenosis were excluded from this study. The detection of a stenotic lesion was based on generally accepted Duplex criteria^{20,21}. The existence of stenotic lesions may lead to early pulse wave reflections^{22,23}, probably disturbing the determination of a normal mechanical profile, by increasing pulse pressure proximal from the stenotic lesion.

Measurements of the elastic properties

The elastic properties of the graft and host artery were assessed with the use of Vessel Wall Doppler Tracking (VWDT). A detailed description of this system is given elsewhere (chapter 3 and 4)^{19,24}. From distension (in μm), diameter (in mm) and simultaneously recorded blood pressure (Finapres, in kPa) RD, DC and CC were calculated. Relative distension (RD) gives the fractional diameter increase. The distensibility coefficient (DC) represents the local stress-strain relationship of the arterial wall and is a measure for the local intrinsic elastic properties of the vessel. The compliance coefficient represents the capacity of the vessel to store blood volume during a cardiac cycle²⁵. From each recording, containing information of 3-5 heartbeats, the average values of the different parameters were used in this study. The

variation coefficient within the recordings was always less than 5%. In a previously conducted study, we observed short-term variations in the elastic properties of the common femoral artery in young healthy subjects²⁶. The variation coefficients of elastic parameters like DC and CC ranged from 10-15% in this medium-sized muscular artery. These variations are probably caused by the contractile state of the vascular smooth muscle cells and probably contribute to the maintenance of pressure and flow homeostasis. In the common carotid artery, an artery of the elastic type, the variation coefficient was found to be between 5-10%. So far, we do not know whether these short-term variations also occur in diseased vessels or saphenous vein grafts. However, the variations in vascular tone in diseased vessels and saphenous vein grafts are probably limited, and certainly do not exceed variations as observed in healthy muscular arteries. Although a variation coefficient of 5-10% has to be considered, we believe the measurements performed in the grafts are valid.

Figure 1: Sites of measurement in the graft and in- and outflow arterial segments.

The Finapres device has been validated for blood pressure values in the brachial artery of healthy volunteers²⁷. However, a validation study comparing Finapres and intra-arterial blood pressure values at the lower limb in patients with diseased vessels has not been reported so far. We therefore decided to assess the elastic properties of the common femoral artery in a group of patients ($n = 20$) with severe atherosclerotic lesions in the lower limb. DC and CC values were calculated with the use of pulse pressure values obtained at the middle finger using Finapres and intra-arterially

obtained pulse pressure values at the common femoral artery. Consistently, DC and CC values were higher when Finapres values were used in the calculation of these parameters (mean DC 8.1 vs 6.0 10^{-3} /kPa and mean CC: 0.331 vs 0.238 mm²/kPa). Regression analysis revealed an excellent correlation between the two ways to calculate DC and CC (for DC: $r = 0.93$, $p < 0.001$ and for CC: $r = 0.91$, $p < 0.001$). We therefore considered the use of Finapres values in the calculation of DC and CC justified. In each graft measurements were performed proximal (1 cm distal to the proximal anastomosis), in the middle segment, and distal (1 cm proximal to the distal anastomosis). The inflow artery was measured 1 cm proximal to the proximal anastomosis, the outflow artery 1 cm distal to the distal anastomosis (Figure 1).

Patients were measured in the supine position after an acclimatisation period of 20 minutes. All patients gave a written informed consent before entering the study. The study was approved by the medical ethical committee of our Hospital.

ANALYSIS

Data are presented as medians and 25th and 75th percentile intervals. Within the groups the values of RD, DC, CC and diameter measured at the defined sites in the graft and in- and outflow arterial segment, respectively, were compared using the Wilcoxon signed rank test. Since patients were not randomly assigned to either of the groups, a direct comparison of in situ and reversed grafts at the defined sites was not performed. A p-value of less than 0.05 was accepted as level of statistical significance.

RESULTS

The patient characteristics of both groups are summarized in Table I. Surprisingly, in the in situ group most patients were female, in contrast to the reversed group. Of importance is the large fraction of proximal anastomoses on the superficial femoral artery in the in situ group. The values of diameter, RD, DC and CC for the reversed and in situ group are summarized in table II and depicted in figure 2.

Table I: Patient characteristics.

	Reversed VSM (N=21)	In Situ (N=11)
Male	16 (76.2%)	3 (27.3%)
Female	5 (23.8%)	8 (72.7%)
Age (Yrs)	71.1 (SD 8.8)	72.6 (SD 7.3)
Diabetic	3 (14.3%)	2 (18.2%)
Clinical stage :		
II	6 (28.6%)	2 (18.2%)
III	8 (38.1%)	3 (27.3%)
IV	7 (33.3%)	6 (54.4%)
Proximal Anastomosis :		
Common Femoral Artery	14 (66.7%)	3 (27.3%)
Superficial Femoral Artery	7 (33.3%)	8 (72.7%)
Distal Anastomosis :		
Popliteal Artery (above knee)	4 (19.0%)	2 (18.2%)
Popliteal Artery (below knee)	9 (42.9%)	2 (18.2%)
Crural	8 (38.1%)	6 (54.5%)
Pedal	0 (0.0%)	1 (9.1%)

Table II: Values (medians and 25th and 75th percentile interval) for diameter (DI), relative distension (RD), distensibility coefficient (DC) and compliance coefficient (CC) assessed at defined sites in the graft and in- and outflow arterial system for the reversed and in situ grafts.

	Inflow	Proximal	Mid	Distal	Outflow
DI (mm):					
Reversed	7.4 (5.5-9.9)	4.3 (3.2-5.7)	5.3 (4.4-6.0)	5.9 (4.6-6.8)	4.8 (3.2-5.3)
In situ	5.7 (4.8-7.7)	4.5 (3.9-5.6)	4.1 (3.6-5.5)	3.2 (2.6-4.3)	3.5 (2.4-6.8)
RD (%):					
Reversed	2.6 (1.7-3.7)	3.8 (2.1-4.6)	1.9 (1.3-3.5)	1.6 (1.1-2.7)	1.6 (1.1-2.6)
In situ	3.1 (2.5-4.8)	2.9 (2.2-3.5)	4.8 (3.2-6.6)	2.9 (1.7-4.6)	1.7 (0.5-3.4)
DC (10^{-3} /kPa):					
Reversed	5.3 (2.6-7.2)	5.8 (3.8-9.8)	3.7 (2.2-5.3)	2.8 (1.9-4.1)	2.9 (1.7-4.1)
In situ	4.7 (3.4-8.3)	4.2 (3.0-7.9)	7.4 (4.2-11.3)	5.2 (3.2-7.4)	2.5 (0.8-4.4)
CC (mm^2/kPa):					
Reversed	0.26 (0.16-0.71)	0.09 (0.05-0.16)	0.07 (0.05-0.13)	0.07 (0.05-0.09)	0.03 (0.02-0.04)
In situ	0.15 (0.09-0.40)	0.08 (0.06-0.14)	0.11 (0.05-0.28)	0.04 (0.03-0.06)	0.02 (0.01-0.05)

Figure 2: Bar graphs showing the results of measurement of diameter (DI), relative distension (RD), distensibility coefficient (DC) and compliance coefficient (CC) measured at the proximal (Prox), mid (Mid) and distal (Dist) segments of the grafts and in the inflow (Infl) and outflow (Outfl) arterial segments (-- * --, $p < 0.05$, Wilcoxon signed rank test).

Reversed group. The proximal graft diameter (4.3 mm) was smaller as compared to the diameter of the arterial inflow segments (7.4 mm, $p<0.05$). The effect of reversed tapering is illustrated by a gradual increase in diameter from proximal (4.3 mm) to distal (5.9 mm, $p<0.05$) in these grafts. A difference in diameter was observed around the distal anastomoses (5.9 in the distal part of the grafts vs 4.8 mm in the arterial outflow segments, $p<0.05$). No significant differences in RD were found between the proximal graft segments (2.6 %) as compared to the inflow segments (3.8 %). A gradual decrease in RD was found from proximal (3.8%) to distal (1.6%, $p<0.05$) in the grafts. Around the distal anastomoses no significant difference in RD could be observed. For DC a similar pattern was found; no significant difference in DC around the proximal anastomoses, a gradual decrease in DC from proximal ($5.8 \cdot 10^{-3} \text{ kPa}^{-1}$) to distal ($2.8 \cdot 10^{-3} \text{ kPa}^{-1}$, $p<0.05$) in the grafts and no significant difference around the distal anastomoses. The compliance coefficient (CC) was larger in the inflow arterial segments ($0.26 \text{ mm}^2/\text{kPa}$) as compared to the proximal segments of the grafts ($0.09 \text{ mm}^2/\text{kPa}$, $p<0.05$). CC remained stable from proximal to distal in the grafts. Around the distal anastomoses no significant differences in CC were apparent.

In situ group. No significant difference in diameter was observed between the arterial inflow segments (5.7 mm) and the proximal segments of the graft (4.5 mm) but a decrease in diameter was found from proximal (4.5 mm) to distal (3.2 mm, $p<0.05$) in the grafts, illustrating the natural taper of in situ grafts. Around the distal anastomoses no significant size differences were found. Around the proximal anastomoses no significant differences were apparent in RD and DC. In the course of the vein graft an increase in RD was found between the proximal segments of the grafts and the middle segments (2.9 vs 4.8%, $p<0.05$). In the distal segments of the graft the median values for RD and DC were higher as compared to the arterial outflow segments. In CC no significant difference was observed between the proximal segments and the arterial inflow segments. A decrease in CC was apparent from proximal to distal in the grafts (0.08 vs $0.04 \text{ mm}^2/\text{kPa}$, $p<0.05$). Around the distal anastomoses no significant differences in CC were found.

DISCUSSION

The aim of this study was to obtain profiles of elastic characteristics of newly implanted in situ and reversed saphenous vein grafts. The major finding of this study is the proper elastic properties of the saphenous vein, when subjected to arterial pressure. Along the entire length of either the in situ or reversed graft the strain and the intrinsic elastic properties of the saphenous vein, represented by RD and DC respectively, were comparable with native artery elastic properties. This is in contrast with most studies evaluating saphenous vein mechanical function in the arterial system. In a canine model low passive compliance values of saphenous vein grafts, when subjected to arterial pressures, have been reported²⁸. In a study evaluating compliance in in situ and reversed saphenous vein grafts in humans with the use of a phase-locked

echo tracking system the values of compliance were found to be half those of native arteries¹. However, no direct comparison was made between elastic properties of the graft and the native arteries, which could explain the differences between their observations and ours. Lüscher demonstrated a diminished endothelium dependent relaxation to acetylcholine (ACH) in human vein grafts as compared to internal mammary artery response, suggesting endothelial cell dysfunction of grafted saphenous veins²⁹. This could possibly lead to a reduction in endothelium derived relaxing factor release resulting in smooth muscle contraction. A recent study reported a diminished endothelium dependent relaxation to ACH in newly implanted vein grafts indicating endothelial cell dysfunction. However, in that study a preservation of endothelium dependent relaxation to other agents was demonstrated³⁰. The preservation of active control of the endothelium on relaxation of grafts subjected to arterial pressures is in concordance with our finding that the elastic properties of the saphenous vein under these conditions are good. Most of our patients were investigated within 14 days after operation. Whether the proper elastic properties of the saphenous vein can be maintained with time is unclear. A longitudinal study is needed to elucidate this issue^{28,31}.

The assessment of elastic profiles in the reversed and in situ vein grafts provided valuable information. In the reversed vein grafts the reversal resulted in diameter mismatch around the proximal anastomoses and an increase in diameter along the grafts from proximal to distal. For RD and DC, representing the strain and intrinsic elastic properties of the graft respectively, comparable values were found in the proximal segments and the arterial inflow segments. However, RD and DC values of the proximal segments were higher as compared to distal values. This suggests a compensatory mechanism for the small diameter of the proximal segments in order to achieve a sufficient hemodynamic capacity in this part of the graft. This is supported by the finding that along the grafts a remarkable stable pattern was found in CC. This compensatory mechanism could be initiated by an increased flow velocity in the narrow proximal segment of the grafts³². Elevations in shear stress caused by a high flow velocity leads to a release of relaxing factors (e.g. EDRF) by the endothelium³³⁻³⁶. These relaxing factors may change the local intrinsic elastic properties by alterations in vascular smooth muscle tone resulting in higher RD and DC values in the narrow proximal segments as compared to the distal segments.

The mismatch in compliance around the proximal anastomoses of reversed grafts has to be attributed solely to the diameter reduction, since intrinsic elastic properties of the proximal part of the grafts are sufficient. In reports evaluating sites of stenoses in reversed bypass grafts, 40% of stenotic lesions were found in the proximal part of the graft, just distal to the anastomosis^{37,38}. This might be explained by our finding, that an abrupt decrease in cross-sectional compliance is present in the proximal anastomoses of reversed vein grafts. However, a prospective study is needed to clarify the exact relation between compliance mismatch around the proximal anastomoses of reversed vein grafts and occurrence of stenotic lesions at this site. The mismatch in diameter

between the distal graft segments and the outflow arteries is due to the reversed tapering of the saphenous vein. Since no difference in CC was found around the distal anastomoses, the observed diameter mismatch probably will not have major hemodynamic consequences.

In the in situ vein grafts a congruent transition in elastic properties from inflow segment to the proximal segment of the graft was observed. In diameter as well as in RD, DC and CC no significant differences were found between these sites. The strain and intrinsic elastic properties of the proximal graft segments, represented by RD and DC respectively, are comparable with the values of these parameters in the reversed group. This implicates that the gradual transition in CC has to be attributed to a diameter match rather than differences in intrinsic elastic properties. The congruent pattern in CC around the proximal anastomosis suggests hemodynamic benefit. The in situ group demonstrated a decline in CC from proximal to distal in the graft. This is in contrast to the stable pattern of CC in the reversed graft, indicating intrinsic hemodynamic advantage of reversed vein grafts compared to the in situ graft. In both the in situ as the reversed vein graft the total hemodynamic performance of a graft seems to be restricted by its narrowest segment.

Part of the differences in profiles of in situ and reversed grafts in the present study may originate from differences in proximal and distal anastomotic levels between the groups. In the in situ group, the majority of the proximal anastomoses were constructed at the superficial femoral artery level, resulting in a proper diameter match around the proximal anastomosis. Since CC is strongly diameter dependent, this might partly explain the absence of a difference in CC between arterial inflow segment and proximal graft segment in the in situ group. Relatively more of the distal anastomoses in the in situ group were constructed at crural and pedal level, limiting total graft performance.

The use of pulse pressure values obtained with the use of Finapres may be a limitation of the present study. Our validation study revealed consistently higher DC and CC values calculated with the Finapres derived pulse pressures as compared to values calculated using intra-arterially obtained pulse pressures at the common femoral artery in a group of patients with severe atherosclerotic lesions of the lower limb. As a consequence, the values for DC and CC in the present study probably overestimate the actual compliance and distensibility. In the comparison of DC and CC values within the graft, this difference does not play a significant role, since for different sites in the same graft an equal pulse pressure is likely. The main advantage of the Finapres device compared to other non-invasive techniques to measure blood pressure is that beat-to-beat information is provided. In bypass patients, blood pressure may vary considerably due to sudden changes in cardiac performance and therefore beat-to-beat information instead of fragmentarily sphygmomanometric recordings is preferable.

In conclusion, the present study demonstrates that the intrinsic elastic properties of newly inserted saphenous vein grafts are comparable with native artery elastic properties. In contrast to the in situ grafts, compliance mismatch was found around the

proximal anastomoses of reversed grafts which has to be attributed entirely to differences in diameter caused by the reversal of the graft. This finding suggests hemodynamic benefit of the proximal anastomoses of in situ grafts. Along the graft a remarkable stable pattern in compliance was observed in the reversed graft, whereas the in situ graft showed an important decrease in compliance, indicating intrinsic hemodynamic advantage of the reversed graft. Prospective studies are needed to study the effect of inhomogeneities in elastic properties of saphenous vein grafts on graft complications such as intimal hyperplasia.

REFERENCES

1. Harris PL, How TV, Jones DR. Prospectively randomized clinical trial to compare in situ and reversed saphenous vein grafts for femoropopliteal bypass. *Br J Surg* 1987; 74:252-255.
2. Harris RW, Andros G, Dulawa LB, Oblath RW, Apyan R, Salles-Cunha S. The transition to "in situ" vein bypass grafts. *Surg Gynecol Obstet* 1986; 163:21-28.
3. Leather RP, Shah DM, Chang BB, Kaufman JL. Resurrection of the in situ saphenous vein bypass. 1000 cases later. *Ann Surg* 1988; 208:435-442.
4. Buchbinder D, Rolins DL, Verta MJ, et al. Early experience with in situ saphenous vein bypass for distal arterial reconstruction. *Surgery* 1986; 99:350-356.
5. Harris PL, Veith FJ, Shanik GD, Nott D, Wengerter KR, Moore DJ. Prospective randomized comparison of in situ and reversed infrapopliteal vein grafts. *Br J Surg* 1993; 80:173-176.
6. O'Rourke MF. Steady and pulsatile energy losses in the systemic circulation under normal conditions and in simulated arterial disease. *Cardiovasc Res* 1967; 1:313-326.
7. Cambria RP, Megerman J, Abbott WM. Endothelial preservation in reversed and in situ autogenous vein grafts. *Ann Surg* 1985; 202:50-55.
8. Lüscher TF. Endothelial control of vascular tone and growth. *Clin Exp Hypertens* 1990; 12:897-902.
9. Dzau VJ, Safar ME. Large conduit arteries in hypertension: role of the vascular renin-angiotensin system. *Circulation* 1988; 77:947-954.
10. Laurent S, London GM, Safar ME. Increased response of the brachial arterial diameter to norepinephrine in essential hypertension: Preliminary report. *J Hypertens* 1986; 4:237-239.
11. Abbott WM, Megerman J, Hasson JE, L'Italien G, Warnock DF. Effect of compliance mismatch on vascular graft patency. *J Vasc Surg* 1987; 5:376-382.
12. Bassiouny HS, White S, Glagov S, Choi E, Giddens DP, Zarins CK. Anastomotic intimal hyperplasia: Mechanical injury or flow induced. *J Vasc Surg* 1992; 15:708-717.
13. Clowes AW, Reidy MA. Prevention of stenosis after vascular reconstruction: Pharmacologic control of intimal hyperplasia - A review. *J Vasc Surg* 1991; 6:885-891.
14. Stewart SFC, Lyman DJ. Effects of a vascular graft/natural artery compliance mismatch on pulsatile flow. *J Biomech* 1992; 25:297-310.
15. Kohler TR, Kirkman TR, Kraiss LW, Zierler BK, Clowes AW. Increased blood flow inhibits neointimal hyperplasia in endothelialized vascular grafts. *Circ Res* 1991; 69:1557-1565.
16. Hokanson DE, Strandness DE. Stress-strain characteristics of various arterial grafts. *Surg Gynecol Obstet* 1968; 57-60.
17. Hokanson DE, Mozersky DJ, Sumner DS, Strandness DE. A phase-locked echo tracking system for recording arterial diameter changes in vivo. *J Appl Physiol* 1972; 32:728-733.
18. Lye CR, Sumner DS, Hokanson DE, Strandness DE. The transcutaneous measurement of the elastic properties of the human saphenous vein femoropopliteal bypass graft. *Surg Gynecol Obstet* 1975; 141:891-895.
19. Hoeks APG, Brands PJ, Smeets FAM, Reneman RS. Assessment of distensibility of superficial arteries. *Ultrasound Med Biol* 1990; 16:121-128.
20. Buth J, Disselhoff B, Sommeling C, Stam L. Color-flow duplex criteria for grading stenosis in infrainguinal vein grafts. *J Vasc Surg* 1991; 14:716-728.
21. Chang BB, Leather RP, Kaufman JL, Kupinski AM, Leopold PW, Shah DM. Hemodynamic characteristics of failing infrainguinal in situ vein bypass. *J Vasc Surg* 1990; 12:596-600.
22. O'Rourke MF. Systolic blood pressure: arterial compliance and early wave reflection, and their modification by antihypertensive therapy. *J Hum Hypertens* 1989; 3:47-52.
23. Kelly RP, Gibbs HH, O'Rourke MF, et al. Nitroglycerin has more favourable effects on left ventricular afterload than apparent from measurement of pressure in a peripheral artery. *Eur Heart J* 1990; 11:138-144.

24. Hofstra L, Ermers EJM, Langeveld APM, Hoeks APG, Kitslaar PJEHM. Vessel Wall Doppler Tracking: A new technique to assess vessel wall compliance. *J Vasc Tech* 1993; 17:65-68.
25. Reneman RS, Merode van T, Brands PJ, Hoeks APG. Inhomogeneities in arterial wall properties under normal and pathological conditions. *J Hypertens* 1992; 10:S35-S39.
26. Hofstra L, Willigers JM, Huvers FC, et al. Short-term variations in elastic properties of a muscular artery in humans. *Clin Sci* 1993; 50:472-480.
27. Imholz BPM, Settels JJ, Meiracker van der AH, Wesseling KH, Wieling W. Non-invasive continuous finger blood pressure measurement during orthostatic stress compared to intra-arterial pressure. *Cardiovasc Res* 1990; 24:214-221.
28. Dobrin PB, Littooy FN, Golan J, Blakeman B, Fareed J. Mechanical and histological changes in canine vein grafts. *J Surg Res* 1988; 44:259-265.
29. Lüscher TF, Diederich D, Siebeenman R, et al. Difference between endothelium-dependent relaxation in arterial and venous coronary bypass grafts. *New Eng J Med* 1988; 319:462-467.
30. Ishii T, Okadome K, Komori K, Odashiro T, Sugimachi K. Natural course of endothelium-dependent and -independent responses in autogenous femoral veins grafted into the arterial circulation of the dog. *Circ Res* 1993; 72:1004-1010.
31. Abbott WM, Megerman J. Adaptive responses of arteries to grafting. *J Vasc Surg* 1989; 9:377-379.
32. Belkin M, Mackey WC, McLaughlin R, Umphrey SE, O'Donnell TF. The variation in vein graft flow velocity with luminal diameter and outflow level. *J Vasc Surg* 1992; 15:991-999.
33. Cooke JP, Rossitch E, Andon NA, Loscalzo J, Dzau VJ. Flow activates an endothelial potassium channel to release an endogenous nitrovasodilator. *J Clin Invest* 1991; 88:1663-1671.
34. Buga GM, Gold ME, Fukuto JM, Ignarro LJ. Shear stress-induced release of nitric oxide from endothelial cells grown on beads. *Hypertension* 1991; 17:187-193.
35. Olesen SP, Clapham DE, Davies PF. Hemodynamic shear stress activates a K⁺ current in vascular endothelial cells. *Nature* 1988; 331:168-170.
36. Rubanyi GM, Freay AD, Kauser K, Johns A, Harder DR. Mechanoreception by the endothelium: Mediators and mechanisms of pressure- and flow-induced vascular responses. *Blood Vessels* 1990; 27:246-257.
37. Berkowitz HD, Greenstein S, Barker CF, Perloff LJ. Late failure of reversed vein bypass grafts. *Ann Surg* 1989; 210:782-786.
38. Berkowitz HD, Fox AD, Deaton DH. Reversed vein graft stenosis: Early diagnosis and management. *J Vasc Surg* 1992; 15:130-142.

CHAPTER 6

MECHANICAL PROFILES OF POLY-TETRA-FLUORO-ETHYLENE AND SAPHENOUS VEIN ARTERIAL BYPASS GRAFTS IN HUMANS

L Hofstra, DCJJ Bergmans, APG Hoeks, JHM Tordoir, and PJEHM Kitslaar.
(submitted)

ABSTRACT

Background. Mechanical conditions, like peri-anastomotic mismatch in elastic properties, may be associated with the occurrence of stenosis in peripheral bypass grafts. Quantitative in vivo information on elastic properties in peripheral bypass grafts in humans is scarce. We studied peri-anastomotic elastic properties of poly-tetra-fluoro-ethylene (PTFE) grafts and reversed saphenous vein grafts with a new ultrasound technique.

Methods. In 15 patients with a PTFE bypass graft and 15 matched patients with a reversed saphenous vein bypass graft, the diameter and distension of vessel segments was determined at defined sites in the graft and native arterial system with the use of Vessel Wall Doppler Tracking. From diameter and distension we calculated relative distension (RD in %) and cross-sectional area-increase (AI in mm) for all.

Results. AI and RD values were higher in the saphenous vein grafts as compared to the PTFE grafts for both the proximal and the distal graft segments ($p < 0.05$). Higher RD and AI values were found in the proximal segments of PTFE grafts as compared to the distal segments ($p < 0.05$). In both the reversed saphenous vein and PTFE grafts a decrease AI was observed from the inflow artery to the proximal graft segment ($p < 0.05$). No differences in AI and RD were found around the distal anastomosis of the saphenous vein grafts. In contrast, considerable increase in AI and RD was observed from the distal graft segment to the outflow artery in the PTFE grafts.

Conclusions. Local strain and cross-sectional area increase were higher in the proximal graft segments of PTFE grafts as compared to the distal graft segments. A possible explanation may be that the proximal PTFE graft segment is passively distended by the inflow artery, which is supported by the finding that a positive correlation between distension of the inflow arterial segments and distension of the proximal graft segments was observed ($r = 0.69$, $p = 0.01$).

INTRODUCTION

The implantation of a bypass graft in the peripheral arterial circulation in order to reestablish sufficient flow to the lower limb may lead to vascular regions where sudden changes in elastic properties are present, especially at the site of the anastomoses¹⁻⁷. Differences in elastic properties around the anastomoses, have been associated with the occurrence of intimal hyperplasia at these sites⁸⁻¹⁴. Peri-anastomotic differences in elastic properties may lead to a point of excessive mechanical stretching at the anastomoses leading to vessel wall injury, which may initiate a vascular response. Despite general acceptance of the compliance mismatch theory among vascular surgeons, no conclusive studies have been reported demonstrating a correlation between mismatch and the development of stenoses¹. Attempts to evaluate the effect of mismatch in elastic properties between graft and native vessels on intimal hyperplasia in animal models have failed to demonstrate an association, which may be explained by the fact that occlusive intimal hyperplasia does not develop in animals¹⁵⁻¹⁷. Investigation of a possible association between mismatch in elastic properties and intimal hyperplasia in humans has been hampered by the lack of a suitable system to assess elastic properties at defined sites around the anastomoses of the grafts in vivo^{18,19}. As a consequence, quantitative information on dimensions and elastic properties around the anastomoses is scarce.

With the introduction of Vessel Wall Doppler Tracking (VWDT), distension parameters of large vessels can be assessed at predefined sites^{20,21}. This system includes B-mode imaging, which allows selection of sites for measurements with acceptable accuracy. We investigated mechanical profiles of newly inserted polytetrafluoroethylene (PTFE) and saphenous vein bypass grafts in humans with the use of VWDT. Attention was focused on the assessment of elastic properties around anastomoses to obtain insight in the differences in elastic properties around the anastomoses of the PTFE and saphenous vein bypass grafts. In addition, peak systolic velocity was measured at the same defined sites in the vascular conduit and native arteries.

MATERIALS AND METHODS

Patients

Patients with arterial insufficiency of the lower extremity as manifested by disabling claudication, restpain and/or tissue loss and receiving a PTFE or reversed saphenous vein femorodistal bypass graft were considered candidates for our study. The PTFE and reversed saphenous vein group were matched for sites of anastomosis and clinical stage (Table I). The patients with reversed saphenous vein bypass grafts were selected from a group of 32 consecutive patients. The function of the graft was tested with the use of a Duplex scan before the patients entered the study. Patients with evidence of graft related stenosis were excluded from this study, since a stenosis could alter the hemodynamics of the graft by influencing pulse wave reflections²². Fifteen consecutive patients with a PTFE graft and 15 matched patients with a reversed saphenous vein

graft entered the study. Patients were not randomly assigned to either PTFE or reversed vein implantation. The choice for either PTFE or reversed saphenous vein graft was based on the quality of the veins suitable for grafting and/or the site of the distal anastomoses. Thin-walled ringed 6 mm PTFE grafts were used (Goretex, Flagstaff, Arizona, USA). Both types of grafts were implanted using standard techniques and were inserted in an end-to-side fashion with running 6x0 Prolene sutures.

Table I: Patient characteristics of the PTFE and reversed saphenous vein group. The distribution of proximal and distal anastomoses, of special importance in this mechanical study, is equal.

	PTFE (n=15)	Reversed (n=15)
Male	13	12
Female	2	3
Age (Yrs $\pm$ SD)	68 (7.0)	67 (7.3)
Diabetic	6	5
Clinical stage :		
II	7	7
III	4	3
IV	4	5
Proximal Anastomosis :		
Common Femoral Artery	15	15
Distal Anastomosis :		
Popliteal Artery (above knee)	7	7
Popliteal Artery (below knee)	4	4
Crural	4	4

Measurements of the elastic properties

The dimensions and distension parameters of the graft and native artery were assessed with the use of Vessel Wall Doppler Tracking (VWDT). A detailed description of this system is given elsewhere (chapter 3 and 4)^{20,21}. From distension (in μm) and diameter (in mm), the following parameters were calculated:

$$\text{Relative Distension (RD)} = \frac{\Delta D}{D} * 100\% \quad (1)$$

$$\text{Area - Increase (AI)} = \frac{\pi \Delta d * D}{2} \quad (2)$$

Relative distension (RD) gives the local strain of the vessel wall. Area-increase (AI) represents the capacity of the vessel to store and to release temporarily blood volume during the cardiac cycle²³. Usually these parameters are expressed per unit of pressure, resulting in the distensibility coefficient and compliance coefficient respectively. Non-invasive assessment of pressure values is possible at the finger (Finapres, volumeclamp technique) or at the arm (Dynamap, automatic cuff device). These system have been validated with intra-arterially obtained pressures at the brachial artery, showing good correlations²⁴. However, the estimation of pulse pressure values in the peripheral bypass from pressure values obtained at the finger or the arm in order to calculate compliance and distensibility in the graft is questionable. First of all we do not know to what extent the pulsatile character of pressure changes over diseased vessel segments. Furthermore, the systolic pressure values may change when a small caliber vessel, like the reversed saphenous vein is anastomosed to a large caliber vessel like the femoral artery. The anastomoses may serve as a reflection point, leading to unpredictable behavior of pressure²⁵⁻²⁷. Since invasive assessment of pressure at different sites in the graft is not acceptable, we decided not to express local strain and area-increase per unit of pressure in this study. To investigate the ability of the VWDT system to detect small distensions of vessels, we tested the system in an in vitro setting. A proper distension waveform was obtained with distension values as low as $1 \mu\text{m}^{21}$. This indicates that expected low distension values in PTFE grafts can be properly measured with the use of Vessel Wall Doppler Tracking.

In each graft measurements were performed proximal (1 cm distal to the proximal anastomosis), in the middle segment, and distal (1 cm proximal to the distal anastomosis). The inflow artery was measured 1 cm proximal to the proximal anastomosis, the outflow artery was measured 1 cm distal to the distal anastomosis (Figure 1). Proper selection of the sites of measurements was performed with the use of repeated B-mode imaging. Systolic and diastolic pressure were monitored, using a Finapres device (Ohmeda 2300, Ohmeda Monitoring Systems, Englewood, Colorado, USA), in order to have insight in the general hemodynamic condition of the patients.

Patients showing an unstable hemodynamic condition during the measurements were excluded from the study, since changes in mean pressure might affect the local elastic properties.

Figure 1: Sites of measurements in the graft and in- and outflow arterial segments.

Measurement of peak systolic velocity (PSV)

The use of Duplex ultrasound in the detection of distension and diameter allowed assessment of blood stream velocity at the same predefined sites in the graft and the inflow and outflow arterial segments. Peak systolic velocity was measured by placement of a Doppler sample midstream, with an angle of 60° . Patients were measured in the supine position after an acclimatisation period of 20 minutes. All patients gave a written informed consent before entering the study. The study was approved by the institutional medical ethical committee.

ANALYSIS

Data are presented as medians and 25th and 75th percentile intervals. Within the groups the values of diameter, distension, RD, and AI measured at the defined sites in the graft and in the in- and outflow arterial segments, respectively, were compared using the Wilcoxon-signed-rank test. Between the groups the values of the parameters measured at the defined sites were compared using the Mann-Whitney test. To investigate correlations between different parameters, linear regression analysis was used. A p-value of less than 0.05 was accepted as level of statistical significance.

RESULTS

The patient characteristics of both groups (Table I) were comparable. Of importance in this mechanical study are the sites of proximal and distal anastomoses. Matching for site of proximal and distal anastomoses resulted in an equal distribution of proximal as well as distal anastomotic sites in the groups. The values of diameter, distension, RD, AI and PSV of the PTFE and the reversed saphenous vein group are summarized in table II and depicted in figures 2 and 3.

Table II: Values for diameter (DI), distension (DIST), relative distension (RD), area-increase (AI) and peak systolic velocity (PSV) measured at the inflow- and outflow arterial segments and proximal, mid and distal graft segments in the PTFE and reversed saphenous vein group. Data are presented as medians and 25th and 75th intervals.

	Inflow	Proximal	Mid	Distal	Outflow
DI (mm):					
PTFE	7.9 (6.7-10.4)	4.9 (4.8-5.4)	5.1 (4.8-5.3)	4.8 (4.6-5.3)	4.7 (2.7-5.4)
Reversed	8.6 (7.1-11.2)	4.1 (3.1-5.6)	5.4 (4.1-6.1)	6.0 (4.6-6.8)	4.9 (3.5-5.3)
DIST (µm)					
PTFE	167 (140-282)	96 (17-112)	18 (14-39)	21 (13-39)	78 (50-128)
Reversed	209 (173-271)	139 (102-212)	101 (77-131)	81 (70-123)	75 (61-100)
RD (%):					
PTFE	2.2 (1.6-2.7)	1.9 (0.3-2.3)	0.4 (0.3-0.8)	0.5 (0.3-0.8)	2.2 (1.1-3.0)
Reversed	2.8 (1.8-3.7)	3.5 (2.4-4.6)	1.7 (1.6-2.9)	1.5 (1.1-2.5)	1.8 (1.3-2.5)
AI (mm ²)					
PTFE	2.12 (1.21-4.34)	0.77 (0.14-0.87)	0.17 (0.12-0.30)	0.16 (0.09-0.27)	0.64 (0.20-1.05)
Reversed	3.06 (2.23-4.72)	1.20 (0.63-1.48)	0.90 (0.55-1.17)	0.78 (0.57-1.32)	0.63 (0.32-0.84)
PSV (cm/s)					
PTFE	103 (82-127)	81 (62-109)	81 (60-97)	71 (53-87)	71 (56-105)
Reversed	107 (82-144)	97 (74-149)	80 (58-97)	63 (51-73)	91 (59-133)

Figure 2: Median values and 25th and 75th intervals of diameter (DI), distension (DIST), relative distension (RD), and area-increase (AI) measured at the proximal (Prox), mid (Mid) and distal (Dist) segments of the grafts and the inflow (Infl) and outflow (Outfl) arterial segments, (* , $p < 0.05$).

Figure 3: Median values and 25th and 75th percentile intervals of peak systolic velocity (PSV) for the PTFE and reversed saphenous vein graft, measured at defined sites in the graft (prox, mid and distal) and the inflow (Infl) and outflow (Outfl) arterial segments, (*, $p < 0.05$).

PTFE group. A significant decrease in diameter was observed between the arterial inflow segments (7.9 mm) and the proximal segments of the graft (4.9 mm, $p < 0.05$). As expected, no change in diameter was found along the prosthetic grafts. Around the distal anastomoses no significant size differences were found. Around the proximal anastomoses no significant differences were apparent in RD. In the course of the PTFE grafts a decrease in RD was found between the proximal segments of the grafts and the middle segments (1.9% vs 0.4%, $p < 0.05$), but no further decrease was found from the middle segments to the distal segments. An increase in RD was observed from the distal graft segments to the arterial outflow segments ($p < 0.05$).

A decrease in distension was observed between the inflow segment and the proximal graft segment (167 vs 96 μm) and from the proximal to mid graft (96 vs 18 μm , $p<0.05$). From the mid-graft segments to the distal segments no changes in distension were found. An increase in distension was observed in the distal anastomoses (21 vs 78 μm , $P<0.05$). A decrease in AI was observed between the arterial inflow segments and proximal graft segments (2.1 vs 0.77 mm^2 , $p<0.05$) and a further decrease was encountered from proximal to midgraft (0.77 vs 0.16 mm^2 , $p<0.05$). Around the distal anastomoses an increase in AI was observed (0.16 vs 0.64 mm^2 , $p<0.05$). Peak systolic velocity decreased in the proximal anastomoses (103 vs 81 cm/s , $p<0.05$). A stable pattern of PSV was found along the prosthetic grafts, and no changes were observed around the distal anastomoses.

Reversed saphenous vein group: The proximal graft diameter (4.1 mm) was smaller as compared to the diameter of the arterial inflow segments (8.6 mm, $p<0.05$). The effect of reversed tapering is illustrated by a gradual increase in diameter from proximal (4.1 mm) to distal (6.0 mm, $p<0.05$) in these grafts. A difference in diameter was observed around the distal anastomoses (6.0 in the distal part of the grafts vs 4.9 mm in the arterial outflow segments, $p<0.05$). An increase in RD was found between the inflow arterial segment (2.8%) and proximal graft segments (3.5%, $p<0.05$). A gradual decrease in RD was found from proximal (3.5%) to distal (1.5%, $p<0.05$) in the grafts. Around the distal anastomoses no significant differences in RD could be observed. A decrease in distension was observed between the inflow segment and the proximal graft segment (209 vs 139 μm) and between the proximal graft segment and the mid-graft segment (139 vs 101 μm , $p<0.01$). Between the mid-graft segments and the distal graft segments and around the anastomoses no differences in distension were observed. The Area-increase (AI) was larger in the inflow arterial segments as compared to the proximal segments of the grafts (3.0 vs 1.2 mm^2 , $p<0.05$). No change in AI was observed along the grafts. Around the distal anastomoses no significant differences in AI were apparent. Within the proximal anastomoses of the reversed vein grafts no changes in PSV were observed. A decrease in PSV was observed along the grafts (99 vs 63 cm/s , $p<0.05$). Around the distal anastomoses no changes in PSV were observed.

DISCUSSION

In the present study we investigated the dimensions and elastic properties around the anastomoses of PTFE and saphenous vein grafts in humans with a new ultrasound system, Vessel Wall Doppler Tracking. We observed higher values of distension in the proximal segments of the PTFE grafts compared to the mid- and distal segments which is surprising, since homogeneous intrinsic elastic properties in prosthetic graft material seemed more likely. Due to the high distension values in the proximal grafts segments, the values for RD and AI were also higher in this segment. The values for distension and relative distension in the mid-graft and distal graft segments of the PTFE grafts

are comparable to values from in vitro assessment of distension of PTFE grafts, reporting values of one-sixth of femoral artery values^{5,19,28}. Theoretically, the high values of distension in the proximal graft segment may be the effect of differentiation of pulse wave reflection^{22,27}. The proximal segments of the grafts could be a site where the pulse wave, reflected from the periphery, augments systolic pressure, thereby increasing pulse pressure values at this particular point, leading to higher distensions^{22,26}. However, this mechanism is not likely, since the length of the graft is many times shorter as the wavelength of the pulse wave. Presumably, the increase of peak systolic pressure due to pulse wave reflection will have a general effect in the graft. Another possible explanation may be the interaction between the arterial inflow segment and the proximal graft segment⁵. The distension of the inflow arterial segments may influence the distension of the most proximal graft segments, since these parameters were assessed quite close to the anastomosis, within 1 cm from the proximal anastomoses. To test this hypothesis, we correlated distension values of inflow arteries and proximal graft segments. The result is depicted in figure 4.

Figure 4: Scatterplot of distension values of the inflow arterial segments (X-axis) and the proximal graft segments (Y-axis). Linear regression analysis revealed a positive correlation between distension values measured at these sites.

Linear regression analysis revealed a significant correlation between the distension values measured at these two different sites ($r = 0.69$, $p = 0.01$, $y = 1.31 + 0.31 \cdot x$). This supports the concept that the distension of the proximal graft segments is influenced by the distension of the inflow arterial system. The tendency of lower distension values in the inflow segments of the PTFE group as compared to the reversed vein group is a further indication that elastic properties of the prosthesis and of the inflow arteries close to the anastomosis are influenced by each other, indicating artery-graft interaction. In another study from our institute, the elastic profiles of different graft arteriovenous fistulas were assessed with the use of VWDT in a similar fashion (chapter 7)²⁹. Substantially lower values for AI and RD were found in the brachial artery of patients with a PTFE interposition graft as compared to patients with a saphenous vein interposition graft for the construction of a vascular access. Since in that study the patients were randomly assigned to either the PTFE or saphenous group, differences in the intrinsic elastic properties of the brachial artery are unlikely and cannot explain the lower values for the elastic properties in the PTFE group. Therefore, it seems that the distension of the brachial artery close to the arterial anastomosis of patients with a PTFE graft is restricted by the stiffness of the prosthesis. This observation supports the concept that the vessel segments close to the anastomosis are interrelated with respect to elastic properties, indicating graft-artery interaction.

The consequence of the relatively high distension values at the proximal graft segments is that the differences in elastic properties around the proximal anastomoses of the PTFE grafts are less pronounced than one would expect. The mechanical interaction between native arterial system and the PTFE graft attenuates the difference in intrinsic elastic properties to some extent.

At the distal anastomoses of the PTFE grafts an increase in the elastic parameters was observed. In vivo flow studies demonstrated that a sudden increase in compliance is associated with flow disturbances like increased particle residence time and flow stasis, suggesting low shear^{7,30}. Low shear rate has been associated with increased intimal thickening by numerous investigators^{2,31-34}. Therefore, the increase in the elastic parameters at the distal anastomoses of the PTFE grafts may explain the predisposition of intimal hyperplasia to develop at this site^{9,14}.

In the reversed saphenous vein bypass grafts a gradual decrease in distension was observed along the entire vascular reconstruction. The most prominent finding in this group is the decline in AI in the proximal anastomoses. However, due to the increase in diameter along the graft the capacity to store blood volume, represented by AI, remained stable in the course of the graft. The sudden decrease in AI in the proximal anastomoses of reversed saphenous vein bypasses coincides with the frequent occurrence of stenosis at this site^{11,35}.

The capacity to store blood volume during the cardiac cycle is a measure for the hemodynamic function of the graft. As a vascular conduit, the bypass has to adequately manage pulsatile flow³. This condition seems to be fulfilled by the reversed saphenous vein graft. The decline in AI along the prosthetic graft indicates that the

capacity to store blood volume in the distal part of the graft is low. In the reversed saphenous vein graft also a decrease in PSV values is observed along the grafts, which can be attributed to the increase in diameter along the grafts. With the use of long prosthetic grafts to construct bypasses to infrapopliteal and crural arteries, a further loss of pulsatile energy might occur. This may explain the more pronounced difference in patency rates between PTFE and saphenous vein grafts at the below-knee level. The length of the PTFE graft may be the limiting factor rather than crossing the knee joint³⁶⁻³⁸. In the development of new prosthetic graft materials the ability of maintaining hemodynamic capacity along the entire length of the graft should be of major concern. The hypothesis that a mismatch in elastic properties contributes to the development of intimal hyperplasia stems from the frequent occurrence of this phenomenon at anastomoses of grafts^{1,2,8,9,12,19,39-42}. A mismatch in elastic properties may lead to vessel wall injury, thereby initiating a vascular response¹. In addition, it leads to a site of excessive mechanical stretching at the anastomoses under pulsatile conditions, which may promote smooth muscle cell proliferation⁴³⁻⁴⁴. In the present study notable differences were observed around the anastomoses of the PTFE and reversed saphenous vein bypass graft. At this moment we can only speculate on the influence of the differences around the anastomoses of distension, RD and AI on the occurrence of intimal hyperplastic stenoses. A prospective analysis correlating quantitative differences in elastic properties with the occurrence of stenosis is necessary to address this question.

In summary, the dimensions and elastic properties of PTFE and saphenous vein bypass grafts in humans were assessed in vivo at defined sites in the graft and native arterial system, in order to obtain a mechanical profile. Higher distension values were found in the proximal segments of PTFE grafts as compared to the distal segments. This may be explained by the interaction between the native artery and the proximal graft segments, which is suggested by the positive correlation between distension of the inflow arterial segments and the proximal graft segments. At the distal anastomoses of the PTFE grafts an increase in distension and area-increase was observed. In the reversed saphenous vein grafts a decrease in distension and AI was observed around the proximal anastomoses. Around the distal anastomoses no differences in elastic parameters were observed in the saphenous vein group.

REFERENCES

1. Abbott WM, Megerman J, Hasson JE, L'Italien G, Warnock DF. Effect of compliance mismatch on vascular graft patency. *J Vasc Surg* 1987; 5:376-382.
2. Bassiouny HS, White S, Glagov S, Choi E, Giddens DP, Zarins CK. Anastomotic intimal hyperplasia: Mechanical injury or flow induced. *J Vasc Surg* 1992; 15:708-717.
3. Baird RN, Abbott WM. Pulsatile blood-flow in arterial grafts. *Lancet* 1976; October 30:948-950.
4. Wu MH, Shi Q, Sauvage LR, et al. The direct effect of graft compliance mismatch per se on development of host arterial intimal hyperplasia at the anastomotic interface. *Ann Vasc Surg* 1993; 7:156-168.
5. Rodgers VGJ, Teodori MF, Brant AM, Borovetz HS. Characterization in vitro of the biomechanical properties of anastomosed host artery-graft combinations. *J Vasc Surg* 1986; 4:396-402.
6. Okuhn SP, Connelly DP, Calakos N, Ferrell L, Man-Xiang P, Goldstone J. Does compliance mismatch alone cause neointimal hyperplasia? *J Vasc Surg* 1989; 9:35-45.
7. Stewart SFC, Lyman DJ. Effects of a vascular graft/natural artery compliance mismatch on pulsatile flow. *J Biomech* 1992; 25:297-310.
8. Abbott WM, Megerman J. Adaptive responses of arteries to grafting. *J Vasc Surg* 1989; 9:377-379.
9. Echave V, Koornick AR, Haimov M, Jacobson JH. Intimal hyperplasia as a complication of the use of the polytetrafluoroethylene graft for femoral-popliteal bypass. *Surgery* 1979; 86:791-798.
10. Madras PN, Ward CA, Johnson WR, Singh PI. Anastomotic hyperplasia. *Surgery* 1981; 90:922-923.
11. Berkowitz HD, Greenstein S, Barker CF, Perloff LJ. Late failure of reversed vein bypass grafts. *Ann Surg* 1989; 210:782-786.
12. Swedberg SH, Brown BG, Sigley R, Wight TN, Gordon D, Nicholls SC. Intimal fibromuscular hyperplasia at the venous anastomosis of PTFE grafts in hemodialysis patients. *Circulation* 1989; 80:1726-1736.
13. Jenkins AMcL, Buist TAS, Glover SD. Medium-term follow-up of forty autogenous vein and forty polytetrafluoroethylene (Gore-Tex) grafts for vascular access. *Surgery* 1980; 88:667-672.
14. Sottiturai VS, Sue SL, Feinberg II EL, Bringaze WL, Tran AT, Batson RC. Distal anastomotic intimal hyperplasia: Biogenesis and etiology. *Eur J Vasc Surg* 1988; 2:245-256.
15. Ferrel M, Fuster V, Gold HK, Chesebro JH. Choosing appropriate experimental animal model for the prevention of restenosis. *Circulation* 1992; 85:1630-1631.
16. Clowes AW, Reidy MA. Prevention of stenosis after vascular reconstruction: Pharmacologic control of intimal hyperplasia - A review. *J Vasc Surg* 1991; 6:885-891.
17. McBride W, Lange RA, Hillis LD. Restenosis after successful coronary angioplasty. Pathophysiology and prevention. *New Eng J Med* 1988; 318:1734-1737.
18. Lye CR, Sumner DS, Hokanson DE, Strandness DE. The transcutaneous measurement of the elastic properties of the human saphenous vein femoropopliteal bypass graft. *Surg Gynecol Obstet* 1975; 141:891-895.
19. Hokanson DE, Strandness DE. Stress-strain characteristics of various arterial grafts. *Surg Gynecol Obstet* 1968; 57-60.
20. Hoeks APG, Brands PJ, Smeets FAM, Reneman RS. Assessment of distensibility of superficial arteries. *Ultrasound Med Biol* 1990; 16:121-128.
21. Hofstra L, Ermers EJM, Langeveld APM, Hoeks APG, Kitslaar PJEHM. Vessel Wall Doppler Tracking: A new technique to assess vessel wall compliance. *J Vasc Tech* 1993; 17:65-68.
22. O'Rourke MF. Arterial stiffness, systolic blood pressure, and logical treatment of arterial hypertension. *Hypertension* 1990; 15:339-347.

23. Reneman RS, Merode van T, Brands PJ, Hoeks APG. Inhomogeneities in arterial wall properties under normal and pathological conditions. *J Hypertens* 1992; 10:S35-S39.
24. Imholz BPM, Settels JJ, Meiracker van der AH, Wesseling KH, Wieling W. Non-invasive continuous finger blood pressure measurement during orthostatic stress compared to intra-arterial pressure. *Cardiovasc Res* 1990; 24:214-221.
25. Murgo JP, Westerhof N, Giolma JP, Altobelli SA. Aortic input impedance in normal man: Relationship to pressure wave forms. *Circulation* 1980; 62:105-116.
26. O'Rourke MF. Systolic blood pressure: arterial compliance and early wave reflection, and their modification by antihypertensive therapy. *J Hum Hypertens* 1989; 3:47-52.
27. Kelly RP, Gibbs HH, O'Rourke MF, et al. Nitroglycerin has more favourable effects on left ventricular afterload than apparent from measurement of pressure in a peripheral artery. *Eur Heart J* 1990; 11:138-144.
28. Pourdeyhi B, Wagner D. On the correlation between the failure of vascular grafts and their structural and material properties: A critical analysis. *J Biomed Mat Res* 1986; 20:375-409.
29. Hofstra L, Bergmans DCCJ, Hoeks APG, Kitslaar PJEHM, Leunissen KML, Tordoir JHM. Mismatch in elastic properties around anastomoses of interposition grafts for hemodialysis access. *JASN* 1994; 5:1243-1250.
30. Shu MCS, Noon GP, Hwang NHC. Flow profiles and wall shear stress distribution at a hemodialysis venous anastomosis: preliminary study. *Biorheology* 1987; 24:723-735.
31. Zarins CK, Bomberger RA, Glagov S. Local effects of stenoses: Increased flow velocity inhibits atherogenesis. *Circulation* 1981; 64:II221-II227.
32. Friedman MH, Barger CB, Deters OJ, Hutchins GM, Mark FF. Correlation between wall shear and intimal thickness at a coronary artery branch. *Atherosclerosis* 1987; 68:27-33.
33. Kamiya A, Togawa T. Adaptive regulation of wall shear stress to flow change in the canine carotid artery. *Am J Physiol* 1980; 239:H14-H21.
34. Rittgers SE, Karayannacos PE, Guy JF, et al. Velocity distribution and intimal proliferation in autologous vein grafts in dogs. *Circ Res* 1978; 42:792-801.
35. Berkowitz HD, Fox AD, Deaton DH. Reversed vein graft stenosis: Early diagnosis and management. *J Vasc Surg* 1992; 15:130-142.
36. Prendiville EJ, Yeager A, O'Donnell TF, et al. Long-term results with the above-knee popliteal expanded polytetrafluoroethylene graft. *J Vasc Surg* 1990; 11:517-524.
37. Michaels JA. Choice of material for above-knee femoropopliteal bypass graft. *Br J Surg* 1989; 76:7-14.
38. Veith FJ, Gupta SK, Ascer E, et al. Six year prospective multicenter randomized comparison of autologous saphenous vein and polytetrafluoroethylene grafts in infrainguinal arterial reconstructions. *J Vasc Surg* 1986; 3:104-114.
39. Kidson IG, Abbott WM. Low compliance and arterial graft occlusion. *Circulation* 1978; 58:11-14.
40. Werner GS, Wiegand V, Tebbe U, Kreuzer H. Contrary effects of acetylcholine on coronary arteries and aortocoronary venous grafts in man. *Eur Heart J* 1989; 10:86-91.
41. LoGerfo FW, Quist WC, Nowak MD, Crawshaw HM, Haudenschild CC. Downstream anastomotic hyperplasia. A mechanism of failure in Dacron arterial grafts. *Ann Surg* 1983; 197:479-483.
42. Sottiurai VS, Yao JST, Flinn WR, Batson RC. Intimal hyperplasia and neointima: An ultrastructural analysis of thrombosed grafts in humans. *Surgery* 1983; 93:809-817.
43. Predel HG, Yang Z, Von Segesser L, Turina M, Buhler FR, Luscher TF. Implications of pulsatile stretch on growth of saphenous vein and mammary artery smooth muscle. *Lancet* 1992; 340:878-879.
44. Leung DYM, Glagov S, Mathews MB. Cyclic stretching stimulates synthesis of matrix components by arterial smooth muscle cells in vitro. *Science* 1976; 191:475-477.

CHAPTER 7

MISMATCH IN ELASTIC PROPERTIES AROUND ANASTOMOSES OF INTERPOSITION GRAFTS FOR HEMODIALYSIS ACCESS

L Hofstra MD, DCJJ Bergmans, APG Hoeks PhD, PJEHM Kitslaar MD PhD,
KML Leunissen MD PhD and JHM Tordoir MD PhD

Adapted from the Journal of the American Society of Nephrology 1994; 5: 1243-1250

ABSTRACT

Background: Arterio-venous fistulas for hemodialysis access, constructed with the use of interposition grafts, are often complicated by intimal hyperplastic stenosis, mainly occurring at the venous anastomosis. In the current study we quantified mismatch in elastic properties around the arterial and venous anastomoses of graft AV fistulas in humans in order to find clues for the predisposition of intimal hyperplasia to develop at the venous anastomosis.

Methods: The elastic properties of graft AV fistulas of 31 hemodialysis patients were investigated with the use of Vessel Wall Doppler Tracking, two weeks after construction. Nine saphenous vein (SV) grafts, 8 expanded-PTFE (ePTFE) grafts and 14 stretch-PTFE (sPTFE) grafts were measured at the arterial inflow segment, the proximal graft segment, the distal graft segment and the venous outflow segment. Area-increase (AI), representing the capacity of the vessel wall to store blood volume, and relative distension (RD), representing the strain, were calculated from diameter and distension.

Results: A decrease in AI was observed in the arterial anastomoses of all graft types. An increase in AI was found in the venous anastomosis of ePTFE and sPTFE grafts. Higher values for AI and RD were found at the proximal and distal graft segments of the saphenous vein grafts when compared to the prosthetic grafts. In the sPTFE grafts the level of AI was maintained along the graft, while in the ePTFE grafts a decrease in AI was found.

Conclusion: In the arterial anastomoses of arteriovenous fistulas a decline in the capacity to store blood volume was observed. By contrast, an increase in capacity to store blood volume was found in the venous anastomoses. Since the latter condition is associated with flow disturbances, like enhanced particle residence time and low shear rate, it may contribute to the predisposition of intimal hyperplasia to develop at the venous site.

INTRODUCTION

Patients requiring hemodialysis due to end-stage renal failure depend on a proper function of the vascular access, to provide sufficient flow during dialysis¹. An increasing number of these patients receive an interposition graft in the usually secondary construction of an arterio-venous (AV) fistula¹⁻³. The reported 50% revision rate of the graft AV fistulas during the first post-operative year, illustrates the susceptibility of this type of vascular access to develop complications and leads to an increased hospitalisation^{2,3}. The main cause for the failure of the vascular access is graft stenosis due to intimal hyperplasia, jeopardizing graft flow and causing high venous pressure¹⁻⁴. Stenotic intimal hyperplasia frequently occurs at the site of the venous anastomosis and rarely develops at the arterial anastomosis^{2,3,5-10}. The incidence of stenoses developing at the venous anastomosis may be caused by the release of growth factors in the graft due to puncturing^{3,4,11}. The upstream release of growth factors may lead to proliferation of vascular smooth muscle cells (VSMC) at the site of the venous anastomosis¹². However, since stenoses are not diffusely distributed along the efferent vein but localized at the anastomosis, other contributing factors are likely. A possible mechanism is mismatch in elastic properties around the anastomoses^{7,10,13-15}. Mismatch in elastic properties leads to excessive mechanical stretch at the anastomoses¹³, which may promote smooth muscle cell replication¹⁶⁻¹⁹. Furthermore, the excessive mechanical stretch may lead to localized vessel wall injury^{13,20}, initiating a vascular response. Another way by which compliance mismatch may affect intimal hyperplasia, is the influence on flow. The effect of an abrupt change in compliance on flow characteristics has been studied with the use of flow models²¹⁻²³. A sudden decrease in compliance did not lead to flow disturbances whereas a sudden increase in compliance resulted in enhanced particle residence time and flow separation, suggesting low shear conditions. Numerous reports demonstrated an increased intimal thickening in areas of low shear stress²²⁻²⁸, probably by affecting the growth control function of the endothelium²⁹⁻³¹.

Quantitative information on elastic properties around the anastomoses of interposition grafts is scarce. The elastic properties of graft AV fistulas of hemodialysis patients with the use of Vessel Wall Doppler Tracking^{32,33} were measured, two weeks after construction. Three different graft types were investigated: autologous saphenous vein grafts, expanded polytetrafluoroethylene (ePTFE) grafts, and a new PTFE graft with stretch properties (stretch-PTFE). The elastic properties were assessed at predefined sites in the graft, the arterial inflow segment and the venous outflow segment. The aim of the study was to compare the peri-anastomotic mismatch in elastic properties between the arterial and venous anastomoses in order to find clues for the predisposition of intimal hyperplasia to develop at the venous anastomosis.

MATERIALS AND METHODS

Patients

A total of 31 patients was examined. In 25 patients the construction of a vascular access using interposition grafts was a secondary procedure. Previous arteriovenous fistulas at the usual sites, mostly Brescia Cimino fistulas, had failed. In the remaining 6 patients the use of an interposition graft was a primary choice due to the absence of a suitable vein. The proper function of the graft was tested with the use of a Duplex scan before the patients entered the study. Patients with evidence of graft related stenoses were excluded from this study, since a stenosis could alter the hemodynamics of the graft by influencing pulse wave reflections³⁴. Thirty-one consecutive patients entered the study. In the first half of the study period patients were randomly assigned to either ePTFE (n=8, mean age 51±20 years) or saphenous vein implantation (n=9, mean age: 53±17 years). In the second half of the study period only stretch-PTFE grafts were used (n=14, mean age 56±15 years). All grafts were implanted in a subcutaneous loop tunnel in the forearm between the brachial artery and the vein. The arterial and venous anastomoses were constructed at the site of the elbow in an end-to-side fashion with running 7-0 polypropylene (Prolene) sutures.

Measurements of the elastic properties

The measurements of the elastic properties of the graft AV fistulas were performed two weeks after construction. Patients were measured in the supine position after an acclimatization time of 20 minutes. The elastic properties of the graft, the brachial artery and venous outflow segment were assessed with the use of Vessel Wall Doppler Tracking (VWDT). This system has been described in detail elsewhere (chapter 3 and 4)^{32,33}. From distension (in μm) and diameter (in mm) the following parameters were calculated:

$$\text{Relative Distension (RD)} = \frac{\Delta D}{D} * 100 (\%) \quad (1)$$

$$\text{Area-Increase (AI)} = \frac{\pi \Delta D * D}{2} \quad (2)$$

Relative distension (RD) gives the fractional diameter increase. Area-increase represents the capacity of the vessel wall to store blood volume during cardiac cycle³⁵. Usually, RD and AI are corrected for pulse pressure, resulting in the distensibility coefficient and the cross-sectional compliance, respectively^{36,37}. Animal studies demonstrated a drop in pressure along AV-fistulas from 120 mmHg in the arterial inflow segment to 50 mmHg in the venous outflow segment³⁸. This indicates that non-invasively obtained pressure values with the use of Finapres or Dynamap cannot

represent the pressure values along the graft. Therefore, it was decided to use RD and AI to represent the elastic properties of the arterial wall. RD and AI were averaged from the 3-5 recorded heartbeats per measurements. The variation coefficient for RD and AI within the measurement was always less than 5%. In each graft measurements were performed proximal (1 cm from the proximal anastomosis) and distal (1 cm from the distal anastomosis). The inflow artery was measured 1 cm from the proximal anastomosis, the outflow vein was measured 1 cm from the distal anastomosis (Figure 1).

Figure 1: Sites of measurements in the graft and in- and outflow segments. The measurements were performed within a distance of 1 centimeter from the anastomoses.

ANALYSIS

Data are presented as medians and 25th and 75th percentile intervals. Within the groups the values of diameter, RD, and AI measured at the defined sites in the graft and in the in- and outflow venous segments, respectively, were compared using the Wilcoxon-signed-rank test. Between the groups the values of the parameters measured at the defined sites were compared using the Mann-Whitney test. Linear regression analysis was used to correlate variables. A p-value of less than 0.05 was accepted as level of statistical significance.

RESULTS

The patient characteristics were not different between the groups (Table I). The results of measurement of diameter, RD and AI measured in the three different groups are summarized in Table II and depicted in Figure 2 to 5.

Table I: Patient characteristics of the different groups (SV = Saphenous vein graft, sPTFE = stretch PTFE, ePTFE = expanded PTFE).

	SV Grafts	sPTFE Grafts	ePTFE Grafts
Age (years)	54 (17)	56 (15)	54 (21)
Male/Female	7/2	5/9	5/3

Table II

	Brachial Artery	Proximal	Distal	Vein
Diameter:				
SV	4.8 (4.4 - 5.3)	4.5 (3.2 - 6.6)	3.6 (2.7 - 6.6)	6.3 (2.8 - 8.0)
sPTFE	5.1 (4.1 - 5.6)	4.4 (4.1 - 4.6)	4.5 (4.1 - 4.9)	6.2 (3.4 - 8.6)
ePTFE	5.8 (5.2 - 6.7)	4.9 (4.8 - 5.7)	4.9 (4.7 - 5.7)	4.3 (2.7 - 5.1)
Relative Distension:				
SV	5.7 (4.5 - 9.7)	3.5 (2.7 - 8.0)	9.0 (3.4 - 12.9)	5.6 (3.0 - 15.9)
sPTFE	4.1 (2.5 - 6.0)	2.8 (1.6 - 3.4)	1.6 (1.0 - 3.4)	4.0 (2.0 - 6.4)
ePTFE	3.4 (1.8 - 5.8)	1.8 (1.1 - 2.4)	1.1 (0.5 - 2.3)	4.8 (0.9 - 9.5)
Area Increase:				
SV	2.1 (1.2 - 4.0)	1.1 (0.9 - 2.8)	1.4 (1.3 - 3.3)	3.1 (1.0 - 6.5)
sPTFE	1.8 (1.0 - 2.6)	0.7 (0.4 - 1.1)	0.7 (0.3 - 1.0)	1.8 (1.1 - 3.5)
ePTFE	1.4 (1.0 - 4.3)	0.7 (0.5 - 1.2)	0.4 (0.2 - 0.8)	0.9 (0.3 - 1.6)

Saphenous vein group (n=9).

No differences in diameter were observed around the anastomoses and no changes in diameter along the graft were found (Table II, Figure 2). A large increase in RD from proximal to distal in the graft was observed ($p < 0.05$). A decrease in AI was found in the arterial anastomosis ($p < 0.05$). No change in AI along the graft was observed and no difference in AI around the venous anastomoses was found.

Figure 2: Saphenous vein grafts. Median values and 25th and 75th percentile intervals of diameter (DI), relative distension (RD) and area-increase (AI) measured at the proximal (Prox), mid (Mid) and distal (Dist) segments of the grafts and the artery (Art) and vein segments (* , p<0.05).

e-PTFE group (n=8)

A decrease in diameter was observed in the arterial anastomoses ($p<0.05$); along the graft and around the venous anastomoses no differences in diameter were found (Table II, Figure 3). A decrease in RD in the arterial anastomoses was shown ($p<0.05$); along the graft and around the venous anastomoses no differences were observed. A decrease in AI was observed in the arterial anastomoses ($p<0.05$) and along the graft ($p<0.05$). Around the venous anastomoses an increase in AI was observed ($p<0.05$).

Figure 3: expanded PTFE grafts. Median values and 25th and 75th percentile intervals of diameter (DI), relative distension (RD) and area-increase (AI) measured at the proximal (Prox), mid (Mid) and distal (Dist) segments of the grafts and the artery (Art) and vein segments, (* , $p<0.05$).

Stretch PTFE group (n=14)

Around the arterial anastomoses and along the grafts no differences in diameter were observed, but an increase in diameter was found around the venous anastomoses (Table II, Figure 4). No differences in RD were observed around the arterial anastomoses and along the graft. RD increased around the venous anastomoses ($p<0.05$). A decrease in AI was observed around the arterial anastomoses ($p<0.05$), but an increase was found around the venous anastomoses ($p<0.05$). Along the grafts no changes in AI were shown.

Figure 4: stretch PTFE grafts. Median values and 25th and 75th percentile intervals of diameter (DI), relative distension (RD) and area-increase (AI) measured at the proximal (Prox), mid (Mid) and distal (Dist) segments of the grafts and the artery (Art) and vein segments, (* , $p<0.05$).

COMPARISON BETWEEN THE GROUPS

Relative Distension

RD was higher in the SV group in the proximal and distal graft segment as compared to both the ePTFE group and the stretch-PTFE group ($p<0.05$) (Table II, Figure 5). Between the ePTFE group and the stretch-PTFE group no differences were found in the graft. In the arterial inflow segment RD was higher in the SV group as compared to the ePTFE group ($p<0.05$). Between the SV group and the stretch-PTFE group and between the two types prosthetic grafts no differences were found in RD in the arterial inflow segments. In the venous outflow segments no differences in RD were found between the three groups.

Figure 5: Bar graphs demonstrating the results of measurements of Relative Distension (RD) assessed at the defined sites in the grafts and host vessels, for all graft types (*, $p<0.05$). Data are presented as medians and 25th and 75th percentile intervals. (sPTFE = stretch PTFE, SV = saphenous vein, ePTFE = expanded PTFE)

Relative Distension

Area Increase

AI was higher in the SV group in the proximal and distal graft segments as compared to both the ePTFE group and the stretch-PTFE group ($p<0.05$) (Table II, Figure 6), indicating a better ability to store blood volume during the cardiac cycle. Between the ePTFE group and the stretch-PTFE group no differences were found in the graft. In the arterial inflow segment no differences were found in AI between the groups. In the venous outflow segments AI was lower in the efferent vein segments in the ePTFE group as compared to both the SV-group and the stretch-PTFE group ($p<0.05$).

Figure 6: Bar graphs demonstrating the results of measurements of Area-Increase (AI) assessed at defined sites in the grafts and host vessels, for all graft types (*, p<0.05). Data are presented as medians and 25th and 75th percentile intervals. (sPTFE = stretch PTFE , SV = saphenous vein, ePTFE = expanded PTFE)

Figure 7: Scatterplot showing the relation between vein RD and distal graft RD (only PTFE grafts). A positive correlation was observed (p<0.05), suggesting vein graft interaction.

Graft-vein interaction.

To study a possible interaction between the dimensions and elastic properties of the vein and the PTFE (stretch and expanded) graft, we correlated the RD values of the vein segment with RD values for the distal graft segment (Figure 7). A positive relation between vein RD values and distal graft RD values was found ($p < 0.05$, linear regression). Between vein diameter and distal graft distension a negative correlation was observed ($p < 0.05$).

DISCUSSION

The major finding of the present study is that mismatch in elastic properties around the arterial and venous anastomoses of human AV-fistulas has a different character. In the arterial anastomosis a decline in AI (area increase) was found, indicating a sudden drop in the capacity to store blood volume. By contrast, in the venous anastomosis an increase in the capacity to store blood volume was encountered in the stretch-PTFE and ePTFE group. Since an abrupt increase in the capacity to store blood volume is associated with disturbances in flow, like enhanced particle residence time and flow stasis, it may contribute to the predisposition of intimal hyperplasia to develop at the venous anastomoses²¹. The stasis of flow, caused by the sudden increase in capacity to store blood volume, may lead to a concentration of platelets and upstream released growth factors at a point just distal from the venous anastomoses, possibly initiating smooth muscle cell replication^{3,23,39}. The repeated damage of the graft due to puncturing may enhance the upstream release of growth factors. The expected stagnation of flow at this particular site indicates a low wall shear rate at this point. This condition is associated with increased intimal thickening, as has been demonstrated in many studies²²⁻²⁸. The mechanism by which low shear conditions promote intimal thickening may be of influence on the growth control function of the endothelium²⁹⁻³¹.

A limitation of the use of elastic parameters like AI and RD is that pulse pressure differences are not corrected for. However, the comparison of the elastic parameters within the grafts and around the anastomoses is independent of the local pulse pressure, assuming that pulse pressure does not change over the short segment considered.

A recent study demonstrated that pulsatile stress initiates vascular smooth muscle cell (VSMC) proliferation in human saphenous vein but not in internal mammary artery¹⁶, indicating a difference between the proliferative response of the arterial and venous smooth muscle cells to this physical stress. This suggests that the excessive stretch at the arterial and venous anastomoses due to the mismatch in elastic properties may lead to proliferation of VSMC at the venous anastomosis, but not at the arterial anastomosis. A possible contributing mechanism may be the different effect of elongation on the excessive stress encountered at the arterial and venous anastomosis, respectively.

A noncompliant stiff graft does not have the possibility to elongate during the cardiac cycle. This indicates that elongation of the arterial inflow segment during the systolic phase approximates the arterial segment and the graft, thereby decreasing the effect of mechanical stretch. At the venous anastomoses the elongation leads to separation, leading to increase in mechanical stretch. This indicates that mismatch in compliance may have a different effect on the extent of the excessive mechanical stretching encountered at the arterial and distal anastomoses, respectively, due to the opposite effect of elongation at these sites. The new stretch-PTFE graft has the possibility to elongate during the cardiac cycle and may therefore decrease the effect of excessive mechanical stretching at the venous anastomosis⁴⁰. This may lead to better primary patency rates for stretch-PTFE grafts as compared to non-compliant grafts.

Placement of an interposition graft in the construction of a vascular access leads to an enormous pressure- and flow increase at the site of the efferent vein, which may contribute to the development of vessel wall injury⁴¹. Mismatch around the anastomoses of interposition grafts, leads to impedance mismatch, thereby reducing the amount of pulsatile energy^{20,42}. Therefore, placement of a stiff graft probably reduces the pulsatile stress and flow in the efferent vein, which may prevent the occurrence of intimal stenoses at this site. On the other hand, placement of a stiff graft increases peri-anastomotic mismatch in elastic properties, which may promote intimal thickening at the anastomoses, especially at the venous site, as is indicated by our data. The complexity of hemodynamics in AV fistulas is illustrated by the interaction of the distal end of the graft and the venous outflow segment. Since the elastic properties of the graft and the vein were assessed quite close to the anastomoses (within 1 cm), the elastic properties of the graft may be influenced by the elastic properties of the vein. This is supported by the finding that a positive correlation was found between RD values obtained in the PTFE grafts and in the vein, as depicted in figure 8. The dimensions of the vein may also be of influence on the distension of the distal graft segments. The pulse wave arriving at a large diameter vein will not be reflected at this point. However, when the distal graft segment is anastomosed to a small diameter vein, part of the pulse wave will be reflected, leading to higher distensions in the distal graft. The presence of this type of interaction at the venous anastomosis of graft AV fistulas is suggested by the negative correlation between vein diameter and distal graft relative distension (Figure 8).

Recently, modifications of the conventional techniques for the construction of distal PTFE anastomoses have been introduced in peripheral bypass surgery, using anastomotic vein patches or cuffs^{43,44}. Patency rates of ePTFE grafts anastomosed with these vein patches or cuffs at the crural vessels are promising⁴⁴. Theoretically, the vein patches or cuffs may improve the abrupt change in compliance encountered in conventional anastomoses. However, the interposition of collars or cuffs with a large diameter may result in more flow disturbance due to the increase of hemodynamic capacity (equation 2). Therefore, in the design of modifications in anastomotic vein patches, the achievement of a gradual change in elastic properties in the anastomoses needs to be addressed.

Figure 8: Scatterplot demonstrating the relation between vein diameter and distal graft relative distension. A negative correlation was observed between these parameters ($p < 0.05$).

The higher values in the graft for RD and AI for the SV group as compared to the prosthetic materials indicate advantageous hemodynamic properties of the autologous graft²⁰. These findings confirm the results of a previous study from our institution, comparing ePTFE and saphenous vein bypass grafts in the lower limb. The beneficial elastic properties of the saphenous vein grafts resulted in a better match around the venous anastomosis, which may be an explanation for the better primary patency rates for saphenous vein grafts as compared to ePTFE grafts when used as a vascular access². The new stretch-PTFE graft showed slightly better elastic properties as compared to the ePTFE graft. In the stretch-PTFE graft the area-increase (AI) was maintained along the graft, in contrast to the ePTFE grafts, where a decrease was observed. This indicates a better preservation of pulsatile energy of the stretch-PTFE grafts as compared to the ePTFE grafts. However, this did not result in a better match around the venous anastomoses. A higher AI was found in the venous outflow segments in the stretch-PTFE group, indicating a better management of pulsatile flow at this point. This may contribute to a better hemodynamic environment of this flow segment, possibly leading to a decrease in the development of flow-induced hyperplastic lesions in the outflow segments of stretch-PTFE grafts⁷.

In conclusion, different characteristics were found for peri-anastomotic mismatch in elastic properties between the arterial and venous anastomoses of graft AV-fistulas in hemodialysis patients. A decline in area-increase, representing the capacity of the vessel wall to store blood volume, was found in the arterial anastomosis, while in the venous anastomosis an augmentation in area-increase was observed in the prosthetic grafts. Since the latter condition may lead to flow disturbances, like increased particle residence time and low shear stress, it may contribute to the predisposition of intimal hyperplasia to develop at the venous anastomoses.

REFERENCES

1. Windus DW. Permanent vascular access: a nephrologist's view. *Am J Kidney Dis* 1993; 21:457-471.
2. Jenkins AMcL, Buist TAS, Glover SD. Medium-term follow-up of forty autogenous vein and forty polytetrafluorethylene (Gore-Tex) grafts for vascular access. *Surgery* 1980; 88:667-672.
3. Swedberg SH, Brown BG, Sigley R, Wight TN, Gordon D, Nicholls SC. Intimal fibromuscular hyperplasia at the venous anastomosis of PTFE grafts in hemodialysis patients. *Circulation* 1989; 80:1726-1736.
4. Rehktter M, Nicholls S, Ferguson M, Gordon D. Cell proliferation in human arteriovenous fistulas use for hemodialysis. *Arterioscler Thromb* 1993; 13:609-617.
5. Sottiurai VS, Yao JST, Flinn WR, Batson RC. Intimal hyperplasia and neointima: An ultrastructural analysis of thrombosed grafts in humans. *Surgery* 1983; 93:809-817.
6. Madras PN, Ward CA, Johnson WR, Singh PI. Anastomotic hyperplasia. *Surgery* 1981; 90:922-923.
7. Bassiouny HS, White S, Glagov S, Choi E, Giddens DP, Zarins CK. Anastomotic intimal hyperplasia: Mechanical injury or flow induced. *J Vasc Surg* 1992; 15:708-717.
8. LoGerfo FW, Quist WC, Nowak MD, Crawshaw HM, Haudenschild CC. Downstream anastomotic hyperplasia. A mechanism of failure in Dacron arterial grafts. *Ann Surg* 1983; 197:479-483.
9. Echave V, Koornick AR, Haimov M, Jacobson JH. Intimal hyperplasia as a complication of the use of the polytetrafluoroethylene graft for femoral-popliteal bypass. *Surgery* 1979; 86:791-798.
10. Sottiurai VS, Sue SL, Feinberg II EL, Bringaze WL, Tran AT, Batson RC. Distal anastomotic intimal hyperplasia: Biogenesis and etiology. *Eur J Vasc Surg* 1988; 2:245-256.
11. Zacharias RK, Kirkman TR, Kenagy RD, Bowen-Pope DF, Clowes AW. Growth factor production by polytetrafluoroethylene vascular grafts. *J Vasc Surg* 1988; 7:606-610.
12. Liu MW, Roubin GS, King III SB. Restenosis after coronary angioplasty. Potential biologic determinants and role of intimal hyperplasia. *Circulation* 1989; 79:1374-1387.
13. Abbott WM, Megerman J, Hasson JE, L'Italien G, Warnock DF. Effect of compliance mismatch on vascular graft patency. *J Vasc Surg* 1987; 5:376-382.
14. Okuhn SP, Connelly DP, Calakos N, Ferrell L, Man-Xiang P, Goldstone J. Does compliance mismatch alone cause neointimal hyperplasia? *J Vasc Surg* 1989; 9:35-45.
15. Kidson IG, Abbott WM. Low compliance and arterial graft occlusion. *Circulation* 1978; 58:11-14.
16. Predel HG, Yang Z, Von Segesser L, Turina M, Buhler FR, Luscher TF. Implications of pulsatile stretch on growth of saphenous vein and mammary artery smooth muscle. *Lancet* 1992; 340:878-879.
17. Friedman MH. Some atherosclerosis may be a consequence of the normal adaptive response to shear. *Atherosclerosis* 1990; 82:193-196.
18. Leung DYM, Glagov S, Mathews MB. Cyclic stretching stimulates synthesis of matrix components by arterial smooth muscle cells in vitro. *Science* 1976; 191:475-477.
19. Sottiurai VS, Kollross P, Glagov S, Zarins CK, Mathews MB. Morphologic alteration of cultured arterial smooth muscle cells by cyclic stretching. *J Surg Res* 1983; 35:490-497.
20. Baird RN, Abbott WM. Pulsatile blood-flow in arterial grafts. *Lancet* 1976; October 30:948-950.
21. Stewart SFC, Lyman DJ. Effects of a vascular graft/natural artery compliance mismatch on pulsatile flow. *J Biomech* 1992; 25:297-310.
22. Zarins CK, Bomberger RA, Glagov S. Local effects of stenoses: Increased flow velocity inhibits atherogenesis. *Circulation* 1981; 64:11221-11227.

23. Zarins CK, Giddens DP, Bharadvaj BK, Sottiurai VS, Mabon RF, Glagov S. Carotid bifurcation atherosclerosis. Quantitative correlation of plaque localization with flow velocity profiles and wall shear stress. *Circ Res* 1983; 53:502-514.
24. Asakura T, Karino T. Flow patterns and spatial distribution of atherosclerotic lesions in human coronary arteries. *Circ Res* 1990; 66:1045-1066.
25. Cashin WL, Sanmarco ME, Nessim SA, Blankenhorn DH. Accelerated progression of atherosclerosis in coronary vessels with minimal lesions that are bypassed. *New Eng J Med* 1984; 311:824-828.
26. Sterpetti AV, Cucina A, Santoro L, Cardillo B, Cavallaro A. Modulation of arterial smooth muscle cell growth by hemodynamic forces. *Eur J Vasc Surg* 1992; 6:16-20.
27. Kohler TR, Jawien A. Flow affects development of intimal hyperplasia after arterial injury in rats. *Arterioscler Thromb* 1992; 12:963-971.
28. Kohler TR, Kirkman TR, Kraiss LW, Zierler BK, Clowes AW. Increased blood flow inhibits neointimal hyperplasia in endothelialized vascular grafts. *Circ Res* 1991; 69:1557-1565.
29. Buga GM, Gold ME, Fukuto JM, Ignarro LJ. Shear stress-induced release of nitric oxide from endothelial cells grown on beads. *Hypertension* 1991; 17:187-193.
30. Hsieh HJ, Li NQ, Frangos JA. Shear stress increases endothelial platelet-derived growth factor mRNA levels. *Am J Physiol* 1991; 260:H642-H646.
31. Cooke JP, Rossitch E, Andon NA, Loscalzo J, Dzau VJ. Flow activates an endothelial potassium channel to release an endogenous nitrovasodilator. *J Clin Invest* 1991; 88:1663-1671.
32. Hoeks APG, Brands PJ, Smeets FAM, Reneman RS. Assessment of distensibility of superficial arteries. *Ultrasound Med Biol* 1990; 16:121-128.
33. Hofstra L, Ermers EJM, Langeveld APM, Hoeks APG, Kitslaar PJEHM. Vessel Wall Doppler Tracking: A new technique to assess vessel wall compliance. *J Vasc Tech* 1993; 17:65-68.
34. O'Rourke MF. Arterial stiffness, systolic blood pressure, and logical treatment of arterial hypertension. *Hypertension* 1990; 15:339-347.
35. Reneman RS, Merode van T, Brands PJ, Hoeks APG. Inhomogeneities in arterial wall properties under normal and pathological conditions. *J Hypertens* 1992; 10:S35-S39.
36. Hofstra L, Willigers JM, Huvers FC, et al. Short-term variations in elastic properties of a muscular artery in humans. *Clin Sci* 1993; 50:472-480.
37. Hofstra L, Bergmans DCJJ, Hoeks APG, Tordoir JHM, Kitslaar PJEHM. Inhomogeneities in elastic properties of saphenous vein arterial bypass grafts in humans. *Eur J Vasc Surg* 1994; 40:300-310.
38. Fillinger MF, Reinitz ER, Schwartz RA, et al. Graft geometry and venous intimal-medial hyperplasia in arteriovenous loop grafts. *J Vasc Surg* 1990; 11:556-566.
39. Nerem RM, Girard PR. Hemodynamic influences on vascular endothelial biology. *Toxicologic Pathology* 1990; 18:572-582.
40. Pärsson H, Jundzill W, Hallberg E, Thörne J, Norgren L. Acute thrombogenicity and 4 weeks healing properties of a new stretch-ePTFE graft. *Eur J Vasc Surg* 1993; 7:63-70.
41. Fry DL. Certain chemorheologic considerations regarding the blood vascular interface with particular reference to coronary artery disease. *Circulation* 1969; XXXIX and XI:IV38-IV59.
42. O'Rourke MF. Steady and pulsatile energy losses in the systemic circulation under normal conditions and in simulated arterial disease. *Cardiovasc Res* 1967; 1:313-326.
43. Tyrrell MR, Chester JF, Vipond MN, Clarke GH, Taylor RS, Wolfe JHN. Experimental evidence to support the use of interposition vein collars/patches in distal PTFE anastomoses. *Eur J Vasc Surg* 1990; 4:95-101.
44. Taylor RS, Loh A, McFarland RJ, Cox M, Chester JF. Improved technique for polytetrafluoroethylene bypass grafting: long-term results using anastomotic vein patches. *Br J Surg* 1992; 79:348-354.

CHAPTER 8

THE DIMENSIONS AND ELASTIC PROPERTIES OF HUMAN SAPHENOUS VEIN GRAFTED INTO THE ARTERIAL CIRCULATION ARE STABLE OVER TIME

L Hofstra, DCJJ Bergmans, APG Hoeks, JHM Tordoir, PJEHM Kitslaar.
(submitted)

ABSTRACT

Background. Structural changes of saphenous veins used as a bypass grafts include generalised intimal thickening and replacement of vascular smooth muscle cells by collagen and fibroblasts in the media. This may be the result of adaptation of the vein to arterial pressure and flow conditions. We investigated the possible influence of the structural adaptations on local elastic properties of venous peripheral bypass grafts in patients. In addition, local diameter changes of the grafts with time were measured.

Methods. In 17 patients receiving a reversed saphenous vein bypass graft the local dimensions and elastic properties of the graft and native arteries were assessed with the use of Vessel Wall Doppler Tracking (VWDT). Measurements were performed at 2, 6, 12 and 26 weeks after construction of the bypass.

Results. In local strain (in %) and cross-sectional area increase (mm^2) no changes with time were observed in the graft and native arterial system. In the proximal graft segments an increase in diameter with time was observed (4.6 vs 5.2 mm, $p < 0.05$). Also in the outflow artery an increase in diameter with time was observed (4.9 vs 5.1 mm, $p < 0.05$).

Conclusions. Our data indicate that the elastic properties of the saphenous vein grafted into the arterial circulation remain stable within the first 6 months after implantation. An increase in diameter in the proximal graft segments of the reversed saphenous vein grafts was observed. Since in the most proximal part of the graft the highest flow velocities are present, the structural dilatation may reflect an adaptational response to decrease shear rate at this point.

INTRODUCTION

The grafting of a saphenous vein into the arterial circulation leads to sudden changes in pressure as well as flow conditions. Local increase in pressure and flow occurs, and these physical stresses become pulsatile. This abrupt change in pressure and flow

conditions may lead to functional as well as structural adaptations¹⁻⁴. The ability of large arteries to undergo structural adaption to changed flow and/or pressure conditions has been demonstrated by many investigators⁵⁻⁹. The role of the endothelium in the adaptional responses seems crucial. An increase in flow results in a functional and structural adaptation^{8,9}, initiated by an increase in shear stress, probably sensed by the endothelial cells^{10,11}. In vitro studies indicate that the mechanism by which structural adaptation occurs, may be shear stress induced release of growth factors, like PDGF and TGF- α , by the endothelium, possibly regulated by a shear responsive element¹²⁻¹⁵. Functional changes after alteration in shear rates are suggested by several studies^{6,7,13,14,16}. High shear stress leads to an increase in the release of NO and substance-P by the endothelium, both resulting in vasodilatation. The time course of functional changes of the saphenous vein after grafting have been studied in a canine model^{4,17}. A decrease in endothelium dependent relaxation after the administration of acetylcholine (Ach) was reported indicating endothelial dysfunction of the venous graft, which has also been demonstrated in the saphenous vein when used as coronary bypass in humans^{17,18}. However, responses to the administration of two other endothelium dependent relaxants showed a recovery to control values after 2 weeks, suggesting at least a partial recovery of the endothelial function.

Structural changes of the human saphenous vein after grafting can be divided into acute and chronic changes^{2,19-23}. Acute changes include endothelial denudation and deposition of platelets, fibrin and leucocytes in the intima and edema, smooth muscle cell necrosis and hemorrhage in the media. Chronic changes include generalized intimal thickening and the partial replacement of VSMC by collagen and fibroblasts in the media.

The observed functional and structural responses of the saphenous vein graft may alter the dimensions and the elastic properties with time and therefore may influence the hemodynamic function of the graft¹⁴. The time course of changes in elastic properties of saphenous veins has been reported in animal studies^{2,24}. An upward shift was found in the pressure-diameter relation of veins after grafting with the use of an in vitro technique to assess the elastic properties². In vivo assessment of the elastic properties of veins grafted into the arterial circulation of dogs, demonstrated no apparent change with time. This was confirmed in a clinical study². However, the compliance values assessed at different sites along the graft were averaged, thereby excluding the evaluation of changes in local elastic properties due to tapering effects²⁵.

In previous studies by our group the dimensions and elastic profiles of newly inserted human reversed saphenous vein grafts were reported, as assessed with the use of Vessel Wall Doppler Tracking, at defined sites in the graft and the arterial in- and outflow segments^{25,32}. The use of both a fractional parameter (strain) and a capacitive parameter (compliance) was essential to give a proper mechanical profile of the reversed vein grafts. The aim of the present study was to investigate the local changes of the dimensions and elastic properties of these vein grafts within the first 6 months after implantation.

MATERIALS AND METHODS

Patients

Patients with arterial insufficiency of the lower extremity as manifested by disabling claudication, restpain and/or tissue loss and receiving a reversed saphenous vein bypass graft were considered candidates for our study. Thirty-two consecutive patients received a reversed saphenous vein bypass during the study period (January 1992 - August 1993). In three patients the vein grafts occluded within one month after implantation due to outflow problems. In seven patients a stenotic process occurred within the study period of six months. The presence of a possible stenosis was detected by Duplex scan before every session and was based on generally accepted Duplex criteria²⁶. The diagnosis of a hemodynamically significant stenosis was confirmed by angiogram. Although these grafts were surgically revised to maintain patency, these patients were excluded from the study since a stenosis could alter the hemodynamics of the graft by influencing pulse wave reflections^{27,28}. In five patients the assessment of elastic properties was not possible at the first session due to hematoma formation and/or wound leakage. The remaining 17 patients were measured at strict time intervals after operation (2,6,12 and 26 weeks post-operatively). The autologous grafts were implanted using standard techniques and were inserted in an end-to-side fashion with running 6-0 Prolene sutures.

Measurements of the elastic properties

The elastic properties of the vein graft and host artery were assessed with the use of Vessel Wall Doppler Tracking (VWDT) (chapter 3 and 4)^{29,30}.

From distension (in μm) and diameter (in mm) the following parameters were calculated:

$$\text{Relative Distension (RD)} = \frac{\Delta D}{D} * 100 (\%) \quad (1)$$

$$\text{Area-Increase (AI)} = \frac{\pi \Delta D * D}{2} \quad (2)$$

Relative distension (RD) represents the local strain of the vessel wall. Area-increase (AI) represents the capacity of the vessel to store and to release temporarily blood volume during the cardiac cycle³¹. Usually these parameters are corrected for pulse pressure, resulting in the distensibility coefficient (DC) and compliance coefficient (CC), respectively. In a previous study we correlated DC and CC values of the common femoral artery calculated with local invasive pressure measurements and digital beat-to-beat pressure measurements (Finapres, volume clamp technique), obtained in patients undergoing peripheral bypass surgery (chapter 5)²⁵. Although we observed reasonable correlation between DC values calculated with local invasive

pressure measurements and Digital Finapres measurements ($r=0.93$ for DC and $r=0.92$ for CC) we decided not to correct for pulse pressure, since pulse pressure behavior in a bypass may be unpredictable. Validation studies of the VWD system demonstrated that the variation coefficient for distension of thick-walled medium-sized arteries, like the common femoral artery, lies between 10-15% for young healthy volunteers (chapter 4). This substantial variation in elastic properties probably depends on the vasomotor tone of the active vessel. In diameter less variation was observed in this study, illustrated by a variation coefficient of 2-4%. The variation coefficient for the carotid artery, an elastic artery, lies between 5-10%³². We believe that the variation coefficient of the elastic parameters for diseased femoral arteries and vein bypasses will not exceed the variation of healthy muscular arteries, and probably lies between 5-10%.

In each graft measurements were performed in the proximal graft segment (1 cm from the proximal anastomosis), in the middle segment, and in the distal graft segment (1 cm from the distal anastomosis). The inflow artery was measured 1 cm from the proximal anastomosis, the outflow artery was measured 1 cm from the distal anastomosis. The patients were measured at these sites 2, 6, 12, and 26 weeks after operation.

At the same predefined sites in the graft and the arterial in- and outflow arterial segments, peak systolic velocity was determined by placement of a Doppler sample midstream in the vessel.

Patients were measured in the supine position after an acclimatisation period of 20 minutes. All patients gave a written informed consent before entering the study. The study was approved by the institutional medical ethical committee.

ANALYSIS

Data are presented as medians and 25th and 75th percentile intervals. The changes in time for the values for the different parameters obtained at the defined sites were analysed with the use of the Friedman-test (two-way ANOVA). A p-value of less than 0.05 was accepted as level of statistical significance.

RESULTS

The patient characteristics are summarized in table I. Note that most grafts were anastomosed at a point below-knee.

Table I: Patient characteristics and sites of anastomoses.

		Reversed SV (N=17)
Male		12 (69%)
Female		5 (31%)
Age	(Yrs)	68 (SD 8.8)
Diabetic		5 (29%)
Clinical stage :		
	II	5 (28.6%)
	III	6 (38.1%)
	IV	6 (33.3%)
Proximal Anastomosis :		
	Common Femoral Artery	11 (66.7%)
	Superficial Femoral Artery	6 (33.3%)
Distal Anastomosis :		
	Popliteal Artery (above knee)	4 (19.0%)
	Popliteal Artery (below knee)	11 (66.7%)
	Crural	2 (14.3%)

Diameter

The results of diameter measurement are summarized in table II and depicted in Figure 1. As the graph shows, an increase in diameter with time could be observed in the proximal graft segment ($p=0.05$) and the arterial outflow segment ($p=0.05$, Friedman two-way Anova). In the mid- and proximal graft segments and the arterial inflow segments no change in diameter was observed.

Table II: Results of the measurements of diameter (mm) at the predefined sites in the graft (proximal, mid and distal) and in the in- and outflow arterial segments, assessed at 2, 6, 12 and 26 weeks after implantation of the saphenous vein graft. Data are presented as medians and 25th and 75th percentile intervals. (* = $p<0.05$, significant change with time, Friedman, two-way Anova).

	2 weeks	6 weeks	12 weeks	26 weeks
Site:				
Inflow	7.4 (7.1-10.6)	7.4 (6.0-10.8)	8.1 (5.6-12.0)	8.6 (6.9-10.7)
Proximal *	4.6 (3.3-5.8)	4.3 (3.2-5.1)	4.7 (3.9-5.4)	5.2 (4.7-5.9)
Mid	5.7 (4.8-6.7)	5.0 (4.1-5.7)	5.6 (4.1-6.6)	5.1 (4.4-6.2)
Distal	6.3 (4.9-7.1)	5.2 (4.9-5.8)	6.2 (4.8-7.1)	5.8 (4.7-6.9)
Outflow *	4.9 (3.7-6.1)	5.3 (3.7-8.0)	5.6 (5.3-6.2)	5.1 (4.1-6.0)

Figure 1: Median values of diameter (DI) assessed after 2, 6, 12 and 26 weeks at the proximal (Prox), mid (Mid) and distal (Dist) segments of the grafts and the inflow (Infl) and outflow (Outfl) arterial segments. (* = $p<0.05$, Friedman two-way ANOVA).

Relative distension

In RD, representing the local strain of the graft, no change with time was observed at the three different sites in the grafts (Table III, Figure 2). Also in the arterial inflow- and outflow segments no changes in RD were shown within 6 months after operation. segments no change in diameter was observed.

Table III: Results of the measurements of relative distension (RD in %) at the predefined sites in the graft (proximal, mid and distal) and in the in- and outflow arterial segments, assessed at 2, 6, 12 and 26 weeks after implantation of the saphenous vein graft. Data are presented as medians and 25th and 75th percentile intervals. No significant changes with time were observed.

	2 weeks	6 weeks	12 weeks	26 weeks
Site:				
Inflow	2.6 (1.8-3.9)	2.5 (1.8-3.9)	2.0 (1.4-3.1)	2.4 (1.8-3.7)
Proximal	3.9 (2.4-4.6)	5.2 (2.8-7.5)	3.4 (2.9-5.2)	3.5 (2.2-4.7)
Mid	2.0 (1.5-3.4)	2.7 (1.9-4.4)	3.3 (2.1-4.8)	2.4 (1.5-2.8)
Distal	1.5 (0.9-3.0)	2.1 (1.6-3.6)	1.3 (1.1-2.3)	1.6 (1.0-2.1)
Outflow	1.4 (1.1-2.9)	1.2 (0.7-1.8)	2.0 (1.3-2.3)	1.9 (1.2-2.1)

Figure 2: Bar graphs demonstrating the results of measurements of relative distension (RD) assessed at the defined sites in the grafts and host arteries. Data are presented as medians. No significant changes with time were observed.

Area-increase

In AI, representing the capacity of the vessel wall to store blood volume during the cardiac cycle, no change was observed in the proximal graft segments ($p=0.19$, Friedman two-way Anova). In the mid- and distal graft segments and the arterial in- and outflow segments also no changes in AI with time could be observed (Table IV, Figure 3).

Table IV: Results of the measurements of cross-sectional area-increase (mm^2) at the predefined sites in the graft (proximal, mid and distal) and in the in- and outflow arterial segments, assessed at 2, 6, 12 and 26 weeks after implantation of the saphenous vein graft. Data are presented as medians and 25th and 75th percentile intervals. No significant changes with time were observed.

	2 weeks	6 weeks	12 weeks	26 weeks
Site:				
Inflow	2.2 (1.1-4.0)	2.1 (1.4-3.3)	1.9 (1.0-2.8)	2.3 (1.3-3.6)
Proximal	1.6 (1.1-2.0)	2.0 (1.1-3.4)	1.7 (1.2-2.1)	1.8 (1.2-2.3)
Mid	1.4 (1.0-1.7)	1.3 (1.0-2.3)	1.9 (1.3-2.4)	1.2 (0.8-1.3)
Distal	0.9 (0.6-1.4)	1.0 (0.9-1.9)	0.8 (0.5-1.4)	0.8 (0.6-1.3)
Outflow	0.8 (0.5-1.4)	0.6 (0.3-1.1)	1.2 (0.8-1.3)	0.8 (0.6-1.0)

Figure 3: Bar graphs demonstrating the results of measurements of the cross-sectional Area Increase (AI) assessed at defined sites in the grafts and host arteries. Data are presented as medians. No significant changes with time were observed.

Peak Systolic Velocity

Peak systolic velocity in the proximal graft segments was higher as compared to PSV values in the mid- and distal portions of the grafts ($p < 0.05$, Mann-Whitney). With time, no changes were observed in as well the different sites of the graft as the arterial in- and outflow segments (Figure 4).

Figure 4: Median values of peak systolic velocity (PSV, cm/s) measured at the defined sites in the graft and the in- and outflow arterial segments.

DISCUSSION

The goal of the current study was to evaluate changes over time of the dimensions and the local elastic properties of human saphenous vein after grafting into the arterial circulation. The major finding of our study is that the diameter of the proximal graft segments and of the arterial outflow segments increased within the first 6 months after operation, indicating an adaptation of these segments to the new flow and pressure conditions. Furthermore, the local elastic properties, as represented by relative distension and area-increase, are preserved with time along the entire graft. These findings indicate that the hemodynamic function of the graft is sustained within the first 6 months after operation.

Dimensions

Numerous studies suggested that changes in flow conditions of large arteries result in structural and/or functional adaptation in order to maintain the shear stress at a certain level^{15-9,12-14,16}. High shear stress levels therefore should lead to diameter increase of the

vessel. Due to the reversal of the saphenous vein in the construction of a bypass, the proximal part of the graft is the narrowest segment. This leads to high flow velocities in this part of the graft resulting in high shear stress levels sensed by the endothelial cells³³, which is illustrated by the significantly higher peak systolic velocity values in the proximal graft segment as compared to the mid- and distal segments. Our finding that the diameter of the proximal graft segments increases in the post-operative period, therefore, supports previous findings that vessels tend to maintain the shear rate at a certain level. Studies evaluating structural adaptation due to flow changes were always performed on large conduit arteries. Our data indicate that the human saphenous vein also has the ability to adapt to flow conditions. A possible mechanism may be the shear-stress dependent release of PDGF or TGF by the endothelium¹²⁻¹⁴. In the outflow arterial segments also an increase in diameter was observed, which may also be related to the increase in flow velocity in this segment, due to the restoration of flow after the construction of the bypass. The midgraft- and distal graft segments did not undergo considerable dimensional changes with time, which may be related to the lower flow velocities in these graft segments.

Studies evaluating the chronic structural changes after grafting demonstrated generalized intimal thickening, necrosis and replacement of vascular smooth muscle cells (VSMC) by fibroblasts and collagen^{2,19-23}. This indicates an increase in wall thickness and loss of elastic components in the vessel wall, both of which are associated with a decrease of the elastic properties. Our data suggest that the local elastic properties assessed at predefined sites in the saphenous vein graft do not undergo considerable changes in the first 6 months after placement in the arterial system. The stable elastic properties of the saphenous vein graft with time may indicate that the decrease in elastic properties due to structural changes may be compensated for by an increase due to flow-induced functional adaptation. Another reason for the unchanged elastic properties of the graft with time may be the restricted ability of human saphenous vein to adapt to the new flow and pressure conditions. In studies evaluating the structure of the saphenous vein before grafting, pre-existing disease was observed in a large percentage of patients, consisting of intimal thickening and fibrosis of the media^{34,35}. We observed similar findings in spare saphenous vein segments of patients undergoing peripheral bypass surgery. Functional testing of the saphenous vein segments in an organ chamber between an isometric force transducer revealed that segments showing media fibrosis had a blunted response to serotonin, indicating loss of contractile reactivity of VSMC in these segments (unpublished data). Therefore, functional adaptation in the graft due to flow increase may also be blunted, and may explain the lack of improvement of elastic properties of the grafts. The pre-existing pathology of the saphenous vein may also restrict the ability of the vein graft to undergo structural adaptation with time. In vein grafts with pre-existing media fibrosis proliferation of VSMC may also be limited. Since proliferation of VSMC is regarded as the most important feature in structural adaptation of large vessels, the ability of the diseased segments to adapt to the new environment may therefore be blunted³⁶⁻⁴⁰. Another factor restricting the ability of the saphenous vein bypass grafts to show

functional adaptation to the new hemodynamic environment may be that about one third of the patients were diabetics. Diabetes mellitus leads to non-enzymatic changes of the elastic components of the media, leading to media sclerosis. This process leads to stiffening of vessels, probably limiting functional changes⁴¹.

The finding that the elastic properties of saphenous vein grafts remain stable with time supports the results from animal studies and from a clinical study^{2,24}. In these studies, the fractional increase in diameter per unit of pressure was assessed at different segments of the graft. These values were averaged to obtain a mean value for the graft. No apparent change with time was observed in both the animal and the clinical study. Our data indicate that large differences in RD, a fractional parameter, occur along the graft, due to the effect of reversed tapering. Therefore, we believe that changes of the local strain have to be assessed per graft segment and cannot be averaged from values obtained at different sites in the graft.

Studies investigating the functional changes of canine saphenous vein after grafting, demonstrated a partial recovery of endothelium-dependent relaxation¹⁷. This recovery occurred in the first two weeks after grafting. Since the first measurements of the elastic properties were performed two weeks after operation, we may have missed the early functional improvement. Also in another study recovery of the endothelium dependent relaxations were observed⁴.

In summary, no apparent changes were observed in the local elastic properties of human saphenous vein grafted into the arterial circulation. An increase in diameter with time was observed in the proximal graft segment, suggesting remodeling in this part of the graft. A possible mechanism explaining the increase in diameter may be high shear rate induced structural adaptation. In the arterial outflow segments also an increase in diameter with time was observed, which also may be a flow induced adaptational process. Our data suggest that the hemodynamic function of saphenous vein grafts in humans is stable during the first 6 months after operation.

REFERENCES

1. Dobrin PB. On the roles of deformation, tension, and wall stress as critical stimuli eliciting myointimal/medial hyperplasia. *J Vasc Surg* 1991; 581-582.
2. Dobrin PB, Littooy FN, Golan J, Blakeman B, Fareed J. Mechanical and histological changes in canine vein grafts. *J Surg Res* 1988; 44:259-265.
3. Dobrin PB, Littooy FN, Endean ED. Mechanical factors predisposing to intimal hyperplasia and medial thickening in autogenous vein grafts. *Surgery* 1989; 105:393-400.
4. Miller VM, Reigel MM, Hollier LH, Vanhoutte PM. Endothelium-dependent responses in autogenous femoral veins grafted into the arterial circulation of the dog. *J Clin Invest* 1987; 80:1350-1357.
5. Cashin WL, Sanmarco ME, Nessim SA, Blankenhorn DH. Accelerated progression of atherosclerosis in coronary vessels with minimal lesions that are bypassed. *New Eng J Med* 1984; 311:824-828.
6. Rubanyi GM, Freay AD, Kauser K, Johns A, Harder DR. Mechanoreception by the endothelium: Mediators and mechanisms of pressure- and flow-induced vascular responses. *Blood Vessels* 1990; 27:246-257.
7. Tesfamariam B, Cohen RA. Inhibition of adrenergic vasoconstriction by endothelial cell shear stress. *Circ Res* 1988; 63:720-725.
8. Kamiya A, Togawa T. Adaptive regulation of wall shear stress to flow change in the canine carotid artery. *Am J Physiol* 1980; 239:H14-H21.
9. Langille BL, O'Donnell F. Reductions in arterial diameter produced by chronic decreases in blood flow are endothelium-dependent. *Science* 1986; 231:405-407.
10. Helmlinger G, Geiger RV, Schreck S, Nerem RM. Effects of pulsatile flow on cultured vascular endothelial cell morphology. *J Biomech Eng* 1991; 113:123-131.
11. Nerem RM, Girard PR. Hemodynamic influences on vascular endothelial biology. *Toxicologic Pathology* 1990; 18:572-582.
12. Hsieh HJ, Li NQ, Frangos JA. Shear stress increases endothelial platelet-derived growth factor mRNA levels. *Am J Physiol* 1991; 260:H642-H646.
13. Cooke JP, Rossitch E, Andon NA, Loscalzo J, Dzau VJ. Flow activates an endothelial potassium channel to release an endogenous nitrovasodilator. *J Clin Invest* 1991; 88:1663-1671.
14. Buga GM, Gold ME, Fukuto JM, Ignarro LJ. Shear stress-induced release of nitric oxide from endothelial cells grown on beads. *Hypertension* 1991; 17:187-193.
15. Resnick N, Collins T, Atkinson W, Bonthron DT, Dewey CF, Gimbrone MA. Platelet-derived growth factor B chain contains a cis-acting fluid shear-stress-responsive element. *Proc Natl Acad Sci U S A* 1993; 90:4591-4595.
16. Ralevic V, Milner P, Hudlicka O, Kristek F, Burnstock G. Substance P is released from the endothelium of normal and capsaicin-treated rat hind-limb vasculature, in vivo, by increased flow. *Circ Res* 1990; 66:1178-1183.
17. Ishii T, Okadome K, Komori K, Odashiro T, Sugimachi K. Natural course of endothelium-dependent and -independent responses in autogenous femoral veins grafted into the arterial circulation of the dog. *Circ Res* 1993; 72:1004-1010.
18. Lüscher TF, Diederich D, Siebeenman R, et al. Difference between endothelium-dependent relaxation in arterial and venous coronary bypass grafts. *New Eng J Med* 1988; 319:462-467.
19. Dilley RJ, McGeachie JK, Prendergast FJ. A review of the histologic changes in vein-to-artery grafts, with particular reference to intimal hyperplasia. *Arch Surg* 1988; 123:691-696.
20. Fuchs JCA, Mitchener JS, Hagen PO. Postoperative changes in autologous vein grafts. *Ann Surg* 1978; 188:1-15.
21. Dilley RJ, McGeachie JK, Tennant M. Vein to artery grafts: a morphological and histochemical study of the histogenesis of intimal hyperplasia. *Aust N Z J Surg* 1992; 62:297-303.

22. Adcock GD, Adcock OT, Wheeler JR, et al. Arterialization of reversed autogenous vein grafts: Quantitative light and electron microscopy of canine jugular vein grafts harvested and implanted by standard or improved techniques. *J Vasc Surg* 1987; 6:283-295.
23. Milroy CM, Scott DJA, Beardt JD, Horrocks M, Bradfield JWB. Histological appearances of the long saphenous vein. *J Pathology* 1989; 159:311-316.
24. Cambria RP, Brewster DC, Hasson J, Megerman J, Warnock DF, Abbott WM. The evolution of morphologic and biomechanical changes in reversed and in situ vein grafts. *Ann Surg* 1987; 205:167-174.
25. Hofstra L, Bergmans DCJJ, Hoeks APG, Tordoir JHM, Kitslaar PJEHM. Assessment of inhomogeneities in elastic properties of in situ and reversed saphenous vein grafts in humans. *Eur J Vasc Surg* 1994; 8:100-105.
26. Buth J, Disselhoff B, Sommeling C, Stam L. Color-flow duplex criteria for grading stenosis in infrainguinal vein grafts. *J Vasc Surg* 1991; 14:716-728.
27. O'Rourke MF. Arterial stiffness, systolic blood pressure, and logical treatment of arterial hypertension. *Hypertension* 1990; 15:339-347.
28. Kelly RP, Gibbs HH, O'Rourke MF, et al. Nitroglycerin has more favourable effects on left ventricular afterload than apparent from measurement of pressure in a peripheral artery. *Eur Heart J* 1990; 11:138-144.
29. Hoeks APG, Brands PJ, Smeets FAM, Reneman RS. Assessment of distensibility of superficial arteries. *Ultrasound Med Biol* 1990; 16:121-128.
30. Hofstra L, Ermers EJM, Langeveld APM, Hoeks APG, Kitslaar PJEHM. Vessel Wall Doppler Tracking: A new technique to assess vessel wall compliance. *J Vasc Tech* 1993; 17:65-68.
31. Reneman RS, Merode van T, Brands PJ, Hoeks APG. Inhomogeneities in arterial wall properties under normal and pathological conditions. *J Hypertens* 1992; 10:S35-S39.
32. Hofstra L, Willigers JM, Huvers FC, et al. Short-term variations in elastic properties of a muscular artery in humans. *Clin Sci* 1993; 50:472-480.
33. Belkin M, Mackey WC, McLaughlin R, Umphrey SE, O'Donnell TF. The variation in vein graft flow velocity with luminal diameter and outflow level. *J Vasc Surg* 1992; 15:991-999.
34. Marin ML, Veith FJ, Panetta TF, et al. Saphenous vein biopsy: A predictor of vein graft failure. *J Vasc Surg* 1993; 18:407-415.
35. Panetta TF, Marin ML, Veith FJ, et al. Unsuspected preexisting saphenous vein disease: An unrecognized cause of vein bypass failure. *J Vasc Surg* 1992; 15:102-112.
36. Clowes AW, Reidy MA. Prevention of stenosis after vascular reconstruction: Pharmacologic control of intimal hyperplasia - A review. *J Vasc Surg* 1991; 6:885-891.
37. Clowes AW, Clowes MM, Fingerle J, Reidy MA. Kinetics of cellular proliferation after arterial injury. V. Role of acute distension in the induction of smooth muscle cell proliferation. *Lab Invest* 1989; 60:360-364.
38. Liu MW, Berk BC. Restenosis following coronary balloon angioplasty. Role of smooth muscle cell proliferation. *Trends Cardiovasc Med* 1991; 1:107-111.
39. Liu MW, Roubin GS, King III SB. Restenosis after coronary angioplasty. Potential biologic determinants and role of intimal hyperplasia. *Circulation* 1989; 79:1374-1387.
40. Schwartz SM, Ross R. Cellular proliferation in atherosclerosis and hypertension. *Prog Cardiovasc Dis* 1984; 26:355-372.
41. Cairns SA, Woodcock JP, Marshall AJ. Early arterial lesions in maturity onset diabetes mellitus detected by an ultrasound technique. *Diabetologia* 1978; 14:107-111.

CHAPTER 9

ANASTOMOTIC INTIMAL HYPERPLASIA IN PROSTHETIC ARTERIO-VEINUS FISTULAS FOR HEMODIALYSIS IS ASSOCIATED WITH INITIAL HIGH SHEAR RATE AND NOT WITH MISMATCH IN ELASTIC PROPERTIES

L Hofstra, DCJJ Bergmans, KML Leunissen, APG Hoeks, PJEHM Kitslaar, MJAP Daemen, and JHM Tordoir
(submitted)

ABSTRACT

Background. Stenotic intimal thickening at the venous end of prosthetic arterio-venous (AV) fistulas for hemodialysis has been associated with peri-anastomotic mismatch in elastic properties and low shear rates. In a prospective way, we investigated the role of these factors on the occurrence of intimal hyperplasia in prosthetic AV fistulas in hemodialysis patients.

Methods. In 24 hemodialysis patients, the elastic properties were assessed in the distal graft and the outflow vein post-operatively, with the use of Vessel Wall Doppler Tracking (VWDT), a non-invasive ultrasound technique. In addition, local peak shear rate (PSR) was calculated from diameter (VWDT) and peak systolic velocity. The initial mismatch around the venous anastomoses and local PSR were correlated with the occurrence of stenoses during follow-up (2 years). The detection of a stenosis was performed with both Duplex ultrasound and angiography. Furthermore, surgically derived stenotic tissue samples were histologically examined with respect to endothelial cell coverage of the lumen.

Results. In 4 cases a stenosis developed in the venous anastomosis, in 8 cases in the venous outflow segment and in 4 cases at both sites. A better initial match in elastic properties around the venous anastomosis was observed in the fistulas developing a stenosis at this site as compared to the non-stenotic fistulas ($p < 0.05$). The initial local PSR values at the site of the later stenosis were higher in the fistulas developing a stenosis as compared to the non-stenotic fistulas ($p < 0.05$). Histologic examination of stenotic tissue samples derived from these patients ($n = 11$) revealed only partial coverage of the lumen with endothelial cells (mean 48%).

Conclusions. The occurrence of stenosis in prosthetic AV fistulas for hemodialysis in or adjacent to the venous anastomoses is associated with a high initial peak shear rate

but not with a mismatch in elastic properties. A possible mechanism may be a denuding effect of excessive high shear on venous endothelial cells, thereby initiating intimal thickening.

INTRODUCTION

Arterio-venous fistulas constructed with the use of interposition grafts are often complicated by the development of occlusive intimal thickening, mostly occurring at the venous anastomoses¹⁻⁴. The repair of these stenotic lesions leads to a mean hospitalisation time of one month a year for these patients^{2,3}. The reasons for the high incidence of stenoses developing at the venous anastomoses are unclear. Studies evaluating the influence of varying levels of shear rate on intimal thickening reported conflicting data. In animal studies increased intimal thickening was found in areas with low shear rates, indicating that low shear rates may induce the intimal thickening⁵⁻¹². A possible factor contributing to low shear rates at a site just downstream from the venous anastomosis may be the sudden increase in the capacity to store blood volume at this site¹³, leading to flow disturbances like flow separation and stasis¹⁴. On the other hand, very high shear rates damage the endothelial cells covering the lumen and may therefore result in vessel wall injury, possibly initiating intimal hyperplasia^{15,16}. The latter concept is supported by animal studies evaluating intimal thickening in the venous outflow segments of arterio-venous fistulas. The most severe intimal thickening was observed in areas with the highest flow velocities^{17,18}.

A second mechanical condition affecting intimal growth may be a difference in elastic properties around anastomoses, also called compliance mismatch^{12,19-22}. This leads to points of excessive mechanical stretching, possibly resulting in vessel wall injury. Furthermore, excessive mechanical stretching may induce vascular smooth muscle cell replication, as has been demonstrated in *in vitro* models²³⁻²⁶. However, an association between intimal thickening and compliance mismatch has not been convincingly demonstrated so far²⁰.

In the present study, vessel wall elastic properties were measured around the venous anastomoses of polytetrafluoroethylene (PTFE) arterio-venous fistulas in hemodialysis patients. These measurements were performed within two weeks after the construction of the AV fistula, with the use of Vessel Wall Doppler Tracking (VWDT)^{27,28}. In addition, peak systolic velocity (PSV) was measured at defined sites in the graft and in the inflow- and outflow vessels. From diameter, measured with VWDT, and PSV the peak shear rate (PSR) was calculated, an approximation of the actual shear rate exerted on the endothelial cells covering the lumen of the vessel wall. Furthermore, stenotic tissue specimens were harvested in the patients developing a stenosis during follow-up necessitating surgical repair. These specimens were histologically evaluated with respect to the endothelial cell coverage of the lumen, determined with the use of immunostaining (ULEX Europeus). The aim of the present study was to define the

influence of both mismatch in elastic properties around the anastomosis and peak shear rate on the occurrence of occlusive intimal hyperplasia in or adjacent to the venous anastomoses in hemodialysis patients with a prosthetic AV fistula. In addition, the possible influence of shear rate on the presence of endothelial cells covering the lumen was evaluated.

MATERIALS AND METHODS

Patient clinical profile

Twenty-four consecutive patients receiving a hemodialysis graft AV fistula with the use of a PTFE prosthesis (WL GORE, Flagstaff, Arizona, USA) entered the study (December 1991 - December 1993). Generally, the construction of the fistula with the use of interposition grafts was a secondary procedure. In 4 patients the construction of a graft AV fistula was a primary choice, due to the absence of suitable veins in the forearm for the creation of a Brescia Cimino fistula. In all patients the arterial and venous anastomoses were constructed at the level of the elbow. The brachial artery served as the inflow segment; as outflow segments served the basilic, the cephalic or the cubital veins. The grafts were implanted in a loop configuration in the forearm.

Measurement of the elastic properties.

The diameter and distension of the vessel and graft segments were measured with the use of Vessel Wall Doppler Tracking (VWDT)^{27,28}. This system combines B-mode imaging (ATL, Ultramark IV, Englewood, USA) and off-line data processing, which offers the possibility to assess the distension and diameter accurately at defined sites in the graft, the inflow artery and the efferent vein. In vitro studies showed that VWDT is able to detect distensions as low as 1 μm , indicating that the expected low distension values of the PTFE grafts can be measured properly. The recording time of 4 seconds per registration allows measurement of distension (ΔD) and internal diastolic diameter (D) during 3-5 consecutive heartbeats. The values of these consecutive beats were averaged in this study. From the diameter and distension the following parameters were calculated:

$$\text{Relative Distension (RD)} = \frac{\Delta D}{D} * 100 (\%) \quad (1)$$

$$\text{Area-Increase (AI)} = \frac{\pi \Delta D * D}{2} \quad (2)$$

Relative distension gives the strain of the vessel during the cardiac cycle. Area-increase represents the capacity of the vessel to store blood volume during the cardiac cycle. Generally, the parameters are corrected for pulse pressure resulting in the

distensibility coefficient and compliance coefficient, respectively²⁹. In studies evaluating the pressure distribution in graft AV fistulas it was shown that mean pressure and pulse pressure decrease along the graft and within the anastomoses^{17,18}. The extent of the decrease varies with graft geometry. This implicates that valid pressure values can only be obtained with the use of invasive pressure measurements at all sites investigated. Since this would be unethical, we decided not to use pulse pressure in the equations.

Diameter and distension were assessed in the distal graft 1 cm proximal from the venous anastomoses, in the venous anastomoses, and in the efferent vein, 1 cm distal from the venous anastomosis (Figure 1). The positioning of the cursor line at these particular sites was facilitated by repeated imaging of the anastomotic complex with B-mode.

Figure 1: Schematic drawing of a venous anastomosis of an arteriovenous fistula. Most of the stenoses developed at the site of the anastomoses (A) or in the venous outflow segment just distal from the anastomosis, within 1 cm (V).

Since we attempted to evaluate the influence of mismatch in elastic properties around anastomoses on the occurrence of intimal hyperplasia, we indexed the values for AI and RD around the anastomoses, resulting in the match factor. Also for diameter a match factor was calculated, in order to obtain insight in the size differences around the anastomoses. The match factors for the different parameters were calculated in the following way:

$$\text{Match factor} = \text{value efferent vein} / \text{value distal graft} \quad (3)$$

A perfect match is considered to be present when the match factor is one. A mismatch value higher than 1 implicates an increase in the parameter within the anastomotic area, while a value lower than 1 implicates a decrease of the parameter in the direction of blood flow.

Determination of Peak Shear Rate (PSR)

PSR was calculated from peak systolic velocity (PSV) and diameter. Peak systolic velocity was measured by placement of a Doppler sample midstream in the vessel at an angle of 60°. At the same location diameter was assessed with the use of VWDT. Diameter measurements with the use of VWDT have been validated in a previous study. The variation coefficient for diameter varied between 3 to 5%, indicating low variation in diameter assessment with the use of this system³⁰. PSV and diameter were assessed in the distal graft (1 cm from the anastomosis), within the venous anastomoses, and in the venous outflow segments (within 1 cm from the anastomoses). From PSV and diameter, PSR was calculated with the use of a modification of the Haagen-Poiseuille formula:

$$\text{Peak Shear Rate (PSR)} = n * \text{Peak Systolic Velocity/Diameter} \quad (4)$$

The value for n depends on the shape of the velocity profile. When a parabolic laminar flow pattern is present, the value for n is 4. When the flow pattern is blunted, the value for n increases. Detailed information on velocity patterns in vivo was not available, and we therefore decided to use $n=4$ in the calculation of PSR. Especially in the anastomotic area high blood velocities are present, probably resulting in complex flow patterns. Therefore, PSR can only give an approximation of the actual shear forces exerted on the inner layer of the vessel wall. Some investigators calculate mean shear rate from diameter and volume flow, also assessed with the use of ultrasound^{5,6}. Besides difficulties in calculating volume flow with the use of ultrasound in AV fistulas, we believe that peak shear rate may give a better representation of the pulsatile character of the stress exerted on the endothelial cells³¹⁻³³. In another study (chapter 11) we observed a negative correlation between peak shear rate and subendothelial VSMC proliferation, indicating that peak shear rate may have physiological importance. Therefore it was decided to use this parameter in the current study.

Most patients were measured within 2 weeks after the construction of the AV fistula. In some cases the measurements were delayed due to wound leakage or hematoma formation. These patients were measured within 4 weeks after operation. All patients were measured in the supine position after an acclimatisation period of 20 minutes. At strict time intervals (2, 6 and 12 weeks, and every 3 months thereafter) a Duplex scan of the arteriovenous fistula was performed for the detection of possible stenoses. When a stenosis was detected with Duplex, an angiogram was made to confirm the diagnosis. The study was approved by the medical ethical committee of the University Hospital of Maastricht. All patients gave written informed consent before entering the study.

Histologic examinations

Stenotic tissue was harvested for histologic evaluation in patients developing a stenosis necessitating repair with an interposition graft. These tissue specimens were fixed overnight in 10% phosphate buffered formalin, processed according to routine histologic

procedures and paraffin-embedded. Serial sections (5 μm thick) were cut and placed on gelatin coated glass slides. To identify endothelial cells, Ulex Europeus agglutinin (1:500 dilution, Vector Labs, Burlingame, CA, USA) was used. After blocking of the endogenous peroxidase activity with 0.3% H_2O_2 , sections were incubated with the lectine antibody (1:250, 45 min., DAKO) after predigestion with pepsin (1 mg/ml, 30 min.). Subsequently, the sections were incubated with an anti-Ulex Europeus lectine peroxidase complex (1:100, DAKO). As the chromogen 3,3' diaminobenzidine was used. As a negative control spare saphenous vein segments were used, obtained from patients undergoing peripheral bypass surgery. The degree of endothelial cell coverage was estimated by dividing the circumference of the lumen covered by endothelial cells by the total luminal circumference.

ANALYSIS

Data are presented as medians and 25th and 75th percentile intervals. To compare differences between groups, the Mann-Whitney-U test was used. The local initial mechanical and flow conditions of the fistulas developing a stenosis at either the venous anastomosis or the efferent vein were compared with the conditions of the fistulas without a stenosis at these locations. Since the initial mechanical and flow conditions were assessed at well defined sites in the anastomotic complex, good correlation of these local conditions with the site of the later stenosis was possible. To test which of the mechanical factors showed an independent correlation with the occurrence of intimal thickening, multiple regression analysis was performed. Mechanical factors showing a correlation with the occurrence of intimal thickening with a p-value < 0.10 were included in a stepwise multiple regression model. A p-value lower than 0.05 was accepted as level of statistical significance.

RESULTS

In the 24 patients, a total of 20 stenoses developed in 16 fistulas in the perianastomotic area at the venous site during the follow-up period (mean 12.2 ± 8.7 (SD) months); in 8 patients no fistula stenosis was found during the study period. In 4 fistulas a stenosis occurred only at the venous anastomosis. In 8 cases a stenosis developed only in the efferent vein adjacent to the anastomosis (within 1 cm from the anastomosis, Figure 1) and in 4 fistulas stenoses developed at both sites. This results in a total of 8 stenoses in the venous anastomosis and 12 in the efferent vein adjacent to the anastomosis (Table I). In all patients with a stenosis repair of the stenotic area was necessary to reestablish a sufficient flow for dialysis treatment, indicating that the stenosis was occlusive or subocclusive. Stenoses occurring at sites more downstream of the efferent vein (n=4) were not included in this study, because of the difficulty to correlate the initial mechanical and flow information with the site of these stenoses.

Table I: Patient Characteristics.

Age	61 (39-68)
Sex	11 male/13 female
Primary/Secondary Fistula	4/20

Stenoses at the venous anastomoses

The initial relative distension (RD) values in the venous anastomoses were higher (Table II) in the fistulas developing a stenosis in the venous anastomoses as compared to the fistulas without stenosis development at this site (2.5 vs 1.0 %, $p<0.05$). No differences in initial diameter match and diameter at the anastomoses were observed between the groups. The initial match factors for AI and RD (ARM and RDM respectively, Table II, Figure 2) were closer to 1 in the stenotic fistulas as compared to the fistulas without a stenosis at this site (1.5 vs 5.1 for AI and 1.2 vs 2.6 for RD, $p<0.05$), indicating a better initial match in elastic properties around the venous anastomoses in stenotic fistulas. Higher initial values for PSR were found in the anastomoses of stenotic fistulas as compared to the fistulas without a stenosis (308 vs 212 1/s, $p<0.05$, Figure 3).

Table II: Values for the different parameters measured at the venous anastomosis, two weeks after construction, for the AV fistulas with a later stenosis at this point (stenotic) and the fistulas without a stenosis (non-stenotic) at this site during follow-up. RD = relative distension, DIAM = diameter, ARM = match factor area increase, RDM = match factor relative distension, PSR = Peak shear rate. Values are presented as medians and 25th and 75th percentile intervals.

VENOUS ANASTOMOSIS

	Stenotic (n=8)	Nonstenotic (n=16)	P-value
RD (%)	2.5 (1.5-3.2)	1.0 (0.4-1.6)	0.02
DIAM (mm)	4.1 (3.6-4.4)	3.5 (3.1-4.5)	0.82
ARM	1.5 (0.8-2.7)	5.1 (2.8-8.4)	0.005
RDM	1.2 (0.9-2.2)	2.6 (2.0-5.0)	0.01
PSR (1/s)	308 (261-346)	212 (168-276)	0.02

Figure 2: Bar graph showing the initial values for the match factor in area increase (ARM) for the fistulas developing a stenosis (stenotic) and non-stenotic fistulas. In the fistulas with a later stenosis at the venous anastomoses as well as fistulas with a later stenosis in the venous outflow segment a more ideal match in area increase was observed (*, $p<0.05$, Mann-Whitney).

Table III: Values for the different parameters measured in the efferent vein just downstream the anastomosis, two weeks after construction, for the arterio-venous fistulas with and without a later stenosis at this point. RD = relative distension, DIAM = diameter, ARM = match factor area increase, RDM = match factor relative distension, PSR = Peak shear rate. Values are presented as medians and 25th and 75th percentile intervals.

EFFERENT VEIN

	Stenotic (n=12)	Non-stenotic (n=12)	P-value
RD (%)	4.2 (3.4-14.3)	3.7 (1.3-6.5)	0.34
DIAM (mm)	4.2 (2.9-8.3)	6.3 (4.2-7.9)	0.34
ARM	1.8 (0.7-2.6)	3.7 (2.2-6.2)	0.04
RDM	1.2 (0.9-2.2)	2.5 (1.8-4.2)	0.05
PSR (1/s)	272 (143-331)	99 (60-219)	0.04

Stenoses in the efferent vein

No differences in initial relative distension and vein diameter (Table III) were found between the fistulas developing a stenosis in the efferent vein (within 1 cm from the anastomosis) and the fistulas without a stenosis at this site. Also no differences in diameter match around the venous anastomoses were observed between the groups. The initial match values for AI and RD (ARM and RDM, Table III, Figure 2) around the venous anastomosis were closer to 1 in the fistulas with a later stenosis in the efferent vein as compared to the non-stenotic fistulas (1.8 vs 3.7 for ARM and 1.2 vs 2.5 for RDM, $p < 0.05$). Initial PSR values in the efferent vein were higher in the fistulas with a later stenosis as compared to the fistulas without a stenosis in this area (272 vs 99 1/s, $p < 0.05$, Figure 3).

Figure 3: Bar graph showing the initial values for PSR at the anastomoses and venous outflow segments for the fistulas developing a stenosis (stenotic) and the non-stenotic fistulas. Higher values for the fistulas with a later stenosis were observed at both sites (*, $p < 0.05$, Mann-Whitney).

Regression analysis

Multiple regression analysis showed that an increase in initial PSR values was independently correlated with the occurrence of (sub)-occlusive intimal thickening ($r = -0.55$, $p < 0.05$) at both the venous anastomosis and the efferent vein. In addition, an RD match (RDM) closer to 1 around the venous anastomoses, indicating a better match, was correlated with the occurrence of a stenosis ($r = 0.41$, $p < 0.05$). For match in AI (ARM) no independent correlation was found.

Histologic examinations

In 9 cases the repair of the stenotic venous anastomosis was performed with the use of an interposition graft, which allowed examination of the stenotic lesions. A total of 11 stenotic specimens were derived from the anastomoses or the efferent vein adjacent to the anastomoses. In all samples the stenotic lesion consisted mainly of smooth muscle cells and extra-cellular matrix material. The mean percentage of endothelial cells covering the lumen was 48%. In the control saphenous vein specimens (n=15) a mean percentage of endothelial cells covering the lumen was 95%.

DISCUSSION

The present study contains two major findings. First, the occurrence of stenotic intimal hyperplasia in prosthetic AV fistulas in humans is associated with initial high peak shear rate at the site of the later stenosis, indicating that high shear rates may initiate the development of stenotic intimal hyperplasia at the venous anastomoses of AV fistulas for hemodialysis. Second, a better initial match in strain around the venous anastomosis was found in fistulas developing a stenosis in this area, indicating that mismatch in elastic properties around anastomoses of prosthetic graft AV fistulas does not contribute to the initiation of stenotic intimal hyperplasia.

Peak shear rate and intimal thickening

In the stenotic fistulas higher PSR values were observed at the venous anastomoses and the outflow venous segments as compared to the non-stenotic fistulas. This suggests that high shear rates may initiate the development of intimal thickening in arterio-venous fistulas used for hemodialysis. These observations support the results of earlier studies investigating the effects of flow on cultured endothelial cells^{15,16}. Swelling, deformation, and finally denudation was observed in the cultured EC, suggesting serious EC injury. In animal studies evaluating the effect of varying levels of flow on the occurrence of intimal thickening at the venous outflow segment of AV fistulas, more intimal thickening was observed in fistulas with the highest flow velocities at this site^{18,34}. This indicates that intimal thickening may be initiated by excessive high shear levels, which is also suggested by the data from the present study. This seems to be in contrast with numerous other investigations, reporting that low shear rate promotes intimal thickening^{5-12,35}. Several points have to be considered when comparing the effects of shear rate on intimal thickening, including the range of shear rate investigated, the possible difference between the regulation of intimal thickening in animals and humans, and the different response of arteries and veins to unphysiological shear rate. In some studies, the effect of a decreased flow was compared with the effect of normal arterial flow^{36,37}. This indicates that the range of shear rates investigated lies between normal and low, which is a different range as compared to the level in our study. In normal arteries peak systolic velocities between 60 to 120 cm/s are present, leading to substantially lower shear rate levels. In the venous anastomoses of an AV-fistula, peak systolic velocity values as high as 300-350 cm/s can be measured,

indicating excessive high shear rate levels. This implies that the shear rate magnitude found in the present study is far beyond physiological values. Therefore, comparison of the results of different studies has to be performed with regard to the range of shear rate values investigated.

Another point which has to be considered is the use of animal models in studies investigating the effects of flow on intimal thickening. In animal models intimal hyperplasia does not proceed to subocclusive or occlusive stenotic lesions, but is restricted in its extent^{38,39}. The changes in intimal thickening that are found with flow variation in animal models may be just an adaptational process. The factors involved in the regulation of intimal thickening in experimental animals may therefore be dissimilar to the factors that induce occlusive intimal lesions in humans. Our study overcomes this disadvantage, since we studied (sub)occlusive intimal hyperplasia in hemodialysis patients. A third consideration may be the use of different vessel types in studying the proliferative process. Some investigators use PTFE grafts, and measure the thickness of the neo-intima at different flow levels^{5,35,40}. Another model which is often used is the balloon catheter denudated carotid artery in the rat or baboons^{6,41-43}. In the present study, the intimal stenotic lesions occurred at the venous anastomoses or the venous outflow segment, which is adjacent to the PTFE graft but not in the graft. The excessive high shear rates in arteriovenous fistulas may be tolerated by arteries and arterial endothelial cells, while veins and venous endothelial cells undergoing these shear rate magnitudes may be damaged. This is supported by the data from the histologic examinations of the stenotic samples derived from the venous anastomoses of the fistulas in our patient group. The mean degree of endothelial cell coverage of the lumen was 48 %, indicating partial denudation of endothelial cells in these vessel segments, indicative for a denudating effect of excessive shear rates at the venous anastomoses and efferent vein of prosthetic AV fistulas in humans.

In low shear areas intimal thickening probably is induced by endothelial cells, sensing the forces exerted by blood flow⁴⁴⁻⁵¹. In case of low shear, endothelial cells release growth promoting factors, leading to intimal thickening, in order to reestablish a certain level of shear rate^{8,10,11,36}. The reversed process, structural adaptive vessel dilatation with increased flow, has also been described^{52,53}. Our working hypothesis is that within a certain spectrum of shear rate, intimal thickening (leading to a decrease in vessel diameter) or dilatation in response to changes in flow are physiologic adaptational processes to regulate the shear rate levels sensed by endothelial cells^{10,11,52,53}. However, when shear rate levels exceed the upper limit of this spectrum, denudation may occur, leading to intimal thickening rather than adaptive structural dilatation. Despite similar flow conditions, structural dilatation is observed at the arterial site of AV fistulas whereas intimal thickening develops at the venous site. This suggests that the upper limit of this spectrum probably is higher for arteries than for veins.

Mismatch in elastic properties and intimal thickening

The compliance mismatch hypothesis was first postulated in the seventies, and stems from numerous observations made by vascular surgeons, that anastomotic areas of peripheral bypasses as well as arteriovenous fistulas are predisposed to the development of intimal hyperplasia^{1-4,12,19-22,54}. It was hypothesized that a mismatch in elastic properties leads to a point of excessive mechanical stretching resulting in wall injury and in the initiation of vascular smooth muscle cell (VSMC) replication^{20,23,25,26,55}. Furthermore, an increase in compliance is associated with flow disturbances like flow stasis and enhanced particle residence time, which may promote the development of intimal thickening¹⁴. In a previous study we observed a rise in area-increase within the venous anastomoses, while area-increase dropped in the arterial anastomoses¹³. Since the first condition is associated with flow disturbances and enhanced particle residence time, we postulated that this might be the reason for the predisposition of intimal hyperplasia to develop at the venous anastomoses¹⁴. The results of the present study suggest that the postulated mechanism does not play a role in the development of stenotic intimal hyperplasia at the venous anastomoses of PTFE AV fistulas. Our data clearly indicate that an increased mismatch in elastic properties does not lead to the occurrence of intimal hyperplasia in AV fistulas in humans. On the contrary, we found that an increased mismatch in strain seems to protect against the development of a stenosis, since according to our definition, the peri-anastomotic match in strain (RD) was far from the ideal value in the non-stenotic groups. A possible explanation may be that an advantageous effect of a proper match in elastic properties around the venous anastomoses of stenotic AV fistulas is preponderated by the influence of excessive high peak shear rate resulting from the construction of an arteriovenous fistula.

Usually, relative distension (RD) and area-increase (AI) are corrected for pulse pressure, resulting in cross-sectional compliance and distensibility coefficient, respectively. There are two reasons why it was decided to use RD and AI instead of the corrected parameters. First, invasively obtained pressure measurements in loop AV-fistulas in dogs have demonstrated that a large drop in pressure is present along the grafts and within the anastomoses¹⁷. This indicates that non-invasively obtained pulse pressure values obtained at one site in the vascular system cannot give a proper estimation of pulse pressure in the whole system. A second reason for the use of AI and RD is that the compliance mismatch hypothesis is based on the thought that peri-anastomotic differences in strain and differences in cross-sectional area increase, possibly lead to local mechanical stress and/or flow disturbances²⁰. The force leading to strain and cross-sectional area increase (i.e. pulse pressure) is of a lesser importance. Therefore, it seems justified to use RD and AI to represent the local elastic properties and to calculate peri-anastomotic mismatch in elastic properties.

Clinical Implications

According to the findings of the present study, one should aim at decreasing initial peak shear rate at the venous site in constructing an AV-fistula. This may be achieved

with the use of 4-7 mm tapered grafts which results in decreased peak systolic velocities at the venous end of graft AV fistulas¹⁷. In addition, the use of wide venous cuffs in the construction of the venous anastomosis may help in decreasing peak shear rate in the efferent vein just downstream the anastomosis^{56,57}. Our data indicate that differences in elastic properties around anastomoses cannot be regarded as a causal factor in the occurrence of anastomotic intimal thickening in human graft AV fistulas. Therefore, the use of stiff prosthetic graft materials is not discouraged by the outcome of our study. Application of the findings of the present study to other clinical manifestations of intimal hyperplasia, like restenosis after PTCA and stenosis in peripheral or coronary bypass grafts, has to be performed with caution^{42,58,59}. Flow characteristics in coronary arteries and peripheral bypass grafts differ substantially from AV fistula flow, indicating quite different magnitudes of shear rate. In addition, intimal thickening in coronary arteries following PTCA may differ from intimal thickening observed at graft-vein junctions in AV fistulas, due to differences in structure and function between arteries and veins/PTFE grafts^{26,60}.

In summary, the results of the present study indicate that the occurrence of occlusive or subocclusive intimal thickening at the venous anastomoses and venous outflow segments of AV fistulas for hemodialysis access is associated with initial high peak shear rate at these sites. This suggests that high shear rate may initiate intimal hyperplasia in AV fistulas in humans. A possible mechanism may be an injurious effect of high shear on the venous endothelial cells leading to denudation, which is supported by the finding that the lumen of the stenotic lesions derived from these patients is only partly covered with endothelial cells. In addition, in the fistulas developing stenoses a more ideal initial match in elastic properties around the venous anastomoses was found. This strongly argues against a contribution of mismatch in elastic properties to intimal thickening in graft AV fistulas in humans. To improve patency rates of arteriovenous fistulas with the use of interposition grafts attention has to be focused on decreasing initial peak shear rate at the venous anastomoses and the efferent vein.

REFERENCES

1. Windus DW. Permanent vascular access: a nephrologist's view. *Am J Kidney Dis* 1993; 21:457-471.
2. Jenkins AMcL, Buist TAS, Glover SD. Medium-term follow-up of forty autogenous vein and forty polytetrafluoroethylene (Gore-Tex) grafts for vascular access. *Surgery* 1980; 88:667-672.
3. Swedberg SH, Brown BG, Sigley R, Wight TN, Gordon D, Nicholls SC. Intimal fibromuscular hyperplasia at the venous anastomosis of PTFE grafts in hemodialysis patients. *Circulation* 1989; 80:1726-1736.
4. Tordoir JHM, Herman JM, Kwan TS, Diderich PM. Long-term follow-up of the polytetrafluoroethylene (PTFE) prosthesis as an arteriovenous fistula for hemodialysis. *Eur J Vasc Surg* 1988; 2:3-7.
5. Kohler TR, Kirkman TR, Kraiss LW, Zierler BK, Clowes AW. Increased blood flow inhibits neointimal hyperplasia in endothelialized vascular grafts. *Circ Res* 1991; 69:1557-1565.
6. Kohler TR, Jawien A. Flow affects development of intimal hyperplasia after arterial injury in rats. *Arterioscler Thromb* 1992; 12:963-971.
7. Zarins CK, Bomberger RA, Glagov S. Local effects of stenoses: Increased flow velocity inhibits atherogenesis. *Circulation* 1981; 64:II221-II227.
8. Zarins CK, Zatina MA, Giddens DP, Ku DN, Glagov S. Shear stress regulation of artery lumen diameter in experimental atherogenesis. *J Vasc Surg* 1987; 5:413-420.
9. Zarins CK, Giddens DP, Bharadvaj BK, Sottiurai VS, Mabon RF, Glagov S. Carotid bifurcation atherosclerosis. Quantitative correlation of plaque localization with flow velocity profiles and wall shear stress. *Circ Res* 1983; 53:502-514.
10. Friedman MH, Barger CB, Deters OJ, Hutchins GM, Mark FF. Correlation between wall shear and intimal thickness at a coronary artery branch. *Atherosclerosis* 1987; 68:27-33.
11. Friedman MH. Some atherosclerosis may be a consequence of the normal adaptive response to shear. *Atherosclerosis* 1990; 82:193-196.
12. Bassiouny HS, White S, Glagov S, Choi E, Giddens DP, Zarins CK. Anastomotic intimal hyperplasia: Mechanical injury or flow induced. *J Vasc Surg* 1992; 15:708-717.
13. Hofstra L, Bergmans DCCJ, Hoeks APG, Kitslaar PJEHM, Leunissen KML, Tordoir JHM. Mismatch in elastic properties around anastomoses of interposition grafts for hemodialysis access. *JASN* 1994; 45:45-55.
14. Stewart SFC, Lyman DJ. Effects of a vascular graft/natural artery compliance mismatch on pulsatile flow. *J Biomech* 1992; 25:297-310.
15. Fry DL. Certain chemorheologic considerations regarding the blood vascular interface with particular reference to coronary artery disease. *Circulation* 1969; XXXIX and XI:IV38-IV59.
16. Fry DL. Acute vascular endothelial changes are associated with increased blood velocity gradients. *Circ Res* 1968; 22:165-197.
17. Fillinger MF, Reinitz ER, Schwartz RA, et al. Graft geometry and venous intimal-medial hyperplasia in arteriovenous loop grafts. *J Vasc Surg* 1990; 11:556-566.
18. Fillinger MF, Reinitz ER, Schwartz RA, Resetarits DE, Paskanik AM, Bredenberg CE. Beneficial effects of banding on venous intimal-medial hyperplasia in arteriovenous loop grafts. *Am J Surg* 1989; 158:87-94.
19. Kidson IG, Abbott WM. Low compliance and arterial graft occlusion. *Circulation* 1978; 58:II-14.
20. Abbott WM, Megerman J, Hasson JE, L'Italien G, Warnock DF. Effect of compliance mismatch on vascular graft patency. *J Vasc Surg* 1987; 5:376-382.
21. Abbott WM, Megerman J. Adaptive responses of arteries to grafting. *J Vasc Surg* 1989; 9:377-379.
22. Baird RN, Abbott WM. Pulsatile blood-flow in arterial grafts. *Lancet* 1976; October 30:948-950.

23. Leung DYM, Glagov S, Mathews MB. Cyclic stretching stimulates synthesis of matrix components by arterial smooth muscle cells in vitro. *Science* 1976; 191:475-477.
24. Sumpio BE, Banes AJ, Levin LG, Johnson G. Mechanical stress stimulates aortic endothelial cells to proliferate. *J Vasc Surg* 1987; 6:252-256.
25. Leung DYM, Glagov S, Mathews MB. Elastin and collagen accumulation in rabbit ascending aorta and pulmonary trunk during postnatal growth. Correlation of cellular synthetic response with medial tension. *Circ Res* 1977; 41:316-323.
26. Predel HG, Yang Z, Von Segesser L, Turina M, Buhler FR, Luscher TF. Implications of pulsatile stretch on growth of saphenous vein and mammary artery smooth muscle. *Lancet* 1992; 340:878-879.
27. Hoeks APG, Brands PJ, Smeets FAM, Reneman RS. Assessment of distensibility of superficial arteries. *Ultrasound Med Biol* 1990; 16:121-128.
28. Hofstra L, Ermers EJM, Langeveld APM, Hoeks APG, Kitslaar PJEHM. Vessel Wall Doppler Tracking: A new technique to assess vessel wall compliance. *J Vasc Tech* 1993; 17:65-68.
29. Reneman RS, Merode van T, Brands PJ, Hoeks APG. Inhomogeneities in arterial wall properties under normal and pathological conditions. *J Hypertens* 1992; 10:S35-S39.
30. Hofstra L, Willigers JM, Huvers FC, et al. Short-term variations in elastic properties of a muscular artery in humans. *Clin Sci* 1993; 50:472-480.
31. Ojha M, Johnston KW, Cobbald RS. Evidence of a possible link between poststenotic dilatation and wall shear stress. *J Vasc Surg* 1990; 11:127-135.
32. Okadome K, Yukizane T, Mii S, Fukuda A, Sugimachi K. Correlation of longterm results of femoropopliteal autogenous vein grafts with shear stress variation. In situ or reversed graft? *J Cardiovasc Surg Torino* 1989; 30:932-935.
33. Nerem RM, Girard PR. Hemodynamic influences on vascular endothelial biology. *Toxicologic Pathology* 1990; 18:572-582.
34. Dobrin PB. On the roles of deformation, tension, and wall stress as critical stimuli eliciting myointimal/medial hyperplasia. *J Vasc Surg* 1991; 581-582.
35. Wallenstein S, Zucker C, Fleiss F. Some statistical methods useful in circulation research. *Circ Res* 1980; 47:1-9.
36. Langille BL, O'Donnell F. Reductions in arterial diameter produced by chronic decreases in blood flow are endothelium-dependent. *Science* 1986; 231:405-407.
37. Langille BL, Bendeck MP, Keeley FW. Adaptations of carotid arteries of young and mature rabbits to reduced carotid blood flow. *Am J Physiol* 1989; 256:H931-H939.
38. Ferrel M, Fuster V, Gold HK, Chesebro JH. Choosing appropriate experimental animal model for the prevention of restenosis. *Circulation* 1992; 85:1630-1631.
39. Clowes AW, Reidy MA. Prevention of stenosis after vascular reconstruction: Pharmacologic control of intimal hyperplasia - A review. *J Vasc Surg* 1991; 6:885-891.
40. Okadome K, Miyazaki T, Onohara T, Yamamura S, Sugimachi K. Hemodynamics and the development of anastomotic intimal hyperplasia of the polytetrafluoroethylene grafts in dogs. *Int Angiol* 1991; 10:238-243.
41. Clowes AW, Clowes MM, Fingerle J, Reidy MA. Kinetics of cellular proliferation after arterial injury. V. Role of acute distension in the induction of smooth muscle cell proliferation. *Lab Invest* 1989; 60:360-364.
42. Clowes AW, Clowes MM, Vergel SC, et al. Heparin and cilazapril together inhibit injury-induced intimal hyperplasia. *Hypertension* 1991; 18:1165-1169.
43. Clowes AW, Reidy MA, Clowes MM. Mechanisms of stenoses after arterial injury. *Lab Invest* 1983; 49:208-215.
44. Cooke JP, Rossitch E, Andon NA, Loscalzo J, Dzau VJ. Flow activates an endothelial potassium channel to release an endogenous nitrovasodilator. *J Clin Invest* 1991; 88:1663-1671.

45. Buga GM, Gold ME, Fukuto JM, Ignarro LJ. Shear stress-induced release of nitric oxide from endothelial cells grown on beads. *Hypertension* 1991; 17:187-193.
46. Hsieh HJ, Li NQ, Frangos JA. Shear stress increases endothelial platelet-derived growth factor mRNA levels. *Am J Physiol* 1991; 260:H642-H646.
47. Tesfamariam B, Cohen RA. Inhibition of adrenergic vasoconstriction by endothelial cell shear stress. *Circ Res* 1988; 63:720-725.
48. Rubanyi GM, Freay AD, Kauser K, Johns A, Harder DR. Mechanoreception by the endothelium: Mediators and mechanisms of pressure- and flow-induced vascular responses. *Blood Vessels* 1990; 27:246-257.
49. Resnick N, Collins T, Atkinson W, Bonthron DT, Dewey CF, Gimbrone MA. Platelet-derived growth factor B chain contains a cis-acting fluid shear-stress-responsive element. *Proc Natl Acad Sci U S A* 1993; 90:4591-4595.
50. DePaola N, Gimbrone MA, Davies PF, Dewey CF. Vascular endothelium responds to fluid shear stress gradients. *Arterioscler Thromb* 1992; 12:1254-1257.
51. Shen J, Luscinskas FW, Conolly A, Dewey CF, Gimbrone MA. Fluid shear stress modulates cytosolic free calcium in vascular endothelial cells. *Am J Cell Physiol* 1992; 31:c384-c390.
52. Kamiya A, Ando J, Shibata M, Masuda H. Roles of fluid shear stress in physiological regulation of vascular structure and function. *Biorheology* 1988; 25:271-277.
53. Kamiya A, Togawa T. Adaptive regulation of wall shear stress to flow change in the canine carotid artery. *Am J Physiol* 1980; 239:H14-H21.
54. Madras PN, Ward CA, Johnson WR, Singh PI. Anastomotic hyperplasia. *Surgery* 1981; 90:922-923.
55. Sottiurai VS, Kollross P, Glagov S, Zarins CK, Mathews MB. Morphologic alteration of cultured arterial smooth muscle cells by cyclic stretching. *J Surg Res* 1983; 35:490-497.
56. Taylor RS, Loh A, McFarland RJ, Cox M, Chester JF. Improved technique for polytetrafluoroethylene bypass grafting: long-term results using anastomotic vein patches. *Br J Surg* 1992; 79:348-354.
57. Tyrrell MR, Wolfe JHN. New prosthetic venous collar anastomotic technique: combining the best of other procedures. *Br J Surg* 1991; 78:1016-1017.
58. Liu MW, Roubin GS, King III SB. Restenosis after coronary angioplasty. Potential biologic determinants and role of intimal hyperplasia. *Circulation* 1989; 79:1374-1387.
59. Cashin WL, Sanmarco ME, Nessim SA, Blankenhorn DH. Accelerated progression of atherosclerosis in coronary vessels with minimal lesions that are bypassed. *New Eng J Med* 1984; 311:824-828.
60. Lüscher TF, Diederich D, Siebeenman R, et al. Difference between endothelium-dependent relaxation in arterial and venous coronary bypass grafts. *New Eng J Med* 1988; 319:462-467.

CHAPTER 10

MECHANICAL FACTORS PREDISPOSING TO INTIMAL HYPERPLASIA IN PERIPHERAL BYPASS GRAFTS CONSTRUCTED WITH THE USE OF AUTOLOGOUS VEIN; A PROSPECTIVE ANALYSIS

L Hofstra, APG Hoeks, JHM Tordoir, DCJJ Bergmans, MJAP Daemen,
PJEHM Kitslaar.
(submitted)

ABSTRACT

Background. Failure of peripheral bypass grafts constructed with the use of saphenous vein is mainly caused by the formation of stenosis due to intimal hyperplasia. Mechanical factors such as compliance mismatch around anastomoses and unphysiological shear rate have been implicated to contribute to the development of intimal hyperplasia but their actual role remains uncertain. The present study was conducted to investigate the role of these mechanical factors on the occurrence of intimal hyperplasia in peripheral bypasses.

Methods. Patients receiving a peripheral bypass for occlusive arterial disease of the leg were included (n=43). The bypass was constructed using either the reversed (n=28) or the in situ vein technique (n=15). Within 2 weeks after construction local dimensions and elastic properties of vessel segments were assessed in the graft and adjacent vessel segments at defined sites, with the use of Vessel Wall Doppler Tracking, a non-invasive ultrasound technique. A match factor was calculated by indexing elastic properties and dimensions of graft and native artery around the anastomoses. A perfect match is considered when the match factor is 1. In addition, peak shear rate (PSR) was calculated from diameter and peak systolic velocity (PSV), assessed at corresponding sites. The local initial match in elastic properties and dimensions and PSR values were correlated with the occurrence of stenoses during follow-up. The presence of stenoses was established with Duplex surveillance, made every 3 months after construction of the bypass. The occurrence of a stenosis was always confirmed by an angiogram.

Results. In 7 cases a stenosis developed at the proximal anastomosis, in 4 cases at the distal anastomosis and in 3 cases in the course of the graft. Bypasses without a stenosis at the proximal anastomosis showed a better initial peri-anastomotic match in strain (1.1 vs 1.5, $p<0.05$) and a better initial diameter match (0.68 vs 0.38, $p<0.05$)

as compared to the bypasses with a stenosis at this point during follow-up. Regression analysis showed that only an initial local high PSR was independently correlated with the occurrence of stenosis at the proximal anastomosis. At the distal anastomosis no differences in initial peri-anastomotic match in elastic properties and flow characteristics were observed between the bypasses with and without a stenosis at this site. Histologic examination of the stenoses (n=6) demonstrated only partial covering (50%) of the remaining lumen with endothelial cells, indicating denudation.

Conclusions. Our data suggest that a greater mismatch in strain and a poor diameter match contribute to the development of intimal hyperplasia. In addition, it is suggested that a high initial peak shear rate initiates the development of intimal hyperplasia in the proximal anastomosis of venous femorodistal grafts. A possible mechanism may be a damaging effect of high shear on venous endothelium, leading to partial denudation.

INTRODUCTION

The main cause for failure of peripheral bypass grafts in the first post-operative year is the formation of stenosis due to intimal hyperplasia¹. The development of a stenosis may jeopardize the function of the graft and may finally lead to occlusion². Platelet aggregation^{3,4}, endothelial cell dysfunction⁵, damage of the vessel wall media⁶ and a variety of growth factors^{3,7} have all been alleged to play a role in the development of intimal thickening, at least in animal models. In addition, mechanical factors such as compliance mismatch⁸⁻¹⁰ and unphysiological shear rates¹¹⁻¹⁵ may play a role in the formation of a stenosis.

Since the anastomoses are predisposed to the development of a stenosis, many researchers hypothesized that a mismatch in elastic properties between graft and native arteries might be an important contributing factor^{1,8,16-18}. Excessive mechanical stretch at the anastomoses due to mismatch may lead to vessel wall injury and may initiate the replication of vascular smooth muscle cells⁸ (VSMC), the dominant cell type in the intimal lesion. In animal models a correlation between compliance mismatch and intimal hyperplasia was never demonstrated^{8,19}. However, occlusive intimal hyperplasia has never been observed in animal models, which may explain the absence of a correlation between mismatch in elastic properties and intimal hyperplasia^{20,21}. Another factor affecting intimal growth is flow. A low shear rate in arteries is associated with increased intimal thickening, as has been demonstrated by numerous investigators^{11,12,22-24}. On the other hand, more intimal thickening was observed at the venous site of AV fistulas in dogs in areas with the highest flow velocities^{14,25}. The influence of shear rate on the occurrence of intimal hyperplasia in saphenous vein bypass grafts in humans could not be investigated so far, due to the absence of suitable techniques to assess shear rate in vivo accurately.

The present prospective study was conducted to investigate the influence of both a mismatch in elastic properties and shear rate on the occurrence of intimal hyperplasia in saphenous vein peripheral bypass grafts in humans. In a group of patients who

underwent peripheral bypass surgery, the elastic properties of the saphenous vein grafts and the host arteries were assessed at defined sites in a non-invasive way, with the use of Vessel Wall Doppler Tracking, an ultrasound technique. By using this method, we were able to calculate mismatch in elastic properties around the anastomoses²⁶. Furthermore, peak shear rate, calculated from diameter and peak systolic velocity was assessed at corresponding sites in the graft and adjacent vessel segments²⁷. The measurements took place within 2-4 weeks after the construction of the bypass. The initial mechanical and flow conditions of the vein grafts were correlated with the occurrence of (sub)occlusive intimal hyperplasia in the post operative period. In addition, histologic examination of stenotic lesions was performed to find clues for the etiology of the stenosis.

MATERIALS AND METHODS

Patient clinical profile

Forty-three consecutive patients receiving a saphenous vein bypass graft to reestablish a sufficient blood flow to the lower limb entered the study (august 1991-august 1993). The majority of the patients were operated for limb threatening ischaemia (Table I). In 28 patients the reversed vein technique was used and in 15 patients the in situ technique. Before inclusion in the study, a Duplex examination was performed to ascertain that the the graft was free of stenotic lesions or incompletely removed valves². A stenotic lesion may alter the hemodynamics of the graft by influencing pulse wave reflections^{28,29}. Therefore these patients had to be excluded.

Table I: Characteristics of the patients with and without a stenosis.

	Stenosis	No stenosis	P value
Sex (male/female)	9/5	16/13	0.35
Age (Years)	67(60-74)	71(64-78)	0.24
Fontaine			
II	4	8	0.93
III	5	11	
IV	5	10	
Diabetes	4	12	0.34

Detection of stenoses during the study period

At 6, and 12 weeks and every 3 months thereafter a Duplex scan was performed as part of a surveillance program in the hospital. When the Duplex scan revealed a stenosis, angiography was performed to assess the severity of the stenotic lesion. A decision to perform surgical correction of the stenotic lesion was based on the outcome of both the Duplex scan and angiography.

Measurement of the elastic properties and calculation of match

The diameter and distension of the vessel and graft segments were measured with the use of Vessel Wall Doppler Tracking (VWDT). A detailed description of this ultrasound system is given elsewhere (chapter 3 and 4)³⁰⁻³². From diameter (D in mm) and distension (Δd in μm) the following parameters were calculated:

$$\text{Relative Distension (RD)} = \frac{\Delta D}{D} * 100\% \quad (1)$$

$$\text{Area-increase} \sim \frac{\pi \Delta D * D}{2} \quad (2)$$

Relative distension (RD in %) represents the local strain of the vessel wall during the cardiac cycle. Area-increase (AI in mm^2) is calculated as the increase in cross-sectional area during cardiac cycle and represents the capacity of the vessel to store and release blood volume. With the use of B-mode imaging measurements were performed at defined sites in the graft and adjacent vessel segments by using the anastomosis as a landmark²⁶.

Since we aimed at investigating the influence of mismatch in elastic properties around anastomoses on the development of intimal hyperplastic stenoses, the values for RD, AI and diameter around the anastomoses were indexed. The calculation of the indices were performed as follows:

$$\text{Match factor} = \text{value distal} / \text{value proximal from the anastomosis} \quad (3)$$

A perfect match is considered to be present when the match factor is 1. A value lower than 1 indicates a loss of the value of parameter in the anastomosis, while a value higher than 1 indicates an increase of the value of the parameter within the anastomosis.

Calculation of peak shear rate

A reliable method to perform in vivo shear rate measurements in humans is not available at this moment. Therefore, shear rate can only be estimated from flow parameters and diameter. Usually, mean shear rate is calculated from the Haagen-Poiseuille formula^{11,12}. However, studies evaluating the effect of steady and pulsatile flow on the lining and the shape of endothelial cells demonstrated that the pulsatile component of shear rate may be of more importance than the mean level³³⁻³⁵. Furthermore, in another study evaluating the effect of flow on proliferation in human stenotic tissue, we observed a negative correlation between peak shear rate and proliferation, which also indicates that the pulsatile component of shear rate may be an important determining factor (chapter 11)²⁷. We therefore decided to calculate peak shear rate from a modification of the Haagen-Poiseuille formula in the following way:

$$\text{Peak shear rate (1/s, PSR)} = n * \text{peak systolic velocity/diameter} \quad (4)$$

Peak systolic velocity was determined from the doppler sample placed midstream under an angle of 60°. Diameter was measured with the use of VWDT. The calculation of PSR with the modification of the Haagen-Poiseuille formula is based on the assumption that flow is laminar and the velocity profile is parabolic. Since one can only speculate on the presence of such conditions in the bypass grafts, PSR can only give a rough approximation of the actual shear stresses exerted on the vessel wall. The value for n varies between 4 and 8 for normal arterial flow and increases when flow velocity increases. The determination of diameter with the use of the B-mode image is not very accurate. Therefore, we used the diameter values obtained from VWDT. A validation study showed a low coefficient of variation (2-4%) in the assessment of diameter with this system³².

Histologic examinations

When a stenotic lesion of hemodynamic importance occurred within the graft necessitating surgical repair with the use of interposition grafts, a stenotic tissue sample was obtained for histologic examination. These specimens were processed according to routine histologic procedures and paraffin-embedded. Serial sections were cut and placed on gelatin coated glass slides. To identify endothelial cells, Ulex Europeus agglutinin (1:500 dilution, Vector Labs, Burlingame, CA, USA) was used. After blocking of the endogenous peroxidase activity with 0.3% H₂O₂ and predigestion with pepsin (1mg/ml, 30 min), sections were incubated with the lectin antibody (1:250, 45 min, DAKO). Subsequently, the sections were incubated with an anti-Ulex Europeus lectine peroxidase complex (1:100, DAKO). As the chromogen 3,3'-diamino-benzidine was used. The degree of endothelial cell coverage was estimated by dividing the surface of the lumen with endothelial cells by the total luminal surface.

ANALYSIS

Data are presented as medians and 25th and 75th percentile intervals. To compare results between different groups, the Mann-Whitney-U test was used. Multiple regression analysis was used to determine the relation between the local initial mechanical and flow characteristics and the occurrence of a stenosis at a certain site. A p-value of less than 0.05 was accepted as a level of statistical significance.

RESULTS

During the follow-up period stenotic lesions developed in 14 grafts (Table II). In 7 cases the lesion developed in or adjacent to the proximal anastomoses, in 4 cases within the distal anastomosis and in 3 cases in the course of the graft. Since measurements of the elastic properties and flow focused on locations around the anastomoses, the development of focal stenoses in or directly adjacent to the proximal and distal anastomoses could be correlated with the initial mechanical and flow characteristics. In the remaining 3 cases, the stenoses were randomly distributed along the entire grafts. Reliable correlation of these stenoses with mechanical and flow conditions with respect to localisation is therefore difficult. Since coupling of local flow and mechanical parameters is essential in this study, these stenoses were not included in the study. The majority of the stenoses in the proximal anastomosis occurred in the reversed vein bypass group (84%, Table II). No differences in characteristics were observed between the stenotic and non-stenotic bypass patients (Table I).

Table II: Distribution of stenosis in the reversed and in situ saphenous vein grafts

	Reversed (n=28)	In Situ (n=15)	Total
Proximal Anastomosis	6	1	7
Graft	1	2	3
Distal Anastomosis	2	2	4

Local elastic properties and stenosis

A summary of the results is presented in Table III and IV. No differences in initial RD and AI values measured at the proximal and distal graft segments were observed between the bypasses with and without a stenosis at these sites. A smaller initial diameter of the proximal graft segments was observed in the bypasses with a later stenosis at this site as compared to the non-stenotic bypasses (3.2 vs 4.4 mm, $p < 0.05$).

Table III: The outcome of different mechanical parameters in the patients with and without a stenosis in or adjacent to the proximal anastomosis of the venous bypass. Data are presented as medians and 25 and 75 th percentile intervals.

	Proximal stenosis (n=7)	No stenosis proximal (n=36)	P Value
Vein diameter(mm)	3.2(2.8-4.3)	4.4(3.7-5.6)	0.03
Diameter match	0.38(0.26-0.58)	0.65(0.43-0.84)	0.03
RD(%)	3.5(2.5-4.5)	3.2(1.5-4.1)	0.35
RD match	1.5(1.2-2.4)	1.1(0.6-1.4)	0.03
AI (mm ²)	0.68 (0.52-1.12)	0.92 (0.54-1.38)	0.21
AI match	0.31 (0.08-0.50)	0.36 (0.16-0.79)	0.45
PSV (cm/s)	127(70-211)	105(77-131)	0.39
PSR (1/s)	143(55-338)	99(64-145)	0.32

Table IV: The outcome of different mechanical parameters in the patients with and without a stenosis in or adjacent to the distal anastomosis of the venous bypass. Data are presented as medians and 25 and 75 th percentile intervals.

	Distal stenosis (n=4)	No stenosis distal (n=39)	P Value
Vein diameter(mm)	3.8 (2.4-5.1)	4.6 (3.6-6.4)	0.18
Diameter match	0.6 (0.5-0.9)	0.7 (0.6-0.90)	0.46
RD(%)	2.3 (1.4-4.7)	2.0 (1.2-3.6)	0.67
RD match	0.9 (0.8-1.2)	0.8 (0.4-1.2)	0.32
AI (mm ²)	0.68 (0.22-1.03)	0.68 (0.48-1.14)	0.59
AI match	0.4 (0.2-0.7)	0.3 (0.2-0.5)	0.66
PSV (cm/s)	100 (45-136)	67 (53-83)	0.31
PSR (1/s)	124 (35-224)	56 (39-94)	0.61

Peri-anastomotic match in elastic properties and stenoses

Proximal anastomosis

No difference in initial peri-anastomotic match in AI was observed between the stenotic and non-stenotic bypasses (Table III). A better initial match in RD (1.1 vs 1.5, $p<0.05$, Figure 1) and diameter (0.65 vs 0.38, $p<0.05$, Figure 1) around the proximal anastomosis was observed in the bypasses without a later stenosis at this site as compared to the non-stenotic bypasses. Between the stenotic and non-stenotic bypasses no differences were observed in initial local peak systolic velocity (PSV). Also in initial local peak shear rate (PSR) no differences were observed between the groups (Table III). In the bypass group with a proximal stenosis, higher PSR and PSV values were measured at the site of the later stenoses as compared to the distal segments of the bypass (PSR: 145 vs 62 1/s, $P<0.05$; PSV: 127 vs 80 cm/s, $p<0.05$, Wilcoxon signed rank test). Multiple regression analysis, including mismatch and flow parameters, demonstrated that only a high initial local PSR at the proximal anastomosis independently correlated with the occurrence of a stenosis at this point ($p<0.05$).

Figure 1: Bar graph showing the differences in match in strain and diameter around the proximal anastomosis between the bypasses with and without a stenosis at this point.

Distal anastomosis

No differences in initial peri-anastomotic match in AI, RD and diameter were observed between the bypasses with and without a later stenosis at the distal anastomosis. Also in PSR and PSV, no differences were observed between the bypasses with and without a stenosis at the distal graft end (Table IV).

Histologic examinations

In 6 cases repair of the proximal stenotic lesions was performed with the use of an interposition graft, allowing harvesting of stenotic material. Histologic evaluation revealed a focal stenotic mass consisting mainly of vascular smooth muscle cells and matrix, confirming the hyperplastic nature of the material. The mean coverage of endothelial cells of the lumen was 50%.

DISCUSSION

The goal of the present study was to correlate initial mechanical factors, like perianastomotic mismatch in elastic properties and flow characteristics, with the occurrence of stenosis during follow-up in peripheral saphenous vein bypass grafts. For the proximal anastomosis, a better initial match in local strain and diameter was observed in the bypasses without a stenosis at this site during follow-up as compared to the stenotic bypasses. This suggests that mismatch in strain and diameter may play a role in initiating the development of an anastomotic stenosis. Furthermore, multiple regression analysis showed that a high initial local peak shear rate at the proximal graft segment was associated with the occurrence of a stenosis at this site during follow-up. This suggests that high shear rate initiates intimal hyperplasia in venous grafts.

Mismatch in elastic properties and stenosis

The compliance mismatch hypothesis stems from numerous observations reporting a high incidence of stenosis within or adjacent to the anastomoses in peripheral bypasses and in arteriovenous fistulas^{1,8,14,16,17,25,36-39}. Theoretically, a point of excessive mechanical stretching is created when vessels with different elastic properties are anastomosed⁸, which may lead to vessel wall injury. A greater initial mismatch in strain was found in peripheral bypasses with a stenosis in the proximal anastomosis during follow-up as compared to the bypasses without a stenosis at this point. The local initial strain values were equal in the proximal graft segments in both groups, suggesting that mismatch in strain had to be the contributing factor rather than local mechanical conditions. A point of interest is that strain increased within the proximal anastomoses of the stenotic bypasses. The median value for the mismatch was 1.5, indicating a 50% increase in relative strain. In vitro studies have demonstrated that cyclic stretching of vessel segments results in proliferation of smooth muscle cells and production of extra-cellular matrix, important events in the development of intimal thickening⁴⁰⁻⁴². In addition increased proliferation of VSMC was observed after cyclic stretching, in venous segments as compared to arterial segments, indicating that veins may be more susceptible to repetitive strain than arteries⁴⁰. Thus, the mismatch in relative strain between the inflow artery and the proximal vein segments, resulting in excessive repetitive cyclic strain at the site of the proximal anastomosis, may contribute to the development of intimal thickening in human saphenous vein bypass grafts.

Another contributing factor may be the influence of a mismatch in compliance on flow⁴³. A sudden increase in compliance has been associated with flow disturbances like flow stasis and increased particle residence time, suggesting low shear. In the present study, the majority of stenoses developed at the proximal anastomoses. At this point, a sudden decrease in compliance is present, which is not associated with flow disturbances and probably will not lead to low shear conditions. At the distal anastomosis, no differences were observed in peri-anastomotic mismatch in elastic properties between the bypasses with and without a stenosis at this point. A possible explanation may be that the number of stenosis at this point is too small for statistical analysis.

Smaller vessel diameters were found in the vein grafts with a stenosis at the proximal anastomosis as compared to the non-stenotic bypasses. Diameter mismatch was also greater in the stenotic bypasses. We do not know whether the occurrence of stenotic lesions in segments with a small diameter during the follow-up period is caused by the fact that intimal thickening in these bypasses leads more rapidly to subocclusive or occlusive stenotic lesions. Recurrence of symptoms will return faster and Duplex surveillance will demonstrate higher velocities in narrow bypasses with a certain stenotic mass as compared to large diameter grafts. As a consequence, angiography and surgical intervention may be performed in relatively early stages in the development of the intimal hyperplastic stenoses. The finding that bypasses with a proximal stenosis had smaller initial diameters indicates that in case of a small caliber saphenous vein, the in situ technique may be preferable over the reversed technique.

Flow characteristics and stenosis

In the bypasses with a stenosis at the proximal anastomosis, higher initial peak shear rate values were found at the proximal graft segments as compared to other regions of the bypass. In addition, multiple regression analysis showed that a high initial local peak shear rate value correlated with the occurrence of a stenosis at the proximal anastomosis during follow-up. This suggests that a high initial peak shear rate initiates stenosis due to intimal thickening in saphenous vein bypass grafts. The majority of the stenosis at the proximal anastomosis occurred in the reversed saphenous vein bypass group, which confirms observations from other groups^{1,16}. The highest initial flow velocities and peak shear rates are present in the most proximal part of the graft due to the small diameter at this point (equation 3). Histologic examinations of the stenotic tissue samples showed that the lumen of the stenosis was only partly covered with endothelial cells. This indicates that denudation of endothelial cells was present at the time of harvesting the stenotic sites. Since PSV and PSR values at the site of the stenosis just prior to intervention will even be further increased as compared to the initial high values, the reason for denudation may be excessive high shear rate^{44,45}. This has been previously observed in a similarly conducted study by our group on AV fistulas⁴⁶. The occurrence of stenosis at the venous anastomoses of hemodialysis graft AV fistulas was associated with high initial PSR and PSV values, and also denudation was found in stenotic areas.

Since denudation is an established method to initiate intimal thickening in animal vessels^{47,48}, one might speculate that at least the initiation of intimal thickening is due to denudation mediated by high shear rate. This seems to be in contrast with observations from many animal studies, reporting increased intimal thickening in areas of low shear^{11,12}. However, some differences between experimental animal models and human saphenous vein bypass grafts may explain the discrepancy. First, in most animal models the influence of varying levels of shear on intimal thickening is studied in arteries, whereas in the present study the influence of shear on intimal thickening in veins was studied. Venous endothelium may be more susceptible to high shear as compared to arterial endothelium as has been suggested by some authors^{14,25}. A second reason may be that the biology of human intimal thickening differs from intimal thickening in animals^{20,21,49}. In animal models intimal thickening is limited and never leads to occlusion, which is quite different from occlusive intimal thickening as found after vascular intervention in humans. The observation that the occurrence of stenosis at the proximal anastomosis is associated with initial high peak shear rate may explain why small sized veins have a lower graft patency^{50,51}. Due to a small diameter, peak systolic velocity will be high, resulting in a high peak shear rate level. Therefore, a certain critical vein diameter may result in peak shear rate levels, which are not well tolerated by venous endothelial cells. This may result in endothelial cells damage and eventually denudation, which may initiate intimal thickening.

In conclusion, our data suggest that a high initial mismatch in strain and a poor diameter match may contribute to the development of proximal anastomotic intimal hyperplasia in saphenous vein peripheral bypass grafts in humans. In addition, the development of intimal hyperplasia at the proximal anastomosis of these grafts was associated with increased flow velocities and high peak shear rates, suggesting that high shear contributes to the development of intimal hyperplasia in vein grafts. A possible mechanism may be a damaging effect of high shear on venous endothelium, leading to partial denudation, thereby initiating the development of a stenosis.

REFERENCES

1. Berkowitz HD, Greenstein S, Barker CF, Perloff LJ. Late failure of reversed vein bypass grafts. *Ann Surg* 1989; 210:782-786.
2. Buth J, Disselhoff B, Sommeling C, Stam L. Color-flow duplex criteria for grading stenosis in infrainguinal vein grafts. *J Vasc Surg* 1991; 14:716-728.
3. Liu MW, Roubin GS, King III SB. Restenosis after coronary angioplasty. Potential biologic determinants and role of intimal hyperplasia. *Circulation* 1989; 79:1374-1387.
4. Fingerle J, Johnson R, Clowes AW, Majesky MW, Reidy MA. Role of platelets in smooth muscle cell proliferation and migration after vascular injury in rat carotid artery. *Proc Natl Acad Sci U S A* 1989; 86:8412-8416.
5. Lüscher TF. Endothelial control of vascular tone and growth. *Clin Exp Hypertens* 1990; 12:897-902.
6. Clowes AW, Clowes MM, Fingerle J, Reidy MA. Kinetics of cellular proliferation after arterial injury. V. Role of acute distension in the induction of smooth muscle cell proliferation. *Lab Invest* 1989; 60:360-364.
7. Clowes AW, Reidy MA. Prevention of stenosis after vascular reconstruction: Pharmacologic control of intimal hyperplasia - A review. *J Vasc Surg* 1991; 6:885-891.
8. Abbott WM, Megerman J, Hasson JE, L'Italien G, Warnock DF. Effect of compliance mismatch on vascular graft patency. *J Vasc Surg* 1987; 5:376-382.
9. Abbott WM, Megerman J. Adaptive responses of arteries to grafting. *J Vasc Surg* 1989; 9:377-379.
10. Kidson IG, Abbott WM. Low compliance and arterial graft occlusion. *Circulation* 1978; 58:11-14.
11. Kohler TR, Jawien A. Flow affects development of intimal hyperplasia after arterial injury in rats. *Arterioscler Thromb* 1992; 12:963-971.
12. Kohler TR, Kirkman TR, Kraiss LW, Zierler BK, Clowes AW. Increased blood flow inhibits neointimal hyperplasia in endothelialized vascular grafts. *Circ Res* 1991; 69:1557-1565.
13. Fry DL. Certain chemorheologic considerations regarding the blood vascular interface with particular reference to coronary artery disease. *Circulation* 1969; XXXIX and XI:IV38-IV59.
14. Fillinger MF, Reinitz ER, Schwartz RA, et al. Graft geometry and venous intimal-medial hyperplasia in arteriovenous loop grafts. *J Vasc Surg* 1990; 11:556-566.
15. Sottiurai VS, Kollross P, Glagov S, Zarins CK, Mathews MB. Morphologic alteration of cultured arterial smooth muscle cells by cyclic stretching. *J Surg Res* 1983; 35:490-497.
16. Berkowitz HD, Fox AD, Deaton DH. Reversed vein graft stenosis: Early diagnosis and management. *J Vasc Surg* 1992; 15:130-142.
17. Echave V, Koornick AR, Haimov M, Jacobson JH. Intimal hyperplasia as a complication of the use of the polytetrafluoroethylene graft for femoral-popliteal bypass. *Surgery* 1979; 86:791-798.
18. Madras PN, Ward CA, Johnson WR, Singh PI. Anastomic hyperplasia. *Surgery* 1981; 90:922-923.
19. Okuhn SP, Connelly DP, Calakos N, Ferrell L, Man-Xiang P, Goldstone J. Does compliance mismatch alone cause neointimal hyperplasia? *J Vasc Surg* 1989; 9:35-45.
20. Ferrel M, Fuster V, Gold HK, Chesebro JH. Choosing appropriate experimental animal model for the prevention of restenosis. *Circulation* 1992; 85:1630-1631.
21. McBride W, Lange RA, Hillis LD. Restenosis after successful coronary angioplasty. Pathophysiology and prevention. *New Eng J Med* 1988; 318:1734-1737.
22. Zarins CK, Giddens DP, Bharadvaj BK, Sottiurai VS, Mabon RF, Glagov S. Carotid bifurcation atherosclerosis. Quantitative correlation of plaque localization with flow velocity profiles and wall shear stress. *Circ Res* 1983; 53:502-514.

23. Zarins CK, Bomberger RA, Glagov S. Local effects of stenoses: Increased flow velocity inhibits atherogenesis. *Circulation* 1981; 64:11221-11227.
24. Zarins CK, Zatina MA, Giddens DP, Ku DN, Glagov S. Shear stress regulation of artery lumen diameter in experimental atherogenesis. *J Vasc Surg* 1987; 5:413-420.
25. Fillinger MF, Reinitz ER, Schwartz RA, Resetarits DE, Paskanik AM, Bredenberg CE. Beneficial effects of banding on venous intimal-medial hyperplasia in arteriovenous loop grafts. *Am J Surg* 1989; 158:87-94.
26. Hofstra L, Bergmans DCJJ, Hoeks APG, Tordoir JHM, Kitslaar PJEHM. Assessment of inhomogeneities in elastic properties of in situ and reversed saphenous vein grafts in humans. *Eur J Vasc Surg* 1994; 8:100-105.
27. Hofstra L, Tordoir JHM, Kitslaar PJEHM, Hoeks APG, Daemen MJAP. High shear rate mediated denudation decreases proliferation in human stenotic lesions derived from arterio-venous fistulas and peripheral bypass grafts. *Circulation* 1994; 100:100-105.
28. O'Rourke MF. Arterial stiffness, systolic blood pressure, and logical treatment of arterial hypertension. *Hypertension* 1990; 15:339-347.
29. Kelly RP, Gibbs HH, O'Rourke MF, et al. Nitroglycerin has more favourable effects on left ventricular afterload than apparent from measurement of pressure in a peripheral artery. *Eur Heart J* 1990; 11:138-144.
30. Hoeks APG, Brands PJ, Smeets FAM, Reneman RS. Assessment of distensibility of superficial arteries. *Ultrasound Med Biol* 1990; 16:121-128.
31. Hofstra L, Ermers EJM, Langeveld APM, Hoeks APG, Kitslaar PJEHM. Vessel Wall Doppler Tracking: A new technique to assess vessel wall compliance. *J Vasc Tech* 1993; 17:65-68.
32. Hofstra L, Willigers JM, Huvers FC, et al. Short-term variations in elastic properties of a muscular artery in humans. *Clin Sci* 1993; 50:472-480.
33. Helmlinger G, Geiger RV, Schreck S, Nerem RM. Effects of pulsatile flow on cultured vascular endothelial cell morphology. *J Biomech Eng* 1991; 113:123-131.
34. Ojiha M. Wall shear stress temporal gradient and anastomotic intimal hyperplasia. *Circ Res* 1994; 74:1227-1231.
35. DePaola N, Gimbrone MA, Davies PF, Dewey CF. Vascular endothelium responds to fluid shear stress gradients. *Arterioscler Thromb* 1992; 12:1254-1257.
36. Jenkins AMcL, Buist TAS, Glover SD. Medium-term follow-up of forty autogenous vein and forty polytetrafluoroethylene (Gore-Tex) grafts for vascular access. *Surgery* 1980; 88:667-672.
37. Sottiurai VS, Sue SL, Feinberg II EL, Bringaze WL, Tran AT, Batson RC. Distal anastomotic intimal hyperplasia: Biogenesis and etiology. *Eur J Vasc Surg* 1988; 2:245-256.
38. Swedberg SH, Brown BG, Sigley R, Wight TN, Gordon D, Nicholls SC. Intimal fibromuscular hyperplasia at the venous anastomosis of PTFE grafts in hemodialysis patients. *Circulation* 1989; 80:1726-1736.
39. Tordoir JHM, Herman JM, Kwan TS, Diderich PM. Long-term follow-up of the polytetrafluoroethylene (PTFE) prosthesis as an arteriovenous fistula for hemodialysis. *Eur J Vasc Surg* 1988; 2:3-7.
40. Predel HG, Yang Z, Von Segesser L, Turina M, Buhler FR, Luscher TF. Implications of pulsatile stretch on growth of saphenous vein and mammary artery smooth muscle. *Lancet* 1992; 340:878-879.
41. Leung DYM, Glagov S, Mathews MB. Elastin and collagen accumulation in rabbit ascending aorta and pulmonary trunk during postnatal growth. Correlation of cellular synthetic response with medial tension. *Circ Res* 1977; 41:316-323.
42. Leung DYM, Glagov S, Mathews MB. Cyclic stretching stimulates synthesis of matrix components by arterial smooth muscle cells in vitro. *Science* 1976; 191:475-477.
43. Stewart SFC, Lyman DJ. Effects of a vascular graft/natural artery compliance mismatch on pulsatile flow. *J Biomech* 1992; 25:297-310.

44. Reidy MA. Biology of disease. A reassessment of endothelial injury and arterial lesion formation. *Lab Invest* 1985; 53:513-520.
45. Fry DL. Acute vascular endothelial changes are associated with increased blood velocity gradients. *Circ Res* 1968; 22:165-197.
46. Hofstra L, Bergmans DCJJ, Leunissen KML, Hoeks APG, Kitslaar PJEHM, Tordoir JHM. Anastomotic intimal hyperplasia in prosthetic arterio-venous hemodialysis access fistulas is associated with high shear rate and not with mismatch in elastic properties. *JASN* 1994; 100:100-105.
47. Buga GM, Gold ME, Fukuto JM, Ignarro LJ. Shear stress-induced release of nitric oxide from endothelial cells grown on beads. *Hypertension* 1991; 17:187-193.
48. Daemen MJAP, Lombardi DM, Bosman FT, Schwartz SM. Angiotensin II induces smooth muscle cell proliferation in the normal and injured rat arterial wall. *Circ Res* 1991; 68:450-456.
49. O'Brien ER, Alpers CE, Stewart DK, et al. Proliferation in primary restenotic coronary atherectomy tissue. Implications for antiproliferative therapy. *Circ Res* 1993; 73:223-231.
50. Wengerter KR, Veith KJ, Gupta SK, Ascer E, Rivers SP. Influence of vein size (diameter) on infrapopliteal reversed vein graft patency. *J Vasc Surg* 1990; 11:525-531.
51. Wengerter KR, Yang PM, Veith FJ, Gupta SK, Panetta TF. A twelve-year experience with the popliteal-to-distal artery bypass: The significance and management of proximal disease. *J Vasc Surg* 1992; 15:143-151.

CHAPTER 11

HIGH SHEAR RATE MEDIATED DENUDATION DECREASES PROLIFERATION IN HUMAN STENOTIC LESIONS DERIVED FROM ARTERIO-VEINOUS FISTULAS AND PERIPHERAL BYPASS GRAFTS

L Hofstra, JHM Tordoir, PJEHM Kitslaar, APG Hoeks, MJAP Daemen.
(submitted)

ABSTRACT

Background. Vascular interventions are often complicated by the development of intimal thickening, leading to stenosis. Proliferation is considered to be one of the key events in the development of an intima. There has been extensive research on the growth regulation of the intima in animals, but information on the biology of intimal thickening in humans is scarce.

Methods. We studied the proliferation patterns of 35 human stenotic lesions, derived from either peripheral bypasses (n=18) or hemodialysis AV-fistulas (n=17) with the use of KI-67, an antibody to a proliferating cell nuclear antigen. In addition, proliferation was assessed in 13 saphenous vein control specimens. Furthermore, peak shear rate was calculated from peak systolic velocity and diameter, both assessed non-invasively with the use of ultrasound, in an attempt to study the influence of local flow characteristics on the proliferation patterns in the stenotic lesions.

Results. Proliferation rates were less than 0.2% in control vessels and were elevated in stenotic tissue. In both the AV-fistula and the bypass stenoses, proliferation was higher in the area just below the endothelium as compared to intimal regions adjacent to the media (AV-fistulas: 3.6 vs 1.1% and bypasses: 3.5 vs 2.0%, $p<0.01$). Multiple regression analysis showed that the extent of subendothelial-intimal proliferation was independently correlated with the endothelial cell coverage of the lumen ($r=+0.74$, $p<0.01$). For local peak shear rate, the age of the lesion and graft material (PTFE or vein) no independent correlations with subendothelial proliferation were found. Local peak shear rate was negatively correlated with endothelial cell coverage of the stenosis ($r=-0.61$, $p<0.05$). Proliferation in the different vessel layers showed no differences between the bypass and the AV-fistula group. From the intima to the adventitia a gradually increasing percentage of proliferating cells could be identified as endothelial cells, indicating neovascularisation.

Conclusions. These data indicate that cell proliferation in human stenotic tissue, derived from AV fistulas and peripheral bypass grafts, is increased as compared to control vessels and is concentrated in subendothelial-intimal areas. Furthermore, the extent of subendothelial-intimal proliferation is correlated with the presence of endothelial cell coverage of the lumen. In addition, a high shear rate was correlated with an decreasing endothelial cell coverage of the lumen, suggesting that denudation of EC may be the mechanism by which high shear rate results in decreasing DNA synthesis in the underlying VSMC.

INTRODUCTION

Failures of vascular reconstructions are mainly caused by the development of stenotic lesions, often located at the anastomoses¹⁻⁵. The incidence of stenoses in the first post-operative year varies from 20-30 % in femorodistal bypass grafts to 50-60% in hemodialysis arterio-venous fistulas when constructed with the use of interposition grafts⁶⁻⁸. Proliferation of vascular smooth muscle cells (VSMC) is thought to be the dominant cellular event in the development of an intima^{5,9,10}. The kinetics of this proliferative response are well described in animal studies¹¹. The biology of human intimal hyperplastic lesions may be different and possibly more complex^{9,10,12,13}. However, information on proliferation in human stenotic lesions is scarce.

In recent studies, the proliferative response of VSMC in humans was investigated in restenotic coronary tissue¹³, peripheral artery restenotic lesions¹⁴ and stenoses derived from AV fistulas¹⁰. In comparable stenotic lesions very high (up to 20%) and very low (0-0.5%) proliferation indices were reported. Consequently, the exact contribution of VSMC proliferation to restenosis in humans still remains unclear. A difficulty of the reported studies may be the use of PCNA as the marker for proliferation. Overestimation of the proliferation index with this marker has been demonstrated, and may explain the reported high proliferation indices¹⁵. A difficulty of evaluating atherectomy derived stenotic tissue may be the possible influence of sampling error, giving underestimation of the contribution of proliferation¹³. The use of another proliferation marker, KI-67 may provide an easier and more reliable method to determine proliferative activity in tissue samples¹⁵⁻¹⁸.

One of the factors that influence the growth of an intimal lesion is local flow. A decrease in flow is associated with increased intimal thickening¹⁹⁻²⁵. The mechanism by which flow affects intimal hyperplasia may be its influence on the release of PDGF and EDRF, which has been demonstrated in *in vitro* studies²⁶⁻²⁸. In humans, accelerated atherosclerosis has been reported in bypassed coronary vessels, which have almost zero flow²⁹. However, information on the possible correlation between flow and intimal growth in humans lesions is lacking. Another factor influencing proliferation is the presence of endothelial cells. In animal models, denudation of arteries initiates proliferation of VSMC in the media and subsequently in the intima.^{30,31} On the other hand, in *in vitro* studies elevated proliferation was observed in non-denuded areas with

high levels of EC DNA synthesis, suggesting a stimulatory effect of activated endothelial cells on VSMC proliferation³²⁻³⁵. The role of endothelial cells in proliferation in human stenotic lesions derived from patients with AV fistulas and peripheral bypasses has never been established.

The aim of the present study was to determine the distribution and time course of proliferation in human stenotic lesions. Furthermore, the influence of local flow characteristics on the proliferative response in the stenoses was studied. Since arteriovenous fistulas are an example of a high flow system, and a peripheral bypass an example of a low flow system, we hypothesized that the extent and pattern of replication in stenotic lesions derived from these two types of vascular reconstructions might be different, i.e. a high proliferation in the peripheral bypass and low proliferation in the AV fistula. In a consecutive series of patients receiving an hemodialysis arteriovenous fistula or a femorodistal bypass, flow characteristics were determined 2 weeks after construction at numerous sites in the graft and in the in- and outflow vessel segments, with the use of ultrasound techniques. At a later time point, the local flow characteristics were assessed at the site of the stenosis in patients developing stenoses in the graft or at the anastomoses, necessitating surgical intervention. After harvesting of the stenotic vessel segments, proliferation was assessed in different layers of the stenosis, with the use of KI-67. Local flow characteristics were correlated with the extent of proliferation in the stenotic vessel segments. In addition, endothelial cell coverage of the stenotic lumen was correlated with the extent of proliferation.

MATERIALS AND METHODS

Tissue Specimens

During the study period, from January 1992 to June 1994, 48 reinterventions were performed following construction of either an arteriovenous fistula or a peripheral bypass in the surgery department of the University Hospital of Maastricht or the Catharina Hospital of Eindhoven. In 13 cases the vascular conduit was corrected by patch plasty at the site of the stenosis. Since collection of stenotic tissue in these cases was impossible without performing extra surgical procedures, which might jeopardize the success of the reintervention, it was decided not to obtain tissue specimens from this group of patients. In the remaining 35 cases surgical repair was performed by placement of an interposition graft, allowing the collection of the stenotic process and adjacent vessel segments. In all cases the stenotic lesion was focal and apparently narrowing or occluding the lumen.

PATIENT CLINICAL PROFILE

Patients with an arterio-venous fistula

This group consisted of 9 men and 6 women. Mean age was 56 years, ranging from 37 to 76 years. Repair of the AV fistula was performed after an average time of 10.4 months, varying from 3 to 27 months. The diagnosis of a stenosis was based on Duplex ultrasound criteria and confirmed by fistulography⁸. In all patients the occurrence of a stenosis was associated with an increase in venous pressure and/or low flow of the fistula during hemodialysis, necessitating reintervention. In one patient a second reintervention was performed during the study period and in one patient two stenotic lesions were derived from different sites in the vascular conduit, resulting in a total number of 17 stenotic samples ($n=17$). In 12 cases, the stenotic samples were derived from the efferent vein of either Brescia-Cimmino fistulas ($n=2$) or fistulas constructed with the use of interposition grafts ($n=10$). In 5 cases the stenotic sample was derived from the venous anastomosis of a PTFE interposition graft.

Patients with a peripheral bypass

This group of patients with a femorodistal bypass graft, consisted of 12 men and 4 women. Mean age was 64 years, ranging from 54 to 77 years. Reintervention was performed after an average time of 4.5 months, varying from 1 to 24 months. The diagnosis of a stenosis was based on generally accepted Duplex criteria and confirmed by angiography³⁶. Patients with a graft at risk underwent repair of the stenotic segment. In most of these patients the occurrence of a stenosis was associated with a recurrence of symptoms. In two patients stenotic tissue was derived from two different sites in the graft, making the total number of stenotic vessel specimens in this group 18 ($n=18$). In 10 cases the stenotic tissue was derived from reversed saphenous vein grafts, in 6 cases from in situ saphenous vein grafts and in 2 cases from a PTFE graft. Stenotic samples were derived from the proximal anastomosis ($n=11$), midgraft ($n=5$), and from the distal anastomosis ($n=2$).

Controls

From patients receiving an AV fistula ($n=6$) or a peripheral bypass graft ($n=9$) with the use of saphenous vein, spare vein segments were derived and used for the assessment of proliferation.

Tissue Staining with KI-67

Cell proliferation was determined using KI-67, an antibody to an antigen expressed in all phases of the cell cycle, except G0 and the early part of G1. The KI-67 antigen is localized at the periphery of the chromosome scaffold and cellular cortex¹⁵. The exact function of this antigen in the process of replication has not been determined so far, but a reaction of the KI-67 antibody with nuclear proteins has been established¹⁵. Furthermore, the proliferation indices determined with KI-67 and 3H-thymidine are closely correlated^{16,18}.

Staining Protocol

Immediately after harvesting, the tissue specimens were carefully processed to preserve the sections intact and to prevent loss of endothelial cells covering the lumen. The tissues were fixed overnight in 10% phosphate buffered formalin, processed according to routine histologic procedures and paraffin-embedded, 5- μ m-thick serial sections were cut and placed on gelatin coated glass slides. After deparaffinizing the sections and blocking of endogenous peroxidase activity with 0.3% H₂O₂, the slides were placed in a 0.01 M citrate buffer, pH 6.0 and boiled during 5 minutes in a microwave oven. After subsequent washes in H₂O and tris buffered saline (TBS), the slides were incubated with a mouse anti KI-67 monoclonal antibody (MIB, Immunotech, Apeldoorn, The Netherlands) during 60 minutes at room temperature at a dilution of 1:100, followed by incubation with biotinylated sheep anti-mouse IgG (1:250; DAKO) at room temperature for 30 minutes. Sections were then labeled for 30 minutes with an Avidin/Biotin peroxidase complex (Vectastain - ABC kit, Vector Labs, Burlingame, CA, USA) using 3,3' diaminobenzidine as the chromogen. The sections were counterstained with hematoxylin, dehydrated through graded series of alcohols and mounted. A piece of human small intestine was used as a positive control in all staining sessions. As an internal positive control we used one stenotic AV fistula segment, which showed consistent positive staining during all sessions. As a negative control the staining procedure was performed without addition of the primary antibody. Furthermore, a spare saphenous vein segment was used as a negative control, showing no proliferation throughout the different staining sessions.

To identify endothelial cells, Ulex Europeus agglutinin I (1:500 dilution, Vector Labs, Burlingame, CA, USA) was used. After blocking of endogenous peroxidase activity with 0.3%, H₂O₂ sections were incubated with the lectin antibody (1:250, 45 min., DAKO) after predigestion with pepsin (1mg/ml, 30 min). Subsequently, the sections were incubated for 45 minutes with an anti-Ulex Europeus lectine peroxidase complex (1:100, DAKO). As the chromogen, 3,3'diamino-benzidine was used.

Cell Counting

Tissue sections were investigated by light microscopy at x400 magnification using a standard field size. Cells containing dark nuclear staining with KI-67 were considered as positive cells^{16,17}. In order to obtain insight in the distribution of the proliferative response in the intima, cells were counted separately in subendothelial-intimal and medial-intimal areas. Furthermore, medial and adventitial areas were counted (Figure 1). All cell counting was performed with an ocular grid by two investigators (LH and MD). In each of the vessel layers, 4 at random selected fields were counted. The proliferation index, defined as the total number of positive cells divided by the total cell number, was calculated for each layer. In the stenotic lesions derived from PTFE grafts, no medial proliferation index was calculated. To assess the contribution of angiogenesis in the proliferative response a separate index for positive endothelial cells was calculated. Since cell counting was performed with the use of standard field size, the total number of cells per field can be used as a measure for the cell content in the different vessel layers. The degree of endothelial cell coverage of the lumen was

estimated by using the Ulex Europeus stained sections and was calculated by dividing the surface of the lumen covered by endothelial cells by the total luminal surface. Tissue processing and all cell counting was performed with the investigators blinded to the clinical information.

Measurement of the flow characteristics

Since we attempted to determine the influence of the initial as well as the actual local flow characteristics on proliferation, peak systolic velocity (PSV) and peak shear rate (PSR) were assessed in vivo at numerous sites in the graft and in the in- and outflow segments with special attention to sites with a predisposition to develop intimal lesions, like the anastomoses. PSV was assessed with the use of Duplex ultrasound (ATL, Ultramark IV, Advanced Technology Laboratories, USA) with a doppler sample placed midstream at defined sites in the vascular conduit. The diameter was determined at the same defined sites in the graft and in- and outflow arterial segments with the use of Vessel Wall Doppler Tracking (VWDT). This system has been described in detail elsewhere (chapter 3 and 4)^{37,38}. Validation studies showed a coefficient of variation of 2-4% in the determination of diameter of a medium-sized muscular artery with the use of this system (chapter 4)³⁹.

From PSV and diameter, peak shear rate (PSR) was calculated, based on the Haagen-Poiseuille formula for an assumed symmetric parabolic velocity profile²⁵:

$$\text{PSR} = n * \text{PSV}/\text{diameter} \quad (1/\text{s}) \quad (1)$$

The local flow characteristics, determined 2 weeks after construction of the vascular conduit and 1-2 weeks before surgical repair of the stenosis, were used for analysis and correlation with proliferative patterns.

ANALYSIS

Data are presented as means and standard deviations. To compare results between the groups, the Mann-Whitney-U test was used. To compare proliferation between different cell layers within stenoses, the Wilcoxon test was performed. To test which factors showed an independent correlation with proliferation in the stenosis a multiple regression model was used. Linear regression analysis was used to define correlations between different parameters. When the data suggested a non-linear relation, non-linear regression analysis was used to define the relationship. A p-value of less than 0.05 was accepted as level of statistical significance.

RESULTS

Patient- and flow characteristics

The patient characteristics are summarized in table I. No differences in mean age were observed between the groups. Surgical repair of the vascular conduit in patients with AV-fistulas was performed on the average 10.9 months after reconstruction, and in bypass patients after an average time of 4.5 months ($p=0.002$). Two weeks after operation mean PSV and PSR were higher at the site of stenosis in the AV fistula group as compared to the bypass group ($p=0.002$ and $p=0.05$). However, no differences in PSV and PSR were observed shortly before correction of the stenotic segment.

Table I: Patient- and flow characteristics. Both the initial and final (before harvesting) values of peak systolic velocity (PSV) and peak shear rate (PSR) are presented (mean(SD)).

	Bypass (n=18)	AV fistula (n=17)	P Value
Age (years)	64 (12)	46 (21)	ns
Male/Female	12/4	9/6	ns
Age Graft (months)	4.5 (5.1)*	10.4 (7.6)	$p = 0.002$
Initial PSV (cm/s)	124 (96)*	341 (60)	$p = 0.002$
Final PSV (cm/s)	291 (90)	375 (71)	ns
Initial PSR (1/s)	52 (28)	78 (29)	$p=0.05$
Final PSR (1/s)	132 (59)	115 (60)	ns

Proliferation

In all stenotic samples KI-67 positive cells were observed, indicating a proper expression of this proliferation marker in the vessel specimens (Figure 1). In the control vein segments proliferation was almost zero (less than 0.2%, data not shown) in the different cell layers.

A summary of the results is presented in Table II. The degree of endothelial cell coverage of the lumen was approximately 50% in both groups (5136 cells counted), with a proliferation index in the endothelium covering the stenosis of 2.9% in the AV fistula group and 3.5% in the bypass group ($p=0.31$, Table II). No differences in proliferation index were observed between the AV fistula group and the bypass group (Table II) in the subendothelial-intimal areas (7234 cell counted), medial-intimal areas (8371 cells counted), medial areas (3893 cells counted) and adventitial areas (6124

cells counted). In both the AV fistula group and the bypass group subendothelial-intimal proliferation was higher as compared to medial-intimal proliferation (3.6% vs 1.1 and 3.5 vs 2.0%, respectively, $P<0.05$). In the adventitia, relatively high proliferation indices were found (7.5% for AV fistula stenoses and 4.2% for bypass stenoses, $p=0.26$). For the whole group, endothelial cell (EC) proliferation, increased from 0.3% in the subendothelial sections to 3.5% in the adventitia. This suggests increasing angiogenesis from the lumen to the outer layers of the vessel specimens.

Table II: Labeling fraction (%) of KI-67 positive cells in the different layers of the stenosis. Data are presented as means ($\pm$ SD). The total labeling fraction and the percentage proliferating vascular smooth muscle cells (VSMC) and endothelial cells (EC) are presented.

Proliferation (%)	Bypass (n=18)	AV-fistula (n=17)
subendothelial intima		
total	3.5 (3.9)	3.6 (3.6)
VSMC	3.3 (3.5)	3.3 (3.6)
EC	0.2 (0.6)	0.3 (1.0)
medial intima		
total	2.0 (1.8)	1.1 (1.5)
VSMC	1.2 (1.8)	0.9 (1.2)
EC	0.8 (1.8)	0.2 (0.5)
media		
total	2.6 (2.7)	1.7 (3.9)
VSMC	1.3 (2.1)	0.7 (1.4)
EC	2.3 (2.2)	1.0 (2.9)
adventitia		
total	4.2 (3.5)	7.5 (6.7)
VSMC	2.3 (2.0)	2.6 (2.8)
EC	1.9 (2.0)	4.9 (6.6)

Figure 1: Photomicrograph of an anti-MIB stained section of a stenosis derived from the venous anastomosis of an AV-fistula. Dark-brown stained nuclei are positive.

- a. Detail of the subendothelial intimal area. Note that most of the positively stained cells are localized in the intima adjacent to the lumen.
- b. Detail of the media/adventitia area. Note that most of the positive nuclei are endothelial cells covering small vessels. (L=lumen, I=intima, M=media, A=adventitia, magnification x400).

Figure 2: Photomicrograph of an anti-Ulex European stained section of an AV fistula.

- a. Detail of the intima and adjacent media. Note that the coverage of the lumen with endothelial cells is almost 100% and that the media contains many vessels.
- b. Detail of the media/adventitia area. The stain visualizes the high vascularity of this region. (L=lumen, I=intima, M=media, A=adventitia, magnification x200 and x400 respectively).

Correlations

Since no differences in proliferation patterns were found between AV fistulas and peripheral bypasses, they were treated as one group in correlating different parameters with proliferation. A multiple regression model was used to identify factors independently correlating with subendothelial-intimal proliferation, including endothelial cell coverage of the lumen of the stenosis, age of the lesion, peak shear rate at the site of the stenosis and graft type (vein or PTFE). The only factor independently influencing subendothelial-intimal cell proliferation was endothelial cell coverage (Table III). A positive correlation was found between endothelial coverage and subendothelial-intimal proliferation ($r=0.74$, $p=0.001$, Figure 3a). The final PSR could not be identified as an independent factor. However, when correlating PSR solely with subendothelial-intimal cell proliferation, a negative correlation was observed (non linear regression analysis, $r=-0.67$, $p<0.05$ Figure 3b). Therefore, a relation between endothelial cell coverage and peak shear rate was sought. Linear regression analysis demonstrated a negative correlation between these parameters (Figure 3c), suggesting that high shear results in loss of endothelial cell coverage, i.e., denudation (PTFE excluded since no endothelial cells are present in these specimens). A positive correlation was observed between proliferation of endothelial cells covering the stenosis and subendothelial-intimal cell proliferation ($y = 1.4 + 0.47 * X$, $r = 0.54$, $p<0.01$). The age of the stenosis could not be identified as an independent factor influencing proliferation, indicating that no distinct time course of proliferation is present in human stenotic tissue (Figure 3d).

Table III: Different parameters with a possible influence on subendothelial proliferation were included in a multiple regression model. From the 95% confidence limits it is shown that only the percentage of endothelial cells covering the lumen can be considered as an independent factor influencing subendothelial proliferation (EC = endothelial cell, PSR = peak shear rate, age = age of the stenosis).

factor	slope	95% confidence limits
ET cell coverage	0.06	0.03 - 0.08
log (PSR)	-0.3	-2.0 - 1.3
log (age)	-0.9	-2.1 - 0.2

Figure 3a: The relationship between the degree of coverage of the lumen by endothelial cells (EC) and subendothelial-intimal (SI) proliferation.

Figure 3b: The relationship between peak shear rate (PSR) and subendothelial-intimal (SI) proliferation. Nonlinear regression analysis revealed a decreasing proliferation with increasing shear rate.

Figure 3c: The relationship between PSR and endothelial cell coverage of the lumen within the stenosis (PTFE grafts excluded). With increasing shear rate, endothelial cell coverage becomes less, indicating denudation.

Figure 3d: The time course of subendothelial-intimal (SI) proliferation. The highest values of SI proliferation are observed in early developing stenoses. However, no significant correlation with time was observed.

DISCUSSION

The present study has three interesting findings. First, it shows increased proliferation in human vascular stenotic lesions derived from AV fistulas and peripheral bypass grafts, determined with the use of KI-67. Furthermore, our data suggest a stimulating role of the endothelium on proliferation in the intima, since proliferating KI-67 positive cells are mainly present in the area adjacent to the lumen and the degree of endothelial coverage of the lumen is correlated with subendothelial-intimal proliferation. In addition, with increasing peak shear rates, a decreasing endothelial cell coverage of the lumen was observed, suggesting that excessive high shear rate leads to denudation. This would imply an indirect inhibitory effect of high shear rate on proliferation in subendothelial areas by means of denudation.

Proliferation in human stenotic tissue

The contribution of proliferation of VSMC on the development of stenoses after vascular intervention in humans remains uncertain. In animal models stimulation and inhibition of replication of vascular smooth muscle cells by numerous pharmacological agents has been reported^{11,40-49}. However, administration of these agents in patients undergoing vascular intervention failed to show beneficial effects^{9,12,50}, indicating that extrapolation of findings in animal models to the human situation may be restricted and that proliferation patterns may differ between human and animal stenotic lesions^{9,12,13}. Therefore, more information on cell proliferation in human stenotic lesions is required to address this question. In the present study we defined replication in stenotic tissue with the use of the marker KI-67. This proliferation marker has been validated with the use of BrdU labeling and gains acceptance as a prognostic marker in cancer research¹⁵⁻¹⁷. The advantage of studying surgically harvested stenotic tissue from anastomoses of AV fistulas and peripheral bypasses is that the tissue samples are well defined and cannot be confused with non-stenotic vessel segments. This excludes the possibility of sample errors, as may be encountered in studies evaluating atherectomy derived tissue, reporting very low and very high replication rates^{13,14}. The replication rates in the different vessel layers as found in the present study (1.1 to 3.6% for the intima) are, however, also considerably lower as compared to the ones found in another study evaluating proliferation in AV fistula stenoses¹⁰. In that study the marker PCNA was used and proliferation rates as high as 15-20% were observed in the intima. High proliferation rates were also observed in a study using PCNA in the evaluating of proliferation in restenotic tissue after PTCA¹⁴. The use of PCNA may lead to overestimation of proliferation, as has been observed in validation studies, and may explain the reported high replication rates. The use of KI-67 may give a better representation of proliferation as compared to PCNA^{16,18}.

The endothelium and proliferation

In the present study replication was markedly higher in the subendothelial-intimal areas as compared to the medial-intimal layer, which has also been observed in animal studies²⁵. In addition, a positive correlation was found between the degree of endothelial

cell coverage of the lumen and proliferation indices in the subendothelial area (Figure 3a). This strongly suggests, that the endothelial cells covering the lumen of human stenotic lesions have a stimulating influence on proliferation. The data as depicted in Figure 3a, suggest that below an endothelial cell coverage of 30-40% subendothelial-intimal proliferation is very limited (mean $0.53 \pm 0.86 \%$), while a coverage over 40% is associated with significantly higher proliferation indices (mean $4.54 \pm 3.29 \%$, $p < 0.01$, Mann-Whitney-U test). In vitro studies have demonstrated that endothelial cells can release a variety of growth regulating factors^{11,51-53}. Endothelium-derived-relaxing-factor (EDRF), heparin, and prostacyclin (PGI-2) can be released by endothelial cells and may inhibit VSMC replication^{26-28,54-57}. Growth-promoting factors released by the endothelium are platelet-derived-growth-factor (PDGF) and angiotensin-II (AG-II)^{28,51,58}. The effect of the endothelium on the growth response of VSMC probably depends on the balance between growth stimulating and inhibiting factors^{55,57}. From our observation it can be concluded that the endothelium, when present, promotes replication, indicating a dysbalance between stimulating and inhibiting factors in favor of stimulation. This has been described previously in an in vitro study, using porcine aortic organ cell cultures³². In a recently published study, using human saphenous vein organ culture, also higher proliferation indices were found in the intima with an intact endothelial cell layer as compared to denuded vein cultures³³. The results of this in vitro study are in concordance with the results from the present study. We found a positive correlation between proliferation of endothelial cells covering the stenosis and subendothelial-intimal proliferation. Proliferation of endothelial cells covering the stenosis suggests that these cells are not in their normal quiescent state. This may be due to partial denudation caused by high flow velocities, resulting in a disruption of cell-cell interaction between endothelial cells, initiating proliferation in order to reestablish a confluent monolayer⁵⁹. The proliferative state of the endothelial cells may be related with a loss of growth inhibitory function, resulting in high subendothelial-intimal proliferation as is suggested by the positive correlation between proliferation of endothelial cells covering the lumen of the stenosis and subendothelial-intimal proliferation. This hypothesis is supported by an in vitro study, reporting increased proliferation of VSMC in areas covered with proliferating endothelial cells^{34,35}.

Peak Shear Rate and Proliferation

The influence of flow characteristics on the development of intimal hyperplasia has been extensively investigated in animal models, showing that low shear conditions lead to increased intimal thickening, while high shear rate inhibits the development of intimal thickening¹⁹⁻²⁵. We therefore hypothesized that the extent and distribution of proliferation may differ between stenotic lesions derived from AV fistulas (relatively high-flow system) and peripheral bypasses (relatively low-flow system). However, we found no apparent differences in proliferation between stenotic tissue derived from these two groups, suggesting that volume flow may not be a determining factor in proliferation. The PSR values just before repair of the stenosis (Table I) showed similar values in the group AV fistulas and bypasses. To establish a possible effect of flow on VSMC replication, the PSR values just before harvesting were correlated with

subendothelial proliferation. A negative correlation was found between final PSR and subendothelial proliferation index ($Y = 16 - 2.8 \ln(\text{PSR})$, $r=0.67$, $p<0.05$, Figure 3b), but multiple regression analysis revealed that final PSR was not an independent determinant of subendothelial proliferation. Therefore, we hypothesized that shear rate might have an indirect effect on proliferation through its influence on the endothelial cells.

Linear regression analysis showed that with increasing shear rate, endothelial cell coverage decreases, suggesting a denuding effect of high shear rate. The calculation of PSR, based on the Haagen-Poiseuille formula, can only give a rough estimation of the actual shear forces exerted on the endothelial cell layer, since a parabolic symmetric velocity profile is assumed. It is likely that in the systolic phase the local velocity profile is blunted, especially in narrowed segments. Therefore, the PSR as calculated in this study gives an underestimation of the actual PSR. However, the negative correlation between PSR and endothelial cell coverage of the lumen indicates that PSR may be of physiologic importance. Peak shear rate possibly gives a better characterization of the local pulsatile shear rate conditions than does mean shear rate, which has been used by other investigators²⁴. This is supported by findings in an animal study, where no correlation was observed between mean shear stress and proliferation²⁴. Our data support the hypothesis that high shear rate inhibits the development of intimal thickening. The exact mechanism by which shear rate influences replication is unclear. Many authors and our data suggest a key role of the endothelium in this mechanism, which is supported by the finding that shear rate can initiate the release of EDRF and can lead to an increase in endothelial PDGF mRNA levels²⁶⁻²⁸. In addition, variation in shear rate levels results in modification of the structure of endothelial cells⁶⁰⁻⁶².

No independent significant correlation between proliferation index and age of the conduit was observed. The highest values of proliferation indices in the subendothelial-intimal layer were found in early developed stenoses, but no significant correlation with time was demonstrated ($p=0.10$, Figure 3d), which confirms the findings of other studies evaluating proliferation in human stenotic tissue^{10,13,14}. This suggests that a small but continuously elevated proliferation is present rather than a high peak in proliferation.

In the present study neovascularisation was observed in the stenotic lesion as well as in the media and adventitia. From the lumen to the adventitia, an increasing proportion of proliferating cells were identified as endothelial cells, covering microvessels. Angiogenesis has been observed previously in human stenotic lesions¹⁰. The close correlation of subendothelial-intimal proliferation with the degree of endothelial cell coverage and the large percentage of proliferating endothelial cells covering microvessels in the stenosis, indicate that endothelial cells may be a major source for growth factors influencing proliferation patterns in human stenotic lesions. The exact contribution of angiogenesis to the development of stenotic lesions and atherosclerosis is unclear, but certainly deserves further investigation.

In summary, a 0.9 - 3.7% proliferation rate was observed in well defined human stenotic tissue, derived from severely or occluded peripheral bypasses or AV-fistulas, with the use of KI 67. Subendothelial proliferation in the intima was positively correlated with the presence of endothelial cells covering the lumen, suggesting a stimulating role for endothelial cells on subendothelial-intimal proliferation. In addition, with increasing peak shear rate, a decrease in endothelial cell coverage of the lumen was observed, indicating that denudation occurs in areas of excessive high shear. This may indicate that high shear rate has an indirect inhibitory effect on proliferation by means of denudation.

ACKNOWLEDGEMENT

We are indebted to Jean Willigers for the technical assistance. We thank Petra Aarts and Anique Janssen for excellent technical support. We thank Dr J Buth and Dr M Idu for supplying us with samples from the Catharina Hospital Eindhoven.

REFERENCES

1. Harris PL, Veith FJ, Shanik GD, Nott D, Wengerter KR, Moore DJ. Prospective randomized comparison of in situ and reversed infrapopliteal vein grafts. *Br J Surg* 1993; 80:173-176.
2. Berkowitz HD, Greenstein S, Barker CF, Perloff LJ. Late failure of reversed vein bypass grafts. *Ann Surg* 1989; 210:782-786.
3. Taylor LM, Edwards JM, Porter JM. Present status of reversed vein bypass grafting: Five-year results of a modern series. *J Vasc Surg* 1990; 11:193-206.
4. Jenkins AMcL, Buist TAS, Glover SD. Medium-term follow-up of forty autogenous vein and forty polytetrafluoroethylene (Gore-Tex) grafts for vascular access. *Surgery* 1980; 88:667-672.
5. Swedberg SH, Brown BG, Sigley R, Wight TN, Gordon D, Nicholls SC. Intimal fibromuscular hyperplasia at the venous anastomosis of PTFE grafts in hemodialysis patients. *Circulation* 1989; 80:1726-1736.
6. Berkowitz HD, Fox AD, Deaton DH. Reversed vein graft stenosis: Early diagnosis and management. *J Vasc Surg* 1992; 15:130-142.
7. Harris PL, How TV, Jones DR. Prospectively randomized clinical trial to compare in situ and reversed saphenous vein grafts for femoropopliteal bypass. *Br J Surg* 1987; 74:252-255.
8. Tordoir JHM, Herman JM, Kwan TS, Diderich PM. Long-term follow-up of the polytetrafluoroethylene (PTFE) prosthesis as an arteriovenous fistula for hemodialysis. *Eur J Vasc Surg* 1988; 2:3-7.
9. Liu MW, Roubin GS, King III SB. Restenosis after coronary angioplasty. Potential biologic determinants and role of intimal hyperplasia. *Circulation* 1989; 79:1374-1387.
10. Rehker M, Nicholls S, Ferguson M, Gordon D. Cell proliferation in human arteriovenous fistulas use for hemodialysis. *Arterioscler Thromb* 1993; 13:609-617.
11. Clowes AW, Reidy MA. Prevention of stenosis after vascular reconstruction: Pharmacologic control of intimal hyperplasia - A review. *J Vasc Surg* 1991; 6:885-891.
12. Ferrel M, Fuster V, Gold HK, Chesebro JH. Choosing appropriate experimental animal model for the prevention of restenosis. *Circulation* 1992; 85:1630-1631.
13. O'Brien ER, Alpers CE, Stewart DK, et al. Proliferation in primary restenotic coronary atherectomy tissue. Implications for antiproliferative therapy. *Circ Res* 1993; 73:223-231.
14. Pickering JG, Weir L, Jekanowski J, Kearny MA, Isner JM. Proliferative activity in peripheral and coronary atherosclerotic plaque among patients undergoing percutaneous revascularisation. *J Clin Invest* 1993; 91:1469-1480.
15. Verheyen R, Kuypers HJH, Schlingeman RO, et al. Ki-67 detects a nuclear matrix-associated proliferation-related antigen. *J Cell Science* 1989; 92:123-130.
16. Sasaki K, Matsumura K, Tsuji T, Shinozaki F, Takahashi M. Relationship between labeling indices of Ki-67 and BrdUrd in human malignant tumors. *Cancer* 1988; 62:989-993.
17. Veronesi SM, Gambacorta M, Gottardi O, Scanzi F, Ferrari M, Lampertico P. Proliferation index as a prognostic marker in breast cancer. *Cancer* 1993; 71:3926-3931.
18. Van Dierendonck JH, Wijsman JH, Keyzer R, Van De Velde CJH, Cornelisse CJ. Cell-cycle-related staining patterns of anti-proliferating cell nuclear antigen monoclonal antibodies- Comparison with BRDUrd labeling and KI-67 staining. *Am J Pathol* 1991; 138:1165-1172.
19. Zarins CK, Bomberger RA, Glagov S. Local effects of stenoses: Increased flow velocity inhibits atherogenesis. *Circulation* 1981; 64:11221-11227.
20. Zarins CK, Giddens DP, Bharadvaj BK, Sottiurai VS, Mabon RF, Glagov S. Carotid bifurcation atherosclerosis. Quantitative correlation of plaque localization with flow velocity profiles and wall shear stress. *Circ Res* 1983; 53:502-514.
21. Asakura T, Karino T. Flow patterns and spatial distribution of atherosclerotic lesions in human coronary arteries. *Circ Res* 1990; 66:1045-1066.

22. Friedman MH, Barger CB, Deters OJ, Hutchins GM, Mark FF. Correlation between wall shear and intimal thickness at a coronary artery branch. *Atherosclerosis* 1987; 68:27-33.
23. Sterpetti AV, Cucina A, Santoro L, Cardillo B, Cavallaro A. Modulation of arterial smooth muscle cell growth by hemodynamic forces. *Eur J Vasc Surg* 1992; 6:16-20.
24. Kohler TR, Jawien A. Flow affects development of intimal hyperplasia after arterial injury in rats. *Arterioscler Thromb* 1992; 12:963-971.
25. Kohler TR, Kirkman TR, Kraiss LW, Zierler BK, Clowes AW. Increased blood flow inhibits neointimal hyperplasia in endothelialized vascular grafts. *Circ Res* 1991; 69:1557-1565.
26. Cooke JP, Rossitch E, Andon NA, Loscalzo J, Dzau VJ. Flow activates an endothelial potassium channel to release an endogenous nitrovasodilator. *J Clin Invest* 1991; 88:1663-1671.
27. Buga GM, Gold ME, Fukuto JM, Ignarro LJ. Shear stress-induced release of nitric oxide from endothelial cells grown on beads. *Hypertension* 1991; 17:187-193.
28. Hsieh HJ, Li NQ, Frangos JA. Shear stress increases endothelial platelet-derived growth factor mRNA levels. *Am J Physiol* 1991; 260:H642-H646.
29. Cashin WL, Sanmarco ME, Nessim SA, Blankenhorn DH. Accelerated progression of atherosclerosis in coronary vessels with minimal lesions that are bypassed. *New Eng J Med* 1984; 311:824-828.
30. Clowes AW, Reidy MA, Clowes MM. Mechanisms of stenoses after arterial injury. *Lab Invest* 1983; 49:208-215.
31. Gordon D, Schwartz SM. Cell proliferation in human atherosclerosis. *Trends Cardiovasc Med* 1991; January/February:24-28.
32. Koo EWY, Gotlieb AI. Endothelial stimulation of intimal cell proliferation in a porcine aortic organ culture. *Am J Pathol* 1989; 134:497-503.
33. Allen KE, Varty K, Jones L, Sayers RD, Bell PRF, London NJM. Human venous endothelium can promote intimal hyperplasia in a paracrine manner. *J Vasc Surg* 1994; 19:577-584.
34. Mey de JGR, Dijkstra EH, Vrijdag MJF. Endothelium reduces DNA synthesis in isolated arteries. *Am J Physiol* 1991; 260:H1-H7.
35. De Mey JGR, Schiffers PM. Effects of the endothelium on growth responses in arteries. *J Cardiovasc Pharmacol* 1993; 21:s22-s25.
36. Buth J, Disselhoff B, Sommeling C, Stam L. Color-flow duplex criteria for grading stenosis in infrainguinal vein grafts. *J Vasc Surg* 1991; 14:716-728.
37. Hoeks APG, Brands PJ, Smeets FAM, Reneman RS. Assessment of distensibility of superficial arteries. *Ultrasound Med Biol* 1990; 16:121-128.
38. Reneman RS, Merode van T, Brands PJ, Hoeks APG. Inhomogeneities in arterial wall properties under normal and pathological conditions. *J Hypertens* 1992; 10:S35-S39.
39. Hofstra L, Willigers JM, Huvers FC, et al. Short-term variations in elastic properties of a muscular artery in humans. *Clin Sci* 1993; 50:472-480.
40. Daemen MJAP, Lombardi DM, Bosman FT, Schwartz SM. Angiotensin II induces smooth muscle cell proliferation in the normal and injured rat arterial wall. *Circ Res* 1991; 68:450-456.
41. Powell JS, Clozel JP, Muller RKM, et al. Inhibitors of angiotensin-converting enzyme prevent myointimal proliferation after vascular injury. *Science* 1989; 245:186-188.
42. Vashiht R, Sian M, Franks PJ, O'Malley MK. Long-term reduction of intimal hyperplasia by the selective alpha-1 adrenergic antagonist doxazosin. *Br J Surg* 1992; 79:1285-1288.
43. Powell JS, Rouge M, Muller RK, Baumgartner HR. Cilazapril suppresses myointimal proliferation after vascular injury: effects on growth factor induction in vascular smooth muscle cells. *Basic Res Cardiol* 1991; 86:65-74.
44. Mitsuka M, Nagae M, Berk BC. Na⁺-H⁺ exchange inhibitors decrease neointimal formation after rat carotid injury. Effects on smooth muscle cell migration and proliferation. *Circ Res* 1993; 73:269-275.

45. Kranzhöfer R, Schirmer J, Schömig A, et al. Suppression of neointimal thickening and smooth muscle cell proliferation after arterial injury in the rat by inhibitors of Na⁺-H⁺ exchange. *Circ Res* 1993; 73:264-268.
46. Kleef van EM, Smits JFM, Mey de JGR, et al. α 1-Adrenoreceptor blockade reduces the angiotensin II-induced vascular smooth muscle cell in the rat thoracic aorta and carotid artery. *Circ Res* 1992; 70:1122-1127.
47. Jawien A, Bowen-Pope DF, Lindner V, Schwartz SM, Clowes AW. Platelet-derived growth factor promotes smooth muscle migration and intimal thickening in a rat model of balloon angioplasty. *J Clin Invest* 1992; 89:507-511.
48. Dryjski M, Mikat E, Björnsson TD. Inhibition of intimal hyperplasia after arterial injury by heparins and heparinoid. *J Vasc Surg* 1988; 8:623-633.
49. Clowes AW, Clowes MM, Vergel SC, et al. Heparin and cilazapril together inhibit injury-induced intimal hyperplasia. *Hypertension* 1991; 18:II65-II69.
50. McBride W, Lange RA, Hillis LD. Restenosis after successful coronary angioplasty. Pathophysiology and prevention. *New Eng J Med* 1988; 318:1734-1737.
51. Ross R. The pathogenesis of atherosclerosis - An update. *New Eng J Med* 1986; 314:488-499.
52. Ralevic V, Milner P, Hudlicka O, Kristek F, Burnstock G. Substance P is released from the endothelium of normal and capsaicin-treated rat hind-limb vasculature, in vivo, by increased flow. *Circ Res* 1990; 66:1178-1183.
53. Ross R. Atherosclerosis: A defense mechanism gone awry. *Am J Pathol* 1993; 143:987-1002.
54. Rubanyi GM, Freay AD, Kauser K, Johns A, Harder DR. Mechanoreception by the endothelium: Mediators and mechanisms of pressure- and flow-induced vascular responses. *Blood Vessels* 1990; 27:246-257.
55. Lüscher TF. Imbalance of endothelium-derived relaxing and contracting factors. A new concept in hypertension? *Am J Hypertens* 1990; 3:317-330.
56. Limanni A, Fleming T, Molina R, et al. Expression of genes for platelet-derived growth factor in adult human venous endothelium. A possible non-platelet-dependent cause of intimal hyperplasia in vein grafts and perianastomotic areas of vascular prostheses. *J Vasc Surg* 1988; 7:10-20.
57. Furchgott RF, Vanhoutte PM. Endothelium-derived relaxing and contracting factors. *FASEB J* 1989; 3:2007-2018.
58. Schwartz CJ, Valente AJ, Sprague EA, Kelly JL, Nerem RM. The pathogenesis of atherosclerosis: An overview. *Clin Cardiol* 1991; 14:1-16.
59. Schwartz SM, Liaw L. Growth control and morphogenesis in the development and pathology of arteries. *J Cardiovasc Pharmacol* 1993; 21:s31-s49.
60. Langille BL, Reidy M, Kline R. Injury and repair of endothelium at sites of flow disturbances near abdominal aortic coarctations in rabbits. *Arterioscler Thromb* 1988; 6:146-154.
61. Gottlieb AI, Langille BL, Wong MKK, Kim DW. Structure and function of the endothelial cytoskeleton. *Lab Invest* 1991; 65:123-137.
62. Langille BL, Graham JJ, Kim D, Gottlieb AI. Dynamics of shear induced redistribution of F-actin in endothelial cells in vivo. *Arterioscler Thromb* 1991; 6:1814-1820.

CHAPTER 12

GENERAL DISCUSSION

MECHANICAL AND FLOW CHARACTERISTICS AFFECTING INTIMAL HYPERPLASIA IN PERIPHERAL BYPASS GRAFTS AND GRAFT ARTERIO VENOUS FISTULAS IN HUMANS

The principal goal of the studies described in this thesis was to clarify the role of perianastomotic mismatch in elastic properties and unphysiological flow conditions on the occurrence of stenotic intimal thickening in peripheral bypasses and graft arterio-venous fistulas. Based on the literature, we hypothesized that an increase in perianastomotic mismatch in elastic properties leads to an increase in the occurrence of stenoses. In addition, we hypothesized that a low shear rate in these vascular reconstructions results in stenotic intimal thickening.

The first step to investigate the role of mechanical factors in the occurrence of stenotic intimal thickening in peripheral bypasses and graft AV fistulas was to test the applicability of Vessel Wall Doppler Tracking (VWDT) in the assessment of elastic properties of these vascular reconstructions. The use of the B-mode image in the positioning of the measurements turned out to be crucial in the determination of elastic properties at defined sites in the vascular systems. By using the anastomosis as a landmark, the assessment of differences in elastic properties around anastomoses was possible, allowing the quantification of peri-anastomotic mismatch in elastic properties. For the first time, assessment of elastic profiles of peripheral bypass grafts and graft AV fistulas was possible, indicating that VWDT is a promising and very useful new tool to obtain insight in the mechanics of vascular reconstructions. In the first chapters (chapter 5-8), we studied the elastic behavior of peripheral bypasses and arteriovenous fistulas constructed using different techniques and graft materials. In addition, we tried to assess local flow characteristics in peripheral bypasses and arterio-venous fistulas at corresponding sites. One of the limitations in the measurement of wall shear rate with the use of ultrasound is that the determination of vessel diameter on a B-mode image is inaccurate. Since VWDT provides accurate determination of diameter, the calculation of peak shear rate was possible. This provided a rough estimate of the actual shear stresses exerted on the vessel wall.

A second step was to correlate initial peri-anastomotic mismatch and local flow conditions with the occurrence of stenosis in peripheral bypasses and AV fistulas during follow-up. It was observed that the occurrence of stenotic intimal thickening in arteriovenous fistulas is associated with high shear rate and not with a mismatch in elastic properties (chapter 9). In addition, a high initial local peak shear rate in venous peripheral bypass grafts was associated with the occurrence of stenotic intimal thickening during follow up. A greater initial peri-anastomotic mismatch in strain was observed in the proximal anastomoses of bypasses developing a stenosis as compared with the bypasses without a stenosis.

MISMATCH IN ELASTIC PROPERTIES AND INTIMAL THICKENING

In the AV fistula group no association was observed between a greater peri-anastomotic mismatch in elastic properties and intimal thickening. In the bypass group, a greater peri-anastomotic mismatch in local strain at the proximal anastomosis was associated with stenotic intimal thickening. However, for the distal anastomosis no association between peri-anastomotic mismatch and the occurrence of stenoses was observed. In addition, multiple regression analysis, including flow- and mismatch parameters, demonstrated that only a high initial peak shear rate was correlated with the occurrence of intimal thickening. For a mismatch in peri-anastomotic strain no correlation with the occurrence of intimal lesions was found in the multiple regression model. The observations from the prospective studies in AV fistulas and peripheral bypasses suggest that initial peri-anastomotic mismatch in elastic properties does not play a major role in the occurrence of intimal lesions in these vascular reconstructions. A possible explanation for the absence of a correlation between peri-anastomotic mismatch in elastic properties and stenotic intimal thickening may be that the differences in distension between graft and native vessels are less than one would expect from the intrinsic material properties. The observations in chapter 6 and 7 suggest that the distension of the graft close to the anastomosis is influenced by the distension of the native vessel and vice versa. Placement of a stiff graft in a compliant arterial system leads to a decrease in distension of the arterial segments close to the anastomosis. On the other hand, the distension of the native vessel leads to passive distension of the adjacent graft segment. This suggests graft-native vessel interaction. This interaction between graft and native vessel system was observed when measuring elastic properties at a distance of approximately 1 centimeter from the anastomosis. One might speculate that more interaction between graft and native vessels is present at points closer to the anastomosis. Therefore, the actual difference in distension between graft and native vessel close to the anastomosis may be even less than was observed in our studies.

Another explanation for the absence of a correlation between peri-anastomotic mismatch in elastic properties and stenotic intimal thickening may be that the in vivo strain at the anastomosis is not enough to initiate the replication of VSMC and to

stimulate the production of matrix by these cells. In *in vitro* studies, VSMC started to produce matrix material when exposed to repetitive cyclic stretching. However, the values of strain in these studies were around 20-25%, which is very high when compared to the strain values found in our studies. We observed strain values of 1-5% around the anastomosis, which may not be sufficient to initiate VSMC responses.

FLOW CHARACTERISTICS AND INTIMAL THICKENING

The finding that intimal thickening in the venous outflow segment of AV fistulas and in venous peripheral bypass grafts is associated with initial high peak shear rate levels seems to be in contrast with numerous studies demonstrating that increased intimal thickening occurs mainly in areas of low shear rate. There are several possible reasons for this apparent discrepancy. First of all, most investigators studied the effect of varying shear rates on intimal thickening after the initial injury had already occurred. This is quite different from the initial situation at the venous outflow segment of the AV fistula and a saphenous vein bypass graft, where intact endothelium is present. Another point that has to be considered is the range of shear rates investigated. In some studies a comparison was made between the effect of a decreased flow and the effect of normal arterial flow. This indicates that the range of shear rates investigated lies between normal and low, which is different from the magnitude of the shear rate values encountered in AV fistulas. In the venous anastomoses of AV fistulas peak systolic velocity values as high as 300-350 cm/s are present, indicating excessive high shear rate levels. In normal arteries peak systolic velocity varies between 60-120 cm/s, leading to substantially lower shear rate levels for the same vessel diameter. This implies that the shear rate values as found in AV fistulas are beyond physiological values. Also in the proximal segments of reversed saphenous vein grafts elevated peak systolic velocities may be present, since this is the narrowest segment in the graft. Another point which has to be considered, when comparing the results in this thesis with data from animal studies, is the fact that in animal studies intimal hyperplasia does not proceed to subocclusive or occlusive stenotic lesions, but is restricted in its extent. The changes in intimal thickening as found in the animal models may just be an adaptational process, and may therefore be dissimilar to occlusive intimal hyperplasia as found in humans.

Another consideration may be the use of different vessel types and reconstructions when studying the effect of varying flow on the development of intimal thickening. Balloon denudation in rat models are mostly performed in the carotid artery, whereas intimal thickening in AV fistulas in humans predominantly occurs at the site of the venous anastomoses or the efferent vein. At the arterial site or at the arterial anastomoses of the AV fistulas intimal thickening hardly develops despite similar flow conditions. This suggests that excessive high shear rates may lead to denudation in veins due to a high susceptibility of the venous endothelial cells to the injurious effect of shear stress. Some investigators have used PTFE grafts, and measured the thickness

of the neo-intima within the graft at varying flow levels. Since initially PTFE grafts are not covered with endothelial cells and no media is present as the origin of the proliferating and migrating vascular smooth muscle cells, the mechanism of intimal thickening in this model has to be different. In addition, occlusive intimal hyperplasia associated with PTFE grafts hardly ever occurs within the graft, but develops at the site of the anastomoses. Therefore, studying intimal thickness within PTFE grafts at varying flow levels probably has limited relevance for the clinical situation.

The prospective studies on the influence of the initial flow characteristics on the occurrence of stenoses in AV fistulas and peripheral bypasses (chapter 9 and 10) showed that initial high local peak shear rate is associated with the occurrence of stenoses in the follow-up period. This seems to be in contrast with the observation in the study described in chapter 11, showing a decreased proliferation with increasing local peak shear rate, suggesting an inhibitory effect of high shear rate on the development of intimal thickening. When combining the results from these studies we propose the following hypothesis on the relation between flow characteristics and intimal thickening in human vascular grafts:

- 1. An initial high peak shear rate, as present in the venous anastomoses and efferent vein segments in AV fistulas and in the proximal segment of reversed saphenous vein bypass grafts, results in partial denudation and intimal damage in venous segments, thereby initiating a vascular response leading to local intimal thickening.

2. A local high peak shear rate at the site of the stenosis inhibits proliferation of VSMC in the intima, possibly by a denudating effect on endothelial cells covering the lumen. Thus after the initiation of vascular injury by a high peak shear rate as discussed under 1, the velocity by which the stenosis develops is inhibited by a high peak shear rate.

LIMITATIONS OF THE STUDY

The technique by which the studies were performed, Vessel Wall Doppler Tracking, has been validated in previous studies reported from our institution. In chapter 4 the data on short-term variations in elastic properties in the femoral artery in humans are presented. We found considerable variation with time, which could not be attributed to changes in pressure, indicating that active vessel wall behavior is responsible for the variation. Changes in vascular muscle tone may be the result of changes in flow and/or the activity of the local renine-angiotensin system (RAS) system. This study was performed in healthy young subjects. So far, we do not know whether short term variations in elastic properties are also present in saphenous vein bypass grafts, femoral and popliteal arteries in atherosclerotic patients, and brachial arteries and veins in hemodialysis patients.

A major restriction of the assessment of compliance and distensibility coefficient in peripheral bypass grafts and AV fistulas is the absence of local pulse pressure information. The calculation of compliance and distensibility based on non-invasively

obtained pulse pressure values may be inaccurate. In chapter 5 we have used compliance and distensibility (CC and DC), calculated with the use of digital blood pressure values obtained with Finapres. We observed a good correlation between DC and CC values of the common femoral artery obtained with either local intra-arterial measurements or Finapres values in patients with diseased vessels (see chapter 5). However, the use of blood pressure values remote from the assessment of distension is under discussion and has resulted in the development of new approaches to determine local pressure and vessel wall distensions simultaneously.

Another point is that the compliance mismatch theory is based on the thought that peri-anastomotic differences in strain and cross-sectional area-increase possibly lead to local mechanical stress and/or flow disturbances. The pressure leading to strain and cross-sectional area increase (i.e. pulse pressure) is of a lesser importance. Based on the arguments mentioned above we decided to use parameters not corrected for pulse pressure, like local strain (RD) and cross-sectional area-increase (AI) in the later studies of the thesis.

The assessment of shear rate in the human in vivo situation is difficult. We used peak shear rate, which can only give a rough estimation of the actual peak wall shear rates. With the use of a new ultrasound system to estimate wall shear rate in vivo in humans, we found an almost linear relationship between velocity in the center of the vessel and wall shear rate. These measurements were performed in the carotid artery in healthy volunteers under well-defined conditions. This implies that velocity in the center of the vessel is a major determinant of wall shear rate, and indicates that the use of PSR in this thesis was justified. The scaling factor may vary between different vascular territories and may be influenced by hemodynamic conditions.

CLINICAL IMPLICATIONS

The concept that high shear rates lead to denudation in peripheral saphenous vein bypass grafts and venous outflow segments of AV fistulas has implications for the method of construction of these vascular conduits. Modifications of operation techniques should aim at decreasing peak shear rate levels, thereby reducing the possible injurious effects of excessive high shear rate, at least in venous vessel segments. In constructing peripheral bypasses from saphenous vein material, the vascular surgeon is restricted by the dimensions of the vein and the length of the bypass. Theoretically, the in situ technique, with its natural taper, overcomes the problem of high shear rate in the proximal graft segment and therefore seems to be the most ideal graft in terms of flow. In constructing AV fistulas, a decrease in flow velocity may be accomplished by the use of inversely tapered grafts (i.e. 4 to 7 mm) and the construction of cuffs at the venous anastomosis.

The assessment of proliferation in human stenotic tissue revealed elevated proliferative activity and suggests an important contribution of proliferation of vascular smooth

muscle cells to the development of intimal thickening. This implicates that inhibition of proliferation may help in decreasing the progress of intimal lesions. One might speculate that angiotensin converting enzyme (ACE) inhibitors, which do also decrease the activity of the peripheral RAS system, may help in preventing intimal growth by decreasing proliferation. In this study we also found a high level of proliferating endothelial cells in the stenosis, indicating neovascularisation. This finding certainly needs further investigation in order to clarify the role of neovascularisation in the process of intimal thickening. A better insight in this mechanism may help in defining therapeutic strategies to decrease the development of intimal thickening.

The clinical implication of the findings in this study for other clinical problems related to intimal hyperplasia, such as restenosis after PTCA, is unclear. Although the time course and incidence of intimal thickening in vascular grafts and after PTCA are similar, it is obvious that compliance mismatch cannot play a role in the initiation or perpetuation of intimal thickening after PTCA. The role of flow in the development of restenosis after PTCA in humans is still unclear, but may be further elucidated with the use of intravascular ultrasound techniques.

FUTURE DIRECTIONS

In studying the effect of mismatch in elastic properties more attention has to be focused on the region close to the anastomosis. One interesting observation from our study is that graft-vein or graft-artery interaction may occur in the regions adjacent to the anastomoses. More knowledge on this subject may help in constructing better anastomoses and may improve the development of vascular prostheses.

In the prospective studies, relating initial flow conditions to the occurrence of intimal hyperplasia, a rough approximation of shear rate, the peak shear rate, was used. Improved techniques, like the new ultrasound system to estimate wall shear rate in the human in vivo situation may further clarify the correlation between intimal thickening and flow, since a more detailed picture of velocity patterns is provided.

Another improvement may be the continuous on-line registration of vessel wall distension. This allows investigation of beat-to-beat changes in elastic behavior of vessels, which may help to increase our understanding of the function of medium-sized arteries in pressure and flow homeostasis. A different topic is the assessment of wall thickness with the use of ultrasound techniques. This may increase our knowledge on the natural history of vascular disease leading to intimal thickening. In addition, the effect of pharmacological intervention in patients with vascular disease can be evaluated when accurate wall thickness measurement is possible.

SUMMARY

MECHANICAL AND FLOW CHARACTERISTICS AFFECTING INTIMAL HYPERPLASIA IN PERIPHERAL BYPASSES AND GRAFT AV FISTULAS IN HUMANS

In chapter 1 and 2 the background and principal goals of the thesis are described. The main goal of the studies was to investigate the role of mechanical factors in the development of intimal thickening in peripheral bypass grafts and arteriovenous fistulas in patients. The development of intimal thickening in these arterial reconstructions leads to hemodynamically significant stenoses in a large fraction of patients, often necessitating surgical interventions. This results in considerable morbidity and hospitalisation. More insight in the initiation and development of intimal thickening in peripheral bypass grafts and AV fistulas may help in designing new surgical approaches in order to optimize mechanical and flow conditions, thereby preventing the development of intimal hyperplasia. In addition, better understanding of the biology of intimal thickening in humans may help in developing pharmacological interventions that inhibit the development of intimal thickening.

VESSEL WALL DOPPLER TRACKING

Information on the elastic properties of peripheral bypasses and graft arteriovenous fistulas has been very difficult to obtain, since suitable systems to assess the dimensions at defined sites in the vascular system were absent. With the introduction of Vessel Wall Doppler Tracking (VWDT), combining B-mode imaging and off-line data processing, the in vivo measurement of elastic properties of vessels in humans at defined sites in a vascular system has become possible. This possibility allows for the assessment of elastic properties around the anastomoses of vascular reconstructions. Quantification of peri-anastomotic differences in elastic properties (mismatch) is crucial to investigate its possible role in the development of anastomotic stenoses in vascular reconstructions.

In chapter 3 VWDT is described and the possible application of the technique in vascular research is discussed.

In chapter 4 the short-term variations in elastic properties of the common femoral artery were studied in 10 young volunteers without vascular disease. The principal finding of this study is that the elastic properties of this artery change considerably

within a relatively short period of time. In addition, it was observed that the changes in elastic properties were not related to changes in local blood pressure, suggesting that changes in vascular smooth muscle tone are responsible for the variation of elastic properties of this medium sized artery.

MISMATCH IN ELASTIC PROPERTIES AROUND ANASTOMOSES OF PERIPHERAL BYPASSES AND GRAFT AV FISTULAS IN HUMANS

The function of the graft as a vascular conduit is to transport blood volume without the loss of pulsatile energy to provide sufficient blood flow in the periphery. The transport function of grafts is largely determined by its mechanical properties. In order to obtain basic information on the mechanical profiles of grafts, we started to assess the elastic properties of different graft types at defined sites in the graft and native vessels in peripheral bypasses and AV fistulas with the use of Vessel Wall Doppler Tracking. By using this method we were able to provide the first in vivo data on mismatch in elastic properties around anastomoses of vascular grafts in humans.

In chapter 5 the mechanical profiles of in situ and reversed saphenous vein grafts are presented. One of the principal findings of this study was that in the in situ graft a proper match in compliance was found around the proximal anastomosis in contrast to the reversed graft, where a large drop in compliance around the proximal anastomosis was observed. However, in the course of the graft compliance in the reversed saphenous vein graft was remarkably stable, whereas in the in situ grafts an important drop in compliance was found.

In chapter 6 the elastic properties of PTFE peripheral bypass grafts and matched saphenous vein grafts were compared. As expected, the distension values of saphenous vein grafts were higher as compared to PTFE grafts. An interesting finding in this study is, that distension in the proximal graft segment of PTFE grafts was higher as compared to the distal graft segments, which is remarkable, since PTFE prostheses have homogeneous material properties throughout the graft. A possible explanation may be that distension of the inflow artery leads to a passive distension of the most proximal graft segments. The finding that distension values of the inflow segment are positively correlated with distension values of the proximal segments supports the hypothesis of an artery-graft interaction.

In chapter 7 the differences in peri-anastomotic mismatch in elastic properties between the arterial and venous anastomoses of graft AV fistulas for hemodialysis were compared in order to find clues for the predisposition of intimal hyperplasia to develop at the venous site. The principal finding of this study was that at the arterial anastomoses a decrease in compliance was observed, whereas a sudden increase in compliance was observed at the venous site. The latter condition is associated with increased particle residence time and flow disturbances, suggesting low shear rates.

Since numerous animal studies reported increased intimal thickening in areas of low shear rate, we postulated that the sudden increase in compliance at the venous anastomoses of graft AV fistulas may be the reason for the predisposition of intimal hyperplasia to develop at this site. At the venous anastomosis of the AV fistulas graft-native vessel interaction was observed. We found that an increase in the distension of the outflow vein was associated with an increase in distension of the graft close to the anastomosis, suggesting that distension of the vein leads to a passive distension of the graft at the site of the anastomosis.

In chapter 8 the dimensions and elastic properties of femoro-distal saphenous vein grafts for obstructive arterial disease were studied with time. The goal of this study was to investigate possible structural and functional adaptive responses of the saphenous vein, when grafted into the arterial circulation. The main finding of this study was that the elastic properties of the vein graft are stable within the first 6 months after grafting. A slight increase in the diameter of the proximal graft segments was observed. Since in this segment the highest flow velocities are present, the increase in diameter may be a flow mediated adaptive response in order to decrease the high level of shear rate at this point.

MECHANICAL INFLUENCES ON THE OCCURRENCE OF INTIMAL HYPERPLASIA IN PERIPHERAL BYPASS GRAFTS AND GRAFT AV FISTULAS IN HUMANS

In chapter 9, the role of peri-anastomotic mismatch in elastic properties and flow conditions on the occurrence of stenoses in graft AV fistulas was reported. As stated in chapter 7, one of the hypotheses was that the sudden increase in compliance at the venous anastomosis would cause flow disturbances indicating low shear rate conditions, thereby leading to increased intimal thickening. However, the prospective analysis described in chapter 9, showed that this mechanism does not play a role in the development of intimal thickening in graft AV fistulas. When correlating initial flow characteristics and peri-anastomotic mismatch in elastic properties with the incidence of intimal thickening at the venous anastomosis, we found that the occurrence of intimal hyperplasia was not associated with mismatch in elastic properties. On the contrary, an even better initial match in elastic properties was observed in AV fistulas developing an intimal stenosis as compared to the AV fistulas that did not develop a stenosis. Another major finding in this prospective study was that local initial high shear rates at the site of the venous anastomosis or the efferent vein is associated with the occurrence of intimal thickening leading to stenosis. Histologic examination of the harvested stenotic tissue specimens revealed that the average coverage of the lumen with endothelial cells was approximately 50%, indicating partial denudation of endothelial cells. The finding that denudation is present at the site of the stenosis suggests that high shear rate at the venous site leads to injury and erosion of endothelial cells, possibly initiating the process of intimal thickening.

In chapter 10 the influence of mechanical factors on the occurrence of intimal hyperplasia in saphenous vein peripheral bypass grafts was studied prospectively. The proximal anastomosis of the graft was predisposed to the development of stenoses. In the grafts developing a stenosis at the proximal anastomosis during follow-up, a higher initial mismatch in strain around the anastomosis was observed than in grafts free of stenosis during follow-up. In addition, the initial diameter mismatch at the proximal anastomosis was larger in the grafts developing a stenosis.

Multiple regression analysis showed that a high initial peak shear rate at the proximal graft segment was associated with the occurrence of stenoses. In addition, in the grafts developing a stenosis, the highest initial flow velocities and peak shear rate levels were observed at the site of the later stenosis. This suggests that intimal thickening in venous grafts is associated with high shear rate rather than low shear rate. This observation is in concordance with our findings in AV fistulas (chapter 9). In the stenotic segments harvested from patients needing surgical intervention denudation was found in approximately the same extent as in the stenotic segments harvested from AV fistulas. This indicates that high shear rates also may play a role in the initiation of intimal hyperplasia in saphenous vein peripheral bypass grafts. This supports the concept that in venous segments excessive high shear rates are not very well tolerated and may lead to endothelial cell injury and denudation which in turn causes a response of the vessel wall initiating intimal thickening.

In chapter 11 the results of the assessment of cellular proliferation in stenotic segments derived from human AV fistulas and peripheral bypass grafts are reported. This is the first prospective study correlating initial flow conditions with the extent of proliferation in human stenoses. In addition, the distribution of proliferation in the stenotic segments was determined, since anatomically intact material was studied. The major finding of this study was that the extent of subendothelial proliferation was positively correlated with the percentage coverage of the lumen by endothelial cells, suggesting that the presence of endothelial cells in the stenotic lesion leads to promotion of proliferation rather than inhibition. This suggests that the endothelial cells covering the stenotic segments are dysfunctional and are unable to control cell proliferation in the vessel wall. A positive correlation was observed between proliferation of endothelial cells covering the lumen and subendothelial-intimal proliferation, suggesting that a proliferative state of the endothelial cells is associated with an imbalance between growth inhibition and promotion in favour of the latter. Peak shear rate was negatively correlated with endothelial cell coverage of the lumen, supporting the concept that an excessive high shear rate leads to endothelial cell injury, and even to denudation. Thus, high shear may have an indirect inhibitory effect on proliferation by means of denudation.

SAMENVATTING

DE INVLOED VAN STROMINGSKARAKTERISTIEKEN (FLOW) EN EEN PLOTSELINGE VERANDERING IN ELASTISCHE EIGENSCHAPPEN (MISMATCH) OP HET ONTSTAAN VAN INTIMA HYPERPLASIE IN PERIFERE BYPASSES EN ARTERIO-VENEUZE FISTELS VOOR HEMODIALYSE

In de hoofdstuk 1 en 2 zijn de achtergronden en vraagstellingen van het proefschrift beschreven. Het belangrijkste doel was de rol van mechanische factoren op het ontstaan van intima hyperplasie te onderzoeken bij patienten met een perifere arteriële bypass in het been of een arterio-veneuze fistel voor hemodialyse. De hypothese hierbij was dat door inhomogeniteit in elastische eigenschappen of door plaatselijke bloedstroom condities intima hyperplasie kan worden gestimuleerd. De ontwikkeling van intima hyperplasie in deze arteriële vaatreconstructies leidt tot hemodynamisch significante stenosen in een groot deel van de patienten, hetgeen chirurgische interventie noodzakelijk maakt. Dit resulteert in substantiële morbiditeit en hospitalisatie bij deze patienten. Meer inzicht in de initiatie en ontwikkeling van intima hyperplasie zou kunnen bijdragen tot het ontwerpen van nieuwe chirurgische benaderingen om de mechanische en flow condities van de reconstructies te optimaliseren en dit zal kunnen leiden tot een vermindering van de incidentie van intima hyperplasie. Ook kan een beter inzicht in de pathofysiologische processen welke leiden tot intima verdikking bijdragen aan de ontwikkeling van farmacologische interventies, om de ontwikkeling van het proces te kunnen remmen.

VESSEL WALL DOPPLER TRACKING

Voor het meten van elastische eigenschappen van perifere bypasses en arterio-veneuze fistels ontbraken tot recent geschikte meetsystemen. Met de introductie van Vessel Wall Doppler Tracking (VWDT) is het mogelijk geworden om de dimensies en uitzetting van bloedvaten in vivo bij de mens te meten. Dit ultrageluids systeem combineert B-mode met off-line automatische detectie van de beweging van structuurovergangen, zoals de vaatwand. Hierdoor is het mogelijk om dimensies en elastische eigenschappen op gedefinieerde plaatsen in een vasculair systeem te meten. Op deze wijze kunnen de elastische eigenschappen rond de anastomosen van arteriële reconstructies worden bepaald. Deze informatie is essentieel om de rol van perianastomotische verschillen in elastische eigenschappen in het ontstaan van stenosering te onderzoeken. In hoofdstuk 3 is het VWDT systeem beschreven en wordt de mogelijke rol van het systeem binnen de het vaatonderzoek bediscussieerd.

In hoofdstuk 4 worden de korte termijn variaties in elastische eigenschappen van de arteria femoralis communis bij gezonde jonge vrijwilligers beschreven. De belangrijkste bevinding in het onderzoek is dat de elastische eigenschappen van deze grote musculaire arterie in een periode van omstreeks 30 minuten belangrijk kunnen veranderen. Deze variatie in elasticiteit was niet gerelateerd aan veranderingen in locale bloeddruk, hetgeen suggereert dat veranderingen in vaattonus verantwoordelijk zijn voor dit fenomeen.

MISMATCH IN ELASTISCHE EIGENSCHAPPEN ROND ANASTOMOSEN VAN PERIFERE BYPASSES EN ARTERIO-VENEUZE FISTELS

De belangrijkste functie van een vaattransplantaat (graft) is het transporteren van bloed volume met een zo gering mogelijk verlies van energie, om zo voldoende doorbloeding in de periferie te verzorgen. Deze transport functie wordt grotendeels bepaald door de mechanische eigenschappen van deze grafts. Om basale informatie over de mechanische eigenschappen te verkrijgen, werden de dimensies en de uitzetting van grafts en van het natieve vasculaire stelsel op gedefinieerde plaatsen gemeten met behulp van VWDT. Op deze wijze werden de eerste humane in vivo gegevens over mismatch in elastische eigenschappen rond de anastomosen van perifere bypasses en arterio-veneuze fistels verkregen.

In hoofdstuk 5 zijn de mechanische profielen van 'in situ' en omgekeerde veneuze bypasses in het been beschreven. In de in situ bypass werd een vloeiende overgang in compliantie rond de proximale anastomose gevonden, in tegenstelling tot de omgekeerde veneuze bypass, waar ter plaatse van de proximale anastomose een belangrijk verval in compliantie werd gemeten. Echter, van proximaal naar distaal in de graft was er een stabiel patroon in compliantie in de omgekeerde veneuze graft, terwijl de in situ bypass een belangrijk verval in compliantie liet zien.

In hoofdstuk 6 zijn de elastische eigenschappen van veneuze en prothetische (PTFE) bypass grafts met elkaar vergeleken. In vivo metingen bleken met het VWDT systeem goed mogelijk. Naar verwachting bleek de uitzetting van de veneuze bypass groter te zijn dan die van de PTFE bypass.

In hoofdstuk 7 zijn de verschillen in peri-anastomotische mismatch in elastische eigenschappen vergeleken tussen de arteriële en de veneuze anastomose van interpositie grafts voor hemodialyse. Het merendeel van de stenosen in deze AV fistels ontwikkelt zich in of rond de veneuze anastomose. Gevonden werd dat er in de arteriële anastomose een abrupte vermindering in compliantie bestaat, terwijl in de veneuze anastomose een abrupte toename in compliantie voorkomt. Een abrupte toename in compliantie is geassocieerd met veranderingen in stromingscondities, zoals stase en omkering van bloedstroom. Omdat in verschillende studies is aangetoond dat een trage flow is geassocieerd met een toename van intima hyperplasie, zou dit de verklaring

kunnen zijn voor de predispositie van de veneuze anastomose tot het ontwikkelen van intima verdikking.

In hoofdstuk 8 zijn de dimensies en elastische eigenschappen van veneuze bypasses als functie van de tijd bestudeerd (tot 6 maanden na aanleggen). Gedurende deze periode werd geen verandering in elastische eigenschappen gevonden, hetgeen suggereert dat de elastische eigenschappen van de veneuze graft bewaard blijven gedurende de eerste 6 maanden na de aanleg ervan.

DE INVLOED VAN MECHANISCHE EIGENSCHAPPEN OP HET ONTSTAAN VAN INTIMA HYPERPLASIE IN PERIFERE BYPASSES EN ARTERIO- VENEUZE FISTELS

In hoofdstuk 9 zijn op prospectieve wijze de invloeden van peri-anastomotische mismatch in elastische eigenschappen en van stromingscondities op het ontstaan van intima verdikking in arterio-veneuze hemodialyse fistels bij patienten bestudeerd. Hierbij werden de initiële mismatch in elastische eigenschappen en stromingsparameters (gemeten 2 weken na aanleggen) gecorreleerd met het optreden van stenosering gedurende de follow-up periode tot 2 jaar. Uit de analyse bleek dat mismatch in elastische eigenschappen rond de anastomosen niet geassocieerd is met het ontstaan van stenosering door intima hyperplasie. Tevens bleek dat een initiële hoge systolische bloedstroomsnelheid, gecorrigeerd voor diameter, geassocieerd is met het optreden van stenosering. Stenosen traden voor het overgrote deel op aan de veneuze zijde van de AV-fistels.

In hoofdstuk 10 werd een soortgelijke prospectieve analyse gedaan voor perifere bypasses in het arteriële systeem van het been. Een grote mismatch in relatieve rek en in einddiastolische diameter ter plaatse van de proximale anastomose van veneuze perifere bypasses bleek geassocieerd met het ontstaan van stenosering tijdens de nacontrole. Tevens bleek in de bypasses die een stenose ontwikkelden gedurende de follow-up periode de stenose te ontstaan op de plaats met de initieel hoogste piek bloedstroomsnelheid. Dit laatste bevestigt de resultaten uit hoofdstuk 9. De data suggereren derhalve dat hoge initiële bloedstroomsnelheden in veneuze segmenten zou kunnen leiden tot stenosering door intima hyperplasie.

Proliferatie van gladde spiercellen wordt als een belangrijke stap beschouwd in de ontwikkeling van intima hyperplasie. In hoofdstuk 11 is de cellulaire proliferatie in een aantal stenotische segmenten bepaald, welke werden verkregen uit de patientenpopulaties zoals beschreven in de hoofdstukken 9 en 10. Tevens werd de invloed van locale bloedstroomkarakteristieken op de cellulaire proliferatie bestudeerd. De gemiddelde proliferatie index van cellen in de stenose bleek zowel in de stenosen uit de bypass groep als uit de AV-fistel groep omstreeks 3.5% te zijn, hetgeen veel hoger is dan in controle vaatsegmenten (0-0.5%). Tevens werd een positieve correlatie gevonden

tussen de mate van endotheelcel bekleding van de stenose en de subintimale proliferatie, hetgeen suggereerd dat de aanwezige endotheelcellen proliferatie bevorderen. Een andere belangrijke bevinding was de grote mate van vaatnieuwvorming in de stenotische segmenten. Op dit moment is nog onduidelijk in welke mate deze vaatnieuwvorming bijdraagt aan de ontwikkeling van stenosering.

DANKWOORD

Niet in de laatste plaats maar toch meestal aan het einde van het van het dankwoord wordt het gezin van de promovendus bedankt. Erica, ondanks het feit dat veel heeft moeten wijken voor de promotie heb je me altijd volledig gesteund en heb je ons gezin gepromoveerd met 2 schatten van kinderen, waar ik dit boekje meteen voor zou inleveren.

Prof Dr PJEHM Kitslaar, beste Peter, gaarne wil ik je hartelijk bedanken voor de grote vrijheid en hulp die je me hebt gegeven voor het opzetten van onderzoek, ook al was het vaak naast de oorspronkelijke vraagstelling. Veel baat heb ik gehad bij je enorme nauwgezetheid en je gedrevenheid om het onderzoek te laten slagen. Onvergetelijk waren de ritten in de Peugeot 504 familiale, die helaas door claudicatio aan z'n einde is gekomen.

Prof Dr APG Hoeks, beste Arnold, aanvankelijk waren de denkstappen die je maakte voor mij onnavolgbaar en deed ik maar wat je me opdroeg. Later kwam ik toch wat dichter in de buurt van jouw uitzend frequentie, hetgeen geleid heeft tot zeer vruchtbare discussies over de resultaten van het onderzoek. Daarbij is de door jouw gretig aangehaalde kloof tussen clinici en basiswetenschappers hopelijk wat kleiner geworden.

Dr JHM Tordoir, beste Jan, tijdens de research periode heb ik altijd veel plezier beleefd aan het onderzoek met de AV fistels, het turbomodel voor de ontwikkeling intima hyperplasie, zoals het door jou treffend wordt genoemd. Wat je doet lijkt vaak laconiek, maar blijkt toch zeer toegewijd. Zowel binnen het onderzoek als erbuiten heb ik veel steun aan je gehad, ook wanneer het eens wat slechter ging.

Dr MJAP Damen, beste Mat, jouw expertise op het gebied van (re)-stenose heeft een belangrijke kwaliteit toegevoegd aan het onderzoek. Een zeer goede maar soms zeer lastige eigenschap van je is dat je altijd zeer kritisch blijft, waarbij je als een terrier aan je standpunten vasthoudt. Dat heeft me wel eens wat moedeloosheid bezorgd. Ondanks en ook dankzij die eigenschap zet ik de samenwerking in de toekomst graag met je voort.

JHM Willigers, beste Jean, high-tech onderzoek kan niet zonder high-tech ondersteuning. Bij calamiteiten op technisch gebied stond je altijd weer klaar, ook al was het soms slechts om een stekker in het juiste stopcontact te steken. Met een zeer speciaal gevoel kijk ik terug naar het fitten van sinusfuncties door onze data, het zogenaande 'snaken'. Veel dank voor deze Eysdense kwaliteiten.

P Brands, beste Peter, ook jij stond altijd klaar in geval van technische problemen, met name wanneer Jean het ook niet meer wist. Veel dank voor de onvoorwaardelijke support tijdens het onderzoek.

DCJJ Bergmans, super Dennis, een ieder kan ik zonen van vlaaienbakkers uit Weert aanbevelen (vlaaien graag goed vasthouden). Als student-assistent heb je geweldig veel werk verzet bij het verwerken van de data-stroom en daardoor zeker een jaar 'gewonnen' voor het onderzoek. Niet alleen je werklust werkte aanstekelijk, maar ook je gevoel voor humor. Daarom vind ik het jaar dat jij er was het beste jaar van de research tijd. Ooit zal ik je verslaan met squash.

H Kurvers, beste Harry, 'a young man in a hurry', altijd geef je vol gas (helaas ook in mijn auto). Ondanks het feit dat je niet direct betrokken was bij het onderzoek, heb je er toch veel aan bijgedragen, alleen al door je competitieve natuur. Naast goede discussies hebben we steeds zeer veel pret gemaakt tijdens het onderzoek. Zeer fraaie herinneringen bewaar ik aan onze ski-congressen en de door mij geschreven en aan jouw gerichte fake-liefdesbrieven van vermeende minnaressen uit het verre Amerika.

Frank Huvers, Nicolaas Schaper, Boy Houben en Karel Leunissen, zeer gewaardeerd en ook zeer waardevol is de internistische bijdrage geweest aan het onderzoek. Van de bekende water-vuur verhouding tussen de heelkunde en interne is mij gelukkig niets gebleken.

De dames van het vaatlab, Edith, Yvonne, Monique, Ien en Laura, bedankt voor de jullie steun tijdens het leren echoën en de gezelligheid, waarvan ik met name het jaarlijkse Sinterklaas vieren zeer heb gewaardeerd. Yvonne en Nicole, jullie wil ik graag bedanken voor de altijd attente ondersteuning tijdens de gehele research periode.

De Nederlandse Hartstichting wil ik graag bedanken voor de financiële steun voor projecten als deze. Ook veel dank voor de ondersteuning van de verschillende congresbezoeken, welke mijn wetenschappelijke horizon zeker verbreed hebben.

CURRICULUM VITAE

The author was born on June 25 1963 in Emmen, The Netherlands. After visiting primary school, he attended high school in Emmen (Christelijke Scholengemeenschap). In 1981 he entered Medical School at the University of Groningen; he graduated in 1989. From August 1989 to January 1991 he served as a medical officer in the Royal Dutch Army. From February 1991 to February 1994 he was a research fellow in the Vascular Laboratory, department of Surgery, and the department of Biophysics. During this period the work contained in this thesis was accomplished under the supervision of Prof PJEHM Kitslaar and Prof APG Hoeks. Part of the work was rewarded with European Society for Vascular Surgery Prize 1993, for the best original paper on clinical research and an Award lecture on the 12th Annual Meeting of the International Society of Blood Purification, 1994, Maastricht, The Netherlands. In September 1994 he started his training in Cardiology in the University Hospital of Maastricht (head: Prof HJJ Wellens).

LIST OF PUBLICATIONS

PUBLICATIONS RELATED TO THIS THESIS

Articles

L Hofstra, EJM Ermers, APM Langeveldt, APG Hoeks, PJEHM Kitslaar. Vessel Wall Doppler Tracking: A new technique to assess vessel wall compliance. *J Vasc Tech* 1993; 17:65-68.

L Hofstra, JM Willigers, ADM Kester, PJEHM Kitslaar, APG Hoeks. Short-term variations in elastic properties of a muscular artery in humans. *Clin Science* 1994; 86: 567-574.

L Hofstra, DCJJ Bergmans, APG Hoeks, JHM Tordoir, PJEHM Kitslaar. Compliance and distensibility of the reversed saphenous vein graft. *VASA* 1993; 42: 40-41.

L Hofstra, DCJJ Bergmans, APG Hoeks, JHM Tordoir, PJEHM Kitslaar. Assessment of inhomogeneities in elastic properties of in situ and reversed saphenous vein grafts in humans. *Eur J Vasc Surg* 1994; 8: 670-676.

L Hofstra, DCJJ Bergmans, APG Hoeks, PJEHM Kitslaar, KML Leunissen, JHM Tordoir. Mismatch in elastic properties around anastomoses of interposition grafts for hemodialysis access. *JASN* 1994; 5: 1243-1250.

L Hofstra, JHM Tordoir, PJEHM Kitslaar, APG Hoeks, MJAP Daemen. The influence of peak shear rate on proliferation patterns in human stenotic lesions derived from arterio-venous fistulas and peripheral bypass grafts - A study using KI-67 (submitted).

L Hofstra, DCJJ Bergmans, JHM Tordoir, APG Hoeks, PJEHM Kitslaar. Elastic properties of PTFE and saphenous vein bypass grafts in humans (submitted).

PJ Brands, APG Hoeks, L Hofstra, RS Reneman. A non-invasive method to estimate wall shear rate using ultrasound. *Ultrasound Med Biol* . 1995; 21:171-185.

JHM Tordoir, L Hofstra, KML Leunissen, PJEHM Kitslaar. Early experience with stretch polytetrafluoroethylene grafts for hemodialysis access surgery: results of a prospective randomized study (submitted).

L Hofstra, DCJJ Bergmans, PJEHM Kitslaar, KML Leunissen, MJAP Daemen, JHM Tordoir. The effect of shear rate and anastomotic mismatch in elastic properties on the development of stenosis in AV fistulas (submitted).

Abstracts

L Hofstra, PJEHM Kitslaar, APG Hoeks, KML Leunissen, JHM Tordoir. Vaatwand compliance in PTFE en veneuze grafts bij hemodialyse patienten. Voordracht chirurgendagen, Rotterdam, mei 1992.

L Hofstra, PJEHM Kitslaar, APG Hoeks, JHM Tordoir. Compliance in PTFE and saphenous vein grafts in hemodialysis patients. 16 th World Congress of the International Union of Angiology, Paris, september 1992.

L Hofstra, PJEHM Kitslaar, APG Hoeks. Cyclic variation in compliance in the common femoral artery. 16 th World Congress of the International Union of Angiology, Paris, september 1992.

L Hofstra, APG Hoeks, JHM Tordoir, PJEHM Kitslaar. Compliance en distensibiliteit in de omgekeerde vene graft. Voordracht Ned Vereniging voor Vaatchirurgie, Utrecht, oktober 1992.

F Huvers, L Hofstra, C Lange, A Hoeks, A Houben, P Kitslaar, P de Leeuw, A Krüseman, N Schaper. Carotid and femoral distensibility in uncomplicated non-insulin-dependent diabetes mellitus patients without atherosclerotic changes. *Neth J Med* 1993; 22: A91

F Huvers, L Hofstra, C Lange, A Hoeks, A Houben, P Kitslaar, P de Leeuw, A Krüseman, N Schaper. Venous distensibility and the venodilatory effect of nitroglycerin are not reduced in uncomplicated non-insulin-dependent diabetes mellitus. *Eur J Clin Invest* 1993; 23: A7.

F Huvers, L Hofstra, C Lange, A Hoeks, A Houben, P Kitslaar, P de Leeuw, A Krüseman, N Schaper. Distensibility of the arterial and venous system in non-insulin-dependent diabetes mellitus. Voordracht Ned Vereniging voor Diabetes Onderzoek, mei 1993.

L Hofstra, DCJJ Bergmans, PJEHM Kitslaar, APG Hoeks, JHM Tordoir. Compliance in PTFE and saphenous vein grafts in hemodialysis patients. 4th Vascular symposium, Züri, Austria, februari 1993.

L Hofstra, P Schiffrers, JHM Tordoir, MJAP Daemen, JGR DeMey, PJEHM Kitslaar. Pre-existente afwijkingen in de vena saphena magna: oorzaak voor het vroeg falen van de perifere bypass. Voordracht chirurgendagen, Amsterdam, mei 1993.

L Hofstra, DCJJ Bergmans, APG Hoeks, JHM Tordoir, PJEHM Kitslaar. Compliance and distensibility of the reversed saphenous vein graft. Voordracht Deutschen Gesellschaft für Gefäßchirurgie, Heidelberg, mei 1993.

L Hofstra, DCJJ Bergmans, APG Hoeks, JHM Tordoir, PJEHM Kitslaar. Elastic properties of the saphenous vein graft: A comparison of the in situ and reversed technique in humans. VII th Annual Meeting of the European Society for Vascular Surgery, Barcelona, september 1993.

JHM Tordoir, L Hofstra. Klinische resultaten en in vivo compliance metingen van stretchable PTFE AV protheses. Symposium Nieuwe Technieken en Materialen voor Hemodialyse Toegang en CAPD Behandeling, Utrecht, april 1993.

L Hofstra, DCJJ Bergmans, APG Hoeks, JHM Tordoir, PJEHM Kitslaar. Elastische eigenschappen van veneuze grafts; vergelijking van de in situ en omgekeerde vene graft bij de mens. SEOHS, Nijmegen, november 1993.

DCJJ Bergmans, L Hofstra, APG Hoeks, JHM Tordoir, PJEHM Kitslaar. Compliance in PTFE en veneuze femoro-distale bypasses. Ned Tijdschr Geneesk 1994, 21; 138 (21).

L Hofstra, APG Hoeks, KML Leunissen, DCJJ Bergmans, PJEHM Kitslaar, JHM Tordoir. Mismatch in elastic properties around anastomoses of interposition grafts for hemodialysis access. A Symposium on Hemodialysis Access IV, Orlando, mei 1994.

JHM Tordoir, L Hofstra, DCJJ Bergmans, APG Hoeks, KML Leunissen, PJEHM Kitslaar. Stretch PTFE grafts for hemodialysis access: improved hemodynamic performance and less complications compared to standard ePTFE grafts. A Symposium on Hemodialysis Access IV, Orlando, mei 1994.

L Hofstra, PJEHM Kitslaar, APG Hoeks, JHM Tordoir, MJAP Daemen. Correlation of proliferation and shear rates in human stenotic lesions derived from AV fistulas and femorodistal bypasses. Symposium on Intimal Hyperplasia, Maastricht, mei 1994.

PJEHM Kitslaar, L Hofstra, JHM Tordoir, APG Hoeks, MJAP Daemen. Het effect van peak shear rate op celproliferatie in stenosen in AV dialyse fistels en perifere arteriele bypasses. Chirurgendagen, Veldhoven 1994.

JHM Tordoir, L Hofstra, DCJJ Bergmans, APG Hoeks, KML Leunissen, PJEHM Kitslaar. Stretch PTFE protheses voor hemodialyse toegangschirurgie: verbetering in hemodynamische eigenschappen en vermindering van complicaties in vergelijking tot standaard ePTFE grafts. Chirurgendagen, Veldhoven 1994.

JHM Tordoir, L Hofstra, KML Leunissen. Duplex scanning and compliance measurement of graft AV fistulas. Simposio de Actualizacao na Cirurgia dos Acessos Vasculares Parahemodialise, Lisabon, juli 1994.

L Hofstra, JHM Tordoir, APG Hoeks, PJEHM Kitslaar, MJAP Daemen. The effect of peak shear rate on proliferation patterns in human stenotic lesions derived from AV fistulas and peripheral bypass grafts. VIIth International Symposium on Vascular Biology, Heidelberg, august 1994.

OTHER PUBLICATIONS

Articles

L Hofstra, E Boersma, SJ Ham. Penetrerende letsels van het abdomen op het eiland Curaçao. Ned Tijdschr Geneesk 1990; 134: 1046-48.

L Hofstra, HH van Woerden en R Deutman. Chemonucleolysis in the Herniated L3-L4 Disk. Clin Orthop 1991; 269:151-156.

H de Swart, L Dijkman, L Hofstra, FW Bär, V van Ommen, J Tordoir, HJJ Wellens. A new hemostatic puncture closure device for the immediate sealing of arterial puncture sites. Am J Cardiol 1993;72:445-449.

MH Booster, RMH Wijnen, JP van Hooff, ATMG Tiebosch, HG Peltenburg, PM van den Berg-Loonen, MJPG van Kroonenburgh, T Verschuren, L Hofstra and G Kootstra. The role of the spleen in pancreas transplantation. Transplantation (in press).

Abstracts

HAJM Kurvers, L Hofstra, APG Hoeks, FAJM van de Wildenberg, PJEHM Kitslaar, MJHM Jacobs. Sympathetic dysfunction in Sudeck's dystrophy assessed by Vessel Wall Echo Tracking. Eur J Clin Inv 1992; 22: A8.

L Hofstra, HAJM Kurvers, FAJM van de Wildenberg, MJHM Jacobs, PJEHM Kitslaar. Veranderingen van vaatwandgedrag bij Sudeck-dystrofie. Ned Tijdschr Geneesk 1992; 136: 1428.

H de Swart, L Dijkman, L Hofstra, F Bär, V van Ommen, J Tordoir, H Wellens. A new hemostatic puncture closure device for the immediate sealing of arterial puncture sites. XIV th Congress of the European Society of Cardiology, Barcelona, september 1992.

L Hofstra, HAJM Kurvers, APG Hoeks, FAJM van de Wildenberg, MJHM Jacobs PJEHM Kitslaar. The influence of sympathetic dysfunction on compliance of the brachial artery in Sudeck's dystrophy. 16th World Congress of the International Union of Angiology, Paris, september 1992.

HAJM Kurvers, L Hofstra, FAJM van den Wildenberg, PJEHM Kitslaar, MJHM Jacobs. Bilateral sympathetic involvement in Sudeck's dystrophy assessed by Vessel Wall Doppler Tracking. 16th World Congress of the International Union of Angiology, Paris, september 1992.

L Hofstra, L Dijkman, H de Swart, F Bär, V van Ommen. Resultaten van een nieuwe hemostatische vaatplug bij A. Femoralis-catheterisatie. Ned Tijdschr Geneesk 1993; 137: 627-628.

L Hofstra, HAJM Kurvers, JPL van der Plas, JHM Tordoir, MJHM Jacobs, BK Janevski, PJEHM Kitslaar. Klinische presentatie en diagnostiek van handarterieletsels door stompe traumata. Ned Tijdschr Geneesk 1993; 137: 628.

H de Swart, L Dijkman, L Hofstra, F Bär, V van Ommen, J Tordoir, H Wellens. A new hemostatic puncture closure device for the immediate sealing of arterial puncture sites. Neth J Card 1992; 5:73.

G van Ommen, J Tordoir, F van der Veen, L Hofstra, D Koster, H Wellens. A new hydrodynamic thrombectomy catheter for the removal of intravascular thrombus: First clinical results. XVth Congress of the European Society of Cardiology, Nice, september 1993.

PJ Brands, APG Hoeks, L Hofstra. Een niet-invasieve methode voor shear-rate meting. Voordracht Nederlandse Vereniging Ultrageluid, Utrecht, Oktober 1993.

L Hofstra, P Schiffers, JHM Tordoir, MJAP Daemen, JGR De Mey, PJEHM Kitslaar. Pre-existente afwijkingen in de vena saphena magna; oorzaak voor het vroeg falen van de perifere bypass? Chirurgedagen, Amsterdam, mei 1993.

JHM Tordoir, V van Ommen, EH van der Veen, L Hofstra, D Koster. Eerste resultaten van een nieuwe hydrodynamische trombectomie catheter. Chirurgedagen, Amsterdam, mei 1993.

Het verschijnen van dit proefschrift werd financieel ondersteund door:

W.L. Gore and Associates B.V.

LEO

Zambon

Servier

Wyeth

Pie Medical

Van der Laar Stichting