

Exposure to secondhand aerosol of electronic cigarettes in indoor settings in 12 European countries: data from the TackSHS survey

Citation for published version (APA):

Amalia, B., Liu, X. Q., Lugo, A., Fu, M., Odone, A., van den Brandt, P. A., Semple, S., Clancy, L., Soriano, J. B., Fernandez, E., Gallus, S., & TackSHS Project Investigators (2021). Exposure to secondhand aerosol of electronic cigarettes in indoor settings in 12 European countries: data from the TackSHS survey. *Tobacco Control*, 30(1), 49-56. <https://doi.org/10.1136/tobaccocontrol-2019-055376>

Document status and date:

Published: 01/01/2021

DOI:

[10.1136/tobaccocontrol-2019-055376](https://doi.org/10.1136/tobaccocontrol-2019-055376)

Document Version:

Publisher's PDF, also known as Version of record

Document license:

Taverne

Please check the document version of this publication:

- A submitted manuscript is the version of the article upon submission and before peer-review. There can be important differences between the submitted version and the official published version of record. People interested in the research are advised to contact the author for the final version of the publication, or visit the DOI to the publisher's website.
- The final author version and the galley proof are versions of the publication after peer review.
- The final published version features the final layout of the paper including the volume, issue and page numbers.

[Link to publication](#)

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal.

If the publication is distributed under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license above, please follow below link for the End User Agreement:

www.umlib.nl/taverne-license

Take down policy

If you believe that this document breaches copyright please contact us at:

repository@maastrichtuniversity.nl

providing details and we will investigate your claim.

Exposure to secondhand aerosol of electronic cigarettes in indoor settings in 12 European countries: data from the TackSHS survey

Beladenta Amalia ,^{1,2,3,4} Xiaoqiu Liu ,⁵ Alessandra Lugo ,⁵ Marcela Fu ,^{1,2,3,4} Anna Odone ,⁶ Piet A van den Brandt ,⁷ Sean Semple ,⁸ Luke Clancy ,⁹ Joan B Soriano ,^{4,10} Esteve Fernández ,^{1,2,3,4} Silvano Gallus ,⁵ the TackSHS Project Investigators

► Additional material is published online only. To view please visit the journal online (<http://dx.doi.org/10.1136/tobaccocontrol-2019-055376>).

For numbered affiliations see end of article.

Correspondence to

Esteve Fernández, Tobacco Control Unit, Catalan Institute of Oncology (ICO), 08908 L'Hospitell de Llobregat, Catalonia, Spain; efernandez@iconologia.net

Received 9 September 2019
Revised 12 December 2019
Accepted 31 December 2019
Published Online First
2 March 2020

ABSTRACT

Introduction Exposure to secondhand aerosol from e-cigarette (SHA) may pose harmful effects to bystanders. This study aims to investigate the prevalence, duration and determinants of SHA exposure in various indoor settings in 12 European countries.

Methods In 2017–2018, we conducted a cross-sectional study, the TackSHS survey, on a representative sample of the population aged ≥ 15 years in 12 European countries (Bulgaria, England, France, Germany, Greece, Ireland, Italy, Latvia, Poland, Portugal, Romania and Spain). We described the prevalence and duration of exposure to SHA in several indoor settings among 11 604 e-cigarette non-users. Individual-level and country-level characteristics associated with SHA exposure were also explored using multilevel logistic regression analyses.

Results Overall, 16.0% of e-cigarette non-users were exposed to SHA in any indoor setting at least weekly, ranging from 4.3% in Spain to 29.6% in England. The median duration of SHA exposure among those who were exposed was 43 min/day. 'Other indoor settings' (eg, bar and restaurant) was reported as the place where most of e-cigarette non-users were exposed (8.3%), followed by workplace/educational venues (6.4%), home (5.8%), public transportation (3.5%) and private transportation (2.7%). SHA exposure was more likely to occur in certain groups of non-users: men, younger age groups, those with higher level of education, e-cigarette past users, current smokers, those perceiving SHA harmless and living in countries with a higher e-cigarette use prevalence.

Conclusions We found inequalities of SHA exposure across and within European countries. Governments should consider extending their tobacco smoke-free legislation to e-cigarettes to protect bystanders, particularly vulnerable populations such as young people.

Trial registration number NCT02928536.

INTRODUCTION

Electronic cigarette (e-cigarette) use has increased in many parts of the world. In the USA, with Juul's extraordinary growth and marketing strategy, e-cigarette use has been declared as an epidemic in youth by the US Surgeon General as it substantially increased by 78% from 2017 to 2018.^{1,2} According to the Eurobarometer surveys, the prevalence of adults who had at least tried e-cigarettes in 28

European countries has grown from 12% in 2014 to 15% in 2017.³

The growing use of e-cigarettes has raised concerns as the product is potentially harmful both to users and bystanders.^{4,5} While some studies showed that e-cigarettes emit lower levels of some toxic chemicals compared with smoke from conventional cigarettes, other studies revealed that e-cigarette aerosol contains comparable or higher levels of other harmful constituents, such as nicotine and metals.^{6–9} It has been also shown that bystanders absorb nicotine from e-cigarette aerosol at levels comparable with secondhand tobacco smoke (SHS).¹⁰ Additionally, e-cigarette aerosol may expose non-users to toxic chemicals, including particulate matter and carcinogens, such as volatile organic compounds, polycyclic aromatic hydrocarbons, formaldehyde, acetaldehyde and tobacco-specific nitrosamines.^{11–14} Secondhand aerosol (SHA) from e-cigarettes has been found to cause acute reduced lung function and associated with higher odds of asthma exacerbations, which might reflect more adverse health effects with longer period of exposure.^{15,16} Exposure to SHA from e-cigarette may renormalise tobacco smoking, induce relapse to smoking for those who have quit smoking and trigger initiation of e-cigarette use among non-smokers, particularly young people.^{17–21} The above evidence suggests that appropriate regulations are needed to prevent involuntary exposure to SHA.

The WHO recommends to Parties of the Framework Convention on Tobacco Control to consider the prohibition of e-cigarette use in indoor settings or at least in those places where smoking is already banned.²² In Europe, e-cigarette use has been frequently observed in indoor places where smoking is normally banned, such as workplaces, bars, restaurants and train and metro stations.^{23–25} Evading smoke-free regulation has been reported by e-cigarette users as one of the main reasons for the use of e-cigarettes.^{26–28} To the best of our knowledge, to date there have been 28 European countries regulating the use of e-cigarettes, but mostly in selected public places only.²⁹

While public debate about the risks and benefits of e-cigarette use continues to arise, evidence on the extent of the population's exposure to the SHA has been documented.³⁰ According to the 2015



© Author(s) (or their employer(s)) 2021. No commercial re-use. See rights and permissions. Published by BMJ.

To cite: Amalia B, Liu X, Lugo A, et al. *Tob Control* 2021;**30**:49–56.

National Youth Tobacco Survey data, exposure to SHA in indoor or outdoor public places was reported by one in four middle-school and high-school students in the USA, including 4.4 million who were e-cigarette non-users and one million not exposed to SHS.³¹ Recent data from six European countries indicated that 37% of smokers (e-cigarette non-users) were exposed to SHA, ranging from 18% in Spain to 63% in Greece.²³ However, there has been no study on exposure to SHA from e-cigarettes among the general population in Europe.

This paper aims to assess the prevalence and duration of exposure to SHA from e-cigarettes in various indoor settings among e-cigarette non-users aged 15 years or older in 12 European countries. We also explored the sociodemographic factors at the individual and country level that were associated with SHA exposure.

METHODS

Data source

This is a questionnaire-based cross-sectional study using data from the TackSHS survey, conducted in 12 selected European countries (Bulgaria, England, France, Germany, Greece, Ireland, Italy, Latvia, Poland, Portugal, Romania and Spain). The detailed methods of the TackSHS survey, including the questionnaire development, have been explained elsewhere.^{32 33} Sampling methods varied across countries, including multistage sampling (in Bulgaria, Greece, Italy, Latvia, Poland and Romania), cluster sampling with quotas (in England and France) and stratified random sampling (in Germany, Ireland, Portugal and Spain). In each country, we sampled around 1000 people representative of the general population in terms of age, sex, geographical area, and in most of countries, socioeconomic characteristics. In total, the survey included 11902 subjects aged 15 years or older from 12 European countries, representing 79.2% of the whole EU population. A pilot study was conducted in Italy in November 2016 while the fieldwork in other countries was conducted between June 2017 (in Romania) and October 2018 (in Latvia), using the same questionnaire administered with computer-assisted personal interviewing in all 12 countries. The questionnaire included information on socioeconomic and demographic characteristics, cigarette smoking, e-cigarette use, SHS and SHA exposures in various indoor and outdoor settings, and attitudes and perception towards SHS and SHA exposures.³³

For the purpose of this study, only e-cigarette non-users were included. Thus, the total sample size in this study was 11604 subjects.

Measures

Respondents who reported that they had never used e-cigarette during their lifetime or had stopped using it at least for 30 days before the time of the survey were considered as e-cigarette non-users (ie, never and ex-users). From a question 'On average, how much time per day do you think you are exposed to e-cigarette aerosol in each of the following sites?', interviewees indicated one or more of the indoor settings where they experienced SHA exposure. Five indoor settings were considered: home, workplace (or educational venues for students), public transportation (eg, train, tram, bus and subway), private transportation and 'other indoor places' (eg, cafeterias, bars, restaurants and leisure facilities). For each indoor setting, e-cigarette non-users reported the average exposure time (in min/day) during a working and non-working day. An e-cigarette non-user was defined as exposed to SHA in a certain setting, if (s)he was exposed in that setting at least 1 min per day in a working and/or non-working day.

The prevalence (%) of exposure (at least weekly) was computed for each setting and overall. Duration of SHA exposure was computed as the weighted daily average minutes of exposure in working and non-working days among subjects exposed to SHA in each setting.

Ever smokers were defined as respondents who reported smoking at least 100 cigarettes (including hand-rolled cigarettes) during their lifetime. Among ever smokers, current smokers were participants who reported current smoking at the time they participated in this survey, while ex-smokers were those who had stopped smoking by the time they participated in this survey.³⁴

Information on harm perception from SHA exposure was obtained by asking respondents 'Do you agree or disagree with the following sentence? Exposure to e-cigarette vapour is harmful to my health, with five possible answer options: (1) strongly agree; (2) moderately agree; (3) moderately disagree; (4) strongly disagree; (5) does not know or does not answer'. Options 1 and 2 were categorised as 'harmful', whereas options 3 and 4 were categorised as 'harmless'.

Level of education was constructed by taking country-specific tertiles of schooling years as low, intermediate and high. The 12 countries were classified by their geographical area into Northern Europe, Western Europe, Southern Europe and Eastern Europe according to United Nations M49 Standard,³⁵ by the World Bank GDP per capita,³⁶ by their score in the 2016 Tobacco Control Scale,³⁷ by country's smoking prevalence and by country's e-cigarette use prevalence. The latter two were estimated from the TackSHS survey data.

Statistical analysis

We reported proportion, and median estimates of the SHA exposure among e-cigarette non-users across countries and sociodemographic subpopulations. We used the median of the minutes exposed as point of estimates for the duration of SHA due to extremely right-skewed distribution of the data.

A multilevel logistic regression model, allowing for clustering of observations at the country level was fitted to examine the relationship between SHA exposure status (as a binary dependent variable) and sociodemographic characteristics at individual and country levels (independent variables). Adjusted OR (aOR) and their corresponding 95% CI were estimated after adjusting for sex, age, level of education, e-cigarette use status and smoking status.

Statistical weights were used to generate representative estimates of the general population of each country (individual weight). To calculate results for the entire sample, we applied 'country weights', which combined individual weights with an additional weighting factor, with each country contributing in proportion to its population aged 15 years or over, obtained by Eurostat.³⁸ Analyses were performed with STATA V.14.0.

RESULTS

The sample sociodemographic characteristics are presented in online supplementary table 1. Among 11604 e-cigarette non-users, 16.0% (95% CI: 15.3% to 16.7%) were exposed at least weekly to SHA from e-cigarettes in any indoor setting, and ranged from 4.3% (95% CI: 3.2 to 5.7) in Spain to 29.6% (95% CI: 26.7 to 32.6) in England, with significant differences among men and women (17.2% vs 15.0%, $p < 0.001$) for the 12 countries combined (table 1). The highest prevalence of at least weekly SHA exposure was reported in England for both men and women (31.8% and 27.8%, respectively). Overall, the median duration of SHA exposure for e-cigarette non-users who had

Table 1 Country-specific prevalence (%) of e-cigarette secondhand aerosol (SHA) exposure (at least weekly), overall and by sex in e-cigarette non-users of the European population aged ≥ 15 years.* TackSHS survey, 2017–2018

Country	N†	% exposure to SHA from e-cigarettes (95% CI)		
		Total	Men	Women
Bulgaria	1035	14.9 (12.8 to 17.2)	14.3 (11.4 to 17.6)	15.4 (12.6 to 18.7)
England	940	29.6 (26.7 to 32.6)	31.8 (27.5 to 36.3)	27.8 (24.1 to 31.8)
France	974	26.3 (23.6 to 29.1)	27.5 (23.6 to 31.7)	25.2 (21.6 to 29.2)
Germany	1000	11.1 (9.2 to 13.2)	12.6 (10.0 to 15.9)	9.5 (7.3 to 12.4)
Greece	959	28.9 (26.1 to 31.8)	30.3 (26.4 to 34.6)	27.4 (23.6 to 31.6)
Ireland	916	22.1 (19.6 to 24.9)	24.6 (20.8 to 28.8)	19.8 (16.4 to 23.7)
Italy	1045	12.8 (10.9 to 15.0)	15.7 (12.8 to 19.2)	10.1 (7.9 to 13.0)
Latvia	1009	5.6 (4.4 to 7.2)	6.7 (4.8 to 9.3)	4.7 (3.2 to 6.8)
Poland	718	12.3 (10.1 to 14.9)	13.9 (10.5 to 18.1)	11.0 (8.2 to 14.5)
Portugal	991	11.4 (9.6 to 13.5)	12.5 (9.8 to 15.8)	10.5 (8.1 to 13.4)
Romania	999	10.0 (8.3 to 12.0)	10.9 (8.5 to 14.0)	9.2 (6.9 to 12.0)
Spain	1018	4.3 (3.2 to 5.7)	2.2 (1.2 to 3.9)	6.3 (4.5 to 8.7)
Total	11 604	16.0 (15.3 to 16.7)	17.2 (16.2 to 18.2)	15.0 (14.1 to 15.9)

* Individual-level weight was applied to all estimates. For total estimates of the entire sample, country-level weight was applied with each country contributing in proportion to its population aged 15 years or over.³⁸

† Sample size (N) is the unweighted country-specific number of e-cigarette non-users.

been exposed to SHA was 43 min/day (Q1–Q3: 14–130). The duration of SHA exposure ranged from 2 min/day (Q1–Q3: 1–7) in Spain to 103 min/day (Q1–Q3: 21–240) in Italy (figure 1).

Table 2 shows the country-specific prevalence and duration of SHA exposure in various indoor settings. SHA exposure among e-cigarette non-users mostly occurred in ‘other indoor settings’ (8.3%), followed by workplace/educational venues (6.4%), home (5.8%), public transportation (3.5%) and private transportation (2.7%). France had the highest prevalence of SHA exposure at home (12.0%), workplace/educational venues (13.2%) and private vehicles (5.9%) compared with other countries, while the highest prevalence of SHA exposure in public transportation was in England (7.9%) and in ‘other indoor settings’ in Greece (19.0%). The longest median duration of SHA exposure was 43 min/day taking place at home and workplace, while the




Figure 1 Median (Q1–Q3) duration of exposure to SHA (minutes/day) from e-cigarettes. TackSHS survey, 2017–2018. Median estimates were calculated among e-cigarette non-users who had been exposed to SHA at any indoor setting. Q1, first quartile; Q3, third quartile.

shortest one was in public transportation with a median exposure of 14 min/day. Despite the low prevalence of SHA exposure (1.8%) among Latvian e-cigarette non-users in ‘other indoor places’, they reported a 2-hour-per-day of SHA exposure in these venues.

Table 3 shows the proportion of SHA exposure and the corresponding aOR according to selected individual-level characteristics. At least weekly SHA exposure was more frequent in men (aOR: 1.13; 95% CI: 1.01 to 1.25) than in women and in the young (aOR for < 25 vs ≥ 65 years: 3.15; 95% CI: 2.52 to 3.94; p for trend < 0.001). The higher the level of education, the more likely e-cigarette non-users were exposed to SHA (aORs for intermediate level of education: 1.19; 95% CI: 1.05 to 1.35, and for high-level of education: 1.26; 95% CI: 1.10 to 1.44; p for trend < 0.001). Higher odds of SHA exposure were related with being an e-cigarette past user (compared with never users aOR: 1.49; 95% CI: 1.14 to 1.95) and being a current smoker (compared with never smokers, aOR: 1.54; 95% CI: 1.36 to 1.74). Those who perceived SHA exposure as harmful were less likely to be exposed to SHA (vs harmless; aOR: 0.69; 95% CI: 0.61 to 0.78).

Compared with Northern Europe, SHA exposure was lower among e-cigarette non-users living in Southern (aOR: 0.27; 95% CI: 0.11 to 0.68) and Eastern Europe (aOR: 0.35; 95% CI: 0.13 to 0.94) (table 4). E-cigarette non-users living in countries with higher prevalence of e-cigarette use were more likely to be exposed to SHA (vs $< 1\%$ e-cigarette use prevalence; aOR for 1%–4% group: 1.64, 95% CI: 1.05 to 2.56; aOR for $> 4\%$ group: 4.35, 95% CI: 2.72 to 6.96; p for trend < 0.001).

DISCUSSION

Sixteen percent of e-cigarette non-users in 12 European countries were exposed to SHA at least weekly in any indoor setting, reporting a median of 43 min/day of exposure. Most of their exposure took place in ‘other indoor settings’ that include restaurants and bars, but, importantly, the exposure of longest duration occurred at home and workplace (43 min/day). It is also evident that variability in SHA exposure exists across countries and among different sociodemographic groups—men, the youngest, highly educated, past e-cigarette users, current smokers, those perceiving SHA as harmless and living in a country with high e-cigarette use prevalence were among individuals who were more likely to be exposed to SHA.

The highest prevalence of SHA exposure (more than one in four non-users, England) does not correspond to the longest duration of SHA exposure (103 min/day, Italy). The discrepancy might be partly due to lower time-sensitisation towards duration of SHA exposure among bystanders in countries where SHA exposure was more common; they perceived shorter duration of SHA exposure because they had already accustomed to it. However, the discrepancy highlights the importance of monitoring both measures, prevalence and duration of SHA exposure, in a population. There is no evidence on the safety levels of SHA exposure, while for SHS, there has been established evidence showing that there is no risk-free level of SHS.^{39–41} However, it has been shown that 2 hours/day of exposure to exhaled aerosol of e-cigarettes for a week may significantly increase urinary and salivary cotinine among bystanders living in homes with e-cigarette users.¹⁰ Another study also found that after an SHA exposure of 1 hour, the serum cotinine concentrations increased at similar levels as in subjects exposed to SHS.⁴² That indicates bystanders may systematically absorb the nicotine from acute exposure to SHA.

Table 2 Country-specific prevalence (%) and duration (minutes/day) of e-cigarette secondhand aerosol (SHA) exposure in selected indoor settings among e-cigarette non-users of the European population aged ≥15 years. * TackSHS survey, 2017–2018

Country	N†	Home		Workplace/educational venues		Public transportation		Private transportation		Other indoor places	
		%	Median‡ (min/day)	%	Median‡ (min/day)	%	Median‡ (min/day)	%	Median‡ (min/day)	%	Median‡ (min/day)
Bulgaria	1035	4.6	64	4.6	43	2.8	17	1.3	43	10.8	43
England	940	7.6	30	10.9	14	7.9	7	5.1	12	14.2	17
France	974	12.0	34	13.2	48	5.1	24	5.9	17	14.2	48
Germany	1000	2.3	34	2.8	43	2.6	30	1.4	27	8.0	26
Greece	959	8.1	60	10.8	46	3.4	43	1.6	60	19.0	60.0
Ireland	916	8.8	31	9.4	14	3.8	7	2.3	10	11.6	10
Italy	1045	5.6	60	6.3	43	3.3	60	3.0	60	5.1	60
Latvia	1009	2.1	60	2.4	43	0.3	21	0.3	14	1.8	120
Poland	718	6.6	69	4.8	21	2.9	14	0.9	19	3.7	33
Portugal	991	4.4	60	4.2	21	0.3	6	2.3	17	6.8	18
Romania	999	4.1	60	4.4	43	1.4	15	2.5	21	3.4	24
Spain	1018	1.5	10	0.5	4	0.9	1	0.0	0	1.9	2
Total	11 604	5.8	43	6.4	43	3.5	14	2.7	21	8.3	33

*Individual weight was applied to all estimates in each country. For total estimates of the entire sample, country weight are applied with each country contributing in proportion to its population aged 15 years or over.³⁸

†Sample size (N) is the unweighted, country-specific number of e-cigarette non-users.

‡Median estimates were calculated among e-cigarette non-users who had been exposed to SHA at the corresponding indoor setting.

A previous study, conducted among smokers in six European countries (Germany, Greece, Hungary, Poland, Romania and Spain) from June to September 2016, also identified differences in SHA exposure prevalence across countries, with Spain having the lowest exposure (18%) and Greece having the highest one (63%).²³ The variation of SHA exposure across countries may reflect a diverse country's e-cigarette use prevalence in Europe. Spain, for instance, was within the lowest e-cigarette use prevalence group (<1%) and had the lowest SHA exposure among others (4.3%). Indeed, the higher odds of SHA exposure in countries with higher e-cigarette use prevalence were evident from our regression analysis as we would expect, especially, if the use of the device is unregulated. The regression analysis revealed that country's e-cigarette use prevalence was an independent factor of SHA exposure among e-cigarette non-users, suggesting the need for countries to restrict the place of e-cigarette use. The policy for e-cigarette use restriction can be included in the country's current tobacco control strategy as, our study has shown, the current score of Tobacco Control Scale was still irrelevant to SHA exposure status. Moreover, a strong association found between SHA exposure and geographical area of the 12 countries might be attributable to the widespread 'vape-free' policy from one country to the neighbouring countries, as has been shown in the policy diffusion theory for local and national smoking ban regulations.^{43 44}

Similar to what has been described with SHS exposure, each country's regulatory environment may also affect the differences in SHA exposure among countries.^{45–48} Among the 12 countries included in this study, only Greece had introduced a 'vape-free' policy in all indoor settings by the time this study was conducted.²⁹ Despite the extensive coverage of 'vape-free' policy in Greece, non-users in the country were still markedly more exposed to SHA in indoor settings compared with other countries without any national 'vape-free' policy, like Bulgaria, Germany, Latvia and Romania.²⁹ In workplaces, including school and university, France, a country which already banned e-cigarette use in such settings, had the highest prevalence of SHA exposure.²⁹ This finding underscores the importance of

implementing and enforcing existing policies on e-cigarette use in indoor places. Most of the SHA exposure occurred in 'other indoor settings', which include bars and restaurants where smoking, but not e-cigarette use is prohibited in all the 12 countries examined.⁴⁹ A previous European study indicated a 20% prevalence of e-cigarette use in indoor places where smoking was banned.²³ The greater opportunity of using e-cigarette compared with smoking conventional cigarettes in enclosed spaces, including pubs, bars and restaurants, has been mentioned as one of the motivations of using e-cigarettes in such settings.^{27 50} That opportunity may encourage e-cigarette users, most of whom are dual users, to use e-cigarettes as an alternative to smoking in places where smoking is banned, as it is the case in 'other indoor settings'.^{3 27 50} Moreover, the already prevalent social norm of smoking in certain recreational facilities, including bars and restaurants, could also drive e-cigarette use in these settings.⁵¹ Thus, they are important factors to be considered in future public policies.

E-cigarette use in homes and private vehicles is a source of involuntary exposure to SHA for vulnerable populations, especially children. Despite the low prevalence of SHA exposure in homes shown in this study, an intense SHA exposure (43 min/day) occurred in such setting. In the UK, less than 10% of e-cigarette users forbid e-cigarette use in their homes, while a study in the USA indicates that about one in five e-cigarette users reported banning e-cigarette use inside their homes and cars.⁵² We also identified sociodemographic discrepancies in SHA exposure. Men, young, highly educated, current smokers and e-cigarette past users were more likely to be exposed to SHA in indoor settings. These determinants of SHA exposure were also true for smokers as has been shown in a study among six European countries.²³ Being in the youngest age groups or the higher educational level were also positive determinants for e-cigarette use and awareness about e-cigarettes.^{53–55} This peculiarity might be explained by the diffusion of innovation theory which states that early adopters of new behaviours tend to be males and those from higher socioeconomic status.⁵⁶ Accordingly, our data also found that SHA exposure was associated with highly educated

Table 3 Proportion (%) and adjusted OR (aOR) for at least weekly exposure to e-cigarette secondhand aerosol (SHA) and corresponding 95% CI according to selected individual-level characteristics among e-cigarette non-users of European population aged ≥ 15 years. * TackSHS survey, 2017–2018

Individual-level characteristics	N†	At least weekly exposed to SHA from e-cigarettes	
		%	aOR (95% CI)‡
Sex			
Women	6122	15.0	1§
Men	5482	17.2	1.13 (1.01 to 1.25)
Age group (years)			
<25	1401	20.9	3.15 (2.52 to 3.94)
25–44	3955	19.3	2.69 (2.20 to 3.30)
45–64	4218	16.4	2.23 (1.83 to 2.73)
≥ 65	2030	6.2	1§
<i>P for trend</i>			<0.001
Level of education¶			
Low	4381	13.4	1§
Intermediate	4064	17.5	1.19 (1.05 to 1.35)
High	3156	17.8	1.26 (1.10 to 1.44)
<i>P for trend</i>			0.001
E-cigarette use status			
Never user	11 299	15.6	1§
Past user	305	32.9	1.49 (1.14 to 1.95)
Smoking status			
Never smoker	6478	14.2	1§
Former smoker	1943	15.2	1.12 (0.96 to 1.31)
Current smoker	3183	20.9	1.54 (1.36 to 1.74)
Perception of SHA exposure harm¶			
Harmless	2104	22.8	1§
Harmful	7662	14.6	0.69 (0.61 to 0.78)

*Country weight was applied with each country contributing in proportion to its population aged 15 years or over.²⁸

†Sample size (N) is the unweighted number of e-cigarette non-users for each corresponding individual-level characteristic.

‡aORs were estimated multilevel logistic random-effects models, to take into account the heterogeneity between the countries. The study country effects were considered as random intercepts, and sex, age, level of education, e-cigarette use status and smoking status as adjusting variables. Estimates in bold are statistically significant at 0.05 level.

§Reference category.

¶The sum does not add to the total because of missing values.

non-users, as it is likely that users and bystanders are peers and they socialise together.

Exposure to SHA has its impact on social norm and using e-cigarette. Constant SHA exposure among the youth may increase their susceptibility to using e-cigarettes and tobacco products, as well as decreased their harm perception of e-cigarettes.^{19 57} A higher likelihood of SHA exposure among e-cigarette past users (compared with never users) found in this study may pose a risk of relapse for those who have quit using e-cigarette. An experimental study reported that passive exposure to e-cigarette significantly increased desire to use e-cigarette.²¹ Additionally, exposure to SHA may put current smokers at a risk of being dual users, as they might start using e-cigarettes.^{58 59} Thus, more preventive campaigns are needed to avoid initiation, relapse and dual use in such vulnerable populations.

In line with a study among youth in the USA,⁵⁷ our study found that those who perceived SHA as harmful were less likely to report SHA exposure. Generally, people viewed SHA as less

Table 4 Proportion (%) and adjusted OR (aOR) for at least weekly exposure to e-cigarette secondhand aerosol (SHA) and corresponding 95% CI according to selected country-level characteristics among e-cigarette non-users of European population aged ≥ 15 years. * TackSHS survey, 2017 to 2018

Country-level characteristics	N†	At least weekly exposure to SHA from e-cigarettes	
		%	aOR (95% CI)‡
Geographical area			
Northern Europe	2865	28.2	1§
Western Europe	1974	17.6	0.52 (0.22 to 1.27)
Southern Europe	4013	10.9	0.27 (0.11 to 0.68)
Eastern Europe	2752	11.9	0.35 (0.13 to 0.94)
Gross domestic product per capita			
$\leq 25,000\text{€}$	5711	13.7	1§
$> 25,000\text{€}$	5893	16.7	1.22 (0.51 to 2.89)
Tobacco control scale score			
≤ 50	5712	12.8	1§
> 50	5892	18.0	1.31 (0.62 to 2.79)
Total population smoking prevalence (%)			
<20	2901	20.4	1§
20 to 30	2727	11.4	0.52 (0.22 to 1.25)
> 30	5976	16.4	0.58 (0.25 to 1.37)
<i>P for trend</i>			<0.266
Total population e-cigarette use prevalence (%)			
<1	2727	8.3	1§
1 to 4	6004	11.9	1.64 (1.05 to 2.56)
> 4	2873	27.8	4.35 (2.72 to 6.96)
<i>P for trend</i>			<0.001

*Country weight was applied with each country contributing in proportion to its population aged 15 years or over.³⁸

†Sample size (N) is the unweighted number of e-cigarette non-users for each corresponding country-level characteristic.

‡aOR were estimated multilevel logistic random-effects models, to take into account the heterogeneity between the countries. Estimates in bold are statistically significant at 0.05 level.

§Reference category.

¶Geographical area was categorised into Northern Europe (Ireland, Latvia and England), Western Europe (France and Germany), Southern Europe (Italy, Greece, Portugal and Spain) and Eastern Europe (Bulgaria, Poland and Romania) according to United Nations M49 Standard,³⁵ by the World Bank gross domestic product (GDP) per capita into GDP per capita³⁶ $\leq 25,000\text{€}$ (Bulgaria, Latvia, Romania, Poland, Portugal and Greece) and GDP per capita³⁶ $> 25,000\text{€}$ (England, France, Germany, Ireland, Italy and Spain), by score of³⁷ Tobacco Control Scales ≤ 50 (Bulgaria, Poland, Portugal, Latvia, Greece and Germany) and Tobacco Control Scales > 50 (England, Ireland, France, Romania, Italy and Spain), by country's total smoking prevalence into $< 20\%$ (Ireland, Italy and England), $20\%–30\%$ (Germany, Latvia and Poland) and $> 30\%$ (Bulgaria, France, Greece, Portugal, Romania and Spain)³³ and by country's total population e-cigarette use prevalence into $< 1\%$ (Poland, Portugal and Spain), $1\%–4\%$ (Bulgaria, Germany, Ireland, Italy, Latvia and Romania) and $> 4\%$ (France, Greece and England). The latter two were estimated from the TackSHS survey data.

harmful than SHS.⁶⁰ A parental interview data in the USA has shown that, while compared with smoke-free policy at homes and cars, there were fewer parents who enforced 'vape-free' homes and cars, suggesting that parents perceived e-cigarette aerosol was safe for their children.⁶¹ Therefore, increasing awareness of the potential harmful effects might decrease SHA exposure.

This study was limited by the inherent nature of the cross-sectional study design and the use of self-reported data by respondents. The accuracy of responses, indeed, relies on participants' perception to sense the passive exposure itself. Moreover,

our question did not define the specific sign of SHA exposure (eg, smell, visibility of the cloud, etc) as it may freely capture all possible indicators of SHA exposure. A similar question has also been used by the ITC six European Country survey.²³ Another strength associated with using self-reported exposure is that the respondents assign it to specific setting, which cannot be ascertained when using personal biomarkers of exposure. As the design of our questionnaire does not have a separate question for educational venues, we were unable to estimate specific exposure at such setting. However, we believe this would not undermine our results given the low proportion of student participants (less than 10%) in this study. The questionnaire gathered information on SHA in working and non-working days separately, thus preventing potential information bias derived from using longer times of recall but it cannot ascertain daily prevalence. We have computed prevalence of 'at least weekly' exposure that in addition to be reliable, is useful, given the relatively low exposure to SHA.

There was relatively small sample size in each country (approximately 1000 subjects), but the total sample size is large enough to draw an overall inference. Finally, this study had some differences in sampling methods across countries.³³ However, we ensured the representativeness of the sample in proportion to each country's population aged 1 ≥5 years by applying the weight factors into the analyses.

To our knowledge, this is the first study that investigates self-reported exposure to SHA at the population level in European countries using a standardised questionnaire that allows comparison among countries. The duration of SHA exposure described in this study may offer an alternative measure of SHA exposure burden apart from the prevalence. Additionally, countries selected in this study enable us to understand the variation of SHA exposure in countries with diverse e-cigarette regulatory environment.

In conclusion, we found that there was a substantial proportion and duration of exposure to SHA among non-users of e-cigarettes in indoor settings in European countries, with heterogeneity of exposure across countries and among socio-demographic groups. Thus, governments are strongly recommended to include e-cigarettes in smoke-free laws and tailor such legislation to be specifically targeted to vulnerable groups, particularly young people and former users, to protect them from the harms of SHA exposure and the temptation to (re)fall into nicotine addiction. Enforcement to increase compliance with existing e-cigarette use legislation is needed. Finally, future work should include repeated cross-sectional and/or longitudinal studies on SHA exposure to monitor its burden over time.

What this paper adds

- ▶ The growing use of e-cigarettes has raised concerns as the product is potentially harmful both to users and to bystanders. Yet, e-cigarette use has often been observed in indoor places where smoking is prohibited.
- ▶ Little is known about population exposure to secondhand aerosol from e-cigarette (SHA) in indoor settings in European countries.
- ▶ Our study found that there was a notable proportion and duration of exposure to SHA among non-users in indoor settings in 12 European countries, with variability of exposure across and within countries.

Author affiliations

- ¹Tobacco Control Unit and WHO Collaborating Center for Tobacco Control, Catalan Institute of Oncology (ICO), L'Hospitalet de Llobregat, Catalonia, Spain
- ²Tobacco Control Research Group, Bellvitge Institute for Biomedical Research (IDIBELL), L'Hospitalet de Llobregat, Catalonia, Spain
- ³Department of Clinical Sciences, University of Barcelona (UB), L'Hospitalet de Llobregat, Catalonia, Spain
- ⁴CIBER Respiratory Diseases (CIBERES), Madrid, Spain
- ⁵Department of Environmental Health Sciences, IRCCS - Istituto di Ricerche Farmacologiche "Mario Negri", Milano, Italy
- ⁶School of Public Health, Università Vita-Salute San Raffaele, Milano, Lombardia, Italy
- ⁷CAPHRI-School for Public Health and Primary Care, Maastricht University Medical Center, Maastricht, The Netherlands
- ⁸Institute for Social Marketing, University of Stirling, Stirling, Scotland, United Kingdom
- ⁹TobaccoFree Research Institute Ireland (TFRI), Dublin, Ireland
- ¹⁰Respiratory DepartmentH, Hospital Universitario de la Princesa, Madrid, Spain

Twitter Beladenta Amalia @beladenta, Esteve Fernández @stfvdz, Silvano Gallus @SilvanoGallus and the TackSHS Project Investigators @tackshsproject

Acknowledgements Authors would like to thank Liliane Chatenoud, Tobias Effertz, Filippos T Filippidis, Gergana Geshanova, Sheila Keogan, Hristo Ivanov, José Precioso, Krzysztof Przewozniak, Polina Starchenko, Vergina K Vyzikidou for their help in the preparation, translation and/or submission of the study protocol to local ethics committees.

Collaborators Catalan Institute of Oncology (ICO); Bellvitge Biomedical Research Institute (IDIBELL), Spain: Esteve Fernández, Yolanda Castellano, Marcela Fu, Montse Ballbè, Beladenta Amalia, Olena Tigova; Public Health Agency of Barcelona (ASPB), Spain: Maria José López, Xavier Continente, Teresa Archavala, Elisabet Henderson; Istituto di Ricerche Farmacologiche Mario Negri IRCCS (IRFMN), Italy: Silvano Gallus, Alessandra Lugo, Xiaojie Liu, Cristina Bosetti, Elisa Borroni, Enrico Davoli; Istituto DOXA, Worldwide Independent Network/Gallup International Association, Italy: Paolo Colombo; University of Stirling (UNISTIR), the UK: Sean Semple, Rachel O'Donnell, Ruairaidh Dobson; TobaccoFree Research Institute Ireland (TFRI), Ireland: Luke Clancy, Sheila Keogan, Hannah Byrne; Hellenic Cancer Society - George D. Behrakis Research Lab (HCS), Greece: Panagiotis Behrakis, Anna Tzortzi, Constantine Vardavas, Vergina Konstantina Vyzikidou, Gerasimos Bakelas, George Mattiampa; Fondazione IRCCS Istituto Nazionale dei Tumori (INT), Italy: Roberto Boffi, Ario Ruprecht, Cinzia De Marco, Alessandro Borgini, Chiara Veronese, Martina Bertoldi, Andrea Tittarelli; Istituto per lo Studio, la Prevenzione, e la Rete Oncologica (ISPRO), Italy: Giuseppe Gorini, Giulia Carreras, Barbara Cortini, Simona Verdi, Alessio Lachi, Elisabetta Chellini; Polytechnic University of Cartagena (UPCT), Spain: Ángel López Nicolás, Marta Traperro-Bertran, Daniel Celdrán Guerrero; European Network on Smoking and Tobacco Prevention (ENSP), Belgium: Cornel Radu-Loghin, Dominick Nguyen, Polina Starchenko; Fundación para la Investigación Biomédica del Hospital Universitario La Princesa (IISP), Spain: Joan B Soriano, Julio Ancochea, Tamara Alonso, María Teresa Pastor, Marta Erro, Ana Roca, Patricia Pérez, Elena García Castillo.

Contributors BA, EF and SG had the original idea for the study; SG, XL and AL contributed to the finalisation of the survey questionnaire; BA and XL carried out the statistical analysis with the supervision of AL; BA wrote the first draft of the article in collaboration with XL, AL, MF, EF and SG; AO, SS, LC, JBS and PAvdB made substantial contributions to conception, design and interpretation of data; all the authors contributed to manuscript preparation and approved its final version prior to submission. SG and EF are the guarantors.

Funding This project has received funding from the European Union's Horizon 2020 research and innovation programme under grant agreement No 681040. BA received the support of a fellowship from 'La Caixa' Foundation (ID 100010434; Fellowship code: LCF/BQ/IN17/11620013). AL was supported by a fellowship from the Italian Association for Cancer Research (AIRC). The Tobacco Control Research Group at ICO-IDIBELL (BA, EF and MF) is partly supported by the Ministry of Universities and Research, Government of Catalonia (2017SGR319) and thanks CERCA Programme Generalitat de Catalunya for the institutional support to IDIBELL. EF is partly supported by the Instituto de Salud Carlos III, Government of Spain, cofunded by the European Regional Development Fund (FEDER; INT16/00211 and INT17/00103). The work of SG was partially funded by the Italian League Against Cancer (LILT, Milan).

Disclaimer This paper reflects only the authors' views and the European Commission is not responsible for any use that may be made of the information it contains.

Competing interests None declared.

Patient consent for publication Not required.

Ethics approval We obtained the approval from a local ethics committee in each of the 12 countries. The study protocol has been registered in ClinicalTrials.gov (ID: NCT02928536). All respondents received detailed information about the survey before they provided their consent to participate.

Provenance and peer review Not commissioned; externally peer reviewed.

Data availability statement Data are available upon reasonable request. Please see at www.TackSHS.eu the conditions of use and how to request the data.

ORCID iDs

Beladenta Amalia <http://orcid.org/0000-0002-9718-2693>

Xiaoqiu Liu <http://orcid.org/0000-0001-8620-6592>

Alessandra Lugo <http://orcid.org/0000-0002-4297-6496>

Marcela Fu <http://orcid.org/0000-0002-4962-0927>

Anna Odone <http://orcid.org/0000-0002-5657-9774>

Piet A van den Brandt <http://orcid.org/0000-0001-8781-8099>

Sean Semple <http://orcid.org/0000-0002-0462-7295>

Luke Clancy <http://orcid.org/0000-0003-2407-2263>

Joan B Soriano <http://orcid.org/0000-0001-9740-2994>

Esteve Fernández <http://orcid.org/0000-0003-4239-723X>

Silvano Gallus <http://orcid.org/0000-0002-8967-0400>

REFERENCES

- Cullen KA, Ambrose BK, Gentzke AS, et al. Notes from the Field: Use of Electronic Cigarettes and Any Tobacco Product Among Middle and High School Students — United States, 2011–2018. *MMWR Morb Mortal Wkly Rep* 2018;67:1276–7.
- Huang J, Duan Z, Kwok J, et al. Vaping versus JUULing: how the extraordinary growth and marketing of JUUL transformed the US retail e-cigarette market. *Tob Control* 2019;28:146–51.
- European Commission. *Special Eurobarometer 458 Report Attitudes of Europeans towards tobacco and electronic cigarettes Fieldwork March 2017 May 2017 Survey requested by the European Commission, Special Eurobarometer 458 Report Attitudes of Europeans towards tobacco and e-cigs 2017*.
- Hess I, Lachireddy K, Capon A. A systematic review of the health risks from passive exposure to electronic cigarette vapour. *Public Health Research & Pr* 2016;26:1–9.
- Logue JM, Sleiman M, Montesinos VN, et al. Emissions from electronic cigarettes: assessing vapers' intake of toxic compounds, secondhand exposures, and the associated health impacts. *Environ Sci Technol* 2017;51:9271–9.
- Ramôa CP, Hiler MM, Spindle TR, et al. Electronic cigarette nicotine delivery can exceed that of combustible cigarettes: a preliminary report. *Tob Control* 2016;25:e6–9.
- Wagener TL, Floyd EL, Stepanov I, et al. Have combustible cigarettes Met their match? the nicotine delivery profiles and harmful constituent exposures of second-generation and third-generation electronic cigarette users. *Tob Control* 2017;26:e23–8.
- Olmedo P, Goessler W, Tanda S, et al. Metal concentrations in e-cigarette liquid and aerosol samples: the contribution of metallic coils. *Environ Health Perspect* 2018;126:027010.
- Williams M, Bozhilov K, Ghai S, et al. Elements including metals in the atomizer and aerosol of disposable electronic cigarettes and electronic hookahs. *PLoS One* 2017;12:e0175430.
- Ballbè M, Martínez-Sánchez JM, Sureda X, et al. Cigarettes vs. e-cigarettes: passive exposure at home measured by means of airborne marker and biomarkers. *Environ Res* 2014;135:76–80.
- Grana R, Benowitz N, Glantz SA. E-Cigarettes: a scientific review. *Circulation* 2014;129:1972–86.
- Goniewicz ML, Knysak J, Gawron M, et al. Levels of selected carcinogens and toxicants in vapour from electronic cigarettes. *Tob Control* 2014;23:133–9.
- Fernández E, Ballbè M, Sureda X, et al. Particulate matter from electronic cigarettes and conventional cigarettes: a systematic review and observational study. *Curr Opin Health Rpt* 2015;2:423–9.
- Martínez-Sánchez JM, Ballbè M, Pérez-Ortuño R, et al. Secondhand exposure to aerosol from electronic cigarettes: pilot study of assessment of tobacco-specific nitrosamine (NNAL) in urine. *Gac Sanit* 2019;33:575–8.
- Tzortzi A, Teloniatis S, Matlampa G, et al. Passive exposure to e-cigarette emissions: immediate respiratory effects. *Tob. Prev. Cessation* 2018;4:1–8.
- Bayly JE, Bernat D, Porter L, et al. Secondhand exposure to aerosols from electronic nicotine delivery systems and asthma exacerbations among youth with asthma. *Chest* 2019;155:88–93.
- King AC, Smith LJ, Fridberg DJ, et al. Exposure to electronic nicotine delivery systems (ends) visual imagery increases smoking urge and desire. *Psychol Addict Behav* 2016;30:106–12.
- Faletau J, Glover M, Nosa V, et al. Looks like smoking, is it smoking?: Children's perceptions of cigarette-like nicotine delivery systems, smoking and cessation. *Harm Reduct J* 2013;10:30.
- Agaku IT, Perks SN, Odani S, et al. Associations between public e-cigarette use and tobacco-related social norms among youth. *Tob Control* 2020;29:332–40.
- Mirbolouk M, Charkhchi P, Orimoloye OA, et al. E-Cigarette use without a history of Combustible cigarette smoking among U.S. adults: behavioral risk factor surveillance system, 2016. *Ann Intern Med* 2019;170:76.
- King AC, Smith LJ, McNamara PJ, et al. Passive exposure to electronic cigarette (e-cigarette) use increases desire for combustible and e-cigarettes in young adult smokers. *Tob Control* 2015;24:501–4.
- WHO FCTC COP 7. *WHO report: electronic nicotine delivery systems and electronic non-nicotine delivery systems (ENDS / ENNDS)*, 2016: 7–12.
- Tigova O, Amalia B, Castellano Y, et al. Secondhand exposure to e-cigarette aerosols among smokers: a cross-sectional study in six European countries of the EUREST-PLUS ITC Europe surveys. *Tob Induc Dis* 2018;16:A11.
- Liu X, Lugo A, Gallus S. Passive exposure to electronic cigarette aerosol in Italy: data from the TackSHS pan-European survey. *Tob Induc Dis* 2018;16:A714.
- Matilla-Santander N, Fu M, Ballbè M, et al. Use of electronic cigarettes in public and private settings in Barcelona (Spain). *Environ Res* 2017;158:685–90.
- Windsor-shellard B. Adult smoking habits in the UK : 2015 2017;2016:1–13.
- Patel D, Davis KC, Cox S, et al. Reasons for current E -cigarette use among U.S. adults. *Prev Med* 2016;93:14–20.
- Simonavicius E, McNeill A, Arnott D, et al. What factors are associated with current smokers using or stopping e-cigarette use? *Drug Alcohol Depend* 2017;173:139–43.
- Amalia B. Regulation for e-cigarette use in public and private areas within European countries. *Tobacco Prevention & Cessation*. 2019;5(supplement):A80.
- Liu X, Lugo A, Davoli E, et al. Electronic cigarettes in Italy: a tool for harm reduction or a gateway to smoking tobacco? *Tob Control* 2020;29:148–52.
- Wang TW, Marynak KL, Agaku IT, et al. Secondhand exposure to electronic cigarette aerosol among US youths. *JAMA Pediatr* 2017;171:490–2.
- Fernández E, López MJ, Gallus S, et al. Tackling second-hand exposure to tobacco smoke and aerosols of electronic cigarettes: the TackSHS project protocol. *Gac Sanit* 2020;34:77–82.
- Gallus S, Lugo A, Liu X, et al. Who smokes in Europe? data from 12 European countries in the TACKSHS survey (2017-2018). *J Epidemiol* 2020. In Press.
- Gallus S, Lugo A, La Vecchia C, et al. Pricing policies and control of tobacco in Europe (PPACTE) project: cross-national comparison of smoking prevalence in 18 European countries. *Eur J Cancer Prev* 2014;23:177–85.
- United Nations. UNSD — methodology. Available: <https://unstats.un.org/unsd/methodology/m49/> [Accessed 13 Jun 2019].
- World Bank. GDP per capita (current US\$) | Data. Available: <https://data.worldbank.org/indicator/ny.gdp.pcap.cd> [Accessed 1 Jul 2019].
- Joossens L, Raw M. The tobacco control scale 2016 in Europe, 2017. Available: www.europecancerleagues.org [Accessed 1 Jul 2019].
- European Commission. Database - Eurostat, 2017. Available: <https://ec.europa.eu/eurostat/data/database> [Accessed 15 Jul 2019].
- Potera C. Smoking and secondhand smoke. study finds no level of shs exposure free of effects. *Environ Health Perspect* 2010;118:A474.
- Strulovici-Barel Y, Omberg L, O'Mahony M, et al. Threshold of biologic responses of the small airway epithelium to low levels of tobacco smoke. *Am J Respir Crit Care Med* 2010;182:1524–32.
- Centers for Disease Control and Prevention. The Health Consequences of Smoking - 50 Years of Progress: A Report of the Surgeon General. Atlanta, 2014. Available: www.cdc.gov/tobacco [Accessed 18 Jun 2019].
- Flouris AD, Chorti MS, Pouliantiti KP, et al. Acute impact of active and passive electronic cigarette smoking on serum cotinine and lung function. *Inhal Toxicol* 2013;25:91–101.
- Shipan CR, Volden C. The diffusion of local antismoking policies. Available: <https://pdfs.semanticscholar.org/55ec/5bfb3e9a4b236fd9e54f997d96bbecaffae4.pdf> [Accessed 28 Jun 2019].
- Nykiforuk CIJ, Eyles J, Campbell HS. Smoke-free spaces over time: a policy diffusion study of bylaw development in Alberta and Ontario, Canada. *Health Soc Care Community* 2008;16:64–74.
- Filippidis FT, Agaku IT, Girvalaki C, et al. Relationship of secondhand smoke exposure with sociodemographic factors and smoke-free legislation in the European Union. *Eur J Public Health* 2016;26:344–9.
- Bauld L, Amos A, Ayres J, et al. The impact of smokefree legislation in England: evidence review, 2011. Available: https://assets.publishing.service.gov.uk/government/uploads/system/uploads/attachment_data/file/216319/dh_124959.pdf [Accessed 1 Jul 2019].
- Sureda X, Martínez-Sánchez JM, Fu M, et al. Impact of the Spanish smoke-free legislation on adult, non-smoker exposure to Secondhand smoke: cross-sectional surveys before (2004) and after (2012) legislation. *PLoS One* 2014;9:e89430.
- Martínez-Sánchez JM, Blanch C, Fu M, et al. Do smoke-free policies in work and public places increase smoking in private venues? Table 1. *Tob Control* 2014;23:204–7.
- Smoke Free Partnership. Smokefree Map. Available: <https://smokefreepartnership.eu/our-policy-work/smokefree-map> [Accessed 1 Jul 2019].
- Wadsworth E, Neale J, McNeill A, et al. How and why do smokers start using e-cigarettes? qualitative study of Vapers in London, UK. *Int J Environ Res Public Health* 2016;13:661.

- 51 Trotter L, Wakefield M, Borland R. Socially cued smoking in bars, nightclubs, and gaming venues: a case for introducing smoke-free policies. *Tob Control* 2002;11:300–4.
- 52 Gentzke AS, Homa DM, Kenemer JB, *et al.* Rules to prohibit the use of electronic vapor products inside homes and personal vehicles among adults in the U.S., 2017. *Prev Med* 2018;114:47–53.
- 53 Gallus S, Lugo A, Pacifici R, *et al.* E-Cigarette awareness, use, and harm perceptions in Italy: a national representative survey. *Nicotine Tob Res* 2014;16:1541–8.
- 54 King BA, Patel R, Nguyen KH, *et al.* Trends in awareness and use of electronic cigarettes among US adults, 2010–2013. *Nicotine Tob Res* 2015;17:219–27.
- 55 Vardavas CI, Filippidis FT, Agaku IT. Determinants and prevalence of e-cigarette use throughout the European Union: a secondary analysis of 26 566 youth and adults from 27 Countries. *Tob Control* 2015;24:442–8.
- 56 Rogers EM. *Diffusion of innovations*. 5th ed. New York: Simon & Schuster, 2003.
- 57 Bayly JE, Bernat D, Porter L, *et al.* Prevalence and characteristics of secondhand smoke and secondhand vapour exposure among youth. *Tob Control* 2019;28:305–10.
- 58 Brandon T, Martinez U, Simmons V, *et al.* Dual use of combustible and electronic cigarettes: patterns and associations between products. *Tob Induc Dis* 2018;16:A238.
- 59 Robertson L, Hoek J, Blank M-L, *et al.* Dual use of electronic nicotine delivery systems (ends) and smoked tobacco: a qualitative analysis. *Tob Control* 2019;28:13–19.
- 60 Tan ASL, Bigman CA, Mello S, *et al.* Is exposure to e-cigarette communication associated with perceived harms of e-cigarette secondhand vapour? results from a national survey of US adults. *BMJ Open* 2015;5:e007134.
- 61 Drehmer JE, Nabi-Burza E, Hipple Walters B, *et al.* Parental smoking and e-cigarette use in homes and cars. *Pediatrics* 2019;143:e20183249.