

Longitudinal associations between inflammatory markers and fatigue up to two years after colorectal cancer treatment

Citation for published version (APA):

Querido, N. R., Kenkhuis, M. F., van Roekel, E. H., Breukink, S. O., van Duijnhoven, F. J. B., Janssen-Heijnen, M. L., Keulen, E. T. P., Ueland, P. M., Vogelaar, F. J., Wesselink, E., Bours, M. J., & Weijenberg, M. P. (2022). Longitudinal associations between inflammatory markers and fatigue up to two years after colorectal cancer treatment. *Cancer Epidemiology Biomarkers & Prevention*, 31(8), 1638-1649. <https://doi.org/10.1158/1055-9965.EPI-22-0077>

Document status and date:

Published: 01/08/2022

DOI:

[10.1158/1055-9965.EPI-22-0077](https://doi.org/10.1158/1055-9965.EPI-22-0077)

Document Version:

Accepted author manuscript (Peer reviewed / editorial board version)

Please check the document version of this publication:

- A submitted manuscript is the version of the article upon submission and before peer-review. There can be important differences between the submitted version and the official published version of record. People interested in the research are advised to contact the author for the final version of the publication, or visit the DOI to the publisher's website.
- The final author version and the galley proof are versions of the publication after peer review.
- The final published version features the final layout of the paper including the volume, issue and page numbers.

[Link to publication](#)

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal.

If the publication is distributed under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license above, please follow below link for the End User Agreement:

www.umlib.nl/taverne-license

Take down policy

If you believe that this document breaches copyright please contact us at:

repository@maastrichtuniversity.nl

providing details and we will investigate your claim.

LONGITUDINAL ASSOCIATIONS BETWEEN INFLAMMATORY MARKERS
AND FATIGUE UP TO TWO YEARS AFTER COLORECTAL CANCER TREATMENT

Nadira R. Querido¹, Marlou-Floor Kenkhuis¹, Eline H. van Roekel¹, Stéphanie O. Breukink^{2,3,4}, Fränzel J.B. van Duijnhoven⁵, Maryska L.G. Janssen-Heijnen^{1,6}, Eric T.P. Keulen⁷, Per Magne Ueland⁸, F. Jeroen Vogelaar¹, Evertine Wesselink⁵, Martijn J.L. Bours¹, Matty P. Weijenberg¹

1 Department of Epidemiology, GROW School for Oncology and Reproduction, Maastricht University, Maastricht, the Netherlands

2 Department of Surgery, Maastricht University Medical Centre+, Maastricht, the Netherlands

3 GROW School for Oncology and Reproduction, Maastricht University Medical Centre+, Maastricht, the Netherlands

4 NUTRIM School of Nutrition and Translational Research in Metabolism, Maastricht University Medical Centre+, Maastricht, the Netherlands

5 Division of Human Nutrition and Health, Wageningen University & Research, Wageningen, the Netherlands

6 Department of Clinical Epidemiology, Viecuri Medical Center, Venlo, the Netherlands

7 Department of Internal Medicine and Gastroenterology, Zuyderland Medical Centre, Sittard-Geleen, the Netherlands

8 BEVITAL, 5021 Bergen, Norway

The authors declare no potential conflicts of interest.

Corresponding author: Matty P. Weijnenberg, Maastricht University, Department of Epidemiology, GROW School for Oncology and Reproduction, P.O. BOX 616, 6200 MD Maastricht, the Netherlands. Phone: +31 43 3882393 E-mail: mp.weijnenberg@maastrichtuniversity.nl

Running title: Longitudinal associations between inflammation and fatigue

Keywords: Inflammation, cytokines, fatigue, colorectal cancer, longitudinal

Registration: The EnCoRe study was registered at trialregister.nl as NL6904 (former ID: NTR7099).

Abstract

Background: Fatigue is often reported by colorectal cancer survivors (CRC) and largely impacts their quality of life. Inflammation has been linked to fatigue mainly in breast cancer patients. Therefore, we investigated how inflammation is longitudinally associated with fatigue in CRC survivors, up to 2 years post-treatment.

Methods: A total of 257 patients from the ongoing Energy for life after ColoRectal cancer (EnCoRe) cohort study were included in the analysis. Plasma levels of IL6, IL8, IL10, TNF α , hsCRP, and fatigue were measured at 6 weeks, 6, 12, and 24 months post-treatment. Fatigue was measured through the validated Checklist Individual Strength (CIS total, 20-140), consisting of four subscales – subjective fatigue (8-56), motivation (4-28), physical activity (3-21), and concentration (5-35), and the EORTC QLQ-C30 fatigue subscale (0-100). Linear mixed-models were used to assess the confounder-adjusted longitudinal associations between inflammatory markers and overall fatigue along with the subscales.

Results: Mean levels of CIS fatigue decreased from 62.9 at 6 weeks to 53.0 at 24 months. In general, levels of inflammatory markers also decreased over time. No statistically significant longitudinal associations were found between IL6, IL8, IL10, TNF α , and fatigue. Higher levels of hsCRP were associated with more CIS fatigue (β per SD 3.21, 95% CI 1.42; 5.01) and EORTC fatigue (β 2.41, 95% CI 0.72; 4.10).

Conclusion: Increased levels of hsCRP are longitudinally associated with more post-treatment fatigue in CRC survivors.

Impact: These findings suggest that low-grade inflammation may play a role in fatigue reported by CRC survivors up to 2 years post-treatment.

1. Introduction

Population ageing, screening programs, early detection, and more effective treatments have led to an increase in the number of colorectal cancer (CRC) survivors (1). In 2020, worldwide, over 5 million individuals had a CRC diagnosis in the past 5 years (2). Due to the rising number of CRC survivors, it becomes increasingly important to address factors that impact their health-related quality of life (HRQoL) post-treatment. There are several chronic or late effects caused by both CRC and its treatment, such as fatigue, pain, bowel dysfunction, and emotional distress, all of which can affect a patient's HRQoL (3, 4).

Fatigue is a common and debilitating symptom experienced by CRC survivors during and post-treatment (5, 6). Reported rates of fatigue among CRC survivors range from 12-69.7% depending on the measurement instrument used and time elapsed since treatment (6-10). Results from prospective studies, including ours, and a systematic review showed that fatigue peaked between 6 weeks and 6 months post-treatment but persisted up to two years post-treatment (9, 11, 12). Many factors, such as treatment, comorbidities, and physical and psychological factors, possibly contribute to cancer-related fatigue (5, 13). Furthermore, there is an increasing interest in the underlying biological mechanisms of fatigue (14).

Inflammation has been mainly identified as an underlying mechanism in post-treatment cancer-related fatigue, with the majority of studies performed in breast cancer survivors (13, 15, 16). Current thought is that production of proinflammatory cytokines in the periphery stimulates the brain resulting in fatigue, among other sickness behaviors (17, 18). Indeed, elevated circulating levels of pro-inflammatory markers, such as interleukin (IL)-6, tumor necrosis factor- α (TNF α) and C-reactive protein (CRP), have been linked to more fatigue in breast cancer survivors (15, 19, 20). In contrast, anti-inflammatory cytokines, such as IL10, may attenuate sickness behavior, but little is known in relation to cancer-related fatigue (18,

21). Most longitudinal studies exploring the association between cancer-related fatigue and inflammation in breast cancer survivors focus on the period during or up to 6 months post-treatment and therefore have not assessed longer term effects (22, 23). In addition, there are important differences between breast cancer and CRC survivors regarding several characteristics, namely age, sex, and treatment, that can differentially affect fatigue. Some studies point to sex differences in both immune response and reporting of fatigue (24-27). Thus, despite the evidence of links between inflammation and fatigue in breast cancer survivors, a further exploration of this association is needed in CRC survivors.

Few studies, with differing methodologies, have explored the link between inflammation and fatigue in CRC survivors (9, 28-30). These methodological differences include the measurement instruments used to assess fatigue, the start (pre- or post-treatment) and duration of follow-up time, and the availability of repeated measurements for both the inflammatory markers and fatigue. To our knowledge, only one study investigated the association between several inflammatory markers, excluding hsCRP, and fatigue up to two years post-treatment with repeated measurements over time (9).

Investigating how post-treatment inflammation is related to post-treatment fatigue over time will help to better understand the role of inflammation in the progression of cancer-related fatigue in CRC survivors. Therefore, the primary aim of this study is to determine how plasma levels of inflammatory markers, namely IL6, IL8, IL10, TNF α , high-sensitivity C-reactive protein (hsCRP), are longitudinally associated with overall fatigue, as well as different dimensions of fatigue (subjective fatigue, motivation, physical activity, and concentration) in CRC survivors followed up from 6 weeks until 2 years post-treatment.

2. Methods

2.1. *Study design and population*

Data analysis was performed with longitudinal data collected from April 18th, 2012 up until November 1st, 2016, from the Energy for life after ColoRectal cancer (EnCoRe) study. The EnCoRe study is an ongoing prospective cohort study with patient recruitment at three participating centers: Maastricht University Medical Center+, VieCuri Medical Center, and Zuyderland Medical Centre (31). Eligible for participation were men and women above the age of 18, diagnosed with stage I-III CRC. Exclusion criteria were stage IV CRC, inability to understand and speak Dutch, residential address outside of the Netherlands, or the presence of comorbidities that could impede a successful study participation, including cognitive and visibility/hearing disorders (31).

Patients were enrolled at diagnosis and followed up with repeated measurements at 6 weeks (n=237), 6 months (n=184), 12 months (n=150), and 24 months post-treatment (n=63). Study measurements were performed during home visits. In case participants were ill (e.g. the flu) or hospitalized, home visits were postponed. Participation rate at diagnosis was 46% and >90% at all post-treatment follow-up visits (Supplemental Figure 1). The main reason for the decrease in sample size as follow-up time increases was that not all participants included at diagnosis had reached the subsequent follow-up points on November 1st, 2016. The EnCoRe study was approved by the Medical Ethics Committee of the Academic Hospital Maastricht and Maastricht University, the Netherlands (Netherlands Trial Register no. NL6904). The study was conducted in accordance with the principles of the Declaration of Helsinki (version 7, October 2008).

2.1. *Plasma inflammatory markers (exposure)*

Fasting blood samples collected during home visits at 6 weeks, 6, 12, and 24 months post-treatment were used to assess plasma levels of inflammatory markers. After collection into

EDTA tubes, blood samples were centrifuged, aliquoted into plasma, and stored in a freezer at -80°C until analysis (32). A custom-made multiplex assay using electrochemiluminescence detection (Meso Scale Diagnostics, Rockville, MD, USA) was used to measure plasma concentration (pg/ml) of IL6, IL8, IL10, and TNF α . Assay plates were analysed on a QuickPlex SQ 120 plate reader (Meso Scale Diagnostics), according to manufacturer's instructions, at Wageningen University & Research, as described previously (32). Alongside the calibration curve, three quality controls were included per plate. All samples were analysed in duplicates and the sample mean was accepted if the coefficient of variation (CV) was <40% (32). Inter- and intra-assay CVs were <8%, with reported values deviating less than 15% from target values (32). Levels of hsCRP were measured at 6 weeks, 6 and 12 months post-treatment. Plasma concentration (μ g/ml) of hsCRP was determined through an immuno-MALDI (matrix-assisted laser desorption/ionization) mass spectrometry method (BEVITAL, Bergen, Norway) (33). The inter-assay CV ranged from 3-6%. hsCRP is used to measure lower levels of CRP which reflect low-grade systemic inflammation (34, 35).

Summary inflammatory z-scores were calculated to group the inflammatory markers and improve statistical efficiency (32, 36). Higher z-scores indicate higher levels of inflammation. First, normalized z-scores from each inflammatory marker were calculated as $z = (x_{ij} - \mu_j) / \sigma_j$, in which x is the participant's (i) inflammatory marker value at a given visit (j), μ is the study population mean, and σ is the study standard deviation, both at given visits (j) (32). Two summary inflammatory z-scores were computed for each participant, at each time point, to use all available data. One was calculated by summing the normalized z-scores of IL6, IL8, TNF α , hsCRP, and subtracting IL10, and thus only includes patients with measurements up to 12 months post-treatment. The other summary inflammatory z-score excluded hsCRP thereby including patients with data available at all post-treatment time points.

2.2. *Fatigue (outcome)*

The validated Checklist Individual Strength (CIS) and the European Organisation for Research and Treatment of Cancer Quality of Life Questionnaire-Core 30 (EORTC QLQ-C30) fatigue subscale were used to measure fatigue at 6 weeks, 6, 12, and 24 months post-treatment. The CIS is a 20-item questionnaire composed of 4 subscales – subjective fatigue (8-56), motivation (4-28), physical activity (3-21), and concentration (5-35) (37). The total fatigue score (20-140) was obtained by summing all item scores. Higher scores represent higher levels of fatigue. The EORTC QLQ-C30 fatigue subscale contains 3 items and ranges from 0-100 (38).

Although initially developed for patients with chronic fatigue syndrome (37), the CIS has been used to measure fatigue in cancer survivors (39). In a study among working people, the CIS was able to adequately distinguish fatigued and non-fatigued individuals (40). A recent study assessed the construct validity of the CIS subjective fatigue subscale and the EORTC QLQ-C30 fatigue subscale in cancer survivors (n=320) and found a high Spearman rank correlation coefficient of 0.77 (41).

2.3. *Other relevant variables*

At the time of diagnosis, patients reported sex and birth date, which was used to calculate the age at each post-treatment time point (11). Data on treatment, such as chemotherapy and radiotherapy, were obtained from clinical records. The number of comorbidities at each post-treatment time point was determined using the 13-item Self-Administered Comorbidity Questionnaire (42). Height and weight, measured by trained dietitians, were used to calculate body mass index (BMI) at every time point. Current smoking status at each time point was self-reported. Information on use of non-steroidal anti-inflammatory drugs (NSAIDs) during the 6 months prior to the follow-up time point was collected using self-reported questionnaires (32). Physical activity was evaluated using the

validated Short QUestionnaire to ASsess Health-enhancing physical activity (SQUASH) (43). Ainsworth's Compendium of Physical Activities was used to give activities a metabolic equivalent of task (MET) value (44). Activities were categorized as light-physical activity (LPA) (<3 MET) or moderate-to-vigorous physical activity (MVPA) (≥ 3 MET), and total time spent in each activity was calculated as hours/week (11).

2.4. *Statistical analysis*

Descriptive analyses were performed to describe patient characteristics at 6 weeks (i.e. the baseline for longitudinal analyses). Categorical variables were presented as frequencies with percentages, and continuous variables as the mean with standard deviation (SD) or medians with interquartile range (IQR) for normally and non-normally distributed data, respectively. Data on inflammatory markers, summary inflammatory z-scores, and fatigue, including the subscales, were presented for all post-treatment time points.

Linear mixed model regression was used to investigate the longitudinal associations between levels of inflammatory markers and fatigue (45). The regression coefficients obtained are a weighted average of the inter-individual (between-subject) differences and intra-individual (within-subject) changes (45). Therefore, separate hybrid models were used to disentangle the intra- and inter-individual components (46). To estimate the intra-individual association, the deviation of an individual's level of inflammatory marker from the person-mean was modelled. The regression coefficient from this model represents changes in fatigue over time in relation to a one-unit change in levels of inflammatory markers over time within individuals. To estimate the inter-individual association, a centered person-mean value of an inflammatory marker – difference between a subject's mean value of the inflammatory marker and the sample mean – was modelled to obtain a regression coefficient which indicates the average difference over time in fatigue between individuals in relation to a one-unit difference in mean levels of inflammatory markers between individuals.

To improve interpretability, levels of inflammatory markers were divided by their SD at 6 weeks to obtain regression coefficients that represented the difference in fatigue per SD increase of the inflammatory marker. The first model included age at measurement (years), sex (men/women), and time since diagnosis (days). The second model included additional potential confounders, selected *a priori* based on the available literature: NSAIDs (yes/no), BMI (kg/m²), physical activity (LPA and MVPA - hours/week), comorbidities (0, 1, ≥2), treatment (chemotherapy/radiotherapy, yes/no), and smoking status (yes/no). The likelihood ratio test was used to evaluate whether including a random slope improved the model fit (45). The false discovery rate (FDR) method ($q < 0.05$) was used to correct for multiple testing of the various exposures with the outcome (47). This was applied separately for each outcome (CIS: total fatigue, subjective fatigue, motivation, physical activity, concentration; EORTC fatigue). No correction was performed for the inflammatory z-scores since they are correlated with the inflammatory markers.

Post-hoc subgroup analyses were performed for sex to explore the longitudinal associations in men and women separately; testing for interaction was done by including a product-term for the inflammatory marker and sex in each model. Sensitivity analyses excluding participants with recurrence and participants who died were performed.

To further explore the role of hsCRP, linear mixed model regression was performed with hsCRP categorized to represent normal values (≤ 3 mg/L), low-grade inflammation (3-10mg/L), and acute inflammation (> 10 mg/L) (35, 48).

Statistical analyses were conducted in STATA (version 15). P-values below 0.05 (two-sided) after correction for multiple testing were considered statistically significant.

2.5. *Data availability*

Data analysed in the manuscript, code book, and analytic code will be made available upon request pending (e.g., application and approval, payment, other) to co-author Dr. Martijn Bours.

3. Results

3.1. *Participant characteristics*

Data on fatigue and IL6, IL8, IL10, and TNF α was available for 237 participants at 6 weeks, 184 at 6 months, 150 at 12 months, and 63 at 24 months post-treatment. Data on fatigue and hsCRP was available for 200 participants at 6 weeks, 148 at 6 months, and 114 at 12 months post-treatment. Participants were on average 67 years old, and the majority were men (68.8%) (Table 1). There were some differences between men and women, notably a higher percentage of women had two or more comorbidities compared to men (women: 68.9%; men: 47.9%). Women also reported higher median levels of LPA than men (women: 14.0 hours/week, IQR 7.0-24.5; men: 6.3 IQR 1.2-12.0), but men reported higher median levels of MVPA than women (men: 9.0 hours/week, IQR 3.5-16.3; women: 4.1 IQR 1.5-7.0). The percentage of men who received radiotherapy was higher than women (men: 30.7%, women: 18.9%) and more men were diagnosed with rectum cancer (men: 42.3%, women: 29.7%).

3.2. *Fatigue and inflammatory markers*

Total fatigue was highest at 6 weeks post-treatment (62.9, SD 26.5) and decreased over time, with the largest decrease occurring between 6 and 12 months post-treatment (Figure 1, Table 2). Across all time points, women reported higher levels of fatigue compared to men. This was also observed in the subjective fatigue subscale, reduced motivation, reduced concentration, being most pronounced for subjective fatigue (Table 2, Supplemental Figure 2). The CIS total fatigue and EORTC fatigue subscale were significantly correlated at all time

points (range: 0.68-0.76). Median levels of IL6, IL10, and TNF α slightly decreased over the course of 24 months post-treatment, and for levels of hsCRP up to 12 months post-treatment, while IL8 increased between the 12 and 24 months time points (Figure 1, Table 2). Pearson correlation coefficients indicated weak to moderated correlations between the inflammatory markers at 6 weeks (range: -0.02 to 0.45), with similar ranges at following time points (Supplemental Figure 3).

3.3. *Longitudinal associations between inflammatory markers and fatigue*

The coefficients presented represent the change in fatigue score for one SD increase of the inflammatory markers (Figure 2, Table 3, Supplemental Figure 4). In the fully adjusted models after FDR correction, there were no statistically significant overall, intra- or inter-individual associations between IL6, IL8, IL10, TNF α , and CIS total fatigue, as well as the subscales. Similar results were observed in the analyses with IL6, IL8, IL10, TNF α and the EORTC fatigue subscale.

After fully adjusting the model and FDR correction, higher levels of hsCRP were longitudinally associated with more CIS total fatigue (β 3.21, 95% CI 1.42; 5.01), subjective fatigue (β 1.82, 95% CI 0.94; 2.70), reduced motivation (β 0.85, 95% CI 0.41; 1.29), and EORTC fatigue (β 2.41, 95% CI 0.72; 4.10). Applying hybrid models revealed a significant inter-individual association between hsCRP and CIS total fatigue (β 5.44, 95% CI 1.61; 9.27). Additionally, higher levels of hsCRP were longitudinally associated with higher scores, both between- and within-subjects, in the subjective fatigue and reduced motivation subscales. The sensitivity analyses indicate that associations were similar after excluding participants who had a recurrence or died (Supplemental Table 1).

Analyses with the summary inflammatory z-score including hsCRP indicated that more inflammation was associated with more CIS total fatigue (β 2.42, 95% CI 0.06; 4.79) and

EORTC fatigue (β 4.49, 95% CI 1.97; 7.01). For CIS total fatigue, an inter-individual association was observed (β 6.71, 95% CI 2.43; 11.00) while the intra-individual association was small and non-significant (β 0.69, 95% CI -2.07; 3.45). In the analyses with EORTC fatigue, both the inter-individual (β 5.75, 95% CI 1.39; 10.12) and intra-individual (β 3.92, 95% CI 0.93; 6.91) associations were statistically significant. The summary inflammatory z-score excluding hsCRP was associated with more EORTC fatigue (β 2.29, 95% CI 0.34; 4.24) but not with CIS total fatigue (β 0.74, 95% CI -1.13; 2.60). To ensure the 24-month time point was not responsible for the different results between the inflammatory z-scores including and excluding hsCRP, extra analyses excluding the 24-month time point were performed for the inflammatory z-score excluding hsCRP. The results led to the same conclusions with similar effect sizes between the inflammatory score excluding hsCRP and fatigue in which all time points were considered.

Results from the exploratory analysis indicated that survivors with levels of hsCRP between 3-10mg/L, and levels >10mg/L experienced more fatigue compared to those with levels \leq 3mg/L (Figure 3, Supplemental Table 2). In subgroup analysis, the statistically significant associations after FDR correction for the individual inflammatory markers were only observed in men (Supplemental Table 3). Additionally, only 5 out of 84 interaction terms were statistically significant.

4. Discussion

No statistically significant associations were found between IL6, IL8, IL10, TNF α and CIS and EORTC fatigue after FDR correction. Higher levels of hsCRP were longitudinally associated with more fatigue from 6 weeks to 12 months post-treatment. Statistically significant inter-individual associations were observed, indicating that CRC survivors with higher mean levels of hsCRP over time reported higher scores of total fatigue. Similar trends were observed

in the subjective and reduced motivation subscales, where both inter- and intra-individual associations for hsCRP were statistically significant. Additionally, statistically significant associations were found between the summary inflammatory score including hsCRP and both CIS and EORTC fatigue. Together these findings suggest that higher levels of low-grade inflammation are associated with more fatigue in CRC survivors.

Findings from longitudinal CRC studies are scarce and inconsistent, the latter likely due to methodological differences in the timing and frequency of measurements for the inflammatory markers and fatigue, the duration of follow-up time, and the types of measurement instruments used to assess the inflammatory markers and fatigue (22, 23). A recent study of 236 stage I-IV CRC survivors did not find statistically significant associations between levels of IL6, IL8, TNF α , CRP measured pre-surgery, and fatigue measured pre-surgery and at 6 and 12 months post-surgery (30). However, unlike this study, only pre-operative inflammatory markers were used, and fatigue was only measured using the EORTC QLQ-C30 fatigue subscale, which mainly measures physical fatigue. A study in patients with localized CRC found weak correlations of IL6, IL8, and IL10 with fatigue at 6 (ρ -0.16 to -0.20) and 24 months (ρ -0.16 to -0.30) after treatment, but not with TNF α (9). From the inflammatory markers we investigated, excluding hsCRP which was not measured, only IL8 was longitudinally inversely associated with more fatigue. Another study on CRC (n=50) and esophageal cancer (n=53) patients found a significant association between IL6 and a component score of fatigue-centered symptom cluster, but not between IL6, IL10, and fatigue severity (29). In the latter study, fatigue was measured weekly for 13 weeks after treatment initiated and the inflammatory markers were measured pre-treatment, during the 5-6 weeks of treatment, and 1 month post-treatment. Since IL10 is considered to have anti-inflammatory properties, it was expected to be inversely associated with fatigue (49, 50). We observed an inverse association

between patients, for both CIS and EORTC fatigue, but this was non-significant after FDR correction. Although evidence is still scarce and inconsistent, higher levels of pro-inflammatory markers seem to be associated with more fatigue in CRC survivors.

A cross-sectional study in 299 disease-free breast cancer survivors, at 4 years post-diagnosis on average, reported a significant association between levels of hsCRP and fatigue (20). Other inflammatory markers, such as IL6, were analyzed but no statistically significant associations were found. Similar results were found in a longitudinal study in breast (n=28) and prostate cancer (n=20) survivors during radiotherapy (51). Both studies argued that pro-inflammatory cytokines, such as IL6, are produced in low quantities and thus harder to detect, possibly explaining the lack of association, as seen in our study (20, 51).

No other studies have used summary inflammatory z-scores to assess the association between inflammatory markers and fatigue. In our study, a significant association was found between the inflammatory z-score excluding hsCRP and EORTC fatigue. This association was not observed in the analysis with CIS fatigue and this difference is possibly explained by the weaker association between IL8 and CIS fatigue compared to EORTC fatigue. Higher levels of the inflammatory z-score including hsCRP were statistically significantly associated with more fatigue. This association is likely driven by levels of hsCRP since the association between the inflammatory z-score excluding hsCRP and fatigue remained non-significant, and with similar effect sizes, after excluding the 24-month time point.

In summary, results from the main analyses add to the existing body of literature on inflammation and fatigue and suggest a link between hsCRP and fatigue. hsCRP can detect low CRP in the blood, and thus can be used to evaluate low-grade inflammation (34, 52). Low-grade inflammation can reduce cellular energy availability and increase energy expenditure, creating an imbalance which possibly explains persistent fatigue (53). CRP is an acute-phase protein mainly upregulated by IL6, and therefore considered a downstream marker for IL6

activity (52). Other cytokines such as $\text{TNF}\alpha$, IL1, IL1 β are also involved in the production of acute-phase proteins (54, 55), and thus require further research as to whether they could be potential targets for intervention (52, 56-58).

In terms of clinical relevance, the observed effect sizes from the fully adjusted models were smaller than the minimal clinically important difference (MCID) defined as 9.3 points for CIS total fatigue (59) and 9 points for EORTC fatigue (60, 61). The largest effect sizes were observed in the analysis with categories of hsCRP where levels $>10\text{mg/L}$ were associated with a 6.14 point (95% CI 0.10; 12.19) increase in EORTC fatigue score, compared to levels $\leq 3\text{mg/L}$. Results from these analyses provide a better comparison to the MCID as the cut-off values chosen are more clinically relevant than the SD increments used in the main analysis (62, 63). Despite not reaching the MCID, the results provide evidence for a longitudinal association between higher levels of hsCRP and an increase in post-treatment fatigue in CRC survivors.

Results from subgroup analysis indicated that the association between hsCRP and fatigue was only present in men. However, this should be interpreted with caution as the analysis in men had twice the sample size as the women's analysis, rendering the associations in women less stable. Furthermore, most of the interaction terms were non-significant.

One of the strengths of this study was the availability of repeated measurements for both inflammatory markers and fatigue, as well as potential confounders. Additionally, the use of hybrid models to disentangle between- and within-individual associations was important to understand how changes in inflammation within-individuals are, on average, related to fatigue over time. To date, this approach has not been attempted by any of the studies exploring an association between inflammatory markers and fatigue.

A limitation of the present study is its observational nature which does not allow for any causal inference. Moreover, patients with higher levels of fatigue at time of diagnosis may view

the measurements involved (i.e. filling out questionnaires and blood collection) as being too burdensome. Thus, patients with higher levels of fatigue could be underrepresented in the study, in part explaining the 45% participation rate at diagnosis and potentially causing an underestimation of the true association. Although the participation rate at diagnosis was 45%, our interest was in the association between inflammation and fatigue, specifically in the post-treatment phase, and all follow-up participation rates were high ($\geq 90\%$). The decrease in sample size as follow-up time increased was mainly due to patients not reaching those time points at the time of data-freeze. Therefore, most participants with missing data are likely missing at random. The smaller sample sizes decrease the power to detect true associations and provide less information on the long-term post-treatment associations of inflammation and fatigue. Additionally, to minimize the potential impact of time of sampling on hsCRP values, which exhibits diurnal variations (64, 65), all samples were collected in fasting individuals during the morning period before breakfast after an overnight fast.

In conclusion, the current study found that higher levels of hsCRP were longitudinally associated with more fatigue in CRC survivors up to 12 months post-treatment. Further longitudinal studies with larger sample sizes will help provide stronger evidence on the long-term association between low-grade inflammation and fatigue post-treatment.

Acknowledgements: The EnCoRe study was supported by a grant from Kankeronderzoekfonds Limburg as part of Health Foundation Limburg (grant 00005739, recipient - M.P. Weijenberg), from Stichting Alpe d'Huzes within the research program “Leven met kanker” of the Dutch Cancer Society grants UM 2010-4867 (recipient - M.P. Weijenberg) and UM 2012-5653 (recipient – M.P. Weijenberg), and by ERA-NET on Translational Cancer Research (TRANSCAN: Dutch Cancer Society (UM 2014-6877, recipient – M.P. Weijenberg)). The measurement of inflammatory markers was funded by a grant from Alpe

d'Huzes/ Dutch Cancer Society (UW2013-6397, recipient – F.J.B. van Duijnhoven). M. Kenkhuis is supported by a grant from Wereld Kanker Onderzoek Fonds (WKOF) / World Cancer Research Fund International (WCRF) (grant number 2017/1619 – recipient M.J.L. Bours). E.H. van Roekel is supported by the Wereld Kanker Onderzoek Fonds (WKOF), as part of the World Cancer Research Fund International grant program (grant number 2016/1620 – recipient M.P. Weijenberg).

We would like to thank all participants of the EnCoRe study and the health professionals in the three hospitals involved in the recruitment of participants of the study: Maastricht University Medical Centre+, VieCuri Medical Centre, and Zuyderland Medical Centre. We would also like to thank the MEMIC centre for data and information management for facilitating the logistic processes and data management of our study. We thank Michiel Balvers and Nhien Ly at Wageningen University & Research for their work on the inflammation markers. Finally, we would like to thank the research dietitians and research assistant who are responsible for patient inclusion and follow-up, performing home visits, as well as data collection and processing.

5. References

1. Ait Ouakrim D, Pizot C, Boniol M, Malvezzi M, Boniol M, Negri E, et al. Trends in colorectal cancer mortality in Europe: retrospective analysis of the WHO mortality database. *BMJ*. 2015;351:h4970.
2. IARC, WHO. Colorectal cancer fact sheet 2020 [Available from: https://gco.iarc.fr/today/data/factsheets/cancers/10_8_9-Colorectum-fact-sheet.pdf].
3. Bours MJ, van der Linden BW, Winkels RM, van Duijnhoven FJ, Mols F, van Roekel EH, et al. Candidate Predictors of Health-Related Quality of Life of Colorectal Cancer Survivors: A Systematic Review. *Oncologist*. 2016;21(4):433-52.
4. Marventano S, Forjaz M, Grosso G, Mistretta A, Giorgianni G, Platania A, et al. Health related quality of life in colorectal cancer patients: state of the art. *BMC Surg*. 2013;13 Suppl 2:S15.
5. Aapro M, Scotte F, Bouillet T, Currow D, Vigano A. A Practical Approach to Fatigue Management in Colorectal Cancer. *Clin Colorectal Cancer*. 2017;16(4):275-85.
6. Thong MSY, Mols F, Wang XS, Lemmens VEPP, Smilde TJ, van de Poll-Franse LV. Quantifying fatigue in (long-term) colorectal cancer survivors: a study from the population-based patient reported outcomes following initial treatment and long term evaluation of survivorship registry. *Eur J Cancer*. 2013;49(8):1957-66.
7. Wang XS, Zhao F, Fisch MJ, O'Mara AM, Cella D, Mendoza TR, et al. Prevalence and characteristics of moderate to severe fatigue: a multicenter study in cancer patients and survivors. *Cancer*. 2014;120(3):425-32.
8. Husson O, Mols F, van de Poll-Franse LV, Thong MSY. The course of fatigue and its correlates in colorectal cancer survivors: a prospective cohort study of the PROFILES registry. *Support Care Cancer*. 2015;23(11):3361-71.

9. Vardy JL, Dhillon HM, Pond GR, Renton C, Dodd A, Zhang H, et al. Fatigue in people with localized colorectal cancer who do and do not receive chemotherapy: a longitudinal prospective study. *Ann Oncol.* 2016;27(9):1761-7.
10. van Baar H, Bours MJL, Beijer S, van Zutphen M, van Duijnhoven FJB, Kok DE, et al. Body composition and its association with fatigue in the first 2 years after colorectal cancer diagnosis. *J Cancer Surviv.* 2021;15(4):597-606.
11. van Roekel EH, Duchateau J, Bours MJL, van Delden L, Breedveld-Peters JJL, Koole JL, et al. Longitudinal associations of light-intensity physical activity with quality of life, functioning and fatigue after colorectal cancer. *Qual Life Res.* 2020;29(11):2987-98.
12. Rutherford C, Müller F, Faiz N, King MT, White K. Patient-reported outcomes and experiences from the perspective of colorectal cancer survivors: meta-synthesis of qualitative studies. *J Patient Rep Outcomes.* 2020;4(1):27.
13. Yang S, Chu S, Gao Y, Ai Q, Liu Y, Li X, et al. A Narrative Review of Cancer-Related Fatigue (CRF) and Its Possible Pathogenesis. *Cells.* 2019;8(7).
14. Bower JE. Cancer-related fatigue--mechanisms, risk factors, and treatments. *Nat Rev Clin Oncol.* 2014;11(10):597-609.
15. Bower JE, Lamkin DM. Inflammation and cancer-related fatigue: mechanisms, contributing factors, and treatment implications. *Brain Behav Immun.* 2013;30 Suppl:S48-57.
16. LaVoy ECP, Fagundes CP, Dantzer R. Exercise, inflammation, and fatigue in cancer survivors. *Exerc Immunol Rev.* 2016;22:82-93.
17. Bower JE. The role of neuro-immune interactions in cancer-related fatigue: Biobehavioral risk factors and mechanisms. *Cancer.* 2019;125(3):353-64.
18. Dantzer R, O'Connor JC, Freund GG, Johnson RW, Kelley KW. From inflammation to sickness and depression: when the immune system subjugates the brain. *Nature reviews Neuroscience.* 2008;9(1):46-56.

19. Collado-Hidalgo A, Bower JE, Ganz PA, Cole SW, Irwin MR. Inflammatory Biomarkers for Persistent Fatigue in Breast Cancer Survivors. *Clin Cancer Res.* 2006;12(9):2759-66.
20. Orre IJ, Reinertsen KV, Aukrust P, Dahl AA, Fosså SD, Ueland T, et al. Higher levels of fatigue are associated with higher CRP levels in disease-free breast cancer survivors. *J Psychosom Res.* 2011;71(3):136-41.
21. Dantzer R. Cytokine-induced sickness behaviour: a neuroimmune response to activation of innate immunity. *Eur J Pharmacol.* 2004;500(1):399-411.
22. Saligan LN, Kim HS. A systematic review of the association between immunogenomic markers and cancer-related fatigue. *Brain Behav Immun.* 2012;26(6):830-48.
23. Schubert C, Hong S, Natarajan L, Mills PJ, Dimsdale JE. The association between fatigue and inflammatory marker levels in cancer patients: a quantitative review. *Brain Behav Immun.* 2007;21(4):413-27.
24. Husain AF, Stewart K, Arseneault R, Moineddin R, Cellarius V, Librach SL, et al. Women experience higher levels of fatigue than men at the end of life: a longitudinal home palliative care study. *J Pain Symptom Manage.* 2007;33(4):389-97.
25. Klein SL, Flanagan KL. Sex differences in immune responses. *Nature Reviews Immunology.* 2016;16(10):626-38.
26. Lasselin J, Lekander M, Axelsson J, Karshikoff B. Sex differences in how inflammation affects behavior: What we can learn from experimental inflammatory models in humans. *Front Neuroendocrinol.* 2018;50:91-106.
27. Valentine RJ, McAuley E, Vieira VJ, Baynard T, Hu L, Evans EM, et al. Sex differences in the relationship between obesity, C-reactive protein, physical activity, depression, sleep quality and fatigue in older adults. *Brain Behav Immun.* 2009;23(5):643-8.

28. Wesselink E, van Baar H, van Zutphen M, Tibosch M, Kouwenhoven EA, Keulen ETP, et al. Inflammation Is a Mediating Factor in the Association between Lifestyle and Fatigue in Colorectal Cancer Patients. *Cancers (Basel)*. 2020;12(12).
29. Wang XS, Williams LA, Krishnan S, Liao Z, Liu P, Mao L, et al. Serum sTNF-R1, IL-6, and the development of fatigue in patients with gastrointestinal cancer undergoing chemoradiation therapy. *Brain Behav Immun*. 2012;26(5):699-705.
30. Himbert C, Ose J, Lin T, Warby CA, Gigic B, Steindorf K, et al. Inflammation- and angiogenesis-related biomarkers are correlated with cancer-related fatigue in colorectal cancer patients: Results from the ColoCare Study. *Eur J Cancer Care (Engl)*. 2019;28(4):e13055.
31. van Roekel EH, Bours MJ, de Brouwer CP, Ten Napel H, Sanduleanu S, Beets GL, et al. The applicability of the international classification of functioning, disability, and health to study lifestyle and quality of life of colorectal cancer survivors. *Cancer Epidemiol Biomarkers Prev*. 2014;23(7):1394-405.
32. Wesselink E, Balvers M, Bours MJL, de Wilt JHW, Witkamp RF, van Baar H, et al. The association between circulating levels of vitamin D and inflammatory markers in the first 2 years after colorectal cancer diagnosis. *Therap Adv Gastroenterol*. 2020;13:1756284820923922.
33. Meyer K, Ueland PM. Targeted quantification of C-reactive protein and cystatin c and its variants by immuno-MALDI-MS. *Anal Chem*. 2014;86(12):5807-14.
34. Dinh KM, Kaspersen KA, Mikkelsen S, Pedersen OB, Petersen MS, Thørner LW, et al. Low-grade inflammation is negatively associated with physical Health-Related Quality of Life in healthy individuals: Results from The Danish Blood Donor Study (DBDS). *PLoS One*. 2019;14(3):e0214468-e.

35. Kushner I, Samols D, Magrey M. A unifying biologic explanation for "high-sensitivity" C-reactive protein and "low-grade" inflammation. *Arthritis Care Res (Hoboken)*. 2010;62(4):442-6.
36. Hopkins MH, Owen J, Ahearn T, Fedirko V, Flanders WD, Jones DP, et al. Effects of supplemental vitamin D and calcium on biomarkers of inflammation in colorectal adenoma patients: a randomized, controlled clinical trial. *Cancer Prev Res (Phila)*. 2011;4(10):1645-54.
37. Vercoulen JH, Swanink CM, Fennis JF, Galama JM, van der Meer JW, Bleijenberg G. Dimensional assessment of chronic fatigue syndrome. *J Psychosom Res*. 1994;38(5):383-92.
38. Aaronson NK, Ahmedzai S, Bergman B, Bullinger M, Cull A, Duez NJ, et al. The European Organization for Research and Treatment of Cancer QLQ-C30: a quality-of-life instrument for use in international clinical trials in oncology. *J Natl Cancer Inst*. 1993;85(5):365-76.
39. Koole JL, Bours MJL, van Roekel EH, Breedveld-Peters J JL, van Duijnhoven FJB, van den Ouweland J, et al. Higher Serum Vitamin D Concentrations Are Longitudinally Associated with Better Global Quality of Life and Less Fatigue in Colorectal Cancer Survivors up to 2 Years after Treatment. *Cancer Epidemiol Biomarkers Prev*. 2020;29(6):1135-44.
40. Beurskens AJ, Bültmann U, Kant I, Vercoulen JH, Bleijenberg G, Swaen GM. Fatigue among working people: validity of a questionnaire measure. *Occup Environ Med*. 2000;57(5):353-7.
41. Worm-Smeitink M, Gielissen M, Bloot L, van Laarhoven HWM, van Engelen BGM, van Riel P, et al. The assessment of fatigue: Psychometric qualities and norms for the Checklist individual strength. *J Psychosom Res*. 2017;98:40-6.
42. Sangha O, Stucki G, Liang MH, Fossel AH, Katz JN. The Self-Administered Comorbidity Questionnaire: a new method to assess comorbidity for clinical and health services research. *Arthritis Rheum*. 2003;49(2):156-63.

43. Wendel-Vos GC, Schuit AJ, Saris WH, Kromhout D. Reproducibility and relative validity of the short questionnaire to assess health-enhancing physical activity. *J Clin Epidemiol.* 2003;56(12):1163-9.
44. Ainsworth BE, Haskell WL, Leon AS, Jacobs DR, Jr., Montoye HJ, Sallis JF, et al. Compendium of physical activities: classification of energy costs of human physical activities. *Med Sci Sports Exerc.* 1993;25(1):71-80.
45. Twisk JWR. *Applied Longitudinal Data Analysis for Epidemiology: A Practical Guide*: Cambridge University Press; 2013.
46. Twisk JWR, de Vente W. Hybrid models were found to be very elegant to disentangle longitudinal within- and between-subject relationships. *J Clin Epidemiol.* 2019;107:66-70.
47. Benjamini Y, Hochberg Y. Controlling the False Discovery Rate: A Practical and Powerful Approach to Multiple Testing. *Journal of the Royal Statistical Society Series B (Methodological).* 1995;57(1):289-300.
48. Imhof A, Fröhlich M, Loewel H, Helbecque N, Woodward M, Amouyel P, et al. Distributions of C-reactive protein measured by high-sensitivity assays in apparently healthy men and women from different populations in Europe. *Clin Chem.* 2003;49(4):669-72.
49. Couper KN, Blount DG, Riley EM. IL-10: The Master Regulator of Immunity to Infection. *The Journal of Immunology.* 2008;180(9):5771-7.
50. Iyer SS, Cheng G. Role of interleukin 10 transcriptional regulation in inflammation and autoimmune disease. *Crit Rev Immunol.* 2012;32(1):23-63.
51. Bower JE, Ganz PA, Tao ML, Hu W, Belin TR, Sepah S, et al. Inflammatory biomarkers and fatigue during radiation therapy for breast and prostate cancer. *Clin Cancer Res.* 2009;15(17):5534-40.
52. Pepys MB, Hirschfield GM. C-reactive protein: a critical update. *The Journal of clinical investigation.* 2003;111(12):1805-12.

53. Lacourt TE, Vichaya EG, Chiu GS, Dantzer R, Heijnen CJ. The High Costs of Low-Grade Inflammation: Persistent Fatigue as a Consequence of Reduced Cellular-Energy Availability and Non-adaptive Energy Expenditure. *Front Behav Neurosci.* 2018;12:78-.
54. Ansar W, Ghosh S. Inflammation and Inflammatory Diseases, Markers, and Mediators: Role of CRP in Some Inflammatory Diseases. *Biology of C Reactive Protein in Health and Disease.* New Delhi: Springer India; 2016. p. 67-107.
55. Ferrucci L, Ble A, Bandinelli S, Lauretani F, Suthers K, Guralnik JM. A flame burning within. *Aging Clin Exp Res.* 2004;16(3):240-3.
56. Sproston NR, Ashworth JJ. Role of C-Reactive Protein at Sites of Inflammation and Infection. *Front Immunol.* 2018;9:754-.
57. Weinhold B, Bader A, Poli V, Rüther U. Interleukin-6 is necessary, but not sufficient, for induction of the human C-reactive protein gene in vivo. *Biochem J.* 1997;325 (Pt 3)(Pt 3):617-21.
58. Ridker PM. From C-Reactive Protein to Interleukin-6 to Interleukin-1: Moving Upstream To Identify Novel Targets for Atheroprotection. *Circ Res.* 2016;118(1):145-56.
59. Rebelo P, Oliveira A, Andrade L, Valente C, Marques A. Minimal Clinically Important Differences for Patient-Reported Outcome Measures of Fatigue in Patients With COPD Following Pulmonary Rehabilitation. *Chest.* 2020;158(2):550-61.
60. Cocks K, King MT, Velikova G, Martyn St-James M, Fayers PM, Brown JM. Evidence-based guidelines for determination of sample size and interpretation of the European Organisation for the Research and Treatment of Cancer Quality of Life Questionnaire Core 30. *J Clin Oncol.* 2011;29(1):89-96.
61. Jayadevappa R, Cook R, Chhatre S. Minimal important difference to infer changes in health-related quality of life - a systematic review. *J Clin Epidemiol.* 2017;89:188-98.

62. Carrero JJ, Andersson Franko M, Obergfell A, Gabrielsen A, Jernberg T. hsCRP Level and the Risk of Death or Recurrent Cardiovascular Events in Patients With Myocardial Infarction: a Healthcare - Based Study. *J Am Heart Assoc.* 2019;8(11):e012638.
63. Salazar J, Martínez MS, Chávez M, Toledo A, Añez R, Torres Y, et al. C-Reactive Protein: Clinical and Epidemiological Perspectives. *Cardiol Res Pract.* 2014;2014:605810.
64. Bogaty P, Dagenais GR, Joseph L, Boyer L, Leblanc A, Bélisle P, et al. Time Variability of C-Reactive Protein: Implications for Clinical Risk Stratification. *PLoS One.* 2013;8(4):e60759.
65. Koc M, Karaarslan O, Abali G, Batur MK. Variation in high-sensitivity C-reactive protein levels over 24 hours in patients with stable coronary artery disease. *Tex Heart Inst J.* 2010;37(1):42-8.

Table 1 – Demographic, lifestyle, and clinical characteristics of stage I to III colorectal cancer survivors at 6 weeks post-treatment, overall and according to sex.

Baseline characteristics	Total population (n = 237)	Men (n=163)	Women (n=74)
Age, mean (SD)	66.8 (9.2)	66.3 (8.8)	68.1 (9.9)
BMI (kg/m ²) ^a , median (IQR)	27.3 (24.4-30.3)	27.3 (24.4-30.4)	27.7 (24.6-29.9)
Use of NSAIDs ^b (yes), n (%)	20 (9.8)	14 (9.8)	6 (9.8)
Physical activity (hours/week), median (IQR)			
LPA	7.5 (2.0-16.5)	6.3 (1.2-12.0)	14.0 (7.0-24.5)
MVPA	7.0 (2.7-14.3)	9.0 (3.5-16.3)	4.1 (1.5-7)
Smoking status (yes), n (%)	22 (9.3)	16 (9.8)	6 (8.1)
Comorbidities n (%)			
0	49 (20.6)	42 (25.8)	7 (9.5)
1	59 (24.8)	43 (26.4)	16 (21.6)
≥ 2	129 (54.4)	78 (47.9)	51 (68.9)
Chemotherapy (yes), n (%)	89 (37.6)	62 (38.0)	27 (36.5)
Radiotherapy (yes), n (%)	64 (27.0)	50 (30.7)	14 (18.9)
Cancer type, n (%)			
Colon cancer	146 (61.6)	94 (57.7)	52 (70.3)
Rectum cancer	91 (38.4)	69 (42.3)	22 (29.7)

^a Data on BMI is missing for 1 person.

^b Thirty-three participants have missing data for use of NSAIDs 6 weeks prior to measurement.

Abbreviations: standard deviation (SD), interquartile range (IQR), non-steroidal anti-inflammatory drug (NSAID), light physical activity (LPA), moderate-to-vigorous physical activity (MVPA).

Table 2 – Fatigue and plasma inflammatory markers in stage I to III colorectal cancer survivors at 6 weeks, 6, 12 and 24 months post-treatment, overall and according to sex.

	Post-treatment follow-up measurements							
	6 weeks n=237 ^a		6 months n=184 ^a		12 months n=150 ^a		24 months n=63 ^a	
	Total Population		Total population		Total population		Total population	
	Men n=163	Women n=74	Men n=123	Women n=61	Men n=104	Women n=46	Men n=44	Women n=19
Checklist Individual Strength (four subscales), mean (SD)								
Total fatigue	62.9 (26.5)		59.9 (27.3)		53.2 (26.3)		53.0 (25.2)	
20-140	61.5 (25.9)	66.0 (27.9)	57.4 (26.6)	65.1 (28.0)	51.2 (25.4)	57.7 (27.9)	47.9 (23.7)	64.7 (25.3)
Subjective fatigue	27.3 (13.4)		25.3 (12.9)		22.3 (12.4)		22.2 (13.1)	
8-56	26.4 (13.3)	29.3 (13.6)	24.1 (12.6)	28.0 (13.2)	21.0 (11.8)	25.4 (13.2)	19.0 (11.9)	29.5 (13.0)
Reduced motivation	12.3 (6.1)		12.1 (6.2)		10.7 (6.1)		10.7 (6.0)	
4-28	11.9 (5.8)	13.0 (6.7)	11.3 (5.7)	13.6 (7.0)	10.1 (5.4)	11.9 (7.5)	9.7 (5.7)	13.1 (6.2)
Reduced physical activity	10.5 (5.2)		9.6 (5.1)		8.4 (4.9)		8.2 (4.9)	
3-21	10.7 (5.0)	10.2 (5.5)	9.6 (5.2)	9.7 (4.8)	8.5 (4.9)	8.0 (4.9)	7.7 (4.9)	9.5 (4.8)
Reduced concentration	12.8 (7.2)		12.9 (7.2)		11.8 (6.7)		11.8 (6.3)	
5-35	12.6 (7.2)	13.4 (7.2)	12.4 (7.3)	13.9 (6.9)	11.6 (6.9)	12.4 (6.3)	11.5 (6.0)	12.6 (7.0)
EORTC QLQ-C30, Fatigue, mean (SD)	29.1 (22.7)		23.6 (22.0)		21.3 (23.6)		20.3 (22.4)	
0-100	28.3 (23.8)	30.9 (20.2)	22.0 (23.6)	27.0 (18.3)	18.2 (22.8)	28.3 (24.3)	14.6 (19.1)	33.3 (24.3)
Inflammatory markers, median (IQR)								
IL6 (pg/ml)	1.5 (0.8-2.2)		1.3 (0.8-2.1)		0.9 (0.6-1.4)		0.9 (0.5-1.5)	
	1.4 (0.8-2.3)	1.5 (0.8-2.1)	1.3 (0.9-2.3)	1.1 (0.8-1.9)	0.9 (0.6-1.5)	0.8 (0.5-1.2)	0.8 (0.4-1.6)	1.0 (0.6-1.3)
IL8 (pg/ml)	5.6 (4.4-7.3)		5.3 (4.4-7.0)		3.9 (3.1-4.8)		4.9 (3.8-7.0)	
	5.5 (4.3-6.8)	5.9 (4.5-8.1)	5.1 (4.4-7.1)	5.5 (4.6-7.0)	3.9 (3.2-4.8)	3.9 (2.9-4.7)	4.8 (3.6-5.9)	5.2 (4.3-7.7)
IL10 (pg/ml)	0.4 (0.3-0.5)		0.4 (0.2-0.5)		0.3 (0.2-0.4)		0.2 (0.1-0.3)	
	0.4 (0.3-0.5)	0.4 (0.3-0.5)	0.4 (0.2-0.5)	0.4 (0.3-0.5)	0.3 (0.2-0.4)	0.2 (0.2-0.3)	0.2 (0.1-0.3)	0.2 (0.1-0.4)
TNFα (pg/ml)	2.9 (2.4-3.8)		2.8 (2.3-3.6)		2.0 (1.6-2.5)		2.0 (1.6-2.9)	
	2.9 (2.4-3.6)	3.0 (2.2-4.0)	2.8 (2.3-3.7)	2.8 (2.3-3.4)	2.0 (1.6-2.4)	2.0 (1.5-2.6)	2.0 (1.6-2.7)	2.3 (1.6-3.4)

Summary inflammatory z-score ^b excluding hsCRP, median (IQR)	-0.3 (-0.7-0.2)		-0.4 (-0.8-0.2)		-0.3 (-0.8-0.3)		-0.5 (-1.2-1.4)	
	-0.3 (-0.7-0.2)	-0.3 (-0.7-0.2)	-0.3 (-0.8-0.3)	-0.4 (-0.8-0.1)	-0.4 (-0.7-0.3)	-0.3 (-0.8-0.1)	-0.6 (-1.2-0.9)	-0.1 (-1.2-1.8)
	Patients with data on hsCRP							
	n=200 ^c		n=148 ^c		n=114 ^c		n=0 ^c	
hsCRP (mg/L), median (IQR)	2.1 (1.1-5.3)		2.1 (0.8-4.6)		1.7 (0.8-5.3)			
	2.0 (1.0-5.5)	2.8 (1.7-5.1)	2.0 (0.8-4.8)	2.2 (0.7-4.4)	1.7 (0.7-5.7)	1.7 (0.8-4.7)		
Summary inflammatory z-score including hsCRP ^d , median (IQR)	-0.6 (-1.0-0.5)		-0.5 (-1.1-0.5)		-0.7 (-1.3-0.5)			
	-0.6 (-1.1-0.5)	-0.6 (-0.9-0.5)	-0.5 (-1.1-0.5)	-0.5 (-1.0-0.5)	-0.7 (-1.3-0.4)	-0.7 (-1.3-0.5)		

^a Participants with data available on fatigue, IL6, IL8, IL10, and TNF α .

^b The inflammatory z-score was calculated as $z = (x - \mu) / \sigma$, in which x is the participant's inflammatory marker value at a given visit, μ is the study population mean, and σ is the study standard deviation, both at given visits. The summary inflammatory z-score for each participant was computed by summing the z-scores of IL6, IL8, TNF α and subtracting IL10.

^c Participants with data available on fatigue, IL6, IL8, IL10, TNF α , and hsCRP.

^d The inflammatory z-score was calculated as $z = (x - \mu) / \sigma$, in which x is the participant's inflammatory marker value at a given visit, μ is the study population mean, and σ is the study standard deviation, both at given visits. The summary inflammatory z-score for each participant was computed by summing the z-scores of IL6, IL8, TNF α , hsCRP and subtracting IL10.

Abbreviations: standard deviation (SD), interquartile range (IQR), interleukin (IL), tumor necrosis factor (TNF), high-sensitivity C-reactive protein (hsCRP), European Organization for the Research and Treatment of Cancer Quality of Life Questionnaire-Core 30 (EORTC QLQ-C30).

Table 3 - Longitudinal associations between inflammatory markers and fatigue in stage I to III colorectal cancer survivors followed-up from 6 weeks to 2 years post-treatment.

	EORTC QLQ-C30, Fatigue β (95%CI)	CIS Total fatigue β (95%CI)	Subjective fatigue β (95%CI)	Reduced motivation β (95%CI)	Reduced physical activity β (95%CI)	Reduced concentration β (95%CI)	
Model I							
IL6	Overall association ^a	1.85 (0.23; 3.47)	2.07 (0.38; 3.76)	0.97 (0.12; 1.83)	0.49 (0.07; 0.90)	0.23 (-0.13; 0.60)	0.48 (-0.00; 0.97)
	Intra-individual ^b	1.50 (-0.35; 3.35)	1.40 (-0.44; 3.24)	0.60 (-0.34; 1.54)	0.35 (-0.11; 0.82)	0.08 (-0.34; 0.50)	0.37 (-0.18; 0.91)
	Inter-individual ^c	3.04 (-0.35; 6.44)	5.44 (1.31; 9.57)	2.67 (0.66; 4.67) *	1.00 (0.09; 1.92)	0.74 (-0.01; 1.50)	0.94 (-0.13; 2.01)
	Model II						
	Overall association ^a	1.69 (0.08; 3.32)	1.67 (-0.03; 3.36)	0.71 (-0.14; 1.56)	0.36 (-0.05; 0.77)	0.12 (-0.24; 0.49)	0.48 (-0.02; 0.98)
	Intra-individual ^b	1.74 (-0.11; 3.60)	1.28 (-0.60; 3.16)	0.46 (-0.49; 1.41)	0.30 (-0.16; 0.77)	0.05 (-0.37; 0.48)	0.44 (-0.13; 1.01)
Inter-individual ^c	1.56 (-1.74; 4.86)	3.34 (-0.56; 7.24)	1.71 (-0.19; 3.61)	0.58 (-0.29; 1.45)	0.32 (-0.40; 1.03)	0.62 (-0.44; 1.68)	
Model I							
IL8	Overall association ^a	1.85 (-0.05; 3.76)	0.08 (-1.95; 2.11)	0.36 (-0.66; 1.39)	-0.24 (-0.73; 0.25)	0.08 (-0.35; 0.51)	-0.06 (-0.64; 0.52)
	Intra-individual ^b	1.55 (-0.81; 3.91)	-0.37 (-2.73; 1.98)	0.18 (-1.02; 1.38)	-0.40 (-0.99; 0.18)	0.13 (-0.40; 0.66)	-0.28 (-0.98; 0.42)
	Inter-individual ^c	2.41 (-0.81; 5.64)	1.40 (-2.62; 5.43)	0.84 (-1.11; 2.79)	0.14 (-0.75; 1.03)	-0.03 (-0.76; 0.70)	0.44 (-0.59; 1.47)
	Model II						
	Overall association ^a	3.34 (0.77; 5.90)	0.74 (-2.10; 3.57)	0.68 (-0.73; 2.08)	-0.18 (-0.84; 0.49)	-0.03 (-0.61; 0.55)	0.37 (-0.44; 1.18)
	Intra-individual ^b	3.68 (0.36; 7.01)	0.25 (-3.21; 3.71)	0.54 (-1.20; 2.28)	-0.43 (-1.27; 0.41)	0.07 (-0.69; 0.83)	0.15 (-0.87; 1.18)
Inter-individual ^c	3.01 (-0.24; 6.26)	1.40 (-2.50; 5.30)	0.84 (-1.06; 2.75)	0.10 (-0.77; 0.97)	-0.12 (-0.84; 0.60)	0.61 (-0.45; 1.67)	
Model I							
IL10	Overall association ^a	-0.12 (-1.87; 1.62)	0.62 (-1.16; 2.41)	0.12 (-0.79; 1.03)	0.13 (-0.31; 0.57)	0.53 (0.14; 0.92)	-0.33 (-0.85; 0.19)
	Intra-individual ^b	0.65 (-1.25; 2.55)	1.41 (-0.48; 3.31)	0.47 (-0.50; 1.44)	0.32 (-0.16; 0.79)	0.77 (0.35; 1.19) *	-0.14 (-0.70; 0.42)
	Inter-individual ^c	-4.31 (-8.69; 0.06)	-5.83 (-11.20; -0.44)	-2.45 (-5.06; 0.17)	-1.04 (-2.24; 0.15)	-0.77 (-1.76; 0.22)	-1.49 (-2.89; -0.10)
	Model II						
	Overall association ^a	-0.07 (-1.80; 1.66)	0.77 (-1.01; 2.54)	0.20 (-0.70; 1.10)	0.14 (-0.30; 0.57)	0.08 (-0.31; 0.48)	-0.29 (-0.82; 0.24)
	Intra-individual ^b	0.76 (-1.09; 2.61)	1.53 (-0.34; 3.39)	0.54 (-0.40; 1.49)	0.33 (-0.13; 0.79)	0.80 (0.03; 1.57)	-0.13 (-0.69; 0.44)
Inter-individual ^c	-5.90 (-10.74; -1.06)	-6.71 (-12.51; -0.92)	-2.95 (-5.79; -0.11)	-1.40 (-2.69; -0.11)	-0.81 (-1.87; 0.25)	-1.57 (-3.14; -0.00)	
Model I							
TNFα	Overall association ^a	0.32 (-1.90; 2.54)	0.43 (-1.89; 2.75)	0.44 (-0.74; 1.61)	0.39 (-0.17; 0.95)	0.07 (-0.42; 0.57)	-0.41 (-1.08; 0.25)
	Intra-individual ^b	-0.05 (-2.61; 2.52)	0.17 (-2.39; 2.73)	0.39 (-0.92; 1.70)	0.27 (-0.37; 0.91)	-0.01 (-0.59; 0.56)	-0.47 (-1.23; 0.29)
	Inter-individual ^c	1.39 (-2.97; 5.76)	1.58 (-3.85; 7.0)	0.63 (-2.00; 3.25)	0.81 (-0.38; 2.01)	0.33 (-0.65; 1.31)	-0.23 (-1.62; 1.16)
	Model II						
	Overall association ^a	-0.34 (-2.52; 1.85)	-0.13 (-2.43; 2.16)	0.16 (-0.99; 1.31)	0.23 (-0.33; 0.78)	-0.08 (-0.58; 0.41)	-0.45 (-1.13; 0.22)
	Intra-individual ^b	0.10 (-2.45; 2.65)	0.20 (-2.37; 2.78)	0.44 (-0.86; 1.74)	0.28 (-0.36; 0.91)	-0.05 (-0.64; 0.53)	-0.42 (-1.19; 0.36)
Inter-individual ^c	-1.56 (-5.84; 2.72)	-1.45 (-6.57; 3.67)	-0.86 (-3.36; 1.64)	0.07 (-1.07; 1.20)	-0.16 (-1.09; 0.77)	-0.56 (-1.94; 0.82)	

	Model I						
	Overall association ^a	2.36	1.13	0.73	0.25	0.03	0.33
	Intra-individual ^b	1.71	-0.07	0.15	-0.05	-0.22	0.05
IZ excluding hsCRP ^d	Inter-individual ^c	4.28	5.89	2.76	1.18	0.68	1.17
	Model II						
	Overall association ^a	2.29	0.74	0.42	0.20	-0.16	0.40
	Intra-individual ^b	2.01	-0.10	0.01	-0.01	-0.31	0.23
	Inter-individual ^c	3.14	3.76	1.79	0.82	0.19	0.89
	Model I						
	Overall association ^a	3.11	3.20	1.75	0.98	0.56	0.14
	Intra-individual ^b	2.37	2.22	1.33	0.77	0.37	-0.25
hsCRP ^e	Inter-individual ^c	5.68	8.25	3.76	1.73	1.18	1.59
	Model II						
	Overall association ^a	2.41	3.21	1.82	0.85	0.38	0.24
	Intra-individual ^b	2.08	2.57	1.62	0.73	0.26	-0.08
	Inter-individual ^c	3.38	5.44	2.53	1.15	0.65	1.10
	Model I						
	Overall association ^a	5.24	2.82	1.72	0.84	0.35	0.33
	Intra-individual ^b	4.06	0.67	0.78	0.30	-0.06	-0.35
IZ including hsCRP ^{d,f}	Inter-individual ^c	8.01	9.71	4.47	2.03	1.20	1.90
	Model II						
	Overall association ^a	4.49	2.42	1.40	0.72	0.04	0.46
	Intra-individual ^b	3.92	0.69	0.68	0.30	-0.27	-0.01
	Inter-individual ^c	5.75	6.71	3.22	1.47	0.53	1.39

Levels of inflammatory markers were divided by their SD at 6 weeks (IL6: SD = 3.15, IL8: SD = 19.57, IL10: SD = 0.86, TNF α : SD = 3.14, IZ excluding hsCRP: SD = 2.08, CRP: SD = 8.19, IZ including hsCRP: SD = 2.62).

Model I: adjusted for age (years), sex (men/women), time since diagnosis (days).

Model II: adjusted for age (years), sex (men/women), time since diagnosis (days), use of NSAIDs (yes/no), BMI (kg/m²), light physical activity (hours/week), moderate-to-vigorous physical activity (hours/week), comorbidities (0, 1, \geq 2), chemotherapy (yes/no), radiotherapy (yes/no), and smoking status (yes/no).

Asterisk (*) represents a significant association after false discovery rate correction for multiple testing.

^a The beta-coefficient represents the overall longitudinal difference in fatigue score per SD difference of the inflammatory marker. It is a weighted average of the intra- and inter-individual associations.

^b The beta-coefficient represents the change in fatigue score over time within-individuals per SD increase of the inflammatory marker.

^c The beta-coefficient represents the difference in fatigue score between-individuals per SD difference of the inflammatory marker.

^d False discovery rate adjustment for multiple testing not performed.

^e hsCRP was measured at 6 weeks, 6, and 12 months post-treatment.

^f Analysis includes patients with data available at 6 weeks, 6, and 12 months post-treatment.

Abbreviations: European Organization for the Research and Treatment of Cancer Quality of Life Questionnaire-Core 30 (EORTC QLQ-C30), Checklist Individual Strength (CIS), beta-coefficient (β), confidence interval (CI), interleukin (IL), tumor necrosis factor (TNF), high-sensitivity C-reactive protein (hsCRP), summary inflammatory z-score (IZ).

Figure Legends

Figure 1. Median levels of inflammatory markers (A, B, C, D, E) and fatigue score (F, G) in stage I to III colorectal cancer survivors from 6 weeks to 24 months post-treatment, overall and according to sex.

Figure 2. Forest plots demonstrating beta coefficients and corresponding 95% CI of overall longitudinal associations, including intra- and inter-individual associations, between inflammatory markers and CIS total fatigue (A) and EORTC QLQ-C30 fatigue (B) in colorectal cancer survivors followed-up at 6 weeks, 6, 12, and 24 months after treatment. Asterisk (*) indicates statistically significant associations after false discovery rate correction for multiple testing.

Figure 3. Overall longitudinal associations between levels of hsCRP increments ($\leq 3\text{mg/L}$, $3\text{-}10\text{mg/L}$, and $>10\text{mg/L}$) with CIS total fatigue (A), EORTC QLQ-C30 fatigue (B), and the CIS subscales – subjective fatigue (C), reduced motivation (D), reduced physical activity level (E), reduced concentration (F), in colorectal cancer survivors followed-up at 6 weeks, 6, and 12 months after treatment. Checklist Individual Strength (CIS) ranges: total fatigue, 20-140; subjective fatigue, 8-56; motivation, 4-28; physical activity level, 3-21; concentration, 5-35. European Organization for the Research and Treatment of Cancer Quality of Life Questionnaire-Core 30 (EORTC QLQ-C30) ranges from 0-100.










Figure 1










Figure 2



Downloaded from <http://aeofjournals.org/cebp/article-pdf/doi/10.1158/1055-9965.EPI-22-0077/3154737/epi-22-0077.pdf> by Maastricht University user on 29 November 2022

Figure 3