

Body Composition, Physical Activity and Quality of Life in end-stage renal disease patients

Citation for published version (APA):

Broers, N. J. H. (2016). *Body Composition, Physical Activity and Quality of Life in end-stage renal disease patients: a phenotypical approach*. [Doctoral Thesis, Maastricht University]. Proefschriftmaken.nl || Uitgeverij BOXPress. <https://doi.org/10.26481/dis.20161103nb>

Document status and date:

Published: 01/01/2016

DOI:

[10.26481/dis.20161103nb](https://doi.org/10.26481/dis.20161103nb)

Document Version:

Publisher's PDF, also known as Version of record

Please check the document version of this publication:

- A submitted manuscript is the version of the article upon submission and before peer-review. There can be important differences between the submitted version and the official published version of record. People interested in the research are advised to contact the author for the final version of the publication, or visit the DOI to the publisher's website.
- The final author version and the galley proof are versions of the publication after peer review.
- The final published version features the final layout of the paper including the volume, issue and page numbers.

[Link to publication](#)

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal.

If the publication is distributed under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license above, please follow below link for the End User Agreement:

www.umlib.nl/taverne-license

Take down policy

If you believe that this document breaches copyright please contact us at:

repository@maastrichtuniversity.nl

providing details and we will investigate your claim.

VALORISATION ADDENDUM

VALORISATION ADDENDUM

Reports showed that at the end of 2013 approximately 3.200.000 patients were treated for ESRD worldwide. More or less than 2.522.000 of these patients were undergoing dialysis. Thereby it was suggested that these numbers will annually increase by 6 to 7 percent¹. In 2010 chronic kidney disease (CKD) was the 18th cause of total number of deaths worldwide, as compared with the 27th cause in 1990². These numbers clearly show the rapid increase of CKD worldwide. Additionally, not only the risk of mortality is high in ESRD patients, also the risk of comorbidity is very high in this patient group and has been shown to have a negatively associated with survival^{3,4}.

Costs for hemodialysis were \$87.945 and peritoneal dialysis were \$71.630 per patient per year respectively in 2011⁵, and it is known that these cost per patient will rise every year. Therefore, dialysis treatment is a high economic burden.

Next to high treatment costs, the burden of disease is high in dialysis patients. Numerous studies showed impaired health-related quality of life (HRQOL) scores in dialysis patients in both physical as well as mental health⁶⁻⁹, which are also directly related to physical activity (PA)^{10,11} and body composition (BC)¹². In addition, results showed strong associations between regular exercise and better HRQOL scores and survival¹⁰, and higher muscle mass and better physical functioning and QOL¹² in dialysis patients. Regardless of these findings, HRQOL is consistently shown to be lower in ESRD patients as compared with the general population^{6,8}, showing the undesirable effects of ESRD in these patients. Thus, even though dialysis is a life-saving treatment, and despite continuous technical developments in dialysis therapy, the burden of kidney disease remains high.

Both these economic and social effects of ESRD stress out the need for interventions that can slow down the progression of CKD and its complications. The current approach for treatment of CKD in the out-patient clinic is mainly based on delaying or halting the progression of renal dysfunction, as well as cardiovascular and metabolic complications. This treatment often consists of blood pressure control, lipid-lowering therapy, dietary advice on salt and protein intake, as well as phosphate control and vitamin D treatment¹³. Additionally, it is well known that exercise on a regular basis is beneficial in relation with outcome in CKD patients¹⁴⁻¹⁶. The guideline for diagnostics and treatment for CKD patients by the Dutch federation of nephrology recommend PA as an intervention goal for CKD patients with mild to moderate loss of kidney function, to reduce the risk of progression of kidney failure and/or cardiovascular complications¹⁷. Also the 'K/DOQI Clinical Practice Guidelines for Cardiovascular Disease in Dialysis Patients', state that PA should be encouraged by nephrologist and dialysis staff to increase PA levels. Next to that, PA and physical functioning should be evaluated every six months¹⁸. Nevertheless, in the current nephrology practice not much attention is addressed to self-management with regard to PA¹⁴, despite the recommendations. So, not implementing stimulation of PA into routine

patient care by nephrologists/dialysis staff, may contribute to low levels of PA and poor physical functioning¹⁴. Furthermore, low levels of PA and poor physical functioning are proven to affect HRQOL and survival¹⁴. All of this is already suggested since 1994, when the Life Options Rehabilitation Advisory Council (LORAC) promoted exercise in their renal rehabilitation report¹⁹. Still, PA programs are not overall implemented in the daily dialysis care, underscoring the need for awareness of this issue in the current renal care.

The lack regarding the promotion of PA is not only a pitfall of the current CKD care. On the one hand, making patients aware, and stimulating them to exercise, is a major advantage for successful rehabilitation of ESRD patients. On the other hand, high costs for PA programs make it difficult to keep patients exercise with the help of professionals. Therefore, one of the important concerns is the support of health insurance to make such programs easier accessible for patients, since ESRD is already a burden for the patient's economic status, due to loss of income and high costs of health care.

Ongoing research already focuses on the long term outcomes of PA programs for dialysis- and transplantation patients and the cost-effectiveness of group rehabilitation. Still, not many interventional studies focus on the earlier stages of CKD. This gives us the opportunity to investigate the effects of PA on parameters such as HRQOL, in CKD patients who are not on dialysis yet. Next to that it would be of great importance to study if the start of dialysis can be delayed by slowing down the progression of CKD in the early stages by stimulating PA.

New technologies, such as activity trackers could facilitate reminders to stay physically active, which may be the future for stimulating PA in patients with chronic diseases. As a recent study already showed, multidimensional PA feedback using graphics and visualizations may be very motivational, and support behavioral changes in patients with increased risk of chronic diseases²⁰.

Knowing that every postponement of starting dialysis, but also the prevention of comorbid diseases is cost effective, and as this thesis showed, what the importance is of PA in ESRD patients, and in line with that its relations with HRQOL, above mentioned interventions may provide beneficial outcomes of PA in CKD patient care, and a decline in the economic burden of kidney disease on the long term. This makes it very important to support research concerning PA, BC and HRQOL in the early CKD stages, but also during the dialysis phase.

REFERENCES

1. Fresenius Medical Care. *ESRD Patients in 2013: A Global Perspective*. 2013 [cited 2016 03-02-2016]; Available from: http://www.vision-fmc.com/files/ESRD_Patients_in_2013.pdf.
2. Lozano R, Naghavi M, Foreman K, Lim S, Shibuya K, Aboyans V, et al. Global and regional mortality from 235 causes of death for 20 age groups in 1990 and 2010: a systematic analysis for the Global Burden of Disease Study 2010. *Lancet* 2012;380:2095-2128.
3. Khan IH, Catto GR, Edward N, Fleming LW, Henderson IS, MacLeod AM. Influence of coexisting disease on survival on renal-replacement therapy. *Lancet* 1993;341:415-418.
4. Gomez AT, Kiberd BA, Royston JP, Alfaadhel T, Soroka SD, Hemmelgarn BR, et al. Comorbidity burden at dialysis initiation and mortality: A cohort study. *Can J Kidney Health Dis* 2015;2:34.
5. US Renal Data System. *USRDS 2013 Annual Data Report. Cost of ESRD*. . 2013 12-04-2016]; Available from: http://www.usrds.org/2013/pdf/v2_ch11_13.pdf.
6. Mazairac AH, de Wit GA, Penne EL, van der Weerd NC, de Jong B, Grooteman MP, et al. Changes in quality of life over time--Dutch haemodialysis patients and general population compared. *Nephrol Dial Transplant* 2011;26:1984-1989.
7. Walters BA, Hays RD, Spritzer KL, Fridman M, Carter WB. Health-related quality of life, depressive symptoms, anemia, and malnutrition at hemodialysis initiation. *Am J Kidney Dis* 2002;40:1185-1194.
8. DeOreo PB. Hemodialysis patient-assessed functional health status predicts continued survival, hospitalization, and dialysis-attendance compliance. *Am J Kidney Dis* 1997;30:204-212.
9. Broers NJ, Usvyat LA, Kooman JP, van der Sande FM, Lacson E, Jr., Kotanko P, et al. Quality of Life in Dialysis Patients: A Retrospective Cohort Study. *Nephron* 2015;130:105-112.
10. Tentori F, Elder SJ, Thumma J, Pisoni RL, Bommer J, Fissell RB, et al. Physical exercise among participants in the Dialysis Outcomes and Practice Patterns Study (DOPPS): correlates and associated outcomes. *Nephrol Dial Transplant* 2010;25:3050-3062.
11. Lopes AA, Lantz B, Morgenstern H, Wang M, Bieber BA, Gillespie BW, et al. Associations of self-reported physical activity types and levels with quality of life, depression symptoms, and mortality in hemodialysis patients: the DOPPS. *Clin J Am Soc Nephrol* 2014;9:1702-1712.
12. Martinson M, Ikizler TA, Morrell G, Wei G, Almeida N, Marcus RL, et al. Associations of body size and body composition with functional ability and quality of life in hemodialysis patients. *Clin J Am Soc Nephrol* 2014;9:1082-1090.
13. Jha V, Garcia-Garcia G, Iseki K, Li Z, Naicker S, Plattner B, et al. Chronic kidney disease: global dimension and perspectives. *Lancet* 2013;382:260-272.
14. Painter P, Roshanravan B. The association of physical activity and physical function with clinical outcomes in adults with chronic kidney disease. *Curr Opin Nephrol Hypertens* 2013;22:615-623.
15. Johansen KL, Painter P. Exercise in individuals with CKD. *Am J Kidney Dis* 2012;59:126-134.
16. Painter P. Physical functioning in end-stage renal disease patients: update 2005. *Hemodial Int* 2005;9:218-235.

17. Nederlandse federatie voor Nefrologie. *Richtlijn Diagnostiek en Behandeling van Patiënten met Chronische Nierschade*. 2015 20-06-2016]; Available from: http://www.nefro.nl/uploads/_I/Mi/_IMi1CJhguCZucn976lhyA/Concept-NfN-richtlijn-Diagnostiek-en-Behandeling-chronische-nierschade-2015.pdf.
18. K/DOQI clinical practice guidelines for cardiovascular disease in dialysis patients. *Am J Kidney Dis* 2005;45:S1-153.
19. Life Options Rehabilitation Advisory Council: *Renal Rehabilitation: Bridging the Barriers*. Madison, WI:Medical Education Institute 1994;
20. Western MJ, Peacock OJ, Stathi A,Thompson D. The understanding and interpretation of innovative technology-enabled multidimensional physical activity feedback in patients at risk of future chronic disease. *PLoS One* 2015;10:e0126156.