

# Intramyocellular lipid content and molecular adaptations in response to a 1-week high-fat diet

## Citation for published version (APA):

Schrauwen-Hinderling, V. B., Kooi, M. E., Hesselink, M. K. C., Kornips, C. F. P., Schaart, G., Mustard, K. J., Hardie, D. G., Saris, W. H. M., Nicolay, K., & Schrauwen, P. (2005). Intramyocellular lipid content and molecular adaptations in response to a 1-week high-fat diet. *Obesity Research*, 13(12), 2088-2094. <https://doi.org/10.1038/oby.2005.259>

## Document status and date:

Published: 01/01/2005

## DOI:

[10.1038/oby.2005.259](https://doi.org/10.1038/oby.2005.259)

## Document Version:

Publisher's PDF, also known as Version of record

## Please check the document version of this publication:

- A submitted manuscript is the version of the article upon submission and before peer-review. There can be important differences between the submitted version and the official published version of record. People interested in the research are advised to contact the author for the final version of the publication, or visit the DOI to the publisher's website.
- The final author version and the galley proof are versions of the publication after peer review.
- The final published version features the final layout of the paper including the volume, issue and page numbers.

[Link to publication](#)

## General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal.

If the publication is distributed under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license above, please follow below link for the End User Agreement:

[www.umlib.nl/taverne-license](http://www.umlib.nl/taverne-license)

## Take down policy

If you believe that this document breaches copyright please contact us at:

[repository@maastrichtuniversity.nl](mailto:repository@maastrichtuniversity.nl)

providing details and we will investigate your claim.

## Intramyocellular Lipid Content and Molecular Adaptations in Response to a 1-Week High-Fat Diet

Vera B. Schrauwen-Hinderling,\*†|| Marianne Eline Kooi,\* Matthijs K.C. Hesselink,†§  
Esther Moonen-Kornips,†‡ Gert Schaart,†§ Kirsty J. Mustard,¶ D. Grahame Hardie,¶ Wim H.M. Saris,†‡  
Klaas Nicolay,|| and Patrick Schrauwen†‡

### Abstract

SCHRAUWEN-HINDERLING, VERA B., MARIANNE ELINE KOOI, MATTHIJS K.C. HESSELINK, ESTHER MOONEN-KORNIPS, GERT SCHAART, KIRSTY J. MUSTARD, D. GRAHAME HARDIE, WIM H.M. SARIS, KLAAS NICOLAY, AND PATRICK SCHRAUWEN. Intramyocellular lipid content and molecular adaptations in response to a 1-week high-fat diet. *Obes Res.* 2005;13:2088–2094.

**Objective:** To investigate molecular adaptations that accompany the elevation of intramyocellular lipid (IMCL) content on a high-fat (HF) diet for 1 week.

**Research Methods and Procedures:** Ten subjects consumed a normal-fat (NF) diet for 1 week, followed by an HF diet for another week. After both dietary periods, we determined the IMCL content by proton magnetic resonance spectroscopy in the vastus lateralis muscle and quantified changes in gene expression, protein content, and activity in biopsy samples. We investigated genes involved in carbohydrate and fatty acid handling [lipoprotein lipase, acetyl-coenzyme A carboxylase (ACC) 2, hormone-sensitive lipase, hexokinase II, and glucose transporter 4] and measured protein levels of CD36 and phosphorylated and un-

phosphorylated ACC2 and the activity of adenosine monophosphate-activated kinase.

**Results:** IMCL content was increased by 54% after the HF period. Lipoprotein lipase mRNA concentration was increased by 33%, whereas ACC2 mRNA concentration tended to be increased after the HF diet. Hexokinase II, glucose transporter 4, and hormone-sensitive lipase mRNA were unchanged after the HF diet. ACC2 and CD36 protein levels, phosphorylation status of ACC2, and adenosine monophosphate-activated kinase activity did not change in response to the HF diet.

**Discussion:** We found that IMCL content in skeletal muscle increased after 1 week of HF feeding, accompanied by molecular adaptations that favor fat storage in muscle rather than oxidation.

**Key words:** human, intramuscular triglycerides, dietary fat, skeletal muscle, magnetic resonance spectroscopy

### Introduction

High-fat (HF)<sup>1</sup> diets, which are widely consumed in western societies, have been related to the development of chronic illnesses and obesity (1). Unlike carbohydrate and protein, fat does not stimulate its own oxidation, and it is well known that the adaptation of fat oxidation is slow (2,3), favoring fat storage when subjects switch from a low-fat to an HF diet. Although the major part of the excessive fat will be stored in white adipose tissue, triglyceride content in skeletal muscle has also been reported to increase after HF feeding (4–8). With proton magnetic resonance spectroscopy;

Received for review March 17, 2005.

Accepted in final form September 27, 2005.

The costs of publication of this article were defrayed, in part, by the payment of page charges. This article must, therefore, be hereby marked "advertisement" in accordance with 18 U.S.C. Section 1734 solely to indicate this fact.

\*Department of Radiology, University Hospital Maastricht, Maastricht, The Netherlands; †Nutrition and Toxicology Research Institute Maastricht, and Departments of ‡Human Biology and §Movement Sciences, Maastricht University, Maastricht, The Netherlands; ¶Division of Molecular Physiology, University of Dundee, Dundee, United Kingdom; and ||Department of Biomedical Engineering, Eindhoven University of Technology, Eindhoven, The Netherlands.

Address correspondence to Vera B. Schrauwen-Hinderling, Department of Radiology, University Hospital Maastricht, P.O. Box 5800, 6202 AZ Maastricht, The Netherlands.

E-mail: vhi@rdia.azm.nl

Copyright © 2006 NAASO

<sup>1</sup> Nonstandard abbreviations: HF, high fat; <sup>1</sup>H-MRS, proton magnetic resonance spectroscopy; IMCL, intramyocellular lipid; VLDL, very low-density lipoprotein; NF, normal fat; FFA, free fatty acid; LPL, lipoprotein lipase; HSL, hormone-sensitive lipase; HKII, hexokinase II; GLUT4, glucose transporter 4; ACC, acetyl-coenzyme A carboxylase; RT, reverse transcription; PCR, polymerase chain reaction; AU, arbitrary unit(s); AMPK, adenosine monophosphate-activated kinase.

copy ( $^1\text{H-MRS}$ ), separation between intramyocellular lipid (IMCL) content and contaminating adipose tissue (extramyocellular lipids) can be made, and IMCL can be quantified in a volume of a few cubic centimeters. Using this method, so far only the very acute effect of HF diets (2 to 3 days) has been investigated (5,6). Here, we aimed to investigate IMCL content in m. vastus lateralis by  $^1\text{H-MRS}$  after 1 week of an HF diet with a macronutrient composition similar to our previous work, in which we determined the effect of an HF diet on fatty acid oxidation (see below) (3).

An increase in IMCL content, as observed on an HF diet, could be simply the consequence of a positive fat balance, due to the slow adaptation of fat oxidation to increased fat intake. However, it is interesting to note that endurance training also increases IMCL content (9), which is not due to a positive fat balance. With training, the oxidation of non-plasma-derived fatty acids [very low-density lipoprotein (VLDL) and/or IMCL] is also increased (10,11), suggesting that the increase in IMCL serves to increase IMCL oxidation. Indeed, some authors found intramuscular triglyceride content to be positively related to IMCL use during exercise (6,7,12,13). Therefore, instead of being simply the result of a positive fat balance, an alternative explanation could be that an increased content of IMCL on an HF diet could lead to increased IMCL breakdown and, therefore, increased availability of intramuscular lipid substrate. In this context, we previously showed that whole-body fat oxidation was increased after 1 week of an HF diet and that the increase in fat oxidation was completely accounted for by an increased oxidation of VLDL and/or IMCL (14).

Therefore, to examine whether the increase in IMCL content on an HF diet is simply due to a positive fat balance or serves to increase fat oxidation, we investigated molecular markers of lipid storage and oxidation in skeletal muscle. We recently investigated the same molecular markers after a short-term endurance training period, which was shown to result in a 42% increase in IMCL content (15) and found a 29% decrease in ACC2 mRNA expression, indicating an improved fat oxidative capacity (V. Schrauwen-Hinderling, unpublished data). In the present study, we quantified IMCL content by  $^1\text{H-MRS}$  after 1 week of normal-fat (NF) diet and 1 week of HF diet, and we investigated mRNA and protein levels and protein activity in muscle biopsies at the end of both dietary periods.

## Research Methods and Procedures

### Subjects

Ten young and healthy male subjects [age (years) (mean  $\pm$  SD), 25.0  $\pm$  6.2] participated in this study. Subjects were recruited by advertisement, and the study was approved by the institutional Medical Ethics Committee. Subjects gave their written informed consent after the nature of the procedure was explained.

### Experimental Protocol

Subjects were provided with the experimental diet for 1 week at a time (NF diet and subsequently HF diet). In the morning of the 7th day, a biopsy was taken in the fasted state, and in the afternoon of the same day, a  $^1\text{H-MRS}$  scan was performed to determine the IMCL content in the m. vastus lateralis. To exclude acute effects of exercise on IMCL content, subjects refrained from exercise the 2 days preceding the  $^1\text{H-MRS}$  measurements.

### Diets

The macronutrient composition of the NF diet was such that 30% of the energy was consumed as fat, 55% as carbohydrate, and 15% as protein. In the HF diet, 60% of the energy was consumed as fat, 25% as carbohydrate, and 15% as protein.

According to the self-reported physical activity level, subjects were given 1.6 to 1.7 times their basic metabolic rate [based on Harris and Benedict equations (16)]. Additionally, subjects received snacks with the same macronutrient composition as the whole diet, which they could eat if needed. The diet consisted of all normal ready-to-use food. Every subject received a personal schedule for every day, listing the provided food items for breakfast, lunch, dinner, and snacks.

### Procedures

**MRS Measurement.** Image-guided localized single-voxel  $^1\text{H-MRS}$  was performed as described earlier (15) in the m. vastus lateralis after the NF and the HF dietary period. MRS measurements were performed on only eight subjects, and for one of them, the spectra were of insufficient quality to be analyzed reliably. Therefore, IMCL content is reported from seven subjects.

**Post-processing.** The spectra were fitted in the time domain using a non-linear least-squares algorithm [AMARES (17)] in the jmrui software package (18) (<http://www.mruui.uab.es/mruui/>) using prior knowledge as described earlier (15). Six peaks were fitted in total, namely three peaks for extramyocellular lipids and three peaks for IMCL. The signals were corrected for T1 and T2 relaxation using the T1 and T2 relaxation times as determined by Schick et al. (19). The corrected area of the  $\text{CH}_2$  peak of IMCL was expressed relative to the area of the corrected water peak. The reproducibility of the quantification of IMCL in the m. vastus lateralis was determined in the setting of another study and reported earlier (15).

**Body Weight.** Body weight was measured in the morning in the fasted state after both dietary periods on a digital scale (E1200; Sauter, Albstadt-Elbingen, Germany) with an accuracy of 0.01 kg.

### Analysis

**Blood Analyses.** For the determination of free fatty acids (FFAs) and triacylglycerol plasma concentrations, blood

was collected in tubes containing 30  $\mu$ l of 0.2 M EDTA. Plasma was immediately centrifuged at high speed, frozen in liquid nitrogen, and stored at  $-80^{\circ}\text{C}$  for later analyses. Triacylglycerol (GPO-trinder 337B; Sigma Diagnostics, St. Louis, MO) and FFA (Wako NEFA-C kit, Wako Chemicals, Neuss, Germany) concentrations were analyzed using the COBAS FARA semiautomatic analyser (Roche, Basel, Switzerland).

**Muscle Biopsy Sampling and Analysis.** Muscle biopsies were taken from the midhigh region of m. vastus lateralis according to the technique of Bergström (20). The biopsy was used for isolation of total RNA using the acid phenol method of Chomczynski and Sacchi (21), with concomitant acid phenol extraction and ethanol precipitation. The mRNA concentrations of lipoprotein lipase (LPL), hormone-sensitive lipase (HSL), hexokinase II (HKII), glucose transporter 4 (GLUT4), and acetyl-coenzyme A carboxylase (ACC) 2 were quantified by reverse transcription (RT)-competitive polymerase chain reaction (PCR) (22). For the assays, the RT reaction was performed from 0.2  $\mu$ g of skeletal muscle total RNA in the presence of a thermostable reverse transcriptase (Tth; Promega, Madison, WI) by use of one of the specific antisense primers. The competitive PCR assays were performed as previously described (23–25). To improve the quantification of the amplified products, fluorescent dye-labeled sense oligonucleotides were used. The PCR products were separated and analyzed on an ALF express DNA sequencer (GE Healthcare, Little Chalfont, Buckinghamshire, United Kingdom) with the Fragment Manager Software. Total RNA preparations and RT-competitive PCR assays of both skeletal muscle samples from the same individual (after NF and HF diets) were performed simultaneously. The mRNA data are reported in arbitrary units (AUs), and the average of the baseline value is set to 1. CD36 protein levels were determined by Western blotting, using a monoclonal antibody (MO25) as previously described (26). Phosphorylated and unphosphorylated ACC2 protein levels were determined as previously described (27). ACC2 protein levels are reported as the ratio of ACC2 to actin, whereas the phosphorylation status of ACC2 is reported as the ratio of phosphorylated to unphosphorylated ACC2 protein. Adenosine monophosphate-activated kinase (AMPK) activity has been investigated as described earlier (27). Total AMPK activity was measured in the tissue lysates by immunoprecipitating with a combination of  $\alpha$  1- and  $\alpha$  2-specific antibodies. A peptide kinase activity could be detected in immunoprecipitates using a variation of the method of Davies et al. (28). Assays were conducted using the AMARAASAAALARRR peptide (29). Activities are presented in units per milligram protein, where unit is defined as nanomoles of phosphate incorporation into the AMARA peptide per minute.

**Statistics.** Results are reported as mean  $\pm$  SE. Statistical analyses were performed with SPSS for Windows 10.0.0




Figure 1: IMCL content after the NF and HF dietary periods ( $n = 7$ ). The IMCL content was significantly higher after the HF diet ( $p = 0.03$ ). \* Significantly different from the NF dietary period.

software (SPSS Inc., Chicago, IL). Differences after the two dietary periods were detected with paired Student's  $t$  tests. Results were considered significant if  $p < 0.05$ .

## Results

### Energy Intake and Body Weight

By design, energy intake during the two dietary periods was identical [ $13.7 \pm 0.7$  MJ/d vs.  $13.7 \pm 0.7$  MJ/d on the NF and the HF diets, respectively (not significant)]. The fat intake was  $29.7 \pm 0.2\%$  and  $60.1 \pm 0.2\%$  of total energy intake on the NF and HF diets, respectively. Body weight was unaffected by the type of diet [ $80.1 \pm 4.4$  and  $79.9 \pm 4.6$  kg after the NF and HF diets, respectively (not significant)].

### IMCL

IMCL content was significantly increased by 53.8% after the HF period, compared with the NF period ( $0.48\% \pm 0.07\%$  of the water resonance after the NF diet vs.  $0.73\% \pm 0.15\%$  after the HF diet, see Figure 1). If values were converted to values in millimoles per kilogram of muscle wet weight as described by Boesch et al. (30) IMCL content was  $6.12 \pm 0.94$  and  $9.4 \pm 1.87$  mmol/kg after the NF and HF diets, respectively.

### mRNA

Concentrations of HKII ( $1.00 \pm 0.24$  vs.  $0.90 \pm 0.20$  AU after NF and HF diets, respectively), Glut4 ( $1.00 \pm 0.32$  vs.  $1.00 \pm 0.25$  AU after NF and HF diets, respectively) and HSL mRNA ( $1.00 \pm 0.26$  vs.  $1.49 \pm 0.48$  AU after NF and HF diets, respectively) were not different after the two dietary periods. LPL mRNA concentration was increased




Figure 2: LPL mRNA concentrations increased significantly after the HF dietary period compared with the NF dietary period ( $p = 0.04$ ). \* Significantly different from the NF dietary period.

significantly by 36% (Figure 2), and ACC2 mRNA tended to be increased by 33% after the HF dietary period ( $1 \pm 0.22$  vs.  $1.33 \pm 0.24$  AU after NF and HF diets, respectively,  $p = 0.08$ ).

#### Protein Levels and Activities

ACC2 protein level (ACC2/actin) was not different after the HF diet ( $1.07 \pm 0.14$  vs.  $1.02 \pm 0.24$  AU after NF and HF diets, respectively). The phosphorylation status (pACC2/ACC2) indicating the activation of the protein did not change either and was  $2.97 \pm 0.47$  vs.  $3.33 \pm 0.50$  AU after the NF and the HF diets, respectively (ratio of the signals obtained with the phosphospecific antibody and streptavidin). AMPK activity was not different after the two dietary periods ( $0.0136 \pm 0.0022$  and  $0.0173 \pm 0.0054$  U/mg after NF and HF diets, respectively). The CD 36 protein level was not different after the two dietary periods ( $88 \pm 11$  AU after NF diet vs.  $87 \pm 12$  AU after HF diet).

#### Plasma Parameters

Fasting FFA concentrations increased after the HF diet compared with the NF diet, but there was no difference in triacylglycerol concentration after the two diets (see Table 1).

### Discussion

In contrast to carbohydrates, the intake of fat does not acutely stimulate its own oxidation, and consumption of an HF diet often leads to a positive fat balance. In addition to white adipose tissue, fat can also be stored in non-adipose tissues, such as heart, liver, and muscle. Here, we show that after 1 week of HF feeding, IMCL content of the vastus

**Table 1.** Plasma concentrations of FFAs and triacylglycerols

	Normal-fat diet	HF diet
FFAs ( $\mu\text{M}$ )	$251 \pm 42$	$334 \pm 54^*$
Triacylglycerols ( $\mu\text{M}$ )	$961 \pm 136$	$733 \pm 175$

FFA, free fatty acid; HF, high fat.  
\* Significantly different from normal-fat diet ( $p = 0.02$ ).

lateralis muscle was significantly increased. It can be suggested that increased IMCL content on an HF diet is simply the result of the positive fat balance that inevitably occurs when switching to an HF diet. Alternatively, it can be suggested that the increased IMCL content on the HF diet could lead to an increase in fat oxidation. Therefore, we examined in the present study the expression of genes involved in lipid handling in skeletal muscle and protein content and activity in response to an HF diet of 1 week.

The reported increase in IMCL content in the present study (54%) is well in line with earlier results in a range of muscles and with different methodologies [36% to 57% increase in IMCL content after 2 to 3 days of an HF diet (5–7) and 86% to 130% increase after 4 to 5 weeks (8,31)]. When the increased IMCL content on an HF diet serves to allow increased IMCL oxidation, an up-regulation of HSL would be anticipated. HSL is necessary for the hydrolysis of intramuscular lipid to allow mitochondrial uptake and subsequent oxidation (32,33). To the best of our knowledge, no data are available on the effect of HF feeding on HSL mRNA expression or activity in human skeletal muscle. Presently, we found that HF feeding did not affect skeletal muscle HSL mRNA expression.

Furthermore, we examined the effect of HF feeding on ACC2 mRNA and protein expression, as well as ACC2 phosphorylation and AMPK activity. ACC2 has been reported to play a critical role in regulating fat oxidation in skeletal muscle. Its unphosphorylated form catalyzes the carboxylation of acetyl-coenzyme A to form malonyl-CoA, an intermediate that inhibits the activity of carnitine-palmitoyl-transferase-1. Carnitine-palmitoyl-transferase-1 catalyzes the rate limiting step in the transfer of fatty acids into mitochondria, where they undergo oxidation. Therefore, a reduction in ACC2 or an increase in the phosphorylation of the protein would increase fat oxidation. Indeed, it was shown that ACC2 knockout mice are characterized by a strongly increased fat oxidation (34). Similarly, fat oxidation is stimulated when the activity of AMPK, which phosphorylates ACC2, increases. In humans, we have previously shown that a 3-month training program resulted in an increase in triacylglycerol (IMCL and/or VLDL)-derived fat oxidation and a reduction in ACC2 mRNA expression (11).

Therefore, when the increased IMCL content on the HF diet would serve to allow increased fat oxidation, we anticipated finding a reduction in ACC2 mRNA and protein level and/or increased phosphorylation of the protein after 1 week of HF feeding. In the current study, however, we report a tendency ( $p = 0.08$ ) to increased ACC2 mRNA concentrations and no change in protein level, rather than the anticipated decrease. Also, AMPK activity and the phosphorylation of its downstream target, ACC2, were unaffected after the HF dietary period. This again suggests that it is unlikely that the increase in IMCL content on an HF diet serves to increase fat oxidation but points toward a role for IMCL in storing excessive fat when consuming an HF diet. This might explain the negative effect of HF diet on insulin sensitivity (35) because IMCL content has been positively associated with the development of insulin resistance (36) and is increased in patients with type 2 diabetes (37,38). However, we have recently shown that 2 weeks of endurance training resulted in a 42% increase in IMCL content, without any detrimental effect on insulin sensitivity (15). Interestingly, the increase in IMCL content was accompanied by a down-regulation of ACC2 mRNA expression (V. Schrauwen-Hinderling, unpublished data). This points toward an important role for ACC2 in determining the effect of IMCL on insulin sensitivity and suggests that high IMCL content does not affect insulin sensitivity as long as it is combined with improved fat oxidative capacity, as was also recently suggested by Goodpaster et al. (37). Interestingly, it was recently shown that HF feeding of wild-type mice resulted in the development of a diabetic phenotype, whereas ACC2 knockout mice were resistant against HF diet-induced obesity and diabetes (34). Therefore, ACC2 seems to be an important target in the treatment of type 2 diabetes and obesity.

An alternative source for fat oxidation could be VLDL. Helge et al. (4) showed that on an HF diet, VLDL oxidation was significantly increased during aerobic exercise. Triacylglycerols from VLDL particles have to be hydrolyzed by LPL located on the endothelial cells lining the blood vessels to be taken up by the skeletal muscles. Because the availability of triacylglycerols increases on an HF diet, a feasible response would be an increased expression of LPL. Indeed, we found the LPL mRNA concentration to be increased by 36% after the HF diet. The increase in LPL suggests that more fatty acids from VLDL are taken up into myocytes.

Although FFA can enter the cells by passive diffusion (39), facilitated diffusion has been described, and proteins with a high fatty acid binding affinity, such as CD36, have been characterized (40). It has been shown that increased fatty acid uptake may result in increased lipid accumulation in muscle (41). Therefore, we hypothesized that after the HF diet, the increased fat availability would result in increased levels of CD36, a key fatty acid translocase. We did, however, not find changes in the protein level of CD36 after

increased dietary fat. In contrast, in trained athletes, CD36 was increased after an HF diet (42). Because training profoundly improves fat oxidative capacity and, hence, warrants increased trans-sarcolemmal transport of fatty acids, the discrepancy could be due to the differing training status of the subjects in the two studies.

To investigate whether 1 week of an HF diet affected genes involved in glucose metabolism, we investigated markers of carbohydrate metabolism (HKII and GLUT4). However, these markers did not change in response to the HF diet. Because the time course of the adaptations to an HF diet of the genes under investigation is presently unknown, it cannot be excluded that transient changes in mRNA might have been missed in the present investigation. However, the protein data reported here are in line with the lack of effect on fat oxidative capacity after the 1-week dietary intervention.

In conclusion, we investigated the adaptive response to an HF diet, which is currently associated with an unhealthy lifestyle. We found that IMCL content was increased after 1 week of HF feeding, which was not accompanied by molecular adaptations that improve fat oxidative capacity.

### Acknowledgments

P.S. was supported by a grant from the Royal Netherlands Academy of Arts and Sciences. K.J.M. and D.G.H. were aided by the pharmaceutical companies supporting the Division of Signal Transduction Therapy Unit at the University of Dundee (AstraZeneca, Boehringer-Ingelheim, Glaxo-SmithKline, Merck & Co. Inc, Merck KgaA, and Pfizer). We thank Niels Boon for practical help with the study and J. Glatz and N.N. Tandon for kindly providing the MO25 antibody. We also thank H. Vidal (Institut National de la Santé et de la Recherche Médicale U.449, R. Laennec Faculty of Medicine Claude Bernard, University of Lyon, Lyon, France) for kindly providing the competitors. The MRUI software package was kindly provided by the participants of the EU Network programs: Human Capital and Mobility, CHRX-CT94-0432, and Training and Mobility of Researchers, ERB-FMRX-CT970160.

### References

1. Hill JO, Peters JC. Environmental contributions to the obesity epidemic. *Science*. 1998;280:1371–4.
2. Hill JO, Peters JC, Reed GW, et al. Nutrient balance in humans: effects of diet composition. *Am J Clin Nutr*. 1991; 54:10–7.
3. Schrauwen P, van Marken Lichtenbelt WD, Saris WH, Westerterp KR. Changes in fat oxidation in response to a high-fat diet. *Am J Clin Nutr*. 1997;66:276–82.
4. Helge JW, Watt PW, Richter EA, Rennie MJ, Kiens B. Fat utilization during exercise: adaptation to a fat-rich diet increases utilization of plasma fatty acids and very low density lipoprotein-triacylglycerol in humans. *J Physiol*. 2001;537: 1009–20.

5. **Bachmann OP, Dahl DB, Brechtel K, et al.** Effects of intravenous and dietary lipid challenge on intramyocellular lipid content and the relation with insulin sensitivity in humans. *Diabetes*. 2001;50:2579–84.
6. **Johnson NA, Stannard SR, Mehalski K, et al.** Intramyocellular triacylglycerol in prolonged cycling with high- and low-carbohydrate availability. *J Appl Physiol*. 2003;94:1365–72.
7. **Zderic TW, Davidson CJ, Schenk S, Byerley LO, Coyle EF.** High-fat diet elevates resting intramuscular triglyceride concentration and whole-body lipolysis during exercise. *Am J Physiol Endocrinol Metab*. 2004;86:E217–25.
8. **Vogt M, Puntchart A, Howald H, et al.** Effects of dietary fat on muscle substrates, metabolism, and performance in athletes. *Med Sci Sports Exerc*. 2003;35:952–60.
9. **Hoppeler H.** Exercise-induced ultrastructural changes in skeletal muscle. *Int J Sports Med*. 1986;7:187–204.
10. **Phillips SM, Green HJ, Tarnopolsky MA, et al.** Effects of training duration on substrate turnover and oxidation during exercise. *J Appl Physiol*. 1996;81:2182–91.
11. **Schrauwen P, van Aggel-Leijssen DP, Hul G, et al.** The effect of a 3-month low-intensity endurance training program on fat oxidation and acetyl-CoA carboxylase-2 expression. *Diabetes*. 2002;51:2220–6.
12. **Standl E, Lotz N, Dixel T, Janka HU, Kolb HJ.** Muscle triglycerides in diabetic subjects: effect of insulin deficiency and exercise. *Diabetologia*. 1980;18:463–9.
13. **Steffensen CH, Roepstorff C, Madsen M, Kiens B.** Myocellular triacylglycerol breakdown in females but not in males during exercise. *Am J Physiol Endocrinol Metab*. 2002;282:E634–42.
14. **Schrauwen P, Wagenmakers AJ, van Marken Lichtenbelt WD, Saris WH, Westerterp KR.** Increase in fat oxidation on a high-fat diet is accompanied by an increase in triglyceride-derived fatty acid oxidation. *Diabetes*. 2000;49:640–6.
15. **Schrauwen-Hinderling VB, Schrauwen P, Hesselink MK, et al.** The increase in intramyocellular lipid content is a very early response to training. *J Clin Endocrinol Metab*. 2003;88:1610–6.
16. **Harris JA, Benedict FG.** *A Biometric Study of Basal Metabolism in Man*. Washington, DC: Carnegie Institution of Washington; 1919.
17. **Vanhamme L, van den Boogaart A, Van Huffel S.** Improved method for accurate and efficient quantification of MRS data with use of prior knowledge. *J Magn Reson*. 1997;129:35–43.
18. **Naressi A, Couturier C, Devos JM, et al.** Java-based graphical user interface for the MRUI quantitation package. *Magma*. 2001;12:141–52.
19. **Schick F, Eismann B, Jung WI, et al.** Comparison of localized proton NMR signals of skeletal muscle and fat tissue in vivo: two lipid compartments in muscle tissue. *Magn Reson Med*. 1993;29:158–67.
20. **Bergstrom J, Hermansen L, Hultman E, Saltin B.** Diet, muscle glycogen and physical performance. *Acta Physiol Scand*. 1967;71:140–50.
21. **Chomczynski P, Sacchi N.** Single-step method of RNA isolation by acid guanidinium thiocyanate-phenol-chloroform extraction. *Anal Biochem*. 1987;162:156–9.
22. **Auboeuf D, Vidal H.** The use of the reverse transcription-competitive polymerase chain reaction to investigate the in vivo regulation of gene expression in small tissue samples. *Anal Biochem*. 1997;245:141–8.
23. **Laville M, Auboeuf D, Khalfallah Y, et al.** Acute regulation by insulin of phosphatidylinositol-3-kinase, Rad, Glut 4, and lipoprotein lipase mRNA levels in human muscle. *J Clin Invest*. 1996;98:43–9.
24. **Millet L, Vidal H, Andreelli F, et al.** Increased uncoupling protein-2 and -3 mRNA expression during fasting in obese and lean humans. *J Clin Invest*. 1997;100:2665–70.
25. **Andreelli F, Laville M, Vega N, Riou JP, Vidal H.** Regulation of gene expression during severe caloric restriction: lack of induction of p85 alpha phosphatidylinositol 3-kinase mRNA in skeletal muscle of patients with type II (non-insulin-dependent) diabetes mellitus. *Diabetologia*. 2000;43:356–63.
26. **Keizer HA, Schaart G, Tandon NN, Glatz JF, Luiken JJ.** Subcellular immunolocalisation of fatty acid translocase (FAT)/CD36 in human type-1 and type-2 skeletal muscle fibres. *Histochem Cell Biol*. 2004;121:101–7.
27. **Hawley SA, Boudeau J, Reid JL, et al.** Complexes between the LKB1 tumor suppressor, STRAD alpha/beta and MO25 alpha/beta are upstream kinases in the AMP-activated protein kinase cascade. *J Biol*. 2003;2:28.
28. **Davies SP, Carling D, Hardie DG.** Tissue distribution of the AMP-activated protein kinase, and lack of activation by cyclic-AMP-dependent protein kinase, studied using a specific and sensitive peptide assay. *Eur J Biochem*. 1989;186:123–8.
29. **Dale S, Wilson WA, Edelman AM, Hardie DG.** Similar substrate recognition motifs for mammalian AMP-activated protein kinase, higher plant HMG-CoA reductase kinase-A, yeast SNF1, and mammalian calmodulin-dependent protein kinase I. *FEBS Lett*. 1995;361:191–5.
30. **Boesch C, Slotboom J, Hoppeler H, Kreis R.** In vivo determination of intra-myocellular lipids in human muscle by means of localized 1H-MR-spectroscopy. *Magn Reson Med*. 1997;37:484–93.
31. **Helge JW, Wulff B, Kiens B.** Impact of a fat-rich diet on endurance in man: role of the dietary period. *Med Sci Sports Exerc*. 1998;30:456–61.
32. **Langfort J, Ploug T, Ihlemann J, et al.** Expression of hormone-sensitive lipase and its regulation by adrenaline in skeletal muscle. *Biochem J*. 1999;340:459–65.
33. **Oscari LB, Essig DA, Palmer WK.** Lipase regulation of muscle triglyceride hydrolysis. *J Appl Physiol*. 1990;69:1571–7.
34. **Abu-Elheiga L, Oh W, Kordari P, Wakil SJ.** Acetyl-CoA carboxylase 2 mutant mice are protected against obesity and diabetes induced by high-fat/high-carbohydrate diets. *Proc Natl Acad Sci USA*. 2003;100:10207–12.
35. **Lovejoy JC.** The influence of dietary fat on insulin resistance. *Curr Diab Rep*. 2002;2:435–40.
36. **Jacob S, Machann J, Rett K, et al.** Association of increased intramyocellular lipid content with insulin resistance in lean non-diabetic offspring of type 2 diabetic subjects. *Diabetes*. 1999;48:1113–9.

37. **Goodpaster BH, He J, Watkins S, Kelley DE.** Skeletal muscle lipid content and insulin resistance: evidence for a paradox in endurance-trained athletes. *J Clin Endocrinol Metab.* 2001;86:5755–61.
38. **Levin K, Daa Schroeder H, Alford FP, Beck-Nielsen H.** Morphometric documentation of abnormal intramyocellular fat storage and reduced glycogen in obese patients with Type II diabetes. *Diabetologia.* 2001;44:824–33.
39. **Hamilton JA, Kamp F.** How are free fatty acids transported in membranes? Is it by proteins or by free diffusion through the lipids? *Diabetes.* 1999;48:2255–69.
40. **Abumrad NA, el-Maghrabi MR, Amri EZ, Lopez E, Grimaldi PA.** Cloning of a rat adipocyte membrane protein implicated in binding or transport of long-chain fatty acids that is induced during preadipocyte differentiation: homology with human CD36. *J Biol Chem.* 1993;268:17665–8.
41. **Hegarty BD, Cooney GJ, Kraegen EW, Furler SM.** Increased efficiency of fatty acid uptake contributes to lipid accumulation in skeletal muscle of high fat-fed insulin-resistant rats. *Diabetes.* 2002;51:1477–84.
42. **Cameron-Smith D, Burke LM, Angus DJ, et al.** A short-term, high-fat diet up-regulates lipid metabolism and gene expression in human skeletal muscle. *Am J Clin Nutr.* 2003; 77:313–8.