

On the nucleolus of neighbor games

Citation for published version (APA):

Hamers, H., Klijn, F., Solymosi, T., Nakabeppu, Y., & Vermeulen, A. J. (2003). On the nucleolus of neighbor games. *European Journal of Operational Research*, 146, 1-18. [https://doi.org/10.1016/S0377-2217\(02\)00240-0](https://doi.org/10.1016/S0377-2217(02)00240-0)

Document status and date:

Published: 01/01/2003

DOI:

[10.1016/S0377-2217\(02\)00240-0](https://doi.org/10.1016/S0377-2217(02)00240-0)

Document Version:

Publisher's PDF, also known as Version of record

Please check the document version of this publication:

- A submitted manuscript is the version of the article upon submission and before peer-review. There can be important differences between the submitted version and the official published version of record. People interested in the research are advised to contact the author for the final version of the publication, or visit the DOI to the publisher's website.
- The final author version and the galley proof are versions of the publication after peer review.
- The final published version features the final layout of the paper including the volume, issue and page numbers.

[Link to publication](#)

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal.

If the publication is distributed under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license above, please follow below link for the End User Agreement:

www.umlib.nl/taverne-license

Take down policy

If you believe that this document breaches copyright please contact us at:

repository@maastrichtuniversity.nl

providing details and we will investigate your claim.



Stochastics and Statistics

On the nucleolus of neighbor games

Herbert Hamers ^a, Flip Klijn ^{b,*}, Tamás Solymosi ^{c,1}, Stef Tijs ^a,
Dries Vermeulen ^d

^a CentER and the Department of Econometrics and Operations Research, Tilburg University, Tilburg, Netherlands

^b CODE and Departament d'Economia i d'Història Econòmica, Universitat Autònoma de Barcelona, Barcelona, Spain

^c Department of Operations Research, Budapest University of Economic Sciences and Public Administration, Budapest, Hungary

^d Department of Quantitative Economics, University of Maastricht, Maastricht, Netherlands

Received 19 September 2001; accepted 7 February 2002

Abstract

Assignment problems are well-known problems in practice. We mention house markets, job markets, and production planning. The games of interest in this paper, the neighbor games, arise from a special class of assignment problems. We focus on the nucleolus [D. Schmeidler, *SIAM J. Appl. Math.* 17 (1969) 1163–1170], one of the most prominent core solutions. A core solution is interesting with respect to neighbor games because it divides the profit of an optimal matching in a stable manner. This paper establishes a polynomial bounded algorithm of quadratic order in the number of players for calculating the nucleolus of neighbor games.

© 2002 Elsevier Science B.V. All rights reserved.

Keywords: Game theory; Neighbor games; Assignment games; Nucleolus

1. Introduction

Neighbor games, which were introduced in Klijn et al. (1999), form the intersection of assignment games (cf. Shapley and Shubik, 1972) and component additive games (cf. Curiel et al., 1994). This property implies that many problems can be analyzed using neighbor games. The following two examples describe situations that can be handled by neighbor games.

In the first example we consider a sequencing situation in which customers are lined up in a queue and waiting for a taxi. The taxi company that provides the service has two types of cars: one that transports only one customer (type A) and one that can only transport two customers (type B). The first customer in

* Corresponding author. Tel.: +34-93-581-1720; fax: +34-93-581-2012. Part of this research was carried out while this author was Ph.D. student at CentER and the Department of Econometrics and Operations Research, Tilburg University, Netherlands.

E-mail address: fklijn@pareto.uab.es (F. Klijn).

¹ This author's work has been supported by CentER and the Department of Econometrics and Operations Research, Tilburg University, Netherlands. Grants from AMFK and OTKA (T030945) are also gratefully acknowledged.

the queue can decide to pick a taxi of type A or wait for the next customer in the queue. In the latter case they decide both to share a taxi of type B or the second customer will wait for the third customer. In the latter case the first customer has to pick a taxi of type A. This procedure is repeated until all customers are transported in a taxi. Since the costs of sharing a taxi of type B are lower than taking two taxis of type A, it is obvious that the customers can save costs by sharing a taxi of type B. However, each customer faces the problem that the cost of a taxi (of type B) is not fixed, because it depends on the trip to bring the customers to the right locations. Hence, we have that only customers that are neighbors in the queue can obtain cost savings, and customers that take a taxi of type A have cost savings equal to zero. All customers in the queue want to choose a combination of taxis of type A and B such that their cost savings are maximized. Moreover, they are looking for an allocation of the cost savings that is ‘stable’.

The second example can be viewed as a restricted matching problem. Suppose a river runs through a number of regions. To be able to utilize this cheap transportation possibility, harbors have to be built. Because of financial restrictions, each country is able to build at most one harbor. Neighbor regions might join to build a harbor at their border (which then can serve both regions) and save costs. The regions are interested in maximizing their cost savings and finding some proper allocation of the cost savings.

Because of the mentioned intersection property and the fact that component additive games are Γ -component additive games (cf. Potters and Reijnierse, 1995) neighbor games have many appealing properties: the core is a non-empty set and coincides with the set of competitive equilibria (Shapley and Shubik, 1972), the core coincides with the bargaining set, and the nucleolus coincides with the kernel (Potters and Reijnierse, 1995). Moreover, neighbor games satisfy the CoMa-property, i.e., the core is the convex hull of all marginal vectors that are in the core (cf. Hamers et al., 2002).

This paper provides an $\mathcal{O}(p^2)$ algorithm for calculating the nucleolus (Schmeidler, 1969) of p -person neighbor games. In literature, the computation of the nucleolus of assignment games and component additive games has been discussed extensively. Solymosi and Raghavan (1994) presented an $\mathcal{O}(p^4)$ algorithm for calculating the nucleolus of $p + p$ -person assignment games. An $\mathcal{O}(p^4)$ algorithm for calculating the nucleolus of p -person balanced connected games was provided by Solymosi et al. (1998). The class of balanced connected games contains the class of component additive games, and thus the class of neighbor games. This paper provides an $\mathcal{O}(p^2)$ algorithm for calculating the nucleolus of p -person neighbor games. Although the algorithm can be considered as a common specialization of the two algorithms mentioned above, we present it on its own right, since it exhibits special features that neither of the two more general algorithms does. Besides, we give a different line of arguments to see the correctness of the algorithm from those which were used to justify the mentioned more general algorithms.

In Section 2 we provide some preliminaries on cooperative games. Then, in Section 3 we recall the definition of neighbor games and present an $\mathcal{O}(p^2)$ algorithm for finding the nucleolus.

2. Preliminaries

A *cooperative game with transferable utilities* (or game, for short) is a pair (P, v) where $P = \{1, \dots, p\}$ is a finite set of players and $v : 2^P \rightarrow \mathbb{R}$ is a map that assigns to each coalition $S \in 2^P$ a real number $v(S)$, such that $v(\emptyset) = 0$. Here, 2^P is the collection of all subsets (coalitions) of P .

Let (P, v) be a game with a non-empty imputation set $I(P, v) := \{x \in \mathbb{R}^P : x_i \geq v(i) \text{ for all } i \in P \text{ and } x(P) = v(P)\}$, where $x(P) := \sum_{i \in P} x_i$. For an imputation $x \in I(P, v)$ and a coalition $S \in 2^P \setminus \{\emptyset\}$ we call $f(S, x) := x(S) - v(S)$ the *satisfaction* of S . Next, let $F(x) := (f(S, x))_{\emptyset \neq S \subseteq N}$ be the vector of satisfactions and let $\theta(F(x))$ denote the vector of satisfactions with its elements arranged in non-decreasing order, i.e., $\theta(F(x))_1 \leq \theta(F(x))_2 \leq \dots \leq \theta(F(x))_{|2^P|-1}$. The *nucleolus* (Schmeidler, 1969) is then defined by

$$n(P, v) := \{x \in I(P, v) : \theta(F(x)) \succeq_{\text{lex}} \theta(F(y)) \text{ for all } y \in I(P, v)\},$$

where \succeq_{lex} denotes the lexicographical ordering on $\mathbb{R}^{|2^P|-1}$. Recall that for two vectors $x, y \in \mathbb{R}^{|2^P|-1}$ we have $x \succeq_{\text{lex}} y$ if either $x = y$ or there exists a k such that $x_i = y_i$ for $i = 1, \dots, k$ and $x_{k+1} > y_{k+1}$. Schmeidler (1969) proved that the nucleolus $n(P, v)$ is a singleton. For the sake of convenience, we identify $n(P, v)$ with its unique element.

The nucleolus is an element of the core, whenever the latter is not empty. The core of a game (P, v) consists of all vectors that distribute the gains $v(P)$ obtained by P among the players in such a way that no subset of players can be better off by seceding from the rest of the players and act on their own behalf. Formally, the *core* of a game (P, v) is defined by

$$\text{Core}(P, v) := \{x \in \mathbb{R}^P : f(S, x) \geq 0 \text{ for all } S \subset P \text{ and } f(P, x) = 0\}. \tag{1}$$

A coalition $S \neq \emptyset$ is called *essential* in the game (P, v) if $S = S_1 \cup S_2, S_1 \cap S_2 = \emptyset$, and $S_1 \neq \emptyset \neq S_2$ imply that $v(S) > v(S_1) + v(S_2)$. Otherwise, it is called *inessential*. Note that one-player coalitions are always essential. Note also that the inequality related to an inessential coalition in the definition of the core is redundant, i.e., it can be left out without enlarging the solution set of the remaining inequalities. Therefore, for any collection $\mathcal{G} \subseteq 2^P \setminus \{\emptyset\}$ that contains all essential coalitions in the game (P, v) we can rewrite (1):

$$\text{Core}(P, v) = \{x \in \mathbb{R}^P : f(S, x) \geq 0 \text{ for all } S \in \mathcal{G} \text{ and } f(P, x) = 0\}.$$

Huberman (1980) showed that inessential coalitions can also be omitted in the determination of the nucleolus provided the core of the game is not empty. Since in that case the nucleolus lies in the core, the underlying payoff set can be reduced to the core. More precisely,

$$n(P, v) = \{x \in \text{Core}(P, v) : \theta(G(x)) \succeq_{\text{lex}} \theta(G(y)) \text{ for all } y \in \text{Core}(P, v)\},$$

where $G(x) := (f(S, x))_{S \in \mathcal{G}}$ is the vector of satisfactions of coalitions in G only.

Let (P, v) be a game with a non-empty core. Let $\mathcal{G} \subseteq 2^P \setminus \{\emptyset\}$ contain all essential coalitions in the game. We need to introduce some notions and notation to be able to describe a simplified version of Kohlberg’s (1971) criterion for the nucleolus of such a game. We call a non-empty collection \mathcal{B} of coalitions in P balanced if there are positive numbers $(\lambda_S)_{S \in \mathcal{B}}$ such that $\sum_{S \in \mathcal{B}} \lambda_S e_S = e_P$, where e_S is the vector in \mathbb{R}^P with $(e_S)_i = 1$ if and only if $i \in S$ and 0 otherwise. Given a number $t \geq 0$ and an allocation $x \in \text{Core}(P, v)$ we define

$$G(x, t) := \{S \in \mathcal{G} : f(S, x) \leq t\}$$

to be the collection of all coalitions in \mathcal{G} whose satisfaction is not more than the given level t at the given core allocation x . In light of Huberman’s (1980) simplification it is easy to see that in this setting Kohlberg’s (1971) general criterion can be replaced by the following characterization of the nucleolus.

Lemma 2.1. *Let (P, v) be a game with a non-empty core. Let $\mathcal{G} \subseteq 2^P \setminus \{\emptyset\}$ contain all essential coalitions in (P, v) . Then for $x \in \text{Core}(P, v)$ it holds that $\{x\} = n(P, v)$ if and only if $G(x, t)$ is balanced for all $t \geq 0$.*

3. Neighbor games and the nucleolus

In this section we provide a polynomially bounded algorithm of order p^2 for finding the nucleolus of neighbor games. We present the algorithm of order p^2 for finding the nucleolus of a special subclass of neighbor games. After that, we show that we can calculate the nucleolus of an arbitrary neighbor game by breaking up the game in appropriate subgames, applying the algorithm to the subgames, and combining the nucleoli of the subgames. Moreover, we prove that this procedure does not change the computational complexity. Before we present the algorithm we recall the definition and some special features of neighbor games.

Formally, let P be the player set of size p . For the sake of convenience we assume² that $P = \{1, \dots, p\}$. Without loss of generality we assume that the players are ordered $1 \prec 2 \prec \dots \prec p$. Players i and j are called *neighbors* if $|i - j| = 1$. A *matching* μ for $Q \subseteq P$ is a (possibly empty) collection of disjoint pairs $(i, i + 1)$ of neighboring players (*partners*) in Q . Let $\mathcal{N}(Q)$ denote the set of matchings for Q . For all pairs of neighbors $(i, i + 1)$ let $a_{ii+1} \geq 0$ be given. Then, a *neighbor game* (P, v) is defined by

$$v(Q) := \max \left\{ \sum_{(i,i+1) \in \mu} a_{ii+1} : \mu \in \mathcal{N}(Q) \right\} \text{ for all } Q \subseteq P.$$

Note that since $a_{ii+1} = v(i, i + 1)$ a neighbor game is completely determined by the values of the pairs of neighbors. Note also that $v(i) = 0$ for all $i \in P$. A matching $\mu \in \mathcal{N}(Q)$ is called *optimal* for Q if $\sum_{(i,i+1) \in \mu} a_{ii+1} = v(Q)$. It is called *minimal* for Q if $a_{ii+1} > 0$ for all $(i, i + 1) \in \mu$. Throughout this paper and with a slight abuse of notation, we identify a (possibly non-matched) pair $(i, i + 1)$ of neighbors in P with the two-person coalition $\{i, i + 1\}$. Let $Q \subseteq P$ and $\mu \in \mathcal{N}(Q)$. Let $i \in P$. If $(i - 1, i) \in \mu$ or $(i, i + 1) \in \mu$ then player i is called *matched* (with respect to μ), otherwise he is called *isolated* (with respect to μ).

Example 3.1. Let $P = \{1, 2, 3, 4\}$ be the player set. Take $a_{12} = 10, a_{23} = 20$, and $a_{34} = 30$. Then the corresponding neighbor game (P, v) is depicted in Table 1. The matching $\mu = \{(1, 2), (3, 4)\}$ is optimal and minimal for P .

From the definition of neighbor games it immediately follows that the class of neighbor games is the intersection of the class of assignment games and component additive games. It is also evident that neighbor games are monotonic game (i.e., $v(S) \leq v(T)$ for all $S \subseteq T \subseteq P$) and superadditive (i.e., $v(S \cup T) \geq v(S) + v(T)$ for all $S, T \subseteq P$ with $S \cap T = \emptyset$).

Since neighbor games are special assignment games, the results of Shapley and Shubik (1972) on the core of assignment games apply to the core of neighbor games. In particular, the core of neighbor games is not empty. Furthermore, it is determined by the inequalities induced by the one player coalitions and the pairs of neighbors. In other words, for any neighbor game (P, v) the collection

$$\mathcal{G} := \{\{i\} : i \in P\} \cup \{(i, i + 1) : i \in P \setminus \{p\}\} \quad (2)$$

contains all essential coalitions of any neighbor game on P . Henceforth, whenever we speak of a coalition it is a singleton or a pair of neighbors.

For an optimal matching μ of P we denote, with a slight abuse of notation, by P^+ the set of players that are matched by μ . Define $P^- := P \setminus P^+$, the set of isolated players. The following lemma is a straightforward consequence of a result of Shapley and Shubik (1972).

Lemma 3.2. *Let (P, v) be a neighbor game. Let μ be an optimal matching of P . Let $x \in \mathbb{R}^P$. Then, $x \in \text{Core}(P, v)$ if and only if the following four conditions are satisfied:*

- (i) $x_i + x_{i+1} = v(i, i + 1)$ for all $(i, i + 1) \in \mu$;
- (ii) $x_i + x_{i+1} \geq v(i, i + 1)$ for all $(i, i + 1) \notin \mu$;
- (iii) $x_i = 0$ for all players $i \in P^-$;
- (iv) $x_i \geq 0$ for all players $i \in P^+$.

Next, we present an algorithm for finding the nucleolus for a special class of neighbor games. Let us consider the subclass of neighbor games (P, v) with an even number of players such that the pairs

² Nevertheless, in some cases (Corollary 3.12 and further) we use a different set of players P to define a neighbor game (P, v) .

Table 1
A neighbor game (P, v)

S	$\{1,2\}$	$\{2,3\}$	$\{3,4\}$	$\{1,2,3\}$	$\{1,2,4\}$	$\{1,3,4\}$	$\{2,3,4\}$	$\{1,2,3,4\}$
$v(S)$	10	20	30	20	10	30	30	40

$(1, 2), \dots, (p - 1, p)$ form the unique optimal matching for P . Consequently, we must have $v(1, 2), \dots, v(p - 1, p) > 0$.

Initial step of algorithm for the nucleolus for neighbor games

Input

A neighbor game (P, v) for which p is even and $\mu = \{(2k - 1, 2k) : k = 1, \dots, \frac{p}{2}\}$ is the unique optimal matching.

The initial allocation

Compute recursively for $k = 1, \dots, \frac{p}{2}$

$$x_{2k-1} := \max\{v(2k - 2, 2k - 1) - x_{2k-2}, 0\}$$

$$x_{2k} := v(2k - 1, 2k) - x_{2k-1},$$

where $v(0, 1) := 0 =: x_0$. The allocation $x \in \mathbb{R}^P$ is the initial allocation.

Satisfaction

Calculate the initial satisfactions of the singletons $f_k := f(\{k\}, x) = x_k$ and the initial satisfactions of the even–odd pairs $f_{2k, 2k+1} := f(\{2k, 2k + 1\}, x) = x_{2k} + x_{2k+1} - v(2k, 2k + 1)$.

Set $\alpha := 0$.

Qualification

Call a coalition

- *settled* if it is an odd–even pair of neighbors;³
- *unsettled* otherwise.

Call an unsettled coalition

- *active* if its satisfaction equals α ;
 - *inactive* otherwise.
-

Before we go on to the inductive step we prove that the initial payoff allocation x is the even friendly core allocation and that the minimum satisfaction level of all unsettled coalitions is $\alpha (= 0)$.

Lemma 3.3. *For the initial payoff allocation x we have*

- (i) $x_i = v(1, \dots, i) - v(1, \dots, i - 1)$ for all $i \in P$ where $v(1, 0) := 0$;
- (ii) $x_i \geq 0$ for every odd player i and $x_i > 0$ for every even player i .

Proof. First note that in case i is even, we have

$$v(1, \dots, i) = v(1, \dots, i - 2) + v(i - 1, i), \tag{3}$$

and in case i is odd, we have

$$v(1, \dots, i) = \max\{v(1, \dots, i - 1), v(1, \dots, i - 2) + v(i - 1, i)\}. \tag{4}$$

³ Henceforth, we say odd–even pair for short. Likewise for even–odd pairs of neighbors.

(i) The proof is by induction on the players. Obviously, the lemma holds for $i = 1, 2$. Suppose that the lemma holds for players $1, \dots, k$ with $k \geq 2$. We will prove that the lemma holds for $i = k + 1$. We distinguish between two cases.

Suppose i is even. Then,

$$x_i = v(i-1, i) - x_{i-1} = v(i-1, i) - [v(1, \dots, i-1) - v(1, \dots, i-2)] = v(1, \dots, i) - v(1, \dots, i-1),$$

where the first equality follows from the definition of x_i , the second equality from the induction hypothesis, and the third equality from (3).

Suppose i is odd. Then,

$$\begin{aligned} x_i &= \max\{v(i-1, i) - x_{i-1}, 0\} \\ &= \max\{v(i-1, i) - [v(1, \dots, i-1) - v(1, \dots, i-2)], 0\} \\ &= \max\{v(1, \dots, i-2) + v(i-1, i) - v(1, \dots, i-1), 0\} \\ &= \max\{v(1, \dots, i-2) + v(i-1, i), v(1, \dots, i-1)\} - v(1, \dots, i-1) \\ &= v(1, \dots, i) - v(1, \dots, i-1), \end{aligned}$$

where the first equality follows from the definition of x_i , the second equality from the induction hypothesis. The third and fourth equalities are obtained by elementary rewriting. Finally, the last equality follows from (4).

(ii) It immediately follows from the definition of x_i that $x_i \geq 0$ for every odd player i . For an even player i it holds that

$$\begin{aligned} x_i &= v(1, \dots, i) - v(1, \dots, i-1) \\ &= v(1, \dots, i-2) + v(i-1, i) - v(1, \dots, i-1) \\ &= v(1, \dots, i-2) + v(i-1, i) - \max\{v(1, \dots, i-2), v(1, \dots, i-3) + v(i-2, i-1)\} \\ &> 0, \end{aligned}$$

where the first equality follows from (i). The second and third equalities from (3) and (4), respectively. Finally, the inequality follows from the fact that μ is the unique optimal matching (here we use that $v(1, 2), v(3, 4), \dots, v(p-1, p) > 0$). \square

Lemma 3.4. *The initial payoff allocation x is a core allocation. Moreover, it is the even friendly core allocation, i.e., for all $y \in \text{Core}(P, v)$ and for all even players i it holds that $x_i \geq y_i$.*

Proof. One easily verifies that x satisfies the conditions of Lemma 3.2. This shows that $x \in \text{Core}(P, v)$.

Suppose there is a core allocation $y \in \text{Core}(P, v)$ and an even player i such that $x_i < y_i$. Since $(i-1, i) \in \mu$,

$$y_{i-1} = v(i-1, i) - y_i = x_{i-1} - (y_i - x_i) < x_{i-1}, \quad (5)$$

where the equalities follow from Lemma 3.2(i) for $x, y \in \text{Core}(P, v)$ and the inequality from $x_i < y_i$.

If $x_{i-1} < (y_i - x_i)$, then it follows from (5) that $y_{i-1} < 0 = v(i-1)$, contradicting $y \in \text{Core}(P, v)$. So, $x_{i-1} \geq (y_i - x_i) > 0$. Then, by definition of x_{i-1} , we have $x_{i-1} = v(i-2, i-1) - x_{i-2}$. So,

$$x_{i-2} = v(i-2, i-1) - x_{i-1} < v(i-2, i-1) - y_{i-1} \leq y_{i-2},$$

where the first inequality follows from (5) and the second inequality from $y \in \text{Core}(P, v)$.

So, $x_{i-2} < y_{i-2}$. We can repeat the same argument until we conclude that $y_1 < x_1 = v(1)$, contradicting $y \in \text{Core}(P, v)$. \square

Corollary 3.5. *The minimum satisfaction level of all unsettled coalitions with respect to the initial allocation x equals $\alpha = 0$.*

Proof. Follows immediately from Lemma 3.4 and $f_1 = x_1 = 0$. \square

During the inductive step of the algorithm, which will be spelt out next, we settle singletons and pairs that have not been settled yet. The satisfaction of a coalition that is settled will no longer change during the remainder of the algorithm. The algorithm terminates when all singletons are settled.

In every step of the algorithm we deal with a collection of settled and unsettled coalitions. We refine the collection of singletons and pairs that have not been settled yet in two subcollections: a collection of active coalitions and a collection of inactive coalitions. An unsettled singleton or pair is called active if it has the minimum satisfaction among all coalitions that are unsettled. Otherwise it is called inactive.

We define a *component* to be a maximal set of consecutive players in which each pair of neighbors is settled or active. Note that since odd–even pairs are settled, a component always starts with an odd player and ends with an even player.

The idea of the inductive step is the following. The initial allocation is the even friendly core allocation (Lemma 3.4). Hence, in order to obtain the nucleolus we should decrease the satisfaction of the even players and increase the satisfaction of the odd players. For every player i we determine a coefficient d_i that indicates in what direction and with which factor the satisfaction of player i is going to change. After that, a positive number β is determined. The number β depends on the unsettled even players and the inactive (even–odd) pairs. For every player i , the satisfaction is now updated by adding $d_i\beta$ to his current satisfaction. The minimum satisfaction of all unsettled players and pairs is increased with β . Finally, some singletons and pairs may become settled. If this is the case, we verify whether there are still unsettled singletons. If there are no unsettled singletons left, then we are done and the allocation corresponding to the final satisfactions is the nucleolus. Otherwise, we repeat the inductive step.

Inductive step of the algorithm for the nucleolus for neighbor games

Input

A neighbor game (P, v) for which p is even and $\mu = \{(2k - 1, 2k) : k = 1, \dots, \frac{p}{2}\}$ is the unique optimal matching.

The satisfactions $f_S, S \in \mathcal{G}$ that correspond with the initial allocation x .

The initial minimal satisfaction level of all unsettled coalitions $\alpha (= 0)$.

The initial qualification of the coalitions in \mathcal{G} as settled, active, or inactive.

As long as there is an unsettled singleton repeat the following procedure.

If all singletons are settled, then **STOP**, $n(P, v) = (f_i)_{i \in P}$.

Beginning of the procedure

1. Coefficients

Compute for $k = 0, \dots, \frac{p}{2} - 1$,

$$d_{2k+1} := \begin{cases} 0 & \text{if } 2k + 1 \text{ is settled;} \\ 1 & \text{if } 2k + 1 \text{ is active and } 2k + 1 = 1; \\ 1 & \text{if } 2k + 1 \text{ is active and } (2k, 2k + 1) \text{ is inactive;} \\ -d_{2k} + 1 & \text{if } 2k + 1 > 1 \text{ and } (2k, 2k + 1) \text{ is active.} \end{cases} \quad (6)$$

$$d_{2k+2} := -d_{2k+1}. \quad (7)$$

Compute for $k = 1, \dots, \frac{p}{2} - 1$,

$$d_{2k, 2k+1} := d_{2k} + d_{2k+1}. \quad (8)$$

2. Increase of minimum satisfaction of unsettled coalitions

Compute

$$\beta := \min \left\{ \frac{f_S - \alpha}{1 - d_S} : S \text{ is an inactive coalition and } d_S \leq 0 \right\}. \quad (9)$$

Now, update the satisfactions of all unsettled coalitions:

$$f_S := f_S + d_S \beta \quad \text{for all unsettled } S. \quad (10)$$

Update $\alpha := \alpha + \beta$.

3. Qualification

If an *unsettled even player* i becomes active, i.e., $f_i = \alpha$, then settle all coalitions from the component's left most (odd) player upto and including player i .

If an *inactive pair* $(i, i + 1)$ becomes active and player $i + 1$ was already settled, then settle all coalitions from the component's left most (odd) player upto and including coalition $(i, i + 1)$.

End of the procedure

In the next example we illustrate the algorithm.

Example 3.6. Let $P = \{1, \dots, 8\}$ be a player set. In Fig. 1, nodes depict the players and the number above an edge denotes the value of the corresponding pair of players. The thick edges correspond with the matched pairs in the optimal matching. The essential information of the neighbor game (P, v) induced by Fig. 1 is represented in the first two rows of Table 2. The players and pairs are put in the first row. The values of the pairs are given by the numbers in the second row. We calculate the nucleolus of (P, v) in Table 2. We first calculate the initial allocation x in row 3 using the initial step. In row 4 we turn to the inductive step. For an explanation of the concise notation in the table, let us consider rows 4–6.

In row 4 we depict the satisfactions of all coalitions. If a particular satisfaction is in a box, then the corresponding coalition has already been settled. If a satisfaction has an asterisk, then the satisfaction equals α , the minimum satisfaction of the unsettled coalitions. In row 5 we put the coefficients, which are determined by (6)–(8). If a satisfaction and the coefficient below are in boldface, then the corresponding coalition determines the number β , using (9). Finally, in row 6 we update the satisfactions using (10). Finally, we settle players using the qualification.

We repeat the inductive step until all singletons are settled. The final allocation $(3, 3, 2, 5, 6, 3, 1, 1)$ is the nucleolus of the game (P, v) . Note that coalition $(2, 3)$ has not been settled by the algorithm.

The next lemma shows, among others, that the inductive step is well-defined and that the algorithm terminates in a finite number of steps ((e), (f), and (g)). The lemma will also be used to prove Theorem 3.9, which states that the resulting allocation of the algorithm is the nucleolus.




Fig. 1. A neighbor game (P, v) .

Table 2
Calculating the nucleolus of (P, v)

	1	12	2	23	3	34	4	45	5	56	6	67	7	78	8
v		6		1		7		9		9		2		2	
x	0		6		0		7		2		7		0		2
$\alpha = 0, f$	0*	0	6	5	0*	0	7	0*	2	0	7	5	0*	0	2
$\beta = 1, d$	+1	0	-1	0	+1	0	-1	+1	+2	0	-2	-1	1	0	-1
$\alpha = 1, f$ settling	1*	0	5	5	1*	0	6	1*	4	0	5	4	1*	0	1*
$\alpha = 1, f$ $\beta = 1, d$	1*	0	5	5	1*	0	6	1*	4	0	5	4	1	0	1
$\alpha = 2, f$ settling	2*	0	4	5	2*	0	5	2*	6	0	3	2*	1	0	1
$\alpha = 2, f$ $\beta = 1, d$	2*	0	4	5	2*	0	5	2*	6	0	3	2*	1	0	1
$\alpha = 3, f$ settling	3*	0	3*	4	2	0	5	2	6	0	3	2	1	0	1
$\alpha = 3, f$	3	0	3	4	2	0	5	2	6	0	3	2	1	0	1

Lemma 3.7. *In the inductive step of the algorithm:*

- (a) *The coefficients of the odd players in (6) are well-defined, that is:*
 - (i) *player 1 is not inactive;*
 - (ii) *for $k > 1$, player $2k + 1$ and pair $(2k, 2k + 1)$ are not both inactive.*
- (b) *The coefficients defined in (6), (7), and (8) satisfy:*
 - (i) $d_{2k+1} \geq 1$ *for an unsettled odd player $2k + 1$;*
 - (ii) $d_{2k+2} \leq -1$ *for an unsettled even player $2k + 2$;*
 - (iii) $d_{2k, 2k+1} = 1$ *for an active pair $(2k, 2k + 1)$;*
 - (iv) $d_{2k, 2k+1} \leq 0$ *for an inactive pair $(2k, 2k + 1)$ in the first inductive step; $d_{2k, 2k+1} \leq 1$ for an inactive pair $(2k, 2k + 1)$ in all other inductive steps.*
- (c) $0 < \beta < \infty$.
- (d) *The updated minimum satisfaction level of all unsettled coalitions is $\alpha + \beta$.*
- (e) *At least one unsettled even player gets settled or one inactive pair becomes active.*
- (f) *Active pairs do not become inactive.*
- (g) *Settled coalitions stay settled and the sum of the payoffs of its members does not change.*
- (h1) *If an unsettled even–odd pair gets settled, then it has satisfaction $\alpha + \beta$.*
- (h2) *Suppose that the left most (odd) player $2k + 1$ of a maximal connected set of players that get settled⁴ is unsettled. Then, player $2k + 1$ has satisfaction $\alpha + \beta$.*

Proof. We prove the lemma by induction on the number of steps. First we prove that (a)–(h2) hold for step 1. After that, we assume that (a)–(h2) hold for step $1, \dots, t - 1$ of the algorithm and that there are still unsettled singletons. Then, we will prove that (a)–(h2) also hold for step t of the algorithm.

⁴ That is, there is no strict superset that is connected and that gets settled as well.

(a) (i) Initially player 1 is not settled and $f_1 = x_1 = 0 = \alpha$. Hence, player 1 is active. (ii) Suppose that $(2k, 2k + 1)$ is inactive, i.e., $x_{2k} + x_{2k+1} - v(2k, 2k + 1) > 0$. So, $x_{2k+1} > v(2k, 2k + 1) - x_{2k}$. Then, by definition of the initial allocation, $x_{2k+1} = 0$. Hence, $x_{2k+1} = 0 = \alpha$. So, player $2k + 1$ is active.

(b) (i) Follows from (6) and (7). (ii) Since player $2k + 1$ is not settled, we have by (i) that $d_{2k+1} \geq 1$. Then, (7) implies $d_{2k+2} \leq -1$. (iii) Since $(2k, 2k + 1)$ is active, we have according to (6) that $d_{2k+1} = -d_{2k} + 1$. By (8) the result follows. (iv) Since $(2k, 2k + 1)$ is inactive, we have by (a)(ii) that $2k + 1$ is active. Hence, $d_{2k+1} = 1$ by (6). By (b)(ii) we have $d_{2k} \leq -1$. So, the result follows by (8).

(c) The coalitions that are relevant to calculate β are the even–odd pairs $(2k, 2k + 1)$ that are inactive and the inactive even players, since the odd–even pairs are settled and for inactive players $2k + 1$ it holds that $d_{2k+1} \geq 1 > 0$ by (b)(i).

Let $(2k, 2k + 1)$ be an inactive pair. Then, $d_{2k,2k+1} \leq 0$ by (b)(iv). Since $(2k, 2k + 1)$ is inactive, $f_{2k,2k+1} - \alpha = f_{2k,2k+1} > 0$. So,

$$\frac{f_{2k,2k+1} - \alpha}{1 - d_{2k,2k+1}} \in (0, \infty).$$

For an inactive even player $2k$ we have $d_{2k} \leq -1$ (by (b)(ii)) and $f_{2k} - \alpha = f_{2k} = x_{2k} > 0$ (by Lemma 3.3 (ii)). So, $((f_{2k} - \alpha)/(1 - d_{2k})) \in (0, \infty)$.

Now note that there is at least one unsettled even player and (c) follows.

(d),(e) We write f'_S for the satisfaction of coalition S after the update.

From (c) it follows that there is an active coalition S that is either an inactive pair $(2k, 2k + 1)$ or an even player $2k$ with

$$\beta = \frac{f_S - \alpha}{1 - d_S} = \frac{f_S}{1 - d_S},$$

since $\alpha = 0$. Then,

$$f'_S = f_S + d_S \beta = f_S + d_S \frac{f_S}{1 - d_S} = \beta = \alpha + \beta.$$

So, it is sufficient to prove that for all unsettled coalitions S we have $f'_S \geq \alpha + \beta$.

Let $2k + 1$ be an (unsettled) odd player. By (b) (i), $d_{2k+1} \geq 1$. So, $f'_{2k+1} = f_{2k+1} + d_{2k+1} \beta \geq f_{2k+1} + \beta \geq \alpha + \beta$, where the last inequality follows since player $2k + 1$ is unsettled.

Let $2k$ be an (unsettled) even player. By (b)(ii), $d_{2k} \leq -1$. By definition of β ,

$$\beta \leq \frac{f_{2k} - \alpha}{1 - d_{2k}}.$$

So, $f'_{2k} = f_{2k} + d_{2k} \beta \geq \alpha + \beta$.

Suppose $(2k, 2k + 1)$ is an active pair. By (b)(iii), $d_{2k,2k+1} = 1$. So,

$$f'_{2k,2k+1} = f_{2k,2k+1} + d_{2k,2k+1} \beta = f_{2k,2k+1} + \beta = \alpha + \beta,$$

where the last equality follows since $(2k, 2k + 1)$ is active.

Suppose $(2k, 2k + 1)$ is an inactive pair. By (b)(iv), $d_{2k,2k+1} \leq 0$. Then from

$$\beta \leq \frac{f_{2k,2k+1} - \alpha}{1 - d_{2k,2k+1}}$$

it follows that

$$f'_{2k,2k+1} = f_{2k,2k+1} + d_{2k,2k+1} \beta \geq \alpha + \beta.$$

(f) Follows from (b)(iii) and (d).

(g) By the settling procedure in the initial step, the only settled coalitions are the odd–even pairs. By definition of the settling procedure they stay settled in the first recursive step. It follows from (7) and (8) that the sum of the payoffs of the players in an odd–even pair does not change.

(h1) Suppose an unsettled even–odd pair $(2k, 2k + 1)$ gets settled. Then, by definition of the settling procedure, $(2k, 2k + 1)$ is active. From (b)(iii) it then follows that the satisfaction of $(2k, 2k + 1)$ is equal to $\alpha + \beta$.

(h2) Let $2k + 1$ be the left most (odd) player of a maximal connected set of players that get settled. By the initial settling procedure, player $2k + 1$ is unsettled. If $2k + 1 = 1$, then the satisfaction of player $2k + 1$ is equal to $\alpha + \beta$ by (6). If $2k + 1 \neq 1$, then, by definition of the settling procedure, $(2k, 2k + 1)$ is not active. So, by (a)(ii), player $2k + 1$ is active. Hence, by (6), the satisfaction of player $2k + 1$ is equal to $\alpha + \beta$.

Now assume that (a)–(h) hold for steps $1, \dots, t - 1$ of the algorithm. Assume that there are still unsettled singletons. We prove that (a)–(h) also hold for step t of the algorithm.

(a) As for (i), suppose that player 1 is not settled in step t . Then it follows from the Induction Hypothesis (IH, for short) ((g) and (a)(i)) that player 1 was active in step $t - 1$. Then, by (6), in step $t - 1$ we had $d_1 = 1$. So, player 1 got the minimal increase of satisfactions of unsettled coalitions in step $t - 1$. So, by (d) for step $t - 1$, player 1 is also active in step t .

As for (ii), suppose that in step t player $2k + 1$ is unsettled and $(2k, 2k + 1)$ is inactive. Then it follows from IH ((f) and (g)) that $(2k, 2k + 1)$ was inactive in step $t - 1$. From IH ((g) and (a)(ii)) it follows that player $2k + 1$ was active in step $t - 1$. So, player $2k + 1$ got the minimal increase of satisfactions of unsettled coalitions in step $t - 1$. So, by (d) for step $t - 1$, player $2k + 1$ is also active in step t .

(b)(i) Follows from (6) and (7). (ii) Follows from (6), (7), and the fact that if player $2k$ is not settled, then player $2k - 1$ is also not settled. (iii) Follows from (6) and (8). As for (iv), note that $d_{2k+1} \leq 1$ by (6). Furthermore, $d_{2k} \leq 0$ by (6) and (7). So, $d_{2k,2k+1} = d_{2k} + d_{2k+1} \leq 0 + 1$ by (8).

(c) Suppose $(i, i + 1)$ is an inactive pair. Then, $(i, i + 1) = (2k, 2k + 1)$ for some k (by IH (g) for steps $1, \dots, t - 1$). Then, $d_{2k,2k+1} \leq 1$ by (b)(iv). Since $(2k, 2k + 1)$ is inactive, $f_{2k,2k+1} - \alpha > 0$. Since we compute

$$\frac{f_{2k,2k+1} - \alpha}{1 - d_{2k,2k+1}}$$

only if $d_{2k,2k+1} < 1$, this ratio is positive and finite.

For an unsettled even player $2k$ we have $d_{2k} \leq -1$ (by (b)(ii)) and $f_{2k} - \alpha > 0$ (by IH (d) for steps $1, \dots, t - 1$ and the definition of the settling procedure). So,

$$\frac{f_{2k} - \alpha}{1 - d_{2k}} \in (0, \infty).$$

By assumption there are still unsettled singletons. Then, it follows from the definition of the settling procedure and IH (g) for steps $1, \dots, t - 1$ that there is at least one unsettled even player. This proves (c).

(d),(e) The proof is almost a copy of the proof of (d),(e) for step 1, except for the part in which we prove that for every inactive pair $(2k, 2k + 1)$ it holds that $f'_{2k,2k+1} \geq \alpha + \beta$. Take an inactive pair $(2k, 2k + 1)$. By (b)(iv), $d_{2k,2k+1} \leq 1$. If $d_{2k,2k+1} = 1$, then $f'_{2k,2k+1} = f_{2k,2k+1} + d_{2k,2k+1}\beta = f_{2k,2k+1} + \beta > \alpha + \beta$, where the inequality follows from the fact that $(2k, 2k + 1)$ is inactive and IH (d) for step $t - 1$.

If $d_{2k,2k+1} < 1$, then from

$$\beta \leq \frac{f_{2k,2k+1} - \alpha}{1 - d_{2k,2k+1}}$$

it follows that $f'_{2k,2k+1} = f_{2k,2k+1} + d_{2k,2k+1}\beta \geq \alpha + \beta$.

(f) Follows from (b)(iii) and (d).

(g) By definition of the settling procedure the settled coalitions stay settled.

Suppose an odd player $2k + 1$ is settled. By (6), $d_{2k+1} = 0$. So, the payoff of player $2k + 1$ does not change.

Suppose an even player $2k$ is settled. By definition of the settling procedure, player $2k - 1$ is also settled. Then, it follows from (6) and (7) that $d_{2k} = 0$. So, the payoff of player $2k$ does not change.

Suppose an even–odd pair $(2k, 2k + 1)$ is settled. By definition of the settling procedure, players $2k$ and $2k + 1$ are also settled. Then, it follows from the above that the payoffs of players $2k$ and $2k + 1$ do not change. So, the sum of the payoffs of players $2k$ and $2k + 1$ does not change either.

Finally, let $(2k - 1, 2k)$ be a (settled) odd–even pair. By (6) and (7), the sum of the payoffs of players $2k - 1$ and $2k$ does not change.

(h1) Suppose that an unsettled even–odd pair $(2k, 2k + 1)$ gets settled. By definition of the settling procedure, $(2k, 2k + 1)$ is active. From (b)(iii) it then follows that the satisfaction of $(2k, 2k + 1)$ is equal to $\alpha + \beta$.

(h2) Let $2k + 1$ be the left most (odd) player of a maximal connected set of players that get settled. Suppose that player $2k + 1$ is unsettled. If $2k + 1 = 1$, then, by (a)(i), player $2k + 1$ is active. From (6) it then follows that the satisfaction of player $2k + 1$ is equal to $\alpha + \beta$. If $2k + 1 \neq 1$, then, by definition of the settling procedure, $(2k, 2k + 1)$ is not active. So, by (a) (ii), player $2k + 1$ is either settled or active. Since we have assumed that player $2k + 1$ is unsettled, it follows that player $2k + 1$ is active. Hence, by (6) the satisfaction of player $2k + 1$ is equal to $\alpha + \beta$. \square

The following lemma will be used to show that the outcome of the algorithm is the nucleolus.

Lemma 3.8. *Let (P, v) be a neighbor game. Let $\mathcal{B} \subseteq \mathcal{G}$ be a non-empty collection of essential coalitions (see (2)). Then, \mathcal{B} is balanced if and only if for every $T \in \mathcal{B}$ there is a partition \mathcal{C} of P such that $T \in \mathcal{C} \subseteq \mathcal{B}$.*

Proof. First we prove the ‘if’-part. For each $T \in \mathcal{B}$, let \mathcal{C}_T be a partition of P such that $T \in \mathcal{C}_T \subseteq \mathcal{B}$. Let us count how many times a coalition $U \in \mathcal{B}$ appears in the partitions \mathcal{C}_T , $T \in \mathcal{B}$, and let u denote this number. Clearly, $1 \leq u \leq |\mathcal{B}|$. Now it is straightforward to check that the weights $\lambda_U := u/|\mathcal{B}|$ balance the collection \mathcal{B} .

To prove the ‘only if’-part, let \mathcal{B} be balanced. Take any $T \in \mathcal{B}$. Let i denote the right most player in T , i.e., $T = \{i\}$ or $\{i - 1, i\}$. If $i < p$, then there must be a coalition $U \in \mathcal{B}$ with left most player $i + 1$, since the weight of coalition $\{i, i + 1\}$ – if it is in \mathcal{B} at all – is strictly less than 1, so player $i + 1$ must also be covered by coalitions in \mathcal{B} disjoint from T . Let j denote the right most player in U , i.e., $j = i + 1$ or $i + 2$. If $j < p$, then we repeat the argument. Eventually we select disjoint coalitions from \mathcal{B} that cover all the players from $i + 1$ upto and including p . A similar argument to the left gives that there are disjoint coalitions in \mathcal{B} that cover all the players on the left of T . We conclude that the condition in the lemma is indeed satisfied. \square

Theorem 3.9. *The final allocation of the algorithm is the nucleolus.*

Proof. Let (P, v) be a neighbor game for which p is even and $\mu = \{(2k - 1, 2k) : k = 1, \dots, \frac{p}{2}\}$ is the unique optimal matching. Let z be the final allocation of the algorithm.

We prove that $z = n(P, v)$ by using Lemma 2.1, i.e., we check the balancedness of $G(z, t)$ for all $t \geq 0$. Actually, we verify the balancedness of $G(z, f(S, z))$ for all $S \in \mathcal{G}$. Take $S \in \mathcal{G}$. We have to show that $G(z, f(S, z))$ is balanced. By Lemma 3.8 it is sufficient to show that for every $T \in G(z, f(S, z))$ there is a partition \mathcal{C} of P such that $T \in \mathcal{C} \subseteq G(z, f(S, z))$. So, take $T \in G(z, f(S, z))$. We distinguish between the case in which T gets settled and the case in which it does not get settled in the algorithm.

Case 1: Coalition T gets settled in the algorithm.

Suppose T gets settled in the initial step. Then T is an odd–even pair. Since also all other odd–even pairs get settled in the initial step, the collection of these pairs is a partition with the desired property for T .

Now we suppose that T gets settled during the inductive step of the algorithm. We distinguish between the two events in the inductive step that cause T to get settled.

Subcase 1.a: an unsettled even player j becomes active.

Let $f'_j = \alpha + \beta > 0$ be his satisfaction. We settle all coalitions from the component's left most (odd) player i upto and including player j . Note that T is one of these coalitions.

By Lemma 3.7 (h1), (h2), and (g), the coalitions $\{i\}, \{i + 1, i + 2\}, \{i + 3, i + 4\}, \dots, \{j - 2, j - 1\}, \{j\}$ have satisfactions $\leq \alpha + \beta$. This collection together with some odd–even pairs forms a partition, and hence a partition with the desired property for T if $T \in \{\{i\}, \{i + 1, i + 2\}, \{i + 3, i + 4\}, \dots, \{j - 2, j - 1\}, \{j\}\}$.

If $T \notin \{\{i\}, \{i + 1, i + 2\}, \{i + 3, i + 4\}, \dots, \{j - 2, j - 1\}, \{j\}\}$, then T is some singleton $\{k\} \subseteq \{i, \dots, j\}$, since T is not an odd–even pair. Note that $f'_T \geq \alpha + \beta > 0$ because all unsettled coalitions have satisfaction $\geq \alpha + \beta$. Now $T = \{k\}$ forms together with some odd–even pairs and $\{k + 1, k + 2\}, \{k + 3, k + 4\}, \dots, \{j - 2, j - 1\}, \{j\}$ a partition with the desired property for T if k is odd. If k is even, then $T = \{k\}$ together with some odd–even pairs and $\{i\}, \{i + 1, i + 2\}, \{i + 3, i + 4\}, \dots, \{k - 2, k - 1\}$ is a partition with the desired property for T .

Subcase 1.b: an inactive pair $\{j, j + 1\}$ becomes active and coalition $j + 1$ is already settled.

Let $f'_{(j,j+1)} = \alpha + \beta$ be the satisfaction of the pair $\{j, j + 1\}$. We settle all coalitions from the component's left most (odd) player i upto and including coalition $\{j, j + 1\}$. Note that T is one of these coalitions.

By Lemma 3.7 (h1), (h2), and (g), the coalitions $\{i\}, \{i + 1, i + 2\}, \{i + 3, i + 4\}, \dots, \{j, j + 1\}$ have satisfactions $\leq \alpha + \beta$. Since $\{j + 1\}$ is settled, there exists some even player l for which the coalitions $\{j + 1\}, \{j + 2, j + 3\}, \{j + 4, j + 5\}, \dots, \{l - 2, l - 1\}, \{l\}$ are settled and all have a fixed satisfaction $< f'_{(j,j+1)}$. The collection that consists of the coalitions $\{i\}, \{i + 1, i + 2\}, \{i + 3, i + 4\}, \dots, \{j, j + 1\}, \{j + 2, j + 3\}, \{j + 4, j + 5\}, \dots, \{l - 2, l - 1\}, \{l\}$ together with some odd–even pairs forms a partition and hence a partition with the desired property for T if $T \in \{\{i\}, \{i + 1, i + 2\}, \{i + 3, i + 4\}, \dots, \{j, j + 1\}\}$.

If $T \notin \{\{i\}, \{i + 1, i + 2\}, \{i + 3, i + 4\}, \dots, \{j, j + 1\}\}$, then T is some singleton $\{k\} \subseteq \{i, \dots, j\}$, since T is not an odd–even pair. Now $T = \{k\}$ forms together with some odd–even pairs and $\{k + 1, k + 2\}, \{k + 3, k + 4\}, \dots, \{l - 2, l - 1\}, \{l\}$ a property with the desired property for T if k is odd. If k is even, then $T = \{k\}$ together with some odd–even pairs and $\{i\}, \{i + 1, i + 2\}, \{i + 3, i + 4\}, \dots, \{k - 2, k - 1\}$ is a partition with the desired property for T .

Case 2: Coalition T does not get settled in the algorithm. Since all singletons and all odd–even pairs get settled, T is an even–odd pair. So, $T = \{2k, 2k + 1\}$ for some k . Note that $\{2k, 2k + 1\}$ is inactive (otherwise we would have settled $\{2k, 2k + 1\}$ in the last step of the algorithm). Since the coalitions $\{2k\}$ and $\{2k + 1\}$ are already settled, it follows from the definition of the settling procedure and Lemma 3.7(h2) that the satisfaction of $\{2k, 2k + 1\}$ is greater than the satisfaction of both $\{2k\}$ and $\{2k + 1\}$.

Then, by combining the partitions for $\{2k\}$ and $\{2k + 1\}$ appropriately, it readily follows that there is also a partition with the desired property for $\{2k, 2k + 1\}$. \square

Proposition 3.10. *Let (P, v) be a neighbor game for which p is even and $\mu = \{(2k - 1, 2k) : k = 1, \dots, \frac{p}{2}\}$ is the unique optimal matching. Then, the algorithm determines the nucleolus of (P, v) in $\mathcal{O}(p^2)$ time.*

Proof. Initially, there are at most $p - 1$ inactive singletons and pairs ($p/2$ even players and $(p/2) - 1$ even–odd pairs and odd players (by Lemma 3.7(a)(ii))). By Lemma 3.7 (e) and (f), in every inductive step at least one unsettled even player gets settled or one inactive pair becomes active (and does not become inactive anymore). So, the algorithm terminates after at most $p - 1$ inductive steps.

Since both the initial and the inductive steps take $\mathcal{O}(p)$ time, the algorithm determines the nucleolus in $\mathcal{O}(p^2)$ time. \square

Now we will show that we can calculate the nucleolus of an arbitrary neighbor game by breaking up the game in appropriate subgames, applying the algorithm to the subgames, and constructing the nucleolus out

of the nucleoli of the subgames. Moreover, we will prove that this procedure does not change the computational complexity.

For this, let (P, v) be a neighbor game. Let us first consider the relation between the nucleolus and the kernel (Davis and Maschler, 1965) of the game (P, v) . Since the class of neighbor games is a subclass of the class of component additive games it immediately follows from Potters and Reijnierse (1995) that the nucleolus coincides with the kernel. Moreover, Corollary of Theorem 5 of Potters and Reijnierse (1995) gives

$$\{n(P, v)\} = \{x \in \text{Core}(P, v) : s_{i+1}(x) = s_{i+1i}(x) \text{ for all } i = 1, \dots, p-1\}, \quad (11)$$

where $s_{i+1}(x) = \min\{x(S) - v(S) : i \in S \subseteq P \setminus \{i+1\}, S \text{ connected}\}$ (and $s_{i+1i}(x)$ defined similarly). Note that $s_{i+1}(x), s_{i+1i}(x) \geq 0$ by the fact that $x \in \text{Core}(P, v)$. Let μ be an optimal matching for P . To simplify expression (11), we make the following two remarks for a core allocation $x \in \text{Core}(P, v)$. First, from Lemma 3.2(iii) it follows that $s_{i+1}(x) = 0$ for $i \in P^-, i \neq p$. For such i , we have that $i+1 \in P^-$ or $(i+1, i+2) \in \mu$. One can verify that in both cases $s_{i+1i}(x) = 0$. Second, if $\{i, i+1\} \subseteq P^+$, but $(i, i+1) \notin \mu$, then $s_{i+1}(x) = 0 = s_{i+1i}(x)$ by using Lemma 3.2(i) and taking $S = \{i-1, i\}$ and $S = \{i+1, i+2\}$, respectively. From these two remarks it follows that (11) can be reduced further to

$$n(P, v) = \{x \in \text{Core}(P, v) : s_{i+1}(x) = s_{i+1i}(x) \text{ for all } (i, i+1) \in \mu\}. \quad (12)$$

Note, however, that (12) does not directly help us a great deal in calculating the nucleolus. This is because the equations $s_{i+1}(x) = s_{i+1i}(x)$ contain a lot of cumbersome minimization operations, already for a small number of players.

Nevertheless, expression (12) together with the next lemma shows that in order to calculate the nucleolus of a neighbor game, it suffices to calculate the nucleolus for the subgames with possible isolated players on the extremes and no isolated players in the middle.

Lemma 3.11. *Let (P, v) be a neighbor game. Let $x \in \text{Core}(P, v)$. Let $i \in P^-$ and $k \in \{i, \dots, p\}$. Then, $s_{kk+1}(x) = \min\{x(j, \dots, k) - v(j, \dots, k) : i \leq j \leq k\}$.*

Proof. Take some connected set $S \supseteq \{i, \dots, k\} =: S'$ with $k+1 \notin S$. Let μ_S be an optimal matching for S . Define a matching $\mu_{S'}$ for S' by $(j, j+1) \in \mu_{S'}$ if and only if $(j, j+1) \in \mu_S$ and $j \geq i$. Then,

$$\begin{aligned} (x(S) - v(S)) - (x(S') - v(S')) &\geq \left(\sum_{j \in S} x_j - \sum_{(j,j+1) \in \mu_S} v(j, j+1) \right) - \left(\sum_{j \in S'} x_j - \sum_{(j,j+1) \in \mu_{S'}} v(j, j+1) \right) \\ &= \sum_{j \in S \setminus S'} x_j - \sum_{(j,j+1) \in \mu_S, j < i} v(j, j+1) \\ &= \sum_{j \in \{i\} \cup (S \setminus S')} x_j - \sum_{(j,j+1) \in \mu_S, j < i} v(j, j+1) \\ &\geq \sum_{j \in \{i\} \cup (S \setminus S')} x_j - v(\{i\} \cup (S \setminus S')) \geq 0. \end{aligned}$$

The first inequality follows from the fact that μ_S is an optimal matching for S and $\mu_{S'}$ is a matching for S' . The first equality follows from the definition of $\mu_{S'}$. The second equality follows from $i \in P^-$. The second inequality follows from the fact that $\{(j, j+1) : (j, j+1) \in \mu_S, j < i\}$ defines a (possibly non-optimal) matching for $\{i\} \cup (S \setminus S')$. The third inequality follows from the fact that $x \in \text{Core}(P, v)$. \square

Corollary 3.12. *Let (P, v) be a neighbor game. Suppose there is a player $i \in P^-$ with $1 < i < p$. Let $S^1 := \{1, \dots, i\}$ and $S^2 := \{i, \dots, p\}$. Then,⁵*

$$n(P, v) = (n_1(S^1, v_{|S^1}), \dots, n_{i-1}(S^1, v_{|S^1}), 0, n_{i+1}(S^2, v_{|S^2}), \dots, n_p(S^2, v_{|S^2})).$$

Now let us consider neighbor games with possible isolated players on the extremes and no isolated players in the middle. In the next lemmas we show that we can make a further reduction by proving that it suffices to calculate the nucleolus of a neighbor game that slightly differs from the original game and in which we leave out the isolated players of the original game. In Lemma 3.13 we consider the case in which only the first player is isolated. In Lemma 3.14 we consider the case in which only the last player is isolated. And finally, in Lemmas 3.15 and 3.16 we consider the case in which both the first and the last player are isolated. Only the proof of Lemma 3.13 is given; the proofs of Lemmas 3.14, 3.15, and 3.16 run similarly.

Lemma 3.13. *Let (P, v) be a neighbor game with $|P| \geq 3$. Suppose that the matching $\mu = \{(2, 3), \dots, (p-1, p)\}$ is optimal. Define a neighbor game (\bar{P}, \bar{v}) , by setting $\bar{P} := P \setminus \{1\}$, $\bar{v}(2, 3) := v(2, 3) - v(1, 2)$, and $\bar{v}(i, i+1) := v(i, i+1)$ for $3 \leq i \leq p-1$. Then, $n_1(P, v) = 0$, $n_2(P, v) = n_2(\bar{P}, \bar{v}) + v(1, 2)$, and $n_i(P, v) = n_i(\bar{P}, \bar{v})$ for $i \in P$, $i \geq 3$.*

Proof. Note that $\bar{v}(2, 3) \geq 0$, since $v(2, 3) \geq v(1, 2)$ by optimality of μ . So, (\bar{P}, \bar{v}) is indeed a neighbor game.

Let $x = n(\bar{P}, \bar{v})$ and define $y \in \mathbb{R}^P$ by $y_1 := 0$, $y_2 := n_2(\bar{P}, \bar{v}) + v(1, 2)$, and $y_i := n_i(\bar{P}, \bar{v})$ for $i \in P$, $i \geq 3$. By Lemma 3.2, $y \in \text{Core}(P, v)$. One easily verifies that $f(\{1\}, y) = 0$, $f(\{1, 2\}, y) = f(\{2\}, x)$, and $f(\{2\}, y) \geq f(\{2\}, x) = f(\{1, 2\}, y)$. Further, it is clear that $f(S, y) = f(S, x)$ for any other singleton and pair of neighbors. Using Lemmas 2.1 and 3.8 one verifies that $y = n(P, v)$. \square

Lemma 3.14. *Let (P, v) be a neighbor game with $|P| \geq 3$. Suppose that the matching $\{(1, 2), \dots, (p-2, p-1)\}$ is optimal. Define a neighbor game (\bar{P}, \bar{v}) , by setting $\bar{P} := P \setminus \{p\}$, $\bar{v}(p-2, p-1) := v(p-2, p-1) - v(p-1, p)$, and $\bar{v}(i-1, i) := v(i-1, i)$ for $1 \leq i \leq p-2$. Then, $n_p(P, v) = 0$, $n_{p-1}(P, v) = n_{p-1}(\bar{P}, \bar{v}) + v(p-1, p)$, and $n_i(P, v) = n_i(\bar{P}, \bar{v})$ for $i \in P$, $i \leq p-2$.*

Lemma 3.15. *Let (P, v) be a neighbor game with $|P| = 4$. Suppose that the matching $\{(2, 3)\}$ is optimal. Define a neighbor game (\bar{P}, \bar{v}) , by setting $\bar{P} := P \setminus \{1, 4\}$ and $\bar{v}(2, 3) := v(2, 3) - v(1, 2) - v(3, 4)$. Then, $n_1(P, v) = n_4(P, v) = 0$, $n_2(P, v) = n_2(\bar{P}, \bar{v}) + v(1, 2)$, and $n_3(P, v) = n_3(\bar{P}, \bar{v}) + v(3, 4)$.*

Lemma 3.16. *Let (P, v) be a neighbor game with $|P| > 4$. Suppose that the matching $\{(2, 3), \dots, (p-2, p-1)\}$ is optimal. Define a neighbor game (\bar{P}, \bar{v}) , by setting $\bar{P} := P \setminus \{1, p\}$, $\bar{v}(2, 3) := v(2, 3) - v(1, 2)$, $\bar{v}(p-2, p-1) := v(p-2, p-1) - v(p-1, p)$, and $\bar{v}(i, i+1) := v(i, i+1)$ for $3 \leq i \leq p-3$. Then, $n_1(P, v) = n_p(P, v) = 0$, $n_2(P, v) = n_2(\bar{P}, \bar{v}) + v(1, 2)$, $n_{p-1}(P, v) = n_{p-1}(\bar{P}, \bar{v}) + v(p-1, p)$, and $n_i(P, v) = n_i(\bar{P}, \bar{v})$ for $3 \leq i \leq p-2$.*

Now we will describe an extension of the algorithm for finding the nucleolus of an arbitrary neighbor game.

⁵ Here and in the remainder of this section we consider neighbor games (Q, v) that differ slightly from the definition in the sense that there is no $q \in \mathbb{N}$ such that $Q = \{1, \dots, q\}$.

Extended algorithm for the nucleolus for neighbor games

Let (P, v) be a neighbor game. Let μ be an optimal and minimal matching for P . In view of Corollary 3.12 we break the game (P, v) up in (overlapping) subgames that are still neighbor games, but have no longer isolated players in the middle. Then, in view of Lemmas 3.13–3.15, and 3.16 we remove the isolated players from the subgames that are not zero two-person games. By doing this the induced matchings of μ for the obtained games may not all be minimal any longer. (See Example 3.17 for an illustration of this.) By taking an optimal and minimal matching for the games in which this occurs, we repeat the above procedure to remove the new isolated players. Eventually we have only games that satisfy either the assumptions made for the algorithm or are zero two-person games. Once we have calculated the nucleolus of every subgame we use Corollary 3.12 and Lemmas 3.13–3.15, and 3.16 to construct the nucleolus of the game (P, v) .

Below we present an example in which the procedure above is used to calculate the nucleolus of an 11-person neighbor game.

Example 3.17. Consider the neighbor game (P, v) , where $P = \{1, \dots, 11\}$ and v is given by Fig. 2. The nodes depict the players and the number above an edge denotes the value of the corresponding pair of players. The numbers below the nodes are the payoffs of the nucleolus of the corresponding game. The thick edges correspond with the matched pairs in the optimal matchings. We calculate the nucleolus of (P, v) .

In view of Corollary 3.12, we break the game (P, v) up in the subgames $(\{1, \dots, 6\}, v_1)$ and $(\{6, \dots, 11\}, v_5)$. Then, in view of Lemma 3.16 we reduce $(\{1, \dots, 6\}, v_1)$ to $(\{2, 3, 4, 5\}, v_2)$. We take the (unique) optimal and minimal matching $\{(4, 5)\}$ for the game $(\{2, 3, 4, 5\}, v_2)$. In view of Corollary 3.12 we reduce the game $(\{2, 3, 4, 5\}, v_2)$ to the game $(\{3, 4, 5\}, v_3)$ and the zero two-person game $(\{2, 3\}, 0)$. In view of Lemma 3.13 we reduce the game $(\{3, 4, 5\}, v_3)$ to the game $(\{4, 5\}, v_4)$. In view of Lemma 3.16 we reduce the subgame $(\{6, \dots, 11\}, v_5)$ to the game $(\{7, 8, 9, 10\}, v_6)$.

The nucleolus of the game $(\{4, 5\}, v_4)$ is $(1, 1)$. So, by Lemma 3.13 the nucleolus of $(\{3, 4, 5\}, v_3)$ is $(0, 4, 1)$. Hence, the nucleolus of $(\{2, 3, 4, 5\}, v_2)$ is $(0, 0, 4, 1)$. Then, by Lemma 3.16, the nucleolus of $(\{1, \dots, 6\}, v_1)$ is $(0, 2, 0, 4, 3, 0)$. By a similar reasoning and the use of the algorithm or Proposition 3.19, the nucleolus of $(\{6, \dots, 11\}, v_5)$ is $(0, \frac{5}{3}, \frac{10}{3}, \frac{4}{3}, \frac{5}{3}, 0)$. Finally, from Corollary 3.12 it follows that the nucleolus of (P, v) is $n(P, v) = (0, 2, 0, 4, 3, 0, \frac{5}{3}, \frac{10}{3}, \frac{4}{3}, \frac{5}{3}, 0)$.




Fig. 2. Subsequent break-ups of the neighbor game (P, v) .

In Proposition 3.18 we show that the extended algorithm has the same computational complexity as the original algorithm.

Proposition 3.18. *The extended algorithm determines for a neighbor game (P, v) with p players the nucleolus in $\mathcal{O}(p^2)$ time.*

Proof. It takes $\mathcal{O}(p^2)$ time to find the subgames to which we apply the algorithm. Let p_1, \dots, p_k be the cardinalities of the player sets of these subgames. Note, $p_1 + \dots + p_k \leq p$. By Proposition 3.10 it takes $\mathcal{O}(p_i^2)$ time to calculate the nucleolus of the l th subgame. Since $p_1^2 + \dots + p_k^2 \leq p^2$, it takes $\mathcal{O}(p^2)$ time to calculate the nucleoli of all subgames. Finally, note that the construction of the nucleolus of (P, v) out of the nucleoli of the subgames takes $\mathcal{O}(p^2)$ time. We conclude that we determine the nucleolus of (P, v) in $\mathcal{O}(p^2)$ time. \square

We conclude the paper with closed formulas for the nucleolus of neighbor games in case there are four or less players involved.

Proposition 3.19. *Let (P, v) be a two-person neighbor game, where $P = \{1, 2\}$ and the characteristic function v is induced by $a_{12} = a \geq 0$. Then $n(P, v) = (\frac{a}{2}, \frac{a}{2})$.*

Let (P, v) be a three-person neighbor game, where $P = \{1, 2, 3\}$ and the characteristic function v is induced by $a_{12} = a \geq 0$ and $a_{23} = b \geq 0$. Assume, without loss of generality, that $a \geq b$. Then,

- (i) *if $b \in [0, \frac{a}{2}]$, then $n(P, v) = (\frac{a}{2}, \frac{a}{2}, 0)$,*
- (ii) *if $b \in (\frac{a}{2}, a]$, then*

$$(P, v) = \left(\frac{a-b}{2}, \frac{a-b}{2} + b, 0 \right).$$

Let (P, v) be a 4-person neighbor game, where $P = \{1, 2, 3, 4\}$ and the characteristic function v is induced by $a_{12} = a \geq 0$, $a_{23} = b \geq 0$, and $a_{34} = c \geq 0$. Assume, without loss of generality, that $a \geq c$. Then,

- (i) *if $b \in [0, \frac{c}{2}]$, then $n(P, v) = (\frac{a}{2}, \frac{a}{2}, \frac{c}{2}, \frac{c}{2})$;*
- (ii) *if $b \in [\frac{c}{2}, \frac{2a-c}{2}]$, then*

$$n(P, v) = \left(\frac{2a-2b+c}{4}, \frac{2a+2b-c}{4}, \frac{c}{2}, \frac{c}{2} \right);$$

- (iii) *if $b \in [\frac{2a-c}{2}, a+c]$, then*

$$n(P, v) = \left(\frac{a-b+c}{3}, \frac{2a+b-c}{3}, \frac{-a+b+2c}{3}, \frac{a-b+c}{3} \right);$$

- (iv) *if $b \in [a+c, \infty)$, then*

$$n(P, v) = \left(0, \frac{b+a-c}{2}, \frac{b-a+c}{2}, 0 \right).$$

Proof. Follows straightforwardly from Lemma 2.1. \square

References

Curiel, I., Potters, J., Rajendra Prasad, V., Tijs, S., Veltman, B., 1994. Sequencing and cooperation. *Operations Research* 42, 566–568.
 Davis, M., Maschler, M., 1965. The kernel of a cooperative game. *Naval Research Logistics Quarterly* 12, 223–259.
 Hamers, H., Klijn, F., Solymosi, T., Tijs, S., Villar, J.P., 2002. Assignment games satisfy the CoMa-property. *Games and Economic Behavior* 38, 231–239.

- Huberman, G., 1980. The nucleolus and the essential coalitions. In: Bensoussan, A., Lions, J.L. (Eds.), *Analysis and Optimization of Systems*. In: *Proceedings of the Fourth International Conference on Analysis and Optimization of Systems*, Versailles, December 16–19. Springer, Berlin, pp. 416–422.
- Klijn, F., Vermeulen, D., Hamers, H., Solymosi, T., Tijs, S., Villar, J.P., 1999. Neighbor games and the leximax solution. CentER DP 99110, Tilburg University, Netherlands.
- Kohlberg, E., 1971. On the nucleolus of a characteristic function game. *SIAM Journal on Applied Mathematics* 20, 62–66.
- Potters, J., Reijnen, H., 1995. Γ -component additive games. *International Journal of Game Theory* 24, 49–56.
- Schmeidler, D., 1969. The nucleolus of a characteristic function game. *SIAM Journal on Applied Mathematics* 17, 1163–1170.
- Shapley, L., Shubik, M., 1972. The assignment game I: The core. *International Journal of Game Theory* 1, 111–130.
- Solymosi, T., Aarts, H., Driessen, T., 1998. On computing the nucleolus of a balanced connected game. *Mathematics of Operations Research* 23, 983–1009.
- Solymosi, T., Raghavan, T., 1994. An algorithm for finding the nucleolus of assignment games. *International Journal of Game Theory* 23, 119–143.