

Socio-demographic disparities in health-related quality of life after hip fracture in China: evidence from the China Health and Retirement Longitudinal Study

Citation for published version (APA):

Xu, L. Z., Chen, M. S., Peng, K., Hiligsmann, M., Jan, S. P., & Si, L. (2023). Socio-demographic disparities in health-related quality of life after hip fracture in China: evidence from the China Health and Retirement Longitudinal Study. *Archives of Osteoporosis*, 18(1), Article 29. <https://doi.org/10.1007/s11657-023-01220-2>

Document status and date:

Published: 10/02/2023

DOI:

[10.1007/s11657-023-01220-2](https://doi.org/10.1007/s11657-023-01220-2)

Document Version:

Publisher's PDF, also known as Version of record

Document license:

Taverne

Please check the document version of this publication:

- A submitted manuscript is the version of the article upon submission and before peer-review. There can be important differences between the submitted version and the official published version of record. People interested in the research are advised to contact the author for the final version of the publication, or visit the DOI to the publisher's website.
- The final author version and the galley proof are versions of the publication after peer review.
- The final published version features the final layout of the paper including the volume, issue and page numbers.

[Link to publication](#)

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal.

If the publication is distributed under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license above, please follow below link for the End User Agreement:

www.umlib.nl/taverne-license

Take down policy

If you believe that this document breaches copyright please contact us at:

repository@maastrichtuniversity.nl

providing details and we will investigate your claim.

Download date: 15 Jan. 2025



Socio-demographic disparities in health-related quality of life after hip fracture in China: evidence from the China Health and Retirement Longitudinal Study

Lizheng Xu^{1,2} · Mingsheng Chen^{3,4,5} · Ke Peng⁶ · Mickael Hiligsmann⁷ · Stephen Jan¹ · Lei Si^{8,9}

Received: 16 August 2022 / Accepted: 30 January 2023 / Published online: 10 February 2023
© International Osteoporosis Foundation and Bone Health and Osteoporosis Foundation 2023

Abstract

Summary This study analyzed the impact of hip fractures on people's health-related quality of life and its socio-demographic disparities in China.

Purpose Hip fractures cause high mortality and worsened health-related quality of life (HRQoL). This study aimed to investigate whether socio-demographic-related inequities in post-hip fracture participants' HRQoL exist in China.

Methods Data from the China Health and Retirement Longitudinal Study (waves 2013, 2015, and 2018) were used. The measurement of HRQoL in this study focused on 5 health dimensions: depression, body pain, mobility, basic activities of daily living, and instrumental activities of daily living. A difference-in-differences (DID) analysis with multiple time periods was performed to gauge the impact of hip fracture on HRQoL. A multivariate regression approach was used to explore socio-demographic-related factors associated with inequities of HRQoL.

Results A total of 23,622 individuals were included, and 341 participants reported hip fracture events during the survey period. In participants with hip fracture, the presentation rate of body pain increased by 14% ($p < 0.01$) and the HRQoL of other health dimensions worsened ($p \leq 0.01$) after hip fracture. The DID analysis showed that hip fracture had a negative impact on all HRQoL dimensions ($p < 0.01$). Socioeconomic-related factors of HRQoL inequities included school education level and location of residence. Study participants with hip fracture with greater educational attainment or living in urban areas had higher ($p < 0.05$) levels of HRQoL. In addition, comorbidities also correlated with a worse HRQoL ($p < 0.05$).

Conclusion Hip fracture significantly affects people's HRQoL in China, and the impact is more profound for those with lower educational attainment or living in rural areas. Targeted interventions should be designed to narrow this inequity.

Keywords Hip fracture · Health-related quality of life · Disparities · China

✉ Mingsheng Chen
cms@njmu.edu.cn

¹ The George Institute for Global Health, Sydney, NSW, Australia

² UNSW Medicine, UNSW Sydney, Sydney, NSW, Australia

³ School of Health Policy and Management, Nanjing Medical University, No. 101, Longmian Avenue, Nanjing 211166, China

⁴ Creative Health Policy Research Group, Nanjing Medical University, Nanjing, China

⁵ Center for Global Health, Nanjing Medical University, Nanjing, China

⁶ National Clinical Research Center for Cardiovascular Diseases, Fuwai Hospital Chinese Academy of Medical Sciences, Shenzhen, China

⁷ Department of Health Services Research, CAPHRI Care and Public Health Research Institute, Maastricht University, Maastricht, the Netherlands

⁸ School of Health Sciences, Western Sydney University, Campbelltown, Australia

⁹ Translational Health Research Institute, Western Sydney University, Penrith, Australia

Introduction

There are 2 types of fracture—namely, traumatic and non-traumatic. Age is an unavoidable risk factor for non-traumatic fractures; hence, non-traumatic fractures should be paid more attention to in the era of population aging [1]. Osteoporosis and osteopenia are leading causes of fractures in older adults. Approximately one-third of women and one-fifth of men aged ≥ 50 years worldwide were expected to experience an osteoporotic fracture in their remaining lifespan [2]. The residual lifetime fracture risk in the Chinese population is even higher due to the high prevalence of osteoporosis and low bone mineral density [3]. China has the largest aging population worldwide, and it was estimated that the number of osteoporotic fractures nationwide will double by 2035 [4].

Given osteoporotic fracture is a major cause of morbidity, mortality, and disability [5–8], it is not surprising that the impacts on both physical and mental health-related quality of life (HRQoL) can persist long after the event [9–11]. Numerous studies on HRQoL loss after fracture have been conducted in Western countries, most of which highlighted the major negative impact of the event on participants' HRQoL [11, 12] often varying by age, fracture history, comorbidity burden, and fracture site [13]. Among all types of fractures, hip fractures were evidenced to have the most profound impact on HRQoL [11, 14, 15].

Research on equity in health outcomes, health resource utilization, and HRQoL has been broadly conducted in many chronic diseases, such as chronic obstructive pulmonary disease (COPD) and cancer [16, 17], revealing HRQoL disparities across socioeconomic groups, such as the location of residence [16]. While there is strong evidence of the impact of fracture on the HRQoL on individuals, little is known about disparities in HRQoL. The aim of this study is therefore to conduct an equity analysis to determine socio-demographic disparities in HRQoL in China.

Methods

Data source and participants

Data from the China Health and Retirement Longitudinal Study (CHARLS) was used. CHARLS is a nationally representative longitudinal study that included participants aged ≥ 45 years in which details of study participants' social, economic, and health status; type of health insurance; and health care service use were collected [18]. The first wave survey was conducted in 2011, including 17,705 participants

from 28 provinces in mainland China. The follow-up surveys were conducted in 2013, 2015, and 2018. In this study, we used only data collected in 2013 or later; as no hip fracture-related information was collected in the 2011 survey.

Participants who sustained a hip fracture during the study were identified by the question “have you fractured your hip since the last interview?” with the fracture time then recorded. Study participants who answered “yes” in the 2015 wave were considered to have experienced a hip fracture from 2013–2015. Similarly, those who answered “yes” in the 2018 wave were considered to have experienced a hip fracture from 2015 to 2018. Hip fracture occurrence was self-reported by participants. All study participants were divided into a post-hip fracture group covering study participants who reported hip fracture in 2015 and 2018 waves and a control group containing study participants with no history of hip fracture throughout the research period.

Measures

While CHARLS did not use a generic HRQoL instrument, there were 31 questions deployed related to HRQoL across different health domains. We selected the HRQoL questions applied in CHARLS that reflected the 5 health domains in the EuroQoL 5-dimension (EQ-5D) questionnaire, which is the most commonly used generic HRQoL instrument in the field of hip fracture [19].

HRQoL questions that assessed mobility, self-care, usual activities, pain/discomfort, and anxiety/depression [20] were thus included in our study. Specifically, 9 items related to simple physical moving activities, such as walking, climbing stairs, stooping, carrying weights, and extending arms, were included to reflect the HRQoL in the area of mobility. Each item was scored from 1 (“no, do not have any difficulty”) to 4 (“cannot do it”) points. The sum of the scores of all 9 items ranged from 9–36 points, with a lower score indicating a better HRQoL in the area of mobility.

Usual activities were measured using the Activities of Daily Living (ADL) questionnaire, which covered Basic ADL (BADL, the ability to perform fundamental activities to care for oneself) and Instrumental ADL (IADL, the ability to perform complex activities in daily life) [21]. Possible scores ranged from 6 to 24 points for BADL and IADL. Similar to the total score for mobility, a lower score indicated a better HRQoL in the area of usual activities.

The existence of pain or discomfort was assessed by asking “are you often troubled with any body pains?” The answer was coded as a dichotomous variable to reflect whether the participant reported pain or not.

Depression was measured using the Center for Epidemiologic Studies Depression (CES-D-10) scale, which is a reliable and validated instrument to screen for depression

[22]. There are 10 items in the CES-D-10, and each item investigates the frequency of depression-related symptoms. Each response ranges from 1 (rarely or none of the time) to 4 (most or all the time) points, and the total score of the CES-D-10 scale ranges from 10 to 40 points. A lower CES-D-10 score indicated a better HRQoL in the area of depression. Details on the item selection are provided in Supplementary Table 1.

Control variables

Previous studies found that several demographic, socioeconomic, and health factors were associated with HRQoL among adults with fractures [14, 23–26]. Those variables informed the selection of control variables in our study. Demographic variables included sex and age. Socioeconomic variables included economic status and school educational level. Economic status was assessed using annual household consumption expenditure per capita [27]. School education was categorized into 4 groups: (1) no school education, (2) primary school, (3) middle school, and (4) high school and above. Region of residence was geographically divided into rural and urban. Health insurance was classified as (1) Urban Employee Basic Medical Insurance (UEBMI), (2) Urban–Rural Residents Basic Medical Insurance (URRBMI), (3) other health insurance (including commercial health insurance), and (4) no health insurance. The presence of comorbidities was also included as a control variable [28]. CHARLS respondents were asked whether they had been diagnosed with one or more diseases from 14 pre-defined chronic conditions (hypertension; dyslipidemia; diabetes or high blood sugar; cancer or malignant tumor; chronic lung diseases; liver disease; heart disease; stroke; kidney disease; stomach or other digestive diseases; emotional, nervous, or psychiatric problems; memory-related disease; arthritis or rheumatism; and asthma). In our study, the presence of comorbidities was categorized into 5 groups by the total number of chronic conditions (0, 1, 2, 3, and ≥ 4 chronic diseases).

Statistical analysis

A descriptive analysis was used to summarize the characteristics of all study participants. In study participants with a hip fracture, difference tests on matched data and McNemar's chi-squared test were performed to evaluate the change in HRQoL between before and after hip fracture. Furthermore, a generalized difference-in-differences (DID) analysis with multiple time periods was conducted to quantify the effects of hip fracture on HRQoL between post-hip fracture group and the control group. The generalized DID analysis was performed using the following formula:

$$y_{it} = \alpha + \delta \text{fracture}_{it} + \beta X_{it} + \varepsilon_{it}$$

where y_{it} is the HRQoL of individual i at time t , fracture_{it} is a dummy variable indicating whether individual i experienced fracture at time t , X_{it} denotes a set of control variables, α is a constant term, and ε is a residual term. δ captures the impact of fracture on HRQoL.

To explore the socioeconomic disparity in HRQoL, a multivariate regression analysis was conducted for the post-hip fracture participants. The dependent variables were changes of each HRQoL dimension and the independent variables were the control variables. For depression, mobility, BADL, and IADL dimensions, a multivariate linear regression analysis was performed while for the pain dimension, a multivariate logistic regression analysis was used. Statistical significance was determined if the 2-tailed p value was < 0.05 . STATA (version 15.0; Stata Corporation, College Station, TX, USA) was used for statistical analyses.

Results

A total of 23,622 study participants from 3 waves of CHARLS were included in this study and 341 respondents were identified as study participants with hip fracture, among which 210 participants had hip fracture between 2013 and 2015, and 131 had hip fracture between 2015 and 2018. The basic characteristics of the respondents are summarized in Table 1. The mean age of the total sample was 59.8 (standard deviation, 10.9) years, and there were more women than men enrolled (52.3% vs. 47.8%). Most of the respondents had a low school education level. Approximately 41% did not complete a primary school education, and 13% had at least a high school education. More than half of the respondents were from rural areas, and 73.6% of the respondents were covered by the URRBMI, while 8% of the study participants did not have any health insurance. About 70% of the study participants reported chronic diseases, with 43.1% suffering from ≥ 2 chronic diseases. Study participants with hip fracture had less household expenditure ($p < 0.05$) and lower educational attainment ($p < 0.05$) and suffered from more chronic diseases ($p < 0.05$), compared with the non-hip fracture group. Besides, more study participants with hip fracture were living in rural areas ($p < 0.05$).

Figure 1 displays the HRQoL values and the proportions of participants who reported any issue by health domain and before/after hip fracture. All of the 5 HRQoL dimensions were worsened after hip fracture ($p < 0.05$). The values for CES-D-10, mobility, BADL, and IADL were increased by 0.61 ($p = 0.01$), 1.50 ($p < 0.01$), 0.60 ($p < 0.01$), and 0.87 ($p < 0.01$) points for participants with a post-fracture status. The prevalence of body pain increased by 14% ($p < 0.01$) after hip fracture.

Table 1 Characteristics of participants #, n (%)

Variables		Total sample		Non-hip fracture participants		Hip fracture participants	
		n	%	n	%	n	%
Age (mean ± SD)		59.81 ± 10.85		59.72 ± 10.80		60.21 ± 9.65	
Sex	Male	11,279	47.75	11,109	0.48	170	49.85
	Female	12,343	52.25	12,172	0.52	171	50.15
Annual household expenditure (RMB, mean ± SD)*		15,341 ± 27,618		15,400 ± 23,889		11,281 ± 10,636	
Educational attainment*	Lower than primary school	9776	41.39	9589	0.41	187	54.84
	Primary school	6018	25.48	5943	0.26	75	21.99
	Middle school	4756	20.13	4699	0.20	57	16.72
	High school and above	3072	13.00	3050	0.13	22	6.45
Number of chronic diseases*	0	7429	31.45	7350	0.32	79	23.17
	1	6006	25.43	5912	0.25	94	27.57
	2	4378	18.53	4299	0.18	79	23.17
	3	2693	11.40	2654	0.11	39	11.44
	≥ 4	3116	13.19	3066	0.13	50	14.66
Rural–urban*	Rural	13,513	57.21	13,279	0.57	234	68.62
	Urban	10,109	42.79	10,002	0.43	107	31.38
Health insurance	UEBMI	2855	12.09	2824	0.12	31	9.09
	URRBMI	17,390	73.62	17,125	0.74	265	77.71
	Other health insurance	1483	6.28	1467	0.06	16	4.69
	No insurance	1894	8.02	1865	0.08	29	8.50

For participants who had enrolled in more than one wave of CHARLS, only their first year’s information was used in the summary statistics. * P-value of the difference test between non-hip fracture group and hip fracture group is lower than 0.05. UEBMI, Urban Employee Basic Medical Insurance; URRBMI, Urban–Rural Residents Basic Medical Insurance

Fig. 1 HRQoL scores and proportions of participants who reported any issue both before and after a hip fracture (percent (score)). # $p=0.01$. * $p < 0.01$




Figure 2 displays the effect of hip fracture on the 5 HRQoL dimensions. After controlling covariates, hip fracture displayed a significantly negative effect ($p < 0.01$) on all the fracture participants’ HRQoL domains. Specifically, the CES-D-10 score increased by 2.2 points after hip fracture compared to the scores of respondents without the occurrence of hip fracture. Similarly, the hip fracture group also showed increased scores for morbidity, BADL, and IADL by about 2.3, 1.3, and 1.6 points, respectively. Furthermore,

likelihood of study participants with hip fracture suffering from body pain increased by 1.5-fold after hip fracture.

The results presented in Table 2 reveal the disparities in HRQoL in each dimension. Specifically, an older age was significantly associated with lower HRQoL values. Study participants with hip fracture living in urban areas had higher depression ($\beta = -1.39, p < 0.05$) and IADL values ($\beta = -0.60, p < 0.05$) compared to their rural counterparts. Besides, a higher school education level was positively




Fig. 2 Effect of hip fracture in each domain of HRQoL among study participants with hip fracture. Note: The results were extracted from generalized DID analysis for all 5 HRQoL domains. Depression, mobility, BADL, and IADL were continuous variables; hence, the effects were determined using β values. Meanwhile, the pain dimen-

sion was a dichotomous variable, so an odds ratio (OR) was used to capture the effect. Sex, age, economic status, education level, health insurance coverage, presence of chronic disease, and region of residence were included as control variables. All of the β values and the OR were statistically significant ($p < 0.01$)

related with higher values for depression, BADL, and IADL and a lower occurrence of pain ($p < 0.05$). Furthermore, the presence of chronic diseases was associated with a low level of HRQoL, though statistical significance was only found among those with ≥ 4 chronic diseases ($p < 0.05$) across all 5 HRQoL domains. Body pain was more prevalent among those with ≥ 2 chronic diseases.

Discussion

This study quantifies the effect of hip fracture on HRQoL and evaluates socio-demographic disparities among hip fracture people in China. Using data from the CHARLS, we found that hip fracture had a negative impact on people's physical and mental HRQoL and led to a higher likelihood of pain. Besides, the impact on HRQoL was not evenly distributed across socio-demographic groups in terms of educational attainment and location of residence. This disparity in HRQoL was evident across different HRQoL dimensions, where mental health was more closely related to education and the location of residence, while physical domains were associated with participants' age and health status.

Previous studies have generally shown that HRQoL can gradually recover after hip fracture but does not return to the pre-fracture level [12, 23]. A multinational study demonstrated that hip fracture was associated with HRQoL declines of 115.6%, 36.4%, and 22.1% immediately, 4 months, and 12 months after fracture, respectively [8]. A systematic review documented a 19–34% HRQoL loss among participants even 2 years after a hip fracture [11]. Our study results

support such previous findings; even 2–3 years after hip fracture, participants still suffered from a 2.2 points reduction in mental health and a 2.5 points reduction in physical mobility. The proportion of study participants with hip fracture who reported body pain was increased by 14% compared to that before fracture.

Hip fracture has varied impacts in different HRQoL domains. We found that the change in score for depression was smallest after a hip fracture, reflecting the lowest depression occurrence and severity among all dimensions of HRQoL. This finding echoed those of many previous studies [10, 29], indicating that hip fracture impacts physical health more than mental health. However, other studies have reported conflicting results, in which a larger impact on psychological health compared to physical health was evident [9, 30, 31].

We obtained a similar result, determining that a large proportion of our sample suffered from body pain after hip fracture [32]. Moreover, pain was not only present in the short term but could last for several months after fracture [33, 34]. Persistent pain in study participants with hip fracture can also adversely affect other HRQoL dimensions, leading to reduced physical activity [35] and depression symptoms [36]. Pain management is therefore of major importance in improving post-hip fracture participants' overall quality of life.

Socioeconomic aspects, such as education and the location of residence, can contribute to disparities in HRQoL among elderly study participants with hip fracture. We found a close relationship between less educational attainment and a lower level of HRQoL in those who had suffered a hip

Table 2 Socio-demographic inequities in HRQoL for study participants with hip fracture

Variables	Depression		Pain		Mobility		BADL		IADL	
	β	95% CI	OR	95% CI	β	95% CI	β	95% CI	β	95% CI
Age	0.04	(-0.02,0.10)	1.00	(0.98,1.02)	0.18**	(0.14,0.23)	0.05**	(0.03,0.07)	0.09**	(0.06,0.12)
Sex										
Male										
Female	0.58	(-0.62,1.78)	1.34	(0.94,1.92)	1.65*	(0.68,2.62)	0.15	(-0.29,0.59)	0.29	(-0.32,0.90)
Income	0.25	(-1.15,1.65)	0.84	(0.53,1.34)	0.09	(-0.99,1.17)	0.37	(-0.10,0.84)	0.10	(-0.54,0.75)
Education										
None										
Primary school	-1.83*	(-3.34,-0.32)	0.58*	(0.38,0.90)	-0.93	(-2.05,0.18)	-0.37	(-0.80,0.06)	-0.85*	(-1.54,-0.16)
Middle school	-0.64	(-2.26,0.98)	1.13	(0.69,1.86)	-0.30	(-1.57,0.97)	-0.07	(-0.69,0.56)	-0.68	(-1.42,0.06)
High school and above	-2.24*	(-4.24,-0.24)	0.56	(0.26,1.21)	-0.98	(-2.56,0.61)	-0.72*	(-1.27,-0.16)	-1.26**	(-2.17,-0.35)
Insurance										
UEBMI										
URRBMI	0.97	(-0.85,2.78)	1.93	(0.95,3.89)	1.37	(-0.03,2.78)	0.19	(-0.37,0.75)	0.59	(-0.26,1.43)
Other	-1.36	(-4.02,1.30)	1.23	(0.52,2.91)	2.15	(-0.09,4.38)	1.01*	(0.12,1.90)	1.48*	(0.15,2.81)
None	0.99	(-1.70,3.69)	1.17	(0.48,2.85)	2.68**	(0.75,4.60)	0.72	(-0.18,1.63)	1.29*	(0.12,2.45)
Location of residence										
Rural										
Urban	-1.39*	(-2.70,-0.07)	0.79	(0.52,1.18)	-0.41	(-1.41,0.59)	-0.31	(-0.73,0.12)	-0.60*	(-1.19,0.00)
Chronic diseases										
0										
1	-0.93	(-2.58,0.71)	1.45	(0.86,2.44)	-0.60	(-1.87,0.67)	-0.25	(-0.75,0.25)	-0.58	(-1.31,0.16)
2	0.28	(-1.33,1.88)	1.84*	(1.11,3.04)	-0.06	(-1.35,1.24)	-0.08	(-0.58,0.42)	-0.15	(-0.98,0.68)
3	1.24	(-0.68,3.16)	2.86**	(1.58,5.18)	0.93	(-0.51,2.36)	0.03	(-0.58,0.65)	-0.22	(-1.11,0.67)
≥4	5.25*	(3.29,7.21)	8.91**	(4.75,16.72)	3.50**	(2.06,4.95)	1.18**	(0.55,1.80)	1.03*	(0.07,1.98)

* $p < 0.05$; ** $p < 0.01$; UEBMI, Urban Employee Basic Medical Insurance; URRBMI, Urban-Rural Residents Basic Medical Insurance

fracture in both the areas of psychological health, as measured by depression symptoms, and physical functions, such as people's mobility and daily activities. Studies that investigated educational disparity found that there was an additional 7–11% loss in HRQoL among those with a lower educational level [23, 26, 37]. A similar association was found in other disease areas, such as cardiovascular diseases and COPD [16, 38]. Potential explanations for this trend include the existence of less access to disease-prevention knowledge, less guidance regarding a healthy lifestyle among lower educated individuals, and consequent less access to good quality care and rehabilitation [39]. However, most previous research mainly focused on overall HRQoL or physical functions; a single study that analyzed post-fracture mental HRQoL indicated a higher level of HRQoL loss in a lower- or medium-educated population but not those having higher educational levels [40]. The observed educational disparity in our study supported these findings and helps to direct attention toward mental HRQoL in hip fracture research.

Comorbidities are prevalent among the elderly, as over half of them have ≥ 2 chronic diseases [41], and they have been widely reported to be negatively correlated with individual HRQoL in either chronic disease people or the general population [41]. As comorbidities usually predict a worse treatment outcome, less function recovery, and more health care needs [42], understanding their association with HRQoL is of major importance for targeted policy-making and intervention implementation. In hip fracture research, an obviously higher risk of mortality in participants having comorbidities was reported by investigators [43]. Our study adds to the evidence on HRQoL, reporting that study participants with hip fracture had an increased risk by 5 times for depression, 9 times for body pain, 4 times for mobility difficulty, and 1 time for daily activities ability loss, respectively, if they suffered from more chronic diseases. Particular attention should be paid to individuals suffering from both a past hip fracture and comorbidities to improve their health status and life quality.

Another interesting finding in this study was the observed rural–urban disparities in HRQoL after hip fracture. We recorded a substantially higher HRQoL among people living in urban places compared to those residing in rural areas after controlling other covariates, especially in the areas of depression symptoms and IADL. A similar disparity has been found in the context of other chronic diseases like hypertension and cancer [17, 44] and in a general Chinese elderly adult population, where rural residents had lower HRQoL [45]. Urbanization usually comes with timely medical diagnosis and treatment, better access to rehabilitation care, and more resources of health management knowledge [46], potentially facilitating the provision of high-level health care to study participants with hip fracture. With some evidence from other diseases, however, studies

investigating rural–urban disparities in hip fracture have not been consistent with these results. Notwithstanding well-documented evidence that hip fracture incidence rates are higher in urban regions [47], data from developed countries like Norway, Canada, and the USA showed no rural–urban difference [48] or even a lower mortality rate in rural areas [35, 49]. However, considering the cultural differences and region characteristics across these countries and the indicators used to evaluate health outcomes, a cross-national comparison should be performed very cautiously. As there is no agreement on rural–urban differences in HRQoL after hip fracture, future studies are needed to provide clarification.

A key strength of this study is that we included pre-fracture HRQoL data so that the real condition of participants before and after their fracture could be evaluated. As seen from previous studies, most HRQoL measurement tools lack the ability to compare to baseline [50]. Therefore, most prior studies did not report pre-fracture quality of life data, while other scholars used retrospective ones [8], potentially leading to recall bias [50]. Using a longitudinal survey, we acquired more accurate HRQoL data before the hip fracture. Another advantage is the DID analysis method used in this study. Compared to previous studies that only reported correlations, this study explored the causality of the significant impact of hip fracture in all HRQoL dimensions.

Still, limitations to our study should be noted. First, this study failed in evaluating the precise HRQoL changes at specific time-points after hip fracture due to data unavailability, instead, we only know that hip fracture occurred within last 2–3 years; therefore, we could not treat time after fracture as a continuous variable. Second, we could not identify whether a reported hip fracture was the first fracture or a re-fracture, and these events might have different impacts on HRQoL. If a participant reported a hip fracture in both the 2015 and 2018 waves, we identified them as 2 independent hip fracture participants, which could cause misestimation or bias. Third, the CHARLS only included study participants aged ≥ 45 years, so our results cannot be generalized to the younger population. Nevertheless, as hip fracture is more prevalent with age, our results might still represent the hip fracture population. Fourth, the study participants of CHARLS are representative for general middle-aged and elderly people in China [18]. However, considering that hip fracture is not evenly distributed in the general population, data used in this study may not completely represent the hip fracture population in China. Fifth, this study did not use a standard HRQoL instrument to evaluate the post-hip-fracture HRQoL, and 31 items of HRQoL questions were selected from the CHARLS to reflect those on the EQ-5D questionnaire. The reliability and validity of the selected questions as an HRQoL instrument need to be tested in future studies. While a similar approach has been employed in the previous study [51], the validity in evaluating the

composite HRQoL in study participants with hip fracture is to be tested. Finally, the hip fracture occurrence was self-reported by study participants. Hip fracture usually causes symptoms and severe health consequences, therefore most people are hospitalized or receive medical treatment after a hip fracture. There is little risk of reporting bias to identify study participants with hip fracture.

In order to gain a better understanding of the socio-demographic disparities of post-hip fracture HRQoL, further studies should be performed. First, HRQoL disparity at specific time points after hip fracture should be further analyzed to provide more accurate evidence about the potential effective intervention implementation. Second, as many findings were gathered from Western countries [8, 40], additional studies in China as well as cross-national comparisons should also be performed to identify country-specific characteristics for the purpose of developing targeted policies.

Practice implications

Hip fractures have remarkably negative effects on individual HRQoL even after 2–3 years. We found that a lower HRQoL existed among people living in rural areas, with less educational attainment, and with more comorbidities. Hip fracture people who meet these conditions usually have less material resources and less access to health knowledge and professional medical care [52]. Targeted policies should strive to improve HRQoL equities—that is, to prevent more physical function reduction or mental health loss. For instance, integration of Urban Residents Basic Medical Insurance which covered urban population and New Rural Cooperative Medical Schemes covering rural residents has been shown to promote both financial and geographic access to healthcare utilization and improve its equity between rural and urban areas in China [53]; nonetheless, more effort should be made to enhance their access to high-quality care. For hip fracture people who suffered from other chronic diseases, integrated health services, instead of a simple treatment of diseases episode by episode, are of great importance. Multidisciplinary healthcare teams should be developed to enable coordinated management and treatment of coexisting diseases [54]. Lessons from other diseases could also be borrowed. Policies increasing healthcare access in rural areas that could potentially improve its utilization [55], health education programs targeting the elderly aimed at equipping them with essential knowledge to control disease [56] and disease-screening strategies for older adults [57] have been proven to remarkably improve health outcomes such as reducing mortality, and to achieve better HRQoL. However, due to the limited available literature, more evidence is still needed concerning the intervention effect in post-hip fracture people.

Supplementary Information The online version contains supplementary material available at <https://doi.org/10.1007/s11657-023-01220-2>.

Author contribution Conceptualization and methodology: Lizheng Xu, Stephen Jan, Mingsheng Chen, and Lei Si; study supervision: Lei Si; formal analysis and investigation: Lizheng Xu; Lizheng Xu prepared the original draft; all authors reviewed, edited, and approved the final manuscript.

Funding Lizheng Xu is supported by an Australian Government Research Training Program Scholarship. Lei Si is supported by a WSU Research Theme Funding. This study was funded by the National Natural Science Foundation of China (grant number: 71874086, 72174093).

Data availability The data that support the findings of this study are openly available in China Health and Retirement Longitudinal Study (CHARLS) website at <http://charls.pku.edu.cn/>.

Declarations

Ethics approval and consent to participate The current study is a secondary analysis of the de-identified China Health and Retirement Longitudinal Study (CHARLS) public data. The original CHARLS was approved by the Ethical Review Committee of Peking University, and all participants signed the informed consent at the time of participation.

Competing interest Lizheng Xu, Mingsheng Chen, Ke Peng, Mickael Hilgsmann, Stephen Jan, and Lei Si declare that they have no conflict of interest.

References

1. Kanis JA, Odén A, McCloskey EV, Johansson H, Wahl DA, Cooper C (2012) A systematic review of hip fracture incidence and probability of fracture worldwide. *Osteoporos Int* 23:2239–2256
2. Hernlund E, Svedbom A, Ivergård M, Compston J, Cooper C, Stenmark J, McCloskey EV, Jönsson B, Kanis JA (2013) Osteoporosis in the European Union: medical management, epidemiology and economic burden. A report prepared in collaboration with the International Osteoporosis Foundation (IOF) and the European Federation of Pharmaceutical Industry Associations (EFPIA). *Arch Osteoporos* 8:136
3. Si L, Winzenberg TM, Chen M, Jiang Q, Palmer AJ (2015) Residual lifetime and 10 year absolute risks of osteoporotic fractures in Chinese men and women. *Curr Med Res Opin* 31:1149–1156
4. Si L, Winzenberg TM, Jiang Q, Chen M, Palmer AJ (2015) Projection of osteoporosis-related fractures and costs in China: 2010–2050. *Osteoporos Int* 26:1929–1937
5. Guzon-Illescas O, Perez Fernandez E, Crespí Villarias N, Quirós Donate FJ, Peña M, Alonso-Blas C, García-Vadillo A, Mazzucchelli R (2019) Mortality after osteoporotic hip fracture: incidence, trends, and associated factors. *J Orthop Surg Res* 14:203
6. Katsoulis M, Benetou V, Karapetyan T et al (2017) Excess mortality after hip fracture in elderly persons from Europe and the USA: the chances project. *J Intern Med* 281:300–310
7. Bliuc D, Nguyen ND, Milch VE, Nguyen TV, Eisman JA, Center JR (2009) Mortality risk associated with low-trauma osteoporotic fracture and subsequent fracture in men and women. *JAMA* 301:513–521
8. Svedbom A, Borgstöm F, Hernlund E et al (2018) Quality of life for up to 18 months after low-energy hip, vertebral, and

- distal forearm fractures-results from the ICUROS. *Osteoporos Int* 29:557–566
9. van de Ree CLP, Landers MJF, Kruithof N, de Munter L, Slaets JPI, Gosens T, de Jongh MAC (2019) Effect of frailty on quality of life in elderly patients after hip fracture: a longitudinal study. *BMJ Open* 9:e025941
 10. Gjertsen JE, Baste V, Fevang JM, Furnes O, Engesaeter LB (2016) Quality of life following hip fractures: results from the Norwegian hip fracture register. *BMC Musculoskelet Disord* 17:265
 11. Si L, Winzenberg TM, de Graaff B, Palmer AJ (2014) A systematic review and meta-analysis of utility-based quality of life for osteoporosis-related conditions. *Osteoporos Int* 25:1987–1997
 12. Peeters CM, Visser E, Van de Ree CL, Gosens T, Den Oudsten BL, De Vries J (2016) Quality of life after hip fracture in the elderly: A systematic literature review. *Injury* 47:1369–1382
 13. Gold T, Williams SA, Weiss RJ, Wang Y, Watkins C, Carroll J, Middleton C, Silverman S (2019) Impact of fractures on quality of life in patients with osteoporosis: a US cross-sectional survey. *J Drug Assess* 8:175–183
 14. Shyu YI, Chen MC, Liang J, Lu JF, Wu CC, Su JY (2004) Changes in quality of life among elderly patients with hip fracture in Taiwan. *Osteoporos Int* 15:95–102
 15. Koh DTS, Chen JY, Yew AKS, Chong HC, Hao Y, Pang HN, Tay DKJ, Chia SL, Lo NN, Yeo SJ (2019) Functional outcome and quality of life in patients with hip fracture after total knee arthroplasty. *J Orthop Surg (Hong Kong)* 27:2309499019852338
 16. Miravittles M, Naberan K, Cantoni J, Azpeitia A (2011) Socioeconomic status and health-related quality of life of patients with chronic obstructive pulmonary disease. *Respiration* 82:402–408
 17. Moss JL, Pinto CN, Mama SK, Rincon M, Kent EE, Yu M, Cronin KA (2021) Rural-urban differences in health-related quality of life: patterns for cancer survivors compared to other older adults. *Qual Life Res* 30:1131–1143
 18. Zhao Y, Hu Y, Smith JP, Strauss J, Yang G (2014) Cohort profile: the China Health and Retirement Longitudinal Study (CHARLS). *Int J Epidemiol* 43:61–68
 19. Parsons N, Griffin XL, Achten J, Costa ML (2014) Outcome assessment after hip fracture: is EQ-5D the answer? *Bone Joint Res* 3:69–75
 20. Balestroni G, Bertolotti G (2012) EuroQol-5D (EQ-5D): an instrument for measuring quality of life. *Monaldi Arch Chest Dis* 78:155–159
 21. (2002) Occupational therapy practice framework: domain and process. *Am J Occup Ther* 56(6):609–639
 22. Björqvinnson T, Kertz SJ, Bigda-Peyton JS, McCoy KL, Aderka IM (2013) Psychometric properties of the CES-D-10 in a psychiatric sample. *Assessment* 20:429–436
 23. Abimanyi-Ochom J, Watts JJ, Borgström F et al (2015) Changes in quality of life associated with fragility fractures: Australian arm of the International Cost and Utility Related to Osteoporotic Fractures Study (AusICUROS). *Osteoporos Int* 26:1781–1790
 24. Zhao H, Wei P, Feng Y (2022) Association between frailty and clinical outcomes and quality of life in older adults following hip fracture surgery: a retrospective cohort study. *Anesth Analg* 134(5):1035–1042
 25. Chiu MH, Hwang HF, Lee HD, Chien DK, Chen CY, Lin MR (2012) Effect of fracture type on health-related quality of life among older women in Taiwan. *Arch Phys Med Rehabil* 93:512–519
 26. Guirant L, Carlos F, Curriel D, Kanis JA, Borgström F, Svedbom A, Clark P (2018) Health-related quality of life during the first year after a hip fracture: results of the Mexican arm of the International Cost and Utility Related to Osteoporotic Fractures Study (MexICUROS). *Osteoporos Int* 29:1147–1154
 27. Zhao Y, Atun R, Oldenburg B, McPake B, Tang S, Mercer SW, Cowling TE, Sum G, Qin VM, Lee JT (2020) Physical multimorbidity, health service use, and catastrophic health expenditure by socioeconomic groups in China: an analysis of population-based panel data. *Lancet Glob Health* 8:e840–e849
 28. Gu J, Chao J, Chen W, Xu H, Zhang R, He T, Deng L (2018) Multimorbidity and health-related quality of life among the community-dwelling elderly: A longitudinal study. *Arch Gerontol Geriatr* 74:133–140
 29. Amarilla-Donoso FJ, López-Espuela F, Roncero-Martín R, Leal-Hernandez O, Puerto-Parejo LM, Aliaga-Vera I, Toribio-Felipe R, Lavado-García JM (2020) Quality of life in elderly people after a hip fracture: a prospective study. *Health Qual Life Outcomes* 18:71
 30. Amarilla-Donoso FJ, Roncero-Martín R, Lavado-García JM, Toribio-Felipe R, Moran-García JM, Lopez-Espuela F (2020) Quality of life after hip fracture: a 12-month prospective study. *PeerJ* 8:e9215
 31. Milte R, Crotty M, Miller MD, Whitehead C, Ratcliffe J (2018) Quality of life in older adults following a hip fracture: an empirical comparison of the ICECAP-O and the EQ-5D-3 L instruments. *Health Qual Life Outcomes* 16:173
 32. Ekström W, Al-Ani AN, Sääf M, Cederholm T, Ponzer S, Hedström M (2013) Health related quality of life, reoperation rate and function in patients with diabetes mellitus and hip fracture—a 2 year follow-up study. *Injury* 44:769–775
 33. Elboim-Gabyzon M, Andrawus Najjar S, Shtarker H (2019) Effects of transcutaneous electrical nerve stimulation (TENS) on acute postoperative pain intensity and mobility after hip fracture: A double-blinded, randomized trial. *Clin Interv Aging* 14:1841–1850
 34. Herrick C, Steger-May K, Sinacore DR, Brown M, Schechtman KB, Binder EF (2004) Persistent pain in frail older adults after hip fracture repair. *J Am Geriatr Soc* 52:2062–2068
 35. Solbakken SM, Magnus JH, Meyer HE et al (2019) Urban-Rural Differences in Hip Fracture Mortality: A Nationwide NOREPOS Study. *JBMR Plus* 3:e10236
 36. Sanzone AG (2016) Current Challenges in Pain Management in Hip Fracture Patients. *J Orthop Trauma* 30(Suppl 1):S1-5
 37. Marques A, Lourenço Ó, da Silva JA (2015) The burden of osteoporotic hip fractures in Portugal: costs, health related quality of life and mortality. *Osteoporos Int* 26:2623–2630
 38. Tchicaya A, Lorentz N (2016) Socioeconomic inequalities in health-related quality of life between men and women, 5 years after a coronary angiography. *Health Qual Life Outcomes* 14:165
 39. JieAnNaMu XuX, You H, Gu H, Gu J, Li X, Cui N, Kou Y (2020) Inequalities in health-related quality of life and the contribution from socioeconomic status: evidence from Tibet. *China BMC Public Health* 20:630
 40. Valentin G, Pedersen SE, Christensen R, Friis K, Nielsen CP, Bhimjiyani A, Gregson CL, Langdahl BL (2020) Socio-economic inequalities in fragility fracture outcomes: a systematic review and meta-analysis of prognostic observational studies. *Osteoporos Int* 31:31–42
 41. Makovski TT, Schmitz S, Zeegers MP, Stranges S, van den Akker M (2019) Multimorbidity and quality of life: Systematic literature review and meta-analysis. *Ageing Res Rev* 53:100903
 42. Fisher K, Griffith L, Gruneir A, Panjwani D, Gandhi S, Sheng LL, Gafni A, Chris P, Markle-Reid M, Ploeg J (2016) Comorbidity and its relationship with health service use and cost in community-living older adults with diabetes: A population-based study in Ontario, Canada. *Diabetes Res Clin Pract* 122:113–123
 43. González-Zabaleta J, Pita-Fernandez S, Seoane-Pillado T, López-Calviño B, Gonzalez-Zabaleta JL (2016) Comorbidity as

- a predictor of mortality and mobility after hip fracture. *Geriatr Gerontol Int* 16:561–569
44. Hou Y, Wu Q, Zhang D, Jin X, Wu W, Wang X (2020) The differences in self-perceptions of aging, health-related quality of life and their association between urban and rural Chinese older hypertensive patients. *Health Qual Life Outcomes* 18:154
 45. Zhou Z, Zhou Z, Gao J, Lai S, Chen G (2018) Urban-rural difference in the associations between living arrangements and the health-related quality of life (HRQOL) of the elderly in China—Evidence from Shaanxi province. *PLoS One* 13:e0204118
 46. Zhao Y, Xu X, Dupre ME, Xie Q, Qiu L, Gu D (2020) Individual-level factors attributable to urban-rural disparity in mortality among older adults in China. *BMC Public Health* 20:1472
 47. Brennan SL, Pasco JA, Urquhart DM, Oldenburg B, Hanna FS, Wluka AE (2010) The association between urban or rural locality and hip fracture in community-based adults: a systematic review. *J Epidemiol Community Health* 64:656–665
 48. Weller I, Wai EK, Jaglal S, Kreder HJ (2005) The effect of hospital type and surgical delay on mortality after surgery for hip fracture. *J Bone Joint Surg Br* 87:361–366
 49. Miller BJ, Cai X, Cram P (2012) Mortality rates are similar after hip fractures for rural and urban patients. *Clin Orthop Relat Res* 470:1763–1770
 50. Lawson A, Tan AC, Naylor J, Harris IA (2020) Is retrospective assessment of health-related quality of life valid? *BMC Musculoskelet Disord* 21:415
 51. Wang L, Liu W (2022) Effects of Family Doctor Contract Services on the Health-Related Quality of Life Among Individuals With Diabetes in China: Evidence From the CHARLS. *Front Public Health* 10:865653
 52. Kim JH, Park EC (2015) Impact of socioeconomic status and subjective social class on overall and health-related quality of life. *BMC Public Health* 15:783
 53. Li C, Tang C, Wang H (2019) Effects of health insurance integration on health care utilization and its equity among the mid-aged and elderly: evidence from China. *Int J Equity Health* 18:166
 54. Egede LE (2006) Disease-focused or integrated treatment: diabetes and depression. *Med Clin North Am* 90:627–646
 55. Arcury TA, Preisser JS, Gesler WM, Powers JM (2005) Access to transportation and health care utilization in a rural region. *J Rural Health* 21:31–38
 56. Chen Y, Li X, Jing G, Pan B, Ge L, Bing Z, Yang K, Han X (2020) Health education interventions for older adults with hypertension: A systematic review and meta-analysis. *Public Health Nurs* 37:461–469
 57. Si L, Winzenberg TM, Chen M, Jiang Q, Neil A, Palmer AJ (2016) Screening for osteoporosis in Chinese post-menopausal women: a health economic modelling study. *Osteoporos Int* 27:2259–2269

Publisher's note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Springer Nature or its licensor (e.g. a society or other partner) holds exclusive rights to this article under a publishing agreement with the author(s) or other rightsholder(s); author self-archiving of the accepted manuscript version of this article is solely governed by the terms of such publishing agreement and applicable law.