

# Shifting the perspective

Citation for published version (APA):

Boselie, J. J. L. M., & Peters, M. L. (2023). Shifting the perspective: how positive thinking can help diminish the negative effects of pain. *Scandinavian Journal of Pain*, 23(3), 452-463.  
<https://doi.org/10.1515/sjpain-2022-0129>

**Document status and date:**

Published: 26/07/2023

**DOI:**

[10.1515/sjpain-2022-0129](https://doi.org/10.1515/sjpain-2022-0129)

**Document Version:**

Publisher's PDF, also known as Version of record

**Document license:**

Taverne

**Please check the document version of this publication:**

- A submitted manuscript is the version of the article upon submission and before peer-review. There can be important differences between the submitted version and the official published version of record. People interested in the research are advised to contact the author for the final version of the publication, or visit the DOI to the publisher's website.
- The final author version and the galley proof are versions of the publication after peer review.
- The final published version features the final layout of the paper including the volume, issue and page numbers.

[Link to publication](#)

**General rights**

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal.

If the publication is distributed under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license above, please follow below link for the End User Agreement:

[www.umlib.nl/taverne-license](http://www.umlib.nl/taverne-license)

**Take down policy**

If you believe that this document breaches copyright please contact us at:

[repository@maastrichtuniversity.nl](mailto:repository@maastrichtuniversity.nl)

providing details and we will investigate your claim.

## Topical Review

Jantine J.L.M. Boselie\* and Madelon L. Peters

# Shifting the perspective: how positive thinking can help diminish the negative effects of pain

<https://doi.org/10.1515/sjpain-2022-0129>

Received September 1, 2022; accepted February 2, 2023;

published online February 20, 2023

### Abstract

**Objectives:** The field of pain psychology has taken significant steps forward during the last decades and the way we think about how to treat chronic pain has radically shifted from a biomedical perspective to a biopsychosocial model. This change in perspective has led to a surge of accumulating research showing the importance of psychological factors as determinants for debilitating pain. Vulnerability factors, such as pain-related fear, pain catastrophizing and escape/avoidant behaviours may increase the risk of disability. As a result, psychological treatment that has emerged from this line of thinking has mainly focused on preventing and decreasing the adverse impact of chronic pain by reducing these negative vulnerability factors. Recently, another shift in thinking has emerged due to the field of positive psychology, which aims to have a more complete and balanced scientific understanding of the human experience, by abandoning the exclusive focus on vulnerability factors towards including protective factors.

**Methods:** The authors have summarised and reflected on the current state-of-the-art of pain psychology from a positive psychology perspective.

**Results:** Optimism is an important factor that may in fact buffer and protect against pain chronicity and disability. Resulting treatment approaches from a positive psychology perspective are aimed at increasing protective factors, such as optimism, to increase resilience towards the negative effects of pain.

**Conclusions:** We propose that the way forward in pain research and treatment is the inclusion of both *vulnerability* and *protective factors*. Both have unique roles in modulating

the experience of pain, a finding that had been neglected for too long. Positive thinking and pursuing valued goals can make one's life gratifying and fulfilling, despite experiencing chronic pain.

**Keywords:** (chronic) pain; biopsychosocial model; optimism; positive affect; positive psychology; protective factor.

## From a biomedical to a biopsychosocial perspective

Our thinking about how to treat chronic pain has radically shifted. For a long time, the experience of chronic pain was thought of solely as a medical problem. Pain was considered a direct sign of an underlying injury, tissue damage, or organic pathology and, as such, should be treated medically. However, numerous studies illustrate that the report of pain, either acute or chronic, is not strongly correlated with the presence of tissue damage. For instance, in a meta-analysis, findings showed that many degenerative spine findings (e.g. disc bulges) are also found in uninjured individuals who do not report lower back pain, with the prevalence rising with age [1]. This suggests that medical findings are not always related to the pain experience. Furthermore, most individuals who experience chronic pain and seek help from a primary care provider rarely have a previously undiagnosed serious pathology or experienced an injury [2]. Moreover, similar pain phenotypes occur in individuals in the absence of an identifiable aetiological trigger (i.e. injury or medical disease), such as fibromyalgia, irritable bowel syndrome, and tension headaches, which may reflect changes in pain sensitivity due to central sensitization (i.e. an amplification of neural signalling within the central nervous system that elicits pain hypersensitivity) [3, 4]. After the discovery of central sensitization, it became clear that experiencing a noxious stimulus is not necessary to produce a pain experience [3].

These medically unexplained experiences of pain fuelled a shift in thinking from a solely biomedical perspective to a biopsychosocial perspective, resulting in influential

\*Corresponding author: Jantine J.L.M. Boselie, Clinical Psychology, Open Universiteit, Heerlen, The Netherlands, E-mail: jantine.boselie@ou.nl

<https://orcid.org/0000-0003-2490-3741>

Madelon L. Peters, Clinical Psychological Science, Maastricht University, Maastricht, The Netherlands. <https://orcid.org/0000-0003-2969-6286>

biopsychosocial models such as the cognitive-behavioural fear-avoidance model [5]. Central in the fear-avoidance model is the dysfunctional interpretation of an acute pain experience. People who tend to have catastrophic thoughts concerning their pain are more likely to get caught in a downward vicious circle of pain-related fear, defensive behaviours, disuse and disability, and, in turn, increased pain [5]. This shift from a biomedical to a biopsychosocial conceptualization of pain led to a surge of research that however, predominantly focused on negative factors, such as negative cognitions (e.g. pain catastrophizing, negative attention bias, fear-avoidance beliefs) and behaviours (e.g. avoidance).

This focus on negative factors has guided numerous studies and increased our knowledge about vulnerability factors that contribute to or maintain chronic pain. As a result, psychological treatment has concentrated on decreasing the adverse impact of chronic pain by reducing these negative vulnerability factors, such as by means of cognitive behavioural therapy (CBT) or exposure techniques [6, 7]. CBT is mainly directed at changing maladaptive cognitions into more benign or helping cognitions [7, 8]. A systematic review [9] indicated that CBT is an effective treatment for adult chronic pain, leading to decreased disability, distress, and pain. However, the effect sizes across all outcomes are modest, ranging from very small to small. In exposure therapy, patients are asked to perform an activity to challenge the validity of their catastrophic expectancies, and to encourage explorative behaviour, enabling recovery [10]. Exposure treatments can effectively reduce pain-related fear and catastrophic expectancies about physical activities and have been found cost-effective [11–14]. Nevertheless, effect sizes for treatment outcomes are small [15] and exposure treatments are associated with high drop-out rates of 30–50% [6]. In sum, there are effective psychological treatments developed for chronic pain, but effect sizes suggest that there is room for improvement.

## From a negative towards a positive focus

At the turn of the century, a new way of thinking emerged that contributed to a new field within psychology: positive psychology [16]. The aim of positive psychology is to have a more complete and balanced scientific understanding of the human experience by abandoning the exclusive focus on vulnerability factors and ‘fixing what is wrong’ and including protective factors and ‘building what is strong’ [17, 18]. This new approach also found its way into the field

of pain [19, 20]. Protective factors, for instance, dampen pain intensity, diminish the negative side effects of a pain experience, protect against developing chronic pain, or promote adaptation to chronic pain. Positive affect, self-efficacy (i.e. perceived ability to learn or perform behaviour to attain designated goals) [21], hope (i.e. belief that one can find pathways to desired goals and become motivated to use those pathways) [22], and optimism (i.e. generalised and global positive outcome expectancies) [23] are examples of protective factors that are shown to promote pain resilience [19, 24–30]. A review of the empirical literature shows that self-efficacy, hope and optimism are structurally distinct and differentially related to important life outcomes, including psychological adjustment [31]. Better chronic pain adjustment by increasing positive factors is in line with the influential broaden-and-build theory of Fredrickson [32]. This theory states that experiencing and strengthening positive behaviours, emotions, and cognitions cause an upward, mutually reinforcing spiral to ensue, in which broadened thinking patterns and building of durable resources result in further positive emotions, in turn enhancing emotional well-being [32]. Examining both vulnerability and protective factors will help to disentangle what leads to outcomes of recovery, sustainability (perseverance in valued activities despite chronic pain), and growth (benefit finding due to adversity, such as experiencing chronic pain) [33].

It should be noted that the fear-avoidance model of chronic pain [5] specifies not only a negative downward spiral, but also a positive upward spiral. When a non-threatening appraisal of pain is made that does not elicit fear, this is assumed to result in confrontation and engagement behaviour, resulting in functional recovery. Originally, the model did not specify any factor, besides the absence of fear that could elicit this upward spiral to recovery. However, since its conception, suggestions to expand the fear-avoidance model have been proposed. In fact, the authors themselves never presumed the model was finished, but saw it as a theory-based heuristic to guide further research in an attempt to gain a better understanding of pain disability and improving existing treatment protocols [10]. It has been argued that in order to understand chronic pain, examining the motivation of an individual and taking the (social) context into account are essential [34–37]. Chronic pain is not experienced in isolation. Chronic pain always occurs within a context of daily life, with various constraints and demands of its own, often resulting in competing goals. Fear and avoidance behaviour operate in this context of multiple competing goals, influencing an individual’s motivation to pursue certain valued activities at one point or abandon them at other times to cope with a chronic pain experience [34]. Protective factors may promote the pursuit

of valued life goals, in contrast to a persistent attempt to control or to avoid pain [34, 38, 39]. For an extensive review of the attenuating effect of positive affect on pain, see Hanssen et al. [40]. In this topical review, we will focus on optimism, given the theoretical underpinnings of the protective effect of optimism in the context of pain. We will first review the evidence that optimism can help alleviate the negative impact of pain and whether it is possible to increase optimism. Then, we will discuss the clinical implications of increasing optimism in chronic pain patients and possible interventions.

## How optimism helps diminish the negative effects of pain

Dispositional optimism is characterised by generalised and global positive outcome expectancies [23]. In other words, individuals with high levels of optimism have the tendency to expect that good things will happen in the future. This definition is based on the expectancy-value model of motivation [41]. It assumes that behaviour reflects the pursuit of goals, which is defined as internal representations of a desired state [42]. The model proposes two aspects that influence goal-directed behaviour. First, if a goal is perceived as more important, the *value* that is given is greater, resulting in more effort displayed to attain the goal. Second, motivation is determined by the *expectancy* of goal attainment. Doubt may increase withdrawal from effort towards goal attainment, while confidence in reaching a goal will foster perseverance and continued effort, even when it is difficult to carry on. Optimism is in this case distinct from self-efficacy in that self-efficacy is only one potential source for the expectancy of goal attainment. Other possible sources that may contribute to these expectancies of goal attainment are for instance help from others, fate or a benevolent god.

There are several possible explanations why optimism can diminish the negative and debilitating impact of chronic pain. The most intuitive explanation is that optimism reduces the experienced intensity of pain [43]. Recently, a meta-analysis [44] was conducted to examine the relationship between optimism and pain-related outcomes. Results showed that optimism was related to lower pain severity, better physical functioning, and less psychological dysfunction (i.e. depressive or anxiety symptoms). A systematic review [25] was conducted on experimental and clinical studies investigating the relation between optimism and the experience of pain. Almost 70% of the studies found a significant beneficial association between optimism and at least one pain characteristic (i.e. pain intensity, frequency or

unpleasantness, pain tolerance thresholds, habituation, temporal summation, conditioned pain modulation, or placebo/nocebo effects). Furthermore, optimism has been shown to increase the responsiveness to placebo-induced pain relief [45].

Optimism may also prevent individuals from being caught in a downward spiral of fear-avoidance that fuels negative pain-related outcomes. Optimism is associated with less pain catastrophizing [19, 33], which may prevent the downward spiral through a non-threatening interpretation of pain-related stimuli [46, 47]. Vancleef and Peters [48] reported that optimistic individuals make less negative or catastrophic interpretations of ambiguous information in various domains, such as pain-related sensations, other bodily sensations, and social and general situations. Alternatively, optimism may broaden attention by reducing pain-induced distressing thoughts and rumination that consume attention [49–52]. As optimism is associated with less worry, rumination, and pain catastrophizing [46, 53–55], all available attentional resources can be dedicated towards continuing pursuing valued activities and/or actively coping with the pain experience. Research findings have indeed indicated that the association between optimism and the experience of less pain is mediated by coping strategies. Optimism is associated with more active coping, which, in turn, is associated with low levels of pain, anxiety, depression, and impairment. Pessimism is related to the use of passive coping, which is associated with high levels of pain, anxiety, depression, and impairment [56].

Another potential mechanism of the beneficial effects of optimism is increased goal attainment. Optimism may diminish the negative impact of chronic pain on an individual's life by increasing motivation towards continuing effort and engagement towards (valued) goal attainment [57]. Self-regulation resources are considered to be limited. Self-regulation is the ability to control or alter emotions and thoughts, enabling behaviour to vary adaptively from moment to moment [41, 58]. When an individual is confronted with a pain experience, resources are directed towards the pain in order to cope with this aversive experience. However, directing these resources towards the pain demand will render them unavailable for additional demands, causing resources to be fatigued [59–61]. Optimism may increase motivation to continue allocating resources towards valued activities, despite fewer resources being available [59, 62]. Optimists indeed seem to invest more effort to reach personal and health goals despite pain or limitations [63, 64]. But this increased motivation by optimism may only occur when the activity is perceived as important and personally relevant [37]. Interestingly, when the decision to avoid potential injury is outweighed by

the motivation to pursue a conflicting goal that is perceived as important and personally relevant, top-down processes in the brain may inhibit nociception (i.e. the neurophysiological process of encoding noxious stimuli that produce actual or potential tissue injury) [65] via the release of opioid peptides. In contrast, if the individual decides to ignore the conflicting goal and instead respond to the noxious stimulus, top-down facilitation of nociception can occur [66].

The ability to adjust goals when goals have become unattainable due to the chronic pain experience is also important [67, 68]. One might argue that this higher tendency to pursue goal achievement may in fact be potentially harmful for chronic pain patients. Pain itself is often not alleviated; thus, pursuing the goal to abolish the pain experience may lead to disappointment and increased likelihood of experiencing goal conflict [69]. Goal conflict arises when pursuing one goal is at the cost of other goals [69] and results in decrements in overall well-being [70, 71]. However, optimism is associated with more flexible goal adjustment [72–75]. Chronic pain patients who report higher levels of optimism seem to be more flexible when confronted with competing goals, re-engage more easily with new attainable goals, and behave tenaciously with valued goals [76]. This tendency of optimistic individuals to show goal adjustment strategies such as re-engagement, flexibility, and tenacity is associated with higher levels of purpose in life, which, in turn, are strongly associated with adaptation in patients with chronic pain [76]. Additionally, optimistic individuals are more flexible in applying different coping strategies [57, 62, 77, 78]. It is proposed that optimists in general are more inclined to focus their attention on positive aspects of the situation, but focus on the pain whenever it is relevant for their well-being and action is required, resulting in active and problem-focused tendencies [27, 57, 78–80]. In an eye-tracker study, results showed that optimists indeed tended to turn away from negative information (i.e. angry faces) and gazed more at positive information (i.e. joyful faces) [81]. However, when faced with a serious threat to their well-being (e.g. an operation or cancer), optimists tend to focus on the context and pain instead of withdrawing from them [82]. Furthermore, studies have indicated that optimists are less vulnerable to experiencing disappointment, even after receiving a cancer diagnosis [83], undergoing cancer treatment [84, 85], or experiencing a failed *in vitro* fertilisation attempt [86].

## Increasing optimism

In the previous section, we made a case for the importance of optimism as a protective factor in the context of pain.

However, one can wonder about the practical and clinical implications of the knowledge that optimism can act as a protective factor. Do chronic pain sufferers with a more pessimistic inclination just have bad luck, or can we help them to think in a more optimistic way? We know that both heritable [87, 88] and environmental factors (e.g. secure attachment to caregivers, self-esteem, and financial security) [89–91] contribute to one's level of optimism. Although optimism has been defined as an enduring personality trait, optimism levels can change over time [92, 93]. More importantly, optimism is modifiable by psychological interventions [94]. One of the most powerful and best researched interventions to increase optimism is the Best Possible Self (BPS) exercise [95]. The BPS is a positive future-thinking technique in which participants are asked to write about and imagine a life in the future where everything turned out for the best. Numerous studies and a recent meta-analysis demonstrated that the BPS manipulation is effective in increasing optimism and positive affect [47, 96, 97] and decreasing goal ambivalence, which hampers positive emotions [98]. The BPS has been applied in different contexts (i.e. online or face-to-face), with different delivery modes (i.e. individually or in a group) and with different components, such as writing and imagery components or combining both [96, 99]. So far, the BPS manipulation has been predominately used in experimental studies to investigate the causal effects of optimism in the context of (experimentally) induced acute pain. For instance, in the study by Hanssen et al. [47], participants who received the BPS manipulation reported less pain intensity during a cold-water task compared to participants who received a neutral manipulation.

Optimism has also been demonstrated to protect against pain-induced executive task performance deterioration [100, 101]. Executive functioning is an umbrella term that describes multiple high-level cognitive processes and behavioural abilities [102]. It encompasses the ability to actively monitor behaviour, inhibit or facilitate certain responses, and optimise one's approach to unfamiliar circumstances [102]. Numerous studies have demonstrated that experimentally induced pain impairs executive task performance [e.g. 103–106]. Pain seems to be especially disruptive to task performance when the pain is novel and the threat-value and pain intensity are high [107, 108]. The first study to examine the effects of optimism in this context explored whether increasing optimism can counteract this sustained deteriorating effect of pain on executive task performance. Participants received either the BPS manipulation or a control task before the pain induction. Results showed that although pain significantly impaired executive task performance in the control condition, this sustained

deteriorating effect of pain on task performance was abolished in the optimism condition [100]. In another study, results showed that experimentally induced heat pain deteriorated concurrent task-shifting performance, but that inducing a temporary optimistic state also protected against this negative pain-induced concurrent task performance [101].

## Increasing optimism in chronic pain patients

Is the BPS exercise a suitable technique to use in patients with chronic pain? One could argue that writing and thinking about an ideal future may be daunting for a person suffering from chronic pain because it might trigger worries that the pain will get worse and becomes more disabling. This may in turn increase pain awareness [109]. Thus, one could imagine that the BPS exercise may potentially even be harmful for chronic pain patients when it increases the perceived discrepancy between the actual self (i.e. who you are now) and the ideal self (i.e. who you want to be) and decreases the perceived discrepancy between the actual self and the feared self (i.e. who you fear to be) [110, 111]. Self-discrepancies are associated with higher levels of pain, depression, anxiety, and distress, and lower levels of acceptance of pain in chronic pain patients [112–114]. This is especially true when the ideal self is conditional on the absence of pain (i.e. high self-pain enmeshment) [115–117]. Therefore, the BPS exercise may potentially increase the risk of maladjustment towards pain when self-discrepancies cause effortful attempts to pursue the (often) unachievable goal to abolish pain. Several studies have examined whether the BPS can be used in patients with chronic pain. Molinari and García-Palacios [118] used a multimedia system to deliver the BPS or a control intervention to 71 patients with fibromyalgia. No negative effects of BPS on patients' well-being were observed. Instead, at 3 months follow-up, patients reported higher levels of optimism and lower negative affect compared to patients in the control condition.

An intervention for chronic pain that has embedded the BPS exercise in a larger package of positive psychology exercises is called 'happy despite pain' [119]. In this intervention, the BPS followed after three other positive exercises that aimed at creating more openness in patients to engage with the BPS by increasing self-compassion and positive affect. The 'happy despite pain' intervention consists of four modules, namely self-compassion, positive focus, savouring,

and optimism, which are offered during an 8 weeks period. Module 1 targets self-compassion, which refers to fully accepting oneself, the ability to treat oneself with kindness rather than self-criticism, and recognising that everyone experiences failures [120, 121]. Module 2 requires patients to practice the 'three good things' exercise on a daily basis. Patients write down three good things that happened that day and why they happened. This exercise aims to shift the focus from a negative orientation towards a more positive one, by raising awareness of the good things in life [122, 123]. Module 3 contains savouring techniques that promote the frequency and intensity of positive experiences in daily life. Patients actively plan pleasant activities, so-called 'mini vacations' of 20 min. In these mini vacations, patients do something for themselves that they enjoy. In addition, patients are asked to actively recall pleasant memories and to write down everyday things that they enjoy in a savouring diary. First, patients formulate and write down their concrete and achievable goals, wishes and expectations for the future. Next, patients visualise this ideal future every day for 5 min, alternating between the three domains of their choice. Some precautions were taken to ward off potential negative consequences of the BPS. First, to prevent a heightened focus on pain and disability, patients were asked to focus on three out of eight pre-determined life domains (i.e. family life, romantic life, social life, professional life, leisure and hobby, educational life, societal commitment, and personal growth). Importantly, patients were asked to write down a detailed narrative of what their best possible future in these domains would look like, *despite pain*. Formulating a BPS narrative despite pain may increase the acceptance of the pain experience, decrease perceived self-discrepancies, and increase engagement in meaningful and valued activities [67].

Two successive studies in samples that predominantly consisted of fibromyalgia patients showed that the 'happy despite pain' intervention was effective in increasing optimism [119, 124]. In addition, happiness, positive affect, self-compassion, and ability to live a desired life despite pain increased, and pain catastrophizing, depression, and anxiety decreased. These effects lasted at least until the 6 months follow-up assessment period [119]. The intervention did not decrease pain intensity and disability, which is not an uncommon outcome when examining psychological treatments for chronic pain [125, 126]. Indeed, as implied by the name, the therapeutic goal of 'happy despite pain' is not pain relief, but disengaging from the pursuit to abolish the pain experience and fostering an individual's ability to live a desired life despite the presence of pain.

## Resilience building interventions

In addition to the interventions that (mainly) focus on optimism, other positive psychology interventions (PPIs) have been developed that focus on building resilience in chronic pain patients. In a systematic review [127], eight studies were identified that tested the effect of PPIs in chronic pain patients. They concluded that PPIs led to improvements in happiness, well-being, hope, life-satisfaction, and self-efficacy in dealing with pain. Müller and Gertz [128] examined the effects of an individually tailored PPI in which they matched the exercises that participants received with their interests by having them complete the Person-Activity Fit Diagnostic (PAFD). In total, there were 10 possible exercises, including the best possible future self-exercise. Each participant was assigned four exercises based on their personal activity fit. A first feasibility study in patients with chronic pain showed that this individually tailored PPI led to increased life satisfaction, positive affect, and pain control and decreased pain intensity, depression, pain interference, and pain catastrophizing, with small to moderate effect sizes. Most benefits remained until 2.5 months follow-up. However, for many of the outcomes the control group showed similar benefits, and at follow-up there were no longer significant differences between the groups [128]. A larger follow-up study with participants experiencing chronic pain secondary to spinal cord injury largely replicated these results [129]. Compared to an active control group, pain intensity was significantly lower after the tailored PPI. Pain catastrophizing and pain interference also decreased and pain control increased, but this was not significantly different from the control intervention. Nevertheless this study is remarkable in that it shows a reduction in pain intensity after a PPI.

## Integrating positive psychology interventions in traditional treatment approaches

The above-mentioned PPIs indicate that not addressing pain cognitions or pain behaviours specifically can still have beneficial effects for patients with chronic pain. In fact, in a direct comparison, ‘happy despite pain’ was as effective as CBT in decreasing catastrophic cognitions about pain. CBT directly targets maladaptive cognitions such as pain catastrophizing by teaching patients to identify these cognitions, challenge them, and replace them with more helpful cognitions. PPIs take a more transdiagnostic focus:

they teach people to lead a good and fulfilling life despite any adversity they encounter. As such, their beneficial effects may extend beyond reducing pain-related suffering and can foster personal growth in general. There is also evidence that patients prefer PPIs over traditional CBT and report to be more stimulated and motivated to change [130]. PPIs might be preferred over traditional pain-focused approaches because they are easily accessible and engaging and come with possibly less stigma than traditional behavioural treatment approaches [131, 132]. Moreover, many of these interventions have a self-help format and do not require the involvement of a mental health professional [133]. As the studies above have shown, PPIs can be effective treatments for chronic pain, even as stand-alone interventions. Nevertheless, it should be admitted that, so far, evidence is still sparse. This is not surprising as it is an emerging field. Yet, as scholars in the field have argued, even greater benefits might be gained by combining PPIs with more traditional approaches [133, 134].

One such approach was employed, for instance, by Herrero et al. [135]. They used virtual reality (VR) to induce positive affect in conjunction with an activity management programme in patients with fibromyalgia. The VR was given with the aim to increase motivation related to the performance of the chosen activities. During the VR component, a positive nature scenario was projected in the VR surrounding, including music with positive valence and high arousal. This VR component was successful in improving self-efficacy and motivation related to meaningful activities, increasing positive emotions, and reducing negative emotions and dysfunctional coping strategies. As this was a pilot study, no control condition was added. In a follow-up study [136], the activity management intervention with VR component was compared to a treatment-as-usual condition. Results showed that participants in the VR condition achieved significantly greater improvements in perceived disability, quality of life, and functional coping strategies. There were no differences regarding pain intensity, pain interference, and depression symptoms.

The recognition that including or increasing protective factors is important has also been incorporated in CBT, of which acceptance-based CBTs (e.g. acceptance and commitment therapy (ACT) and mindfulness-based cognitive therapy) are examples. All these therapies have in common that they do not focus on controlling or fighting the pain experience, but on the acceptance of pain. They commonly use certain acceptance techniques (e.g. mindfulness, meditation) to facilitate a separation between the ‘self’ and one’s thoughts, feelings and the actual pain experience. Moreover, patients are encouraged to base their actions on their values as opposed to their immediate feelings, thoughts and pain

[137]. In a meta-analysis on the effects of acceptance-based therapies on the mental and physical health in patients with chronic pain, it was found that acceptance-based therapies have small to medium effects on decreasing pain intensity, depression, and anxiety and increasing physical well-being and quality of life. The authors conclude that based on these findings, acceptance-based interventions and CBT produce small but equivalent effects [137]. In a recent systematic review, there was no evidence of a difference between ACT and an active control for pain, disability, or distress at treatment end [9]. Another approach is mindfulness-based stress reduction (MBSR), which also focusses on the acceptance of moment-to-moment experiences, but additionally includes increasing bodily and emotional awareness, and learning self-regulation strategies and more adaptive responses to stress [138, 139]. A systematic review comparing MBSR to CBT showed that both caused beneficial changes in physical functioning, pain intensity, and depression relative to control conditions, but there was no evidence that MBSR and CBT differed in their impact on these outcomes [140]. Thus, these acceptance-based CBT interventions are helpful, but not more effective than traditional CBT.

Another new development that aims to integrate positive psychology within CBT is what has been termed positive CBT. Whereas acceptance-based CBT interventions do not focus on increasing positive emotions per se, positive CBT was developed using a solution-focused framework, amplified with positive psychology exercises [130]. Increasing positive emotions and positive mental health is a specific aim of this intervention. In a crossover within-subjects design study on patients with depressive disorders, positive CBT was compared to traditional, problem-focused CBT. Results showed that the positive CBT led to significantly higher rates of clinically significant or reliable change for depression, negative affect, and happiness [141]. Although not examined, this treatment may be a promising approach for chronic pain.

A treatment approach incorporating positive psychology techniques that is specifically developed for pain patients is Mindfulness-Oriented Recovery Enhancement (MORE) [142]. This approach has been driven by the notion that because positive emotions and rewarding experiences can produce analgesia, they should be intentionally cultivated for their therapeutic value. See Garland [142] for more details, but in essence, it is argued that given the observation that chronic pain and the overuse and misuse of opioid analgesics can impair reward learning and reward responsiveness and exacerbate pain-related hedonic dysregulation, psychological treatment should focus on modulating the positive affective/reward-related mechanisms as means of addressing chronic pain and opioid misuse. Garland [142]

proposes three core positive affective/reward mechanisms that should be activated in treatment: savouring (i.e. focusing on pleasant features and associated emotions and bodily responses of naturally rewarding stimuli), noticing and generating pleasant internal states (i.e. cultivating interoceptive awareness of pleasurable sensations, and using imagination to superimpose pleasure on painful body sites), and cultivating meaning and self-transcendence (i.e. shifting patients' thinking from a narrowed focus on pain to a much broader awareness of how pleasant and painful life experiences interact to produce meaning in life). MORE encompasses these three core therapeutic mechanisms and has been shown to significantly decrease chronic pain, opioid misuse, and craving [143–145].

## A final note on the way forward

This review summarised the current state-of-the-art of pain psychology from a positive psychology perspective and has shown that interventions targeting positive and protective factors are feasible and acceptable for chronic pain patients. Interestingly, most of these positive focussed interventions have demonstrated to increase positive factors, such as positive affect, optimism, quality of life and well-being despite pain, as well as decrease negative risk factors for pain disability, such as pain catastrophizing and anxiety. The finding that pain mostly persists after treatment validates the importance of having self-help techniques to reduce the burden of living with chronic pain. Moreover, the improvements of PPIs specifically were achieved via minimal therapeutic guidance and relatively low attrition rates (all <25%) [119, 124, 128], making them an attractive treatment option for chronic pain patients, either as stand-alone treatment or in conjunction with other treatment protocols. Especially when considering that positive and negative affect may have unique roles in modulating the experience of pain [19, 146], a treatment combining insights from both research areas may have additional benefits. As discussed earlier, optimism may increase the confidence regarding the attainability of a desired goal (e.g. treatment goal), which fosters action and continued effort towards this goal, resulting in more motivation and treatment adherence.

We propose that the way forward in pain research and treatment is the inclusion of both *vulnerability* and *protective factors*, since they have unique roles in modulating the experience of pain, a finding that had been neglected for too long. The effort to identify both vulnerability and protective factors in the context of chronic pain has obvious theoretical and scientific benefits. But maybe the most important contribution of these types of studies is that they raise



awareness among clinicians of the important role of protective factors in the context of pain. This does not mean that existing approaches and theories have to be abandoned. For instance, clinicians may use the fear-avoidance model to explain patient's pain condition and how their vulnerability factors can instigate a vicious downward spiral that results in maladjustment to pain. But clinicians should not stop there. Together with their clients, they may explore their resilience factors and explain how these may incite a positive upward spiral, fuelling positive adaptation to pain.

There are still important steps to take. Many of the findings related to protective factors such as optimism come from studies in which experimental pain was induced. It is well known that experimentally induced pain in most cases is predictable, temporary, and evaluated as less threatening than chronic pain [147]. Chronic pain has far greater emotional and motivational significance [147]. Therefore, most researchers see findings regarding experimental pain as a proof of concept, which should be replicated in pain populations. However, such studies are rarely performed, likely due to the difficulties inherent in research with chronic pain patients, such as recruitment difficulties, time constraints, and ethical objections. There is a clear need to validate research findings in clinical samples. This is, of course, true not only for research findings related to protective factors, but because this is a relatively new field, much work remains to be done.

Although these are early days for application of PPIs in chronic pain treatment, exciting new developments are ongoing. The transdiagnostic focus on what makes life worthwhile is a recognition of the fact that a good and meaningful life is not dependent on the absence of physical ailments such as pain. Positive thinking, feeling, acting and pursuing valued goals can make one's life gratifying and fulfilling, despite experiencing chronic pain.

**Research funding:** Authors state no funding involved.

**Author contributions:** All authors have accepted responsibility for the entire content of this manuscript and approved its submission.

**Competing interests:** Authors state no conflict of interest.

**Informed consent:** Not applicable.

**Ethical approval:** Not applicable.

## References

1. Brinjikji W, Diehn F, Jarvik J, Carr C, Kallmes DF, Murad MH, et al. MRI findings of disc degeneration are more prevalent in adults with low back pain than in asymptomatic controls: a systematic review and meta-analysis. *Am J Neuroradiol* 2015;36:2394–9.
2. Henschke N, Maher CG, Refshauge KM, Herbert RD, Cumming RG, Bleasel J, et al. Prevalence of and screening for serious spinal pathology in patients presenting to primary care settings with acute low back pain. *Arthritis Rheum* 2009;60:3072–80.
3. Woolf CJ. Central sensitization: implications for the diagnosis and treatment of pain. *Pain* 2011;152:S2–15.
4. Nijs J, George SZ, Clauw DJ, Fernández-de-las-Peñas C, Kosek E, Ickmans K, et al. Central sensitisation in chronic pain conditions: latest discoveries and their potential for precision medicine. *Lancet Rheumatol* 2021;3:383–92.
5. Vlaeyen JWS, Linton SJ. Fear-avoidance and its consequences in chronic musculoskeletal pain: a state of the art. *Pain* 2000;85:317–32.
6. Vlaeyen J, Morley S, Linton SJ, Boersma K, de Jong J. Pain-related fear: exposure based treatment for chronic pain. Washington: IASP press; 2012.
7. Eccleston C, Morley S, Williams AdC. Psychological approaches to chronic pain management: evidence and challenges. *Br J Anaesth* 2013;111:59–63.
8. Morley S. Efficacy and effectiveness of cognitive behaviour therapy for chronic pain: progress and some challenges. *Pain* 2011;152: S99–106.
9. de C Williams AC, Fisher E, Hearn L, Eccleston C. Psychological therapies for the management of chronic pain [excluding headache] in adults. *Cochrane Database Syst Rev* 2020;8:1–170.
10. Vlaeyen JW, Crombez G. Behavioral conceptualization and treatment of chronic pain. *Annu Rev Clin Psychol* 2020;16:187–212.
11. Leeuw M, Goossens ME, van Breukelen GJ, de Jong JR, Heuts PH, Smeets RJ, et al. Exposure in vivo versus operant graded activity in chronic low back pain patients: results of a randomized controlled trial. *Pain* 2008;138:192–207.
12. den Hollander M, Goossens M, de Jong J, Ruijgrok J, Oosterhof J, Onghena P, et al. Expose or protect? A randomized controlled trial of exposure in vivo vs pain-contingent treatment as usual in patients with complex regional pain syndrome type 1. *Pain* 2016;157:2318–29.
13. den Hollander M, Heijnders N, de Jong JR, Vlaeyen JW, Smeets RJ, Goossens ME. Exposure in vivo versus pain-contingent physical therapy in complex regional pain syndrome type I: a cost-effectiveness analysis. *Int J Technol Assess Health Care* 2018;34: 400–9.
14. de Jong JR, Vlaeyen JW, van Eijdsden M, Loo C, Onghena P. Reduction of pain-related fear and increased function and participation in work-related upper extremity pain [WRUEP]: effects of exposure in vivo. *Pain* 2012;153:2109–18.
15. Silva Guerrero AV, Maujean A, Campbell L, Sterling M. A systematic review and meta-analysis of the effectiveness of psychological interventions delivered by physiotherapists on pain, disability and psychological outcomes in musculoskeletal pain conditions. *Clin J Pain* 2018;34:838–57.
16. Seligman MEP, Csikszentmihalyi M. Positive psychology – an introduction. *Am Psychol* 2000;55:5–14.
17. Seligman ME, Parks AC, Steen T. A balanced psychology and a full life. *Philos Trans R Soc B* 2004;359:1379–81.
18. Seligman ME, Csikszentmihalyi M. Positive psychology: an introduction. *Flow and the foundations of positive psychology*. New York City: Springer; 2014:279–98 pp.
19. Finan PH, Garland EL. The role of positive affect in pain and its treatment. *Clin J Pain* 2015;31:177.
20. Garland EL, Fredrickson B, Kring AM, Johnson DP, Meyer PS, Penn DL. Upward spirals of positive emotions counter downward spirals of negativity: insights from the broaden-and-build theory and affective

- neuroscience on the treatment of emotion dysfunctions and deficits in psychopathology. *Clin Psychol Rev* 2010;30:849–64.
21. Bandura A. Self-efficacy: toward a unifying theory of behavior change. *Psychol Rev* 1977;84:191–215.
  22. Snyder CR, Rand KL, Sigmon DR. Hope theory. *Handbook of positive psychology*. New York City: Oxford University Press; 2002:257–76 pp.
  23. Scheier MF, Carver CS. Optimism, coping, and health: assessment and implications of generalized outcome expectancies. *Health Psychol* 1985;4:219.
  24. Basten-Günther J, Peters M, Lautenbacher S. Optimism and the experience of pain: a systematic review. *Behav Med* 2019;45:323–39.
  25. Pulvers K, Hood A. The role of positive traits and pain catastrophizing in pain perception. *Curr Pain Headache Rep* 2013;17:1–11.
  26. Garofalo JP. Perceived optimism and chronic pain. In: Gatchel RJ, Weisberg JN, editors. *Personality characteristics of patients with pain*. Washington, DC US: American Psychological Association; 2000:203–17 pp.
  27. Goodin BR, Bulls HW. Optimism and the experience of pain: benefits of seeing the glass as half full. *Curr Pain Headache Rep* 2013;17:329.
  28. Ong AD, Thoemmes F, Ratner K, Ghezzi-Kopel K, Reid MC. Positive affect and chronic pain: a preregistered systematic review and meta-analysis. *Pain* 2020;161:1140–9.
  29. Ong AD, Zautra AJ, Reid MC. Chronic pain and the adaptive significance of positive emotions. *Am Psychol* 2015;70:283–4.
  30. Curtin KB, Norris D. The relationship between chronic musculoskeletal pain, anxiety and mindfulness: adjustments to the Fear-Avoidance Model of chronic pain. *Scand J Pain* 2017;17:156–66.
  31. Rand KL. Hope, self-efficacy, and optimism: conceptual and empirical differences. In: Gallagher MW, Lopez SJ, editors. *The Oxford handbook of hope*. New York City: Oxford University Press; 2018:45–58 pp.
  32. Fredrickson BL. The role of positive emotions in positive psychology: the broaden-and-build theory of positive emotions. *Am Psychol* 2001;56:218.
  33. Sturgeon JA, Zautra AJ. Resilience: a new paradigm for adaptation to chronic pain. *Curr Pain Headache Rep* 2010;14:105–12.
  34. Crombez G, Eccleston C, Van Damme S, Vlaeyen JW, Karoly P. Fear-avoidance model of chronic pain: the next generation. *Clin J Pain* 2012;28:475–83.
  35. Leeuw M, Goossens ME, Linton SJ, Crombez G, Boersma K, Vlaeyen JW. The fear-avoidance model of musculoskeletal pain: current state of scientific evidence. *J Behav Med* 2007;30:77–94.
  36. Vlaeyen JW, Crombez G, Linton SJ. The fear-avoidance model of pain. *Pain* 2016;157:1588–9.
  37. Van Damme S, Legrain V, Vogt J, Crombez G. Keeping pain in mind: a motivational account of attention to pain. *Neurosci Biobehav Rev* 2010;34:204–13.
  38. Boselie JJ, Vlaeyen JW. Broadening the fear-avoidance model of chronic pain? *Scand J Pain* 2017;17:176–7.
  39. Wideman TH, Asmundson GG, Smeets RJM, Zautra AJ, Simmonds MJ, Sullivan MJ, et al. Re-thinking the fear avoidance model: toward a multi-dimensional framework of pain-related disability. *Pain* 2013;154:2262.
  40. Hanssen MM, Peters ML, Boselie JJ, Meulders A. Can positive affect attenuate [persistent] pain? State of the art and clinical implications. *Curr Rheumatol Rep* 2017;19:1–9.
  41. Carver CS, Scheier MF. *On the self-regulation of behavior*. New York, NY US: Cambridge University Press; 1998.
  42. Austin JT, Vancouver JB. Goal constructs in psychology: structure, process, and content. *Psychol Bull* 1996;120:338.
  43. Mahler H, Kulik J. Optimism, pessimism and recovery from coronary bypass surgery: prediction of affect, pain and functional status. *Psychol Health Med* 2000;5:347–58.
  44. Shanahan ML, Fischer IC, Hirsh AT, Stewart JC, Rand KL. Hope, optimism, and clinical pain: a meta-analysis. *Ann Behav Med* 2021;55:815–32.
  45. Geers AL, Wellman JA, Fowler SL, Helfer SG, France CR. Dispositional optimism predicts placebo analgesia. *J Pain* 2010;11:1165–71.
  46. Goodin BR, Glover TL, Sotolongo A, King CD, Sibille KT, Herbert MS, et al. The association of greater dispositional optimism with less endogenous pain facilitation is indirectly transmitted through lower levels of pain catastrophizing. *J Pain* 2013;14:126–35.
  47. Hanssen MM, Peters ML, Vlaeyen JW, Meevissen YM, Vancleef LM. Optimism lowers pain: evidence of the causal status and underlying mechanisms. *Pain* 2013;154:53–8.
  48. Vancleef LM, Peters ML. Examining content specificity of negative interpretation biases with the Body Sensations Interpretation Questionnaire [BSIQ]. *J Anxiety Disord* 2008;22:401–15.
  49. Segerstrom SC, Tsao JC, Alden LE, Craske MG. Worry and rumination: repetitive thought as a concomitant and predictor of negative mood. *Cognit Ther Res* 2000;24:671–88.
  50. McCracken LM, Iverson GL. Predicting complaints of impaired cognitive functioning in patients with chronic pain. *J Pain Symptom Manage* 2001;21:392–6.
  51. Kane MJ, Brown LH, McVay JC, Silvia PJ, Myin-Germeys I, Kwapil TR. For whom the mind wanders, and when: an experience-sampling study of working memory and executive control in daily life. *Psychol Sci* 2007;18:614–21.
  52. Eccleston C, Crombez G, Aldrich S, Stannard C. Worry and chronic pain patients: a description and analysis of individual differences. *Eur J Pain* 2001;5:309–18.
  53. Hanssen MM, Vancleef LM, Vlaeyen JW, Peters ML. More optimism, less pain! The influence of generalized and pain-specific expectations on experienced cold-pressor pain. *J Behav Med* 2014;37:47–58.
  54. McGregor BA, Bowen DJ, Ankerst DP, Andersen MR, Yasui Y, McTiernan A. Optimism, perceived risk of breast cancer, and cancer worry among a community-based sample of women. *Health Psychol* 2004;23:339.
  55. Tucker RP, Wingate LR, O’Keefe VM, Mills AC, Rasmussen K, Davidson CL, et al. Rumination and suicidal ideation: the moderating roles of hope and optimism. *Pers Individ Differ* 2013;55:606–11.
  56. Ramírez-Maestre C, Esteve R, López AE. The role of optimism and pessimism in chronic pain patients adjustment. *Spanish J Psychol* 2012;15:286–94.
  57. Carver CS, Scheier MF, Segerstrom SC. Optimism. *Clin Psychol Rev* 2010;30:879–89.
  58. Baumeister RF, Heatherton TF, Tice DM. *Losing control*. San Diego, CA: Academic Press; 1994.
  59. Solberg Nes L, Carlson CR, Crofford LJ, de Leeuw R, Segerstrom SC. Individual differences and self-regulatory fatigue: optimism, conscientiousness, and self-consciousness. *Pers Individ Differ* 2011;50:475–80.
  60. Evans DR, Boggero IA, Segerstrom SC. The nature of self-regulatory fatigue and “ego depletion” lessons from physical fatigue. *Pers Soc Psychol Rev* 2016;20:291–310.
  61. Solberg Nes L, Carlson CR, Crofford LJ, de Leeuw R, Segerstrom SC. Self-regulatory deficits in fibromyalgia and temporomandibular disorders. *Pain* 2010;151:37–44.

62. Segerstrom SC, Nes LS. When goals conflict but people prosper: the case of dispositional optimism. *J Res Pers* 2006;40:675–93.
63. Affleck G, Tennen H, Zautra A, Urrows S, Abeles M, Karoly P. Women's pursuit of personal goals in daily life with fibromyalgia: a value-expectancy analysis. *J Consult Clin Psychol* 2001;69:587–96.
64. Duke J, Leventhal H, Brownlee S, Leventhal EA. Giving up and replacing activities in response to illness. *J Gerontol B Psychol Sci Soc Sci* 2002; 57B:P367–76.
65. Gilam G, Gross JJ, Wager TD, Keefe FJ, Mackey SC. What is the relationship between pain and emotion? Bridging constructs and communities. *Neuron* 2020;107:17–21.
66. Fields H. State-dependent opioid control of pain. *Nat Rev Neurosci* 2004;5:565–75.
67. McCracken LM, Carson JW, Eccleston C, Keefe FJ. Acceptance and change in the context of chronic pain. *Pain* 2004;109:4–7.
68. Van Damme S, Crombez G, Eccleston C. Coping with pain: a motivational perspective. *Pain* 2008;139:1–4.
69. Solberg Nes L, Segerstrom SC. Dispositional optimism and coping: a meta-analytic review. *Pers Soc Psychol Rev* 2006;10:235–51.
70. Reimers S, Maylor EA. Task switching across the life span: effects of age on general and specific switch costs. *Dev Psychol* 2005;41:661–71.
71. Falletti MG, Maruff P, Collie A, Darby DG. Practice effects associated with the repeated assessment of cognitive function using the CogState battery at 10-minute, one week and one month test-retest intervals. *J Clin Exp Neuropsychol* 2006;28:1095–112.
72. Wrosch C, Scheier MF, Carver CS, Schulz R. The importance of goal disengagement in adaptive self-regulation: when giving up is beneficial. *Self Ident* 2003;2:1–20.
73. Wrosch C, Scheier MF. Personality and quality of life: the importance of optimism and goal adjustment. *Qual Life Res* 2003;12(1 Suppl): 59–72.
74. Wrosch C. Self-regulation of unattainable goals and pathways to quality of life. In: Folkman S, editor. *The Oxford handbook of stress, health, and coping*. Oxford library of psychology. New York, NY US: Oxford University Press; 2011:319–33 pp.
75. Rasmussen HN, Wrosch C, Scheier MF, Carver CS. Self-regulation processes and health: the importance of optimism and goal adjustment. *J Pers* 2006;74:1721–47.
76. Ramírez-Maestre C, Esteve R, López-Martínez AE, Serrano-Ibáñez ER, Ruiz-Párraga GT, Peters M. Goal adjustment and well-being: the role of optimism in patients with chronic pain. *Ann Behav Med* 2019;53: 597–607.
77. Hanssen M, Vancleef L, Vlaeyen J, Hayes A, Schouten E, Peters M. Optimism, motivational coping and well-being: evidence supporting the importance of flexible goal adjustment. *J Happiness Stud* 2015;16: 1525–37.
78. Geers AL, Wellman JA, Helfer SG, Fowler SL, France CR. Dispositional optimism and thoughts of well-being determine sensitivity to an experimental pain task. *Ann Behav Med* 2008;36:304–13.
79. Geers AL, Wellman JA, Lassiter GD. Dispositional optimism and engagement: the moderating influence of goal prioritization. *J Pers Soc Psychol* 2009;96:913.
80. Geers AL, Wellman JA, Seligman LD, Wuyek LA, Neff LA. Dispositional optimism, goals, and engagement in health treatment programs. *J Behav Med* 2010;33:123–34.
81. Peters ML, Vieler JS, Lautenbacher S. Dispositional and induced optimism lead to attentional preference for faces displaying positive emotions: an eye-tracker study. *J Posit Psychol* 2016;11:258–69.
82. Luo J, Isaacowitz DM. How optimists face skin cancer information: risk assessment, attention, memory, and behavior. *Psychol Health* 2007; 22:963–84.
83. Epping-Jordan JE, Compas BE, Osowiecki DM, Oppedisano G, Gerhardt C, Primo K, et al. Psychological adjustment in breast cancer: processes of emotional distress. *Health Psychol* 1999;18:315.
84. Allison PJ, Guichard C, Gilain L. A prospective investigation of dispositional optimism as a predictor of health-related quality of life in head and neck cancer patients. *Qual Life Res* 2000;9:951–60.
85. Carver CS, Pozo-Kaderman C, Harris SD, Noriega V, Scheier MF, Robinson DS, et al. Optimism versus pessimism predicts the quality of women's adjustment to early stage breast cancer. *Cancer* 1994;73: 1213–20.
86. Litt MD, Tennen H, Affleck G, Klock S. Coping and Cognitive factors in adaptation to in vitro fertilization failure. *J Behav Med* 1992;15:171–87.
87. Mosing MA, Zietsch BP, Shekar SN, Wright MJ, Martin NG. Genetic and environmental influences on optimism and its relationship to mental and self-rated health: a study of aging twins. *Behav Genet* 2009;39: 597–604.
88. Plomin R, Scheier MF, Bergeman CS, Pedersen NL, Nesselrode JR, McClearn GE. Optimism, pessimism and mental health: a twin/ adoption analysis. *Pers Individ Differ* 1992;13:921–30.
89. Heinonen K, Räikkönen K, Keltikangas-Järvinen L. Dispositional optimism: development over 21 years from the perspectives of perceived temperament and mothering. *Pers Individ Differ* 2005;38:425–35.
90. Heinonen K, Räikkönen K, Keltikangas-Järvinen L. Self-esteem in early and late adolescence predicts dispositional optimism–pessimism in adulthood: a 21-year longitudinal study. *Pers Individ Differ* 2005;39: 511–21.
91. Heinonen K, Räikkönen K, Matthews KA, Scheier MF, Raitakari OT, Pulkki L, et al. Socioeconomic status in childhood and adulthood: associations with dispositional optimism and pessimism over a 21-year follow-up. *J Pers* 2006;74:1111–26.
92. Atienza AA, Stephens MAP, Townsend AL. Role stressors as predictors of changes in women's optimistic expectations. *Pers Individ Differ* 2004; 37:471–84.
93. Segerstrom SC. Optimism and resources: effects on each other and on health over 10 years. *J Res Pers* 2007;41:772–86.
94. Malouff JM, Schutte NS. Can psychological interventions increase optimism? A meta-analysis. *J Posit Psychol* 2017;12:594–604.
95. Peters ML, Flink IK, Boersma K, Linton SJ. Manipulating optimism: can imagining a best possible self be used to increase positive future expectancies? *J Posit Psychol* 2010;5:204–11.
96. Carrillo A, Rubio-Aparicio M, Molinari G, Enrique Á, Sánchez-Meca J, Baños RM. Effects of the best possible self intervention: a systematic review and meta-analysis. *PLoS One* 2019;14:e0222386.
97. Boselie JLM. Filling the glass: optimism as a protective factor against pain-induced executive functioning impairments. Maastricht: Datawyse/Universitaire Pers Maastricht; 2017.
98. Heekerens JB, Eid M, Heinitz K. Dealing with conflict: reducing goal ambivalence using the best-possible-self intervention. *J Posit Psychol* 2020;15:325–37.
99. Loveday PM, Lovell GP, Jones CM. The best possible selves intervention: a review of the literature to evaluate efficacy and guide future research. *J Happiness Stud* 2018;19:607–28.
100. Boselie JJ, Vancleef LM, Smeets T, Peters ML. Increasing optimism abolishes pain-induced impairments in executive task performance. *Pain* 2014;155:334–40.

101. Boselie JJ, Vancleef LM, Peters ML. Increasing optimism protects against pain-induced impairment in task-shifting performance. *J Pain* 2017;18:446–55.
102. Miyake A, Friedman NP, Emerson MJ, Witzki AH, Howerter A. The unity and diversity of executive functions and their contributions to complex 'frontal lobe' tasks: a latent variable analysis. *Cognit Psychol* 2000;41:49–100.
103. Ryckeghem D, Crombez G, Eccleston C, Legrain V, Damme S. Keeping pain out of your mind: the role of attentional set in pain. *Eur J Pain* 2013;17:402–11.
104. Legrain V, Crombez G, Verhoeven K, Mouraux A. The role of working memory in the attentional control of pain. *Pain* 2011;152:453–9.
105. Vancleef LM, Peters ML. The interruptive effect of pain on attention. *J Pain* 2006;7:21–2.
106. Moore DJ, Keogh E, Eccleston C. The interruptive effect of pain on attention. *Q J Exp Psychol* 2012;65:565–86.
107. Buhle J, Wager TD. Performance-dependent inhibition of pain by an executive working memory task. *Pain* 2010;149:19–26.
108. Eccleston C, Crombez G. Pain demands attention: a cognitive–affective model of the interruptive function of pain. *Psychol Bull* 1999;125:356–66.
109. Eccleston C, Crombez G. Worry and chronic pain: a misdirected problem solving model. *Pain* 2007;132:233–6.
110. Carver CS, Lawrence JW, Scheier MF. Self-discrepancies and affect: incorporating the role of feared selves. *Pers Soc Psychol Bull* 1999;25:783–92.
111. Higgins ET. Self-discrepancy theory: what patterns of self-beliefs cause people to suffer? In: Leonard B, editor. *Advances in experimental social psychology*. Cambridge: Academic Press; 1989, vol 22:93–136 pp.
112. Goossens ME, Kindermans HP, Morley SJ, Roelofs J, Verbunt J, Vlaeyen JW. Self-discrepancies in work-related upper extremity pain: relation to emotions and flexible-goal adjustment. *Eur J Pain* 2010;14:764–70.
113. Waters SJ, Keefe FJ, Strauman TJ. Self-discrepancy in chronic low back pain: relation to pain, depression, and psychological distress. *J Pain Symptom Manage* 2004;27:251–9.
114. Kindermans HP, Huijnen IP, Goossens ME, Roelofs J, Verbunt JA, Vlaeyen JW. "Being" in pain: the role of self-discrepancies in the emotional experience and activity patterns of patients with chronic low back pain. *Pain* 2011;152:403–9.
115. Morley S. The self in pain. *Rev Pain* 2010;4:24–7.
116. Morley S, Davies C, Barton S. Possible selves in chronic pain: self-pain enmeshment, adjustment and acceptance. *Pain* 2005;115:84–94.
117. Sutherland R, Morley S. Self-pain enmeshment: future possible selves, sociotropy, autonomy and adjustment to chronic pain. *Pain* 2008;137:366–77.
118. Molinari G, García-Palacios A, Enrique Á, Roca P, Fernández-Llanio Comella N, Botella C. The power of visualization: back to the future for pain management in fibromyalgia syndrome. *Pain Med* 2018;19:1451–68.
119. Peters ML, Smeets E, Feijge M, van Breukelen G, Andersson G, Buhrman M, et al. Happy despite pain: a randomized controlled trial of an 8-week internet-delivered positive psychology intervention for enhancing well-being in patients with chronic pain. *Clin J Pain* 2017;33:962.
120. Neff KD, Germer CK. A pilot study and randomized controlled trial of the mindful self-compassion program. *J Clin Psychol* 2013;69:28–44.
121. Neff KD, Vonk R. Self-compassion versus global self-esteem: two different ways of relating to oneself. *J Pers* 2009;77:23–50.
122. Seligman ME, Rashid T, Parks AC. Positive psychotherapy. *Am Psychol* 2006;61:774.
123. Seligman ME, Steen TA, Park N, Peterson C. Positive psychology progress: empirical validation of interventions. *Am Psychol* 2005;60:410.
124. Boselie JJLM, Vancleef LMG, Peters ML. Filling the glass: effects of a positive psychology intervention on executive task performance in chronic pain patients. *Eur J Pain* 2018;22:1268–80.
125. Eccleston C, Williams ACdC, Morley S. Psychological therapies for the management of chronic pain [excluding headache] in adults. *Cochrane Database Syst Rev* 2009;8:1–170.
126. Williams AC, Eccleston C, Morley S. Psychological therapies for the management of chronic pain [excluding headache] in adults. *Cochrane Database Syst Rev* 2012;11:CD007407.
127. Iddon JE, Dickson JM, Unwin J. Positive psychological interventions and chronic non-cancer pain: a systematic review of the literature. *Int J Appl Posit Psychol* 2016;1:133–57.
128. Müller R, Gertz KJ, Molton IR, Terrill AL, Bombardier CH, Ehde DM, et al. Effects of a tailored positive psychology intervention on well-being and pain in individuals with chronic pain and a physical disability: a feasibility trial. *Clin J Pain* 2016;32:32–44.
129. Lyubomirsky S. *The how of happiness: a scientific approach to getting the life you want*. New York City: Penguin; 2008.
130. Geschwind N, Bosgraaf E, Bannink F, Peeters F. Positivity pays off: clients' perspectives on positive compared with traditional cognitive behavioral therapy for depression. *Psychotherapy* 2020;57:366.
131. Rini C, Williams DA, Broderick JE, Keefe FJ. Meeting them where they are: using the internet to deliver behavioral medicine interventions for pain. *Transl Behav Med* 2012;2:82–92.
132. Stabolidis A. All in their mind? The stigma of pain. *Women's Health* 2019;21.
133. Hassett AL, Finan PH. The role of resilience in the clinical management of chronic pain. *Curr Pain Headache Rep* 2016;20:39.
134. Müller R, Segerer W, Ronca E, Gemperli A, Stirnimann D, Scheel-Sailer A, et al. Inducing positive emotions to reduce chronic pain: a randomized controlled trial of positive psychology exercises. *Disabil Rehabil* 2020;44:1–14.
135. Herrero R, García-Palacios A, Castilla D, Molinari G, Botella C. Virtual reality for the induction of positive emotions in the treatment of fibromyalgia: a pilot study over acceptability, satisfaction, and the effect of virtual reality on mood. *Cyberpsychol Behav Soc Netw* 2014;17:379–84.
136. García-Palacios A, Herrero R, Vizcaíno Y, Belmonte MA, Castilla D, Molinari G, et al. Integrating virtual reality with activity management for the treatment of fibromyalgia. *Clin J Pain* 2015;31:564–72.
137. Veehof MM, Oskam M-J, Schreurs KM, Bohlmeijer ET. Acceptance-based interventions for the treatment of chronic pain: a systematic review and meta-analysis. *Pain* 2011;152:533–42.
138. Kabat-Zinn J. Mindfulness-based stress reduction [MBSR]. *Constr Hum Sci* 2003;8:73.
139. Kabat-Zinn J. Mindfulness-based interventions in context: past, present, and future. *Clin Psychol Sci Pract* 2003;10:144–56.
140. Khoo E-L, Small R, Cheng W, Hatchard T, Glynn B, Rice DB, et al. Comparative evaluation of group-based mindfulness-based stress reduction and cognitive behavioural therapy for the treatment and management of chronic pain: a systematic review and network meta-analysis. *Evid Base Ment Health* 2019;22:26–35.
141. Geschwind N, Arntz A, Bannink F, Peeters F. Positive cognitive behavior therapy in the treatment of depression: a randomized order

- within-subject comparison with traditional cognitive behavior therapy. *Behav Res Ther* 2019;116:119–30.
142. Garland EL. Psychosocial intervention and the reward system in pain and opioid misuse: new opportunities and directions. *Pain* 2020;161:2659–66.
143. Garland EL, Manusov EG, Froeliger B, Kelly A, Williams JM, Howard MO. Mindfulness-oriented recovery enhancement for chronic pain and prescription opioid misuse: results from an early-stage randomized controlled trial. *J Consult Clin Psychol* 2014;82:448.
144. Garland EL, Hanley AW, Kline A, Cooperman NA. Mindfulness-oriented recovery enhancement reduces opioid craving among individuals with opioid use disorder and chronic pain in medication assisted treatment: ecological momentary assessments from a stage 1 randomized controlled trial. *Drug Alcohol Depend* 2019;203:61–5.
145. Garland EL, Hanley AW, Riquino MR, Reese SE, Baker AK, Salas K, et al. Mindfulness-oriented recovery enhancement reduces opioid misuse risk via analgesic and positive psychological mechanisms: a randomized controlled trial. *J Consult Clin Psychol* 2019;87:927.
146. Davis MC, Zautra AJ, Smith BW. Chronic pain, stress, and the dynamics of affective differentiation. *J Pers* 2004;72:1133–60.
147. Edens JL, Gil KM. Experimental induction of pain: utility in the study of clinical pain. *Behav Ther* 1995;26:197–216.