

Oropharyngeal dysphagia and its psychiatric comorbidities

Citation for published version (APA):

Verdonschot, RJCG. (2018). *Oropharyngeal dysphagia and its psychiatric comorbidities: the prevalence of affective symptoms and the unmet clinical need for integrated care in medically unexplained symptoms*. Maastricht University. <https://doi.org/10.26481/dis.20181115rv>

Document status and date:

Published: 01/01/2018

DOI:

[10.26481/dis.20181115rv](https://doi.org/10.26481/dis.20181115rv)

Document Version:

Publisher's PDF, also known as Version of record

Please check the document version of this publication:

- A submitted manuscript is the version of the article upon submission and before peer-review. There can be important differences between the submitted version and the official published version of record. People interested in the research are advised to contact the author for the final version of the publication, or visit the DOI to the publisher's website.
- The final author version and the galley proof are versions of the publication after peer review.
- The final published version features the final layout of the paper including the volume, issue and page numbers.

[Link to publication](#)

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal.

If the publication is distributed under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license above, please follow below link for the End User Agreement:

www.umlib.nl/taverne-license

Take down policy

If you believe that this document breaches copyright please contact us at:

repository@maastrichtuniversity.nl

providing details and we will investigate your claim.

Valorization addendum

The studies presented in this thesis were designed to provide insight in psychiatric (co)morbidity in patients with OD and to provide guidance for OD caregivers. OD, which basically means the inability to eat and drink normally, affects over 40 million people in Europe.⁴¹ The financial impact of OD is substantial.^{42,43} OD has various underlying etiologies, including stroke, progressive degenerative neurological diseases, and oncological diseases, and may cause severe complications.^{1-4,44} In recent decades, this problem has been gaining attention worldwide and high-quality research has been done, mainly concerning the diagnostic trajectory and treatment strategies for OD patients. Although the literature suggests a high impact on OD patients' health-related quality of life, knowledge about mental distress in OD patients is scarce and discussions of the psychological burden remain speculative.

The recommended integrated approach to the OD patient is a multidisciplinary one, involving different caregivers including medical specialists (otorhinolaryngologists, neurologists, psychiatrists, radiologists, oncologists, etc.) and allied health professionals (speech-language pathologists, dieticians, nurses, etc.). In this thesis, we propose developing an integrated psychosomatic approach by adding a staff-guided mental-healthcare worker to this multidisciplinary team. To that end, the results of this thesis are not only of relevance for ENT specialists and psychiatrists but for all professionals involved in the long-lasting trajectory of OD care.

This thesis offers deeper insight into the high prevalence of mental health issues in OD patients, but it also draws attention to psychiatric problems in patients with various ENT complaints. However, prospective longitudinal studies are needed to understand mental distress in the ENT population more fully. Better understanding must underpin efforts to improve patients' healthcare status and rehabilitation outcome, which will in turn lead to decreased healthcare utilization and costs. Unfortunately, financial resources and grants are scarce in this field; consequently, little fundamental research or research of clinical relevance has been done. We hope that this thesis will lead to greater awareness among caregivers of the mental health issues that accompany OD, to more research in this field, and to better management strategies, all in order to better help this complex category of patients. Future models of integrated OD care certainly deserve to be considered for research grants.